

Existence of infinitely many solutions for a nonlocal elliptic PDE involving singularity

Sekhar Ghosh & Debajyoti Choudhuri *

Department of Mathematics, National Institute of Technology Rourkela

Emails: sekharghosh1234@gmail.com & dc.iit12@gmail.com

Abstract

In this article, we will prove the existence of infinitely many positive weak solutions to the following nonlocal elliptic PDE.

$$\begin{aligned} (-\Delta)^s u &= \frac{\lambda}{u^\gamma} + f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \end{aligned}$$

where Ω is an open bounded domain in \mathbb{R}^N with Lipschitz boundary, $N > 2s$, $s \in (0, 1)$, $\gamma \in (0, 1)$. We will employ variational techniques to show the existence of infinitely many weak solutions of the above problem.

keywords: Elliptic PDE, Genus, PS condition, Mountain Pass Theorem.

AMS classification: 35J20, 35J35, 35J60, 35J75.

1 Introduction

We consider the following nonlocal problem involving singularity.

$$\begin{aligned} (-\Delta)^s u &= \frac{\lambda}{u^\gamma} + f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \\ u &> 0 \text{ in } \Omega, \end{aligned} \tag{1.1}$$

where,

$$(-\Delta)^s u(x) = -\frac{1}{2} \int_{\mathbb{R}^N} \frac{(u(x+y) + u(x-y) - 2u(x))}{|x-y|^{N+2s}} dy, \text{ for } x \in \mathbb{R}^N,$$

$\lambda > 0$, $s \in (0, 1)$, $\gamma \in (0, 1)$ and Ω be an open, bounded subset of \mathbb{R}^N , $N \geq 2$.

The study of nonlocal elliptic PDEs is important to both, from the mathematical research as well as from the real world application, point of views. Some of the applications are in the

*Corresponding author: dc.iit12@gmail.com

probability theory, the obstacle problem, optimization, finance, phase transitions, soft thin films, conservation laws, minimal surfaces, material science and water waves. The application in probability theory, in particular to the Lévy process, can be found in [2] and that in the field of finance, one can refer to [10]. For a fruitful note on the application of fractional Laplacian one may refer [29] and the references therein. Recently, the study of singular elliptic PDE has drawn a great attention to many researchers. One of the earliest study on the existence and regularity of a weak solution was made by Lazer and Mckenna [21] to the following problem.

$$\begin{aligned} -\Delta u &= \frac{p(x)}{u^\gamma} \text{ in } \Omega, \\ u &= 0 \text{ in } \partial\Omega, \\ u &> 0 \text{ in } \Omega, \end{aligned} \tag{1.2}$$

where, $p : \Omega \rightarrow \mathbb{R}$ is a nonnegative bounded function. In [21], the authors proved that for a $C^{2+\gamma}$ boundary, the problem (1.2) has a $H_0^1(\Omega)$ solution iff $\gamma < 3$ and if $\gamma > 1$, the problem cannot have $C^1(\bar{\Omega})$ solution. The following problem have been studied for existence, uniqueness and regularity of solutions for $p = 2$ in [14] and for $1 < p < \infty$ in [8], where $0 < s < 1$.

$$\begin{aligned} (-\Delta_p)^s u &= \lambda \frac{a(x)}{u^\gamma} \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \\ u &> 0 \text{ in } \Omega, \end{aligned} \tag{1.3}$$

where, $a : \Omega \rightarrow \mathbb{R}$ is a nonnegative bounded function. The author in [14], guaranteed the existence of a unique $C^{2,\alpha}(\Omega)$, ($0 < \alpha < 1$) solution for $\lambda a(x) \equiv 1$. Canino et al. [8], had proved the existence and uniqueness of solution to the problem (1.3) by dividing λ into three cases $0 < \lambda < 1$, $\lambda = 1$ and $\lambda > 1$. A few more noteworthy study involving singularity for both Laplacian and fractional Laplacian operators can be found in [5, 11, 17, 24] and the references therein.

Multiplicity of solutions to the following type of problem has been widely studied by many authors, a few of them are in [15, 16, 23, 25] and the references therein.

$$\begin{aligned} (-\Delta_p)^s u &= \lambda \frac{a(x)}{u^\gamma} + f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \\ u &> 0 \text{ in } \Omega. \end{aligned} \tag{1.4}$$

Here $N > ps$, $M \geq 0$, $a : \Omega \rightarrow \mathbb{R}$ is a nonnegative bounded function. The authors in [16, 25], have used a variational technique to guarantee the existence of multiple solutions. Nehari manifold method has been used to prove the multiplicity result in [15, 23]. In most of these studies, the authors obtained two distinct weak solutions.

The existence results of infinitely many solutions to both Laplacian and fractional Laplacian with a nonsingular, nonlinear data have been studied widely with Dirichlet boundary condition. In most of these studies the authors proved the existence result with the help of the

symmetric Mountain Pass Theorem [19, 20]. One of the earliest attempt to show the existence of infinitely many solutions was made by Ambrosetti and Rabinowitz [1] to the following problem.

$$\begin{aligned} -\Delta u &= f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \partial\Omega, \\ u &> 0 \text{ in } \Omega, \end{aligned} \tag{1.5}$$

where, f is superlinear but subcritical near infinity. The authors in [19] have used the symmetric Mountain Pass Theorem to guarantee the existence. A few more similar type of studies made are in [20, 22, 30] and the references therein.

Recently, Gu et al. [18] has guaranteed the existence of infinitely many solutions to a nonlocal problem of the following type with a sublinear growth of f .

$$\begin{aligned} (-\Delta)^s u &= f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \\ u &> 0 \text{ in } \Omega. \end{aligned} \tag{1.6}$$

For further details to the problem (1.6), we refer the readers to [3, 4, 13, 27, 28] and the references therein.

In the literature the study to obtain infinitely many solutions for the problems of the type (1.5), (1.6), the authors have considered either a sublinear or a superlinear growth on f . To our knowledge, the study of the problem (1.1) is very new to the literature due to the presence of a singular term $u^{-\gamma}$. Motivated from [18], we will prove the existence of infinitely many weak solutions to the problem (1.1). We assume the following growth conditions on f .

(A1) $f \in C(\Omega \times \mathbb{R}, \mathbb{R})$ and there exists a $\delta > 0$ such that $\forall x \in \Omega$ and $|t| \leq \delta$, $f(x, -t) = -f(x, t)$.

(A2) $\lim_{t \rightarrow 0} \frac{f(x, t)}{t} = +\infty$ uniformly on Ω .

(A3) There exists $r > 0$ and $\alpha \in (1 - \gamma, 2)$ such that $\forall, x \in \Omega$ and $|t| \leq r$, $tf(x, t) \leq \alpha F(x, t)$, where $F(x, t) = \int_0^t f(x, \tau) d\tau$.

Prior to stating the main theorem, we will define the necessary function spaces and the associated notations. Consider the space $(X, \|\cdot\|_X)$ which is defined as

$$X = \left\{ u : \mathbb{R}^N \rightarrow \mathbb{R} \text{ is measurable, } u|_{\Omega} \in L^2(\Omega) \text{ and } \frac{|u(x) - u(y)|}{|x - y|^{\frac{N+2s}{2}}} \in L^2(Q) \right\}$$

equipped with the Gagliardo norm

$$\|u\|_X = \|u\|_2 + \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy \right)^{\frac{1}{2}}.$$

where $\Omega \subset \mathbb{R}^N$, $Q = \mathbb{R}^{2N} \setminus ((\mathbb{R}^N \setminus \Omega) \times (\mathbb{R}^N \setminus \Omega))$. Here, $\|u\|_2$ refers to the L^2 -norm of u . A frequently used space in this article will be the subspace X_0 of X defined as

$$X_0 = \{u \in X : u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega\}$$

equiped with the norm

$$\|u\| = \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy \right)^{\frac{1}{2}}.$$

The space $(X_0, \|\cdot\|)$ is a Hilbert space [26]. The best Sobolev constant is defined as

$$S = \inf_{u \in X_0 \setminus \{0\}} \frac{\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy}{\left(\int_{\Omega} |2_s^*| dx \right)^{\frac{2}{2_s^*}}} \quad (1.7)$$

We now define a weak solution to the problem (1.1).

Definition 1.1. *A function $u \in X_0$ is a weak solution to the problem (1.1), if $u > 0$, $\phi u^{-\gamma} \in L^1(\Omega)$ and*

$$\int_Q \frac{(u(x) - u(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s}} dx dy - \int_{\Omega} \left(\frac{\lambda}{u^\gamma} + f(x, u) \right) \phi dx = 0, \quad \forall \phi \in X_0. \quad (1.8)$$

The energy functional $I : X_0 \rightarrow (-\infty, \infty]$ associated with the problem (1.1) is defined as

$$I(u) = \frac{1}{2} \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \frac{\lambda}{1 - \gamma} \int_{\Omega} u^{1-\gamma} dx - \int_{\Omega} F(x, u) dx \quad (1.9)$$

We will now state our main result

Theorem 1.2. *Let the assumptions (A1) – (A3) hold, then there exists $\Lambda < \infty$ and for every $\lambda \in (0, \Lambda)$, the problem (1.1) has a sequence of nonnegative weak solutions $\{u_n\} \subset X_0 \cap L^\infty(\Omega)$ such that $I(u_n) < 0$, $I(u_n) \rightarrow 0^-$ and $u_n \rightarrow 0$ in X_0 .*

We will use the symmetric Mountain Pass Theorem version due to Clark [9], to guarantee the existence of distinct infinitely many weak solutions. There exists two versions of the symmetric Mountain Pass Theorem. One of them gives a sequence of critical values diverging to infinity for the superlinear data. The other one provides a sequence of critical values converging to zero for the sublinear data. In the present article, we will use the theorem for the sublinear case. To state the symmetric Mountain Pass Theorem, we will define the notion of genus, which will be used to prove our main Theorem 1.2.

Definition 1.3 (Genus). *Let X be a Banach space and $A \subset X$. A set A is said to be symmetric if $u \in A$ implies $-u \in A$. Let A be a close, symmetric subset of X such that $0 \notin A$. We define a genus $\gamma(A)$ of A by the smallest integer k such that there exists an odd continuous mapping from A to $\mathbb{R}^k \setminus \{0\}$. We define $\gamma(A) = \infty$, if no such k exists.*

Let us consider the following set,

$$\Gamma_n = \{A_n \subset X : A_n \text{ is closed, symmetric and } 0 \notin A_n \text{ such that the genus } \gamma(A_n) \geq n\}.$$

The following version of the symmetric Mountain Pass Theorem has been taken from [20].

Theorem 1.4. *Let X be an infinite dimensional Banach space and $\tilde{I} \in C^1(X, \mathbb{R})$ satisfies the following*

(i) \tilde{I} is even, bounded below, $\tilde{I}(0) = 0$ and \tilde{I} satisfies the $(PS)_c$ condition.

(ii) For each $n \in \mathbb{N}$, there exists an $A_n \in \Gamma_n$ such that $\sup_{u \in A_n} \tilde{I}(u) < 0$.

Then for each $n \in \mathbb{N}$, $c_n = \inf_{A \in \Gamma_n} \sup_{u \in A} \tilde{I}(u) < 0$ is a critical value of \tilde{I} .

In order to apply the symmetric Mountain Pass Theorem, we modify the problem (1.1) to

$$\begin{aligned} (-\Delta)^s u &= \lambda \frac{\text{sign}(u)}{|u|^\gamma} + f(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \end{aligned} \quad (1.10)$$

We define the energy functional $J : X_0 \rightarrow (-\infty, \infty]$ associated with the problem (1.10) as

$$J(u) = \frac{1}{2} \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \frac{\lambda}{1 - \gamma} \int_\Omega |u|^{1-\gamma} dx - \int_\Omega F(x, u) dx \quad (1.11)$$

We now give the definition of a weak solution to the problem (1.10).

Definition 1.5. *A function $u \in X_0$ is a weak solution of (1.10), if $\phi|u|^{-\gamma} \in L^1(\Omega)$ and*

$$\int_Q \frac{(u(x) - u(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s}} dx dy - \int_\Omega \left(\frac{\lambda \text{sign}(u)}{|u|^\gamma} + f(x, u) \right) \phi dx = 0, \quad (1.12)$$

for all $\phi \in X_0$.

It is easy to see that, if u is a weak solution to the problem (1.10) and $u > 0$ a.e., then u is also a weak solution to the problem (1.1). We use a *cutoff* technique given in [9] to guarantee the existence of infinitely many positive weak solutions to the problem (1.10). Let us choose l to be small, such that $0 < l \leq \frac{1}{2} \min\{\delta, r\}$, where δ and r are same as in the assumptions on f . Let us define a C^1 function $\xi : \mathbb{R} \rightarrow \mathbb{R}^+$ such that $0 \leq \xi(t) \leq 1$ and

$$\xi(t) = \begin{cases} 1, & \text{if } |t| \leq l \\ \xi \text{ is decreasing,} & \text{if } l \leq t \leq 2l \\ 0, & \text{if } |t| \geq 2l. \end{cases}$$

Since our main objective is to prove the existence of positive solutions, we will define $f(x, t) = 0$ for $t \leq 0$. Let us consider the following *cutoff* problem.

$$\begin{aligned} (-\Delta)^s u &= \lambda \frac{\text{sign}(u)}{|u|^\gamma} + \tilde{f}(x, u) \text{ in } \Omega, \\ u &= 0 \text{ in } \mathbb{R}^N \setminus \Omega, \end{aligned} \quad (1.13)$$

where,

$$\tilde{f}(x, u) = \begin{cases} f(x, u)\xi(u), & \text{for } u \geq 0 \\ 0, & \text{for } u \leq 0. \end{cases}$$

One can easily see that if u is a weak solution to (1.13) with $\|u\|_\infty \leq l$, then u is also a weak solution to (1.10). We will investigate the existence of infinitely many weak solutions to the problem (1.13). Moreover, to achieve our goal we will prove that $\|u\|_\infty \leq l$ and the solutions to (1.13) are positive.

The energy functional $\tilde{I} : X_0 \rightarrow (-\infty, \infty]$ associated with the problem (1.13) is defined as

$$\tilde{I}(u) = \frac{1}{2} \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \frac{\lambda}{1 - \gamma} \int_\Omega |u|^{1-\gamma} dx - \int_\Omega \tilde{F}(x, u) dx \quad (1.14)$$

A weak solution to the problem (1.13) is given by the following definition.

Definition 1.6. *A function $u \in X_0$ is a weak solution of (1.13), if $\phi|u|^{-\gamma} \in L^1(\Omega)$ and*

$$\int_Q \frac{(u(x) - u(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s}} dx dy - \int_\Omega \left(\frac{\lambda \text{sign}(u)}{|u|^\gamma} + \tilde{f}(x, u) \right) \phi dx = 0, \quad (1.15)$$

for all $\phi \in X_0$.

Henceforth, a weak solution will be referred to as a solution.

2 Existence and Multiplicity of solutions

We begin this section by proving that

$$\Lambda = \inf\{\lambda > 0 : \text{The problem (1.1) has no weak solution}\}.$$

is a finite, nonnegative real number.

Lemma 2.1. *Assume $0 < \gamma < 1$ and (A1) – (A3) holds. Then $0 \leq \Lambda < \infty$.*

Proof. It is clear that $\Lambda \geq 0$ from its definition. Let λ_1 be the principal eigenvalue of the fractional Laplacian operator $(-\Delta)^s$ in Ω and let $\phi_1 > 0$ be the associated eigenfunction [6]. Therefore, we have

$$\begin{aligned} (-\Delta)^s \phi_1 &= \lambda_1 \phi_1 \text{ in } \Omega, \\ \phi_1 &> 0 \text{ in } \Omega, \\ \phi &= 0 \text{ in } \mathbb{R}^N \setminus \Omega. \end{aligned} \quad (2.16)$$

By choosing ϕ_1 as a test function in the Definition 1.1, we get

$$\begin{aligned}\lambda_1 \int_{\Omega} u \phi_1 dx &= \int_{\Omega} (-\Delta)^s \phi_1 u dx \\ &= \int_{\Omega} \left(\frac{\lambda}{u^\gamma} + f(x, u) \right) \phi_1 dx.\end{aligned}\quad (2.17)$$

Let us now choose, any arbitrary constant $\tilde{\Lambda}$ such that $\tilde{\Lambda} t^{-\gamma} + f(x, t) > 2\lambda_1 t \ \forall t > 0$. This contradicts to the equation (2.17). Hence, we get $\Lambda < \infty$. \square

Remark 2.2. *In fact, we will finally prove that $\Lambda > 0$.*

We will now prove the following Lemma, to obtain one of the hypothesis of the symmetric Mountain Pass Theorem.

Lemma 2.3. *The functional \tilde{I} is bounded below and satisfies $(PS)_c$ condition.*

Proof. By using the definition of ξ and Hölder's inequality, we get

$$\begin{aligned}\tilde{I}(u) &\geq \frac{1}{2} \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \lambda C \|u\|^{1-\gamma} - C_1 \\ &\geq \frac{1}{2} \|u\|^2 - \lambda C \|u\|^{1-\gamma} - C_1\end{aligned}$$

where, C, C_1 are non negative constants. This implies that \tilde{I} is coercive and bounded below in X_0 . Let $\{u_n\} \subset X_0$ be a Palais Smale sequence for \tilde{I} . Then $\{u_n\}$ is bounded in X_0 due to the coerciveness of \tilde{I} . Therefore, we may assume, $u_n \rightharpoonup u$ in X_0 upto a subsequence. Thus, we have

$$\int_Q \frac{(u_n(x) - u_n(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s}} dx dy \longrightarrow \int_Q \frac{(u(x) - u(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s}} dx dy \quad (2.18)$$

for all $\phi \in X_0$. By the embedding result [26], we can assume

$$u_n \longrightarrow u \text{ in } L^p(\Omega), \quad (2.19)$$

$$u_n(x) \longrightarrow u(x) \text{ a.e. } L^p(\Omega). \quad (2.20)$$

Therefore, from Lemma A.1 [31], we get that there exists $g \in L^p(\Omega)$ such that

$$|u_n(x)| \leq g(x) \text{ a.e. in } \Omega, \forall n \in \mathbb{N}. \quad (2.21)$$

Hence, by using (2.19), (2.20), (2.21) and the Lebesgue dominated convergence theorem, we get

$$\int_{\Omega} \tilde{f}(x, u_n) u dx \rightarrow \int_{\Omega} \tilde{f}(x, u) u dx \text{ and } \int_{\Omega} \tilde{f}(x, u_n) u_n dx \rightarrow \int_{\Omega} \tilde{f}(x, u) u dx \quad (2.22)$$

Again, on using the Hölder's inequality and passing to the limit $n \rightarrow \infty$, we get

$$\begin{aligned} \int_{\Omega} u_n^{1-\gamma} dx &\leq \int_{\Omega} u^{1-\gamma} dx + \int_{\Omega} |u_n - u|^{1-\gamma} dx \\ &\leq \int_{\Omega} u^{1-\gamma} dx + C \|u_n - u\|_{L^2(\Omega)}^{1-\gamma} \\ &= \int_{\Omega} u^{1-\gamma} dx + o(1) \end{aligned} \quad (2.23)$$

Similarly, we have

$$\begin{aligned} \int_{\Omega} u^{1-\gamma} dx &\leq \int_{\Omega} u_n^{1-\gamma} dx + \int_{\Omega} |u_n - u|^{1-\gamma} dx \\ &\leq \int_{\Omega} u_n^{1-\gamma} dx + C \|u_n - u\|_{L^2(\Omega)}^{1-\gamma} \\ &= \int_{\Omega} u_n^{1-\gamma} dx + o(1) \end{aligned} \quad (2.24)$$

Therefore,

$$\int_{\Omega} u_n^{1-\gamma} dx = \int_{\Omega} u^{1-\gamma} dx + o(1) \quad (2.25)$$

Now, since $\langle \tilde{I}(u_n), u_n \rangle \rightarrow 0$, we have

$$\int_Q \frac{|u_n(x) - u_n(y)|^2}{|x - y|^{N+2s}} dx dy - \lambda \int_{\Omega} |u_n|^{1-\gamma} dx - \int_{\Omega} \tilde{f}(x, u_n) u_n dx \rightarrow 0 \quad (2.26)$$

Therefore, by (2.22), (2.25) and (2.26), we get

$$\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy \rightarrow \lambda \int_{\Omega} |u|^{1-\gamma} dx - \int_{\Omega} \tilde{f}(x, u) u dx \quad (2.27)$$

Moreover,

$$\begin{aligned} \langle \tilde{I}(u_n), u \rangle &= \int_Q \frac{(u_n(x) - u_n(y))(u(x) - u(y))}{|x - y|^{N+2s}} dx dy \\ &\quad - \lambda \int_{\Omega} \text{sign}(u_n) |u_n|^{-\gamma} u dx - \int_{\Omega} \tilde{f}(x, u_n) u dx \end{aligned} \quad (2.28)$$

Note that

$$\langle \tilde{I}(u_n), u \rangle \longrightarrow 0, \text{ as } n \rightarrow \infty.$$

Hence, taking $\phi = u$ in (2.18) and by using (2.25)-(2.27), we get

$$\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy = \lambda \int_{\Omega} |u|^{1-\gamma} dx + \int_{\Omega} \tilde{f}(x, u) u dx \quad (2.29)$$

Therefore, we conclude $\|u_n\| \rightarrow \|u\|$ and this completes the proof. \square

We will now prove a Lemma which will guarantee that, for each $n \in \mathbb{N}$, the set $\Gamma_n \neq \phi$, where ϕ is the empty set.

Lemma 2.4. *For any $n \in \mathbb{N}$, there exists a closed, symmetric subset $A_n \subset X_0$ with $0 \notin A_n$ such that the genus $\gamma(A_n) \geq n$ and $\sup_{u \in A_n} \tilde{I}(u) < 0$.*

Proof. We will first guarantee an existence of a closed, symmetric subset A_n over every finite dimensional subspace of X_0 such that $\gamma(A_n) \geq n$. Let X_k be a subspace of X_0 such that $\dim(X_k) = k$. We know that every norm over a finite dimensional norm linear space are equivalent. Therefore, there exists a positive constant $L = L(k)$ such that $\|u\| \leq L\|u\|_{L^2(\Omega)}$ for all $u \in X_k$.

Claim: There exists a positive constant R such that

$$\frac{1}{2} \int_{\Omega} |u|^2 dx \geq \int_{\{|u|>l\}} |u|^2 dx, \quad \forall u \in X_k \text{ such that } \|u\| \leq R. \quad (2.30)$$

The proof is by contradiction. Let $\{u_n\}$ be a sequence in $X_k \setminus \{0\}$ such that $u_n \rightarrow 0$ in X_0 and

$$\frac{1}{2} \int_{\Omega} |u_n|^2 dx < \int_{\{|u_n|>l\}} |u_n|^2 dx. \quad (2.31)$$

Choose, $v_n = \frac{u_n}{\|u_n\|_{L^2(\Omega)}}$. Then

$$\frac{1}{2} < \int_{\{|u_n|>l\}} |v_n|^2 dx. \quad (2.32)$$

Now, since $\dim(X_k) = k$, we can assume $v_n \rightarrow v$ in X_0 upto a subsequence. Therefore, $v_n \rightarrow v$ also in $L^2(\Omega)$. Further, observe that,

$$m\{x \in \Omega : |u_n| > l\} \rightarrow 0 \text{ as } n \rightarrow \infty$$

since, $u_n \rightarrow 0$ in X_0 . This is a contradiction to (2.32). Therefore, the claim is proved. Now, from the assumption (A2), we can choose $0 < l \leq 1$ such that,

$$\tilde{F}(x, t) = F(x, t) \geq 2L^2 t^2, \quad \forall (x, t) \in \Omega \times [0, l].$$

Hence, for all $u \in X_k \setminus \{0\}$ such that $\|u\| \leq R$ and by using (2.30), we get

$$\begin{aligned} \tilde{I}(u) &\leq \frac{1}{2} \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \frac{\lambda}{1 - \gamma} \int_{\Omega} |u|^{1-\gamma} dx - \int_{\{|u| \leq l\}} \tilde{F}(x, u) dx \\ &\leq \frac{1}{2} \|u\|^2 - \frac{\lambda}{1 - \gamma} \int_{\Omega} |u|^{1-\gamma} dx - 2L^2 \int_{\{|u| \leq l\}} |u|^2 dx \\ &= \frac{1}{2} \|u\|^2 - \frac{\lambda}{1 - \gamma} \int_{\Omega} |u|^{1-\gamma} dx - 2L^2 \left(\int_{\Omega} |u|^2 dx - \int_{\{|u| > l\}} |u|^2 dx \right) \\ &\leq \frac{1}{2} \|u\|^2 - \frac{\lambda}{1 - \gamma} \int_{\Omega} |u|^{1-\gamma} dx - L^2 \int_{\Omega} |u|^2 dx \\ &\leq -\frac{1}{2} \|u\|^2 - \frac{\lambda}{1 - \gamma} \int_{\Omega} |u|^{1-\gamma} dx \\ &< 0. \end{aligned}$$

Let us now choose, $0 < \rho \leq R$ and $A_n = \{u \in X_n : \|u\| = \rho\}$. This serves the purpose of showing that $\Gamma_n \neq \phi$. Since A_n is symmetric, closed with $\gamma(A_n) \geq n$ such that $\sup_{u \in A_n} \tilde{I}(u) < 0$.

This completes the proof. \square

The following Lemmas will be proved to guarantee the boundedness of the solutions to the problem (1.13).

Lemma 2.5. *Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a convex C^1 function. Then for every $a, b, A, B \in \mathbb{R}$ with $A, B > 0$ the following inequality holds.*

$$(g(a) - g(b))(A - B) \leq (a - b)(Ag'(a) - Bg'(b)) \quad (2.33)$$

Proof. Since, g is convex, we have

$$g(b) - g(a) \geq g'(a)(b - a) \text{ and } g(a) - g(b) \geq g'(b)(a - b) \quad (2.34)$$

Therefore, using (2.34), we get

$$\begin{aligned} (a - b)[Ag'(a) - Bg'(b)] &= A(a - b)g'(a) - B(a - b)g'(b) \\ &\geq A[g(a) - g(b)] - B[g(a) - g(b)] \\ &= (A - B)(g(a) - g(b)) \end{aligned}$$

\square

Lemma 2.6. *Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function, then for $a, b, \tau \in \mathbb{R}$ with $\tau \geq 0$ we have*

$$[H(a) - H(b)]^2 \leq (a - b)(h(a) - h(b)) \quad (2.35)$$

where, $H(t) = \int_0^t \sqrt{h'(\tau)} d\tau$, for $t \in \mathbb{R}$.

Proof.

$$\begin{aligned} (a - b)(h(a) - h(b)) &= (a - b) \int_b^a h'(\tau) d\tau \\ &= (a - b) \int_b^a (H'(\tau))^2 d\tau \\ &\geq \left(\int_b^a H'(\tau) d\tau \right)^2 \text{ by Jensen's inequality} \\ &= [H(a) - H(b)]^2 \end{aligned}$$

This completes the proof \square

Lemma 2.7. *Let $u \in X_0$ be a positive weak solution to the problem in (1.13), then $u \in L^\infty(\Omega)$.*

Proof. We follow the steps from Brasco and Parini [6]. For every small $\epsilon > 0$, let us define the smooth function

$$g_\epsilon(t) = (\epsilon^2 + t^2)^{\frac{1}{2}}$$

Observe that the function g_ϵ is convex and Lipschitz. For all positive $\psi \in C_c^\infty(\Omega)$, we take $\phi = \psi g'_\epsilon(u)$ to be the test function in (1.13). By choosing $a = u(x), b = u(y), A = \psi(x)$ and $B = \psi(y)$ in Lemma 2.5, we have

$$\int_Q \frac{(g_\epsilon(u(x)) - g_\epsilon(u(y)))(\psi(x) - \psi(y))}{|x - y|^{N+2s}} dx dy \leq \int_\Omega \left(|\lambda u^{-\gamma} + \tilde{f}(x, u)| \right) |g'_\epsilon(u)| \psi dx \quad (2.36)$$

The function $g_\epsilon(t) \rightarrow |t|$ as $t \rightarrow 0$ and hence $|g'_\epsilon(t)| \leq 1$. Therefore, on using Fatou's Lemma and passing the limit $\epsilon \rightarrow 0$ in (2.36), we obtain

$$\int_Q \frac{(|u(x)| - |u(y)|)(\psi(x) - \psi(y))}{|x - y|^{N+2s}} dx dy \leq \int_\Omega \left(|\lambda u^{-\gamma} + \tilde{f}(x, u)| \right) \psi dx \quad (2.37)$$

for all $\psi \in C_c^\infty(\Omega)$ with $\psi > 0$. The inequality (2.37) remains true for all $\psi \in X_0$ with $\psi \geq 0$. We define the *cutoff* function $u_k = \min\{(u - 1)^+, k\} \in X_0$ for $k > 0$. Now for any given $\beta > 0$ and $\delta > 0$, we choose $\psi = (u_k + \delta)^\beta - \delta^\beta$ as the test function in (2.37) and get

$$\begin{aligned} & \int_Q \frac{(|u(x)| - |u(y)|)((u_k(x) + \delta)^\beta - (u_k(y) + \delta)^\beta)}{|x - y|^{N+2s}} dx dy \\ & \leq \int_\Omega \left(|\lambda u^{-\gamma} + \tilde{f}(x, u)| \right) ((u_k + \delta)^\beta - \delta^\beta) dx \end{aligned} \quad (2.38)$$

Now applying the Lemma 2.6 to the function $h(u) = (u_k + \delta)^\beta$, we get

$$\begin{aligned} & \int_Q \frac{|((u_k(x) + \delta)^{\frac{\beta+1}{2}} - (u_k(y) + \delta)^{\frac{\beta+1}{2}})|^2}{|x - y|^{N+2s}} dx dy \\ & \leq \frac{(\beta + 1)^2}{4\beta} \int_Q \frac{(|u(x)| - |u(y)|)((u_k(x) + \delta)^\beta - (u_k(y) + \delta)^\beta)}{|x - y|^{N+2s}} dx dy \\ & \leq \frac{(\beta + 1)^2}{4\beta} \int_\Omega \left(|\lambda u^{-\gamma} + \tilde{f}(x, u)| \right) ((u_k + \delta)^\beta - \delta^\beta) dx \\ & \leq \frac{(\beta + 1)^2}{4\beta} \int_\Omega \left(|\lambda u^{-\gamma}| + |\tilde{f}(x, u)| \right) ((u_k + \delta)^\beta - \delta^\beta) dx \\ & = \frac{(\beta + 1)^2}{4\beta} \int_{\{u \geq 1\}} \left(|\lambda u^{-\gamma}| + |\tilde{f}(x, u)| \right) ((u_k + \delta)^\beta - \delta^\beta) dx \\ & \leq \frac{(\beta + 1)^2}{4\beta} \int_{\{u \geq 1\}} (|\lambda| + (|c_1| + |c_2||u|^\alpha)) ((u_k + \delta)^\beta - \delta^\beta) dx \end{aligned}$$

$$\begin{aligned}
&\leq C_1 \frac{(\beta+1)^2}{4\beta} \int_{\{u \geq 1\}} (1+|u|^\alpha) ((u_k + \delta)^\beta - \delta^\beta) dx \\
&\leq 2C_1 \frac{(\beta+1)^2}{4\beta} \int_{\{u \geq 1\}} |u|^\alpha ((u_k + \delta)^\beta - \delta^\beta) dx \\
&\leq C \frac{(\beta+1)^2}{4\beta} |u|_{2_s^*}^\alpha |(u_k + \delta)^\beta|_q
\end{aligned} \tag{2.39}$$

where, $q = \frac{2_s^*}{2_s^* - \alpha}$ and $C = \max\{1, |\lambda|\}$. Now by using the Sobolev inequality for X_0 , given by [12], we get

$$\int_Q \frac{|((u_k(x) + \delta)^{\frac{\beta+1}{2}} - (u_k(y) + \delta)^{\frac{\beta+1}{2}})|^2}{|x-y|^{N+2s}} dx dy \geq C_{N,s} |(u_k + \delta)^{\frac{\beta+1}{2}} - \delta^{\frac{\beta+1}{2}}|_{2_s^*}^2 \tag{2.40}$$

where, $C_{N,s}$ is the embedding constant. The triangle inequality and $(u_k + \delta)^{\beta+1} \geq \delta(u_k + \delta)^\beta$ implies

$$\left[\int_\Omega \left((u_k + \delta)^{\frac{\beta+1}{2}} - \delta^{\frac{\beta+1}{2}} \right)^{2_s^*} dx \right]^{\frac{2}{2_s^*}} \geq \frac{\delta}{2} \left(\int_\Omega (u_k + \delta)^{\frac{2_s^* \beta}{2}} dx \right)^{\frac{2}{2_s^*}} - \delta^{\beta+1} |\Omega|^{\frac{2}{2_s^*}} \tag{2.41}$$

Therefore, on using (2.41) in (2.40) and then applying (2.40) to the last inequality of (2.39), we get

$$\begin{aligned}
|(u_k + \delta)^{\frac{\beta}{2}}|_{2_s^*}^2 &\leq C \left(\frac{2(\beta+1)^2}{4\beta\delta C_{N,s}} |u|_{2_s^*}^\alpha |(u_k + \delta)^\beta|_q + \delta^\beta |\Omega|^{\frac{2}{2_s^*}} \right) \\
&\leq C \left(\frac{2(\beta+1)^2}{4\beta\delta C_{N,s}} |u|_{2_s^*}^\alpha |(u_k + \delta)^\beta|_q + \frac{(\beta+1)^2}{4\beta} |\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} |(u_k + \delta)^\beta|_q \right) \\
&\leq C \frac{(\beta+1)^2}{4\beta} |(u_k + \delta)^\beta|_q \left(\frac{|u|_{2_s^*}^\alpha}{\delta C_{N,s}} + |\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)
\end{aligned} \tag{2.42}$$

where, $C = C(N, s) > 0$.

We will now choose, $\delta = \frac{|u|_{2_s^*}^\alpha}{\delta C_{N,s} |\Omega|^{1-\frac{1}{q}-\frac{2s}{N}}} > 0$ and $\beta \geq 1$ such that $\left(\frac{\beta+1}{2\beta}\right)^2 \leq 1$. Let us now choose $\eta = \frac{2_s^*}{2q} > 1$ and $\tau = q\beta$ and then rewrite the inequality (2.42) as

$$|(u_k + \delta)|_{\eta\tau} \leq \left(C |\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)^{\frac{q}{\tau}} \left(\frac{\tau}{q} \right)^{\frac{q}{\tau}} |(u_k + \delta)|_\tau \tag{2.43}$$

We now iterate the inequality (2.43) by setting the following sequence.

$$\tau_0 = q \quad \text{and} \quad \tau_{n+1} = \eta\tau_n = \eta^{n+1}q$$

Hence, after n iteration, the inequality (2.43) reduces to

$$|(u_k + \delta)|_{\tau_{n+1}} \leq \left(C |\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)^{\left(\sum_{i=0}^n \frac{q}{\tau_i} \right)} \prod_{i=0}^n \left(\frac{\tau_i}{q} \right)^{\frac{q}{\tau_i}} |(u_k + \delta)|_q \tag{2.44}$$

Since $\eta > 1$, we have

$$\sum_{i=0}^{\infty} \frac{q}{\tau_i} = \sum_{i=0}^{\infty} \frac{1}{\eta^i} = \frac{\eta}{\eta - 1}$$

and

$$\prod_{i=0}^{\infty} \left(\frac{\tau_i}{q} \right)^{\frac{q}{\tau_i}} = \eta^{\frac{\eta}{(\eta-1)^2}}.$$

Now, by taking limit $n \rightarrow \infty$ in (2.44), we get

$$|u_k|_{\infty} \leq \left(C|\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)^{\frac{\eta}{\eta-1}} \eta^{\frac{\eta}{(\eta-1)^2}} |(u_k + \delta)|_q \quad (2.45)$$

Therefore, by using $u_k \leq (u - 1)^+$ and the triangle inequality in (2.45), we get

$$|u_k|_{\infty} \leq C\eta^{\frac{\eta}{(\eta-1)^2}} \left(|\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)^{\frac{\eta}{\eta-1}} \left(|(u - 1)^+|_q + \delta|\Omega|^{\frac{1}{q}} \right) \quad (2.46)$$

Now letting $k \rightarrow \infty$ in (2.46), we have

$$|(u - 1)^+|_{\infty} \leq C\eta^{\frac{\eta}{(\eta-1)^2}} \left(|\Omega|^{1-\frac{1}{q}-\frac{2s}{N}} \right)^{\frac{\eta}{\eta-1}} \left(|(u - 1)^+|_q + \delta|\Omega|^{\frac{1}{q}} \right) \quad (2.47)$$

Hence, we conclude that $u \in L^{\infty}(\Omega)$. \square

Proof of Theorem 1.2. From the definition of ξ and the assumption (A1), we have \tilde{I} is even and $\tilde{I}(0) = 0$. Therefore, Lemma 1.4, Lemma 2.3 and Lemma 2.4 guarantees that \tilde{I} has sequence of critical points $\{u_n\}$ such that $\tilde{I}(u_n) < 0$ and $\tilde{I}(u_n) \rightarrow 0^-$.

Claim: Suppose u_n is a critical point of \tilde{I} , then for each $n \in \mathbb{N}$, $u_n \geq 0$ a.e. in X_0 .

Proof. Let us consider, $\Omega = \Omega^+ \cup \Omega^-$, where $\Omega^+ = \{x \in X_0 : u_n(x) \geq 0\}$ and $\Omega^- = \{x \in X_0 : u_n(x) < 0\}$. We define $u_n(x) = u_n^+ - u_n^-$, where $u_n^+(x) = \max\{u_n(x), 0\}$ and $u_n^-(x) = \max\{-u_n(x), 0\}$. Suppose, $u_n < 0$ a.e. in Ω , then on taking, $\phi = u_n^-$ as the test function in the equation (1.6) in conjunction with the inequality $(a-b)(a^- - b^-) \leq -(a^- - b^-)^2$, we get

$$\begin{aligned} \int_{\Omega} \left(\lambda \frac{\text{sign}(u_n)u_n^-}{|u_n|^{\gamma}} + \tilde{f}(x, u_n)u_n^- \right) dx &= \int_Q \frac{(u_n(x) - u_n(y))(u_n^-(x) - u_n^-(y))}{|x - y|^{N+2s}} dx dy \\ \Rightarrow \lambda \int_{\Omega^-} |u_n^-|^{1-\gamma} dx &\leq -\|u_n^-\|^2 < 0. \end{aligned}$$

This implies $|\Omega^-| = 0$, which is a contradiction to the assumption $u_n < 0$ a.e. in Ω . This proves our claim. \square

Moreover, from the definition of \tilde{I} , we have

$$\begin{aligned}
\frac{1}{\alpha} \langle \tilde{I}'(u_n), u_n \rangle - \tilde{I}(u_n) &= \frac{1}{\alpha} \left[\|u_n\|^2 - \int_{\Omega} \left(\lambda \frac{\text{sign}(u_n)u_n}{|u_n|^\gamma} + \tilde{f}(x, u_n)u_n \right) dx \right] \\
&\quad - \left[\frac{1}{2} \|u_n\|^2 - \int_{\Omega} \left(\frac{\lambda}{1-\gamma} |u_n|^{1-\gamma} + \tilde{F}(x, u_n) \right) dx \right] \\
&= \left(\frac{1}{\alpha} - \frac{1}{2} \right) \|u_n\|^2 - \lambda \left(\frac{1}{\alpha} - \frac{1}{1-\gamma} \right) \int_{\Omega} |u_n|^{1-\gamma} dx \\
&\quad + \frac{1}{\alpha} \int_{\Omega} (\alpha \tilde{F}(x, u_n) - \tilde{f}(x, u_n)) dx \\
&\geq \left(\frac{1}{\alpha} - \frac{1}{2} \right) \|u_n\|^2 + \lambda \left(\frac{1}{1-\gamma} - \frac{1}{\alpha} \right) \int_{\Omega} |u_n|^{1-\gamma} dx \\
&\geq \left(\frac{1}{\alpha} - \frac{1}{2} \right) \|u_n\|^2
\end{aligned}$$

It can be easily seen that, since

$$\begin{aligned}
\frac{1}{\alpha} \langle \tilde{I}'(u_n), u_n \rangle - \tilde{I}(u_n) &= o_n(1) \\
\Rightarrow \left(\frac{1}{\alpha} - \frac{1}{2} \right) \|u_n\|^2 &\leq o_n(1),
\end{aligned}$$

hence, we get $u_n \rightarrow 0$ in X_0 . Now by using Moser iteration and Lemma 2.7, we can assume that as $n \rightarrow \infty$, $\|u_n\|_{L^\infty(\Omega)} \leq l$. Therefore, the problem (1.10) has infinitely many solutions. Further, due to the nonnegativity of u_n and $\tilde{I}(u_n) < 0$, we conclude that the problem (1.1) has infinitely many weak solutions and hence the Theorem 1.2 is proved. \square

Remark 2.8. $\Lambda > 0$, because the solution to the problem (1.1) exists for some $\lambda > 0$.

Acknowledgement

The author S. Ghosh, thanks the Council of Scientific and Industrial Research (C.S.I.R), India, for the financial assistantship received to carry out this research work. Both the authors thanks the research facilities received from the Department of Mathematics, National Institute of Technology Rourkela, India. The authors thank the anonymous reviewers for their constructive comments and suggestions.

References

- [1] Ambrosetti A. and Rabinowitz P. H., Dual variational methods in critical point theory and applications. *Journal of functional Analysis*, 14(4), 349-381, 1973.
- [2] Bertoin J., Lévy processes, volume 121 of cambridge tracts in mathematics, *Cambridge Tracts in Mathematics, vol. 121, Cambridge University Press*, 1998.

- [3] Binlin Z., Bisci G. M. and Servadei R., Superlinear nonlocal fractional problems with infinitely many solutions. *Nonlinearity*, 28(7), 2247-2264, 2015.
- [4] Bisci G. M., Repovš D. and Servadei R., Nontrivial solutions of superlinear nonlocal problems. In *Forum Mathematicum*, 28, 1095-1110, 2016.
- [5] Boccardo L. and Orsina L., Semilinear elliptic equations with singular nonlinearities. *Calculus of Variations and Partial Differential Equations*, 37(3), 363-380, 2010.
- [6] Brasco L. and Parini E., The second eigenvalue of the fractional p -laplacian. *Advances in Calculus of Variations*, 9(4), 323-355, 2016.
- [7] Brezis H. and Lieb E., A relation between pointwise convergence of functions and convergence of functionals. *Proceedings of the American Mathematical Society*, 88(3), 486-490, 1983.
- [8] Canino A., Montoro L., Sciunzi B. and Squassina, M., Nonlocal problems with singular nonlinearity. *Bulletin des Sciences Mathématiques*, 141(3), 223-250, 2017.
- [9] Clark D. and Gilbarg D., A variant of the lusternik-schnirelman theory. *Indiana University Mathematics Journal*, 22(1), 65-74, 1972.
- [10] Cont R., Financial modelling with jump processes *Chapman and Hall/CRC Financial Mathematics Series*. Chapman and Hall/CRC, 2003.
- [11] Crandall M. G., Rabinowitz P. H. and Tartar L., On a dirichlet problem with a singular nonlinearity. *Communications in Partial Differential Equations*, 2(2) 193-222, 1977.
- [12] Di Nezza E., Palatucci G. and Valdinoci E., Hitchhiker's guide to the fractional sobolev spaces. *Bulletin des Sciences Mathématiques*, 136(5), 521-573, 2012.
- [13] Dipierro S., Medina M. and Valdinoci E., *Fractional elliptic problems with critical growth in the whole of \mathbb{R}^n* . Lecture Notes Series (15), Springer, 2017.
- [14] Fang Y., Existence, uniqueness of positive solution to a fractional laplacians with singular nonlinearity. *arXiv preprint arXiv:1403.3149*, 2014.
- [15] Ghanmi A. and Saoudi K., The nehari manifold for a singular elliptic equation involving the fractional laplace operator. *Fractional Differential Calculus*, 6(2), 201-217, 2016.
- [16] Saoudi K., Ghosh S., and Choudhuri D., Multiplicity and Hölder regularity of solutions for a nonlocal elliptic PDE involving singularity. *arXiv preprint arXiv:1808.02469*, 2018.
- [17] Ghosh S., Choudhuri D. and Giri R. K., Singular nonlocal problem involving measure data. *Bulletin of the Brazilian Mathematical Society, New Series*, 2018. (In press).
- [18] Gu G., Zhang W. and Zhao F., Infinitely many positive solutions for a nonlocal problem. *Applied Mathematics Letters*, 84, 49-55, 2018.

- [19] Heinz H. P., Free ljusternik-schnirelman theory and the bifurcation diagrams of certain singular nonlinear problems. *Journal of differential equations*, 66(2), 263-300, 1987.
- [20] Kajikiya R., A critical point theorem related to the symmetric mountain pass lemma and its applications to elliptic equations. *Journal of Functional Analysis*, 225(2), 352-370, 2005.
- [21] Lazer A. C. and McKenna P. J., On a singular nonlinear elliptic boundary-value problem. *Proceedings of the American Mathematical Society*, 111(3), 721-730, 1991.
- [22] Liu Z. and Wang Z., On clark's theorem and its applications to partially sublinear problems. In *Annales de l'Institut Henri Poincare (C) Non Linear Analysis*, 32, 1015-1037, 2015.
- [23] Mukherjee T. and Sreenadh K., On dirichlet problem for fractional p -laplacian with singular non-linearity. *Advances in Nonlinear Analysis*, 2016.
- [24] Rosen G., Minimum value for c in the sobolev inequality $|\varphi^3| \leq c|\nabla\varphi|^3$. *SIAM Journal on Applied Mathematics*, 21(1), 30-32, 1971.
- [25] Saoudi K. A critical fractional elliptic equation with singular nonlinearities. *Fractional Calculus and Applied Analysis*, 20(6), 1507-1530, 2017.
- [26] Servadei R. and Valdinoci E., Mountain pass solutions for non-local elliptic operators. *Journal of Mathematical Analysis and Applications*, 389(2), 887-898, 2012.
- [27] Servadei R. and Valdinoci E., Variational methods for non-local operators of elliptic type. *Discrete Contin. Dyn. Syst*, 33(5), 2105-2137, 2013.
- [28] Servadei R. and Valdinoci E., The brezis-nirenberg result for the fractional laplacian. *Transactions of the American Mathematical Society*, 367(1), 67-102, 2015.
- [29] Valdinoci E. From the long jump random walk to the fractional laplacian. *Bol. Soc. Esp. Mat. Apl. SeMA*, 49, 33-44, 2009.
- [30] Wang Z., Nonlinear boundary value problems with concave nonlinearities near the origin. *Nonlinear Differential Equations and Applications NoDEA*, 8(1), 15-33, 2001.
- [31] Willem M., *Minimax theorems*, 24, Springer Science & Business Media, 1997.