

3D TQFT AND HOMFLYPT HOMOLOGY

A. OBLOMKOV AND L. ROZANSKY

ABSTRACT. We describe a family of 3d topological B-models whose target spaces are Hilbert schemes of points in \mathbb{C}^2 . The interfaces separating theories with different numbers of points correspond to braid strands. The Hilbert space of the picture of a closed braid is the HOMFLY-PT homology of the corresponding link.

1. INTRODUCTION

In our previous papers we proposed a construction of the HOMFLY-PT link homology which is based on an interaction between two quiver varieties: the cotangent bundle to the flag variety and the Hilbert scheme of points in \mathbb{C}^2 . For $r, n \geq 0$ denote $\mathcal{M}_{n,r}$ the moduli space of n $U(r)$ instantons in \mathbb{C}^2 . Equivalently, $\mathcal{M}_{n,r}$ is the Nakajima quiver variety of the framed \tilde{A}_0 quiver:



(1.1)

Following [KRS09] and [KR10] we associate to $\mathcal{M}_{n,r}$ the 2-category $\mathring{\text{Cat}}(\mathcal{M}_{n,r})$. We consider a special flag-related object \mathcal{F}_n in this category. Using our previous results [OR18e] we construct a monoidal functor

$$\Phi : \mathfrak{B}\mathfrak{t}_n \longrightarrow \text{Hom}(\mathcal{F}_n, \mathcal{F}_n), \quad \beta \mapsto \Phi(\beta),$$

where $\mathfrak{B}\mathfrak{t}_n$ is a braid group on n strands.

Our main knot theoretic result from [OR18e],[OR20] can be recast into

Theorem 1.0.1. *If β is an ordinary braid, then in case of $r = 1$ there is an isomorphism*

$$(1.2) \quad \mathcal{H}om_{\text{Hom}(\mathcal{F}_n, \mathcal{F}_n)}(\Phi(1_n), \Phi(\beta) \otimes \Lambda^\bullet V_n) \cong \text{HHH}(L(\beta)).$$

Here 1_n is the identity braid, $V_n = \mathbb{C}^n$ is the defining representation of $\text{GL}(n)$, $L(\beta)$ is the link constructed by closing β and $\text{HHH}(L(\beta))$ is its HOMFLY-PT homology.

In this paper we consider this construction of link homology from a 3-category perspective. We outline a construction of a 3-category $\mathring{\mathring{\text{Cat}}}(Q)$ for any \tilde{A}_n -type quiver Q . In the case $Q = \tilde{A}_0$ we provide a conjectural description of a part of 3-categorical structure by relating our setting to the theory of monoidal 2-categories of Kapranov and Voevodsky

[KV94]. The 2-category $\mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;r})$ emerges as the 2-category of morphisms between the objects $\tilde{A}_0(n;r)$ and $\tilde{A}_0(0;0)$, that is $\mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;r}) = \mathbb{H}\text{om}(\tilde{A}_0(n;r), \tilde{A}_0(0;0))$.

The objects of the 3-category $\mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\tilde{A}_0)$ are $\tilde{A}_0(n;r)$ or, equivalently, pairs of numbers $(n;r)$. For two pairs $(n;r)$ and $(n+k;r)$ we define two ‘box’ functors:

$$(1.3) \quad \hat{\square}_k, \hat{\blacksquare}_k : \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;r}) \longrightarrow \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n+k;r}).$$

The functors represent generalized Lagrangian correspondences between $\mathcal{M}_{n;r}$ and $\mathcal{M}_{n+k;r}$, and those originate from the edges of the second Nakajima quiver which is ‘transversal’ to the \tilde{A}_0 quiver(1.1) in the sense of Gaiotto-Witten defects [GW09] in Kapustin-Witten theory [KW07], or brane arrangements in the superstring theory, see section 2.

The quiver variety $\mathcal{M}_{0;r}$ is just a point, hence its 2-category $\mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{0;r})$ has a canonical object \mathcal{O}_\emptyset , and we show in proposition 5.2.1 that \mathcal{F}_n is the result of applying $\hat{\square}_1$ n times to this object:

$$\mathcal{F}_n = \underbrace{\hat{\square}_1 \cdots \hat{\square}_1}_n \mathcal{O}_\emptyset \in \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;r}).$$

Each individual $\hat{\square}_1$ corresponds to a braid strand, and an elementary braid σ_i which braids the i -th and the $(i+1)$ -st string, becomes a natural self-transformation of the composition:

$$\sigma_i \rightsquigarrow \hat{\square}_1 \hat{\square}_1 : \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{i-1;r}) \longrightarrow \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{i+1;r}).$$

When $r = 0$ we define two more ‘box’ functors

$$(1.4) \quad \check{\square}_k, \check{\blacksquare}_k : \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;0}) \longrightarrow \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n+k;0}).$$

These functors are related to Cherkis bow edges of the transversal ‘arrow-bow’ quiver. For an n -box, m -row Young diagram λ with rows $\lambda_1 \geq \cdots \geq \lambda_m \geq 0$ we consider an object

$$\mathcal{S}_\lambda = \check{\square}_{\lambda_1} \cdots \check{\square}_{\lambda_m} \mathcal{O}_\emptyset \in \mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(\mathcal{M}_{n;r})$$

related to the equivariant Slodowy slice of a nilpotent matrix corresponding to λ . The category of morphisms between \mathcal{F}_n and \mathcal{S}_λ is the category of sheaves on (the S-dual of) the Nakajima quiver variety related to the weight space λ in the tensor product of n fundamental representations of $\text{GL}(m)$.

Most recently, the related categories were studied by Nakajima, Takayama [NT17] in the context of 3D mirror symmetry (see section 6.2 for more discussion). The case of $\mathcal{S}_{\frac{n}{2}, \frac{n}{2}}$ $m = 2$ is related to the equivariant resolution of the Slodowy slice for the two-block nilpotent orbit in \mathfrak{gl}_{2n} . In [AN16] it was conjectured that the corresponding category can be used to construct the annular $\text{GL}(2)$ link homology [APS04]. We explain a precise statement of the conjecture and its TQFT interpretation in section 6.2.

The box functors have a natural interpretation as defects in 3d B-models considered by Kapustin-Rozansky-Saulina [KRS09],[KR10]. This allows us to expand our results in three directions.

First, we provide details of the mathematical construction of some the structural morphisms of three-category $\mathring{\mathcal{C}}\mathring{\mathcal{A}}\mathring{\mathcal{T}}(Q)$ for a linear quiver.

Second, we use basic TQFT axioms to give a physical interpretation of the Chern functor from our previous work [OR18d] and thus provide an explanation for the appearance of the Hilbert scheme of points on \mathbb{C}^2 in knot homology [OR20].

Third, we unify several homology theories of the closures of an affine braid $\beta \in \mathfrak{Br}_n^{\text{aff}}$ by interpreting them as the TQFT partitions functions on the annulus with external boundary consisting of n NS5 defects braided into the braid β . More precisely, by setting an appropriate configuration of the defects on the internal boundary of the annulus we obtain: Hochschild homology of the Rouquier complex [KR08], the Hilbert scheme version of the triply braided homology [OR20] and the trace of the braid action on \mathfrak{sl}_2 conformal blocks.

While exploring the categorical foundations of Kapustin-Rozansky-Saulina (KRS) theory [KRS09], we avoid working with 3-category of KRS directly. Instead, we work with the de-looped version $\mathring{\text{Cat}}(Q)$ of the three-category. The 3-categorical nature of the 2-category $\mathring{\text{Cat}}(Q)$ is reflected in the 2-categorical monoidal structure of $\mathring{\text{Cat}}(Q)$. The categorical details can be found in section 4, in the rest of the introduction we discuss the last two directions that are concerned the most topologically motivated case TQFT that is based on the three category $\mathring{\text{Cat}}(\hat{A}_0)$.

1.1. Partition-function evaluation for HOMFLY-PT homology. The initial construction in [OR18e] is based on the homomorphism from the braid group to a special category of matrix factorizations which appears as the category of endomorphism of the object $\mathcal{F}_n \in \mathring{\text{Cat}}(\mathcal{M}_{n,1})$ $\text{MF}^{\text{st}} := \text{Hom}(\mathcal{F}_n, \mathcal{F}_n)$:

$$(1.5) \quad \mathfrak{Br}_n \rightarrow \text{MF}^{\text{st}} = \text{MF}_{\text{GL}_n}((\mathfrak{gl}_n \times \mathbb{C}^n \times \text{T}^*\text{Fl} \times \text{T}^*\text{Fl})^{\text{st}}, W)^1, \quad W = \mu_1 - \mu_2,$$

where Fl is the flag variety, μ is the moment map of the GL_n action on T^*Fl and the index ‘st’ stands for the stability condition.

The object \mathcal{F}_n appears as a boundary condition for the KRS theory with the target

$$\mathcal{M}_{n,r} = (\text{T}^*\mathfrak{gl}_n \times \mathbb{C}^{n \times r})^{\text{st}} // \text{GL}_n.$$


The cases $r = 0$ or $r = 1$ are the most relevant for the link homology. We refer to the first case as *unframed* and to the second case as *framed*. We choose the space-time of all our 3d TQFTs to be $S^2 \times \mathbb{R}$.

The total space-time $S^2 \times \mathbb{R}$ can have ‘defect surfaces’ separating different models, that is, the connected components of the complement of defect surfaces are labeled by integers n . The defects can intersect along curves and curves can intersect at points. In the framed category, the defects have orientation and in both settings the curves of the intersection of the defects have signs.

The simplest case of the defect picture is when the defect is equal to $\text{Def} = C \times \mathbb{R} \subset S^2 \times \mathbb{R}$ where C is a submersed curve (possibly with many connected components). In this case we define the partition-function evaluation Z^\bullet where \bullet is \emptyset in the unframed case and $\bullet = \text{f}$ in the framed case. Our choices in the construction of the partition-function evaluation are driven by the topological applications [OR18e] of the homomorphism (1.5):

¹In the original paper we worked with the affine version of this category of matrix factorizations.

Theorem 1.1.1. *Let $S^1 \times 0 \subset S^2 \times \mathbb{R}$ be an embedded circle such that it intersect the defect $\text{Def} = C \times \mathbb{R}$ transversally at $2n$ points and the labels of the connected components of $S^1 \setminus S^1 \cap \text{Def}$ are $0, 1, 2, \dots, n, n-1, \dots, 2, 1$, as for example, in the picture below:*



then

$$Z(S^1) = \text{MF}_{\text{GL}_n}(\mathfrak{gl}_n \times T^*\text{Fl} \times T^*\text{Fl}, W).$$

Moreover, if the first n intersection points of $S^1 \cap \text{Def}$ are oriented up and the rest down, as in the above picture, then

$$(1.6) \quad Z^f(S^1) = \text{MF}_{\text{GL}_n}((\mathfrak{gl}_n \times \mathbb{C}^n \times T^*\text{Fl} \times T^*\text{Fl})^{\text{st}}, W).$$

In [OR18d] we constructed the pair of adjoint functors:

$$(1.7) \quad \begin{array}{ccc} & \text{CH}_{\text{loc}}^{\text{st}} & \\ & \curvearrowright & \\ \text{MF}^{\text{st}} & & \text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2)) \\ & \curvearrowleft & \\ & \text{HC}_{\text{loc}}^{\text{st}} & \end{array},$$

where $\text{Hilb}_n(\mathbb{C}^2)$ is the Hilbert scheme of n points on \mathbb{C}^2 , while $\text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2))$ is the derived category of two-periodic $\mathbb{T}_{q,t}$ -equivariant complexes on the Hilbert scheme. We also showed that

$$(1.8) \quad \text{HHH}(\beta) = \mathcal{H}om(\text{CH}_{\text{loc}}^{\text{st}}(\beta), \Lambda^\bullet V_n).$$

The TQFT picture gives a natural interpretation of the isomorphism (1.8) as the result of gluing the same disc in two different ways. The appearance of $\text{Hilb}_n(\mathbb{C}^2)$ is due to the following:

Theorem 1.1.2. *Let $S^1 \times 0 \subset S^2 \times \mathbb{R}$ be an embedded circle which does not intersect the defect $\text{Def} = C \times \mathbb{R}$ and lies in the connected component of the complement of the defect with the label n (see Figure 4 for $n = 2$ example) then*

$$Z^f(S^1) = \text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2)).$$

Moreover, if D is a disc in the complement of the defect, such that $\partial D = S^1$, then


$$Z^f(D) = \mathcal{O} \in \text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2)).$$

To explain the appearance of the exterior powers of \mathcal{B} we introduce a special line of defect in our theory: $0 \times \mathbb{R} \subset S^2 \times \mathbb{R}$, $S^2 = \mathbb{R}^2 \cup \infty$. We assume that this line of defect does not intersect the surface of defect. For a small disc intersecting this line of defect we have:

$$(1.9) \quad Z^f(D) = \Lambda^\bullet V_n \in \text{Hilb}_n(\mathbb{C}^2).$$

The link homology emerges as the vector space associated to a disc which intersects defect surfaces and a defect line:

Theorem 1.1.3. *Suppose that the curve $C \subset S^2 = \mathbb{R}^2 \cup \infty$ in the defect $\text{Def} = C \times \mathbb{R}$ is the picture of the natural closure of the braid $\beta \in \mathfrak{B}\mathfrak{r}_n$, the closure goes around the line of defect. Let us also assume the infinite point has the label 0, the connected component of $S^2 \times \mathbb{R} \setminus \text{Def}$ has label n and the labels change by 1 as we cross the surface of defect as on the picture below:*



where the green dot represents a cross-section of the defect line. Then

$$(1.10) \quad \mathbf{Z}^f(S^2) = \text{HHH}(\beta).$$

Thus the formulas (1.2) and (1.8) correspond to two ways to present S^2 as a gluing of two D^2 along their common boundary S^1 . The formula (1.2) is given by cutting along S^1 with S^1 as in theorem 1.1.1 and the formula (1.8) is given by cutting along S^1 that is boundary of a tubular neighborhood of the line of defect.


Note that theorem 1.1.2 interprets $\text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2))$ as the category of endomorphisms of the identity functor in the two category $\mathbf{Z}^f(\mathfrak{p}_n)$:

$$\text{Hom}(\mathbb{I}d, \mathbb{I}d) = \text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2))$$

where the point \mathfrak{p}_n lies in complement of the defect with the label n and $\mathbb{I}d$ is the identity endomorphism of two-category $\mathbf{Z}^f(\mathfrak{p}_n)$. Thus it is reasonable to call $\text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2))$ the Drinfeld center of two-category $\mathbf{Z}^f(\mathfrak{p}_n)$. Since it is not common to work with the Drinfeld centers of two-categories, we spell out the expected property of such a center in corollary 5.1.1.

1.2. Unified perspective on traces. The equation (1.10) can be interpreted as a categorification of the Jones-Oceanu trace [Jon87]. A pictorial presentation of the configuration

of the defects that participate in the formula (1.10) is the annulus with two boundary circles:



(1.11)

here $\alpha \in \mathfrak{B}\mathfrak{r}_n$, the numeric labels indicate the type TQFT in the connected component. In the picture we have n NS5 defects following the outer boundary of the annulus and braid themselves into the braid α . The red domain contains the inner circle boundary defect configuration \mathbb{B} . In the case treated by the equation (1.10) the boundary configuration is the domain with the line defect that carries the vector bundle $\Lambda^\bullet V_n$. The line defect is reflected by the equation (1.9).

It turns out that by varying the configuration of defects \mathbb{B} in the center we can categorify several other interesting trace evaluations on the braid group as well as algebra of the braid graphs $\mathfrak{B}\mathfrak{r}_n^b$. Let us fix notations for the trace we plan to discuss. Soergel explained [Soe01] a realization of the braid graphs $\gamma \in \mathfrak{B}\mathfrak{r}_n^b$ inside the additive category of Soergel bimodules Sbim_n . Thus the Hochschild homology functor HH_* yields a trace on $\mathfrak{B}\mathfrak{r}_n^b$:

$$\Phi^S : \mathfrak{B}\mathfrak{r}_n \rightarrow \text{Sbim}_n, \quad \text{HH}_{\text{alg}}(\gamma) = \text{HH}_*(\Phi^S(\gamma)) = \mathcal{H}om(\Phi^S(\beta), \Phi^S(I)).$$

The last example that we discuss is related to the space of conformal blocks. Let us denote by V_λ and L_λ the Verma module and its irreducible quotient for quantum group $U_q(\mathfrak{sl}_2)$ and $\lambda \in \mathbb{C} = \mathfrak{h}^*$. In particular, $L_1 = \mathbb{C}^2$ is the vector representation of $U_q(\mathfrak{sl}_2)$ and L_λ is irreducible for not integer λ and $L_\lambda = V_\lambda$. The space of conformal on the cylinder is defined as

$$H_{1^n, \mu}^\lambda = \text{Hom}_{U_q(\mathfrak{sl}_2)}(L_\lambda, L_1 \otimes \cdots \otimes L_1 \otimes L_\mu),$$

where we assume that λ, μ are not integers.

The R -matrix yields the braid group $\mathfrak{B}\mathfrak{r}_n$ action on the last space. Let's denote the trace of the action by $\text{Tr}(\beta)[H_{1^n, \mu}^\lambda]$. The space $H_{1^n, \mu}^\lambda$ is trivial unless $|\lambda - \mu| \leq n$. On the other hand if $\lambda - \mu = k$, $|k| \leq n$ then the above mentioned trace does not depend on the value of λ as long as λ is not integer.

The vector space $H_{1^n, \mu}^\lambda$ has two gradings that originate from the weight grading on L_μ and the decomposition of $L_1^{\otimes n}$ on the irreducible components. The braid action respects these gradings. We propose a construction of doubly graded vector space $\mathcal{T}r(\beta)[H_{1^n, \mu}^\lambda]$ that categorifies the above trace.

The three constructions are partition-function evaluations of defect configuration as in the picture and the internal disc defect \mathbb{B} as in the table

²The elements of $\mathfrak{B}\mathfrak{r}_n$ are related to the braid graphs by the MOY relations [MOY98].

\mathbb{B}	$\Lambda \bullet V_n$	$\text{NS5}^{(n)}$	$\text{D5}^{(k)} + \text{D5}^{(n-k)}$
α	$\mathfrak{B}\mathfrak{r}_n$	$\mathfrak{B}\mathfrak{r}_n^b$	$\mathfrak{B}\mathfrak{r}_n$
r	1	1	0
Z	$\text{HHH}(\alpha)$	$\text{HHH}_{\text{alg}}(\alpha)$	$\mathcal{T}r(\alpha)[H_{1^n, \lambda+k}^\lambda]$

where $\text{NS5}^{(k)}$, $\text{D5}^{(k)}$ are NS5 and D5 defects of charge k discussed in the section 3.6. The first two columns of the table reflect mathematically proven statements and the last column seems to be new mathematical construction. We explain the mathematical details of the relations in the table in section 6.

In section 3 we recall some basics of the constructions in [KR10]. In the section 4 we explain how our particular example of TQFT fits into the setting of [KR10]. In section 5 we construct the partition-function Z^f and prove the results that we mentioned in the introduction. We also discuss the Drinfeld center subtleties in the subsection 5.3.

Acknowledgments We would like to thank Dmitry Arinkin, Tudor Dimofte, Eugene Gorsky, Sergey Gukov, Tina Kanstrup, Ivan Losev, Roman Bezrukavnikov and Andrei Neguț for useful discussions. The authors also extremely grateful to an anonymous referee for many corrections and important suggestion on the structure of the paper. The work of A.O. was supported in part by the NSF CAREER grant DMS-1352398, NSF FRG grant DMS-1760373 and Simons Fellowship. The work of L.R. was supported in part by the NSF grant DMS-1760578.

2. PHYSICS BACKGROUND

2.1. TQFT perspective.

2.1.1. *Kapustin-Witten theory.* We study the $U(n)$ Kapustin-Witten (KW) theory [KW07], that is, a 4-dimensional $N = 4$ supersymmetric $U(n)$ Yang-Mills (SYM) theory with a GL-twist. We select $t = i$ which means that the theory represents the Galois side of the geometric Langland duality. We consider the KW theory on the 4-manifold

$$(2.1) \quad M^4 = \mathbb{R}_0 \times \Sigma \times S_3^1,$$

where Σ is a 2-dimensional surface, $\mathbb{R}_{i_1 \dots i_k}^k$ denotes \mathbb{R}^k with coordinates x_{i_1}, \dots, x_{i_k} and S_i^1 denotes the circle S^1 with the periodic coordinate x_i . In fact, we will be mostly interested in the case of $\Sigma = \mathbb{R}_{12}^2$, which is a local form of the general Σ .

We consider the KW theory in the presence of NS5 and D5-interfaces introduced by Gaiotto and Witten [GW09]. The interfaces split M^4 into separate domains, and each domain has its own value of n .

We place two interface arrangements simultaneously in M^4 . The first (basic) arrangement consists of parallel NS5 and D5-interfaces, each interface sweeping a 3-dimensional subspace $\mathbb{R}_0 \times \Sigma \times \{s_I\}$, where $s_I \in S_3^1$ is a point on the circle corresponding to $x_3 = s_I$. We assume that the values of n are the same on both sides of each D5-interface, but they can vary across NS5-interfaces.

An interface I of the second (additional) arrangement has the form $\mathbb{R}_0^1 \times \alpha_I \times S_3^1$, where $\alpha_I \subset \Sigma$ is a curve. The curves of different additional interfaces may intersect.

2.1.2. *3-dimensional B-models.* Fix a point $b \in \mathbb{R}_0$. Our goal is to describe the category corresponding to a 2-dimensional surface $\mathcal{C} \subset M^4$ which has the form $\mathcal{C} = \{b\} \times \beta \times S_3^1$, where $\beta \subset \Sigma$ is a curve possibly intersecting the curves α_I of additional interfaces, and to determine the (Hilbert) vector space of a 3-dimensional section $\{b\} \times \Sigma \times S_3^1$. In the full rigorous approach this would require the description of 2-dimensional defects appearing at the intersections of interfaces, in particular, the intersections of each additional interface with all basic interfaces.

In this paper we pursue a heuristic but easier approach. We compactify M^4 on S_3^1 , thus reducing it to $M^3 = \mathbb{R}_0 \times \Sigma$. M^3 is split by additional interfaces into domains. Each 3-dimensional domain is the result of squeezing the 4-dimensional domains separated by basic interfaces. Hence in each 3-dimensional domain lives a 3-dimensional topological ‘Coulomb-twisted’ $N = 4$ SYM theory with matter fields. Its Higgs branch yields a B-model whose target is the affine (\tilde{A}_{m-1}) Nakajima quiver variety, where m is the number of NS5-interfaces in the basic arrangement.

The description of boundary conditions (including interfaces by folding trick) of 3-dimensional B-models has been worked out in [KRS09] and [KR10], so our goal is to conjecture the form of NS5 and D4 (formerly D5, the new name being due to string theory conventions) interfaces between 3-domains populated by affine Nakajima quivers. We will present the conjecture for the general NS5 interface and we will also describe the D4 interfaces in a particular case when the basic arrangement consists of a single interface of NS5 type.

2.2. String Perspective. A standard IIB string theory construction with two Hanany-Witten type brane arrangements yields the Kapustin-Witten model with interfaces, and we just recall it here briefly.

Consider the IIB string theory on $M_{\text{IIB}}^{10} = \mathbb{R}_{0123}^4 \times \mathbb{R}_{4567}^4 \times \mathbb{R}_{89}^2$, where $\mathbb{R}_{i_1 \dots i_k}^k$ denotes \mathbb{R}^k with coordinates x_{i_1}, \dots, x_{i_k} . The twisting identifies the first two factors, so that the $\text{SO}(4)$ rotation of \mathbb{R}_{0123}^4 is accompanied by the matching rotation of \mathbb{R}_{4567}^4 . D3-branes are stretched along \mathbb{R}_{0123}^4 . Consider an orthogonal splitting $\mathbb{R}_{0123}^4 = \mathbb{R}_{\text{br}}^3 \oplus \mathbb{R}^1$, where \mathbb{R}_{br}^3 subspace is tangent to an NS5 or a D5 brane, and the corresponding splitting $\mathbb{R}_{4567}^4 = \mathbb{R}_{\text{br}}^3 \oplus \mathbb{R}^1$. For *this choice* of the brane directions within the stack of D3-branes, the D3-transverse subspace \mathbb{R}_{456789}^6 splits into the sum of the Coulomb subspace $\mathbb{R}_{\text{c}}^3 = \mathbb{R}_{\text{br}}^3$ and the Higgs subspace $\mathbb{R}_{\text{h}}^3 = \mathbb{R}^1 \times \mathbb{R}_{89}^2$. Now an NS5-brane must be parallel to $\mathbb{R}_{\text{br}}^3 \times \mathbb{R}_{\text{c}}^3$, while a D5-brane must be parallel to $\mathbb{R}_{\text{br}}^3 \times \mathbb{R}_{\text{h}}^3$.

After the topological twisting, the physics of the stack of D3-branes spanning \mathbb{R}_{0123}^3 is described by the Kapustin-Witten model [KW07] of electro-magnetic geometric Langlands duality at $t = i$ (i.e., at Galois side) in the presence of D5 and NS5 Gaiotto-Witten interfaces [GW09].

Now replace \mathbb{R}_{0123}^4 with M^4 of (2.1), so that the IIB space-time is

$$M_{\text{IIB}}^{10} = \mathbb{R}_0 \times \mathbb{R}_{12}^2 \times S_3^1 \times \mathbb{R}_{4567}^4 \times \mathbb{R}_{89}^2.$$

The first (basic) HW brane arrangement corresponds to the choice $\mathbb{R}_{\text{br}}^3 = \mathbb{R}_{012}^3$, so its m NS5-branes stretch along $\mathbb{R}_{012}^3 \times \mathbb{R}_{456}^3$ and its D5-branes stretch along $\mathbb{R}_{012}^3 \times \mathbb{R}_{789}^3$.

The second (additional) HW arrangement has the form $\mathbb{R}_{\text{br}}^3 = \mathbb{R}_0^1 \times \mathbb{R}_{\text{dr}}^1 \times S_3^1$, where the ‘directional’ subspace $\mathbb{R}_{\text{dr}}^1 \subset \mathbb{R}_{12}^2$ is tangent to the brane interface which is represented by its curve α_I in the \mathbb{R}_{12}^2 plane.

In sting theory the compactification of S_3^1 within the KW theory is a result of performing T -duality on S_3^1 . This duality turns the IIB theory into IIA theory and the stack of D3-branes turns into the stack of D2-branes stretched along \mathbb{R}_{012}^3 . Now their movements are described by a 3d TQFT. Since the circle S_3^1 intersects m NS5-branes of the basic HW arrangement, the T -duality turns their orthogonal space $S_3^1 \times \mathbb{R}_{789}^3$ into an m -centered Taub-NUT space $m\text{TN}$, so the IIA space-time is

$$M_{\text{IIA}}^{10} = \mathbb{R}_0 \times \mathbb{R}_{12}^2 \times \mathbb{R}_{456}^3 \times m\text{TN}.$$

D5-branes of the first HW arrangement become D6-branes, which wrap $\mathbb{R}_{012}^3 \times m\text{TN}$.

After the T -duality, the NS5 and D5 branes of the additional HW arrangement turn into NS5 and D4-branes of IIA, which separate the stack of D2-branes into domains. In other words, they become the interfaces of the 3d B-model and hence they are described by Lagrangian correspondences. The form of the Lagrangian correspondence for an additional brane depends on which 2-cycle this brane wraps inside the Taub-NUT space $m\text{TN}$.

3. MATHEMATICAL PERSPECTIVE

This section provides a motivation for the definition of our main working category. We do not provide details for the constructions in this section. The main two-category that we work with appears in section 3.4 and the details of the main working category are spelled out in this section.

In [KRS09] and [KR10] a two-category $\ddot{\text{Cat}}(\mathcal{Y})$ was associated to a symplectic variety \mathcal{Y} . This construction admits a generalization to the equivariant case, namely, a 2-category $\ddot{\text{Cat}}^G(\mathcal{Y})$ emerges when a group G has a hamiltonian action on \mathcal{Y} . We present an three-categorical description of this category.

3.1. Large three-category $\ddot{\text{Cat}}_{\text{sym}}$: a proposal. The objects of $\text{Obj}(\ddot{\text{Cat}}_{\text{sym}})$ are holomorphic symplectic manifolds. The morphisms between two such manifolds $\mathbf{X}, \mathbf{Y} \in \text{Obj}(\ddot{\text{Cat}}_{\text{sym}})$ form a 2-category $\text{Hom}(\mathbf{X}, \mathbf{Y})$ whose objects are fibrations with lagrangian bases:

$$(3.1) \quad (F, L, f: F \rightarrow L), \quad L \subset \mathbf{X} \times \mathbf{Y} \text{ is a Lagrangian subvariety.}$$

As hinted in [KRS09] the fibration of F over the Lagrangian L is a categorical analog of a local system over the Lagrangian in the Fukaya category theory.

The composition of $(F, L, f) \in \text{Hom}(\mathbf{X}, \mathbf{Y})$ and $(G, L', g) \in \text{Hom}(\mathbf{Y}, \mathbf{W})$ is defined to be (H, L'', h) where

$$H := (F \times \mathbf{W}) \times_{\mathbf{Y}} (\mathbf{X} \times G),$$

while $h: H \rightarrow \mathbf{X} \times \mathbf{W}$ is the natural projection and $L'' := h(H)$. The composition is not always defined since the L'' is not always a Lagrangian.

The morphisms between the objects $(F, L, f), (F', L', f') \in \text{Hom}(\mathbf{X}, \mathbf{Y})$ is defined as follows:

$$\text{Hom}((F, L, f), (F', L', f')) := \text{Qcoh}(F \times_{\mathbf{X} \times \mathbf{Y}} F').$$

The category of quasi-coherent sheaves can be enriched to $(\infty, 1)$ -category. The study of the structure of these categories as well as $(2, \infty)$ -category of such $(\infty, 1)$ -categories is initiated in [GR17]. The above mentioned work is extremely technical and we do not attempt to use infinity categories in our work. Instead, we use the previous definition as a guiding principle.

Let us also remark that the category $\check{\check{\check{\text{Cat}}}}_{\text{sym}}$ contains a final object pt which is just a point. Thus for every $\mathbf{X} \in \text{Obj}(\check{\check{\check{\text{Cat}}}}_{\text{sym}})$ there is a related two-category

$$\check{\check{\check{\text{Cat}}}(\mathbf{X})} := \mathbb{H}om(pt, \mathbf{X}).$$

3.2. Small three-category: first steps. In this paper we construct smaller category $\check{\check{\check{\text{Cat}}}}_{\text{man}}$ that has fewer objects and fewer morphisms. As a pay-back we get a mathematically satisfactory theory for two and three-morphisms in such categories, without use of heavy machinery the derived algebraic geometry mentioned above.

The objects of the category $\check{\check{\check{\text{Cat}}}}_{\text{man}}$ are smooth algebraic varieties. The two-category of morphisms $\mathbb{H}om(\mathcal{X}, \mathcal{Y})$ between two objects $\mathbf{X}, \mathbf{Y} \in \text{Obj}(\check{\check{\check{\text{Cat}}}}_{\text{man}})$ has objects

$$(\mathcal{Z}, W), \quad \mathcal{Z} \text{ is algebraic manifold, } \quad W \in \mathbb{C}[\mathbf{X} \times \mathcal{Z} \times \mathbf{Y}].$$

For $\mathbf{X}, \mathbf{Y}, \mathbf{W} \in \text{Obj}(\check{\check{\check{\text{Cat}}}}_{\text{man}})$ and $(\mathcal{Z}, W) \in \mathbb{H}om(\mathbf{X}, \mathbf{Y})$, $(\mathcal{Z}', W') \in \mathbb{H}om(\mathbf{Y}, \mathbf{W})$ the composition is defined by:

$$(\mathcal{Z}', W') \circ (\mathcal{Z}, W) = (\mathcal{Z} \times \mathcal{Y} \times \mathcal{Z}', W' - W) \in \mathbb{H}om(\mathbf{X}, \mathbf{W}).$$

The category of morphisms $\text{Hom}((\mathcal{Z}, W), (\mathcal{Z}', W'))$ between the objects $(\mathcal{Z}, W), (\mathcal{Z}', W') \in \mathbb{H}om(\mathbf{X}, \mathbf{Y})$ is a triangulated one-category of matrix factorizations

$$\text{Hom}((\mathcal{Z}, W), (\mathcal{Z}', W')) = \text{MF}(\mathbf{X} \times \mathcal{Z} \times \mathcal{Z}' \times \mathbf{Y}, W' - W).$$

The objects of the homotopy category $\text{MF}(\mathcal{Z}, W)$, $W \in \mathbb{C}[\mathcal{Z}]$ are pairs

$$(M, D), \quad M = M_0 \oplus M_1, \quad D \in \text{Hom}_{\mathbb{C}[\mathcal{Z}]}(M, M), \quad D^2 = W,$$

where D is \mathbb{Z}_2 -graded morphism: $D(M_i) = M_{i+1}$. If we think of the matrix factorizations as two periodic curved complexes, then the homotopy is defined in exactly the same way as for usual complexes.

The standard convolution product \star (see the next section for the details) yields the monoidal structure:

$$\star : \text{MF}(\mathcal{Z} \times \mathcal{Z}', W - W') \times \text{MF}(\mathcal{Z}' \times \mathcal{Z}'', W' - W'') \rightarrow \text{MF}(\mathcal{Z} \times \mathcal{Z}'', W - W'').$$

For two objects $\mathcal{F} = (M, D)$, $\mathcal{G} = (M', D') \in \text{MF}(\mathcal{Z}, W)$ the space of morphisms is defined by:

$$\text{Hom}(\mathcal{F}, \mathcal{G}) = \{\phi \in \text{Hom}_{\mathbb{C}[\mathcal{Z}]}(M, M') \mid D \circ \phi = \phi \circ D'\}.$$

We also define $D^{\text{per}}(\mathcal{Z})$ to be a derived category of the two-periodic complexes of $\mathbb{C}[\mathcal{Z}]$ -modules. Given an element $\mathcal{F} = (M, D) \in \text{MF}(\mathcal{Z}, W)$, by inverting the sign of the component of the differential from M_0 to M_1 we obtain an element of $\text{MF}(\mathcal{Z}, -W)$ which

we denote \mathcal{F}^* . Since the potentials of the matrix factorizations add if take tensor product, we have:

$$\mathcal{H}om(\mathcal{F}, \mathcal{G}) = \mathcal{E}xt(\mathcal{F}, \mathcal{G}) = \mathcal{G} \otimes \mathcal{F}^* \in \mathcal{D}^{\text{per}}(\mathcal{Z}).$$

There is a special class of invertible morphisms that we want to discuss separately. These morphisms provide an explicit realization of the Knörrer periodicity functor [Kno87]. Namely, if $V \rightarrow \mathcal{Z}$ is a finite rank vector bundle over \mathcal{Z} and V^* is its dual then there is a canonical bilinear function

$$\mathbf{Q}_V \in \mathbb{C}[V \times V^*].$$

For any $W \in \mathbb{C}[V]$ there is a invertible functor [Kno87]:

$$\text{KN}_V : \text{MF}(\mathcal{Z}, W) \rightarrow \text{MF}(V \times V^*, W + \mathbf{Q}_V), \quad \text{KN}_V(\mathcal{F}) = \mathcal{KN}_Q \otimes \mathcal{F},$$

$$\mathcal{KN}_Q = (\Lambda^\bullet V_\theta, D = \sum v_i \frac{\partial}{\partial \theta_i} + v_i^* \theta),$$

where $\Lambda^\bullet V_\theta$ is an exterior algebra with the generators θ_i, v_i^* and v_i are dual coordinates on the spaces V and V^* .

That is $\mathcal{KN}_Q \in \text{Hom}((\mathcal{Z}, W), (V \times V^*, W + \mathbf{Q}_V)), (\mathcal{Z}, W), (V \times V^*, W + \mathbf{Q}_V) \in \mathbb{H}om(\mathbf{X}, \mathbf{Y})$. The functor KN_V is an equivalence of categories [Kno87]. To construct the inverse we need a functor:

$$\text{KN}_V^* : \text{MF}(V \times V^*, W + \mathbf{Q}_V) \rightarrow \text{MF}(\mathcal{Z}, W), \quad \text{KN}_V^*(\mathcal{F}) = \text{Hom}(\mathcal{F}, \mathcal{KN}_Q).$$

The composition of functors KN_V and KN_V^* is the duality functor \mathcal{D} . In general, for a matrix factorization $\mathcal{F} \in \text{MF}(\mathcal{Z}, W)$ the dual matrix factorization $\mathcal{D}(\mathcal{F}) \in \text{MF}(\mathcal{Z}, -W)$ is defined by

$$\mathcal{D}(\mathcal{F}) = \text{Hom}(\mathcal{F}, \mathcal{O}_{\mathcal{Z}}).$$

Since, KN_V^* and \mathcal{D} are functors we can enlarge $\text{Hom}((V \times V^*, W + \mathbf{Q}_V), (\mathcal{Z}, W))$ by including KN_V^* and \mathcal{D} . Thus the Knörrer functor internally invertible in our category.

3.3. Relation between the three-categories. There is a functor $j_3 : \overset{\dots}{\text{Cat}}_{\text{man}} \rightarrow \overset{\dots}{\text{Cat}}_{\text{sym}}$ which acts on the objects as

$$\mathbf{X} \mapsto T^*\mathbf{X}.$$

The embedding at the level of morphism is based on describing lagrangian submanifolds by generating functions [Arn89]. Let us recall the basic facts. Given a (complex) manifold \mathcal{Z} and function $W : \mathbf{X} \times \mathcal{Z} \rightarrow \mathbb{C}$ we define a subvariety $F_W \subset T^*(\mathbf{X}) \times \mathcal{Z}$ by the equations

$$\partial_{z_i} W(x, p, z) = 0, \quad \partial_{x_i} W(x, p, z) = p_i,$$

where z_i are local coordinates along \mathcal{Z} , x_i are local coordinates along \mathbf{X} and p_i are the coordinates on the cotangent space that are dual to the coordinates x_i . As shown in [Arn89], the image L_w of F_w in $T^*\mathbf{X}$ under the natural projection π is (generally) a Lagrangian subvariety.

Thus the functor j_3 at the level of homomorphisms between the objects is defined as

$$(\mathcal{Z}, W) \mapsto (F_W, \pi, L_W).$$

The real problem arises at the level of morphisms between the morphisms of objects. It is tempting to say that we have a functor

$$\mathrm{MF}(\mathbf{X} \times \mathcal{Z} \times \mathcal{Z}' \times \mathbf{Y}, W' - W) \longrightarrow \mathrm{DG}_{\mathrm{coh}}^{\mathrm{per}}(F_w \times_{T^*(\mathbf{X} \times \mathbf{Y})} F_{w'}).$$

It is not clear to the authors how one could construct such a functor in a canonical way. One option here is to use the functor $\mathcal{H}om$, if we fix some element $\mathcal{F} \in \mathrm{MF}(\mathbf{X} \times \mathcal{Z} \times \mathcal{Z}' \times \mathbf{Y}, W' - W)$; then we obtain a functor

$$\mathcal{H}om(\mathcal{F}, \cdot): \mathrm{MF}(\mathbf{X} \times \mathcal{Z} \times \mathcal{Z}' \times \mathbf{Y}, W' - W) \rightarrow \mathrm{DG}_{\mathrm{coh}}^{\mathrm{per}}(\mathbf{X} \times \mathcal{Z} \times \mathcal{Z}' \times \mathbf{Y}).$$

However, it is not clear whether we can make a choice of \mathcal{F} canonical.

3.4. Equivariant version of the three-category $\check{\mathrm{C}}\check{\mathrm{a}}\check{\mathrm{t}}_{\mathrm{man}}$. If \mathbf{X} is an affine variety with an action of G and $T^*\mathbf{X}$, then an object in the 2-category $\check{\mathrm{C}}\check{\mathrm{a}}\check{\mathrm{t}}(T^*\mathbf{X})$ is a pair (\mathcal{Z}, W) , where the auxiliary variety \mathcal{Z} is an affine variety with the action of G , while W is a superpotential: $W \in \mathbb{C}[\mathbf{X} \times \mathcal{Z}]^G$. Respectively, for \mathbf{X} with action of G and \mathbf{Y} with action of H the space of morphisms $\mathbb{H}\mathrm{om}(\mathbf{X}, \mathbf{Y})$ consists of an the pairs (\mathcal{Z}, W) , $W \in \mathbb{C}[\mathbf{X} \times \mathcal{Z} \times \mathbf{Y}]^{G \times H}$ and \mathcal{Z} has an action of $G \times H$.

For a pair $(\mathcal{Z}^1, W^1), (\mathcal{Z}^2, W^2) \in \mathbb{H}\mathrm{om}(\mathbf{X}, \mathbf{Y})$ we define the category of morphisms as category of $G \times H$ -equivariant matrix factorizations.

$$\mathrm{Hom}((\mathcal{Z}^1, W^1), (\mathcal{Z}^2, W^2)) = \mathrm{MF}_{G \times H}(\mathbf{X} \times \mathcal{Z}^1 \times \mathcal{Z}^2 \times \mathbf{Y}, W^2 - W^1).$$

In most of the cases we omit the group from the notation, on the other hand if we want to emphasize the presence of the group action we use notation (\mathbf{X}, G) for the object with G -action. Similarly, sometimes we include the group in the notation of the composition of morphisms, that is $(\mathbf{X}, G), (\mathbf{Y}, H), (\mathbf{W}, K) \in \check{\mathrm{C}}\check{\mathrm{a}}\check{\mathrm{t}}_{\mathrm{man}}$ and $(\mathcal{Z}_{xy}, W_{xy}) \in \mathbb{H}\mathrm{om}((\mathbf{X}, G), (\mathbf{Y}, H)), (\mathcal{Z}_{yw}, W_{yw}) \in \mathbb{H}\mathrm{om}((\mathbf{Y}, H), (\mathbf{W}, K))$ then

$$(3.2) \quad (\mathcal{Z}_{xw}, W_{xw}) = (\mathcal{Z}_{yw}, W_{yw}) \circ_H (\mathcal{Z}_{xy}, W_{xy}) = (\mathcal{Z}_{yw} \times \mathbf{Y} \times \mathcal{Z}_{xy}/H, W_{yw} - W_{xy})$$

Furthermore, for

$$(3.3) \quad \mathcal{F}_{xy}^{12} \in \mathrm{Hom}((\mathcal{Z}_{xy}^1, W_{xy}^1), (\mathcal{Z}_{xy}^2, W_{xy}^2)), \quad \mathcal{F}_{yw}^{12} \in \mathrm{Hom}((\mathcal{Z}_{yw}^1, W_{yw}^1), (\mathcal{Z}_{yw}^2, W_{yw}^2))$$

the vertical composition is defined by

$$\mathcal{F}_{xw}^{12} = \mathcal{F}_{yw}^{12} \circ_H \mathcal{F}_{xy}^{12} := (\pi_{Y\mathbf{W}}^*(\mathcal{F}_{yw}^{12}) \otimes \pi_{XY}^*(\mathcal{F}_{xy}^{12}))^{H^2},$$

where π_{yw}, π_{xy} are the projections from $(\mathbf{X} \times \mathcal{Z}_{yw}^1 \times \mathcal{Z}_{yw}^2 \times \mathbf{Y}) \times (\mathbf{Y} \times \mathcal{Z}_{xy}^1 \times \mathcal{Z}_{xy}^2 \times \mathbf{W})$ to the first and the last factors, respectively and

$$(3.4) \quad (\mathcal{Z}_{xy}^i, W_{xy}^i) \in \mathbb{H}\mathrm{om}((\mathbf{X}, G), (\mathbf{Y}, H)), \quad (\mathcal{Z}_{yw}^i, W_{yw}^i) \in \mathbb{H}\mathrm{om}((\mathbf{Y}, H), (\mathbf{W}, K)).$$

Similarly, we define the horizontal composition for \mathcal{F}_{xy}^{12} as before and

$$(3.5) \quad \mathcal{F}_{xy}^{23} \in \mathrm{Hom}((\mathcal{Z}_{xy}^2, W_{xy}^2), (\mathcal{Z}_{xy}^3, W_{xy}^3)),$$

here we assume (3.4). In more details, we have

$$\mathcal{F}_{xy}^{13} = \mathcal{F}_{xy}^{23} \star \mathcal{F}_{xy}^{12} = \pi_{13*}(\pi_{12}^*(\mathcal{F}_{xy}^{23}) \otimes \pi_{23}^*(\mathcal{F}_{xy}^{12})),$$

where $\pi_{ij}: \mathbf{X} \times \mathcal{Z}_1 \times \mathcal{Z}_2 \times \mathcal{Z}_3 \times \mathbf{Y} \rightarrow \mathbf{X} \times \mathcal{Z}_i \times \mathcal{Z}_j \times \mathbf{Y}$.

We often omit the group super-script when the group is clear from the context. Finally, let us observe that the vertical composition satisfies the exchange rule:

Proposition 3.4.1. *For any $(\mathbf{X}, G), (\mathbf{Y}, H), (\mathbf{W}, K) \in \check{\text{Cat}}_{\text{man}}$ and $(\mathcal{L}_{xy}, W_{xy}^i), (\mathcal{L}_{yw}, W_{yw}^i)$ as in (3.4) and $\mathcal{F}_{xy}^{12}, \mathcal{F}_{yw}^{12}$ as in (3.3), \mathcal{F}_{xy}^{23} as in (3.5) and*


$$\mathcal{F}_{yw}^{23} \in \text{Hom}((\mathcal{L}_{yw}^2, W_{yw}^2), (\mathcal{L}_{yw}^3, W_{yw}^3))$$

there is a three-isomorphism:

$$E(\mathcal{L}_{xy}^\bullet; \mathcal{L}_{yw}^\bullet; \mathcal{F}_{xy}^{\bullet\bullet}; \mathcal{F}_{yw}^{\bullet\bullet}) : (\mathcal{F}_{yw}^{12} \circ \mathcal{F}_{xy}^{12}) \star (\mathcal{F}_{yw}^{23} \circ \mathcal{F}_{xy}^{23}) \rightarrow (\mathcal{F}_{yw}^{12} \star \mathcal{F}_{yw}^{23}) \circ (\mathcal{F}_{xy}^{12} \star \mathcal{F}_{xy}^{23}).$$

Both sides of the last equation are functors between one-categories and the three-isomorphism E is a natural transformation of between these functors.


Proof. In the proof below we use the short hand notation $\mathcal{L}_{\bullet\bullet}^{i_1 \dots i_k} = \mathcal{L}_{\bullet\bullet}^{i_1} \times \dots \times \mathcal{L}_{\bullet\bullet}^{i_k}$. First we observe that $(\mathcal{F}_{yw}^{12} \circ \mathcal{F}_{xy}^{12}) \star (\mathcal{F}_{yw}^{23} \circ \mathcal{F}_{xy}^{23})$ is computed with the push-forwards and pull-backs along the maps in the diagram and taking H^2 -quotient:



here all the maps are the natural projections. In more details, the matrix factorizations $\mathcal{F}_{\bullet\bullet}^{ij}$ are the matrix factorizations on the spaces in the boxes. Thus to obtain $(\mathcal{F}_{yw}^{12} \circ \mathcal{F}_{xy}^{12}) \star (\mathcal{F}_{yw}^{23} \circ \mathcal{F}_{xy}^{23})$ we pull-back and push-forward these matrix factorizations along the solid arrow until we reach the space in the center of the bottom row.

The dashed arrow maps indicate the natural projections that make the diagram commute. By functoriality of the pull-back we can obtain $(\mathcal{F}_{yw}^{12} \circ \mathcal{F}_{xy}^{12}) \star (\mathcal{F}_{yw}^{23} \circ \mathcal{F}_{xy}^{23})$ by first pulling back along the dashed arrow and then pushing forward along the central down arrow.

Similarly, we draw the diagram for the maps that participate in the construction of $(\mathcal{F}_{yw}^{12} \star \mathcal{F}_{yw}^{23}) \circ (\mathcal{F}_{xy}^{12} \star \mathcal{F}_{xy}^{23})$:



here the maps are the natural projections and

$$\star = \mathbf{X} \times \mathcal{L}_{xy}^{13} \times \mathbf{Y}^2 \times \mathcal{L}_{yw}^{13} \times \mathbf{W}.$$

We obtain $(\mathcal{F}_{yw}^{12} \star \mathcal{F}_{yw}^{23}) \circ (\mathcal{F}_{xy}^{12} \star \mathcal{F}_{xy}^{23})$ by the sequence of pull-backs and push-forwards that starts from the boxed spaces and ends at \star . By setting

$$\ast = \mathbf{X} \times \mathcal{L}_{xy}^{123} \times \mathbf{Y}^2 \times \mathcal{L}_{yw}^{123} \times \mathbf{W}$$

we can complete the diagram to the commuting diagram where the dashed arrows are the natural projections.

The dashed arrow maps are the same maps as in the previous diagram. Thus to complete our proof we need to show that $(\mathcal{F}_{yw}^{12} \star \mathcal{F}_{yw}^{23}) \circ (\mathcal{F}_{xy}^{12} \star \mathcal{F}_{xy}^{23})$ can be obtained by the pull-back along the dashed arrows followed with the push-forward along the vertical dashed arrow.

The last statement follows from a repeated application the base change relation. Indeed, first we apply base change to the commuting squares that have as sides the vertical dashed arrow and the dotted arrows. Finally, we apply the functoriality of the pull-backs since the composition of the pull-backs along the vertical arrow and the dotted arrows are equal to the pull-back along the corresponding non-vertical dashed arrows.

Each base change in the construction is facilitated by a particular invertible morphism between the matrix factorizations. The composition of these morphisms yields the three-morphism $E(\mathcal{L}_{xy}^\bullet, \mathcal{L}_{yw}^\bullet; \mathcal{F}_{xy}^\bullet, \mathcal{F}_{yw}^\bullet)$. The base change morphisms are natural transformations. Hence E is a natural transformation. \square

In the cases when the first group of arguments of $E(\dots)$ is clear we abbreviate the notation as $E(; \mathcal{F}_{xy}^{12}, \mathcal{F}_{xy}^{23}, \mathcal{F}_{yw}^{12}, \mathcal{F}_{yw}^{23})$.

Let us also discuss the exchange relation for two-morphisms and three-morphisms. Let us fix the objects (\mathbf{X}, G) , (\mathbf{Y}, K) and let

$$(3.6) \quad (\mathcal{L}^i, W^i) \in \mathbb{H}\text{om}(\mathbf{X}, \mathbf{Y}), \quad \mathcal{F}^{ii+1}, \mathcal{G}^{ii+1}, \mathcal{H}^{ii+1} \in \text{Hom}((\mathcal{L}^i, W^i), (\mathcal{L}^{i+1}, W^{i+1})).$$

Respectively, we define $\mathcal{F}^{ii+2} = \mathcal{F}^{ii+1} \star \mathcal{F}^{i+1, i+2}$, $\mathcal{G}^{ii+2} = \mathcal{G}^{ii+1} \star \mathcal{G}^{i+1, i+2}$.

Let us also fix notation for the three-morphisms

$$(3.7) \quad u_{fg}^{ii+1} \in \text{Hom}(\mathcal{F}^{ii+1}, \mathcal{G}^{ii+1}), \quad u_{gh}^{ii+1} \in \text{Hom}(\mathcal{G}^{ii+1}, \mathcal{H}^{ii+1}).$$

We denote by \bullet for the composition of morphisms in the category of matrix factorizations, for example $u_{gh}^{12} \bullet u_{fg}^{12} \in \mathcal{H}\text{om}(\mathcal{F}^{12}, \mathcal{H}^{12})$.

On the other hand we can define the horizontal composition $u_{fg}^{23} \star u_{fg}^{12} \in \mathcal{H}\text{om}(\mathcal{F}^{13}, \mathcal{G}^{13})$ by

$$(3.8) \quad u_{fg}^{23} \star u_{fg}^{12} = \pi_{13*}(\mathcal{G}^{12} \otimes \pi_{23}^*(u_{fg}^{23}) \bullet \pi_{12}^*(u_{fg}^{12}) \otimes \mathcal{F}^{23}),$$

$$\pi_{12}^*(u_{fg}^{12}) \otimes \mathcal{F}^{23} = \pi_{12}^*(u_{fg}^{12}) \otimes \text{Id}_{\pi_{23}^*(\mathcal{F}^{23})} \in \mathcal{H}\text{om}(\pi_{12}^*(\mathcal{F}^{12}) \otimes \pi_{23}^*(\mathcal{F}^{23}), \pi_{12}^*(\mathcal{G}^{12}) \otimes \pi_{23}^*(\mathcal{F}^{23})),$$

$$\mathcal{G}^{12} \otimes \pi_{23}^*(u_{fg}^{23}) = \text{Id}_{\pi_{12}^*(\mathcal{G}^{12})} \otimes \pi_{23}^*(u_{fg}^{23}) \in \mathcal{H}\text{om}(\pi_{12}^*(\mathcal{G}^{12}) \otimes \pi_{23}^*(\mathcal{F}^{23}), \pi_{12}^*(\mathcal{G}^{12}) \otimes \pi_{23}^*(\mathcal{G}^{23})).$$

The exchange rule in this setting is given by

Proposition 3.4.2. *In notations of (3.6),(3.7)(3.8) we have*

$$(u_{gh}^{23} \star u_{gh}^{12}) \bullet (u_{fg}^{23} \star u_{fg}^{12}) = (u_{gh}^{23} \bullet u_{fg}^{23}) \star (u_{gh}^{12} \bullet u_{fg}^{12}).$$

Proof. To show that the morphisms of the matrix factorizations are equal it is enough to check that the morphisms are equal as morphisms of stalks at a point $z \in \mathbf{X} \times \mathcal{Z}^1 \times \mathcal{Z}^3 \times \mathbf{Y}$. The convolution \star intertwines $i_{(x,z_1,y)}^* \times i_{(x,z_3,y)}^*$ with i_z^* where $z = (x, z_1, z_3, y)$. Thus it is enough to show the statement in the case $\mathbf{X} = \mathbf{Y} = \mathcal{Z}_1 = \mathcal{Z}_3 = \mathfrak{p}$. In this case the statement of the theorem is equivalent to the functoriality of the push-forward.

Indeed, in this situation $\mathcal{F}^{12}, \mathcal{G}^{12}, \mathcal{H}^{12} \in \text{MF}(\mathcal{Z}^2, W)$, $\mathcal{F}^{23}, \mathcal{G}^{23}, \mathcal{H}^{23} \in \text{MF}(\mathcal{Z}^2, -W)$. Respectively, the convolutions $\mathcal{F}^{12} \star \mathcal{F}^{23} = H^*(\mathcal{F}^{12} \otimes \mathcal{F}^{23})$, $\mathcal{G}^{12} \star \mathcal{G}^{23} = H^*(\mathcal{G}^{12} \otimes \mathcal{G}^{23})$, $\mathcal{H}^{12} \star \mathcal{H}^{23} = H^*(\mathcal{H}^{12} \otimes \mathcal{H}^{23})$ are the derived push-forwards $R\pi_*$ to the point, $R\pi_* : \mathcal{Z} \rightarrow \mathfrak{p}$.


Since the derived push-forward is a functor, the convolutions of the endomorphism are defined by means of functoriality $u_{gh}^{23} \star u_{gh}^{12} = R\pi_*(u_{gh}^{23} \otimes u_{gh}^{12})$, $u_{fg}^{23} \star u_{fg}^{12} = R\pi_*(u_{fg}^{23} \otimes u_{fg}^{12})$, $(u_{gh}^{23} \bullet u_{fg}^{23}) \star (u_{gh}^{12} \bullet u_{fg}^{12}) = R\pi_*((u_{gh}^{23} \bullet u_{fg}^{23}) \otimes (u_{gh}^{12} \bullet u_{fg}^{12}))$. Thus the statement of the proposition is equivalent to the relation:

$$R\pi_*(u_{gh}^{23} \otimes u_{gh}^{12}) \bullet R\pi_*(u_{fg}^{23} \otimes u_{fg}^{12}) = R\pi_*((u_{gh}^{23} \bullet u_{fg}^{23}) \otimes (u_{gh}^{12} \bullet u_{fg}^{12}))$$

which follows from the functoriality of $R\pi_*$. \square

3.5. Categories of linear quivers and NS5 interfaces. In this paper we consider the subcategories $\check{\text{C}}\text{at}(\tilde{A}_m)$ (resp. $\check{\text{C}}\text{at}(A_m)$) of $\check{\text{C}}\text{at}$ formed by Nakajima quiver varieties of (Dynkin diagram) type \tilde{A}_m (resp. A_m) and describe some generalized Lagrangian correspondences connecting them. These correspondences are of two distinct types: NS5 and D4 (or D5). We will describe mostly the NS5 correspondences (1.3) for $m = 1$. Their auxiliary varieties originate from arrows similar to the ones appearing in Nakajima quivers, which could be expected given that both come from NS5 branes in the super-string theory. We will also conjecture the D4 Lagrangian correspondences connecting the so-called commuting varieties, that is, Nakajima varieties of the \tilde{A}_0 quiver without framing (1.4).


3.5.1. *NS5 interfaces between \tilde{A}_{m-1} quiver varieties.* The \tilde{A}_{m-1} quiver has the form



where $i = 1, \dots, m$. It is ‘colored’ by the numbers $\mathbf{n} = (n_1, \dots, n_m)$ and $\mathbf{r} = (r_1, \dots, r_m)$. We denote this quiver by $\tilde{A}_m(\mathbf{n}; \mathbf{r})$. An edge between the i -th and $(i+1)$ -st circles corresponds to the i -th basic, in the sense of section 2, NS5 interface, the number n_i refers to the gauge group $U(n_i)$ between the $(i-1)$ -st and i -th NS5-interfaces, and there are r_i D5-interfaces there.

Remark 3.5.2. A ‘linear’ A_m quiver appears as a particular case of the \tilde{A}_{m-1} quiver if one sets $n_m = r_m = 0$.

In order to describe additional interfaces as Lagrangian correspondences, we have to present the quiver varieties as cotangent bundles. Hence we choose the orientation of quiver edges (for example, clockwise and from circles to boxes) and denote the corresponding maps as X_i and v_i :



The quiver variety $\mathcal{Y}_{\mathbf{n}, \mathbf{r}}$ is the stable part of the cotangent bundle

$$\mathcal{Y}_{\mathbf{n}, \mathbf{r}} = (\mathrm{T}^* \mathcal{X}_{\mathbf{n}, \mathbf{r}})^{st}, \quad \mathcal{X}_{\mathbf{n}, \mathbf{r}} = \left(\prod_{i=1}^m \mathrm{Hom}(\mathbb{C}^{n_i}, \mathbb{C}^{n_{i+1}}) \right) \times \left(\prod_{i=1}^m \mathrm{Hom}(\mathbb{C}^{n_i}, \mathbb{C}^{r_i}) \right).$$

The stable locus is defined by means the hamiltonian action of the group

$$G_{\mathbf{n}} = \prod_{i=1}^m \mathrm{GL}(n_i).$$

In more details, given a character of $\chi = (\chi_1, \dots, \chi_m)$ of $G_{\mathbf{n}}$ the χ -stable locus of $\mathrm{T}^* \mathcal{X}_{\mathbf{n}, \mathbf{r}}$ is the projection of the GIT stable locus of $\mathrm{T}^* \mathcal{X}_{\mathbf{n}, \mathbf{r}} \times \mathbb{C}_{\chi_1}^* \times \dots \times \mathbb{C}_{\chi_m}^*$, see for example [Nak99]. The χ -stable condition can also be described by explicit linear algebra construction as explained in [Nak99]. If \mathbf{r} is zero, we do not impose any stability constraints.

Consider two affine quivers $Q = \tilde{A}_m(\mathbf{n}; \mathbf{r})$ and $Q' = \tilde{A}_m(\mathbf{n}'; \mathbf{r}')$ with the same m but, generally, different dimensions: $(\mathbf{n}; \mathbf{r})$ and $(\mathbf{n}'; \mathbf{r}')$. The NS5-interface auxiliary variety \mathcal{Z}

has an arrow description, that is, it is based on the spaces of linear maps. First of all, we introduce the maps $\varphi_i \in \text{Hom}(\mathbb{C}^{n_i}, \mathbb{C}^{n'_i})$ and $\varphi_i^{\text{fr}} \in \text{Hom}(\mathbb{C}^{r_i}, \mathbb{C}^{r'_i})$. We *fix* the values of all φ_i^{fr} and require that they all have the highest rank. Then for each ‘horizontal square’

$$(3.9) \quad \begin{array}{ccc} n_{i+1} & \xrightarrow{\varphi_{i+1}} & n'_{i+1} \\ \uparrow X_i & \nearrow \psi_i & \uparrow X'_i \\ n_i & \xrightarrow{\varphi_i} & n'_i \end{array} \quad \text{or} \quad \begin{array}{ccc} n_{i+1} & \xrightarrow{\varphi_{i+1}} & n'_{i+1} \\ \uparrow X_i & \nearrow \psi_i & \uparrow X'_i \\ n_i & \xrightarrow{\varphi_i} & n'_i \end{array}$$

\tilde{X}_i \tilde{X}'_i

and for each ‘vertical square’

$$(3.10) \quad \begin{array}{ccc} r_i & \xrightarrow{\varphi_i^{\text{fr}}} & r'_i \\ \uparrow v_i & \nearrow \psi_i^{\text{fr}} & \uparrow v'_i \\ n_i & \xrightarrow{\varphi_i} & n'_i \end{array} \quad \text{or} \quad \begin{array}{ccc} r_i & \xrightarrow{\varphi_i^{\text{fr}}} & r'_i \\ \uparrow v_i & \nearrow \psi_i^{\text{fr}} & \uparrow v'_i \\ n_i & \xrightarrow{\varphi_i} & n'_i \end{array}$$

\tilde{v}_i \tilde{v}'_i


we choose either the first diagonal (on the left) or the second diagonal (on the right) and add the corresponding linear maps $\psi_i, \psi_i^{\text{fr}}$ and, in case of the second diagonal choice we also add maps $\tilde{X}_i, \tilde{X}'_i, \tilde{v}_i$ and \tilde{v}'_i . The variety \mathcal{Z} is formed by all these maps, including φ_i , while the superpotential W is the sum of superpotentials W_{square} of individual squares, the latter being the (alternating) sums of traces along the boundaries of triangles and digons:

$$W_{\text{square}} = \begin{cases} \text{Tr}(\psi_i X'_i \varphi_i) - \text{Tr}(\psi_i \varphi_{i+1} X_i), \\ (\text{Tr}(X'_i \tilde{X}'_i) - \text{Tr}(\tilde{X}'_i \varphi_{i+1} \psi_i)) - (\text{Tr}(\tilde{X}_i X_i) - \text{Tr}(\tilde{X}_i \psi_i \varphi_i)) \\ \text{Tr}(\psi_i^{\text{fr}} v'_i \varphi_i) - \text{Tr}(\psi_i^{\text{fr}} \varphi_i^{\text{fr}} v_i) \\ (\text{Tr}(\tilde{v}'_i v'_i) - \text{Tr}(\psi_i^{\text{fr}} \tilde{v}'_i \varphi_i^{\text{fr}})) - (\text{Tr}(\tilde{v}_i v_i) - \text{Tr}(\psi_i^{\text{fr}} \varphi_i v_i)) \end{cases}$$

We get more interface squares by swapping the quivers Q and Q' in the diagrams (3.9) and (3.10), that is, reversing the directions of horizontal arrows and adjusting the choices of diagonals.

Not all of these choices are compatible with stability conditions on the arrows of the quivers, and some of these choices are Legendre-equivalent if we replace the spaces at the quiver vertices by their dual spaces and, at the same time, reverse the directions of all arrows. For example, the generalized Lagrangian correspondence of the second diagram

of (3.10) can be equivalently presented, if we reverse the directions of vertical arrows:




In our previous work [OR18e],[OR18b],[OR18d] and in the section 5 we work with the last choice of the direction of circle-square edge. In the rest of this section we work with convention of formulas (3.9), (3.10) since this choice is closer the standard quiver description of Grassmannians.

Remark 3.5.3. If $\mathbf{r}' = \mathbf{r}$ while $\mathbf{n} \leq \mathbf{n}'$, that is, $n_i \leq n'_i$ for all i , and one requires all horizontal arrow maps to have highest rank, then the first correspondences of (3.9) and (3.10) describe the ‘standard’ interface which has a simple symplectic interpretation. Vertical arrow maps describe representations of Q and Q' , the horizontal arrows establish the representation of Q as a subrepresentation of Q' , and one takes the conormal bundle to this arrangement within the product of two full quiver varieties.


Now we consider several special cases.

3.5.4. *NS5 interfaces between Grassmannians.* This example was communicated to us by Tudor Dimofte and it represents an interface between two A_1 quivers, obtained from \tilde{A}_1 quivers by setting $n_2 = r_2 = 0$. The corresponding quiver varieties are cotangent bundles $T^*\text{Gr}_{n,r}$ and $T^*\text{Gr}_{n',r}$. If $n \leq n'$, then the first diagram of (3.10) yields a well-known generalized Lagrangian correspondence:



Here φ^{fr} is a fixed isomorphism, while Grassmannian stability conditions require that v and v' have highest ranks (n and n'). The potential and the space of maps $\varphi, \varphi^{\text{fr}}$ realizes NS5 defect of charge $n - n'$, $\text{NS5}^{(n-n')}$ in three category $\check{\text{Cat}}(A_0)$.

The criticality condition for W with respect to ψ^{fr} implies the commutativity of the vertical and horizontal arrows, so the commutative subdiagram



represents the partial flag formed by the subspaces of both Grassmannians: $\mathbb{C}^n \subset \mathbb{C}^{n'} \subset \mathbb{C}^r$. In addition, the criticality condition with respect to φ guarantees that the maps $\psi^{\text{fr}}\varphi^{\text{fr}}$ and $\varphi\psi^{\text{fr}}$ represent the vectors in their cotangent spaces.

3.5.5. *NS5 interfaces between instanton moduli spaces.* Now we consider the interfaces between two \tilde{A}_0 quivers with $n \leq n'$ and $r = r'$ which yield the functors $\hat{\square}_k$ and $\hat{\blacksquare}_k$ of (1.3). The NS5 interfaces $\text{NS5}^{(n-n')}$ of charge $n - n'$ are described by diagrams

(3.11)

and the superpotentials are

$$W_1 = W(X, X') + W^{\text{fr}}, \quad W_2 = \text{Tr}(XY) + \text{Tr}(X'Y') - W(Y, Y') + W^{\text{fr}},$$

where

$$W(X, X') = \text{Tr}(X'\varphi\psi) - \text{Tr}(X\psi\varphi), \quad W^{\text{fr}} = \text{Tr}(\psi^{\text{fr}}v'\varphi) - \text{Tr}(\psi^{\text{fr}}\varphi^{\text{fr}}v).$$

Note that the second interface is the Legendre transform of the first interface with respect to the loop arrows X, X' .

3.6. **NS5 and D5 interfaces between commuting varieties.** The simplest oriented quiver of the \tilde{A}_0 type corresponds to $r = 0$ and, hence, consists of a single circle and a

loop:



For the family of these quivers, in addition to NS5 generalized Lagrangian correspondence we introduce another (conjecturally, D5) correspondence (and its Legendre transform), borrowing them from bow edges of Nakajima-Cherkis quiver varieties. The following discussion is parallel to that of [RR21].

In a bow-arrow quiver Q , to an edge e connecting vertices with indices n_1 and n_2 one assigns a symplectic variety \mathcal{Y}_e with the hamiltonian action of $\mathrm{GL}(n_1) \times \mathrm{GL}(n_2)$, and then the quiver variety \mathcal{X}_Q is the result of the hamiltonian reduction:

$$\mathcal{X}_Q = \left(\prod_e \mathcal{Y}_e \right) // \prod_i \mathrm{GL}(n_i).$$


An edge variety \mathcal{Y}_e determines a generalized Lagrangian correspondence connecting two commuting varieties:

$$(3.12) \quad \begin{array}{c} n_1 \\ \circ \\ \text{---} (\mathcal{Y}_e, W_e) \text{---} \\ \circ \\ n_2 \end{array} \quad , \quad W_e = \mathrm{Tr} X_2 \mu_2 - \mathrm{Tr} X_1 \mu_1.$$

and its Legendre transform


$$\begin{array}{c} n_1 \\ \circ \\ \text{---} (\mathcal{Y}_e, W_e) \text{---} \\ \circ \\ n_2 \end{array} \quad , \quad W_e = (\mathrm{Tr} X_2 Y_2 - \mathrm{Tr} Y_2 \mu_2) - (\mathrm{Tr} X_1 Y_1 - \mathrm{Tr} Y_1 \mu_1).$$

The arrow edge variety is $\mathcal{X}^{\text{arr}} = \text{T}^*\text{Hom}(\mathbb{C}^{n_1}, \mathbb{C}^{n_2})$ and the associated correspondences match those of (3.11) for $r = 0$. The bow variety is more complicated, and its correspondence has the form



$$, \quad W_e = \text{Tr}X'Y' + \text{Tr}XY - \text{Tr}\psi(Y'\varphi - \varphi Y + \sigma\rho).$$

Its Legendre transform has a simpler diagram:



$$, \quad W_e = \text{Tr}\psi(X'\varphi - \varphi X + \sigma\rho).$$

These two correspondences yield the functors $\check{\square}_k$ and $\check{\blacksquare}_k$, $k = n - n'$ of (1.4) which realize the D5 interfaces $D5^{(k)}$ of charge k .

4. CATEGORICAL ASPECTS OF THE THEORY.

In this section we discuss the categorical setting for the proposed TQFT. In particular, we explain how de-looping procedure allows us to state a precise mathematical statement that underlies a construction of the TQFT. The delooping procedure conjecturally produces a monoidal two-category $\check{\text{Cat}}(Q)$ for a quiver Q .

4.1. Delooping. In the previous section discussed the general setting for a construction of the three-category $\check{\check{\text{Cat}}}_{\text{man}}$ as well as the simplest equivariant version of the three-category $\check{\check{\text{Cat}}}(Q)$ for a linear quiver Q . It is natural to conjecture that $\check{\check{\text{Cat}}}_{\text{man}}$ and $\check{\check{\text{Cat}}}(Q)$ are weak three-categories [GG09]. However, checking the conditions for the structural morphisms of a weak three-category is a rather challenging task. We choose a different path for constructing $\check{\check{\text{Cat}}}(Q)$. First, we define a delooped monoidal two-category $\check{\text{Cat}}(Q)$, then we define a large three-category $\check{\check{\text{Cat}}}(Q)$ as category of module two-categories over $\check{\text{Cat}}(Q)$. The three-category $\check{\check{\text{Cat}}}(Q)$ is a subcategory inside $\check{\check{\text{Cat}}}(Q)$.

To construct two-category $\mathring{\mathcal{C}}\text{at}(Q)$ we first reduce the three-category $\mathring{\mathring{\mathcal{C}}}\text{at}(Q)$ to a three-category with one object $\mathbb{O}_Q = \bigoplus_{\mathbf{n}, \mathbf{r}} Q(\mathbf{n}; \mathbf{r})$. Respectively, the category of the endomorphisms $\mathbb{H}\text{om}(\mathbb{O}_Q, \mathbb{O}_Q)$ of \mathbb{O}_Q has objects that are collections of elements

$$(\mathcal{L}_{\mathbf{n}, \mathbf{r}}, W_{\mathbf{n}, \mathbf{r}}) \in \mathbb{H}\text{om}(Q(\mathbf{n}; \mathbf{r}), Q(\phi(\mathbf{n}); \psi(\mathbf{r}))), \quad \mathbf{n}, \mathbf{r} \in \mathbb{Z}_+^Q, \quad \phi, \psi : \mathbb{Z}_+^Q \rightarrow \mathbb{Z}_+^Q.$$

The composition law for the morphisms in $\mathring{\mathcal{C}}\text{at}(Q)$ yields a monoidal structure on $\mathbb{H}\text{om}(\mathbb{O}_Q, \mathbb{O}_Q)$.

Now we declare that the two-category $\mathring{\mathcal{C}}\text{at}(Q)$ has elements of $\mathbb{H}\text{om}(\mathbb{O}_Q, \mathbb{O}_Q)$ as objects and the rest of the structure is inherited from the three-category $\mathring{\mathring{\mathcal{C}}}\text{at}(Q)$. Thus we state our main categorical

Theorem 4.1.1. *The delooped two-category $\mathring{\mathcal{C}}\text{at}(Q)$ has a structure of a monoidal two-category (without a unit) in the sense of Kapranov-Voevodsky [KV94], [BN96].*

The essence of the monoidal two-category is its monoidal structure and the corresponding coherence conditions that are imposed by the compatibility with the monoidal structure. If we ignore the monoidal structure we essentially put ourselves in the standard setting similar to the one considered in [CW07].

Proposition 4.1.2. *The composition \star and the natural composition \bullet of $\mathcal{H}\text{om}$ functors define a structure of a weak two-category on $\mathring{\mathcal{C}}\text{at}(Q)$.*

Proof. It is enough to show that $\mathbb{H}\text{om}(Q(\mathbf{n}, \mathbf{r}), Q(\mathbf{n}', \mathbf{r}'))$ is a weak two-category for a fixed choice $\mathbf{n}, \mathbf{r}, \mathbf{n}', \mathbf{r}'$. For a given $(\mathcal{L}, W) \in \mathbb{H}\text{om}(Q(\mathbf{n}, \mathbf{r}), Q(\mathbf{n}', \mathbf{r}'))$ there is a canonical choice of the convolution unit:

$$\mathbf{1}^{\mathcal{L}} \in \text{Hom}((\mathcal{L}, W), (\mathcal{L}, W)) = \text{MF}_{G(\mathbf{n}) \times G(\mathbf{n}')} (Q(\mathbf{n}, \mathbf{r}) \times \mathcal{L}^2 \times Q(\mathbf{n}', \mathbf{r}'), W^{(1)} - W^{(2)}),$$

where $W^{(1)} = \pi_i^*(W)$ is the pull-back of the potential W along two natural projections to $Q(\mathbf{n}, \mathbf{r}) \times \mathcal{L} \times Q(\mathbf{n}', \mathbf{r}')$. The unit is given by the push-forward of the structure sheaf along the inclusion $1 \times \Delta \times 1$ of $Q(\mathbf{n}, \mathbf{r}) \times \mathcal{L} \times Q(\mathbf{n}', \mathbf{r}')$ into $Q(\mathbf{n}, \mathbf{r}) \times \mathcal{L} \times \mathcal{L} \times Q(\mathbf{n}', \mathbf{r}')$, here Δ is the diagonal inclusion.

To show that $\mathbf{1}^{\mathcal{L}}$ is the convolution unit we need to use the base change transformation we need to use the base change relations in the commuting diagram of maps:

$$\begin{array}{ccccc} \mathcal{L}^3 & \xrightarrow{\Delta \times 1 \times 1} & \mathcal{L}^4 & \xleftarrow{\pi_{12} \times \pi_{23}} & \mathcal{L}^3 \\ \downarrow \pi_{23} & & \downarrow \pi_{124} & & \downarrow \pi_{13} \\ \mathcal{L}^2 & \xrightarrow{\Delta \times 1} & \mathcal{L}^3 & & \mathcal{L}^2 \end{array},$$

here $\pi_{i_1 \dots i_s}$ is the projection on the corresponding product of copies of \mathcal{L} . Thus we have

$$\begin{aligned} \mathbf{1}^{\mathcal{L}} \star \mathcal{F} &= \pi_{13*} \circ (\pi_{12} \times \pi_{23})^* \circ (\Delta \times 1 \times 1)_* \circ \pi_{23}^*(\mathcal{F}) = \\ &= \pi_{13*} \circ (\pi_{12} \times \pi_{23})^* \circ \pi_{124}^* \circ (\Delta \times 1)_*(\mathcal{F}) = \pi_{13*} \circ (\Delta \times 1)_*(\mathcal{F}) = \mathcal{F}. \end{aligned}$$

Here the second equality follows from the base-change along the commuting square and the last two equalities follow from $\pi_{124} \circ \pi_{12} \times \pi_{23} = 1$ and $\pi_{13} \circ (\Delta \times 1) = 1$. Since we use the base-change we obtain an invertible natural transformation $U_l(\mathcal{L})$ between the functor of

the left multiplication by $\mathbf{1}^{\mathcal{Z}}$ and the identity functor. Similarly, one defines the natural transformation $U_r(\mathcal{Z})$ for the right multiplication.

The first group of axioms of the weak two-category that is concerned with the interaction of the vertical composition \star and horizontal composition \circ is exactly the interchange relation from proposition 3.4.1.

Our two-category is weak because in the standard proof of the associativity of the convolution product \star one has to use the base change relation. Thus the base change natural transformations yield the associator natural transformation:

$$A(\mathcal{F}, \mathcal{G}, \mathcal{H}) : (\mathcal{F} \star \mathcal{G}) \star \mathcal{H} \rightarrow \mathcal{F} \star (\mathcal{G} \star \mathcal{H}).$$

The associator in a weak two category satisfies the pentagon axiom and the unitors U_l, U_r satisfy the triangle axioms see for example [EGNO15]. The construction of the associator and the unitors is in terms of base-change relations. Thus we can use the standard argument for convolution of coherent sheaves to show the axioms in our setting. \square

We refer to the original paper [KV94] and subsequent paper [BN96] for a streamlined treatment of the (semi-strict) monoidal two-category. In our case, the composition operation \circ is strictly associative since the equivariant pull-backs in the definition of the composition in (3.2).

Thus the only nontrivial structural two-morphism $R = R(\mathcal{F}, \mathcal{G})$ of our two-category is the two-isomorphism $\circ_{\mathcal{F}, \mathcal{G}}$ that relates the morphisms of pairs of objects $\mathcal{F} \in \text{Hom}(\mathcal{Z}, \mathcal{Z}')$, $\mathcal{G} \in \text{Hom}(\mathcal{X}, \mathcal{X}')$ as in the diagram below:

$$\begin{array}{ccc} \mathcal{Z} \circ \mathcal{X} & \xrightarrow{1_{\mathcal{Z}} \circ \mathcal{G}} & \mathcal{Z} \circ \mathcal{X}' \\ \mathcal{F} \circ 1_{\mathcal{X}} \downarrow & \swarrow R & \downarrow \mathcal{F} \circ 1_{\mathcal{X}'} \\ \mathcal{Z}' \circ \mathcal{X} & \xrightarrow{1_{\mathcal{Z}'} \circ \mathcal{G}} & \mathcal{Z}' \circ \mathcal{X}' \end{array}$$

We define $R(\mathcal{F}, \mathcal{G})$ as composition of exchange morphisms from proposition 3.4.1:

$$E(\mathcal{Z}, \mathcal{Z}, \mathcal{Z}'; \mathcal{X}, \mathcal{X}', \mathcal{X}'; \mathbf{1}_{xy}^{\mathcal{Z}}, \mathcal{F}; \mathcal{G}, \mathbf{1}_{yz}^{\mathcal{X}'}) \circ E^{-1}(\mathcal{Z}, \mathcal{Z}', \mathcal{Z}'; \mathcal{X}, \mathcal{X}, \mathcal{X}'; \mathcal{F}, \mathbf{1}_{xy}^{\mathcal{Z}'}; \mathbf{1}_{yz}^{\mathcal{X}}, \mathcal{G}),$$

where $\mathbf{1}_{\star, \star}^{\mathcal{Y}}$ stands for the identity element in $\text{Hom}(\mathcal{Y}, \mathcal{Y})$.

Proof of theorem. The two-category $\mathring{\text{Cat}}(Q)$ is almost semi-strict. Indeed, the convolution \circ is strictly associative on the objects. In more details, in notations [KV94] \circ corresponds to the composition \otimes . The structural one-morphisms that appear in the definition of the monoidal of [KV94] are $a_{A,B,C} : A \otimes (B \otimes C) \rightarrow (A \otimes B) \otimes C$, in notations of of [KV94]. The analogous one-morphisms in our setting are the identity morphisms. Respectively, the related analogs of two-morphisms $a_{A,B,C,D}, a_{u,A,B}, a_{A,u,B}, a_{A,B,u}$ are also identity in our setting.

The analogs of the structural two-morphisms $\otimes_{u,u',B}$, $\otimes_{A,v,v'}$, $\otimes_{u,v}$ in our setting are expressed in terms of the exchange natural-transformation E :

$$\begin{aligned} \otimes_{u,u',B} R(\mathcal{F}, \mathcal{G}; \mathcal{X}) &= E(\mathcal{Z}, \mathcal{Z}', \mathcal{Z}''; \mathcal{X}, \mathcal{X}; \mathcal{F}, \mathcal{G}; \mathbf{1}_{xy}^{\mathcal{X}}, \mathbf{1}_{yw}^{\mathcal{X}}) \\ \otimes_{A,v,v'} R(\mathcal{Z}; \mathcal{F}, \mathcal{G}) &= E(\mathcal{Z}, \mathcal{Z}, \mathcal{Z}; \mathcal{X}, \mathcal{X}', \mathcal{X}''; \mathbf{1}_{xy}^{\mathcal{Z}}, \mathbf{1}_{yw}^{\mathcal{Z}}; \mathcal{F}, \mathcal{G}) \\ \otimes_{u,v} R(\mathcal{F}, \mathcal{G}) &. \end{aligned}$$

The rest non-trivial two-morphisms in our setting are structural morphisms of the two-category from proposition 4.1.2. One can apply the method of MacLane strictness theorem to turn the 2-categories of 1-morphisms into the strict 2-categories, thus we suppress associator and unitor two-morphism in the discussion below.

The axioms that need to be checked for the structural two-morphisms $R(\bullet, \bullet, \bullet), R(\bullet, \bullet)$, are on the pages 221 and 222 of [KV94]. Let us first explain how one can prove the relations on the page 222 of [KV94] which are also the essential relations of the semi-strict monoidal two-category (see Lemma 4 in [BN96]). The rest of the axioms can be check by a similar computation.

The relation from the top of the page 221 of [KV94] as well as relation (??) in Lemma 4 of [BN96] can be read of the commuting diagram:

$$\begin{array}{ccccc} \mathcal{X} \circ \mathcal{Z} & \xrightarrow{(\mathcal{F}\star\mathcal{F}')\circ\mathbf{1}^{\mathcal{Z}}} & \mathcal{X}'' \circ \mathcal{Z} & & \\ \downarrow \scriptstyle{1^{\mathcal{X}}\circ\mathcal{G}} & \searrow \scriptstyle{\mathcal{F}\circ\mathbf{1}^{\mathcal{Z}}} & \swarrow \scriptstyle{\mathcal{F}'\circ\mathbf{1}^{\mathcal{Z}}} & & \downarrow \scriptstyle{1^{\mathcal{X}''}\circ\mathcal{G}} \\ & \mathcal{X}' \circ \mathcal{Z} & & & \\ \downarrow \scriptstyle{1^{\mathcal{X}'}\circ\mathcal{G}} & & \downarrow \scriptstyle{(\mathcal{F}\star\mathcal{F}')\circ\mathbf{1}^{\mathcal{Z}'}} & & \downarrow \scriptstyle{1^{\mathcal{X}''}\circ\mathcal{G}} \\ \mathcal{X} \circ \mathcal{Z}' & \xrightarrow{\text{dotted}} & \mathcal{X}'' \circ \mathcal{Z}' & & \\ \downarrow \scriptstyle{\mathcal{F}\circ\mathbf{1}^{\mathcal{Z}'}} & & \downarrow \scriptstyle{\mathcal{F}'\circ\mathbf{1}^{\mathcal{Z}'}} & & \\ & \mathcal{X}' \circ \mathcal{Z}' & & & \end{array}$$

here $\mathcal{F} \in \text{Hom}(\mathcal{X}, \mathcal{X}')$, $\mathcal{F}' \in \text{Hom}(\mathcal{X}', \mathcal{X}')$, $\mathcal{G} \in \text{Hom}(\mathcal{Z}, \mathcal{Z}')$ and $\mathcal{Z}, \mathcal{Z}' \in \mathbb{H}\text{om}(\mathbf{Y}, \mathbf{Z})$, $\mathcal{X}, \mathcal{X}', \mathcal{X}'' \in \mathbb{H}\text{om}(\mathbf{X}, \mathbf{Y})$, we omit spaces $\mathbf{X}, \mathbf{Y}, \mathbf{Z}$ from our notations.

There are two dotted paths that connect $\mathcal{X} \circ \mathcal{Z}$ to $\mathcal{X}'' \circ \mathcal{Z}'$. The relation that we are after is the equality of two compositions of two-morphisms that relate these two paths. In more details, the first sequences of two-morphisms is:

$$\begin{aligned} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X} \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}'') &\xrightarrow{R(\mathcal{F}, \mathcal{F}'; \mathcal{Z}')} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X} \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}'), \\ (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X} \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}') &\xrightarrow{R(\mathcal{F}\star\mathcal{F}', \mathcal{G})} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}'' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}'). \end{aligned}$$

The second sequence of two-morphisms is

$$\begin{aligned} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X} \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}'') &\xrightarrow{R(\mathcal{F}, \mathcal{G})} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}'), \\ (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}') &\xrightarrow{R(\mathcal{G}, \mathcal{F}')} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}'), \end{aligned}$$

$$(\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}' \circ \mathcal{Z} \rightarrow \mathcal{X}'' \circ \mathcal{Z} \rightarrow \mathcal{X}'' \circ \mathcal{Z}') \xrightarrow{R(\mathcal{F}, \mathcal{F}'; \mathcal{Z})} (\mathcal{X} \circ \mathcal{Z} \rightarrow \mathcal{X}'' \circ \mathcal{Z}' \rightarrow \mathcal{X}'' \circ \mathcal{Z}').$$

The two morphisms in the sequences are compositions of the base-change two-morphisms of some geometric maps. Let us discuss sequences of two-morphism in details. There is a sequence of maps $g_{i+1/2;k}$, $k = 1, \dots, r$ between spaces $S_{i;k}$, $g_{i+1/2;k} : S_{i+(-\epsilon_{i;k}+1)/2} \rightarrow S_{i+(\epsilon_{i;k}+1)/2}$, $\epsilon_{i;k} = \pm 1$ and the functors $F_k = f_1 \circ \dots \circ f_{m_k}$ where $f_i = g_{i+1/2;k*}$ if $\epsilon_{i;k} = 1$ and $f_i = g_{i+1/2;k}^*$ if $\epsilon_{i;k} = -1$ such that

- (1) $F_1 : (\mathcal{F}, \mathcal{F}', \mathcal{G}) \mapsto ((\mathbf{1}^{\mathcal{X}} \circ \mathcal{G}) \star (\mathcal{F} \circ \mathbf{1}^{\mathcal{Z}'}) \star \mathcal{F}' \circ \mathbf{1}^{\mathcal{Z}'})$,
- (2) There is $s > 1$ such that $F_s : (\mathcal{F}, \mathcal{F}', \mathcal{G}) \mapsto ((\mathbf{1}^{\mathcal{X}} \circ \mathcal{G}) \star (\mathcal{F} \star \mathcal{F}' \circ \mathbf{1}^{\mathcal{Z}'}))$,
- (3) $F_r : (\mathcal{F}, \mathcal{F}', \mathcal{G}) \mapsto ((\mathcal{F} \star \mathcal{F}' \circ \mathbf{1}^{\mathcal{Z}'}) \star (\mathbf{1}^{\mathcal{X}} \circ \mathcal{G}))$,
- (4) F_{k+1} is obtained from F_k either by the base-change morphism or composition of some the push-forwards or or pull-backs in the sequence $f_{i;k}$.
- (5) the corresponding sequence of natural transformations connecting F_1 with F_s is $R(\mathcal{F}, \mathcal{F}'; \mathcal{Z})$ and the sequence connecting F_s and F_r is $R(\mathcal{F} \star \mathcal{F}', \mathcal{G})$.

Similarly, there is a sequence of maps $h_{i+1/2;k}$, $\epsilon'_{i;k} = \pm 1$ and spaces $S'_{i;k}$ $k = 1, \dots, l$ that realizes the sequence of the two-morphisms $R(\mathcal{F}, \mathcal{G})$, $R(\mathcal{G}, \mathcal{F}')$, $R(\mathcal{F}, \mathcal{F}'; \mathcal{Z})$. Moreover, $h_{i+1/2;k} = g_{i+1/2;k'}$, $\epsilon'_{i;k} = \epsilon_{i;k'}$ for $k = k' = 1$ and $k = r$, $k' = l$.

Thus we can paste the sequence of the maps $h_{i+1/2;k}$ and the sequence $g_{i+1/2;k}$ along the sequences $g_{i+1/2;1}$ and $g_{i+1/2;r}$. One half of the pasted diagram yields the compositions of the natural transformations $R(\mathcal{F}, \mathcal{F}'; \mathcal{Z})$ and $R(\mathcal{F} \star \mathcal{F}', \mathcal{G})$ and the other half the composition of $R(\mathcal{F}, \mathcal{G})$, $R(\mathcal{G}, \mathcal{F}')$, $R(\mathcal{F}, \mathcal{F}'; \mathcal{Z})$. The commutativity of the diagram of maps implies the equality of two compositions of the natural transformations.

The next group of the relations on the page 221 of [KV94] are equivalent to the naturality of $R(\mathcal{F}, \mathcal{G})$, in our case the naturality is automatic since $R(\mathcal{F}, \mathcal{G})$ is a composition of natural transformations of base change. The last group of relation relation on that page follows immediately from the exchange relation from proposition 3.4.2.

The first group of relations on the page 222 of [KV94] is equivalent to naturality of $R(\mathcal{F}, \mathcal{G}; \mathcal{Z})$. Finally, the last relation on this page is based on the commuting diagram:

$$\begin{array}{ccc}
 & \mathcal{X}''' \circ \mathcal{Z} & \\
 \mathcal{F} \star \mathcal{F}' \star \mathcal{F}'' \circ \mathbf{1} & \nearrow & \mathcal{F}'' \circ \mathbf{1} \\
 \mathcal{X} \circ \mathcal{Z} & \xrightarrow{\quad} & \mathcal{X}'' \circ \mathcal{Z} \\
 \mathcal{F}' \star \mathcal{F}'' \circ \mathbf{1} & \searrow & \mathcal{F}' \circ \mathbf{1} \\
 & \mathcal{X}' \circ \mathcal{Z} &
 \end{array}$$

There are two compositions of the dotted arrow that start at $\mathcal{X} \circ \mathcal{Z}$ and end at $\mathcal{X}''' \circ \mathcal{Z}$. Respectively, there are two ways to connect these compositions by the two-morphisms which suppose to result into the same two-morphism:

$$R(\mathcal{F}, \mathcal{F}' \star \mathcal{F}''; \mathcal{Z}) \circ R(\mathcal{F}', \mathcal{F}'', \mathcal{Z}) = R(\mathcal{F} \star \mathcal{F}', \mathcal{F}''; \mathcal{Z}) \circ R(\mathcal{F}, \mathcal{F}'; \mathcal{Z}).$$

The last equality follows from the pasting argument as in the case discussed at the beginning of the proof. \square

4.2. **Module category construction of $\check{\check{\check{\text{Cat}}}}(Q)$.** Let us define $\check{\check{\check{\text{Cat}}}}(Q)$ to be a three-category of two-categories with the two-monoidal action of $\check{\check{\check{\text{Cat}}}}(Q)$. This three-category is too big and the three-category $\check{\check{\check{\text{Cat}}}}(Q)$ is a sub three-category inside $\check{\check{\check{\text{Cat}}}}(Q)$. Indeed, the product of two-categories

$$\mathbb{M}_{\mathbf{n},\mathbf{r}} = \prod_{\mathbf{n}',\mathbf{r}'} \mathbb{H}\text{om}(Q(\mathbf{n},\mathbf{r}), Q(\mathbf{n}',\mathbf{r}'))$$

is a two-category with the monoidal action of $\check{\check{\check{\text{Cat}}}}(Q)$, that is an object in $\check{\check{\check{\text{Cat}}}}(Q)$.

Respectively, the two-category $\mathbb{H}\text{om}(Q(\mathbf{n},\mathbf{r}), Q(\mathbf{m},\mathbf{s}))$ is a sub two-category of two-category of morphisms from $\mathbb{M}_{\mathbf{n},\mathbf{r}}$ to $\mathbb{M}_{\mathbf{m},\mathbf{s}}$. Thus we define the three-category $\check{\check{\check{\text{Cat}}}}(Q)$ to be the sub three-category of $\check{\check{\check{\text{Cat}}}}(Q)$ with objects $\mathbb{M}_{\mathbf{n},\mathbf{r}}$ and the morphisms between these objects as above.

4.3. **The case $Q = \tilde{A}_0$.** In the case of general quiver Q it is not immediately clear how to construct a unit in the monoidal two-category $\check{\check{\check{\text{Cat}}}}(Q)$. However, in the case $Q = \tilde{A}_0$ there is a natural candidate for the unit.

$$\mathbb{L}_{\text{id}} = \{\mathbb{L}_{\text{id}}^{n,r}\}_{n,r \geq 0}, \quad \mathbb{L}_{\text{id}}^{n,r} \in \mathbb{H}\text{om}(\tilde{A}_0(n;r), \tilde{A}_0(n;r)),$$

the one-morphism $\mathbb{L}_{\text{id}}^{n,r}$ is defined in the next section where we provide a proof for

Theorem 4.3.1. *The two-category $\check{\check{\check{\text{Cat}}}}(Q)$, $Q = \tilde{A}_0$ is a monoidal two-category with a unit \mathbb{L}_{id} .*

In the notations of the previous section the three-category $\check{\check{\check{\text{Cat}}}}_{\text{gl}}$ is a subcategory of $\check{\check{\check{\text{Cat}}}}(\tilde{A}_0)$ with objects $\tilde{A}_0(n,0)$, $n \geq 0$. We also define the framed version of the three-category $\check{\check{\check{\text{Cat}}}}_{\text{gl}}^f$ in the second half of this section. The later category is a sub-category of $\check{\check{\check{\text{Cat}}}}(\tilde{A}_0)$ with objects $\tilde{A}_0(n,1)$. Implicitly, this category was studied in [OR18e], [OR18c], [OR17] where several results in theory of knot homology were derived. We spell out the details of the categories $\check{\check{\check{\text{Cat}}}}_{\text{gl}}$ and $\check{\check{\check{\text{Cat}}}}_{\text{gl}}^f$ and state some conjectures about these categories.

4.3.2. *Objects and morphisms.* The objects of $\check{\check{\check{\text{Cat}}}}_{\text{gl}}$ are labeled by $\mathbb{Z}_{\geq 0}$:

$$\text{Obj } \check{\check{\check{\text{Cat}}}}_{\text{gl}} = \{\mathbf{n} | \mathbf{n} \in \mathbb{Z}_{\geq 0}\}.$$

The objects in the 2-category of morphisms are pairs (Z, w) , where Z is an algebraic variety with an action of $\text{GL}_n \times \text{GL}_m$:

$$\text{Obj } \mathbb{H}\text{om}(\mathbf{n}, \mathbf{m}) = \{(Z, w), \quad w \in \mathbb{C}[\mathfrak{gl}_n \times Z \times \mathfrak{gl}_m]^{\text{GL}_n \times \text{GL}_m}\}.$$

The composition of morphism $(Z, w) \in \mathbb{H}\text{om}(\mathbf{n}, \mathbf{m})$, $(Z', w') \in \mathbb{H}\text{om}(\mathbf{m}, \mathbf{k})$ is defined as

$$(Z, w) \circ (Z', w') = (Z \times \mathfrak{gl}_m \times Z' / {}_+ \text{GL}_m, w' + w) \in \mathbb{H}\text{om}(\mathbf{n}, \mathbf{k}).$$

Here the quotient is defined via GIT theory as follows. Suppose that X is a variety with a GL_m -action. A character χ of GL_m determines the trivial line bundle L_χ with GL_m -equivariant structure defined by χ . Recall that a point $x \in X$ is semistable (with respect

to L_χ) if there is $m > 0$ and $s \in \Gamma(X, L_\chi)^{GL_m}$ such that $s(x) \neq 0$. Denote

$$X/\chi GL_m := X^{ss}/GL_m.$$

Since the group of characters of GL_m is generated by \det , we introduce short-hand notations:

$$X/\pm GL_m := X/\det^{\pm 1} GL_m, \quad X/0 GL_m := X/\det^0 GL_m.$$

Two-morphism are the objects of the corresponding category of equivariant matrix factorizations. Given $(Z, w), (Z', w') \in \mathbb{H}\text{om}(\mathbf{n}, \mathbf{m})$ we define:

$$\text{Hom}((Z, w), (Z', w')) = \text{MF}_{GL_n \times GL_m}(\mathfrak{gl}_n \times Z \times Z' \times \mathfrak{gl}_m, w - w').$$

The group $GL_n \times GL_m$ is reductive, hence the equivariance of the matrix factorization (M, D) is equivalent to the condition that the group action on M commutes with the differential D .

The space of morphisms $\mathcal{H}\text{om}(\cdot, \cdot)$ between the equivariant matrix factorizations is defined to be the space of morphisms between the underlying matrix factorizations that commute the group action.

4.3.3. *Framed version of the three-category.* We enlarge slightly our category to include the framing. The objects of the new three-category $\mathring{\text{C}\ddot{\text{a}}\text{t}}_{\mathfrak{gl}}^f$ are again labeled by the positive integers:

$$\text{Obj}(\mathring{\text{C}\ddot{\text{a}}\text{t}}_{\mathfrak{gl}}^f) = \{\mathbf{n}^f \mid n \in \mathbb{Z}_{\geq 0}\}.$$

For the space of morphisms we have

$$\mathbb{H}\text{om}(\mathbf{n}^f, \mathbf{m}^f) = (Z, w), \quad w \in \mathbb{C}[V_n \times \mathfrak{gl}_n \times Z \times \mathfrak{gl}_m \times V_m]^{GL_n \times GL_m}$$

here Z is an algebraic variety with an action of $GL_n \times GL_m$ and $V_n = \mathbb{C}^n, V_m = \mathbb{C}^m$ with the standard GL_n and GL_m actions.

The rest of definitions are identical to the constructions from the previous subsection, after we replace \mathfrak{gl}_n with $V_n \times \mathfrak{gl}_n$. For brevity we introduce the following shorthand notation:

$$\mathfrak{gl}_n^f := \mathfrak{gl}_n \times V_n.$$

In general, many definitions in our paper are parallel in framed and unframed cases, in the cases when the definitions are parallel we use \bullet notation to indicate that \bullet could be "f" or empty set.

First of all, note that the category $\mathbb{H}\text{om}(\mathbf{n}^\bullet, \mathbf{m}^\bullet)$ has a natural monoidal structure. Given $\mathcal{F} \in \text{Hom}((Z, w), (Z', w'))$ and $\mathcal{G} \in \text{Hom}((Z', w'), (Z'', w''))$ we define $\mathcal{F} \star \mathcal{G} \in \text{Hom}((Z, w), (Z'', w''))$

$$\mathcal{F} \star \mathcal{G} := \pi_*(\mathcal{F} \otimes \mathcal{G}), \quad \pi: \mathfrak{gl}_n \times Z \times Z' \times Z'' \times \mathfrak{gl}_m \rightarrow \mathfrak{gl}_n \times Z \times Z'' \times \mathfrak{gl}_m$$

4.4. Monoidal properties of $\mathring{\mathcal{C}\text{at}}_{\mathfrak{gl}}$. The objects of $\mathring{\mathcal{C}\text{at}}_{\mathfrak{gl}}$ are labeled by the positive numbers and it seems natural to use the addition for the monoidal structure on the level of objects. It is much harder to extend this monoidal structure to 1, 2, 3-morphisms. Nevertheless, we conjecture that it is possible to construct such extension. Below, we provide detailed motivation for the

Conjecture 4.4.1. *The three-category $\mathring{\mathcal{C}\text{at}}_{\mathfrak{gl}}$ can be enhanced to a monoidal three-category.*

Let us introduce the auxiliary object that facilitates the extension of monoidal structure to level of 1-morphisms. The object is the pair

$$\mathbb{I}_{n,k;n+k} = (T^*\text{GL}_{n+k}, W_{n,k;n+k}),$$

$\mathbb{C}[\mathfrak{gl}_{n+k} \times T^*\text{GL}_{n+k} \times \mathfrak{p}_{n,k}]^{\text{GL}_{n+k} \times P_{n,k}} \ni W_{n,k;n+k}(X, z, X') = \text{Tr}(X\mu_{\mathfrak{gl}_{n+k}}(z)) - \text{Tr}(X'\mu_{\mathfrak{p}_{n,k}})$, here $P_{n,k} \subset \text{GL}_{n+k}$ is the standard parabolic subgroup with the Levi group $\text{GL}_n \times \text{GL}_k$, respectively $\mathfrak{p}_{n,k} = \text{Lie}(P_{n,k})$ and $\mu_{\mathfrak{gl}_{n+k}}$ and $\mu_{\mathfrak{p}_{n,k}}$ are the moment maps for the left and right actions of the corresponding groups.

Given two objects $\mathbb{O}_1 \in \mathbb{H}\text{om}(\mathfrak{m}_1^\bullet, \mathfrak{n}^\bullet)$, $\mathbb{O}_2 \in \mathbb{H}\text{om}(\mathfrak{m}_2^\bullet, \mathfrak{k}^\bullet)$ we define the induced object

$$\mathbb{O}_1 \oplus \mathbb{O}_2 := \mathbb{O}_1 \times \mathbb{O}_2 \circ_{P_{n,k}} \mathbb{I}_{n,k;n+k} \in \mathbb{H}\text{om}(\mathfrak{m}_1^\bullet \times \mathfrak{m}_2^\bullet, (\mathfrak{n} + \mathfrak{k})^\bullet).$$

Thus defined operation extends to the level of two-morphisms. Moreover, the exchange relation from proposition 3.4.1 implies that this operation is monoidal with respect to the monoidal structure \star . In the case $m_1 = m_2$ the above operation was studied in [OR18e] where it is called induction functor and it was used to construct the braid realization discussed in the next section.

We expect that the object $\mathbb{I}_{n,k;n+k}$ will play an important role in the foam extension of our TQFT, this element is the element one needs for defining a value of the partition function on the configurations that contain three planes of defect intersecting along the line of defect. Also this element interacts well with NS5 defects discussed in the next section:

$$\mathbb{I}_{l|n} \times \mathbb{O} \circ_{P_{n,k}} \mathbb{I}_{n,k;n+k} = \mathbb{I}_{\text{id}}^l \times \mathbb{O} \circ_{P_{n,k}} \mathbb{I}_{l,k;l+k} \circ_{\text{GL}_{n+k}} \mathbb{I}_{l+k|n+k}, \quad l \leq n,$$

where $\mathbb{O} \in \mathbb{H}\text{om}(\mathfrak{m}^\bullet, \mathfrak{k}^\bullet)$ is any one-morphism and $\mathbb{I}_{l|n} \in \mathbb{H}\text{om}(\mathfrak{l}^\bullet, \mathfrak{n}^\bullet)$ is NS5 type one-morphism discussed in the next section.

5. DEFECTS AND KNOT INVARIANTS

In this section we explain how we interpret the results of [OR18e] in terms of three-categories $\mathring{\mathcal{C}\text{at}}_{\mathfrak{gl}}$ and $\mathring{\mathcal{C}\text{at}}_{\mathfrak{gl}}^f$. In particular, we make a connection with the theory of foams and provide an explanation for the Chern character construction. We also construct a partition functions \mathbf{Z}^f for the KRS theories the targets $\text{Hilb}_n(\mathbb{C}^2)$, $n \in \mathbb{Z}_{\geq 0}$ with NS5 type defects.

Recall the standard setup of the 3D topological field theory with defects. A 3D QTFT is characterized by its partition-function evaluation \mathbf{Z} . The partition function is an assignment:

$$\text{closed connected three-manifold } X \mapsto \mathbf{Z}(X) \in \mathbb{C},$$

$$\begin{aligned}
& \text{closed connected surface } S \mapsto Z(S) \in \text{Vect}, \\
& \text{three-manifold } X \text{ with boundary } \partial X = \bigcup_i S_i \mapsto Z(X) \in Z(\partial X) = \bigotimes_{i=1}^m Z(S_i), \\
& \text{closed connected curve } C \mapsto Z(C) \in \text{Cat}, \\
& \text{surface with boundary } S \mapsto Z(S) \in Z(\partial S) = \bigotimes_{i=1}^k Z(C_i), \\
& \text{point } p \mapsto Z(p) \in \overset{\bullet}{\text{Cat}}_{\text{gl}}, \\
& \text{interval } I = [b, e] \mapsto Z(I) \in \text{Hom}(Z(b), Z(e)).
\end{aligned}$$

This collection of data behaves naturally under the gluing operation. For example, suppose that a three-manifold X without a boundary is cut into two pieces over a surface S :

$$X = X_1 \cup X_2, \quad \partial(X_1) = S, \quad \partial(X_2) = S^\vee.$$

Then S and S^\vee have opposite orientations, hence $Z(S)$ and $Z(S^\vee)$ are dual vector spaces and the partition function $Z(X)$ is a pairing between their elements:

$$Z(X) = Z(X_1) \cdot Z(X_2),$$

More generally, the formalism of TQFT provides a method for computing values of $Z(Y)$ by cutting Y into pieces, evaluating Z on the pieces and pairing them in a standard way. More details can be found in [Lur08]. To define a 3D TQFT, we need to include into the domain of Z also manifolds with corners and work with more subtle setting of (∞, k) -categories. We postpone the discussion of such extension to our future publication [OR18a].

Often TQFTs may be defined not only on smooth manifolds but also on ‘smooth’ CW-complexes. In particular, a TQFT may have defects (coming from lower-dimensional cells). Topologically, defects are unions of embedded surfaces and curves. Surfaces and curves may intersect. The cuts must be transverse to the defects. All other properties of TQFT without defects are preserved.

The full categories $\overset{\bullet}{\text{Cat}}_{\text{gl}}$ will be used to construct 3D TQFT and we discuss the construction in the forthcoming publication [OR18a] where we construct the corresponding maps Z and Z^\dagger .

In this note we concentrate on \mathbb{R} -invariant part of the TQFT. That is from now on all defects are invariant with respect to shift along last coordinate in $S^2 \times \mathbb{R}$ and thus we can recover a configuration of defects from the two-dimensional cut which is an intersection with $S^2 \times \{0\}$.

We present S^2 as \mathbb{R}^2 compactified by a point ∞ at infinity and we can assume that the defects do not contain ∞ . The two-dimensional cut is $\mathbb{R}^2 \cup \infty$. Studying two-dimensional slice is equivalent to restricting ourselves to the defects of the form $C \times \mathbb{R}$ where C is a curve in \mathbb{R}^2 .

The surfaces of defect intersect our fixed \mathbb{R}^2 along the union of oriented curves. Suppose that the curves on the plane lies in an annulus and their union is a projection of the closure of a braid. By assigning a sign to each intersection, we obtain an interpretation of the union of curves as a projection of a link in \mathbb{R}^3 presented as a closure $L(\beta)$ of a braid $\beta \in \mathfrak{B}\mathfrak{r}_n$. Denote the plane with defect \mathbb{R}_β^2 . Our TQFT provides an isotopy invariant of




FIGURE 1. \mathbb{R}_β^2 for $\beta = \sigma_1^3$.

$L(\beta)$, namely, the vector space $Z^f(\mathbb{R}_\beta^2)$, assigned to \mathbb{R}_β^2 , and we show that this invariant coincides with previously defined invariant for [OR18e].

Now let us give details of the \mathbb{R}^2 -sliced TQFT. A small neighborhood of the plane contains two-dimensional surfaces of defect that are products of the defect curves in \mathbb{R}^2 with an interval. The surfaces of defect divide \mathbb{R}^3 into connected components and each connected component has an integer marking. We choose the markings in such a way that if we move along the (oriented) intersection of a defect surface with \mathbb{R}^2 and the marking on the left is k , then the marking on the right is $k + 1$.

We assume that the curves of intersection are compact, thus their union is contained in a large disc. The marking of the disc complement (‘the infinite marking’) determines all other markings.

In this paper we only consider the theories with the infinite marking equal to 0. Hence, if the intersection of the surface defect and \mathbb{R}^2 is the braid closure then all markings are positive. Thus the picture of the braid β determines the marking of our theory uniquely. Slightly abusing notation, we denote such data as \mathbb{R}_β^2 . The figure is a slice \mathbb{R}_β^2 for $\beta = \sigma_1^3$. The closure of $L(\beta)$ is a trefoil and we explain below how we can compute the homology of this knot.

5.1. Values on points and defect trivial intervals. There is a canonical way to upgrade the marking of \mathbb{R}_β^2 to a categorical marking. Denote by \mathfrak{p}_n a point lying inside a region marked by n . To a pair of points $(\mathfrak{p}_{n_1}, \mathfrak{p}_{n_2})$ we assign a 2-category

$$(5.1) \quad Z^\bullet(\mathfrak{p}_{n_1}, \mathfrak{p}_{n_2}) = \mathbb{H}\text{om}(\mathfrak{n}_1^\bullet, \mathfrak{n}_2^\bullet).$$

Recall that an object of this 2-category is a pair (Z, W) , where Z is a variety with a $(\text{GL}(n_1) \times \text{GL}(n_2))$ -action and $W \in \mathbb{C}[\mathfrak{gl}_{n_1} \times \mathfrak{gl}_{n_2} \times Z]^{\text{GL}(n_1) \times \text{GL}(n_2)}$.

In accordance with the assignment (5.1), to an interval I connecting \mathfrak{p}_{n_1} and \mathfrak{p}_{n_2} (and possibly crossing defect surfaces) we assign an object $Z^\bullet(I)$ of $\mathbb{H}\text{om}(\mathfrak{n}_1^\bullet, \mathfrak{n}_2^\bullet)$, so that if I is the result of gluing together the intervals I_1 and I_2 over the common middle point, then the object of I is the composition:

$$Z^\bullet(I) = Z^\bullet(I_2) \circ Z^\bullet(I_1).$$

This relation implies that if the interval I lies within a single region marked by \mathfrak{p}_n , then the corresponding object $Z^\bullet(I)$ is the identity with respect to the monoidal structure (*i.e.*, the composition) of the category $\mathbb{H}\text{om}(\mathfrak{n}^\bullet, \mathfrak{n}^\bullet)$.

Theorem 4.3.1 states that the following pair is the identity object of the ‘unframed’ 2-category $\mathbb{H}\text{om}(\mathfrak{n}, \mathfrak{n})$:

$$\mathbb{L}_{\text{id}}^{n,0} = \mathbb{L}_{\text{id}}^n = \left(T^*\text{GL}_n, W_{\text{id}} := \text{Tr} \phi(X - \text{Ad}_g(X')) \right)$$

in which $(g, \phi) \in \text{GL}(n) \times \mathfrak{gl}_n \cong T^*\text{GL}(n)$, while $(X, X') \in \mathfrak{gl}_n \times \mathfrak{gl}_n$, and the action of $\text{GL}(n) \times \text{GL}(n)$ on the total variety $\mathfrak{gl}_n \times \mathfrak{gl}_n \times T^*\text{GL}(n)$ is given by the following formula:

$$(a, b) \cdot (X, X', g, \phi) = (\text{Ad}_a X, \text{Ad}_b X', agb^{-1}, \text{Ad}_a \phi)$$

The identity object in the framed 2-category $\mathbb{H}\text{om}(\mathfrak{n}^f, \mathfrak{n}^f)$ is the case $r = 1$ of

$$\mathbb{L}_{\text{id}}^{n,r} = \left(T^*\text{GL}_n \times \text{Hom}(V_n, \mathbb{C}^r), W_{\text{id}}^f := W_{\text{id}}(X, g, \phi, X') + \text{Tr}(w \cdot (v - gv')) \right),$$

where $(X, v), (X, v') \in \mathfrak{gl}_n \times \text{Hom}(\mathbb{C}^r, V_n)$, while $(g, \phi, w) \in T^*\text{GL}(n) \times \text{Hom}(V_n, \mathbb{C}^r)$ and the action of $\text{GL}(n) \times \text{GL}(n)$ on framing related variables is

$$(a, b) \cdot (v, v', w) = (av, bv', aw).$$

Proof of theorem 4.3.1. We need to define the left and right unitizing 1-morphisms

$$\begin{aligned} l(\mathcal{X}) &: \mathcal{X} \rightarrow \mathbb{L}_{\text{id}}^{n;r} \circ \mathcal{X}, \quad \mathcal{X} \in \mathbb{H}\text{om}(\tilde{A}_0(n, r), \tilde{A}_0(n', r')), \\ r(\mathcal{X}) &: \mathcal{X} \rightarrow \mathcal{X} \circ \mathbb{L}_{\text{id}}^{n;r}, \quad \mathcal{X} \in \mathbb{H}\text{om}(\tilde{A}_0(n', r'), \tilde{A}_0(n, r)). \end{aligned}$$

We take the left unitizer to be the tensor product of matrix factorizations

$$l(\mathcal{X}) := \mathbf{1}_{\mathcal{X}} \circ \mathcal{KN}, \quad \mathcal{KN} = [\phi, X - \text{Ad}_g(X')] \otimes [w, v - gv'],$$

where we use the notation from the definition of the unit object as well as the standard notations for the Koszul matrix factorization. The right unitizer is defined by the identical formula.

The matrix factorization $\mathcal{KN}|_{g=1}$ is the matrix factorization that yields the Knörrer’s equivalence of categories [Kno87]. The composition \circ in the product $\mathbb{O} \circ \mathbb{L}_{\text{id}}$ involves taking quotient by the GL_n -action. Since the left and right actions on GL_n on $T^*\text{GL}_n$ is free, we can trade the GL_n -quotient for the gauge fixing $g = 1$. Thus we are in the setting of the Knörrer periodicity. Respectively, the inverse to the Knörrer equivalence is composition of KN^* and the duality functor.

The only non-trivial structural two-morphisms that involves the unit object are

$$\begin{array}{ccc}
\mathcal{X} & \xrightarrow{\mathcal{F}} & \mathcal{X}' \\
\downarrow l(\mathcal{X}) \swarrow l(\mathcal{F}) & & \downarrow l(\mathcal{X}') \\
\mathbb{L} \circ \mathcal{X} & \xrightarrow{1_{\mathbb{L}} \circ \mathcal{F}} & \mathbb{L} \circ \mathcal{X}'
\end{array}
\quad , \quad
\begin{array}{ccc}
\mathcal{X} & \xrightarrow{\mathcal{F}} & \mathcal{X}' \\
\downarrow r(\mathcal{X}) \swarrow r(\mathcal{F}) & & \downarrow r(\mathcal{X}') \\
\mathcal{X} \circ \mathbb{L} & \xrightarrow{\mathcal{F} \circ 1_{\mathbb{L}}} & \mathcal{X}' \circ \mathbb{L}
\end{array}$$

which are defined by

$$l(\mathcal{F}) = R(\mathcal{KN}, \mathcal{F}), \quad r(\mathcal{F}) = R(\mathcal{F}, \mathcal{F}).$$

Thus these structural two-morphisms are compositions of the base-change natural transformations. The relevant conditions for the structural morphism related to the unit object of monoidal two-category are listed on the page 224 of [KV94]. Since the structural two-morphisms in our setting are of geometric origin one prove the conditions by the pasting method used in theorem \square

It follows from the axioms of the monoidal two-category (see for example [BN96]) that the category of endomorphisms of unit object is a universal Drinfeld center, in the following sense

Corollary 5.1.1. *The monoidal category $\text{Hom}(\mathbb{L}_{\text{id}}^{n;r}, \mathbb{L}_{\text{id}}^{n;r})$ is a braided monoidal category. Moreover for any $\mathcal{X} \in \mathbb{H}\text{om}(\tilde{A}_0(n;r), \tilde{A}_0(n';r'))$ there is a monoidal functor to the Drinfeld center of $\text{Hom}(\mathcal{X}, \mathcal{X})$:*

$$\text{HC}[\mathcal{X}] : \text{Hom}(\mathbb{L}_{\text{id}}^{n;r}, \mathbb{L}_{\text{id}}^{n;r}) \rightarrow Z(\text{Hom}(\mathcal{X}, \mathcal{X}), \star).$$

In the special case of $n' = 0$ and $r = r' = 1$ and special choice of \mathcal{X} this functor was studied in [OR18d]. The exact connection is explained in the next subsections. In this work we also define a right adjoint functor to the HC. We expect that the adjoint functor exists in more general setting:

Conjecture 5.1.2. *For any $\mathcal{X} \in \mathbb{H}\text{om}(\tilde{A}_0(n;r), \tilde{A}_0(n';r'))$ there is a functor*

$$\text{CH}[\mathcal{X}] : \text{Hom}(\mathcal{X}, \mathcal{X}) \rightarrow \text{Hom}(\mathbb{L}_{\text{id}}^{n;r}, \mathbb{L}_{\text{id}}^{n;r})$$

that is the right adjoint to $\text{HC}[\mathcal{X}]$ and is a categorical trace

$$\text{CH}[\mathcal{X}](\mathcal{F} \star \mathcal{G}) = \text{CH}[\mathcal{X}](\mathcal{G} \star \mathcal{F}).$$

5.2. Value on the intervals intersecting defects. Now assume that I is a small interval intersecting a curve of defect in a smooth point. Let us assume that the curve of defect separates regions with marking k and l . Denote by (ϕ_{kl}, ϕ_{lk}) the coordinates on $\text{T}^*\text{Hom}(V_k, V_l)$, where $\phi_{kl} \in \text{Hom}(V_k, V_l)$ and $\phi_{lk} \in \text{Hom}(V_l, V_k)$. Also denote by v_l and v_k the coordinates on V_l and V_k and denote by X_l and X_k the coordinates on \mathfrak{gl}_l and \mathfrak{gl}_k . Now the object assigned to the interval \vec{I} in the unframed category is a pair

$$Z(\vec{I}) = \mathbb{I}_{k|l} = (\text{T}^*\text{Hom}(V_k, V_l), \quad W_{k,l} := \text{Tr}(X_k \phi_{kl} \phi_{lk}) - \text{Tr}(X_l \phi_{lk} \phi_{kl})).$$

In the context of the section 3.5 $\mathbb{I}_{k|l}$ corresponds to the NS5 defect of charge $k - l$ with no framing i.e. $r = 0$.

In case of framed 2-categories, if the interval starts at k and ends in l and the shortest path from the head of the vector \vec{I} to the head of the vector of direction of the defect goes clockwise, then we choose

$$Z^f(\vec{I}) = \mathbb{I}_{k|l}^f = (\mathrm{T}^*\mathrm{Hom}(V_k, V_l) \times V_k^*, \quad W_{k,l}^f := \mathrm{Tr}(X_k \phi_{kl} \phi_{lk}) - \mathrm{Tr}(X_l \phi_{lk} \phi_{kl}) + \mathrm{Tr}(\psi(v_k - \phi_{kl} v_l)).$$

In case of opposite orientation we set

$$Z^f(\vec{I}) = (\mathrm{T}^*\mathrm{Hom}(V_k, V_l) \times V_l^*, \quad W_{k,l}^f := \mathrm{Tr}(X_k \phi_{kl} \phi_{lk}) - \mathrm{Tr}(X_l \phi_{lk} \phi_{kl}) + \mathrm{Tr}(\psi(v_l - \phi_{lk} v_k)).$$

To match the objects $\mathbb{I}_{k|l}^f$ matches with the description of NS5 defects of charge $k-l$ with r from section 3.5.5 we need to set $\varphi^{\mathrm{fr}} = 1$ and $\psi = \psi^{\mathrm{fr}}$. The element $Z^\bullet(\vec{I})$ is an element of the two-category $\mathbb{H}\mathrm{om}(\mathbf{k}^\bullet, \mathbf{l}^\bullet)$ thus the composition construction allow us to interpret an element $Z^\bullet(\vec{I})$ as morphism from $\mathbb{H}\mathrm{om}(\mathbf{k}^\bullet, \mathbf{0}^\bullet)$ to $\mathbb{H}\mathrm{om}(\mathbf{l}^\bullet, \mathbf{0}^\bullet)$.

Let us denote the intervals as above by $\vec{I}_{k\uparrow l}$ and $\vec{I}_{k\downarrow l}$. More generally we denote by

$$\vec{I}_{k_1\uparrow k_2\uparrow \dots \uparrow k_l}, \quad \vec{I}_{k_1\downarrow k_2\downarrow \dots \downarrow k_l}$$

the interval that connects the connected components with the labels k_1 and k_l and traverses the domains with the labels k_2, \dots, k_l in between with the orientation of the intersections as indicated by the arrows. We also allow a mixture of down/up orientations of the intersections.

According to the definition of our TQFT we have:

$$(5.2) \quad Z(\vec{I}_{k_1\uparrow k_2 \dots \uparrow k_l}) = Z(\vec{I}_{k_1\uparrow k_2}) \circ Z(\vec{I}_{k_2\uparrow k_3}) \circ \dots \circ Z(\vec{I}_{k_{l-1}\uparrow k_l}).$$

The GIT quotient in the definition of the composition can be made explicit in many important cases:

Proposition 5.2.1. *For any $n \geq 0$ we have:*

$$\begin{aligned} Z^f(\vec{I}_{0\uparrow 1 \dots n}) &= (\mathrm{T}^*\mathrm{Fl}_n \times V_n, w), & Z^f(\vec{I}_{0\downarrow 1 \dots n}) &= (\mathrm{T}^*\mathrm{Fl}_n, w), \\ Z(\vec{I}_{0|1 \dots n}) &= (\mathrm{T}^*\mathrm{Fl}_n, w), & w &= \mu \cdot X \in \mathbb{C}[\mathfrak{gl}_n \times \mathrm{T}^*\mathrm{Fl}_n]^{\mathrm{GL}_n}, \end{aligned}$$

where $\mu : \mathrm{T}^*\mathrm{Fl}_n \rightarrow \mathfrak{gl}_n^*$ is the moment map and X are the coordinates on \mathfrak{gl}_n .

Proof. Let us first prove the last equation, the other equations are analogous and it will be indicated at the end of the proof how one needs to modify the proof to get the first two formulas. We proceed by induction on n . Thus we need to compute the composition:

$$Z(\vec{I}_{0|1 \dots |n-1}) \circ Z(\vec{I}_{n-1|n}).$$

It is convenient to think of $\mathrm{T}^*\mathrm{Fl}_n$ as B_n -quotient because the trace map gives a natural pairing on \mathfrak{gl}_n thus we can think of μ as a map $\mathrm{T}^*\mathrm{Fl}_n \rightarrow \mathfrak{gl}_n$:

$$\mathrm{T}^*\mathrm{Fl}_n = \mathrm{GL}_n \times \mathfrak{n}_n / B_n, \quad \mu(g, Y) = \mathrm{Ad}_g Y,$$

where g and Y are the coordinates on GL_n and \mathfrak{n}_n .

In these notations the composition in question is the pair of the GIT quotient space and a potential:

$$\mathrm{T}^*\mathrm{Fl}_{n-1} \times \mathrm{T}^*\mathrm{Hom}(V_{n-1}, V_n) / {}_+ \mathrm{GL}_{n-1}, w_{n-1,n} = \mathrm{Tr}(X' \mathrm{Ad}_g Y') + \mathrm{Tr}(X' \phi \psi) - \mathrm{Tr}(X'' \psi \phi),$$

where $X' \in \mathfrak{gl}_{n-1}$, $X'' \in \mathfrak{gl}_n$, g, Y are the coordinates along $T^*\mathrm{Fl}_{n-1}$ and $\psi \in \mathrm{Hom}(V_n, V_{n-1})$, $\phi \in \mathrm{Hom}(V_{n-1}, V_n)$ are the coordinates along $T^*\mathrm{Hom}(V_{n-1}, V_n)$.

The GIT quotient in last formula could be made very explicit, we choose to describe the quotient by constructing explicit charts in the quotients. Then we show that in each chart we can apply the Knorrer periodicity to simplify the potential.

The GIT stable locus consists of points where ϕ is injective. Thus we can assume that there is k such that

$$\det(\phi_{\hat{k}}) \neq 0,$$

where $\phi_{\hat{k}}$ is ϕ with k -th row removed. Let us denote the locus where the last inequality holds by U_k . It is clear that the quotient is covered by the charts U_k/GL_{n-1} and we can analyze the potential in each chart.

To simplify notations let us consider the case $k = n$. The natural slice to the GL_{n-1} -action is the closed subset of elements constrained by:

$$\phi_{ij} = \delta_{i,j}, \quad 1 \leq i \leq j \leq n-1.$$

Let us also denote the last row of ϕ by v and the matrix of the first $n-1$ columns of ψ by $\tilde{\psi}$ and the last row of ψ by ψ' . Then the potential $w_{n-1,n}$ becomes:

$$(5.3) \quad \mathrm{Tr}(X' \mathrm{Ad}_g Y') + \mathrm{Tr}(X' \tilde{\psi}) + \mathrm{Tr}(v X' \psi') - \mathrm{Tr}(X'' \phi \psi).$$

The sum of the first three terms is quadratic and we can apply Knorrer reduction. The reduction forces the following vanishing of the coordinates:

$$X' = 0, \quad \mathrm{Ad}_g Y' + \tilde{\psi} + \psi' v = 0.$$

Thus the new coordinates on the Knorrer reduced space are X'', Y', v, ψ' and in these coordinates we have:

$$\phi \psi = \begin{bmatrix} -\psi' v - \mathrm{Ad}_g Y' & \psi' \\ -v \psi' v - v \mathrm{Ad}_g Y' & v \psi' \end{bmatrix}$$

Thus a direct computation shows that the last term of (5.3) is equal to

$$\mathrm{Tr}(X'' \mathrm{Ad}_h Y), \quad \text{with } Y = \begin{bmatrix} Y & g^{-1} \psi' \\ 0 & 0 \end{bmatrix}, \quad h = \begin{bmatrix} g & 0 \\ v g & 1 \end{bmatrix}.$$

Hence we proved the last formula in the chart U_n and the computations in other charts are analogous. The argument in the framed case is basically the same. \square

The last proposition computes the object \mathcal{F}_n from the introduction since $Z^f(\vec{I}_{0 \uparrow 1 \uparrow \dots n}) = \underbrace{\hat{\square}_1 \cdots \hat{\square}_1}_n \mathcal{O}$.

5.3. The categories of closed curves. The choice of defect-related objects $Z^\bullet(\vec{I})$ determines categories assigned to closed curves: a curve C is presented as a gluing of two intervals, then its category $Z^\bullet(C)$ must be the category of morphisms between their objects. Two curves are of special importance for our braid-related constructions.

The first type is a curve that does not intersect any defects. So the curve is a circle that lies inside of the connected component with the marking n . We denote such circle S_n^1 . To a point $p \in S_n^1$ we assign two-category $\mathbb{H}\text{om}(\mathbf{n}^\bullet, \mathbf{0}^\bullet)$. For brevity, we start using notation

$$[\mathbf{n}^\bullet, \mathbf{m}^\bullet] := \mathbb{H}\text{om}(\mathbf{n}^\bullet, \mathbf{m}^\bullet),$$

for the corresponding two-category.

The interval I connecting \mathbf{p}_n to itself get assigned the identity $\mathbf{Z}^\bullet(I) = \mathbb{L}_{\text{id}}^n \in [n, n]$. Since S_n^1 is a result of gluing two such intervals, its category is the category of endomorphisms of the interval object:

$$\mathbf{Z}^\bullet(S_n^1) = \text{Hom}(\mathbb{L}_{\text{id}}^n, \mathbb{L}_{\text{id}}^n).$$

In the introduction explained the matrix factorization realization of the braid group (1.5) and the Drinfeld center construction of the corresponding monoidal category (1.7) from [OR18d]. To relate these constructions to corollary 5.1.1 and motivate the conjecture 5.1.2 we need to study the Drinfeld category from [OR18d]:

$$(5.4) \quad \text{MF}_{\text{Dr}}^\bullet = \text{MF}_{G_n}((\mathfrak{g}^\bullet \times \mathfrak{g}^\bullet \times G)^{\text{st}}, W_{\text{Dr}}^\bullet),$$

$$W_{\text{Dr}}(X, Y, g) = \text{Tr} X(\text{Ad}_g(Y) - Y), \quad W_{\text{Dr}}^f(X, v, Y, u, g) = W_{\text{Dr}}(X, Y, g) + u^* \cdot v - u^* \cdot gv.$$

Proposition 5.3.1. *For $\bullet = \emptyset$, the categories $\text{MF}_{\text{Dr}}^\bullet$ are equivalent to the categories of endomorphisms of the identity objects \mathbb{L}_{id}^n :*

$$\text{MF}_{\text{Dr}}^\bullet = \text{Hom}(\mathbb{L}_{\text{id}}^n, \mathbb{L}_{\text{id}}^n)$$

Proof. We consider only the case $\bullet = \emptyset$ the other case is analogous. The statement follows from the Knorrer periodicity [Kno87]. Indeed, by our definition we have

$$\text{Hom}(\mathbb{L}_{\text{id}}^n, \mathbb{L}_{\text{id}}^n) = \text{MF}_{\text{GL}_n}(\text{T}^*\text{GL}_n \times \mathfrak{gl}_n \times \mathfrak{gl}_n \times \text{T}^*\text{GL}_n/\text{GL}_n, W),$$

$$W = \text{Tr} \phi(X - \text{Ad}_g(X')) - \text{Tr} \phi(X' - \text{Ad}_{g'}(X))$$

where (ϕ, g) and (ϕ', g') are the coordinates on two copies of T^*GL_n and X, X' are the coordinates on two copies of \mathfrak{gl}_n . By setting $g' = 1$ we take slice to the GL_n -orbits. On the slice the second term in the formula for W becomes $\text{Tr} \phi'(X - X')$. Thus the Knorrer periodicity implies that the restriction on the locus $X = X', \phi' = 0$ is the equivalence of the corresponding categories of matrix factorizations. \square

We can linearize the potential W_{Dr}^\bullet by introducing a new variable $U = Yg^{-1}$:

$$W_{\text{lin}}(X, U, g) = \text{Tr}(X[g, U]), \quad W_{\text{lin}}^f(X, v, U, u, g) = W_{\text{lin}}(X, U, g) + u^* \cdot v - u^* \cdot gv.$$

The group G naturally embeds inside its lie algebra \mathfrak{g} , $j_G: G \rightarrow \mathfrak{g}$. Induced by this embedding we have the localization functor:

$$\text{loc}^\bullet: \text{MF}_{\text{Dr}}^\bullet \rightarrow \underline{\text{MF}}_{\text{Dr}}^\bullet = \text{MF}_G((\mathfrak{g}^\bullet \times \mathfrak{g}^\bullet \times \mathfrak{g})^{\text{st}}, W_{\text{lin}}^\bullet).$$

It turns out that in the framed case the localization functor is an isomorphism:

Proposition 5.3.2. [OR18d] *The localization functor loc^f is an isomorphism.*




FIGURE 2. Plane $\mathbb{R}_{\sigma_1}^2$ cut by $\mathbb{R}_{0,1,2,1,0}$

Since the potential W^{lin} is linear along the last copy of \mathfrak{g} , the Koszul duality (see for example [AK15] or [OR18d]) provides an equivalence:

$$(5.5) \quad \text{KSZ: } \underline{\text{MF}}_{\text{Dr}}^f \longrightarrow \text{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2)).$$

Thus we have completed proof of theorem 1.1.2.

The second type of a closed curve is the line that intersects our braid transversally. The line goes through the regions with the marks $0, 1, \dots, n-1, n, n-1, \dots, 0$. Figure 2 gives an example. We denote such a line compactified by a point at infinity as $S_{0\uparrow 1\uparrow \dots \uparrow n\downarrow n-1\downarrow \dots \downarrow 0}^1$. The value \mathbf{Z}^\bullet follows immediately from the proposition 5.2.1:

$$(5.6) \quad \mathbf{Z}^f(S_{0\uparrow 1\uparrow \dots \uparrow n\downarrow n-1\downarrow \dots \downarrow 0}^\bullet) = \text{MF}_{G_n}^{\text{st}}(\mathfrak{gl}_n^\bullet \times \text{T}^*\text{Fl}_n \times \text{T}^*\text{Fl}_n, w_1 - w_2),$$

where "st" indicate that we restrict to the GIT stable locus of the corresponding space.

The categories (5.6) are closely related to the main categories of [OR18e]. Recall that in [OR18e] we worked with $G_n \times B \times B$ -equivariant categories of matrix factorizations on the space


$$\mathcal{X}^\bullet := \mathfrak{gl}_n^\bullet \times (G_n \times \mathfrak{n}_n) \times (G_n \times \mathfrak{n}_n)$$

with the equivariant structure preserving the potential

$$W(X, g_1, Y_1, g_2, Y_2) = \text{Tr } X(\text{Ad}_{g_1} X_1 - \text{Ad}_{g_2} X_2),$$

where X is the coordinate in \mathfrak{gl}_n and g_i, Y_i are the coordinates in G and \mathfrak{n} .

5.4. Values on discs. As a final step of our construction we need to discuss the values of TQFT on discs. The first type of disc is the disc D_\emptyset that bounds S_n^1 and does not contain tautological defect point. The category $\mathbf{Z}^\bullet(S^2) = \text{Hom}(L_{\text{id}}^n, L_{\text{id}}^n)$ is monoidal and $\mathbf{Z}^\bullet(D_\emptyset)$


 FIGURE 3. Decomposing \mathbb{R}_β^2 on two half-planes

represents the identity object in it. Hence we set

$$\mathbf{Z}^f(D_\emptyset) := \mathcal{O} \in \mathbf{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2)).$$

If the disc contains the point of tautological defect then we set

$$\mathbf{Z}^f(D_{\text{taut}}) := \Lambda \bullet \mathcal{B} \in \text{Coh}(\text{Hilb}_n(\mathbb{C}^2)),$$

where \mathcal{B} is the tautological vector bundle.

The other important type of a disc is a half-plane H bordered by the line $S_{0\uparrow 1\uparrow \dots \uparrow n\downarrow n-1\downarrow \dots \downarrow 0}$. Its object $\mathbf{Z}^\bullet(H)$ lies in the monoidal category

$$(5.7) \quad \mathbf{Z}^\bullet(S_{0\uparrow 1\uparrow \dots \uparrow n\downarrow n-1\downarrow \dots \downarrow 0}) = \text{End}(\vec{I}_{0\uparrow 1\uparrow \dots \uparrow n-1\uparrow n}).$$

and it depends on the configuration of defects inside H . Denote by H_1 the simplest configuration which is the collection of non-intersecting curves connecting the points of the same type as in the right half-plane in Figure 3. In this situation \mathbf{Z}^\bullet is the identity object:

$$\mathbf{Z}^\bullet(H_1) = \mathcal{C}_1 \in \text{MF}_n^\bullet.$$

More generally, denote by H_β the half-plane containing a braid β as in the left of Figure 3. The value of $\mathbf{Z}(H_\beta)$ for more complicated configurations of defects can be computed by using the monoidal structure of the category (5.7) through cutting H_β into the union:

$$H_\beta = \bigcup_k S_{\sigma_{j_k}^{\epsilon_k}},$$

where $\beta = \sigma_{j_1}^{\epsilon_1} \dots \sigma_{j_l}^{\epsilon_l}$ and $S_{\sigma_{j_k}^{\epsilon_k}}$ is the disc with the boundary $S_{0\uparrow 1\uparrow \dots \uparrow n\downarrow n-1\downarrow \dots \downarrow 0}$ and defects inside the strip form an elementary braid on the j_k -th and $(j_k + 1)$ -st stands, see the figure below for the case $\beta = \sigma_1^3$. Since

$$\mathbf{Z}^\bullet(H_\beta) = \mathbf{Z}^\bullet(S_{\sigma_{j_1}^{\epsilon_1}}) \star \cdots \star \mathbf{Z}^\bullet(S_{\sigma_{j_l}^{\epsilon_l}}),$$

it is enough to define $\mathbf{Z}^\bullet(S_{\sigma_k^{\pm 1}}) \in \mathbf{MF}_n^\bullet$, as in [OR18e]:

$$\mathbf{Z}^\bullet(S_{\sigma_k^{\pm 1}}) := \mathcal{C}_\pm^{(k)} \in \mathbf{MF}^\bullet,$$

It is shown in [OR18e] that the element $Z(H_\beta)$ only depends on the braid β but not on the braid presentation, thus our disc assignment indeed is a well-defined partition function of TQFT.

Finally, we define the value of Z^f on the half-plane H_1^{taut} containing the unit braid and the tautological point defect as

$$Z^f(H_1^{\text{taut}}) := \mathcal{C}_1 \otimes \Lambda^\bullet \mathcal{B}.$$

We leave the following statement as conjecture and will provide a proof in the forthcoming publication.

Conjecture 5.4.1. *The above assignments of the values of Z are the part of the data of a well-defined 3D TQFT.*

5.5. **Value on \mathbb{R}_β^2 .** There are two ways to cut the plane with a closed braid defect \mathbb{R}_β^2 into two pieces. As a result, the TQFT formalism implies two presentations of the corresponding vector space $Z^f(\mathbb{R}_\beta^2)$ as the space of morphisms between two objects in the category of the cutting line.

The first cut splits \mathbb{R}_β^2 in two half-planes H_1^{taut} and H_β , and the corresponding presentation is

$$Z^f(\mathbb{R}_\beta^2) = \text{Hom}(Z^f(H_1^{\text{taut}}), Z^f(H_\beta)) = (\text{Hom}(\mathcal{C}_\beta, \mathcal{C}_1) \otimes \Lambda^\bullet \mathcal{B})^{B^2 \times G}.$$

The vector space $Z^f(\mathbb{R}_\beta^2)$ is triply-graded and the main result of [OR18e] could be restated as

Theorem 5.5.1. *The triply-graded vector space $Z^f(\mathbb{R}_\beta^2)$ is an isotopy invariant of the closure of the braid β after a special shift of the grading.*

The second cut (see Figure 4) splits \mathbb{R}_β^2 into the inner disc D_{taut} (containing the tautological bundle defect) and its complement D_β^∞ which contains the closed braid defect. The cut goes over a circle, that lies in the region marked by n and does not intersect defect lines, hence its category is

$$(5.8) \quad Z^f(S_n^1) = \mathbf{MF}_{\text{Dr}}^f \cong \mathbf{D}^{\text{per}}(\text{Hilb}_n(\mathbb{C}^2))$$

of (5.4) and (5.5). The object of D_{taut} is just the defect bundle: $Z^f(D_{\text{taut}}) = \Lambda^\bullet \mathcal{B}$. The object of D_β^∞ is determined by the categorical Chern character functor

$$\text{CH}_{\text{loc}}^{\text{fs}} : \mathbf{MF}_n^f \rightarrow Z^f(S_n^1),$$

that is, $Z^f(D_\beta^\infty) = \text{CH}^{\text{fs}}(Z^f(H_\beta))$. Thus we get the second presentation of the vector space $Z^f(\mathbb{R}_\beta^2)$ as the Ext space between two complexes of sheaves within the derived category of 2-periodic sheaves on the Hilbert scheme $\text{Hilb}_n(\mathbb{C}^2)$:

$$(5.9) \quad Z^f(\mathbb{R}_\beta^2) = \mathcal{H}om(\text{CH}^{\text{fs}}(Z^f(H_\beta)), \Lambda^\bullet \mathcal{B})$$




FIGURE 4. Decomposition of S_β^2 on two discs.

The isomorphism between the vector spaces (5.8) and (5.9) is one of the key properties of the functor $\text{CH}_{\text{loc}}^{\text{fs}}$ (one may call it a simple case of the categorical Riemann-Roch formula):

Theorem 5.5.2. [OR18d] *For any $\mathcal{C} \in \text{MF}_n$ we have:*

$$\mathcal{H}om(\Lambda^\bullet \mathcal{B}, \text{CH}_{\text{loc}}^{\text{fs}}(\mathcal{C}))^G = \mathcal{H}om(\mathcal{C}_1 \otimes \Lambda^\bullet \mathcal{B}, \mathcal{C})^{B^2 \times G}.$$

Thus we constructed a complex of sheaves $S_\beta := \text{CH}_{\text{loc}}^{\text{fs}}(\mathcal{C}_\beta)$ such that its global sections are the knot homology. In the language of TQFT the categorical Riemann-Roch and our main theorem is just a gluing property of TQFT: the picture 3 and picture 4 present two different ways of computing the same partition sum $Z^{\text{f}}(\mathbb{R}_\beta^2)$.

6. CATEGORIFICATION OF TRACES

In this section we expand on the table from the introduction. The first column of the table is a summary of the previous sections. The second column is a physical interpretation of the results of [OR20] where a fully-faithful functor from the category of Soergel bimodules to the category of stable matrix factorizations was constructed. The last column is closely related to the work [AN16], [NT17] and forthcoming preprint of the authors [OR22].

6.1. Soergel bimodules. A slight modification of the computation in the proof of proposition 5.2.1 shows that:

$$\tilde{\mathcal{F}}_n = \blacksquare_1 \underbrace{\hat{\square}_1 \cdots \hat{\square}_1}_{n-1} \mathcal{O}, \quad \tilde{\mathcal{F}}_n = ((\tilde{\mathfrak{gl}}_n \times V_n)^{\text{st}}, \tilde{W}_n),$$

where $\tilde{\mathfrak{gl}}_n = \text{GL}_n \times \mathfrak{b}/B$ is the Grothendieck-Springer alteration with the moment map $\mu : \tilde{\mathfrak{gl}}_n \rightarrow \mathfrak{gl}_n$ and $\tilde{W}_n(X, z) = \text{Tr}(X\mu(z))$. The stability condition is given by requiring $\mathbb{C}[\mu(z)]v = V_n$.

Indeed, the computation in the proposition 5.2.1 shows that the composition results into the object the B -equivariant object $(\mathfrak{gl}_n \times \mathfrak{gl}_n \times \mathrm{GL}_n \times \mathfrak{n} \times V_n, W'')$ where $W''(Y', g, X, Y, v) = \mathrm{Tr}(Y'X) + \mathrm{Tr}(X\mathrm{Ad}_g Y)$. The last potential, has a quadratic term:

$$W''(Y', g, X, Y, v) = \mathrm{Tr}((\mathrm{Ad}_g^{-1}X)_{--}(Y + \mathrm{Ad}_g^{-1}Y')) + \mathrm{Tr}((\mathrm{Ad}_g^{-1}X)_+ \mathrm{Ad}_g^{-1}Y).$$

where the sub-indices $--$ and $+$ stand for the strictly lower-triangular and the upper-triangular parts of the matrices. Thus the Knörrer reduced object has potential $\mathrm{Tr}(X' \mathrm{Ad}_g^{-1}Y)$, $X' = (\mathrm{Ad}_g^{-1}X)_+$ and it is isomorphic to the object $\tilde{\mathcal{F}}_n$.

The algebra of the MOY graphs [MOY98] has an algebraic incarnation in terms of the algebra \mathfrak{Br}_n^b of flat braids. The last algebra can be realized by means of endomorphisms of $\tilde{\mathcal{F}}_n$

Proposition 6.1.1. [OR20] *For any n there is a monoidal functor*

$$\Phi_n^b : \mathfrak{Br}_n \rightarrow \mathrm{Hom}(\tilde{\mathcal{F}}_n, \tilde{\mathcal{F}}_n).$$

To connect the previous discussion with the second column of the table from the introduction we examine the picture (1.11) for case when the region \mathbb{B} contains the circular defect of type $\mathrm{NS}5^{(n)}$. In this case, the ray I_{rad} that starts at the center of the circle from the picture (1.11) intersects the domains with labels $0, n, n-1, \dots, 1, 0$ and the corresponding category of morphisms $Z^f(I_{rad}) = \hat{\blacksquare}_{-n}(\tilde{\mathcal{F}})$ is computed in

Proposition 6.1.2. *For any n we have*

$$\hat{\blacksquare}_{-n}(\tilde{\mathcal{F}}_n) = \mathrm{Mod}(\mathbb{C}[y_1, \dots, y_n]).$$

Proof. The functor $\hat{\blacksquare}_{-n}$ corresponds to the object $\tilde{\mathbb{I}}_{0|n} = (\mathfrak{gl}_n, \mathrm{Tr}(ZX)) \in \mathbb{H}\mathrm{om}(\mathfrak{n}^f, \mathbf{0}^f)$. Respectively, the composition with the object $\tilde{\mathcal{F}}_n = ((\tilde{\mathfrak{gl}}_n \times V_n)^{st}, \tilde{W}_n) \in \mathbb{H}\mathrm{om}(\mathbf{0}^f, \mathfrak{n}^f)$ is by the $G \times B$ -equivariant object:

$$(6.1) \quad \hat{\blacksquare}_{-n}(\tilde{\mathcal{F}}_n) = \tilde{\mathbb{I}}_{0|n} \circ \tilde{\mathcal{F}}_n = (\mathfrak{gl}_n \times \tilde{\mathfrak{gl}}_n \times V_n, \mathrm{Tr}(ZX) + \mathrm{Tr}(X\mathrm{Ad}_g Y)).$$

The potential of the last object is quadratic in Z and X thus after the Knörrer reduction we obtain the $G \times B$ equivariant object $G \times B$ -equivariant object $(G \times \mathfrak{n} \times V_n)^{st}$. The action of $G \times B$ on this object is free and a slice to the orbits of the action is defined by the triple $(1, K(y), v^0)$, where $K(y)_{ij} = \sum_{i=1}^n y_i \delta_{i,i} + \sum_{i=1}^{n-1} \delta_{i,j-1}$, $v_i^0 = \delta_{i,n}$. \square

The category of endomorphism of $\mathrm{Mod}(R_n)$, $R_n = \mathbb{C}[y_1, \dots, y_n]$ is the category of two-periodic complexes of R_n -bimodules $\mathbb{B}\mathrm{im}_n$. Respectively, the formula (6.1) implies that there is a monoidal functor:

$$\mathbb{B} : \mathrm{Hom}(\tilde{\mathcal{F}}_n, \tilde{\mathcal{F}}_n) : \rightarrow \mathbb{B}\mathrm{im}_n.$$

Combining the last functor with Φ^b we obtain a realization of the algebra \mathfrak{Br}_n^b inside $\mathbb{B}\mathrm{im}_n$. Thus if we have two radial segments I_{rad}, I'_{rad} with defects between the rays forming $\beta \in \mathfrak{Br}_n^b$, the value of the partition function Z^f on the segment is $\mathbb{B} \circ \Phi^b(\beta)$.

The bimodule realization of \mathfrak{Br}_n^b inside $\mathbb{B}\mathrm{im}$ used for HHH_{alg} is due to Soergel [Soe01]:

$$\Phi^S : \mathfrak{Br}_n^b \rightarrow \mathrm{Sbim}_n \subset \mathbb{B}\mathrm{im}_n, \quad \mathrm{HHH}_{alg}(\beta) := \mathrm{HH}_*(\Phi^S(\beta)),$$

where $\mathrm{HH}_* = \mathrm{Tor}_R$ is functor of Hochschild homology. The Hochschild homology functor is an trace functor $\mathrm{HHH}_{\mathrm{alg}}(\alpha \cdot \beta) = \mathrm{HHH}_{\mathrm{alg}}(\beta \cdot \alpha)$

We show in [OR20] that two realizations of $\mathfrak{B}\mathfrak{r}_n^{\flat}$ actually match and there is a natural isomorphism between the corresponding trace functors:

Theorem 6.1.3. [OR20] *For any n we have a commuting diagram of monoidal functors:*

$$\begin{array}{ccc} \mathfrak{B}\mathfrak{r}_n^{\flat} & \xrightarrow{\Phi^S} & \mathrm{Sbim}_n \\ \downarrow \Phi^{\flat} & & \downarrow \\ \mathrm{Hom}(\tilde{\mathcal{F}}_n, \tilde{\mathcal{F}}_n) & \xrightarrow{\mathbb{B}} & \mathbb{B}\mathrm{im}_n \end{array}$$

Moreover, the functors intertwine the trace functors:

$$\mathrm{HH}^*(\Phi^S(\beta)) = \mathcal{H}\mathrm{om}(\Phi^{\flat}(\beta), \Phi^{\flat}(1) \otimes \Lambda^* V_n).$$

6.2. Annular homology. In this section we explain the relations between the objects in the last column of the table from the introduction. The column has a natural physical explanation in terms of NS5 and D5 branes as we explained before and the mathematical side of the story is mostly contained in the work of Nakajima-Takayama [NT17],[Nak18] and Anno-Nandakumar [AN16].

The work of Nakajima-Takayama [NT17] explains realization of the Slodowy slices and their resolutions in terms of Cherkis quivers. These quiver varieties are related to the objects we study by the linear Koszul duality. We provide more details below.

Let us denote by Fl_{μ} , $\mu_n \leq \mu_{n-1} \leq \dots$, $\sum_{i=1}^m \mu_i = n$ the partial flag variety consisting of nested subspaces of dimension $M_1 = \mu_1$, $M_2 = \mu_1 + \mu_2$, $M_3 = \mu_1 + \mu_2 + \mu_3$, \dots . Similarly, we denote by G_{μ} the product of the groups $\mathrm{GL}_{M_1} \times \mathrm{GL}_{M_2} \times \dots \times \mathrm{GL}_{M_{m-1}}$. Respectively, $\mathcal{N}(\mu) \subset \mathfrak{gl}_n$ is the nilpotent orbit with the Jordan blocks of size μ_i and $\mathcal{S}(\mu)$ is a transversal slice to the the orbit $\mathcal{N}(\mu)$.

Proposition 6.2.1. [NT17, Theorem 7.11] *The $G_{\lambda} \times \mathrm{GL}_n \times G_{\mu}$ -quotient of the quiver $Q^{\mathrm{bow}}(\lambda, \mu)$ pictured below*

$$\begin{array}{ccccccc} \mathbb{C} & & \mathbb{C} & & \dots & & \mathbb{C} \\ \uparrow & \searrow & \uparrow & \searrow & \searrow & \uparrow & \searrow \\ \mathbb{C}^{L_1} & \rightarrow & \mathbb{C}^{L_3} & \rightarrow & \dots & \rightarrow & \mathbb{C}^{L_{l-1}} & \rightarrow & \mathbb{C}^n & \rightleftharpoons & \mathbb{C}^{M_{m-1}} & \rightleftharpoons & \dots & \rightleftharpoons & \mathbb{C}^{M_2} & \rightleftharpoons & \mathbb{C}^{M_1} \end{array}$$

is isomorphic to $\overline{\mathcal{N}}(\mu) \cap \mathcal{S}(\lambda)$ and the quotient of the stable locus is the simultaneous resolution $\mathcal{R}_{\lambda, \mu} = p^{-1}(\overline{\mathcal{N}}(\mu) \cap \mathcal{S}(\lambda)) \subset T^*Fl_{\mu}$ with $p : T^*Fl_{\mu} \rightarrow \mathfrak{g}$ being the Springer resolution map.

The relevant object in the TQFT setting discussed in our paper is a segment $I^{\lambda, \mu}$ with ends in the regions marked with $\mathbf{0}$ and such that the list of defects it traverses is $\mathrm{NS}5^{\lambda_1}, \mathrm{NS}5^{\lambda_2}, \dots, \mathrm{NS}5^{\lambda_l}, \mathrm{D}5^{\mu_m}, \dots, \mathrm{D}5^{\mu_1}$. As we explained in the section 3.6 the value $Z(I^{\lambda, \mu})$ of partition function is given by composition of morphisms

$$Z(I^{\lambda, \mu}) = \check{\square}_{\lambda_1} \circ \check{\square}_{\lambda_2} \circ \dots \circ \check{\square}_{\lambda_l} \circ \hat{\square}_{\mu_m} \circ \dots \circ \hat{\square}_{\mu_2} \circ \hat{\square}_{\mu_1}.$$

Thus the previous proposition gives a geometric description of the morphism

$$\mathbf{Z}(I^{\lambda,\mu}) = (Q^{bow} \times \text{Lie}(G_\lambda \times \text{GL}_n \times G_\mu), W^{\lambda,\mu}(z, x) = \text{Tr}(\mu(z)x)).$$

To specialize to the case treated in the table in the introduction we need to set $\mu = (1^n)$ and $\lambda = (k, n-k)$. Since the braid group $\mathfrak{B}\mathfrak{r}_n$ acts on $\widehat{\square}_1^n$ there is monoidal functor

$$\Phi^\lambda : \mathfrak{B}\mathfrak{r}_n \rightarrow \text{Hom}(\mathbf{Z}(I^{\lambda,1^n}), \mathbf{Z}(I^{\lambda,1^n})).$$

In the setting of the last column of the table from the introduction the radial rays in the picture (1.11) are homotopic to the interval $I^{(k,n-k),(1^n)}$. Thus the last column of the table is equivalent to the following conjecture that we plan to address in the forthcoming paper [OR22]:

Conjecture 6.2.2. *For any n a trace functor $\mathcal{T}r : \mathfrak{B}\mathfrak{r}_n \rightarrow 2 - \text{grVect}$*

$$\mathcal{T}r_{k,n-k}(\beta) = \mathcal{H}om(\Phi_{(k,n-k)}(\beta), \Phi_{(k,n-k)}(1_n))$$

categorifies the trace $\text{Tr}(\beta)[H_{1^n, \lambda+k}^\lambda]$.

The space $H_{1^n, \lambda+k}^\lambda$ is the weight k part of $L_1^{\otimes n}$, it is of dimension $\binom{n}{k}$. To provide an evidence to the conjecture we compute the Euler characteristic of the trace for the identity.

Proposition 6.2.3. *For any n, k we have:*

$$\chi(\mathcal{T}r_{k,n-k}(1_n)) = \binom{n}{k}.$$

Proof. First let us observe that by the linear Koszul duality for the matrix factorizations and by the proposition 6.2.1 the category of morphism $\text{Hom}(\mathbf{Z}(I^{\lambda,\mu}), \mathbf{Z}(I^{\lambda,\mu}))$ is equivalent to the dg category two-periodic complexes $D_{\mathbb{C}^* \times \mathbb{C}^*}^{per}(\mathcal{R}_{\lambda,\mu} \times \mathcal{R}_{\lambda,\mu})$. Respectively, $\Phi_\lambda(1_n)$ is the structure sheaf of the diagonal $\Delta \subset \mathcal{R}_{\lambda,1^n} \times \mathcal{R}_{\lambda,1^n}$.

Thus $\mathcal{T}r(1_n)$ is the homology self-intersection of the diagonal and since $\mathcal{R}_{\lambda,1^n}$ is smooth we get $\mathcal{T}r(1_n) = H^*(\oplus_i \Omega_{\mathcal{R}_{\lambda,1^n}}^i)$. Since the $\mathbb{C}^* \times \mathbb{C}^*$ -fixed locus in $\mathcal{R}_{\lambda,1^n}$ is zero-dimensional, the localization formula implies that Euler characteristics $\chi(\mathcal{T}r(1_n))$ is equal to the number of torus fixed points.

The 3D mirror partner of $\mathcal{R}_{(k,n-k),1^n}$ is the Grassmanian $Gr(k, n)$ [NT17]. The mirror symmetric partners have the same number of the torus fixed points and $Gr(k, n)$ has $\binom{n}{k}$ fixed points. \square

Finally, let us remark that in the work [AN16] the category of coherent sheaves on $\mathcal{R}_{(k,k),(1^{2k})}$ was used to construct the invariants of the affine tangles and relate this invariant to the annular Khovanov homology [APS04]. The under/over-crossing the generators of the affine tangles in [AN16] match with the braid generators used in this paper. The cap/cup of the affine tangle monoid from [AN16] have a natural interpretation in terms of NS5 defects. Thus in the forthcoming paper [OR22] we also address the

Conjecture 6.2.4. *The construction of the trace functor $\mathcal{T}r_{(k,k)}$ can be extended to the affine tangle and the corresponding invariant provides a realization of the sl_2 annular Khovanov homology.*

REFERENCES

- [AK15] S. Arkhipov and T. Kanstrup. Equivariant Matrix Factorizations and Hamiltonian reduction. *Bull. of Korean Math. Soc.*, (5):1803–1825, 2015, 1510.07472v1.
- [AN16] R. Anno and V. Nandakumar. Exotic t-structures for two-block Springer fibers, 2016, 1602.00768v1.
- [APS04] M. Asaeda, J. Przytycki, and A. Sikora. Categorification of the kauffman bracket skein module of bundles over surfaces. *Algebraic and Geometric Topology*, 4(2):1177–1210, Dec 2004.
- [Arn89] V. Arnold. *Mathematical methods of classical mechanics*. Springer, 1989.
- [BN96] J. Baez and M. Neuchl. Higher dimensional algebra. *Advances in Mathematics*, 121(2):196–244, Aug 1996.
- [CW07] A. Caldararu and S. Willerton. The Mukai pairing, I: a categorical approach, 2007, 0707.2052v1.
- [EGNO15] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. Tensor categories. *Mathematical Surveys and Monographs*, Jul 2015.
- [GG09] R. Garner and N. Gurski. The low-dimensional structures formed by tricategories. *Mathematical Proceedings of the Cambridge Philosophical Society*, 146(03):551, Jan 2009.
- [GR17] D. Gaitsgory and N. Rozenblyum. A study in derived algebraic geometry. *Mathematical Surveys and Monographs*, Jul 2017.
- [GW09] D. Gaiotto and E. Witten. Supersymmetric boundary conditions in $\mathcal{N} = 4$ super yang-mills theory. *Journal of Statistical Physics*, 135(5-6):789–855, Feb 2009.
- [Jon87] V. F. R. Jones. Hecke algebra representations of braid groups and link polynomials. *The Annals of Mathematics*, 126(2):335–392, Sep 1987.
- [Kno87] H. Knorrer. Cohen-Macaulay modules on hypersurface singularities. I. *Inventiones Mathematicae*, (1):153–164, 1987.
- [KR08] M. Khovanov and L. Rozansky. Matrix factorizations and link homology II. *Geometry and Topology*, 12:1387–1425, 2008.
- [KR10] A. Kapustin and L. Rozansky. Three-dimensional topological field theory and symplectic algebraic geometry II. *Communications in Number Theory and Physics*, 4(3):463–549, 2010.
- [KRS09] A. Kapustin, L. Rozansky, and N. Saulina. Three-dimensional topological field theory and symplectic algebraic geometry I. *Nuclear Physics B*, 816(3):295–355, Aug 2009.
- [KV94] M. M. Kapranov and V. A. Voevodsky. 2-categories and zamolodchikov tetrahedra equations. *Algebraic Groups and Their Generalizations: Quantum and Infinite-Dimensional Methods*, page 177–259, 1994.
- [KW07] A. Kapustin and E. Witten. Electric-magnetic duality and the geometric langlands program. *Communications in Number Theory and Physics*, 1(1):1–236, 2007.
- [Lur08] J. Lurie. On the classification of topological field theories. *Current Developments in Mathematics*, 2008(1):129–280, 2008.
- [MOY98] H. Murakami, T. Ohtsuki, and Sh. Yamada. HOMFLY polynomial via an invariant of colored plane graphs. *Enseign. Math.*, 44(2):325–360, 1998.
- [Nak99] H. Nakajima. Lectures on Hilbert schemes of points on surfaces. *University Lecture Series*, Sep 1999.
- [Nak18] H. Nakajima. Geometric Satake correspondence for affine Kac-Moody Lie algebras of type A , 2018, 1812.11710v1.
- [NT17] H. Nakajima and Y. Takayama. Cherkis bow varieties and coulomb branches of quiver gauge theories of affine type a . *Selecta Mathematica*, 23(4):2553–2633, Jun 2017.
- [OR17] A. Oblomkov and L. Rozansky. HOMFLYPT homology of Coxeter links, 2017, 1706.00124v1.
- [OR18a] A. Oblomkov and L. Rozansky. 3D TQFT and HOMFLYPT homology. 2018, 1812.06340.
- [OR18b] A. Oblomkov and L. Rozansky. A categorification of a cyclotomic Hecke algebra, 2018, 1801.06201v1.

- [OR18c] A. Oblomkov and L. Rozansky. Affine braid group, JM elements and knot homology. *Transformation Groups*, Jan 2018.
- [OR18d] A. Oblomkov and L. Rozansky. Categorical Chern character and braid groups, 2018, 1811.03257v1.
- [OR18e] A. Oblomkov and L. Rozansky. Knot homology and sheaves on the Hilbert scheme of points on the plane. *Selecta Mathematica*, 24(3):2351–2454, Jan 2018.
- [OR20] A. Oblomkov and L. Rozansky. Soergel bimodules and matrix factorizations, 2020, 2010.14546v1.
- [OR22] A. Oblomkov and L. Rozansky. Matrix factorizations and annular homology. *in preparation*, 2022.
- [RR21] R. Rimanyi and L. Rozansky. New quiver-like varieties and Lie superalgebras, 2021, 2105.11499v1.
- [Soe01] W. Soergel. Langlands’ philosophy and Koszul duality. In *Algebra-Representation Theory (Constanta, 2000)*, pages 379–414, 2001.

A. OBLOMKOV, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS AT AMHERST, LEDERLE GRADUATE RESEARCH TOWER, 710 N. PLEASANT STREET, AMHERST, MA 01003 USA

Email address: `oblomkov@math.umass.edu`

L. ROZANSKY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, CB # 3250, PHILLIPS HALL, CHAPEL HILL, NC 27599 USA

Email address: `rozansky@math.unc.edu`