

Asymptotic behavior of the $W^{1/q,q}$ -norm of mollified BV functions and applications to singular perturbation problems

ARKADY POLIAKOVSKY ¹

Department of Mathematics, Ben Gurion University of the Negev,
P.O.B. 653, Be'er Sheva 84105, Israel

Abstract

Motivated by results of Figalli and Jerison [8] and Hernández [7], we prove the following formula:

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|\eta_\varepsilon * u\|_{W^{1/q,q}(\Omega)}^q = C_0 \int_{J_u} |u^+(x) - u^-(x)|^q d\mathcal{H}^{N-1}(x),$$

where $\Omega \subset \mathbb{R}^N$ is a regular domain, $u \in BV(\Omega) \cap L^\infty$, $q > 1$ and $\eta_\varepsilon(z) = \varepsilon^{-N} \eta(z/\varepsilon)$ is a smooth mollifier. In addition, we apply the above formula to the study of certain singular perturbation problems.

1 Introduction

Figalli and Jerison found in [8] a relationship between the perimeter of a set and a fractional Sobolev norm of its characteristic function. More precisely, for the mollifying kernel $\eta_\varepsilon(z) = \varepsilon^{-N} \eta(z/\varepsilon)$, where $\eta(z)$ denotes the standard Gaussian in \mathbb{R}^N , they showed that there exist constants $C_1 > 0$ and $C_2 > 0$ such that for every set $A \subset \mathbb{R}^N$ of finite perimeter $P(A)$ we have

$$C_1 P(A) \leq \liminf_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|\eta_\varepsilon * \chi_A\|_{H^{1/2}(\mathbb{R}^N)}^2 \leq \limsup_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|\eta_\varepsilon * \chi_A\|_{H^{1/2}(\mathbb{R}^N)}^2 \leq C_2 P(A), \quad (1.1)$$

where χ_A is the characteristic function of A . More recently, Hernández improved this result in [7] as follows. For η_ε as above he showed that there exist a constant $C_0 > 0$ such that for every $u \in BV(\mathbb{R}^N) \cap L^\infty$ we have

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|\eta_\varepsilon * u\|_{H^{1/2}(\mathbb{R}^N)}^2 = C_0 \int_{J_u} |u^+(x) - u^-(x)|^2 d\mathcal{H}^{N-1}(x). \quad (1.2)$$

¹E-mail: poliakov@math.bgu.ac.il

A related result in which the same R.H.S. as in (1.2) appears, was obtained in [13]. More precisely, we showed in [13] that for every radial $\eta \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$ there exists a constant $C = C_\eta > 0$ such that for every $u \in BV(\Omega, \mathbb{R}^d) \cap L^\infty$ we have

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon \|\eta_\varepsilon * u\|_{H^1(\Omega)}^2 = C_\eta \int_{J_u} \left| u^+(x) - u^-(x) \right|^2 d\mathcal{H}^{N-1}(x). \quad (1.3)$$

More recently, we showed in [14] yet another related result:

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^N$ be an open set with bounded Lipschitz boundary and let $u \in BV(\Omega, \mathbb{R}^d) \cap L^\infty(\Omega, \mathbb{R}^d)$. Then, for every $q > 1$ we have*

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \int_{B_\varepsilon(x) \cap \Omega} \frac{1}{\varepsilon^N} \frac{|u(y) - u(x)|^q}{|y - x|} dy dx = C_N \int_{J_u} \left| u^+(x) - u^-(x) \right|^q d\mathcal{H}^{N-1}(x), \quad (1.4)$$

with the dimensional constant $C_N > 0$ defined by

$$C_N := \frac{1}{N} \int_{S^{N-1}} |z_1| d\mathcal{H}^{N-1}(z), \quad (1.5)$$

where we denote $z := (z_1, \dots, z_N) \in \mathbb{R}^N$.

In the present paper we generalize the formula (1.2) in several aspects:

- We allow a general mollifying kernel $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$ (not only the Gaussian as before),
- We allow a general domain $\Omega \subset \mathbb{R}^N$, of certain regularity, while previous results required $\Omega = \mathbb{R}^N$,
- We treat the $W^{1/q,q}(\Omega)$ -norm for any $q > 1$, while previous results were restricted to the case $q = 2$.

Recall that the Gagliardo seminorm $\|u\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}$ is given by

$$\|u\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)} := \left(\int_{\Omega} \left(\int_{\Omega} \frac{|u(x) - u(y)|^q}{|x - y|^{N+1}} dy \right) dx \right)^{\frac{1}{q}}. \quad (1.6)$$

Our first main result is

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^N$ be an open set and let $u \in BV(\mathbb{R}^N, \mathbb{R}^d) \cap L^\infty(\mathbb{R}^N, \mathbb{R}^d)$ be such that $\|Du\|(\partial\Omega) = 0$. For $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$, every $x \in \mathbb{R}^N$ and every $\varepsilon > 0$ define*

$$u_\varepsilon(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta\left(\frac{y - x}{\varepsilon}\right) u(y) dy = (\eta_\varepsilon * u)(x). \quad (1.7)$$

Then, for any $q > 1$ we have

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|u_\varepsilon\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q = 2 \left| \int_{\mathbb{R}^N} \eta(z) dz \right|^q \left(\int_{\mathbb{R}^{N-1}} \frac{dv}{(\sqrt{1 + |v|^2})^{N+1}} \right) \int_{J_u \cap \Omega} \left| u^+(x) - u^-(x) \right|^q d\mathcal{H}^{N-1}(x). \quad (1.8)$$

Theorem 1.2 enables us to prove an upper bound, in the limit $\varepsilon \rightarrow 0^+$, for the following singular perturbation functionals with differential constraints:

(i)

$$E_\varepsilon^{(1)}(v) := \begin{cases} \frac{1}{|\ln \varepsilon|} \|v\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_\Omega W(v, x) dx & \text{if } A \cdot \nabla v = 0 \\ +\infty & \text{otherwise,} \end{cases} \quad (1.9)$$

for $v : \Omega \rightarrow \mathbb{R}^d$;

(ii)

$$E_\varepsilon^{(2)}(v) := \begin{cases} \frac{1}{|\ln \varepsilon|} \left(\|v\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|v\|_{W^{1/q,q}(\mathbb{R}^N \setminus \bar{\Omega}, \mathbb{R}^d)}^q \right) + \frac{1}{\varepsilon} \int_\Omega W(v, x) dx & \text{if } A \cdot \nabla v = 0 \\ +\infty & \text{otherwise,} \end{cases} \quad (1.10)$$

for $v : \mathbb{R}^N \rightarrow \mathbb{R}^d$.

In both cases $A : \mathbb{R}^{d \times N} \rightarrow \mathbb{R}^l$ is a linear operator (possibly trivial). The most important particular cases are the following:

(a) $A \equiv 0$ (i.e., without any prescribed differential constraint),

(b) $d = N$, $l = N^2$ and $A \cdot \nabla v \equiv \text{curl } v := \{\partial_k v_j - \partial_j v_k\}_{1 \leq k, j \leq N}$,

(c) $l = d$ and $A \cdot \nabla v \equiv \text{div } v$.

The Γ -limit of the functional (1.9) in the L^p -topology when $A \equiv 0$, $q = 2$, $N = 1$ and W is a double-well potential was found by Alberti, Bouchitté and Seppecher [1]. The result was generalized to any dimension $N \geq 1$, for the functional (1.10), by Savin and Valdinoci [15].

Note that the functional (1.9) resembles the energy functional in the following singular perturbation problem:

$$\hat{E}_\varepsilon(v) := \begin{cases} \varepsilon^{q-1} \|v\|_{W^{1,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_\Omega W(v, x) dx & \text{if } A \cdot \nabla v = 0 \\ +\infty & \text{otherwise,} \end{cases} \quad (1.11)$$

that attracted a lot of attention by many authors, starting from Modica and Mortola [10], Modica [9], Sternberg [16] and others, who studied the basic special case of (1.11) with $A \equiv 0$, $q = 2$ and W being a double-well potential. The Γ limit of (1.11) with $A \equiv 0$, $q = 2$ and a general $W \in C^0$ that does not depend on x , was found by Ambrosio in [2]. As an example with a nontrivial differential constraint we mention the Aviles-Giga functional, that appear in various applications. It is defined for scalar functions ψ by

$$\tilde{E}_\varepsilon(\psi) := \int_\Omega \left\{ \varepsilon |\nabla^2 \psi|^2 + \frac{1}{\varepsilon} \left(1 - |\nabla \psi|^2 \right)^2 \right\} dx \quad (\text{see [3, 5, 6]}), \quad (1.12)$$

and the objective is to study the Γ -limit, as $\varepsilon \rightarrow 0^+$. This can be seen as a special case of (1.11) if we set $v := \nabla \psi$ and let $A \cdot \nabla v \equiv \text{curl } v$, $q = 2$ and $W(v, x) = (1 - |v|^2)^2$.

Our second result provides an upper bound for the energies (1.9)-(1.10):

Theorem 1.3. *Let $\Omega \subset \mathbb{R}^N$ be an open set and let $W : \mathbb{R}^d \times \mathbb{R}^N \rightarrow \mathbb{R}$ be a Borel measurable nonnegative function, continuous and continuously differentiable w.r.t. the first argument, such that $W(0, \cdot) \in L^1(\Omega, \mathbb{R})$. Assume further that for every $D > 0$ there exists $C := C_D > 0$ such that*

$$|\nabla_b W(b, x)| \leq C_D \quad \forall x \in \mathbb{R}^N, \forall b \in B_D(0). \quad (1.13)$$

Let $u \in BV(\mathbb{R}^N, \mathbb{R}^d) \cap L^\infty(\mathbb{R}^N, \mathbb{R}^d)$ be such that $W(u(x), x) = 0$ a.e. in Ω , $\|Du\|(\partial\Omega) = 0$, and $A \cdot Du = 0$ in \mathbb{R}^N , where $A : \mathbb{R}^{d \times N} \rightarrow \mathbb{R}^l$ is a prescribed linear operator (possibly trivial). Then, for any $q > 1$ there exists a sequence of functions $\{\psi_\varepsilon\}_{\varepsilon > 0} \subset C^\infty(\mathbb{R}^N, \mathbb{R}^d) \cap W^{1,1}(\mathbb{R}^N, \mathbb{R}^d) \cap W^{1,\infty}(\mathbb{R}^N, \mathbb{R}^d)$ such that $A \cdot D\psi_\varepsilon = 0$ in \mathbb{R}^N , $\psi_\varepsilon(x) \rightarrow u(x)$ strongly in $L^p(\mathbb{R}^N, \mathbb{R}^d)$ for every $p \geq 1$, and

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{|\ln \varepsilon|} \left(\|\psi_\varepsilon\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|\psi_\varepsilon\|_{W^{1/q,q}(\mathbb{R}^N \setminus \bar{\Omega}, \mathbb{R}^d)}^q \right) + \frac{1}{\varepsilon} \int_{\Omega} W(\psi_\varepsilon(x), x) dx \right) = \\ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{|\ln \varepsilon|} \|\psi_\varepsilon\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_{\Omega} W(\psi_\varepsilon(x), x) dx \right) = \\ \left(\int_{\mathbb{R}^{N-1}} \frac{2}{(\sqrt{1+|v|^2})^{N+1}} dv \right) \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \end{aligned} \quad (1.14)$$

Moreover, in the case $A \equiv 0$ we can choose ψ_ε to satisfy also

$$\int_{\Omega} \psi_\varepsilon(x) dx = \int_{\Omega} u(x) dx \quad \forall \varepsilon > 0. \quad (1.15)$$

Unfortunately, the upper bound found in Theorem 1.3 is not sharp in the most general case with a nontrivial prescribed differential constraint. For example, in the particular case of (1.9) with $N = 2$, $A \cdot \nabla v \equiv \text{curl } v$, $q > 3$ and $W(v, x) = (1 - |v|^2)^2$, the functional on the R.H.S. of (1.14) is not lower semicontinuous, hence cannot be the Γ -limit (see [3]). However, we still hope that the result of the above theorem could provide the sharp upper bound in some cases with $A = 0$. Indeed, the Γ -limit, computed in [1] for the special case of (1.9) with $A \equiv 0$, $q = 2$, $N = 1$ and W being a double well potential, coincides with the upper bound found in Theorem 1.3. Moreover, since the functional in (1.10) is superior to the functional in (1.9), the Γ -limit, found in [15] (see also [12]) for the energy (1.10) in any dimension $N \geq 1$ with $A \equiv 0$, $q = 2$ and W being a double well potential, coincides again with our upper bound.

The paper is organized as follows. In section 2 we prove our two main results. For the convenience of the reader, in the Appendix we recall some known results on BV functions, needed for the proofs.

2 Proof of the main results

Proposition 2.1. *Let $q > 1$, $\Omega \subset \mathbb{R}^N$ be an open set and $u \in BV(\mathbb{R}^N, \mathbb{R}^d) \cap L^\infty(\mathbb{R}^N, \mathbb{R}^d)$ be such that $\|Du\|(\partial\Omega) = 0$. Let $\eta \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$ and for every $x \in \mathbb{R}^N$ and every $\varepsilon > 0$ define*

$$u_\varepsilon(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy = (\eta_\varepsilon * u)(x). \quad (2.1)$$

Then,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|u_\varepsilon\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q &= \\ 2 \left| \int_{\mathbb{R}^N} \eta(z) dz \right|^q &\left(\int_{\mathbb{R}^{N-1}} \frac{1}{(\sqrt{1+|v|^2})^{N+1}} dv \right) \int_{J_u \cap \Omega} |u^+(x) - u^-(x)|^q d\mathcal{H}^{N-1}(x). \end{aligned} \quad (2.2)$$

Proof. We start with some notations. For every $\nu \in S^{N-1}$ and $x \in \mathbb{R}^N$ set

$$H_+(x, \nu) = \{\xi \in \mathbb{R}^N : (\xi - x) \cdot \nu > 0\}, \quad (2.3)$$

$$H_-(x, \nu) = \{\xi \in \mathbb{R}^N : (\xi - x) \cdot \nu < 0\} \quad (2.4)$$

and

$$H_0(\nu) = \{\xi \in \mathbb{R}^N : \xi \cdot \nu = 0\}. \quad (2.5)$$

Let $R > 0$ be such that $\text{supp } \eta \subset B_R(0)$. For every $x \in \mathbb{R}^N$ and every $\varepsilon > 0$ we rewrite (2.1) as:

$$u_\varepsilon(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy = \int_{\mathbb{R}^N} \eta(z) u(x + \varepsilon z) dz = \int_{B_R(0)} \eta(z) u(x + \varepsilon z) dz. \quad (2.6)$$

By (2.6) we have

$$\begin{aligned} \frac{d}{d\varepsilon} u_\varepsilon(x) &:= -\frac{N}{\varepsilon^{N+1}} \int_{\mathbb{R}^N} \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy - \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \frac{y-x}{\varepsilon^2} \cdot \nabla \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy = \\ &- \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \text{div}_y \left\{ \eta\left(\frac{y-x}{\varepsilon}\right) \frac{y-x}{\varepsilon} \right\} u(y) dy = \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta\left(\frac{y-x}{\varepsilon}\right) \frac{y-x}{\varepsilon} \cdot d[Du(y)]. \end{aligned} \quad (2.7)$$

Moreover, by (1.6) we have

$$\begin{aligned} \|u_\varepsilon\|_{W^{1/q,q}}^q &= \|u_\varepsilon\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q = \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} \chi_\Omega(y) dy \right) \chi_\Omega(x) dx \\ &= \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+z) - u_\varepsilon(x)|^q}{|z|^{N+1}} \chi_\Omega(x+z) \chi_\Omega(x) dz \right) dx, \end{aligned} \quad (2.8)$$

where

$$\chi_\Omega(x) := \begin{cases} 1 & \forall x \in \Omega \\ 0 & \forall x \in \mathbb{R}^N \setminus \Omega \end{cases}. \quad (2.9)$$

Thus,

$$\frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = -\frac{1}{\ln \varepsilon} \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+z) - u_\varepsilon(x)|^q}{|z|^{N+1}} \chi_\Omega(x+z) \chi_\Omega(x) dz \right) dx. \quad (2.10)$$

Since $-\ln \varepsilon \rightarrow +\infty$ as $\varepsilon \rightarrow 0^+$, applying L'Hôpital's rule to the expression in (2.10) yields

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q &= \\ - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{\varepsilon}{|z|^{N+1}} \left(\frac{d}{d\varepsilon} (u_\varepsilon(x+z) - u_\varepsilon(x)) \right) \cdot \nabla F_q(u_\varepsilon(x+z) - u_\varepsilon(x)) \chi_\Omega(x+z) \chi_\Omega(x) dz \right) dx, \end{aligned} \quad (2.11)$$

where $F_q \in C^1(\mathbb{R}^d, \mathbb{R})$ is defined by

$$F_q(h) := |h|^q \quad \forall h \in \mathbb{R}^d. \quad (2.12)$$

Thus, by (2.11), (2.6) and (2.7) we get

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\ & - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\varepsilon}{|z|^{N+1}} \left\{ \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \left(\eta\left(\frac{y-(x+z)}{\varepsilon}\right) \frac{y-(x+z)}{\varepsilon} - \eta\left(\frac{y-x}{\varepsilon}\right) \frac{y-x}{\varepsilon} \right) \cdot d[Du(y)] \right\} \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} \eta(\xi) (u(x+z+\varepsilon\xi) - u(x+\varepsilon\xi)) d\xi \right) \chi_\Omega(x+z) \chi_\Omega(x) dz dx = \\ & - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\varepsilon}{|z|^{N+1}} \frac{1}{\varepsilon^N} \left(\eta\left(\frac{y-(x+z)}{\varepsilon}\right) \frac{y-(x+z)}{\varepsilon} - \eta\left(\frac{y-x}{\varepsilon}\right) \frac{y-x}{\varepsilon} \right) \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} \eta(\xi) (u(x+z+\varepsilon\xi) - u(x+\varepsilon\xi)) d\xi \right) \chi_\Omega(x+z) \chi_\Omega(x) dz dx \cdot d[Du(y)]. \quad (2.13) \end{aligned}$$

Changing variable, $z/\varepsilon \rightarrow z$, in the integration on the R.H.S. of (2.13) gives

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\ & - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \frac{1}{\varepsilon^N} \left(\eta\left(\frac{y-x}{\varepsilon} - z\right) \left(\frac{y-x}{\varepsilon} - z\right) - \eta\left(\frac{y-x}{\varepsilon}\right) \frac{y-x}{\varepsilon} \right) \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} \eta(\xi) (u(x+\varepsilon z + \varepsilon\xi) - u(x+\varepsilon\xi)) d\xi \right) \chi_\Omega(x+\varepsilon z) \chi_\Omega(x) dz dx \cdot d[Du(y)] = \\ & - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x-z)(x-z) - \eta(x)x \right) \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} \eta(\xi) (u(y+\varepsilon z + \varepsilon\xi - \varepsilon x) - u(y+\varepsilon\xi - \varepsilon x)) d\xi \right) \chi_\Omega(y-\varepsilon x + \varepsilon z) \chi_\Omega(y-\varepsilon x) dz dx \cdot d[Du(y)]. \quad (2.14) \end{aligned}$$

Therefore,

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q \\ & = - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x-z)(x-z) - \eta(x)x \right) \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} (\eta(\xi-z) - \eta(\xi)) u(y+\varepsilon\xi - \varepsilon x) d\xi \right) \chi_\Omega(y-\varepsilon x + \varepsilon z) \chi_\Omega(y-\varepsilon x) dz dx \cdot d[Du(y)] \\ & = - \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x-z)(x-z) - \eta(x)x \right) \times \\ & \quad \times \nabla F_q \left(\int_{\mathbb{R}^N} (\eta(\xi+x-z) - \eta(\xi+x)) u(y+\varepsilon\xi) d\xi \right) \chi_\Omega(y-\varepsilon x + \varepsilon z) \chi_\Omega(y-\varepsilon x) dz dx \cdot d[Du(y)]. \quad (2.15) \end{aligned}$$

On the other hand, by (3.1) in the Appendix, for every $x, z \in \mathbb{R}^N$ and \mathcal{H}^{N-1} -a.e. $y \in \mathbb{R}^N$ we have

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \left\{ \int_{\mathbb{R}^N} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) u(y + \varepsilon \xi) d\xi \right\} = \\ & u^+(y) \int_{H_+(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi + u^-(y) \int_{H_-(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi. \end{aligned} \quad (2.16)$$

with $H_{\pm}(x, \nu)$ as defined in (2.3) and (2.4). Thus, since $\|Du\|(\partial\Omega) = 0$, by (2.16) and the Dominated Convergence Theorem we obtain:

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_{\varepsilon}\|_{W^{1/q, q}}^q = \\ & - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x - z)(x - z) - \eta(x)x \right) \nabla F_q \left(u^+(y) \int_{H_+(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right. \\ & \quad \left. + u^-(y) \int_{H_-(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right) \chi_{\Omega}^2(y) dz dx \cdot d[Du(y)] = \\ & - \int_{\Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x - z)(x - z) - \eta(x)x \right) \nabla F_q \left(u^+(y) \int_{H_+(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right. \\ & \quad \left. + u^-(y) \int_{H_-(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right) dz dx \cdot d[Du(y)]. \end{aligned} \quad (2.17)$$

It follows that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_{\varepsilon}\|_{W^{1/q, q}}^q = - \int_{\Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x - z)(x - z) - \eta(x)x \right) \times \\ & \quad \times \nabla F_q \left((u^+(y) - u^-(y)) \int_{H_+(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right. \\ & \quad \left. + u^-(y) \int_{\mathbb{R}^N} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right) dz dx \cdot d[Du(y)] \\ & = - \int_{\Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x - z)(x - z) - \eta(x)x \right) \times \\ & \quad \times \nabla F_q \left((u^+(y) - u^-(y)) \int_{H_+(0, \nu(y))} \left(\eta(\xi + x - z) - \eta(\xi + x) \right) d\xi \right) dz dx \cdot d[Du(y)], \end{aligned} \quad (2.18)$$

where we used in the last step the fact that $\int_{\mathbb{R}^N} \eta(\xi + x - z) d\xi = \int_{\mathbb{R}^N} \eta(\xi + x) d\xi$. Next, by

(2.18) and (2.12) we infer that

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = - \int_{\Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x-z)(x-z) - \eta(x)x \right) \times \\
& \quad \times \nabla F_q \left((u^+(y) - u^-(y)) \left(\int_{H_+(x-z, \nu(y))} \eta(\xi) d\xi - \int_{H_+(x, \nu(y))} \eta(\xi) d\xi \right) \right) dz dx \cdot d[Du(y)] \\
& \quad = \int_{J_u \cap \Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{1}{|z|^{N+1}} \left(\eta(x)x \cdot \nu(y) - \eta(x-z)(x-z) \cdot \nu(y) \right) \times \\
& \quad \times \frac{dG_q}{d\rho} \left(\int_{(x-z) \cdot \nu(y)}^{x \cdot \nu(y)} \int_{H_0(\nu(y))} \eta(t\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) dt \right) dx dz |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y), \quad (2.19)
\end{aligned}$$

where $G_q(\rho) \in C^1(\mathbb{R}, \mathbb{R})$ is defined by

$$G_q(\rho) := |\rho|^q \quad \forall \rho \in \mathbb{R}, \quad (2.20)$$

and $H_0(\nu)$ is defined in (2.5). Therefore,

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\
& \int_{J_u \cap \Omega} \int_{\mathbb{R}^N} \int_{\mathbb{R}} \int_{H_0(\nu(y))} \frac{1}{|z|^{N+1}} \left(\eta(s\nu(y) + \zeta)s - \eta((s-z \cdot \nu(y))\nu(y) + \zeta)(s-z \cdot \nu(y)) \right) \times \\
& \times \frac{dG_q}{d\rho} \left(\int_{s-z \cdot \nu(y)}^s \int_{H_0(\nu(y))} \eta(t\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) dt \right) d\mathcal{H}^{N-1}(\zeta) ds dz |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \\
& \quad = \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}^{N-1}} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{1}{(\sqrt{\tau^2 + |w|^2})^{N+1}} \times \right. \\
& \quad \times \left. \left(\int_{H_0(\nu(y))} \left(\eta(s\nu(y) + \zeta)s - \eta((s-\tau)\nu(y) + \zeta)(s-\tau) \right) d\mathcal{H}^{N-1}(\zeta) \right) \times \right. \\
& \quad \times \left. \frac{dG_q}{d\rho} \left(\int_{s-\tau}^s \int_{H_0(\nu(y))} \eta(t\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) dt \right) d\tau ds dw \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.21)
\end{aligned}$$

Introducing the notation

$$\Lambda(y, a, b) = \int_a^b \int_{H_0(\nu(y))} \eta(t\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) dt \quad (2.22)$$

allows us to rewrite (2.21) as

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\
& \int_{J_u \cap \Omega} \left\{ \int_{\mathbb{R}^{N-1}} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{1}{\tau^2} \frac{1}{|\tau|^{N-1}} \frac{1}{(\sqrt{1 + |w/|\tau|^2})^{N+1}} \times \right. \\
& \quad \left(\int_{H_0(\nu(y))} \left(\eta(s\nu(y) + \zeta)s - \eta((s-\tau)\nu(y) + \zeta)(s-\tau) \right) d\mathcal{H}^{N-1}(\zeta) \right) \times \\
& \quad \times \left. \frac{dG_q}{d\rho} \left(\Lambda(y, s-\tau, s) \right) d\tau ds dw \right\} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.23)
\end{aligned}$$

The change of variables $w/|\tau| \rightarrow v$ in the R.H.S. of (2.23) gives

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\ & D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{1}{\tau^2} \left(\int_{H_0(\nu(y))} \left(\eta(s\nu(y) + \zeta) s - \eta((s-\tau)\nu(y) + \zeta)(s-\tau) \right) d\mathcal{H}^{N-1}(\zeta) \right) \times \right. \\ & \quad \left. \times \frac{dG_q}{d\rho}(\Lambda(y, s-\tau, s)) d\tau ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y), \quad (2.24) \end{aligned}$$

where D_N is the dimensional constant given by

$$D_N := \int_{\mathbb{R}^{N-1}} \frac{1}{(\sqrt{1+|v|^2})^{N+1}} dv. \quad (2.25)$$

Then we rewrite (2.24) as

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\ & \lim_{M \rightarrow +\infty} \left(D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \int_{-M}^M \frac{1}{\tau^2} \left(\int_{H_0(\nu(y))} s \left(\eta(s\nu(y) + \zeta) - \eta((s-\tau)\nu(y) + \zeta) \right) d\mathcal{H}^{N-1}(\zeta) \right) \times \right. \right. \\ & \quad \left. \left. \times \frac{dG_q}{d\rho}(\Lambda(y, s-\tau, s)) d\tau ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \right. \\ & \quad \left. + D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \int_{-M}^M \frac{1}{\tau} \left(\int_{H_0(\nu(y))} \eta((s-\tau)\nu(y) + \zeta) d\mathcal{H}^{N-1}(\zeta) \right) \times \right. \right. \\ & \quad \left. \left. \times \frac{dG_q}{d\rho}(\Lambda(y, s-\tau, s)) d\tau ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \right). \quad (2.26) \end{aligned}$$

Integration by parts of (2.26) and using (2.20) give

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\ & - \lim_{M \rightarrow +\infty} D_N \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q \left(\int_{\mathbb{R}} \int_{-M}^M \frac{1}{\tau^2} |\Lambda(y, s-\tau, s)|^q d\tau ds \right) d\mathcal{H}^{N-1}(y) \\ & + \lim_{M \rightarrow +\infty} D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \int_{-M}^M \frac{1}{\tau^2} |\Lambda(y, s-\tau, s)|^q d\tau ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \\ & + \lim_{M \rightarrow +\infty} \frac{D_N}{M} \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} |\Lambda(y, s-M, s)|^q ds + \int_{\mathbb{R}} |\Lambda(y, s, s+M)|^q ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \\ & = \lim_{M \rightarrow +\infty} \frac{D_N}{M} \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} |\Lambda(y, s-M, s)|^q ds + \int_{\mathbb{R}} |\Lambda(y, s, s+M)|^q ds \right) |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.27) \end{aligned}$$

Therefore, applying L'Hôpital's rule in (2.27), using (2.20), we deduce that

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\
& \lim_{M \rightarrow +\infty} D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, s - M, s) \right) \left(\int_{H_0(\nu(y))} \eta((s - M)\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right. \\
& \quad \left. + \int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, s, s + M) \right) \left(\int_{H_0(\nu(y))} \eta((s + M)\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right) \\
& \quad \times |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.28)
\end{aligned}$$

Changing variables of integration we rewrite (2.28) as

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\
& \lim_{M \rightarrow +\infty} D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, s, s + M) \right) \left(\int_{H_0(\nu(y))} \eta(s\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right. \\
& \quad \left. + \int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, s - M, s) \right) \left(\int_{H_0(\nu(y))} \eta(s\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right) \\
& \quad \times |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \\
& = D_N \int_{J_u \cap \Omega} \left(\int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, s, \infty) \right) \left(\int_{H_0(\nu(y))} \eta(s\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right. \\
& \quad \left. + \int_{\mathbb{R}} \frac{dG_q}{d\rho} \left(\Lambda(y, -\infty, s) \right) \left(\int_{H_0(\nu(y))} \eta(s\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) \right) ds \right) \\
& \quad \times |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.29)
\end{aligned}$$

Applying Newton-Leibniz formula in (2.29) and using (2.20) we obtain that

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln \varepsilon} \|u_\varepsilon\|_{W^{1/q,q}}^q = \\
& 2D_N \int_{J_u \cap \Omega} \left| \int_{-\infty}^{\infty} \int_{H_0(\nu(y))} \eta(t\nu(y) + \xi) d\mathcal{H}^{N-1}(\xi) dt \right|^q |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \\
& = 2D_N \left| \int_{\mathbb{R}^N} \eta(z) dz \right|^q \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y), \quad (2.30)
\end{aligned}$$

and (2.2) follows. \square

Corollary 2.1. *Let $q > 1$ and let $\Omega \subset \mathbb{R}^N$ be an open set. Assume $W : \mathbb{R}^d \times \mathbb{R}^N \rightarrow \mathbb{R}$ is a Borel measurable function such that, $W(0, \cdot) \in L^1(\Omega, \mathbb{R})$ and for every $D > 0$ there exists $C := C_D > 0$ such that*

$$|W(b, x) - W(a, x)| \leq C_D |b - a| \quad \forall x \in \mathbb{R}^N, \forall a, b \in B_D(0). \quad (2.31)$$

Let $u \in BV(\mathbb{R}^N, \mathbb{R}^d) \cap L^\infty(\mathbb{R}^N, \mathbb{R}^d)$ be such that $\|Du\|(\partial\Omega) = 0$ and $W(u(x), x) = 0$ a.e. in Ω . Let $\eta \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$ be such that $\int_{\mathbb{R}^N} \eta(z) dz = 1$ and $\text{supp } \eta \subset B_R(0)$. For every $\rho > 0$ set

$$\eta_\rho(z) := \frac{1}{\rho^N} \eta\left(\frac{z}{\rho}\right) \quad \forall z \in \mathbb{R}^N. \quad (2.32)$$

Finally, for every $x \in \mathbb{R}^N$ and every $\varepsilon > 0$ define

$$u_{\rho,\varepsilon}(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta_\rho\left(\frac{y-x}{\varepsilon}\right) u(y) dy = \int_{\mathbb{R}^N} \eta(z) u(x + \varepsilon\rho z) dz = \int_{B_R(0)} \eta(z) u(x + \varepsilon\rho z) dz. \quad (2.33)$$

Then,

$$\begin{aligned} & \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln(\varepsilon)} \left(\|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N \setminus \overline{\Omega}, \mathbb{R}^d)}^q \right) + \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right) \right\} \\ &= \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln(\varepsilon)} \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right) \right\} \\ &= \left(\int_{\mathbb{R}^{N-1}} \frac{2}{(\sqrt{1+|v|^2})^{N+1}} dv \right) \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.34) \end{aligned}$$

Proof. Since $\int_{\mathbb{R}^N} \eta_\rho(z) dz = 1$, applying Proposition 2.1, first for \mathbb{R}^N , then for $\mathbb{R}^N \setminus \overline{\Omega}$, and finally for Ω , yields, for every $\rho > 0$,

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \frac{1}{-\ln(\varepsilon)} \left(\|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N \setminus \overline{\Omega}, \mathbb{R}^d)}^q \right) \\ &= 2D_N \left(\int_{J_u} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) - \int_{J_u \cap (\mathbb{R}^N \setminus \overline{\Omega})} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) \right) \\ &= 2D_N \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y) = \lim_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln(\varepsilon)} \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q \right), \quad (2.35) \end{aligned}$$

where D_N is the constant defined in (2.25). On the other hand, since $W(u(x), x) = 0$ a.e. in Ω and $u \in L^\infty$, by (2.31) we get that

$$\begin{aligned} \left| \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right| &= \left| \frac{1}{\varepsilon} \int_{\Omega} \left(W(u_{\rho,\varepsilon}(x), x) - W(u(x), x) \right) dx \right| \leq C \int_{\mathbb{R}^N} \frac{1}{\varepsilon} |u_{\rho,\varepsilon}(x) - u(x)| dx \\ &\leq C \int_{B_R(0)} |\eta(z)| \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon} |u(x + \varepsilon\rho z) - u(x)| dx \right) dz \\ &= C\rho \int_{B_R(0)} |z| |\eta(z)| \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon\rho|z|} |u(x + \varepsilon\rho z) - u(x)| dx \right) dz, \quad (2.36) \end{aligned}$$

for some constant $C > 0$, independent of ε and ρ . Thus, taking into account the following well known uniform bound from the theory of BV functions,

$$\int_{\mathbb{R}^N} \frac{1}{\rho\varepsilon|z|} |u(x + \rho\varepsilon z) - u(x)| dx \leq C_0 \|Du\|(\mathbb{R}^N) \quad \forall z \in \mathbb{R}^N, \forall \rho, \varepsilon > 0, \quad (2.37)$$

we obtain that

$$\limsup_{\varepsilon \rightarrow 0^+} \left| \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right| \leq CC_0 \|Du\|(\mathbb{R}^N) \rho \int_{B_R(0)} |z| |\eta(z)| dz = O(\rho). \quad (2.38)$$

By (2.38) and (2.35) we finally derive (2.34). \square

Proof of Theorem 1.3. Let η, η_ρ and $u_{\rho,\varepsilon}$ be defined as in Corollary 2.1. Then $u_{\rho,\varepsilon} \in C^\infty(\mathbb{R}^N, \mathbb{R}^d) \cap W^{1,1}(\mathbb{R}^N, \mathbb{R}^d) \cap W^{1,\infty}(\mathbb{R}^N, \mathbb{R}^d)$ and by Corollary 2.1 we have

$$\begin{aligned} & \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln(\varepsilon)} \left(\|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N \setminus \bar{\Omega}, \mathbb{R}^d)}^q \right) + \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right) \right\} \\ &= \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln \varepsilon} \|u_{\rho,\varepsilon}\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_{\Omega} W(u_{\rho,\varepsilon}(x), x) dx \right) \right\} \\ &= \left(\int_{\mathbb{R}^{N-1}} \frac{2}{(\sqrt{1+|v|^2})^{N+1}} dv \right) \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.39) \end{aligned}$$

Clearly, for every $x \in \mathbb{R}^N$ we have $A \cdot \nabla u_{\rho,\varepsilon}(x) = 0$ and $u_{\rho,\varepsilon}(x) \rightarrow u(x)$ strongly in $L^p(\mathbb{R}^N, \mathbb{R}^d)$ as $\varepsilon \rightarrow 0^+$ for every fixed ρ and p . Therefore, by the above and by (2.39) we can complete the proof of the first assertion of the theorem using a standard diagonal argument.

It remains to show the second assertion of the theorem, namely, that in the case $A \equiv 0$ we can construct ψ_ε satisfying the additional condition (1.15). Let $\varphi \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$ be such that $\int_{\Omega} \varphi(x) dx = 1$. Define

$$\tilde{u}_{\rho,\varepsilon}(x) := u_{\rho,\varepsilon}(x) - \varphi(x) c_{\varepsilon,\rho}, \quad (2.40)$$

where

$$c_{\varepsilon,\rho} := \int_{\Omega} u_{\rho,\varepsilon}(y) dy - \int_{\Omega} u(y) dy. \quad (2.41)$$

In particular,

$$\int_{\Omega} \tilde{u}_{\rho,\varepsilon}(x) dx = \int_{\Omega} u(x) dx, \quad (2.42)$$

and $\lim_{\varepsilon \rightarrow 0^+} c_{\varepsilon,\rho} = 0$. On the other hand, since $W(u(x), x) = 0$ a.e. in Ω , $W(b, x)$ is nonnegative and $W(b, x)$ is differentiable with respect to the b variable, we have

$$\nabla_b W(u(x), x) = 0 \quad \text{a.e. in } \Omega. \quad (2.43)$$

Thus, since $u \in L^\infty$, by (2.40) we get that

$$\begin{aligned}
& \left| \frac{1}{\varepsilon} \int_{\Omega} \left(W(\tilde{u}_{\rho,\varepsilon}(x), x) - W(u_{\rho,\varepsilon}(x), x) \right) dx \right| = \left| \frac{c_{\varepsilon,\rho}}{\varepsilon} \cdot \int_0^1 \int_{\Omega} \nabla_b W \left(u_{\rho,\varepsilon}(x) - s\varphi(x)c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right| \\
& \leq C \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon} |u_{\rho,\varepsilon}(x) - u(x)| dx \right) \left| \int_0^1 \int_{\Omega} \nabla_b W \left(u_{\rho,\varepsilon}(x) - s\varphi(x)c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right| \\
& \leq C \left(\int_{B_R(0)} |\eta(z)| \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon} |u(x + \varepsilon\rho z) - u(x)| dx \right) dz \right) \times \\
& \quad \times \left| \int_0^1 \int_{\Omega} \nabla_b W \left(u_{\rho,\varepsilon}(x) - s\varphi(x)c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right| \\
& = C\rho \left(\int_{B_R(0)} |z| |\eta(z)| \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon\rho|z|} |u(x + \varepsilon\rho z) - u(x)| dx \right) dz \right) \times \\
& \quad \times \left| \int_0^1 \int_{\Omega} \nabla_b W \left(u_{\rho,\varepsilon}(x) - s\varphi(x)c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right|. \quad (2.44)
\end{aligned}$$

On the other hand, taking into account (2.37) and using the Dominated Convergence Theorem and (2.43), we obtain that

$$\begin{aligned}
& \limsup_{\varepsilon \rightarrow 0^+} \left(\int_{B_R(0)} |z| |\eta(z)| \left(\int_{\mathbb{R}^N} \frac{1}{\varepsilon\rho|z|} |u(x + \varepsilon\rho z) - u(x)| dx \right) dz \right) \times \\
& \times \left| \int_0^1 \int_{\Omega} \nabla_b W \left(u_{\rho,\varepsilon}(x) - s\varphi(x)c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right| \leq C_0 \left(\|Du\|(\mathbb{R}^n) \right) \left(\int_{B_R(0)} |z| |\eta(z)| dz \right) \times \\
& \quad \times \left| \int_0^1 \int_{\Omega} \nabla_b W \left(\lim_{\varepsilon \rightarrow 0^+} u_{\rho,\varepsilon}(x) - s\varphi(x) \lim_{\varepsilon \rightarrow 0^+} c_{\varepsilon,\rho}, x \right) \varphi(x) dx ds \right| \\
& = C_0 \left(\|Du\|(\mathbb{R}^n) \right) \left(\int_{B_R(0)} |z| |\eta(z)| dz \right) \left| \int_{\Omega} \nabla_b W \left(u(x), x \right) \varphi(x) dx \right| = 0. \quad (2.45)
\end{aligned}$$

Using (2.45) in (2.44) yields

$$\limsup_{\varepsilon \rightarrow 0^+} \left| \frac{1}{\varepsilon} \int_{\Omega} \left(W(\tilde{u}_{\rho,\varepsilon}(x), x) - W(u_{\rho,\varepsilon}(x), x) \right) dx \right| = 0. \quad (2.46)$$

Plugging (2.46) into (2.39) we get that

$$\begin{aligned}
& \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln(\varepsilon)} \left(\|\tilde{u}_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N, \mathbb{R}^d)}^q - \|\tilde{u}_{\rho,\varepsilon}\|_{W^{1/q,q}(\mathbb{R}^N \setminus \bar{\Omega}, \mathbb{R}^d)}^q \right) + \frac{1}{\varepsilon} \int_{\Omega} W(\tilde{u}_{\rho,\varepsilon}(x), x) dx \right) \right\} \\
& = \lim_{\rho \rightarrow 0^+} \left\{ \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{1}{-\ln \varepsilon} \|\tilde{u}_{\rho,\varepsilon}\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q + \frac{1}{\varepsilon} \int_{\Omega} W(\tilde{u}_{\rho,\varepsilon}(x), x) dx \right) \right\} \\
& = \left(\int_{\mathbb{R}^{N-1}} \frac{2}{(\sqrt{1+|v|^2})^{N+1}} dv \right) \int_{J_u \cap \Omega} |u^+(y) - u^-(y)|^q d\mathcal{H}^{N-1}(y). \quad (2.47)
\end{aligned}$$

Moreover, $\tilde{u}_{\rho,\varepsilon} \rightarrow u$ strongly in $L^p(\mathbb{R}^N, \mathbb{R}^d)$ as $\varepsilon \rightarrow 0^+$ for every fixed ρ and p . Therefore, by the above and (2.47) we complete again the proof by a standard diagonal argument. \square

The next lemma is needed for the proof of Theorem 1.2 (in the general case $\eta \in W^{1,1}$).

Lemma 2.1. *Let $\Omega \subset \mathbb{R}^N$ be an open set and let $u \in BV(\mathbb{R}^N, \mathbb{R}^d) \cap L^\infty(\mathbb{R}^N, \mathbb{R}^d)$. For $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$, every $x \in \mathbb{R}^N$ and every $\varepsilon > 0$ define*

$$u_\varepsilon(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy = \int_{\mathbb{R}^N} \eta(z) u(x + \varepsilon z) dz. \quad (2.48)$$

Then, for every $q > 1$ and for every $\varepsilon \in (0, 1)$ we have

$$\begin{aligned} \frac{1}{\omega_{N-1} |\ln \varepsilon|} \int_{\Omega} \left(\int_{\Omega} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq \frac{2^q \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q}{|\ln \varepsilon|} \\ &+ \frac{(3\|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N)}{(q-1) |\ln \varepsilon|} \\ &+ (3\|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N), \end{aligned} \quad (2.49)$$

where ω_{N-1} denotes the surface area of the unit ball in \mathbb{R}^N .

Proof. Assume first that $\eta(z) \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$. Then, by (2.48) we have

$$\varepsilon \nabla u_\varepsilon(x) = -\frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \nabla \eta\left(\frac{y-x}{\varepsilon}\right) u(y) dy = -\int_{\mathbb{R}^N} \nabla \eta(z) u(x + \varepsilon z) dz. \quad (2.50)$$

By (2.48) and (2.50) we get that

$$\begin{aligned} \|u_\varepsilon\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} + \|\varepsilon \nabla u_\varepsilon\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} &\leq \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})} \quad \text{and} \\ \|u_\varepsilon\|_{L^q(\mathbb{R}^N, \mathbb{R}^d)}^q &\leq \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q \quad \forall \varepsilon > 0, \forall q \in [1, +\infty). \end{aligned} \quad (2.51)$$

Next, for every $\varepsilon \in (0, 1)$ we have

$$\begin{aligned} \int_{\Omega} \left(\int_{\Omega} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx = \\ &\int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|^{N+1}} dy \right) dx = \int_{\mathbb{R}^N} \left(\int_{B_\varepsilon(0)} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|^{N+1}} dy \right) dx \\ &+ \int_{\mathbb{R}^N} \left(\int_{B_1(0) \setminus B_\varepsilon(0)} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|^{N+1}} dy \right) dx + \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N \setminus B_1(0)} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|^{N+1}} dy \right) dx \\ &= \int_{B_\varepsilon(0)} \frac{1}{|y|^{N+1-q}} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|^q} dx \right) dy \\ &+ \int_{B_1(0) \setminus B_\varepsilon(0)} \frac{1}{|y|^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|^q}{|y|} dx \right) dy \\ &+ \int_{\mathbb{R}^N \setminus B_1(0)} \frac{1}{|y|^{N+1}} \left(\int_{\mathbb{R}^N} |u_\varepsilon(x+y) - u_\varepsilon(x)|^q dx \right) dy. \end{aligned} \quad (2.52)$$

On the other hand, (2.51) yields

$$|u_\varepsilon(x+y) - u_\varepsilon(x)| + \frac{\varepsilon |u_\varepsilon(x+y) - u_\varepsilon(x)|}{|x-y|} \leq 3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})} \quad \forall \varepsilon > 0, \forall x, y \in \mathbb{R}^N. \quad (2.53)$$

Thus, inserting (2.53) into (2.52) we deduce that

$$\begin{aligned} \int_{\Omega} \left(\int_{\Omega} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq 2^q \|u_\varepsilon\|_{L^q(\mathbb{R}^N, \mathbb{R}^d)}^q \int_{\mathbb{R}^N \setminus B_1(0)} \frac{dy}{|y|^{N+1}} \\ &+ \frac{(3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1}}{\varepsilon^{q-1}} \int_{B_\varepsilon(0)} \frac{1}{|y|^{N+1-q}} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|}{|y|} dx \right) dy \\ &+ (3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \int_{B_1(0) \setminus B_\varepsilon(0)} \frac{1}{|y|^N} \left(\int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+y) - u_\varepsilon(x)|}{|y|} dx \right) dy. \end{aligned} \quad (2.54)$$

Inserting (2.48) into (2.54) and using the second inequality in (2.51) we infer,

$$\begin{aligned} \int_{\Omega} \left(\int_{\Omega} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq 2^q \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q \int_{\mathbb{R}^N \setminus B_1(0)} \frac{dy}{|y|^{N+1}} \\ &+ \frac{(3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1}}{\varepsilon^{q-1}} \times \\ &\times \int_{B_\varepsilon(0)} \frac{1}{|y|^{N+1-q}} \left(\int_{\mathbb{R}^N} |\eta(z)| \int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+\varepsilon z+y) - u_\varepsilon(x+\varepsilon z)|}{|y|} dx dz \right) dy \\ &+ (3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \times \\ &\times \int_{B_1(0) \setminus B_\varepsilon(0)} \frac{1}{|y|^N} \left(\int_{\mathbb{R}^N} |\eta(z)| \int_{\mathbb{R}^N} \frac{|u_\varepsilon(x+\varepsilon z+y) - u_\varepsilon(x+\varepsilon z)|}{|y|} dx dz \right) dy. \end{aligned} \quad (2.55)$$

Taking into account the following well known uniform bound from the theory of BV functions:

$$\int_{\mathbb{R}^N} \frac{|u(x+\varepsilon z+y) - u(x+\varepsilon z)|}{|y|} dx = \int_{\mathbb{R}^N} \frac{|u(x+y) - u(x)|}{|y|} dx \leq \|Du\|(\mathbb{R}^N) \quad \forall y \in \mathbb{R}^N, \quad (2.56)$$

we rewrite (2.55) as

$$\begin{aligned} \int_{\Omega} \left(\int_{\Omega} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq 2^q \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q \int_{\mathbb{R}^N \setminus B_1(0)} \frac{dy}{|y|^{N+1}} \\ &+ \frac{(3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1}}{\varepsilon^{q-1}} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N) \int_{B_\varepsilon(0)} \frac{dy}{|y|^{N+1-q}} \\ &+ (3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N) \int_{B_1(0) \setminus B_\varepsilon(0)} \frac{dy}{|y|^N}. \end{aligned} \quad (2.57)$$

Computing the integrals on the R.H.S. of (2.57) yields (2.49) in the case $\eta \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$.

Next consider the general case $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$. Thanks to the density of $C_c^\infty(\mathbb{R}^N, \mathbb{R})$ in $W^{1,1}(\mathbb{R}^N, \mathbb{R})$, there exists a sequence $\{\eta_n\}_{n=1}^\infty \subset C_c^\infty(\mathbb{R}^N, \mathbb{R})$ such that

$$\lim_{n \rightarrow +\infty} \|\eta_n - \eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})} = 0. \quad (2.58)$$

Thus if we define

$$u_{n,\varepsilon}(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta_n \left(\frac{y-x}{\varepsilon} \right) u(y) dy = \int_{\mathbb{R}^N} \eta_n(z) u(x + \varepsilon z) dz, \quad (2.59)$$

then

$$\lim_{n \rightarrow +\infty} u_{n,\varepsilon}(x) = u_\varepsilon(x) \quad \forall x \in \mathbb{R}^N, \quad \forall \varepsilon > 0. \quad (2.60)$$

On the other hand, since we proved (2.49) for the case $\eta_n \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$, for every $q > 1$, for every $n = 1, 2, \dots$ and for every $\varepsilon \in (0, 1)$ we have:

$$\begin{aligned} \frac{1}{\omega_{N-1} |\ln \varepsilon|} \int_{\Omega} \left(\int_{\Omega} \frac{|u_{n,\varepsilon}(x) - u_{n,\varepsilon}(y)|^q}{|x-y|^{N+1}} dy \right) dx &\leq \frac{2^q \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta_n\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q}{|\ln \varepsilon|} \\ &+ \frac{(3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta_n\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta_n\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N)}{(q-1) |\ln \varepsilon|} \\ &+ (3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta_n\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta_n\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N). \end{aligned} \quad (2.61)$$

Letting n go to infinity in (2.61), using (2.58) in the R.H.S. and (2.60) together with Fatou's Lemma in the L.H.S., we obtain (2.49) in the general case $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$. \square

Proof of Theorem 1.2. In the case $\eta \in C_c^\infty(\mathbb{R}^N, \mathbb{R})$ the result follows by Proposition 2.1. Next consider the general case $\eta \in W^{1,1}(\mathbb{R}^N, \mathbb{R})$. As before, by the density of $C_c^\infty(\mathbb{R}^N, \mathbb{R})$ in $W^{1,1}(\mathbb{R}^N, \mathbb{R})$, there exists a sequence $\{\eta_n\}_{n=1}^\infty \subset C_c^\infty(\mathbb{R}^N, \mathbb{R})$ such that

$$\lim_{n \rightarrow +\infty} \|\eta_n - \eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})} = 0. \quad (2.62)$$

Next, as before, define

$$u_{n,\varepsilon}(x) := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \eta_n \left(\frac{y-x}{\varepsilon} \right) u(y) dy = \int_{\mathbb{R}^N} \eta_n(z) u(x + \varepsilon z) dz. \quad (2.63)$$

Defining $u_{n,\varepsilon}$ as in (2.59) we get by Proposition 2.1, for all $n \geq 1$ (see (2.25)),

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|} \|u_{n,\varepsilon}\|_{W^{1/q,q}(\Omega, \mathbb{R}^d)}^q = 2D_N \left| \int_{\mathbb{R}^N} \eta_n(z) dz \right|^q \int_{J_u \cap \Omega} |u^+(x) - u^-(x)|^q d\mathcal{H}^{N-1}(x) := L_n, \quad (2.64)$$

and then

$$\lim_{n \rightarrow \infty} L_n = \bar{L} := 2D_N \left| \int_{\mathbb{R}^N} \eta(z) dz \right|^q \int_{J_u \cap \Omega} |u^+(x) - u^-(x)|^q d\mathcal{H}^{N-1}(x). \quad (2.65)$$

On the other hand, by Lemma 2.1, for all $n \geq 1$ and every $\varepsilon \in (0, 1/e)$ we have

$$\begin{aligned} \frac{1}{\omega_{N-1} |\ln \varepsilon|} \int_{\Omega} \left(\int_{\Omega} \frac{1}{|x-y|^{N+1}} \left| (u_{n,\varepsilon}(x) - u_{n,\varepsilon}(y)) - (u_\varepsilon(x) - u_\varepsilon(y)) \right|^q dy \right) dx &= \\ \frac{1}{\omega_{N-1} |\ln \varepsilon|} \int_{\Omega} \left(\int_{\Omega} \frac{1}{|x-y|^{N+1}} \left| (u_{n,\varepsilon}(x) - u_\varepsilon(x)) - (u_{n,\varepsilon}(y) - u_\varepsilon(y)) \right|^q dy \right) dx & \\ \leq 2^q \|u\|_{L^1(\mathbb{R}^N, \mathbb{R}^d)} \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)}^{q-1} \|\eta_n - \eta\|_{L^1(\mathbb{R}^N, \mathbb{R})}^q & \\ + \frac{(3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta_n - \eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta_n - \eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N)}{(q-1)} & \\ + (3 \|u\|_{L^\infty(\mathbb{R}^N, \mathbb{R}^d)} \|\eta_n - \eta\|_{W^{1,1}(\mathbb{R}^N, \mathbb{R})})^{q-1} \|\eta_n - \eta\|_{L^1(\mathbb{R}^N, \mathbb{R})} \|Du\|(\mathbb{R}^N) &:= H_n. \end{aligned} \quad (2.66)$$

Thus, by the triangle inequality we get, for every $n \geq 1$ and every $\varepsilon \in (0, 1/e)$,

$$\frac{1}{|\ln \varepsilon|^{1/q}} \left| \|u_{n,\varepsilon}\|_{W^{1/q,q}} - \|u_\varepsilon\|_{W^{1/q,q}} \right| \leq \frac{\|u_{n,\varepsilon} - u_\varepsilon\|_{W^{1/q,q}}}{|\ln \varepsilon|^{1/q}} \leq (\omega_{N-1} H_n)^{1/q}. \quad (2.67)$$

Then, by (2.67) and (2.64), for all $n \geq 1$ we obtain:

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0^+} \left| \frac{\|u_\varepsilon\|_{W^{1/q,q}}}{|\ln \varepsilon|^{1/q}} - \bar{L}^{1/q} \right| &\leq \limsup_{\varepsilon \rightarrow 0^+} \frac{1}{|\ln \varepsilon|^{1/q}} \left| \|u_{n,\varepsilon}\|_{W^{1/q,q}} - \|u_\varepsilon\|_{W^{1/q,q}} \right| \\ &+ \limsup_{\varepsilon \rightarrow 0^+} \left| \frac{\|u_{n,\varepsilon}\|_{W^{1/q,q}}}{|\ln \varepsilon|^{1/q}} - L_n^{1/q} \right| + |L_n^{1/q} - \bar{L}^{1/q}| \leq (\omega_{N-1} H_n)^{1/q} + 0 + |L_n^{1/q} - \bar{L}^{1/q}|. \end{aligned} \quad (2.68)$$

Letting n go to infinity in (2.68), using (2.65), the definition of \bar{L} in (2.65) and the fact that $\lim_{n \rightarrow +\infty} H_n = 0$, we finally deduce (1.8). \square

3 Appendix: Some known results on BV-spaces

In what follows we present some known definitions and results on BV-spaces; some of them were used in the previous sections. We rely mainly on the book [4] by Ambrosio, Fusco and Pallara.

Definition 3.1. Let Ω be a domain in \mathbb{R}^N and let $f \in L^1(\Omega, \mathbb{R}^m)$. We say that $f \in BV(\Omega, \mathbb{R}^m)$ if the following quantity is finite:

$$\int_{\Omega} |Df| := \sup \left\{ \int_{\Omega} f \cdot \operatorname{div} \varphi \, dx : \varphi \in C_c^1(\Omega, \mathbb{R}^{m \times N}), |\varphi(x)| \leq 1 \, \forall x \right\}.$$

Definition 3.2. Let Ω be a domain in \mathbb{R}^N . Consider a function $f \in L^1_{loc}(\Omega, \mathbb{R}^m)$ and a point $x \in \Omega$.

i) We say that x is an *approximate continuity point* of f if there exists $z \in \mathbb{R}^m$ such that

$$\lim_{\rho \rightarrow 0^+} \frac{\int_{B_\rho(x)} |f(y) - z| \, dy}{\rho^N} = 0.$$

In this case we denote z by $\tilde{f}(x)$. The set of approximate continuity points of f is denoted by G_f .

ii) We say that x is an *approximate jump point* of f if there exist $a, b \in \mathbb{R}^m$ and $\nu \in S^{N-1}$ such that $a \neq b$ and

$$\lim_{\rho \rightarrow 0^+} \frac{\int_{B_\rho(x)} |f(y) - \chi(a, b, \nu)(y)| \, dy}{\rho^N} = 0, \quad (3.1)$$

where $\chi(a, b, \nu)$ is defined by

$$\chi(a, b, \nu)(y) := \begin{cases} b & \text{if } \nu \cdot y < 0, \\ a & \text{if } \nu \cdot y > 0. \end{cases}$$

The triple (a, b, ν) , uniquely determined, up to a permutation of (a, b) and a change of sign of ν , is denoted by $(f^+(x), f^-(x), \nu_f(x))$. We shall call $\nu_f(x)$ the *approximate jump vector* and we shall sometimes write simply $\nu(x)$ if the reference to the function f is clear. The set of approximate jump points is denoted by J_f . A choice of $\nu(x)$ for every $x \in J_f$ determines an orientation of J_f . At an approximate continuity point x , we shall use the convention $f^+(x) = f^-(x) = \tilde{f}(x)$.

Theorem 3.1 (Theorems 3.69 and 3.78 from [4]). *Consider an open set $\Omega \subset \mathbb{R}^N$ and $f \in BV(\Omega, \mathbb{R}^m)$. Then:*

- i) \mathcal{H}^{N-1} -a.e. point in $\Omega \setminus J_f$ is a point of approximate continuity of f .
- ii) The set J_f is σ - \mathcal{H}^{N-1} -rectifiable Borel set, oriented by $\nu(x)$. I.e., the set J_f is \mathcal{H}^{N-1} σ -finite, there exist countably many C^1 hypersurfaces $\{S_k\}_{k=1}^\infty$ such that $\mathcal{H}^{N-1}\left(J_f \setminus \bigcup_{k=1}^\infty S_k\right) = 0$, and for \mathcal{H}^{N-1} -a.e. $x \in J_f \cap S_k$, the approximate jump vector $\nu(x)$ is normal to S_k at the point x .
- iii) $[(f^+ - f^-) \otimes \nu_f](x) \in L^1(J_f, d\mathcal{H}^{N-1})$.

Theorem 3.2 (Theorems 3.92 and 3.78 from [4]). *Consider an open set $\Omega \subset \mathbb{R}^N$ and $f \in BV(\Omega, \mathbb{R}^m)$. Then, the distributional gradient Df can be decomposed as a sum of two Borel regular finite matrix-valued measures μ_f and $D^j f$ on Ω ,*

$$Df = \mu_f + D^j f,$$

where

$$D^j f = (f^+ - f^-) \otimes \nu_f \mathcal{H}^{N-1} \llcorner J_f$$

is called the jump part of Df and

$$\mu_f = (D^a f + D^c f)$$

is a sum of the absolutely continuous and the Cantor parts of Df . The two parts μ_f and $D^j f$ are mutually singular to each other. Moreover, $\mu_f(B) = 0$ for any Borel set $B \subset \Omega$ which is \mathcal{H}^{N-1} σ -finite.

References

- [1] G. Alberti, G. Bouchitté and P. Seppecher, *Un résultat de perturbations singulières avec la norme $H^{1/2}$* , C. R. Acad. Sci. Paris, **319**, Série I (1994), 333–338.
- [2] L. Ambrosio, *Metric space valued functions of bounded variation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **17** (1990), 439–478.
- [3] L. Ambrosio, C. De Lellis and C. Mantegazza, *Line energies for gradient vector fields in the plane*, Calc. Var. PDE **9** (1999), 327–355.

- [4] L. Ambrosio, N. Fusco and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Mathematical Monographs. Oxford University Press, New York, 2000.
- [5] P. Aviles and Y. Giga, *A mathematical problem related to the physical theory of liquid crystal configurations*, Proc. Centre Math. Anal. Austral. Nat. Univ. **12** (1987), 1–16.
- [6] P. Aviles and Y. Giga, *On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields*, Proc. Roy. Soc. Edinburgh Sect. A **129** (1999), 1–17.
- [7] F. Hernández, *Properties of a Hilbertian Norm for Perimeter*, to appear in Pure Appl. Funct. Anal., <https://arxiv.org/abs/1709.08262>.
- [8] A. Figalli and D. Jerison, *How to recognize convexity of a set from its marginals*, J. Funct. Anal., **266** (2014), 1685–1701.
- [9] L. Modica, *The gradient theory of phase transitions and the minimal interface criterion*, Arch. Rational Mech. Anal. **98** (1987), 123–142.
- [10] L. Modica and S. Mortola, *Un esempio di Γ^- -convergenza*, Boll. Un. Mat. Ital. B **14** (1977), 285–299.
- [11] L. Modica and S. Mortola, *Il limite nella Γ -convergenza di una famiglia di funzionali ellittici*, Boll. Un. Mat. Ital. A **14** (1977), 526–529.
- [12] G. Palatucci, O. Savin, E. Valdinoci, *Local and global minimizers for a variational energy involving a fractional norm*, Ann. Mat. Pura Appl. **192** (2013), no. 4, 673–718.
- [13] A. Poliakovsky, *Upper bounds for singular perturbation problems involving gradient fields*, J. Eur. Math. Soc. **9** (2007), 1–43.
- [14] A. Poliakovsky, *Jump detection in Besov spaces via a new BBM formula. Applications to Aviles-Giga type functionals*, Commun. Contemp. Math. **20** (2018), no. 7, 1750096, 36 pp.
- [15] O. Savin and E. Valdinoci, *Γ -convergence for nonlocal phase transitions*, Ann. Inst. H. Poincaré Anal. Non Linéaire **29** (2012), 479–500.
- [16] P. Sternberg, *The effect of a singular perturbation on nonconvex variational problems*, Arch. Rational Mech. Anal. **101** (1988), 209–260.