

Charmonium resonances from 2+1 flavor CLS lattices

Sara Collins^a, Daniel Mohler^{*c,d}, M. Padmanath^a, Stefano Piemonte^{†a}, Sasa Prelovsek^b, Simon Weishäupl^a











^a Institute for Theoretical Physics, University of Regensburg, 93040 Regensburg, Germany

^b Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia

^c Helmholtz-Institut Mainz, 55099 Mainz, Germany

^d Johannes Gutenberg Universität Mainz, 55099 Mainz, Germany

E-mail: stefano.piemonte@ur.de

The 36th Annual International Symposium on Lattice Field Theory - LATTICE2018

22-28 July, 2018

Michigan State University, East Lansing, Michigan, USA.

* 
† 

1. Introduction

Charmonium resonances from 2+1 flavor CLS lattices

ab-initio [1]

Charmonium resonances from 2+1 flavor CLS lattices

2]

[6]

2. Charmonium states in the single-hadron approach

Charmonium states in the single-hadron approach

$$C_{ij}(t) = \langle O_i(t) O_j^\dagger(0) \rangle$$

O_i

Charmonium states in the single-hadron approach

Charmonium states in the single-hadron approach

$$N_f = 2 + 1$$

[4, 5]

$a =$

$$m_\pi \simeq 280 \text{ MeV}$$

$$m_K \simeq$$

$$\times 24^3$$

m_c

D

3. Scattering in a finite box: Lüscher formalism

Scattering in a finite box: Lüscher formalism




Figure 1: Resonance $P(1,0,0)$ fit

$P = (1,0,0)$ (10-6) d

$P = (1,1,0)$ (10-7) f

of δ is fitted with the function $p^{2l+1} \cot(\delta)$. The value of the fit is $\chi^2/d.f. = 2.2$, which is consistent with the $\chi^2/d.f. = 3.1$ of the K -invariant fit.

cc-p 10-6

K -in

3.1 Vector channel

The J/ψ resonance is fitted with the function $\psi(4040), \psi(3700)$ and $\bar{D}D$ states. The fit is shown in Figure 1(b) for the U101 lattice with $\kappa_c = 0.12522$.

J/ψ resonance

$\Psi_{2S}, \psi(3700)$ d

$\bar{D}D$ states

$\bar{D}D$

fit

$$k^2 = s/4 - m_D^2$$

(10-6) d

$\kappa_c = 0.12522$. Fit

$$\frac{k^3 \cot(\delta)}{\sqrt{s}} = A + Bs + Cs^2. \quad (3.1)$$

The J/ψ resonance is fitted with the function $\chi^2/d.f. = 2.2$, which is consistent with the $\chi^2/d.f. = 3.1$ of the K -invariant fit.

J/ψ resonance fit $\chi^2/d.f. = 2.2$, which is consistent with the $\chi^2/d.f. = 3.1$ of the K -invariant fit.

(10-7) f ansatz

$\chi^2/d.f. = 3.1$.




Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

$$k^3 \cot(\delta) = -k^2 \sqrt{k^2} \quad (3.2)$$

Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2


Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2


Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 2: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2



(a) I



(b) I

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2

Figure 3: $\frac{ap^3 \cot(\delta)}{\sqrt{s}}$ vs ak^2




Figure 4: Ω with $\kappa_c = 0.12315$ at $\beta = 1.9$

$\Psi(2S)$, $\psi(3700)$ 3. χ_{c0}
 $\psi(3700)$ 8]

3.2 Scalar channel

χ_{c0} , $X(3950)$ $\chi_{c0}(380)$
 $X(3950)$ $\bar{D}D$
 $\bar{D}_s D_s$ $X(3950)$
 $\bar{D}D$ $\bar{D}_s D_s$
 $J/\Psi - \omega$ $\bar{D}^* D^*$ ++
 K^{-1} s. ansatz

$$\tilde{K}^{-1} = \begin{bmatrix} c_{11}s + b_{11} & b_{12} \\ b_{12} & b_{22} \end{bmatrix}$$

$\chi^2/df \approx$
 \tilde{K}^{-1} (7). Ω Ω
7]

8

$$\Omega(\mu, A(s)) = \frac{\text{tr}(A)}{\text{tr}((\mu^2 + AA^\dagger)^{\frac{1}{2}})}, \quad (3.3)$$

by $A(s)$ is defined as $A(s) = \langle \mathcal{O}(s) \mathcal{O}^\dagger(0) \rangle$ with $\mathcal{O}(s) = \sum_{\mathbf{x}} \psi(\mathbf{x}, s) \bar{\psi}(\mathbf{x}, 0)$ and $\psi(\mathbf{x}, s) = \sum_{\mathbf{y}} U(\mathbf{x}, \mathbf{y}, s) \psi(\mathbf{y}, 0)$.

4. Results




Figure 5: $|T_{11}|$ for $\bar{D}D$ and $D_S D_S$ channels. The vertical axis is $|T_{11}|$ and the horizontal axes are $\text{Re}(s)$ and $\text{Im}(s)$.

$\bar{D}D$ and $D_S D_S$ channels. The vertical axis is $|T_{11}|$ and the horizontal axes are $\text{Re}(s)$ and $\text{Im}(s)$.

The pole positions are determined by the minimum of $|T_{11}|$ in the complex plane. The real part of the pole position is related to the mass of the resonance, and the imaginary part is related to its width.

$$\text{sign}(\text{Im}[k_{m_D}], \text{sign}(\text{Im}[k_{m_{D_S}}])), \quad \text{with} \quad k_{m_i}^2 = s/4 - m_i^2, \quad \text{in } \bar{D}D \text{ and } D_S D_S \text{ channels.}$$

4. Conclusions

The results show that the charmonium resonances are well described by the $\bar{D}D$ and $D_S D_S$ channels. The pole positions are consistent with the experimental data.

$\psi(3700)$ is identified as $\bar{D}D$ and $D_S D_S$.

K is identified as $\bar{D}D$.

arXiv:1512.03443
 hep-lat/1512034

5. Acknowledgements

We thank the organizers of the Workshop on Lattice QCD for their kind invitation to participate in the workshop. This work is supported by the National Natural Science Foundation of China (NSFC) under Grant No. 11465902 and the Strategic Priority Program (PAP) of the Chinese Academy of Sciences (CAS) under Grant No. XDB06000000 (P0035 and P0837).

References

- [1] C. B. Lang, *JHEP* **1509** (2015) 09.
- [2] L. Chang, M. Cheng, H. Jiang, J. He, C. B. Lang, *JHEP* **07** (2012) 126.
- [3] M. Cheng, *JHEP* **105**, 153 (10); *JHEP* **B354**, 531 (19); *JHEP* **M14**, 237 (14).
- [4] M. Cheng *et al.*, *JHEP* **1502** (2015) 043.
- [5] C. B. Lang *et al.*, *JHEP* **94** (2016) 0704501.
- [6] M. Cheng *et al.*, [arXiv:1811.04116](https://arxiv.org/abs/1811.04116) [hep-lat].
- [7] C. B. Lang, *JHEP* **04**, 475072017.
- [8] T. S. Koehn, *JHEP* **03**, 034521 (2017).
- [9] *JHEP*, 112003 (2017).
- [10] *JHEP* **09** (2005), 18002; *JHEP* **101** (2008) 08001; *JHEP* **104** (2010) 02001.
- [11] Z. H. Guo, *JHEP* **115**, n2, 022001 (2015).
- [12] R. K. Choudhury, *JHEP* **140**, 82 (2005).
- [13] S. B. Datta, *JHEP* **14**, 036 (2014); *JHEP* **15**, 036 (2016); *JHEP* **15**, 035 (2016).
- [14] *JHEP* **16**, 361 (2017).
- [15] P. A. Boyle *et al.*, *JHEP* **14**, 021 (2015).