

On a modified Vakhnenko–Parkes equation

SERGEI SAKOVICH

Institute of Physics, National Academy of Sciences of Belarus
sergsako@gmail.com

Abstract

We show that the modified Vakhnenko–Parkes equation, introduced recently by Wazwaz, is an avatar of the sine-Gordon equation.

1 Introduction

Recently, Wazwaz [1] introduced the modified Vakhnenko–Parkes (MVP) equation

$$uu_{xxt} - u_x u_{xt} + u^3 u_t = 0, \quad (1)$$

similar in its form to the original Vakhnenko–Parkes equation [2]

$$uu_{xxt} - u_x u_{xt} + u^2 u_t = 0. \quad (2)$$

Wazwaz [1] showed that the MVP equation (1) passes the Painlevé test for integrability, in the formulation for PDEs [3, 4], and possesses a three-soliton solution. The reliability of the Painlevé test has been empirically verified by analysis of numerous multi-parameter families of nonlinear equations (see, e.g., [5]–[13], to mention a few), while the existence of a three-soliton solution is also generally considered as a clear indicator of integrability. Therefore it is interesting to investigate whether—and in which sense—the MVP equation (1) is integrable.

2 Integrability

The MVP equation (1) is a third-order two-dimensional PDE, therefore its general solution must contain three arbitrary functions depending on one variable each. One of those arbitrary functions is evidently related to the invariance of the MVP equation (1) with respect to the transformation $t \mapsto f(t)$ with any function $f(t)$. In other words, if a function $u(x, t)$ satisfies (1), then $u(x, f(t))$ will do for any $f(t)$. This kind of invariance can be useful to lower the order of a studied equation [14]–[17].

Let us multiply (1) by $2u_{xt}/u^3$ and integrate with respect to x . Then we get

$$\frac{u_{xt}^2}{u^2} + u_t^2 = g(t), \quad (3)$$

where $g(t)$ is an arbitrary function.

If $g(t) \neq 0$ in (3), we can make $g(t) = 1$ by the transformation $t \mapsto f(t)$ with an appropriately chosen function $f(t)$ (of course, we consider complex-valued transformations and variables throughout). Then we get (3) in the form

$$u_{xt} = u(1 - u_t^2)^{1/2}, \quad (4)$$

where we have omitted the \pm sign in the right-hand side because the sign can be changed by $t \mapsto -t$. This equation (4) is related to the sine-Gordon equation

$$v_{xt} = \sin v \quad (5)$$

by the potential transformation

$$u = v_x(x, t). \quad (6)$$

Since the function $g(t)$ in (3) is arbitrary, the corresponding function $f(t)$ in the used transformation $t \mapsto f(t)$ is arbitrary (but non-constant) as well. Consequently, all solutions of the MVP equation (1) which correspond to non-zero $g(t)$ in (3) are given by the expression

$$u = v_x(x, h(t)), \quad (7)$$

where $h(t)$ is an arbitrary non-constant function (inverse to f), and $v(x, t)$ stands for any solution of the sine-Gordon equation (5).

If $g(t) = 0$ in (3), we have

$$u_{xt} \pm iuu_t = 0, \quad (8)$$

where $i^2 = -1$. The potential transformation

$$u = \pm iw_x(x, t) \quad (9)$$

relates this equation (8) to the Liouville equation

$$w_{xt} = \exp w, \quad (10)$$

whose general solution is well known:

$$w = \log \frac{2a_x b_t}{(a + b)^2}, \quad (11)$$

where $a(x)$ and $b(t)$ are arbitrary non-constant functions. Consequently, all solutions of the MVP equation (1) which correspond to $g(t) = 0$ in (3) are given by the expression

$$u = \pm i \left(\frac{a_{xx}}{a_x} - \frac{2a_x}{a + b} \right), \quad (12)$$

where $a(x)$ and $b(t)$ are arbitrary non-constant functions.

At last, when we used the integrating factor $2u_{xt}/u^3$ to obtain (3) from (1), the factor was assumed to be non-zero. If $u_{xt} = 0$, we have $u_t = 0$ from (1), that is

$$u = c(x), \quad (13)$$

where $c(x)$ is an arbitrary function. This class of solutions of (1) is (formally) covered by the class (12) if we set the function $b(t)$ to be a constant.

3 Conclusion

We have shown that any solution of the MVP equation (1) belongs to one of the three classes: (7), (12), and (13). The general solution (7) is determined by the general solution of the sine-Gordon equation and one extra arbitrary function, while the special solutions (12) and (13) are given explicitly. In this sense, the MVP equation (1) is an avatar of the sine-Gordon equation (5). Let us add that the MVP equation appeared also in [14] (see eq. (16) there) as an auxiliary equation, in the transformation way from the cubic Rabelo equation (a.k.a. the short pulse equation) to the sine-Gordon equation.

References

- [1] A.M. Wazwaz, The integrable Vakhnenko–Parkes (VP) and the modified Vakhnenko–Parkes (MVP) equations: multiple real and complex soliton solutions, Chinese J. Phys., DOI:10.1016/j.cjph.2018.11.004
- [2] V.O. Vakhnenko, E.J. Parkes, Approach in theory of nonlinear evolution equations: the Vakhnenko–Parkes equation, Adv. Math. Phys. 2016 (2016) 2916582.
- [3] J. Weiss, M. Tabor, G. Carnevale, The Painlevé property for partial differential equations, J. Math. Phys. 24 (1983) 522–526.
- [4] M. Tabor, Chaos and Integrability in Nonlinear Dynamics: An Introduction, Wiley, New York, 1989.
- [5] H. Harada, S. Oishi, A new approach to completely integrable partial differential equations by means of the singularity analysis, J. Phys. Soc. Jpn. 54 (1985) 51–56.
- [6] A. Karasu-Kalkanlı, Painlevé classification of coupled Korteweg–de Vries systems, J. Math. Phys. 38 (1997) 3616–3622.
- [7] S.Yu. Sakovich, Painlevé analysis of a higher-order nonlinear Schrödinger equation, J. Phys. Soc. Jpn. 66 (1997) 2527–2529.
- [8] S.Yu. Sakovich, Coupled KdV equations of Hirota–Satsuma type, J. Nonlinear Math. Phys. 6 (1999) 255–262; arXiv:solv-int/9901005.
- [9] S.Yu. Sakovich, Addendum to: Coupled KdV equations of Hirota–Satsuma type, J. Nonlinear Math. Phys. 8 (2001) 311–312; arXiv:nlin/0104072.
- [10] S.Yu. Sakovich, T. Tsuchida, Symmetrically coupled higher-order nonlinear Schrödinger equations: singularity analysis and integrability, J. Phys. A 33 (2000) 7217–7226; arXiv:nlin/0006004.
- [11] A. Karasu-Kalkanlı, A. Karasu, S.Yu. Sakovich, Integrability of a generalized Ito system: the Painlevé test, J. Phys. Soc. Jpn. 70 (2001) 1165–1166; arXiv:nlin/0102030.
- [12] A. Karasu-Kalkanlı, A. Karasu, A. Sakovich, S. Sakovich, R. Turhan, A new integrable generalization of the Korteweg–de Vries equation, J. Math. Phys. 49 (2008) 073516; arXiv:0708.3247.
- [13] G.Q. Xu, The integrability for a generalized seventh-order KdV equation: Painlevé property, soliton solutions, Lax pairs and conservation laws, Phys. Scr. 89 (2014) 125201.

- [14] A. Sakovich, S. Sakovich, On transformations of the Rabelo equations, SIGMA 3 (2007) 086; arXiv:0705.2889.
- [15] S. Sakovich, Transformation and integrability of a generalized short pulse equation, Commun. Nonlinear Sci. Numer. Simulat. 39 (2016) 21–28; arXiv:1510.08822.
- [16] S. Sakovich, Integrability of a generalized short pulse equation revisited, Res. Appl. Math. 2 (2018) 101272; arXiv:1612.03105.
- [17] S. Sakovich, On a new avatar of the sine-Gordon equation, Nonlinear Phenom. Complex Syst. 21 (2018) 62–68; arXiv:1703.04678.