

Characterization of the Most Probable Transition Paths of Stochastic Dynamical Systems with Stable Lévy Noise

Yuanfei Huang¹, Ying Chao¹, Shenglan Yuan¹, Jinqiao Duan^{2,*}

¹School of Mathematics and Statistics & Center for Mathematical Sciences & Hubei Key Laboratory of Engineering Modeling and Scientific Computing, Huazhong University of Science and Technology, Wuhan 430074, China

²Department of Applied Mathematics, Illinois Institute of Technology, Chicago, IL 60616, USA

*Author to whom correspondence should be addressed: duan@iit.edu

E-mail: yfhuang@hust.edu.cn, yingchao1993@hust.edu.cn, shanglanyuan@hust.edu.cn, duan@iit.edu.

Abstract. This work is devoted to the investigation of the most probable transition path for stochastic dynamical systems driven by either symmetric α -stable Lévy motion ($0 < \alpha < 1$) or Brownian motion. For stochastic dynamical systems with Brownian motion, minimizing an action functional is a general method to determine the most probable transition path. We have developed a method based on path integrals to obtain the most probable transition path of stochastic dynamical systems with symmetric α -stable Lévy motion or Brownian motion, and the most probable path can be characterized by a deterministic dynamical system.

Keywords: Symmetric α -stable Lévy motions; Stochastic differential equations; Most probable transition path; Path integrals.

1. Introduction

Stochastic differential equations (SDEs) have been widely used to describe complex phenomena in physical, biological, and engineering systems. Transition phenomena between dynamically significant states occur in nonlinear systems under random fluctuations. Hence a practical problem is that given a stochastic dynamical system, how to capture the transition behavior between two metastable states, and then how to determine the most probable transition path. This subject has been the research topic by a number of authors [4, 12, 22, 18, 6, 11, 8].

In this paper, we consider the following SDE in the state space \mathbb{R}^k :

$$dX_t = b(X_t)dt + dL_t, \quad T_0 \leq t \leq T_f, \quad X_{T_0} = X_0, \quad (1.1)$$

where L_t is a k -dimensional symmetric α -stable (non-Gaussian) Lévy motion in the probability space (Ω, \mathbb{P}) . The solution process X_t uniquely exists under appropriate

conditions on the drift term $b(x) : \mathbb{R}^k \rightarrow \mathbb{R}^k$ (see the next section). Moreover, T_0, T_f are the initial and final time instants, respectively. For simplicity, we first consider the one dimensional case ($k = 1$) and will extend to higher dimensional cases ($k > 1$) in Section 3.

We will also compare with the following one dimensional SDE system with (Gaussian) Brownian motion B_t :

$$dX_t = b(X_t)dt + dB_t, \quad T_0 \leq t \leq T_f, \quad X_{T_0} = X_0. \quad (1.2)$$

The Onsager-Machlup function [4] and *Path integrals* [12] are two methods to study the most probable transition path of this system (1.2). The central points of these two methods were to express the transition probability (density) function of a diffusion process by means of a functional integral over paths of the process. That is for the solution process X_t , with initial time and position (T_0, X_0) and final time and position (T_f, X_f) , the transition probability density $p(X_f, T_f | X_0, T_0)$ (or denoted by $p(X_f | X_0)$) is expressed as a path integral

$$p(X_f, T_f | X_0, T_0) = \int_{\substack{x_{T_0} = X_0 \\ x_{T_f} = X_f}} \mathcal{D}x \exp\left\{-\frac{1}{2} \int_{T_0}^{T_f} [(\dot{x}_s - b(x_s))^2 + b'(x_s)] ds\right\}, \quad (1.3)$$

where $\mathcal{L}(x, \dot{x}) = \frac{1}{2}[(\dot{x} - b(x))^2 + b'(x)]$ is called the Lagrangian of (1.1). In Onsager-Machlup's method, $OM(x, \dot{x}) = (\dot{x} - b(x))^2 + b'(x)$ is called the *OM* function. When the path x_t is restricted in continuous functions mapping from $[T_0, T_f]$ into \mathbb{R} , the exponent $S(x) = -\frac{1}{2} \int_{T_0}^{T_f} [(\dot{x} - b(x))^2 + b'(x)] ds$ is called the Onsager-Machlup action functional. Hence finding the most probable transition path is to find a path x_t such that the Lagrangian (OM function or the action functional) to be minimum, which is called the least action principle. This leads to the Euler-Lagrangian equation by means of a variational principle when the path restricted in twice differentiable functions. For more details of *Path integrals* and applications, see [21, 9, 24, 2, 25] and references therein.

In this present paper, we will determine the most probable transition path for the stochastic system with non-Gaussian noise (1.1). The situation is different from the Gaussian case (1.2). If we try to get the exponential form (containing the action functional) for the transition probability density function for the transition path as in the Gaussian case, we need to use the Fourier transformation of the probability density [15, 5] (or characteristic function). But this brings the Fourier integral into the density function, and the essential difficulty is not solved. Instead, in this paper, we develop a method to characterize the most probable transition path, based on the path integral rather than on the action functional (or the Onsager-Machlup function). This is made possible with a new representation [17] for the transition probability density functions of symmetric Lévy motions in terms of two families of metrics. This representation provides an exponential structure of the transition probability density function [16, 17]. It can be further extended in our case, which will be discussed in Section 2.2.

This paper is organized as follows. In Section 2, we recall some preliminaries. In Section 3, we develop a method to characterize the most probable transition paths for

a stochastic system with symmetric α -stable Lévy motion ($0 < \alpha < 1$) or Brownian motion. In Section 4, we extend the results of Section 3 to higher dimensional cases. Finally, in Section 5, we present some examples to illustrate our results.

2. Preliminaries

2.1. Lévy motions

We recall some basic facts about 1-dimensional Lévy motions (or Lévy processes) [3, 10].

Definition 1. *A stochastic process L_t is a Lévy process if*

- (i) $L_0=0$ (a.s.);
- (ii) L_t has independent increments and stationary increments; and
- (iii) L_t has stochastically continuous sample paths, i.e., for every $s \geq 0$, $L(t) \rightarrow L(s)$ in probability, as $t \rightarrow s$.

A Lévy process L_t taking values in \mathbb{R} is characterized by a drift term $\eta \in \mathbb{R}$, a non-negative variance Q and a Borel measure ν defined on $\mathbb{R} \setminus \{0\}$. (η, Q, ν) is called the generating triplet of the Lévy motion L_t . Moreover, the Lévy-Itô decomposition for L_t as follows:

$$L_t = \eta t + B_Q(t) + \int_{|z|<1} z \tilde{N}(t, dz) + \int_{|z|\geq 1} z N(t, dz), \quad (2.4)$$

where $N(dt, dz)$ is the Poisson random measure, $\tilde{N}(dt, dz) = N(dt, dz) - \nu(dz)dt$ is the compensated Poisson random measure, and $\nu(S) \triangleq \mathbb{E}N(1, S)$, here \mathbb{E} denotes the expectation with respect to the probability \mathbb{P} , and $B_Q(t)$ is a Brownian motion with variance Q . The characteristic function of L_t is given by

$$\mathbb{E}[\exp(iuL_t)] = \exp(-t\psi(u)), \quad u \in \mathbb{R}, \quad (2.5)$$

where the function $\psi : \mathbb{R} \rightarrow \mathbb{C}$ is the characteristic exponent

$$\psi(u) = iu\eta + \frac{1}{2}Qu^2 + \int_{\mathbb{R}\setminus\{0\}} (1 - e^{iuz} + iuz\mathbb{I}_{\{|z|<1\}})\nu(dz). \quad (2.6)$$

The Borel measure ν is called the jump measure.

2.2. Asymptotic properties of the probability density functions of α -stable Lévy motions

From now on, we consider a scalar symmetric α -stable Lévy processes. Recall the standard symmetric α -stable random variable has distribution $S_\alpha(1, 0, 0)$. Here $S_\alpha(\sigma, \beta, \mu)$ is the distribution of a stable random variable, with σ the scale parameter, β the skewness parameter and μ the shift parameter. The corresponding probability

density function $f_\alpha(x)$ can be represented as an infinite series [1, 10, 14]

$$f_\alpha(x) = \begin{cases} \frac{1}{\pi\alpha} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k!} \Gamma(\alpha k + 1) |x|^{-\alpha k} \sin\left(\frac{\alpha k \pi}{2}\right), & x \neq 0, 0 < \alpha < 1, \\ \frac{1}{\pi} \int_0^{\infty} e^{-u^\alpha} du, & x = 0, 0 < \alpha < 1, \\ \frac{1}{\pi(1+x^2)}, & \alpha = 1, \\ \frac{1}{\pi\alpha} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k)!} \Gamma\left(\frac{2k+1}{\alpha}\right) x^{2k}, & 1 < \alpha < 2. \end{cases} \quad (2.7)$$

Recall the probability density function $f_b(x)$ for a Brownian random variable $X \sim N(0, \sigma^2)$ is [10]

$$f_b(x) = \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{x^2}{2\sigma^2}}. \quad (2.8)$$

In [5], it was proved that the transition probability density function $p(x_t|x_0)$ of a symmetric α -stable Lévy process is

$$\begin{aligned} p(x_t|x_0) &= \frac{1}{t^{1/\alpha}} f_\alpha\left(\frac{x_t - x_0}{t^{1/\alpha}}\right) \\ &= \frac{1}{t^{1/\alpha}} \exp[-(-\ln f_\alpha(\frac{x_t - x_0}{t^{1/\alpha}}))] \\ &\triangleq \frac{1}{t^{1/\alpha}} \exp[-\theta_t^\alpha(x_t - x_0)], \end{aligned} \quad (2.9)$$

where $\theta_t^\alpha(\cdot)$ is a function maps $[0, \infty)$ to $[0, \infty)$ for any $\alpha \in (0, 2)$ and $t \in (0, \infty)$. Differentiate $\theta_t^\alpha(x)$ with respect to variable x :

$$\begin{aligned} (\theta_t^\alpha(x))' &= (-\ln f_\alpha(\frac{x}{t^{1/\alpha}}))' \\ &= -\frac{f_\alpha'(\frac{x}{t^{1/\alpha}})}{f_\alpha(\frac{x}{t^{1/\alpha}})} \frac{1}{t^{1/\alpha}}, \end{aligned} \quad (2.10)$$

which shows that $\theta_t^\alpha(\cdot)$ is a strict increase function since $f_\alpha'(\frac{x}{t^{1/\alpha}}) \leq 0$ for symmetric α -stable Lévy random variables. Now we focus on the concavity of the function $\theta_t^\alpha(\cdot)$.

$$\begin{aligned} (\theta_t^\alpha(x))'' &= \left(-\frac{f_\alpha'(\frac{x}{t^{1/\alpha}})}{f_\alpha(\frac{x}{t^{1/\alpha}})} \frac{1}{t^{1/\alpha}}\right)' \\ &= -\frac{f_\alpha(\frac{x}{t^{1/\alpha}}) f_\alpha''(\frac{x}{t^{1/\alpha}}) - (f_\alpha'(\frac{x}{t^{1/\alpha}}))^2}{f_\alpha^2(\frac{x}{t^{1/\alpha}})} \frac{1}{t^{2/\alpha}}, \end{aligned} \quad (2.11)$$

In [1, 3], it was proved that if $X \sim S_\alpha(\sigma, \beta, \mu)$ and $\alpha \in (0, 2)$,

$$\begin{aligned} \lim_{y \rightarrow \infty} y^\alpha \mathbb{P}(X > y) &= C_\alpha \frac{1 + \beta}{2} \sigma^\alpha, \\ \lim_{y \rightarrow \infty} y^\alpha \mathbb{P}(X < -y) &= C_\alpha \frac{1 - \beta}{2} \sigma^\alpha, \end{aligned} \quad (2.12)$$

where

$$C_\alpha = \left(\int_0^\infty x^{-\alpha} \sin(x) dx\right)^{-1}. \quad (2.13)$$




Figure 1: The graphs of function $\theta_1^\alpha(x)$ (with $\alpha = 0.5, 1$ and 1.5): (a) on $[0, 10]$; (b) on $[0, 100]$.

We use this result to study the asymptotic behavior of tail probabilities. For y large enough,

$$\begin{aligned}
 y^\alpha \mathbb{P}(X > y) &= C_\alpha \frac{1 + \beta}{2} \sigma^\alpha, \\
 \Leftrightarrow y^\alpha \int_y^\infty f_\alpha(x) dx &= C_\alpha \frac{1 + \beta}{2} \sigma^\alpha, \\
 \Leftrightarrow f_\alpha(y) &= \alpha C_\alpha \frac{1 + \beta}{2} \sigma^\alpha y^{-\alpha-1} \triangleq C y^{-\alpha-1}, \\
 \Leftrightarrow f_\alpha''(y) f_\alpha(y) - (f_\alpha'(y))^2 &= C^2 (\alpha + 1) y^{-2\alpha-4},
 \end{aligned} \tag{2.14}$$

which means that the asymptotic behavior of tail concavity and convexity of $\theta_t^\alpha(\cdot)$ is concave. And the graphs of $\theta_1^\alpha(\cdot)$ are shown in Figure 1.

For Brownian motion case, similar to the symmetric α -stable Lévy motion case, the corresponding exponent is $\theta_t^b(x) = \frac{x^2}{2t}$, which is convex.

We also notice that

$$\begin{aligned}
 \theta_t^\alpha(x - y) &= \theta_t^\alpha(|x - y|) \triangleq \theta_t^\alpha(x, y), \\
 \theta_t^b(x - y) &= \theta_t^b(|x - y|) \triangleq \theta_t^b(x, y).
 \end{aligned} \tag{2.15}$$

2.3. Conditions for the well-posedness of the system (1.1) and (1.2)

For the system (1.1):

$$dX_t = b(X_t)dt + dL_t, \quad T_0 \leq t \leq T_f, \quad X_{T_0} = X_0,$$

it was proved in [23, 7] that if $b(x)$ is locally Lipschitz continuous function and satisfies “one sided linear growth” condition in the following sense:

- **C1 (Locally Lipschitz condition)** For any $r > 0$, there exists $K_1 > 0$ such that, for all $|y_1|, |y_2| \leq r$,

$$|b(y_1) - b(y_2)|^2 \leq K_1 |y_1 - y_2|^2,$$

- **C2 (One sided linear growth condition)** There exists $K_2 > 0$ such that, for all $y \in \mathbb{R}$,

$$2y \cdot b(y) \leq K_2(1 + |y|^2),$$

then there exists a unique global solution to (1.1) and the solution is adapted and càdlàg. These two conditions also guarantee the existence and uniqueness for the solution of (1.2).

3. Method

In this section, we first study the transition behavior of the system without drift term, and then we study the general case for system (1.1) and (1.2). Before we develop the method, we present the following definition.

Definition 2. For a solution path X_t and the time interval $[T_0, T_f]$ with a partition $T_0 = t_0 \leq t_1 \leq \dots \leq t_n = T_f$, define the sequence $\{X_{t_0}, X_{t_1}, \dots, X_{t_n}\}$ as a discretized path of X_t . And a discretized path $\{X_{t_0}, X_{t_1}, \dots, X_{t_n}\}$ is said to be monotonic with respect to the time partition $T_0 = t_0 \leq t_1 \leq \dots \leq t_n = T_f$ if either $X_{t_0} \leq X_{t_1} \leq \dots \leq X_{t_n}$ or $X_{t_0} \geq X_{t_1} \geq \dots \geq X_{t_n}$.

3.1. Most probable transition path in the absence of drift

Theorem 1. (Monotonicity for the most probable transition path in the absence of drift) For system (1.1) and system 1.2 with drift $b \equiv 0$, every discretized path $\{X_{t_0}, X_{t_1}, \dots, X_{t_n}\}$ of the most probable transition path is monotonic with respect to every time partition $T_0 = t_0 \leq t_1 \leq \dots \leq t_n = T_f$.

Proof. For a time interval partition $T_0 = t_0 \leq t_1 \leq t_2 \leq \dots \leq t_n = T_f$, define $t_{i+1} - t_i \triangleq \epsilon = (T_f - T_0)/n$, $i = 0, 1, \dots, n-1$. (This proof is also true for an arbitrary partition.) In path integral method, the transition density function (or Markov transition probability) of X_t of (1.1) with $b \equiv 0$ is

$$\begin{aligned} p_t(X_f, T_f | X_0, T_0) &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} dx_{t_1} \dots dx_{t_{n-1}} p_\epsilon(x_{t_1} | X_0) p_\epsilon(x_{t_2} | x_{t_1}) \dots p_\epsilon(X_f | x_{t_{n-1}}) \\ &= \int \mathcal{D}_n x \exp\left\{-\theta_\epsilon^\alpha(x_{t_1}, X_0) - \sum_{i=1}^{n-2} \theta_\epsilon^\alpha(x_{t_{i+1}}, x_{t_i}) - \theta_\epsilon^\alpha(X_f, x_{t_{n-1}})\right\} \\ &\triangleq \int \mathcal{D}_n x \exp\left\{-\sum_{i=0}^{n-1} \theta_\epsilon^\alpha(x_{t_{i+1}}, x_{t_i})\right\} \\ &\triangleq \int \mathcal{D}_n x \exp\{-\mathcal{S}_n(x)\}, \end{aligned} \tag{3.16}$$

where $\mathcal{D}_n x = \epsilon^{-1/\alpha} \prod_{i=1}^{n-1} \epsilon^{-1/\alpha} dx_{t_i}$. Note that x of $\mathcal{S}_n(x)$ is a path connecting (T_0, X_0) and (T_f, X_f) (*i.e.* it starts at X_0 at the time $t = T_0$ and reaches X_f at the time $t = T_f$). We call $\mathcal{S}_n(x)$ the action quantity of x . The contribution of a path x_t to the transition probability density $p_t(X_f, T_f | X_0, T_0)$ depends on $\mathcal{S}_n(x) = \sum_{i=0}^{n-1} \theta_\epsilon^\alpha(x_{t_{i+1}}, x_{t_i})$.

In order to find the most probable transition path u_t , we are supposed to find it satisfies that

$$\mathcal{S}_n(u_t) = \min_{x_t \in D_{X_0}^{X_f}} \mathcal{S}_n(x_t), \quad (3.17)$$

where $D_{X_0}^{X_f}$ denotes the set of paths that connect (T_0, X_0) and (T_f, X_f) . Equivalently

$$\frac{\mathcal{S}_n(x_t)}{\mathcal{S}_n(u_t)} \geq 1, \quad (3.18)$$

for every path x_t connecting (T_0, X_0) and (T_f, X_f) . Here n goes to infinity, and the $\lim_{n \rightarrow \infty}$ is dropped for now for clarity.

Without loss of generality, we set $X_0 < X_f$. When $n = 2$ (the time partition is $\{T_0 = t_0 < t_1 < t_2 = T_f\}$), for a path x_t connecting (T_0, X_0) and (T_f, X_f) , $\mathcal{S}_2(x_t) = \theta_\epsilon^\alpha(x_{t_1}, X_0) + \theta_\epsilon^\alpha(X_f, x_{t_1})$, if $x_{t_1} < X_0$. Since $\theta_\epsilon^\alpha(\cdot)$ is a strictly increasing function, together with $|x_{t_1} - X_0| > |X_0 - X_0| = 0$ and $|X_f - x_{t_1}| > |X_f - X_0|$, we have $\mathcal{S}_2(x_t) > \theta_\epsilon^\alpha(X_0, X_0) + \theta_\epsilon^\alpha(X_0, X_f)$ where $\{X_0, x_{t_1} = X_f, X_f\}$ is a discretized path of a certain transition path connecting (T_0, X_0) and (T_f, X_f) . The case that $x_{t_1} > X_f$ is similar. When we add time partitions the situation is similar again. Thus the discretized path $\{X_{t_0}, X_{t_1}, \dots, X_{t_n}\}$ of the most probable transition path of X_t is supposed to be monotonic with respect to the time partition, otherwise there exists a path whose action quantity is smaller.

For system (1.2), the proof is similar. □

The most probable transition path might not be unique, *i.e.*, there might be several paths satisfying (3.17) or (3.18). In fact we will see in the following corollary that, for system (1.1), the number of the most probable transition paths is infinity but they can be characterized by a class of paths that have only one jump, and the difference among the paths of this class is the jump time.

Corollary 1. (i) For system (1.1) with $b \equiv 0$, if L_t is a symmetric α -stable Lévy noise with $0 < \alpha < 1$, then the most probable transition path is not unique, and it can be represented as a Heaviside-like function

$$X_t^m = \begin{cases} X_0, & T_0 \leq t < t^*, \\ X_f, & t^* \leq t \leq T_f, \end{cases} \quad (3.19)$$

for every time instant t^* satisfying $T_0 < t^* \leq T_f$;

(ii) For system (1.2) with drift $b \equiv 0$, the most probable transition path is the line segment connecting X_0 and X_f : $X_t^m = X_0 + \frac{t-T_0}{T_f-T_0}(X_f - X_0)$;

Proof. (i) For system (1.1) with drift $b \equiv 0$, as shown in Section 2, we notice that for a time interval partition $T_0 = t_0 \leq t_1 \leq \dots \leq t_n = T_f$, $t_{i+1} - t_i = \epsilon = \frac{T_f - T_0}{n}$,

$$\theta_\epsilon^\alpha(|x - y|) = \theta_1^\alpha\left(\left|\frac{x - y}{\epsilon^{1/\alpha}}\right|\right) = \theta_1^\alpha\left(n^\alpha \left|\frac{x - y}{(T_f - T_0)^{1/\alpha}}\right|\right). \quad (3.20)$$

Define a path space $DM_{X_0}^{X_f} = \{x_t \mid x_t \text{ is a monotonic path connects } (T_0, X_0) \text{ and } (T_f, X_f)\}$. So one should search for the most probable transition path within path space $DM_{X_0}^{X_f}$.

Take a path $x_t \in DM_{X_0}^{X_f}$ and a time partition $\{T_0 = t_0 < t_1 < \dots < t_n = T_f\}$. Assume that $\{\lambda_j\}_{j=0}^{n-1}$ are non-negative constants, and $x_{t_{j+1}} - x_{t_j} = \lambda_j(X_f - X_0)$ ($0 \leq j \leq n-1$). It is easy to see that $\sum_{j=0}^{n-1} \lambda_j = 1$ by the Theorem 1, and $0 \leq \lambda_j \leq 1$. As discussed in Section 2.2, $\theta_1^\alpha(\cdot)$ is concave in $[r, \infty)$ for some constant $r \in \mathbb{R}^+$ (r depending on α).

For $n^{1/\alpha} \lambda_j \left| \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} \right| \triangleq n^{1/\alpha} \lambda_j C_{0f} \geq r$, we obtain

$$\begin{aligned} \theta_\epsilon^\alpha(|x_{t_{j+1}} - x_{t_j}|) &= \theta_1^\alpha\left(n^{1/\alpha} \lambda_j \left|\frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}}\right|\right) \\ &\geq \lambda_j \theta_1^\alpha\left(n^{1/\alpha} \left|\frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}}\right|\right) \\ &= \lambda_j \theta_\epsilon^\alpha(|X_f - X_0|). \end{aligned} \quad (3.21)$$

When $n^{1/\alpha} \lambda_j C_{0f} < r$, we have

$$0 \leq \sum_{n^{1/\alpha} \lambda_j C_{0f} < r} \lambda_j < n \frac{r}{n^{1/\alpha} C_{0f}}, \quad (3.22)$$

and

$$\lim_{n \rightarrow \infty} \sum_{n^{1/\alpha} \lambda_j C_{0f} < r} \lambda_j = 0, \quad \lim_{n \rightarrow \infty} \sum_{n^{1/\alpha} \lambda_j C_{0f} \geq r} \lambda_j = 1. \quad (3.23)$$

Hence

$$\begin{aligned}
 & \lim_{n \rightarrow \infty} \mathcal{S}_n(x_t) / \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) \\
 &= \lim_{n \rightarrow \infty} \sum_{j=0}^{n-1} \theta_1^\alpha(n^{1/\alpha} \lambda_j | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) / \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) \\
 &= \lim_{n \rightarrow \infty} [\sum_{n^\alpha \lambda_j C_{0f} < r} \theta_1^\alpha(n^{1/\alpha} \lambda_j | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) + \sum_{n^\alpha \lambda_j C_{0f} \geq r} \theta_1^\alpha(n^{1/\alpha} \lambda_j | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |)] / \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) \\
 &\geq \lim_{n \rightarrow \infty} \sum_{n^{1/\alpha} \lambda_j C_{0f} \geq r} \theta_1^\alpha(n^{1/\alpha} \lambda_j | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) / \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) \\
 &\geq \lim_{n \rightarrow \infty} \sum_{n^{1/\alpha} \lambda_j C_{0f} \geq r} \lambda_j \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) / \theta_1^\alpha(n^{1/\alpha} | \frac{X_f - X_0}{(T_f - T_0)^{1/\alpha}} |) \\
 &= \lim_{n \rightarrow \infty} \sum_{n^\alpha \lambda_j C_{0f} \geq r} \lambda_j \\
 &= 1.
 \end{aligned} \tag{3.24}$$

This means that the most probable transition path for symmetric α -stable Lévy process ($0 < \alpha < 1$) is a *Heaviside*-like function

$$X_t^m \triangleq \begin{cases} X_f, & t \geq t^*, \\ X_0, & t < t^*, \end{cases} \tag{3.25}$$

or

$$X_t^m \triangleq \begin{cases} X_f, & t > t^*, \\ X_0, & t \leq t^*, \end{cases} \tag{3.26}$$

where t^* satisfies $T_0 \leq t^* \leq T_f$. Hence in this case, the most probable transition path is not unique since the “jump time” t^* can be chosen arbitrarily.

(ii) For system (1.2) with $b \equiv 0$, by Theorem 1 and the fact that $\theta_\epsilon^b(\cdot)$ being convex, we conclude that

$$\begin{aligned}
 \sum_{i=0}^{n-1} \theta_\epsilon^b(x_{t_{i+1}}, x_{t_i}) &= \sum_{i=0}^{n-1} \theta_\epsilon^b(|x_{t_{i+1}} - x_{t_i}|) \\
 &\geq n \theta_\epsilon^b \left(\sum_{i=0}^{n-1} \frac{1}{n} |x_{t_{i+1}} - x_{t_i}| \right) \\
 &= n \theta_\epsilon^b \left(\frac{1}{n} |X_f - X_0| \right).
 \end{aligned} \tag{3.27}$$

The inequality holds if and only if

$$\begin{aligned}
 \theta_\epsilon^b(x_{t_1}, x_{t_0}) &= \theta_\epsilon^b(x_{t_2}, x_{t_1}) = \cdots = \theta_\epsilon^b(x_{t_n}, x_{t_{n-1}}) \\
 \Leftrightarrow x_{t_1} - x_{t_0} &= x_{t_2} - x_{t_1} = \cdots = x_{t_n} - x_{t_{n-1}}.
 \end{aligned} \tag{3.28}$$

So the most probable transition path is the line segment which connects the initial and final points. Therefore, we obtain

$$X_{t_i}^m = \frac{i(X_f - X_0)}{n}, \quad i = 1, 2, \dots, n-1. \quad (3.29)$$

When n goes to infinity, $X_t^m = X_0 + \frac{t-T_0}{T_f-T_0}(X_f - X_0)$. This implies that the most probable transition path is the path for the particle (i.e., solution) moving in constant velocity. \square

3.2. Most probable transition path in the case of non-zero drift

Theorem 2. (Characterization of the most probable transition path with drift term)

For system (1.1) and system (1.2), assume that the transition probability density exists, and that the most probable transition path X_t^m exists and satisfies the integrability condition $|\int_{T_0}^t b(X_s^m)ds| < \infty$ for $t \in [T_0, T_f]$.

(i) For system (1.1) with L_t a symmetric α -stable Lévy motion with ($0 < \alpha < 1$), the most probable transition path X_t^m is determined by the following deterministic dynamical system (i.e., an ordinary differential equation),

$$\begin{cases} dX_t^m - b(X_t^m)dt = 0, & t \in [T_0, T_f] \setminus \{t^*\}, \\ X_{T_0}^m = X_0, & X_{T_f}^m = X_f, \end{cases} \quad (3.30)$$

where the jump time t^* is such that

$$J_{t^*} = \inf_{T_0 < t \leq T_f} J_t. \quad (3.31)$$

Here $J_t = |X_0 + \int_{T_0}^t b(X_s^m)ds - X_f + \int_t^{T_f} b(X_s^m)ds|$ is the jump size.

(ii) For system (1.2) with Brownian motion, the most probable transition path X_t^m is determined by the following deterministic dynamical system (i.e., an integral-differential equation),

$$X_t^m - X_0 - \int_{T_0}^t b(X_s^m)ds = \frac{t - T_0}{T_f - T_0}(X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m)ds). \quad (3.32)$$

Proof. For the system (1.1), the corresponding stochastic integral equation is

$$\begin{aligned} X_t - X_0 &= \int_{T_0}^t b(X_s)ds + L_t - L_{T_0}, \\ Y_t \triangleq X_t - X_0 - \int_{T_0}^t b(X_s)ds &= L_t - L_{T_0}, \end{aligned} \quad (3.33)$$

and the differential form is

$$X_{t_{i+1}} - X_{t_i} - [\gamma b(X_{t_{i+1}}) + (1 - \gamma)b(X_{t_i})]\Delta t = L_{t_{i+1}} - L_{t_i}. \quad (3.34)$$

In Itô interpretation ($\gamma = 0$),

$$X_{t_{i+1}} - X_{t_i} - b(X_{t_i})\Delta t = L_{t_{i+1}} - L_{t_i}. \quad (3.35)$$

The transition probability density function of X_t is

$$\begin{aligned} & p(X_f, T_f | X_0, T_0) \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} dL_1 \cdots dL_{n-1} dL_n p_\epsilon(L_1 | L_{T_0}) p_\epsilon(L_2 | L_1) \cdots p_\epsilon(L_n | L_{n-1}) \delta(x_{t_n} - X_f) \\ &= \int \mathcal{D}_{n+1} x \mathcal{J} \exp\left\{-\sum_{i=0}^n \theta_\epsilon^\alpha(x_{t_{i+1}} - x_{t_i} - b(x_{t_i})\Delta t)\right\} \delta(x_{t_n} - X_f) \\ &= \int \mathcal{D}_{n+1} x \exp\left\{-\sum_{i=0}^n \theta_\epsilon^\alpha(x_{t_{i+1}} - x_{t_i} - b(x_{t_i})\Delta t)\right\} \delta(x_{t_n} - X_f) \\ &= \int \mathcal{D}_{n+1} y \exp\left\{-\sum_{i=0}^n \theta_\epsilon^\alpha(y_{t_{i+1}}, y_{t_i})\right\} \delta(x_{t_n} - X_f), \end{aligned} \quad (3.36)$$

where \mathcal{J} is the Jacobian of the transformation given by

$$\mathcal{J} = \det\left(\frac{\partial L_i}{\partial x_k}\right) = \prod_{i=1}^n \left(1 - \epsilon \gamma \frac{db(x_i)}{dx_i}\right). \quad (3.37)$$

Assume that the most probable transition path of X_t exist, which is denoted by X_t^m .

(i) For system (1.1), in order to determine the most probable transition path X_t^m , we consider the transition of the process Y_t from $Y_{T_0} = 0$ to $Y_{T_f} = X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m) ds$. Since L_t is an α -stable Lévy motion with $0 < \alpha < 1$, by Corollary 1, the most probable transition path Y_t^m is

$$Y_t^m = \begin{cases} 0, & T_0 \leq t < t^*, \\ X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m) ds, & t^* \leq t \leq T_f, \end{cases} \quad (3.38)$$

for some $t^* \in (T_0, T_f]$. Then

$$\begin{cases} X_t^m - X_0 - \int_{T_0}^t b(X_s^m) ds = 0, & T_0 \leq t \leq t^*, \\ X_t^m - X_0 - \int_{T_0}^t b(X_s^m) ds = X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m) ds, & t^* < t \leq T_f. \end{cases} \quad (3.39)$$

This is

$$\begin{cases} dX_t^m - b(X_t^m) dt = 0, \\ X_{T_0}^m = X_0, X_{T_f}^m = X_f, t \in [T_0, T_f] \setminus \{t^*\}. \end{cases} \quad (3.40)$$

It means that the most probable transition path has one jump and the jump size is $J_{t^*} = |X_0 + \int_{T_0}^{t^*} b(X_s^m) ds - X_f + \int_{t^*}^{T_f} b(X_s^m) ds|$, and the jump time t^* is supposed to be the one which satisfies

$$J_{t^*} = \min_{T_0 \leq t \leq T_f} J_t. \quad (3.41)$$

As J_t is continuous in t , the minimizer t^* exists (although it may not be unique).

In other words, the most probable transition path consists of two components: One component is part of the solution of equation

$$\begin{cases} dX_t^m - b(X_t^m)dt = 0, \\ X_{T_0}^m = X_0, \quad t \in [T_0, T_f], \end{cases} \quad (3.42)$$

and the other component is part of the solution of equation:

$$\begin{cases} dX_t^m - b(X_t^m)dt = 0, \\ X_{T_f}^m = X_f, \quad t \in [T_0, T_f]. \end{cases} \quad (3.43)$$

The jump time t^* is the moment at which the distance of the solutions of (3.42) and (3.43) is minimum.

(ii) For system (1.2), the most probable transition path of Y_t (denoted by Y_t^m) can be obtained exactly,

$$\begin{aligned} Y_t^m &= X_t^m - X_0 - \int_{T_0}^t b(X_s^m)ds \\ &= Y_0 + (Y_f - Y_0) \frac{t - T_0}{T_f - T_0} \\ &= \frac{t - T_0}{T_f - T_0} (X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m)ds). \end{aligned} \quad (3.44)$$

This completes the proof of this theorem. \square

Remark 1. For a symmetric α -stable Lévy motion with $1 \leq \alpha < 2$, the equalities (3.23) do not hold. Define the path space $DMS_{X_0}^{X_f} \triangleq \{x_t \mid x_t = \sum_{i=0}^{n-1} a_i \mathbb{I}_{[t_i, t_{i+1})} + a_n \mathbb{I}_{\{T_f\}}, T_0 = t_0 \leq t_1 \leq \dots \leq t_n = T_f, X_0 = a_0 \leq a_1 \leq \dots \leq a_n = X_f\}$, where \mathbb{I}_B is the characteristic function of set $B \subset \mathbb{R}$.

If we search for the most probable transition path within the simple path space $DMS_{X_0}^{X_f}$ for the symmetric α -stable Lévy motion with $1 \leq \alpha < 2$, the similar results of Lemma 1 and Theorem 2 hold. This is because for every fixed simple path in $DMS_{X_0}^{X_f}$, the non-zero λ_j of $\{\lambda_j\}_{j=0}^{n-1}$ are bounded below. Thus the inequality (3.21) holds for every non-negative λ_j , for n large enough.

Remark 2. For simplicity, we call the solution of (3.42) the **initial transition path**, and call the solution of (3.43) the **final transition path**. Consequently, the most probable transition path starts from the initial transition path and jumps to the final transition path at the time that initial transition path and final transition path are closest to each other.

3.3. Existence, uniqueness and numerical simulation for the most probable transition path.

3.3.1. *Non-Gaussian noise: System with a symmetric α -stable Lévy motion with $0 < \alpha < 1$.* The most probable transition path is determined by two “initial” value

problems of a deterministic ordinary differential equation (ODE)

$$\begin{cases} dX_t^m - b(X_t^m)dt = 0, & t \in [T_0, T_f] \setminus \{t^*\}, \\ X_{T_0}^m = X_0, & X_{T_f}^m = X_f, & t^* \in [T_0, T_f]. \end{cases} \quad (3.45)$$

The first initial value problem solves this ODE with $X_{T_0}^m = X_0$, and the second problem solves this ODE backward in time with terminal value condition $X_{T_f}^m = X_f$. The existence and uniqueness of these solutions are ensured by the local Lipschitz continuity of the drift term $b(x)$.

Given a time partition $\{T_0 = t_0 < t_1 < \dots < t_n = T_f\}$, we simulate the most probable transition path as follows: forward Euler scheme

$$x_{t_{i+1}} - x_{t_i} = b(x_{t_i})\Delta t, \quad T_0 \leq t_i < t_{i+1} < t^*, \quad x_{T_0} = X_0, \quad (3.46)$$

and backward Euler scheme

$$x_{t_{i+1}} - x_{t_i} = b(x_{t_{i+1}})\Delta t, \quad t^* \leq t_i < t_{i+1} \leq T_f, \quad x_{T_f} = X_f. \quad (3.47)$$

3.3.2. Gaussian noise: System with a Brownian motion. In this case, the most probable transition path is determined by a deterministic integral-differential equation

$$X_t^m - X_0 - \int_{T_0}^t b(X_s^m)ds = \frac{t - T_0}{T_f - T_0}(X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m)ds). \quad (3.48)$$

Now we study the existence and uniqueness of the solution of (3.48). Differentiate (3.48) with respect to t on both sides:

$$dX_t^m = b(X_t^m)dt + \frac{dt}{T_f - T_0}(X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m)ds), \quad (3.49)$$

notice that $\frac{1}{T_f - T_0}(X_f - X_0 - \int_{T_0}^{T_f} b(X_s^m)ds)$ is independent of t .

For any constant $c \in \mathbb{R}$, the equation

$$U_t' = b(U_t) + c, \quad U_{T_0} = X_0, \quad U_t' = \frac{dU_t}{dt}, \quad (3.50)$$

has a unique solution which is continuous and differentiable since $(b(u) + c)$ is also a locally Lipschitz continuous function. The integral of the solution of 3.50 is finite in time interval $[T_0, T_f]$. If there exists a constant c_0 such that the solution of 3.50 denoted by U_t^* satisfies $U_{T_f}^* = X_f$, then

$$\begin{aligned} U_t^* &= X_0 + \int_{T_0}^t b(U_s^*)ds + c(t - T_0), \\ \Rightarrow X_f &= X_0 + \int_{T_0}^{T_f} b(U_s^*)ds + c_0(T_f - T_0), \\ \Rightarrow c_0 &= \frac{1}{T_f - T_0}(X_f - X_0 - \int_{T_0}^{T_f} b(U_s^*)ds), \end{aligned} \quad (3.51)$$

thus U_t^* satisfies the equation 3.48.

To check that if there exists thus constant c_0 , we assume b is differentiable, and consider the following second order differential equation with two-point boundary value:

$$U_t'' = b'(U_t)U_t', \quad U_{T_0} = X_0, \quad U_{T_f} = X_f. \quad (3.52)$$

If $\tilde{b}(t, u, u') \triangleq b'(u_t)u_t'$ is continuous and satisfies $|\tilde{b}| \leq M$ for $T_0 \leq t \leq T_f$, $|u| \leq R_0$, $|u'| \leq R_1$, let p and X_0 satisfy $T_0 < p \leq T_f$ and

$$\frac{Mp^2}{8} + |X_0| \leq R_0, \quad \frac{Mp}{2} + \frac{X_0}{p} \leq R_1, \quad (3.53)$$

then it was proved in [19] that 3.52 has a solution. The solution of 3.52 also satisfies equation 3.50 which means there exists a constant c_0 such that the solution of 3.48 exists

Thus the existence of solution of (3.48) is guaranteed provided the assumptions above. The uniqueness can be guaranteed if \tilde{b} is continuous for $T_0 \leq t \leq T_f$ and all (u, u') and satisfies a Lipschitz condition with respect to u, u' of the form

$$|\tilde{b}(u_1, u_1') - \tilde{b}(u_2, u_2')| \leq \zeta_0|u_1 - u_2| + \zeta_1(u_1' - u_2'), \quad (3.54)$$

with Lipschitz constants ζ_0, ζ_1 so small that

$$\frac{\zeta_0 p^2}{8} + \frac{\zeta_1 p}{2} < 1, \quad (3.55)$$

which can be found in [19]

Let us consider a numerical method to calculate the most probable transition path under the time interval partition $\{T_0 = t_0 < t_1 < \dots < t_n = T_f, \Delta t = \frac{T_f - T_0}{n}\}$ when the noise is a Brownian motion:

$$\begin{cases} x_{t_{i+1}} - x_{t_i} - b(x_{t_i})\Delta t = x_{t_i} - x_{t_{i-1}} - b(x_{t_{i-1}})\Delta t, & i = 1, 2, \dots, n-1, \\ x_{T_0} = X_0, \quad x_{T_f} = X_f. \end{cases} \quad (3.56)$$

There are $n - 1$ equations to determine $n - 1$ unknowns,

$$\begin{cases} x_{t_i} = \frac{1}{2}[x_{t_{i+1}} + x_{t_{i-1}} - (b(x_{t_i}) - b(x_{t_{i-1}}))\Delta t] \triangleq \Phi_i(x_{t_{i+1}}, x_{t_{i-1}}), & i = 1, 2, \dots, n-1, \\ x_{T_0} = X_0, \quad x_{T_f} = X_f. \end{cases} \quad (3.57)$$

This may be written as $X = \Phi(X)$. When the function Φ is a contraction map with respect to the domain of the state space that contains continuous paths connecting X_0 and X_f , the fixed point theory gives a method to approximate the solution of the equations (3.57), and the existence and uniqueness are also obtained.

4. Higher Dimensional Cases

In this section, we discuss the higher dimensional cases. We consider an SDE system with non-Gaussian noise

$$\begin{cases} dX_{1,t} = b_1(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dL_{1,t}, & X_{1,T_0} = X_{1,0}, \\ dX_{2,t} = b_2(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dL_{2,t}, & X_{2,T_0} = X_{2,0}, \\ \vdots & \vdots \\ dX_{k,t} = b_k(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dL_{k,t}, & X_{k,T_0} = X_{k,0}, \end{cases} \triangleq dX_t = b(X_t)dt + dL_t, \quad X_{T_0} = X_0, \quad (4.58)$$

where $L_{i,t}$ are symmetric α -stable Lévy noises ($0 < \alpha < 1$) and $\{L_{i,t}, \dots, L_{j,t}\}$ are independent, and an SDE system with Gaussian noise

$$\begin{cases} dX_{1,t} = b_1(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dB_{1,t}, & X_{1,T_0} = X_{1,0}, \\ dX_{2,t} = b_2(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dB_{2,t}, & X_{2,T_0} = X_{2,0}, \\ \vdots & \vdots \\ dX_{k,t} = b_k(X_{1,t}, X_{2,t}, \dots, X_{k,t})dt + dB_{k,t}, & X_{k,T_0} = X_{k,0}, \end{cases} \triangleq dX_t = b(X_t)dt + dB_t, \quad X_{T_0} = X_0, \quad (4.59)$$

where $B_{i,t}$ are Brownian motions, and $\{B_{1,t}, \dots, B_{k,t}\}$ are independent.

It was known [20] that the random variables X_1, \dots, X_k are independent if and only if $f(x_1, \dots, x_k) = f_1(x_1) \cdots f_k(x_k)$ for all (x_1, \dots, x_n) except possibly for a Borel subset of \mathbb{R}^k with Lebesgue measure zero. Here f is the probability density of (X_1, \dots, X_k) and f_i is the probability density of X_i ($i = 1, \dots, k$). Hence by the independence of the noises, the probability density function denoted by $f_\alpha^k(x)$ of k -dimensional α -stable Lévy variable $x = (x_1, x_2, \dots, x_k) \in \mathbb{R}^k$ is

$$\begin{aligned} f_\alpha^k(x) &= f_\alpha(x_1)f_\alpha(x_2) \cdots f_\alpha(x_k) \\ &= \exp\left\{-\left[-\sum_{i=1}^k \ln(f_\alpha(x_i))\right]\right\} \\ &= \exp\left\{-\sum_{i=1}^k \theta_1^\alpha(x_i)\right\}. \end{aligned} \quad (4.60)$$

Recall in Theorem 1, we proved that the most probable transition path is supposed to be monotonic with respect to time t . In higher dimensional cases, the transition probability density function has the similar form of (3.16). It implies that every component of the most probable transition path are monotonic with respect to time t .

For the system (4.58) with zero drift term, the time partition $\{T_0 = t_0 < t_1 < \dots < t_n = T_f, t_{i+1} - t_i = \frac{T_f - T_0}{n}\}$, and for a path $x_t = (x_{1,t}, x_{2,t}, \dots, x_{k,t})$ connecting (T_0, X_0)

and (T_f, X_f) monotonically, we denote $x_{i,t_{j+1}} - x_{i,t_j} = \lambda_j^i (X_{i,f} - X_{i,0})$. Thus

$$\begin{aligned}
 \mathcal{S}_n(x_t) &= \sum_{i=1}^k \sum_{j=0}^{n-1} \theta_\epsilon^\alpha (x_{i,t_{j+1}} - x_{i,t_j}) \\
 &= \sum_{i=1}^k \sum_{j=0}^{n-1} \theta_1^\alpha (\lambda_j^i n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}}) \\
 &\geq \sum_{i=1}^k \sum_{\lambda_j^i n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}} \geq r_i} \theta_1^\alpha (\lambda_j^i n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}}) \\
 &\geq \sum_{i=1}^k \sum_{\lambda_j^i n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}} \geq r_i} \lambda_j^i \theta_1^\alpha (n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}}) \\
 &= \sum_{i=1}^k \theta_1^\alpha (n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}}) \sum_{\lambda_j^i n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}} \geq r_i} \lambda_j^i \\
 &\geq \sum_{i=1}^k \theta_1^\alpha (n^{1/\alpha} \frac{X_{i,f} - X_{i,0}}{(T_f - T_0)^{1/\alpha}}) (n \rightarrow \infty).
 \end{aligned} \tag{4.61}$$

This implies the results of Corollary 1 and Theorem 2 in higher dimensional cases and they are similar to 1-dimensional case.

For the system (4.59) with zero drift term,

$$\begin{aligned}
 \mathcal{S}_n(x_t) &= \sum_{i=1}^k \sum_{j=0}^{n-1} \theta_\epsilon^b (x_{i,t_{j+1}} - x_{i,t_j}) \\
 &= \sum_{i=1}^k \sum_{j=0}^{n-1} \theta_\epsilon^b (|x_{i,t_{j+1}} - x_{i,t_j}|) \\
 &\geq \sum_{i=1}^k n \theta_\epsilon^b (\sum_{j=0}^{n-1} \frac{1}{n} |x_{i,t_{j+1}} - x_{i,t_j}|) \\
 &= \sum_{i=1}^k n \theta_\epsilon^b (\frac{1}{n} |X_{i,f} - X_{i,0}|).
 \end{aligned} \tag{4.62}$$

This implies the results of Corollary 1 and Theorem 2 in higher dimensional cases and they are similar to 1-dimensional case.

5. Examples

Let us consider several examples in order to illustrate our results.

Example 1. *Ornstein-Uhlenbeck process*

Consider a linear scalar SDE:

$$dX_t = rdt + dL_t, \quad T_0 \leq t \leq T_f, \quad X_{T_0} = X_0, \quad X_{T_f} = X_f,$$

with r a constant. Let $Y_t = X_t - rt$. Then by Itô formula

$$dY_t = -rdt + dX_t = dL_t.$$

When L_t is a symmetric α -stable Lévy motion with $0 < \alpha < 1$, the most probable transition path of Y_t is

$$Y_t^m = \begin{cases} X_f - rT_f, & t \geq t^* \quad (t > t^*), \\ X_0 - rT_0, & t < t^* \quad (t \leq t^*), \end{cases}$$

for every t^* satisfying $T_0 \leq t^* \leq T_f$. Thus the most probable transition path for X_t is

$$X_t^m = \begin{cases} X_f - r(T_f - t), & t \geq t^* \quad (t > t^*), \\ X_0 + r(t - T_0), & t < t^* \quad (t \leq t^*), \end{cases}$$

where t^* can be chosen arbitrarily in $[T_0, T_f]$.

When L_t is replaced by a Brownian motion,

$$dX_t = rdt + dB_t, \quad T_0 \leq t \leq T_f, \quad X_{T_0} = X_0, \quad X_{T_f} = X_f,$$

the most probable transition path of X_t is (by Theorem 2),

$$\begin{aligned} X_t^m - r(t - T_0) &= X_0 + \frac{t - T_0}{T_f - T_0} (X_f - X_0 - r(T_f - T_0)) \\ \Leftrightarrow X_t^m &= X_0 + \frac{t - T_0}{T_f - T_0} (X_f - X_0). \end{aligned}$$

So in this linear system with drift and Gaussian noise, the most probable transition path is also a line segment.

Figure 2 shows the most probable transition paths of this example.

Example 2. *Geometric Brownian motion*

Consider a linear scalar SDE with multiplicative noise

$$dX_t = rX_t dt + \eta X_t dB_t,$$

where r and η are real constants, and $X_t > 0$, *a.s.*. Setting $Y_t = \ln X_t - (r - \frac{1}{2}\eta^2)t$ and applying Itô formula, we obtain

$$\begin{aligned} dY_t &= d\ln(X_t) - (r - \frac{1}{2}\eta^2)dt \\ &= \frac{1}{X_t} dX_t + \frac{1}{2} \left(-\frac{1}{X_t^2}\right) (dX_t)^2 - (r - \frac{1}{2}\eta^2)dt \\ &= \frac{dX_t}{X_t} - \frac{1}{2}\eta^2 dt - (r - \frac{1}{2}\eta^2)dt \\ &= \eta dB_t. \end{aligned}$$




Figure 2: The most probable transition paths of Ornstein-Uhlenbeck process in Example 1 with $T_0 = 0$, $T_f = 4$, $r = 1$, $X_0 = 0$, $X_f = 5$. We choose $t^* = 2$ and the ‘right-continuous with left limit’ version of the most probable transition path.

By Corollary 1,

$$Y_t^m = Y_0 + \frac{t - T_0}{T_f - T_0}(Y_f - Y_0).$$

Thus

$$X_t^m = \exp\left\{\left(r - \frac{1}{2}\eta^2\right)t + \ln X_0 + \frac{t - T_0}{T_f - T_0}[\ln X_f - \left(r - \frac{1}{2}\eta^2\right)T_f - \ln X_0]\right\}.$$

Figure 3 shows the most probable transition path of this example.

Example 3. *Geometric Lévy process*

Consider the stochastic differential equation

$$dX_t = X_t[\zeta dt + \beta dB(t) + \int_{\mathbb{R}} \gamma(t, z) \bar{N}(dt, dz)],$$

where ζ , β are constants and $\gamma(t, z) \geq -1$, and

$$\bar{N}(dt, dz) = \begin{cases} N(dt, dz) - \nu(dz)dt, & \text{if } |z| < r, \\ N(dt, dz), & \text{if } |z| \geq r, \end{cases}$$

$r \in \mathbb{R}^+$. For simplicity we set $\beta = 0$, $\gamma(t, z) = e^z - 1$. Now define $Y_t = \ln X_t$. By Itô formula, we have

$$\begin{aligned} dY_t &= \zeta dt + \int_{|z| < r} \{\ln(1 + e^z - 1) - (e^z - 1)\} \nu(dz) dt + \int_{\mathbb{R}} \ln(1 + e^z - 1) \bar{N}(dt, dz) \\ &= \zeta dt + \int_{|z| < r} \{z - (e^z - 1)\} \nu(dz) dt + \int_{\mathbb{R}} z \bar{N}(dt, dz). \end{aligned}$$

Let $U_t = Y_t - (\zeta + \int_{|z|<r} \{z - (e^z - 1)\} \nu(dz))t$ be a symmetric α -stable Lévy process, i.e., $dU_t = \int_r z \bar{N}(dt, dz) = dL_t^\alpha$.

Let the jump measure $\nu(dz)$ be the jump measure of an α -stable Lévy process, that is, $\nu(dz) = c_\alpha \frac{dz}{|z|^{1+\alpha}}$ with

$$c_\alpha = \frac{\alpha}{2^{1-\alpha} \sqrt{\pi}} \frac{\Gamma(\frac{1+\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})}.$$

For α with $0 < \alpha < 1$ and $r = 1$,

$$\int_{|z|<1} \{z - (e^z - 1)\} \nu(dz) < \infty.$$

By Corollary 1, the most probable transition path of U_t is

$$U_t^m = \begin{cases} U_0, & T_0 \leq t < t^* \quad (T_0 \leq t \leq t^*), \\ U_f, & t^* \leq t \leq T_f \quad (t^* < t \leq T_f), \end{cases}$$

where $t^* \in [T_0, T_f]$. Thus

$$X_t^m = \begin{cases} \exp(\ln X_0 + (t - T_0)[\zeta + \int_{|z|<1} \{z - (e^z - 1)\} \nu(dz)]), & T_0 \leq t < t^* \quad (T_0 \leq t \leq t^*), \\ \exp(\ln X_f + (t - T_f)[\zeta + \int_{|z|<1} \{z - (e^z - 1)\} \nu(dz)]), & t^* \leq t \leq T_f \quad (t^* < t \leq T_f). \end{cases}$$

Figure 3 also shows the most probable transition path of this example.

Example 4. *1-dimensional Nonlinear SDE: Stochastic double-well system*

Consider the stochastic double-well systems

$$dX_t = (X_t - X_t^3)dt + dL_t,$$

where L_t is a symmetric α -stable Lévy motion with $0 < \alpha < 1$, and

$$dX_t = (X_t - X_t^3)dt + dB_t.$$

By Theorem 2, the most probable transition path of the first system is described as the following deterministic differential equation:

$$\begin{cases} dX_t^m - (X_t^m - (X_t^m)^3)dt = 0, & t \in [T_0, T_f] \setminus \{t^*\}, \\ X_{T_0}^m = X_0, \quad X_{T_f}^m = X_f, & t^* \in [T_0, T_f]. \end{cases}$$

The most probable transition path for the second system is determined by

$$X_t^m - X_0 - \int_{T_0}^t (X_s^m - (X_s^m)^3)ds = \frac{t - T_0}{T_f - T_0} (X_f - X_0 - \int_{T_0}^{T_f} (X_s^m - (X_s^m)^3)ds).$$

By the numerical method in section 3.3, we compute the most probable transition paths as shown in Figure 4.

Example 5. *2-Dimensional Nonlinear SDE: The Maier-Stein model*




Figure 3: The most probable transition paths of Geometric Brownian motion (black solid line) and the initial and final transition path of geometry symmetric α -stable Lévy motion with $\alpha = 0.5$ (grey solid lines, the most probable path starts at initial transition path and jumps to final transition path at time t^* at which the two grey solid lines are closest to each other), for $T_0 = 0, T_f = 2, X_0 = 1, X_2 = 5, r = 1, \eta = 1, \beta = 0$: (a) $\zeta = -1, t^* = 2$; (b) $\zeta = 0, t^* = 2$; (c) $\zeta = 0.5, t^* = 0$; (d) $\zeta = 2, t^* = 0$.

Consider the following SDEs:

$$\begin{cases} du = (u - u^3 - \beta uv^2)dt + dL_{1,t}, \\ dv = -(1 + u^2)vdt + dL_{2,t}, \end{cases}$$

and

$$\begin{cases} du = (u - u^3 - \beta uv^2)dt + dB_{1,t}, \\ dv = -(1 + u^2)vdt + dB_{2,t}, \end{cases}$$

By Theorem 2, the most probable transition path (u_t^m, v_t^m) of the first system is described




Figure 4: The most probable transition paths of stochastic double-well system with Brownian motion (black spots), the initial and final transition paths of symmetric α -stable Lévy motion (grey spots) in time interval $[0, 2]$ and $X_2 = 1$: (a) $X_0 = -1$, t^* can be chosen arbitrarily in $[T_0, T_f]$; (b) $X_0 = -1.5$, $t^* = 2$; (c) $X_0 = -0.5$, $t^* = 0$; (d) $X_0 = 0.2$, $t^* = 2$.

by the following deterministic differential equations:

$$\begin{cases} du_t^m = (u_t^m - (u_t^m)^3 - \beta u_t^m (v_t^m)^2) dt, & t \in [T_0, T_f] \setminus \{t^*\}, \\ dv_t^m = -(1 + (u_t^m)^2) v_t^m dt, & t \in [T_0, T_f] \setminus \{t^*\}, \\ u_{T_0}^m = u_0, \quad u_{T_f}^m = u_f, \quad v_{T_0}^m = v_0, \quad v_{T_f}^m = v_f, & t^* \in [T_0, T_f], \end{cases}$$

and the most probable transition path for the second system is determined by

$$\begin{cases} u_t^m - u_0 - \int_{T_0}^t (u_s^m - (u_s^m)^3 - \beta u_s^m (v_s^m)^2) ds = \frac{t-T_0}{T_f-T_0} (u_f - u_0 - \int_{T_0}^{T_f} (u_s^m - (u_s^m)^3 - \beta u_s^m (v_s^m)^2) ds), \\ v_t^m - v_0 - \int_{T_0}^t (-(1 + (u_s^m)^2) v_s^m) ds = \frac{t-T_0}{T_f-T_0} (v_f - v_0 - \int_{T_0}^{T_f} (-(1 + (u_s^m)^2) v_s^m) ds). \end{cases}$$

Figure 5 shows the most probable transition paths of Maier-Stein Model with Brownian motion, the initial and final transition paths of Maier-Stein Model with independent symmetric α -stable Lévy motions.




Figure 5: The most probable transition paths of Maier-Stein Model with Brownian motion (black spots), the initial and final transition paths of Maier-Stein Model with symmetric α -stable Lévy motion (grey spots): $T_0 = 0$, $T_f = 2$, $u_{T_0} = -1$, $u_{T_f} = 0$, $v_{T_0} = 0$, $v_{T_f} = 1$, $\beta = 10$, and $t^* = 0$.

Acknowledgements

The authors would like to thank Professor Xu Sun, Dr Qiao Huang, Dr Yayun Zheng, Dr Wei Wei, Dr Ao Zhang, and Dr Jianyu Hu for helpful discussions. This work was partly supported by the NSF grant 1620449, and NSFC grants 11531006 and 11771449.

References

- [1] A. Janicki and A. Weron. Simulation and Chaotic Behavior of alpha-Stable Stochastic Processes. New York: Marcel Dekker (1994).
- [2] A. W. C. Lau and T. C. Lubensky. State-dependent diffusion: Thermodynamic consistency and its path integral formulation. *Phys. Rev. E*. 76,011123 (2007).
- [3] D. Applebaum. Lvy Processes and Stochastic Calculus (Cambridge: Cambridge University Press) (2009).
- [4] D. Dürr and A. Bach. The Onsager-Machlup Function as Lagrangian for the Most Probable Path of a Diffusion Process. *Commun.Math. Phys.* 60: 153 (1978).
- [5] D. Janakiraman and K. L. Sebastian, Path-integral formulation for Lvy flights: Evaluation of the propagator for free, linear, and harmonic potentials in the over- and underdamped limits, *Phys. Rev. E*. 86, 061105 (2012).
- [6] D. M. Zuckerman and T. B. Woolf. Efficient dynamic importance sampling of rare events in one dimension. *Phys. Rev. E*. 63, 016702 (2000).
- [7] F. Xi, C. Zhu. Jump type stochastic differential equations with non-Lipschitz coefficients: Non-confluence, Feller and strong Feller properties, and exponential ergodicity. *J. Differential Equations*(2018), <https://doi.org/10.1016/j.jde.2018.10.006>.
- [8] G. Gobbo, A. Laio, A. Maleki, and S. Baroni. Absolute Transition Rates for Rare Events from Dynamical Decoupling of Reaction Variables. *Phys. Rev. Lett.* 109, 150601 (2012).

- [9] H. S. Wio. Path Integrals for Stochastic Processes: An Introduction. World Scientific, New York (2013).
- [10] J. Duan. An Introduction to Stochastic Dynamics (New York: Cambridge University Press) (2015).
- [11] J. Wang, K. Zhang, H. Lu, and E. Wang. Dominant Kinetic Paths on Biomolecular Binding-Folding Energy Landscape. *Phys. Rev. Lett.* 96, 168101 (2006).
- [12] K. L. C. Hunt, and J. Ross. Path integral solutions of stochastic equations for nonlinear irreversible processes: The uniqueness of the thermodynamic Lagrangian. *The Journal of Chemical Physics.* 75, 976 (1981).
- [13] K.-I. Sato. Lvy Processes and Infinite Divisibility. Cambridge University Press (1999).
- [14] M. Shao and C. L. Nikias. Signal processing with fractional lower order moments: Stable processes and their applications. *Proc. of the IEEE.* 81(7): 986-1010 (1993).
- [15] N. Laskin. Fractional quantum mechanics. *Phys. Rev. E.* 66, 056108 (2002).
- [16] N. Jacob. Pseudo-Differential Operators and Markov Processes, vol.1: Fourier Analysis and Semigroups. London: Imperial College Press (2001).
- [17] N. Jacob, V. Knopova, S. Landwehr, and R. Schilling. A geometric interpretation of the transition density of a symmetric Lévy process. *Sci. China Math.* 55(6): 1099C1126 (2012).
- [18] P. Faccioli, M. Sega, F. Pederiva, and H. Orland. Dominant Pathways in Protein Folding. *Phys.Rev. Lett.* 97, 108101 (2006).
- [19] P. Hartman. Ordinary Differential Equations (Second Edition). *SIAM.* Classics in applied mathematics ; 38.
- [20] R. B. Ash. Probability and Measure Theory (A Harcourt Science and Technology Company press) (1999).
- [21] R. P. Feynman and A. R. Hibbs. Quantum Mechanics and Path Integrals (New York: McGraw-Hill) (1965).
- [22] S. Machlup and L. Onsager. Fluctuations and Irreversible Processes. *Phys. Rev.* 91, 1512 (1953).
- [23] S. Albeverio, Z. Brzezniak and J. WU. Existence of global solutions and invariant measures for stochastic differential equations driven by poisson type noise with non-lipschitz coefficients *Journal of Mathematical Analysis and Applications.* 371(1) 309-322 (2010).
- [24] Y. Tang, R. Yuan, P. Ao. Summing over trajectories of stochastic dynamics with multiplicative noise. *J. Chem. Phys.* 141, Article 044125 (2014).
- [25] Z. G. Arenas and D. G. Barci. Functional integral approach for multiplicative stochastic processes. *Phys. Rev. E.* 81, 051113 (2010).