

Robust Approximate Bayesian Inference with Synthetic Likelihood

David T. Frazier*, and Christopher Drovandi†

February 5, 2022

Abstract

Bayesian synthetic likelihood (BSL) is now an established method for conducting approximate Bayesian inference in models where, due to the intractability of the likelihood function, exact Bayesian approaches are either infeasible or computationally too demanding. Similar to other approximate Bayesian methods, such as the method of approximate Bayesian computation, implicit in the application of BSL is the assumption that the data generating process (DGP) can produce simulated summary statistics that capture the behaviour of the observed summary statistics. We demonstrate that this notion of compatibility between the observed and simulated summaries is critical for the performance of BSL and its variants: across several different simulated and empirical examples, we demonstrate that if the assumed DGP used in BSL differs from the true DGP, BSL can yield unreliable parameter inference. To circumvent this issue, we propose two robust versions of BSL that can deliver reliable inference regardless of whether or not the assumed DGP is correctly specified. Simulation results and two real data examples demonstrate the performance of this robust approach to BSL, and its superiority over standard BSL when the assumed model is misspecified.

Keywords: approximate Bayesian computation; likelihood-free inference; model misspecification; robust Bayesian inference; MCMC; slice sampling.

1 Introduction

In situations where the likelihood of the underlying model is intractable, approximate Bayesian methods are often the only feasible solution to conduct Bayesian inference. Indeed, approximate Bayesian methods are an increasingly common tool in the arsenal of the practicing statistician and allow users to conduct reliable inference in models where exact Bayesian inference procedures are either infeasible, or too computationally demanding.

*Department of Econometrics and Business Statistics and the Australian Centre of Excellence for Mathematical and Statistical Frontiers, Monash University (david.frazier@monash.edu).

†School of Mathematical Sciences and the Australian Centre of Excellence for Mathematical and Statistical Frontiers, Queensland University of Technology (c.drovandi@qut.edu.au)

The literature on approximate Bayesian inference now includes several competing approximate methods that are often useful in different scenarios. Arguably, the two most common likelihood-free Bayesian methods are approximate Bayesian computation (ABC) (see, e.g., Marin et al., 2012 for a review) and Bayesian synthetic likelihood (BSL) (Wood, 2010, Price et al., 2018).

Both ABC and BSL eschew calculation of the intractable likelihood via simulation from a model that is assumed to have generated the data, and are applicable to any model in which data can be reliably simulated. This simple requirement has led to the application of ABC and BSL in a host of problems in biology, ecology, genetics, economics, finance, and astrophysics; see Sisson et al., 2018 for several examples.

Generally speaking, ABC and BSL are based on a choice of summary statistics that are used to represent the data, and the goal of the approach is to produce draws from the so-called ‘partial-posterior’ based on these statistics. In effect, ABC uses a nonparametric approach to estimate this posterior (see Blum, 2010 for details), while BSL, in its simplest form, employs a parametric assumption. As a result, ABC based-inference can scale poorly with the dimension of the summary statistics. Indeed, most ABC analyses resort to a low-dimensional summary statistic to maintain reasonable computation times, which could lead to a significant increase in parameter uncertainty.

Following the synthetic likelihood approach of Wood (2010), Price et al. (2018) develop a parametric alternative to ABC by constructing a Bayesian version of synthetic likelihood. Unlike ABC, which implicitly estimates a version of the likelihood for the summaries, BSL directly assumes that the joint density of the summary statistics, conditional on the unknown model parameters, is Gaussian with unknown mean and variance. Using independent simulations obtained from the assumed data generating process (DGP), the mean and variance of the summary statistics are then estimated, and used to construct a (simulated) Gaussian likelihood function that is directly inserted into standard Markov Chain Monte Carlo (MCMC) algorithms. Price et al. (2018) demonstrate the BSL approach across several examples, and show that it often performs well in comparison with ABC.

BSL, and approximate Bayes methods more generally, are most often applied in situations where the complexity of the model that is assumed to have generated the observed data renders exact Bayesian inference infeasible. That is, by the very nature of the problems to which BSL is commonly applied, the model is so complex that we can not easily access the DGP and must instead resort to an approximate inference approach. However, while complicated, highly-structured models that allow for vast complexity allow us to explain critical features of the observed data, it is unlikely that any modeler will be able to construct an entirely accurate model that captures all features of the observed data. In short, all models are wrong and the scientist cannot obtain a “correct” one through excessive elaboration (Box, 1976).

The implications of such a statement are particularly worrying in the context of BSL, where the assumed underlying DGP is often very complex. Indeed, applying the above reasoning of Box, it must be the case that the models to which BSL is routinely applied are misspecified representations of the actual, or true, DGP. In such situations, the application of BSL deserves further scrutiny given the recent results of Frazier et al. (2017), which demonstrate that if the model is misspecified, ABC-based inference can be suspect. Given that the principles underlying ABC and BSL are qualitatively similar, further analysis is needed to ensure that BSL does not suffer the same shortcoming of ABC in cases where the model is misspecified.

Through several simulated and empirical examples, we demonstrate that when the assumed

model is misspecified BSL can deliver unreliable results. To circumvent this issue, we propose two novel versions of BSL that can deliver reliable results regardless of whether the model is correctly specified. These robust versions of BSL are based on adjusting the simulated summaries to ensure a form of compatibility between the assumed model and the chosen summary statistics.

The first approach we propose is based on adjusting the mean of the simulated summaries, while the second approach adjusts the variance. We demonstrate both approaches through a series of Monte Carlo experiments and empirical examples. The results of these experiments yields a striking juxtaposition between our new robust approach and standard BSL: when the model is misspecified, standard BSL inference can fail completely, due to extremely low acceptance rates within MCMC sampling, while the robust versions of BSL proposed herein remain well-behaved even under significant model misspecification. In addition, the improved computational efficiency of our approaches allow practitioners to explore sources of model misspecification to inform further model development when using BSL as the inferential tool.

The remainder of the paper is organized as follows. In Section two we give a brief overview of BSL and examine the consequences of model misspecification in the BSL context. Section three presents our robust approach to BSL and considers the theoretical properties of this new method. In particular, we demonstrate that when the model is correctly specified, the robust version of BSL gives the same results as standard BSL, but behaves in a more regular fashion under model misspecification. Section four contains a mix of Monte Carlo and empirical results that demonstrate the performance of this robust BSL approach on simulated and challenging real data, and the potentially poor performance of standard BSL. Section five concludes.

2 Bayesian Synthetic Likelihood and Compatibility

2.1 Bayesian Synthetic Likelihood Framework

We observe data $\mathbf{y} = (y_1, \dots, y_n)^\top$, $n \geq 1$, and denote by P_0^n the true distribution of the observed sample. The true distribution is unknown and instead we consider that the class of probability measures $\{\theta \in \Theta \subset \mathbb{R}^{d_\theta}, n \geq 1 : P_\theta^n\}$, for some value of θ , have generated the data, and denote the corresponding conditional density as $p_n(\cdot|\theta)$. Given prior beliefs over the unknown parameters in the model θ , represented by the probability measure $\Pi(\theta)$, with its density denoted by $\pi(\theta)$, our aim is to produce draws from the exact posterior density

$$\pi(\theta | \mathbf{y}) \propto p_n(\mathbf{y}|\theta)\pi(\theta).$$

In situations where the likelihood is intractable, sampling from $\pi(\theta | \mathbf{y})$ can be computationally costly or infeasible, however, so-called likelihood-free methods can still be used to conduct inference on the unknown parameters θ . The most common implementations of these methods are approximate Bayesian computation (ABC) and Bayesian synthetic likelihood (BSL). Both ABC and BSL generally degrade the data down to a vector of summary statistics and then perform posterior inference on the unknown θ , conditional only on this vector of summary statistics.

More formally, let $\eta(\cdot) : \mathbb{R}^n \rightarrow \mathbb{R}^{d_\eta}$ denote a d_η -dimensional map that represents the chosen summary statistics, and let $\mathbf{z} := (z_1, \dots, z_n)^\top \sim P_\theta^n$ denote data simulated from the model P_θ^n . For $G_n(\cdot|\theta)$ denoting the projection of P_θ^n under $\eta(\cdot) : \mathbb{R}^n \rightarrow \mathbb{R}^{d_\eta}$, with $g_n(\cdot|\theta)$ its corresponding density, the goal of approximate Bayesian methods is to generate samples from the approximate

or ‘partial’ posterior

$$\pi[\theta \mid \eta(\mathbf{y})] \propto g_n[\eta(\mathbf{y}) \mid \theta]\pi(\theta).$$

However, given the complexity of the assumed model, P_θ^n , it is unlikely that the structure of $G_n(\cdot \mid \theta)$ is any more tractable than the original likelihood function $p_n(\mathbf{y} \mid \theta)$. Therefore, simulation-based sampling schemes must be applied to generate samples from $\pi[\theta \mid \eta(\mathbf{y})]$.

The approximate methods of ABC and BSL differ in how $g_n(\cdot \mid \theta)$ is estimated. ABC forms an implicit nonparametric estimator of $g_n(\cdot \mid \theta)$, while BSL uses a parametric or semi-parametric (An et al., 2018) approximation of $g_n(\cdot \mid \theta)$. Given the nonparametric nature of ABC, it is not surprising that the curse of dimensionality inherent in ABC, with regards to the dimension of the summary statistics (Blum, 2010), as well as in the dimension of the parameters themselves (Frazier et al., 2018), ensures that ABC based approaches are heavily restricted in terms of the number and type of summaries they can choose. Due to this need to use only a small number of summary statistics in the analysis, standard ABC-based inference approaches can often lead to imprecise inference.

In contrast to ABC, BSL employs a parametric approximation to $G_n(\cdot \mid \theta)$: BSL replaces - an estimate of - the (intractable) density $g_n(\cdot \mid \theta)$ by a multivariate Gaussian approximation:

$$\mathcal{N}[\eta; \mu_m(\theta), \Sigma_m(\theta)],$$

where $\mu_m(\theta)$ and $\Sigma_m(\theta)$ are simulated estimates of the mean and variance of the summaries (conditional on θ)

$$\mu_m(\theta) = \frac{1}{m} \sum_{i=1}^m \eta(\mathbf{z}^i), \quad \Sigma_m(\theta) = \frac{1}{m} \sum_{i=1}^m [\eta(\mathbf{z}^i) - \mu_m(\theta)] [\eta(\mathbf{z}^i) - \mu_m(\theta)]^\top,$$

and where each simulated data set \mathbf{z}^i , $i = 1, \dots, m$, are generated iid from P_θ^n . The Gaussian approximation is then directly used within an MCMC sampling scheme to sample from the following approximation to the partial posterior, hereafter referred to as the BSL posterior,

$$\begin{aligned} \hat{\pi}[\theta \mid \eta(\mathbf{y})] &\propto \bar{g}_n[\eta(\mathbf{y}) \mid \theta]\pi(\theta), \\ \bar{g}_n[\eta(\mathbf{y}) \mid \theta] &:= \int \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), \Sigma_m(\theta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m). \end{aligned} \tag{1}$$

Due to the parametric nature of equation (1), BSL can often treat summary statistics of larger dimension than ABC and can lead to sharper inference in some cases. While the validity of the Gaussian approximation is often warranted if the underlying summaries satisfy a central limit theorem (Wood, 2010), even in cases where the summary statistics are far from Gaussian, BSL has displayed some insensitivity to violations of this assumption (Price et al., 2018). However, if the statistics are very far from being Gaussian, this can result in a significant loss of sampling efficiency and accuracy, see An et al. (2018) for a demonstration.

2.2 Model Incompatibility and its Consequences

Approximate Bayesian inference methods based on summary statistics implicitly assume that the model being used to generate the simulated summary statistics $\eta(\mathbf{z})$ can replicate the behavior of the observed summary statistics $\eta(\mathbf{y})$. That is, BSL is not required to match every aspect of

the data, but only those features of the data that are captured via the summary statistics $\eta(\mathbf{y})$. This differs from a standard Bayesian framework based on a likelihood, where, under general regularity conditions, the posterior ultimately gives higher probability mass to values of $\theta \in \Theta$ that ensure the Kullback-Liebler (KL) divergence

$$\mathcal{D}(P_0^n \| P_\theta^n) = \int \log \left\{ \frac{p_0^n(\mathbf{y})}{p_n(\mathbf{y}|\theta)} \right\} dP_0^n(\mathbf{y})$$

is as close to zero as possible. When the model is misspecified, i.e., when $P_0^n \neq P_\theta^n$ for any $\theta \in \Theta$, following Kleijn and Van der Vaart (2012), the posterior eventually places increasing mass on the value $\theta^* \in \Theta$ that minimizes the KL-divergence.

In contrast, BSL is not based on trying to find the P_θ^n that is closest to P_0^n in KL divergence but on trying to match simulated and observed summary statistics. Therefore, the meaningful concept of model misspecification in BSL is not KL divergence but that the choice of the assumed model, allied with our specific choice of summary statistics, are able to replicate the behavior of the observed summary statistic.

Let $b_0 := \text{plim} \eta(\mathbf{y})$ and $b(\theta) := \text{plim} \eta(\mathbf{z})$, where plim denotes probability limit. Using the framework of Marin et al. (2014), and following Frazier et al. (2017), we formalize this notion of misspecification as follows.

Definition 1. *The model $P_\theta^n \times \Pi$ and summary statistic map $\eta(\cdot)$ are **compatible** if*

$$\inf_{\theta \in \Theta} \|b(\theta) - b_0\| = 0.$$

Intuitively, compatibility means that, asymptotically, $\eta(\mathbf{y})$ must be in the support of $\mu_m(\theta)$ and implies that, for some value of θ , $\eta(\mathbf{z})$ can replicate the behavior of $\eta(\mathbf{y})$ when \mathbf{z} is simulated under P_θ^n .

By focusing inference on summary statistics, approximate inference methods have a type of in-built robustness against model misspecification: the goal of these methods is not to find the region of the parameter space Θ that allow us to match all the features of the DGP; instead, approximate methods only seek to find regions of Θ that allow us to match the features of the data measured by $\eta(\cdot)$. Therefore, we argue that the researcher should not be concerned about the correct specification of the entire DGP but should instead concern herself with the satisfaction of the above compatibility condition. If this condition is satisfied, even though the model P_θ^n may be misspecified, we can still, in some sense, adequately capture the features of the data that are described by $\eta(\mathbf{y})$, and which are likely to be of primary interest to the analysis.

However, as recently discussed by Frazier et al. (2017), in the context of ABC, when approximate methods are based on a model and summary statistic combination that are not compatible, the resulting inference can be suspect. Given that ABC and BSL are based on very similar principles, it is highly likely that BSL will suffer from the same issues as ABC when the above compatibility condition is not satisfied. While we demonstrate this with several realistic examples in Section 4, we first consider an artificially simple example where BSL should perform very well, but, due to this incompatibility issue, BSL can deliver unreliable inference.

2.3 Consequences of Incompatibility: Toy Normal Model

To demonstrate the performance of BSL under model/summary incompatibility, we consider an artificially simple example: our goal is to conduct inference on the mean parameter θ in the

model

$$y_i = \theta + v_i, v_i \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2).$$

Assume that we believe with certainty that $\sigma = 1$, and so the model we will use to generate data within BSL is given by

$$z_i = \theta + v_i, v_i \stackrel{iid}{\sim} \mathcal{N}(0, 1).$$

However, we will expressly consider the scenario where the assumption $\sigma = 1$ is false and the true DGP for the observed data maintains $\sigma \neq 1$, which implies that the assumed DGP for \mathbf{z} differs from the actual DGP for \mathbf{y} .

We consider as summary statistics the sample mean $\eta_1(\mathbf{y}) = \frac{1}{n} \sum_{i=1}^n y_i$ and the sample variance $\eta_2(\mathbf{y}) = \frac{1}{n-1} \sum_{i=1}^n (y_i - \eta_1(\mathbf{y}))^2$. When $\sigma \neq 1$, the model is not compatible with the variance summary, and Definition 1 will not be satisfied.

While this example is artificially simple, it will be useful to demonstrate the potential issues with BSL when the model is misspecified. Indeed, regardless of the misspecification, one might think that BSL should perform well: the first summary statistic, $\eta_1(\mathbf{y})$, is Gaussian, the statistic $\eta_2(\mathbf{y})$ satisfies the central limit theorem, and $\eta_1(\mathbf{y})$ is sufficient for θ .

For this experiment, we generate simulated data sets for \mathbf{y} such that each data set corresponds to a different value of σ , with σ taking values from $\sigma = 0.2$ to $\sigma = 2$ with evenly spaced increments of 0.1. Across all the data sets we fix the random numbers used to generate the ‘‘observed’’ data and only change the value of σ to isolate the impact of model misspecification. The sample size across the experiments is taken to be $n = 50$. Our prior for θ is $\theta \sim \mathcal{N}(0, 10)$.

We generate $m = 10,000$ simulated data sets to estimate $\mu_m(\theta)$ and $\Sigma_m(\theta)$, and carry out BSL using a random-walk Metropolis algorithm, where the variance of the proposal has been tuned to be the true posterior variance. The sampler is initialized at the true value of the parameters, $\theta = 1$, and run for 10,000 iterations. Since the chain is started at the truth, we retain all values for inference.

First, we analyze how the BSL posterior mean responds to varying levels of model misspecification. Panel A of Figure 1 plots the posterior means and corresponding 95% credible intervals for each data set and Panel B gives the acceptance rates across the various data sets. From the results in Panel A, it seems that BSL displays resilience to model misspecification, except perhaps for some instability at large levels of misspecification. However, analyzing the results in Panel B, we see that the stability implied by the results in Panel A is not genuine. In particular, the results in Panel B demonstrate that at high levels of misspecification, σ either very small or very large, BSL has a very difficult time exploring the parameter space. As a consequence, the MCMC chain stays stuck for long periods of time and the acceptance rates in BSL plummet from a peak of about 70%, to a low of 0.85% at $\sigma = 2$ despite the large number of model simulations used to estimate the synthetic likelihood. At higher levels of misspecification, the MCMC chain rarely moves and so most values simply reflect the starting value of $\theta = 1$, which gives the results in Panel A of Figure 1 the false sense that BSL is performing well across all levels of misspecification.

In this scenario, the poor performance of BSL can be directly traced to the incompatibility of the second summary statistic $\eta_2(\mathbf{y})$: no matter the value of θ , we can never hope to match the value of $\eta_2(\mathbf{y})$. This gap between $\eta_2(\mathbf{y})$ and $\eta_2(\mathbf{z})$ causes the MCMC chain to stick, and results in unreliable inference. Moreover, with real data it is not clear *a priori* how one could hope to reliably detect such incompatibility issues, since the true DGP is unknown. To circumvent this poor performance of BSL in contexts where the summaries are not compatible, in the following

section, we propose a robust version of BSL that delivers reliable inference when the summaries are not compatible and which is capable of detecting precisely which summaries, if any, are incompatible.




Figure 1: Panel A: BSL Posterior means (blue) across all levels of misspecification. Panel B: acceptance rates across all levels of misspecification.

3 Robust Bayesian Synthetic Likelihood

We propose two possible strategies for conducting inference using BSL when the model and summaries are incompatible. The first approach augments the mean of the simulated summaries with additional free parameters, while the second approach augments the variance of the simulated summaries with additional free parameters. Both specifications will allow us to conduct robust inference on the model parameters and determine which of the summaries may be incompatible with the data. Throughout the remainder, we refer to these approaches as “mean” and “variance” robust BSL (R-BSL).

3.1 Mean Adjustment and Prior Specification

Incompatibility implies that the observed statistics $\eta(\mathbf{y})$ are not in the support of the simulated mean $\mu_m(\theta)$, for any $\theta \in \Theta$, with probability converging to one. Therefore, one approach to create a BSL procedure that is robust to this incompatibility issue is to adjust the vector of simulated means. This can be accomplished by adding to $\mu_m(\theta)$ an additional free parameter Γ , where $\Gamma = (\gamma_1, \dots, \gamma_{d_\eta})^\top \in \mathcal{G} \subset \mathbb{R}^{d_\eta}$, so that $\eta(\mathbf{y})$ is always in the support of this new simulated mean. Defining the joint vector of unknown parameters as $\zeta := (\theta^\top, \Gamma^\top)^\top \in \Theta \times \mathcal{G} \subset \mathbb{R}^{d_\theta} \times \mathbb{R}^{d_\eta}$, we define the vector of simulated means for use in BSL as

$$\phi_m(\zeta) = \mu_m(\theta) + \text{diag}[\Sigma_m^{1/2}(\theta)] \Gamma.$$

Note that, by considering the scaled adjustment term, $\text{diag}[\Sigma_m^{1/2}(\theta)]\Gamma$, we ensure that these components are measured in the same units as $\mu_m(\theta)$, which allows us to treat Γ as if they were unitless.

Given this linear adjustment, and under weak conditions on the summary statistics and the parameter space $\Theta \times \mathcal{G}$, it is simple to see that $\phi_m(\zeta)$ will be compatible with $\eta(\mathbf{y})$ for any prior choice on Γ such that each individual component of Γ has support over \mathbb{R} .

Denote the prior on ζ by $\pi(\zeta)$. Following Price et al. (2018), the augmented BSL target, which we refer to as the Robust BSL-mean (R-BSL-M) posterior, is to generate samples from

$$\hat{\pi}[\zeta \mid \eta(\mathbf{y})] \propto \bar{g}_n[\eta(\mathbf{y}) \mid \zeta] \pi(\zeta),$$

where

$$\bar{g}_n[\eta(\mathbf{y}) \mid \zeta] = \int \mathcal{N}[\eta(\mathbf{y}); \phi_m(\zeta), \Sigma_m(\theta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m). \quad (2)$$

Prior Choice: Laplace Prior

To ensure that the observed summary $\eta(\mathbf{y})$ is in the support of $\phi_m(\zeta)$, our prior on the components of Γ should allow for $\text{diag}[\Sigma_m^{1/2}(\theta)]\Gamma$ to escape the support of $\mu_m(\theta)$ with large probability. However, given that some components of the original $\mu_m(\theta)$ are likely compatible with some components of $\eta(\mathbf{y})$, we want to make sure that $\text{diag}[\Sigma_m^{1/2}(\theta)]\Gamma$ does not unduly perturb the components that are compatible. Therefore, we should choose a prior that places the vast majority of its mass near the origin. In this way, our prior choice for Γ should induce sparsity in the components of Γ : only the components of Γ that correspond to incompatible summaries should receive significant posterior probability away from the origin, while the components of Γ that correspond to compatible summaries should have the majority of their posterior mass near the origin.

With these dual requirements in mind, and given that each component of Γ has the same prior scale, we propose to follow the Bayesian Lasso literature (Park and Casella, 2008) and use independent Laplace (i.e., double-exponential) priors for each component of Γ , with fixed location 0 and common scale $\lambda > 0$:

$$\pi(\Gamma) := \prod_{j=1}^{d_\eta} \lambda e^{-\lambda|\gamma_j|} = \lambda^{d_\eta} e^{-\lambda \sum_{j=1}^{d_\eta} |\gamma_j|}. \quad (3)$$

When convenient, we denote this prior by $\text{La}(0, \lambda)$. The Laplace prior for Γ guarantees that the majority of prior mass for γ_j is near the origin, but has thick enough tails so that $\phi_m(\zeta)$ is compatible with virtually any $\eta(\mathbf{y})$ that would be used in practice.

The hyper-parameter λ should be chosen so that the prior support of $\pi(\Gamma)$ compliments the support of $\mu_m(\theta)$. That is, λ should be chosen so that the prior support of Γ is larger than the support of $\mu_m(\theta)$, which will allow us to detect deviations from compatibility, but not so large as to cause the statistic $\phi_m(\zeta)$ to have heavy tails. Indeed, if the tails of $\phi_m(\zeta)$ are too heavy, the implicit normality assumption made in BSL will be violated and can result in inefficient sampling.

Since there is no reason to believe a priori that θ and Γ are related, we take as our overall prior on ζ in R-BSL-M to be

$$\pi(\zeta) := \pi(\theta) \text{La}(0, \lambda).$$

3.2 Variance Compatibility and Prior Specification

Compatibility requires that $\eta(\mathbf{y})$ be in the support of $\mu_m(\theta)$, for some $\theta \in \Theta$. While one approach to ensure this compatibility is to adjust the mean of the simulated summaries, an alternative is to inflate the variance of the simulated summaries to ensure that $\eta(\mathbf{y})$ is always in the support of $\mu_m(\theta)$.

Under regularity conditions and for fixed m , it is likely to be the case that the centered statistic $Z_{n,m}(\theta) := \{\mu_m(\theta) - b(\theta)\}$ behaves as $Z_{n,m}(\theta) = O_P(1/(m\sqrt{n}))$, with the variance of $Z_{n,m}(\theta)$ decreasing like $1/(mn)$. Consequently, for n (or m) large enough, if for a given value of θ the statistic $\{\eta(\mathbf{y}) - b(\theta)\}$ is more than a few standard deviations (as measured by $\Sigma_m^{1/2}(\theta)$) away from $\{\mu_m(\theta) - b(\theta)\}$, we can effectively view the summaries as being incompatible. In such cases, the BSL posterior is likely to have low acceptance rates in this region of the parameter space.

Given this characterization, an alternative approach to ensure that $\mu_m(\theta)$ is compatible with $\eta(\mathbf{y})$ is to artificially inflate the variance $\Sigma_m(\theta)$ so that the variance of $Z_{n,m}(\theta)$ never completely collapses to zero, and thus $\{\eta(\mathbf{y}) - b(\theta)\}$ can always be found in the support of $Z_{n,m}(\theta)$, albeit perhaps with small probability. More specifically, we propose to artificially inflate the variance used within BSL by adding to $\Sigma_m(\theta)$ the free parameters $\Gamma = (\gamma_1, \dots, \gamma_{d_\eta})^\top$.

Recalling $\zeta = (\theta^\top, \Gamma^\top)^\top$, a robust BSL procedure based on adjusting the variance can be implemented by re-defining the variance of the simulated statistics used within BSL to be

$$V_m(\zeta) := \Sigma_m(\theta) + \begin{pmatrix} [\Sigma_m(\theta)]_{11}\gamma_1^2 & 0 & \dots & 0 \\ 0 & [\Sigma_m(\theta)]_{22}\gamma_2^2 & \dots & 0 \\ \vdots & \dots & \ddots & \vdots \\ 0 & \dots & \dots & [\Sigma_m(\theta)]_{d_\eta d_\eta}\gamma_{d_\eta}^2 \end{pmatrix}, \quad (4)$$

where $[\Sigma_m(\theta)]_{ii}$ denotes the (i, i) element of $\Sigma_m(\theta)$. Given the structure of $V_m(\zeta)$, γ_i can be interpreted as an inflation factor operating on the standard deviations of the original BSL variance. An equivalent interpretation is that the i -th, $i = 1, \dots, d_\eta$, BSL variance is multiplied by the factor $1 + \gamma_i^2$. Using $V_m(\zeta)$ in place of $\Sigma_m(\theta)$ in the BSL posterior target, (1), and for $\pi(\zeta)$ an appropriate prior on ζ , the Robust BSL-variance (R-BSL-V) posterior is given as:

$$\hat{\pi}[\zeta|\eta(\mathbf{y})] \propto \bar{g}_n[\eta(\mathbf{y})|\zeta]\pi(\zeta),$$

where

$$\bar{g}_n[\eta(\mathbf{y})|\zeta] = \int \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), V_m(\zeta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) | \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m). \quad (5)$$

Prior Choice: Exponential Prior

Note that, by considering the standardization in (4), we ensure that each γ_i has the same scale and can be considered as unit-less. Moreover, similar to the case of the mean adjustment BSL approach, there is no reason to believe there is any *a priori* dependence between θ and Γ , so we can consider independent priors, i.e., $\pi(\zeta) := \pi(\theta)\pi(\Gamma)$. While several prior choices exist for Γ , following the arguments for the prior choice in the mean adjustment procedure, we need to choose a prior for the components of Γ so that there is a large amount of prior mass near the origin, and enough mass out in the tails to ensure we can detect incompatible summaries.

To this end, we consider independent exponential priors for each component γ_i , ($i = 1, \dots, d_\eta$), with common rate $\lambda > 0$:

$$\pi(\Gamma) := \prod_{i=1}^{d_\eta} \lambda e^{-\lambda \gamma_i} = \lambda^{d_\eta} e^{-\lambda \sum_{i=1}^{d_\eta} \gamma_i}.$$

The hyper-parameter λ should be chosen so that a large amount of prior mass is close to the origin, so as not to over-inflate the variance of the simulated summaries that are compatible.

While this choice of prior is not, strictly speaking, a shrinkage prior, it is still the case that we should observe some shrinkage like behavior for summaries that are compatible. That is, for the summaries that are compatible, this additional inflation by Γ is unnecessary and we expect that, for appropriate choices of λ , the addition of this component will not greatly affect the corresponding components in the variance. In contrast, for the summaries that are not compatible, this adjustment term is critical to ensure that the variance of the summaries is large enough to contain the observed summary $\eta(\mathbf{y})$.

3.3 Sampling Robust BSL

As demonstrated in Price et al. (2018), the standard BSL target posterior is given by

$$\hat{\pi}[\zeta \mid \eta(\mathbf{y})] \propto \bar{g}_n[\eta(\mathbf{y}) \mid \theta] \pi(\theta), \quad (6)$$

where

$$\bar{g}_n[\eta(\mathbf{y}) \mid \theta] = \int \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), \Sigma_m(\theta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m).$$

Price et al. (2018) use a Metropolis-Hastings algorithm to sample from (6) that proposes $\theta^* \sim q(\cdot \mid \theta)$ according to a Markov transition, and estimating $\bar{g}_n[\eta(\mathbf{y}) \mid \theta^*]$ unbiasedly through a single draw from $\prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta^*]$ and evaluating $\mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), \Sigma_m(\theta)]$. Using pseudo-marginal MCMC arguments of Andrieu and Roberts (2009), substituting this estimator into a Metropolis-Hastings algorithm produces an algorithm that targets (6).

Our robust BSL methods operate on an extended state space over θ and Γ with target distribution

$\hat{\pi}[\zeta \mid \eta(\mathbf{y})] \propto \bar{g}_n[\eta(\mathbf{y}) \mid \zeta] \pi(\theta) \pi(\Gamma)$, where

$$\bar{g}_n[\eta(\mathbf{y}) \mid \zeta] = \int \mathcal{N}[\eta(\mathbf{y}); \phi_n(\zeta), \Sigma_n(\theta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m) \quad (\text{Mean Adjustment}),$$

$$\bar{g}_n[\eta(\mathbf{y}) \mid \zeta] = \int \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), V_m(\zeta)] \left\{ \prod_{i=1}^m g_n[\eta(\mathbf{z}^i) \mid \theta] \right\} d\eta(\mathbf{z}^1) \cdots d\eta(\mathbf{z}^m) \quad (\text{Variance Adjustment}).$$

To sample these target distributions we use a component-wise MCMC algorithm that updates, in turn, θ conditional on Γ and then Γ conditional on θ . The update for θ is the same as in standard BSL, but where the adjusted mean or inflated variance is computed as appropriate using the current value of Γ . As before, the update for θ involves generating m model simulations, $\eta(\mathbf{z}^i)$, $i = 1, \dots, m$.

The update for Γ holds the currently accepted model simulations fixed, and thus $\mu_m(\theta)$ and $\Sigma_m(\theta)$ are fixed within the update step for Γ . Each component of Γ , γ_j , for $j = 1, \dots, d_\eta$, is updated separately, conditional on the current values of the remaining components (denoted $\gamma_{/j}$). The full conditional distribution for γ_j is given by

$$\pi(\gamma_j^* | \theta, \mu_m(\theta), \Sigma_m(\theta), \gamma_{/j}) \propto \mathcal{N}[\eta(\mathbf{y}); \phi_m(\zeta^*), \Sigma_m(\theta)] \pi(\gamma_j^*) \quad (\text{Mean Adjustment})$$

$$\pi(\gamma_j^* | \theta, \mu_m(\theta), \Sigma_m(\theta), \gamma_{/j}) \propto \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta), V_m(\zeta^*)] \pi(\gamma_j^*) \quad (\text{Variance Inflation}),$$

where γ_j^* is a realisation of γ_j and $\zeta^* = [\theta, \gamma_j^*, \gamma_{/j}]$.

We sample this full conditional distribution using a slice sampler, and, in particular, the “stepping out” and “shrinkage” procedures detailed in Neal (2003). The appeal of the slice sampler is that the acceptance probability is one, and thus there are no tuning parameters that can affect the statistical efficiency. However, a stepping out width needs to be selected, which can impact the speed of the slice sampler. Given that the components of Γ are eventually scaled by the summary statistic standard deviation, and given that our prior choices effectively penalise large values, we expect each component of Γ to be $O(1)$. Thus, we set the stepping out width to be 1, except for the lower bound of γ_j in the variance inflation, which is immediately set to 0; and hence the stepping out procedure is not required. We find this choice of width to be suitable, and since updating Γ does not require any model simulations, the slice sampler is very fast. Hence, importantly, our robust BSL methods do not require any additional tuning and the run-time per iteration for non-trivial applications is not noticeably slower. The full MCMC algorithm to sample from the R-BSL posteriors is provided in Algorithm 1.

It is important to note that under either R-BSL approach, we recover the original BSL target when

$$\gamma_1 = \dots = \gamma_{d_\eta} = 0.$$

Therefore, if the model P_θ^n can generate summaries $\eta(\mathbf{z})$ that match $\eta(\mathbf{y})$, under an appropriate prior specification, the posterior $\hat{\pi}[\Gamma | \eta(\mathbf{y})]$ should not differ substantially from the prior, with most of the posterior mass located near the origin.

While obviously related, the two R-BSL adjustment procedures operate on very different principles. The R-BSL-M approach is based on finding joint values (θ, Γ) under which $\|\phi_m(\zeta) - \eta(\mathbf{y})\|$ is “small”, which we can interpret as, for fixed θ , finding values of Γ so that $\mu_m(\theta) + \text{diag}[\Sigma_m^{1/2}(\theta)] \Gamma$ matches $\eta(\mathbf{y})$. In contrast, the R-BSL-V procedure is not required to find values of θ such that $\mu_m(\theta) - \eta(\mathbf{y})$ is small. Rather, for a fixed value of θ , the R-BSL-V approach seeks a value of Γ so that the weighted norm

$$\|\mu_m(\theta) - \eta(\mathbf{y})\|_{V_m(\zeta)} = (\mu_m(\theta) - \eta(\mathbf{y}))^\top V_m^{-1}(\zeta) (\mu_m(\theta) - \eta(\mathbf{y}))$$

is “small”. Therefore, if $\|\mu_m(\theta) - \eta(\mathbf{y})\|$ is already small, no variance inflation is needed, and the resulting value of Γ should be close to zero; if instead $\|\mu_m(\theta) - \eta(\mathbf{y})\|$ is “large”, the term $\|\mu_m(\theta) - \eta(\mathbf{y})\|_{V_m(\zeta)}$ can always be made small by choosing a large value of Γ .

3.4 Theoretical Properties of R-BSL

If the model is compatible, what behavior should we expect from the R-BSL approach? Given the priors used in R-BSL, which place the majority of their mass near the origin, one would hope that when the compatibility condition (Definition 1) is satisfied, the introduction of the additional

Input : Summary statistic of the data, $\eta(\mathbf{y})$, the prior distribution, $\pi(\theta)$, the proposal distribution q , the number of iterations, T , and the initial value of the chain, θ^0 , Γ^0 .

Output: MCMC sample $(\theta^0, \theta^1, \dots, \theta^T)$ and $(\Gamma^0, \Gamma^1, \dots, \Gamma^T)$ from the robust BSL posterior. Some samples can be discarded as burn-in if required.

- 1 Estimate $\mu_m(\theta^0)$ and $\Sigma_m(\theta^0)$ via m independent model simulations at θ^0
- 2 Compute $\phi_m(\zeta^0) = \mu_m(\theta^0) + \text{diag} \left[\Sigma_m^{1/2}(\theta^0) \right] \Gamma^0$ (mean adjustment) or $V_m(\zeta^0)$ (variance inflation) defined in (4).
- 3 Compute robust synthetic likelihood $L^0 = \mathcal{N}[\eta(\mathbf{y}); \phi_m(\zeta^0), \Sigma_m(\theta^0)]$ (mean adjustment) or $L^0 = \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta^0), V_m(\zeta^0)]$ (variance inflation)
- 4 **for** $i = 1$ to T **do**
- 5 | %%%% Obtain Γ^i via the following, which does not require any model simulations
- 6 | **for** $j = 1$ to d_η **do**
- 7 | | Update γ_j^i with a slice sampler with target density
 | | $\pi(\gamma_j^i | \theta^{i-1}, \mu_m(\theta^{i-1}), \Sigma_m(\theta^{i-1}), \gamma_1^i, \dots, \gamma_{j-1}^i, \gamma_{j+1}^{i-1}, \dots, \gamma_{d_\eta}^{i-1}) \propto$
 | | $\mathcal{N}[\eta(\mathbf{y}); \phi_m(\zeta^*), \Sigma_m(\theta^{i-1})] \pi(\gamma_j^i)$ (mean adjustment) or
 | | $\mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta^{i-1}), V_m(\zeta^*)] \pi(\gamma_j^i)$ (variance inflation) where
 | | $\zeta^* = [\theta^{i-1}, \gamma_1^i, \dots, \gamma_{j-1}^i, \gamma_j^i, \gamma_{j+1}^{i-1}, \dots, \gamma_{d_\eta}^{i-1}]$
- 8 | **end**
- 9 | Denote $L^i = \mathcal{N}[\eta(\mathbf{y}); \phi_m([\theta^{i-1}, \Gamma^i]), \Sigma_m(\theta^{i-1})]$ (mean adjustment) or
 | $L^i = \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta^{i-1}), V_m([\theta^{i-1}, \Gamma^i])]$ (variance inflation)
- 10 | %%%% Update θ^i conditional on Γ^i
- 11 | Draw $\theta^* \sim q(\cdot | \theta^{i-1})$
- 12 | Estimate $\mu_m(\theta^*)$ and $\Sigma_m(\theta^*)$ via m independent model simulations at θ^*
- 13 | Compute $\phi_m(\zeta^*)$ (mean adjustment) or $V_m(\zeta^*)$ (variance inflation) defined in (4),
 | where $\zeta^* = [\theta^*, \Gamma^i]$
- 14 | Compute proposed adjusted synthetic likelihood $L^* = \mathcal{N}[\eta(\mathbf{y}); \phi_m(\zeta^*), \Sigma_m(\theta^*)]$ (mean
 | adjustment) or $L^* = \mathcal{N}[\eta(\mathbf{y}); \mu_m(\theta^*), V_m(\zeta^*)]$ (variance inflation)
- 15 | Compute Metropolis-Hastings ratio:
 |
$$r = \frac{L^* \pi(\theta^*) q(\theta^{i-1} | \theta^*)}{L^i \pi(\theta^{i-1}) q(\theta^* | \theta^{i-1})}$$
- 16 | **if** $\mathcal{U}(0, 1) < r$ **then**
- 17 | | Set $\theta^i = \theta^*$, $\mu_m(\theta^i) = \mu_m(\theta^*)$ and $\Sigma_m(\theta^i) = \Sigma_m(\theta^*)$
- 18 | **else**
- 19 | | Set $\theta^i = \theta^{i-1}$, $\mu_m(\theta^i) = \mu_m(\theta^{i-1})$ and $\Sigma_m(\theta^i) = \Sigma_m(\theta^{i-1})$
- 20 | **end**
- 21 **end**

Algorithm 1: Robust MCMC BSL.

parameters Γ should not influence the BSL posterior for θ . We formalize the above intuition and demonstrate that when Definition 1 is satisfied the R-BSL posterior for the adjustment components will converge to the prior, while the components for θ will collapse onto the value in Θ for which $b(\theta) = b_0$.

To simply arguments and derivations, we demonstrate this result in the case of the “idealized” BSL posterior, where we replace $\mu_m(\theta)$ and $\Sigma_m(\theta)$ with their idealized counterparts

$$b(\theta) = \mathbb{E}[\eta(\mathbf{y})|\theta] \text{ and } A_n(\theta) := \left\{ \mathbb{E} \left[\{\eta(\mathbf{y}) - b(\theta)\}^2 \mid \theta \right] \right\}^{1/2}.$$

In this simplified scenario, the “model likelihood” for BSL become

$$\bar{g}_n(\eta|\theta) = \mathcal{N} \left[\eta; b(\theta), A_n(\theta)^\top A_n(\theta) \right].$$

Before presenting the formal result, we must state some notation. Define P_0^n as the true distribution generating \mathbf{y} . The map $\eta : \mathbb{R}^n \rightarrow \mathbb{R}^d$, which defines the summary statistics used in the procedure, satisfies $\eta(\mathbf{y}) \sim G_n^0$, where G_n^0 denotes the projection of P_0^n under the map η , and denote by g_n^0 the density of G_n^0 . Likewise, recall that $G_n[\cdot|\theta]$ denotes the projection of P_n^θ under the map η . For real-valued sequences $\{a_n\}_{n \geq 1}$ and $\{b_n\}_{n \geq 1}$: $a_n \lesssim b_n$ denotes $a_n \leq Cb_n$ for some finite $C > 0$ and all n large, $a_n \gtrsim b_n$ denotes $a_n \geq Cb_n$ for some finite $C > 0$, and $a_n \asymp b_n$ implies $a_n \lesssim b_n$ and $a_n \gtrsim b_n$. The terms O_P and o_P have their usual connotations and the notation \Rightarrow denotes weak convergence in distribution.

We impose the following regularity conditions.

Assumption 1. *There exists a sequence of positive real numbers v_n diverging to ∞ such that, for some distribution Q on \mathbb{R}^d and some vector $b_0 \in \mathbb{R}^d$,*

$$v_n [\eta(\mathbf{y}) - b_0] \Rightarrow Q, \text{ under } P_n^0.$$

Assumption 2. *(i) The sequence $\{v_n\}_{n \geq 1}$ is such that, for all $\theta \in \Theta$ and some n large enough, there exists constants c_1, c_2 , $c_1 \leq c_2$, satisfying: $0 < c_1 \leq \|v_n A_n(\theta)\| \leq c_2 < \infty$; (ii) For all $\theta \in \Theta$ and some n large enough, the $d \times d$ -matrix $A_n(\theta)$ is continuous in θ .*

Assumption 3. *There exists a deterministic map $b : \Theta \rightarrow \mathcal{B}$, such that, for all $\theta \in \Theta$, and for constants $\alpha, u_0 > 0$, for all $0 < u < u_0 v_n$,*

$$G_n [\|v_n \{\eta(\mathbf{y}) - b(\theta)\}\| > u \mid \theta] \leq c(\theta)u^{-\alpha},$$

uniformly for $n \geq 1$ and where $\int_{\Theta} c(\theta)\pi(\theta)d\theta = O(1)$.

Assumption 4. *(i) There exists some $\tau > 0$ such that, for all $0 < u < u_0 v_n$, the prior probability satisfies*

$$\Pi [\|b(\theta) - b_0\| \leq u] \asymp u^\tau.$$

(ii) The prior density $\pi(\theta)$ is continuous and satisfies $\pi(\theta_0) > 0$.

Assumption 5. *(i) The map $\theta \mapsto b(\theta)$ is continuous and injective, with $b(\theta_0) = b_0$ for some $\theta_0 \in \Theta$, and satisfies: $\|\theta - \theta_0\| \leq L\|b(\theta) - b_0\|^\kappa$ on some open neighbourhood of θ_0 with $L > 0$ and $\kappa > 0$.*

Assumption 6. *If Assumption 5 is satisfied, for any $\epsilon > 0$, there exists $u, \delta > 0$ and a set V_n such that for all $\theta \in \{\theta : \|b(\theta) - b_0\| \leq uv_n^{-1}\}$*

$$V_n \subset \{\eta \in \mathbb{R}^d : g_n^0(\eta) \lesssim g_n(\eta | \theta)\} \text{ where } P_n^0(V_n^c) < \epsilon.$$

The above assumptions are similar to those imposed by Marin et al. (2014) in demonstrating the behavior of Bayes factors in situations where inference is conditioned on summary statistics, as opposed to the entire data set. In particular, Assumptions 1, 3 and 6 are virtually identical to their key assumptions, and we refer the interested reader to that paper for specific discussion of these assumptions. However, Assumptions 2, 4 and 5 differ from those in Marin et al. (2014): Assumption 2 is specific to the nature of BSL and requires that the variance matrix used in formulating the BSL posterior be well-behaved, uniformly in θ ; Assumption 3 is a condition on the tails of the prior used in BSL, and is much stronger than the corresponding assumption used in Marin et al. (2014) but simpler to interpret; lastly, Assumption 5 is specific to BSL and is required to ensure the simulated posterior mean is capable of replicating the asymptotic mean of the observed summary statistics. Assumption 1 together with Assumption 5, place conditions on the model that ensure the compatibility condition in Definition 1 is satisfied.

The following result, the proof of which is given in the Appendix, describes the theoretical behavior of the R-BSL posterior under the above assumptions.

Proposition 1. *Under Assumption 1-6, for any $t > 0$:*

$$\Pi[\|\Gamma\| > t | \eta(\mathbf{y})] = \Pi[\|\Gamma\| > t] + o_P(1).$$

Proposition 1 demonstrates that when Definition 1 is satisfied, the R-BSL posteriors for the adjustment components Γ converge to the prior. This implies that under correct model specification the resulting posteriors for the Γ components will not asymptotically concentrate on the origin. As discussed in Bhattacharya et al. (2012), this is not surprising given the relatively mild shrinkage priors placed on the adjustment components. We conjecture that if stronger shrinkage priors were employed, posterior concentration, toward the origin, for these components could also be achieved. However, the use of these more complex priors could create issues within the sampling. Therefore, we believe that these mild shrinkage priors strike a good balance between theoretical guarantees and practical implementation.

A key step in the proof of Proposition 1 is the demonstration that under Definition 1 the BSL posterior for θ concentrates onto θ_0 , the value of θ at which $b_0 = b(\theta)$, at rate $1/v_n$. This result is given as Lemma 1 in the Appendix. We note that this result is of interest in its own right, and complements the existing results on the theoretical behavior of BSL obtained in Nott et al. (2019).

When the compatibility condition in Definition 1 is not satisfied, the R-BSL posterior will deviate from the prior. Therefore, the result of Proposition 1 can be used to determine the level of model misspecification by comparing the difference between the R-BSL posterior for Γ and the prior for Γ . While visual detection will often be enough to determine if any meaningful differences between these two exist, any norm on the space of probability measures could be used to quantify this discrepancy.

By analyzing the posterior elements of Γ that differ from the prior, we can deduce precisely which of the summary statistics the assumed model can not match, i.e., which summaries are not compatible. This information can then be incorporated into subsequent modeling steps to

construct a model that can more adequately capture the observed data. In this way, not only is R-BSL a robust tool for inference, but it can also be used as a model criticism device to help researchers locate discrepancies between the assumed model and the observed data.

4 Examples

In this section, across a variety of examples, we demonstrate the significantly improved inferences and computational efficiency that can be achieved with R-BSL. The first two examples are relatively straightforward and based on simulated data, allowing us to compare with the ground truth and perform more comprehensive simulations. The following two examples are challenging applications in ecology and biology and involve real data.

4.1 Returning to the Normal Example

In this section, we compare the performance of BSL and R-BSL under model misspecification in the toy normal example. Recall that our goal is inference on θ in the model

$$y_i = \theta + v_i, \quad v_i \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2),$$

where we explicitly assume that $\sigma = 1$, and generate data in BSL according to

$$z_i = \theta + v_i, \quad v_i \stackrel{iid}{\sim} \mathcal{N}(0, 1).$$

We consider as summary statistics the sample mean $\eta_1(\mathbf{y}) = \frac{1}{n} \sum_{i=1}^n y_i$ and the sample variance $\eta_2(\mathbf{y}) = \frac{1}{n-1} \sum_{i=1}^n (y_i - \eta_1(\mathbf{y}))^2$. The sample size across all experiments is taken to be $n = 50$, and our prior is $\theta \sim \mathcal{N}(0, 10)$. Recall that, under this setting, we observed that the BSL could lead to unreliable inferences when σ^2 was much larger or smaller than unity.

For the experiments in this section, we use precisely the same “observed” data sets in Section 2.3, but also apply the two different versions of the R-BSL approach, based on the mean and variance adjustments, to the data sets. Similar to BSL in the previous experiment, for both R-BSL procedures we use a random-walk Metropolis algorithm that is initialized at the true value of the parameters, $\theta = 1$, and that we run for 10,000 iterations. Since the chain is started at the truth, we retain all values for inference.

For the mean adjustment approach, our prior on $\Gamma = (\gamma_1, \gamma_2)^\top$ is independent $\text{La}(0, \lambda)$ with $\lambda = 0.5$ across both parameters, which places the vast majority of prior mass between -3 and 3. For the variance adjustment, we give each parameter independent $\text{Exp}(\lambda)$ priors and choose λ so that the prior mean is 0.3. Figure 2 presents the results and mirrors the output in Figure 1 but for the robust versions of BSL.

The results in Panels A and B of Figure 2 demonstrate that the R-BSL approaches displays resilience to model misspecification. Moreover, the results in Panel C demonstrates that the acceptance rate issue that plagued the standard BSL approach is not in evidence for the R-BSL approaches. Indeed, the acceptance rate for the variance adjustment R-BSL approach is nearly unaffected by the level of model misspecification, while the mean adjustment version does display some degradation for $\sigma > 1.5$, but still maintains acceptance rates above 5% in all cases. In comparison, when $\sigma = 2$, the standard BSL acceptance rate was 0.85%.

Figure 3 displays the resulting posterior densities for Γ across the two adjustment procedures, and across all levels of misspecification. Panels A and B give the results for the mean adjustment




Figure 2: Panels A and B give the posterior means and 95% credible sets for the two different R-BSL procedures. Panel C gives the acceptance rates of the different R-BSL procedures and compares them with the standard BSL approach across the different data sets.

approach, and correspond to the components γ_1 and γ_2 , respectively, while panels C and D give the results for the variance adjustment approach. For comparison purposes, the black figure in each panel represents the prior densities, and the color-coding in each figure represents the level of misspecification, and where the red colored density encodes correct model specification (i.e., $\sigma = 1$).

Focusing on Panel A, we see that the posterior densities for the γ_1 component, which captures our ability to match the first observed statistic (the mean), are indistinguishable from the prior across all the observed data sets, which implies that we can match the mean of this model regardless of model misspecification. In contrast, in Panel B we see that the second component, which captures our ability to capture the second observed statistics (the variance), looks nothing like the prior, except perhaps at low levels of misspecification.

Panels C and D describe precisely the same story as in Panels A and B but correspond to γ_1 and γ_2 in the variance adjustment approach. That is, in the case of the variance adjustment BSL approach (R-BSL-V), under correct specification ($\sigma = 1$) the posteriors are indistinguishable from the priors, we are easily able to detect departures from compatibility for the second summary statistic, and the posteriors for the first adjustment term remain indistinguishable from the prior across all data sets. These striking results demonstrate that both R-BSL approaches are capable of producing meaningful inference on the unknown parameter and are capable of determining which features of the model we are not able to replicate.




Figure 3: Marginal posteriors for adjustment components. Panels A and B correspond to the mean adjustment R-BSL approach, while Panels C and D correspond to the variance adjustment R-BSL approach. Panels A and C correspond to the components for the first summary, and panels B and D for the second summary.

4.2 Moving Average Model

To further demonstrate this methodology, we turn to a common toy examples used to demonstrate approximate inference methodology, the moving average model of order two MA(2).

Assume the researcher believes \mathbf{y} is generated according to an MA(q) model:

$$z_t = e_t + \sum_{i=1}^q \theta_i e_{t-i},$$

where, say, $e_t \sim \mathcal{N}(0, 1)$ i.i.d and $\theta_1, \dots, \theta_q$ are such that the roots of the polynomial

$$p(x) = 1 - \sum_{i=1}^q \theta_i x^i$$

all lie outside the unit circle. Specializing this model to the case where $q = 2$, we have that

$$z_t = e_t + \theta_1 e_{t-1} + \theta_2 e_{t-2}, \tag{7}$$

and the unknown parameters $\theta = (\theta_1, \theta_2)^\top$ are assumed to obey

$$-2 < \theta_1 < 2, \theta_1 + \theta_2 > -1, \theta_1 - \theta_2 < 1. \quad (8)$$

Our prior information on $\theta = (\theta_1, \theta_2)^\top$ is uniform over the invertibility region in (8). A useful choice of summary statistics for the MA(2) model are the sample autocovariances $\eta_j(\mathbf{z}) = \frac{1}{T} \sum_{t=1+j}^T z_t z_{t-j}$, for $j = 0, 1, 2$. Letting $\eta(\mathbf{z})$ denote the summaries $\eta(\mathbf{z}) = (\eta_0(\mathbf{z}), \eta_1(\mathbf{z}), \eta_2(\mathbf{z}))^\top$ and define their probability limit as $b(\theta) := \text{plim} \eta(\mathbf{z})$. Then, under the DGP in equations (7)-(8), the limit map $\theta \mapsto b(\theta)$ is given by

$$b(\theta) = (1 + \theta_1^2 + \theta_2^2, \theta_1(1 + \theta_2), \theta_2)^\top.$$

Since we are interested in examining the ability of BSL and the robust variants proposed herein to deal with model incompatibility, we consider that, while the researcher believes the data is generated according to an MA(2) model, equation (7), the actual DGP for \mathbf{y} evolves according to the stochastic volatility (SV) model

$$\begin{aligned} y_t &= \exp(h_t/2)u_t \\ h_t &= \omega + \rho h_{t-1} + v_t \sigma_v \end{aligned} \quad (9)$$

where $0 < \rho < 1$, $0 < \sigma_v < 1$, u_t and v_t and both iid standard Gaussian. In this case, if one takes $\eta(\mathbf{y}) := (\eta_0(\mathbf{y}), \eta_1(\mathbf{y}), \eta_2(\mathbf{y}))^\top$ it follows that, under the DGP in (9),

$$\eta(\mathbf{y}) \xrightarrow{P} b_0 := \left(\exp\left(\frac{\omega}{1-\rho} + \frac{1}{2} \frac{\sigma_v^2}{1-\rho^2}\right), 0, 0 \right)^\top. \quad (10)$$

For any value of ω, σ_v and ρ such that

$$\exp\left(\frac{\omega}{1-\rho} + \frac{1}{2} \frac{\sigma_v^2}{1-\rho^2}\right) \neq 1,$$

it follows that $(P_\theta^n \times \Pi, \eta)$ is not compatible. From the definition of $b(\theta)$ and b_0 , it also follows that the value that minimizes $\|b(\theta) - b_0\|_{\Sigma(\theta)}$ is $\theta^* = (0, 0)^\top$, and is the value we would expect the BSL posterior to concentrate onto in the limit.

To understand how BSL and the R-BSL alternatives proposed herein perform under this misspecified model, we consider the following Monte Carlo analysis: we generate $n = 100$ observations from the SV model in (9) and use BSL, R-BSL-M, and R-BSL-V to conduct inference on the unknown parameters θ in the misspecified MA(2) model. Each BSL approach uses $m = 10$ simulated data sets to estimate the mean and the variance.

Given the form of incompatibility in (10), we would expect that the R-BSL-M and R-BSL-V procedures would detect incompatibility in the first summary statistic, the sample variance, while the corresponding adjustment components for the other summaries would not be indistinguishable from the prior. This analysis follows from noting that, under the true DGP in (9), the first two autocorrelations are zero for all values of $(\omega, \rho, \sigma_v)^\top$, and hence the MA(2) parameters should concentrate posterior mass around zero.

For the R-BSL-M and R-BSL-V procedures, we consider the same priors specification as for the simple normal example, and the exact same choice of hyper-parameters: namely, for R-BSL-M we consider that $\gamma_i \sim \text{La}(0, \lambda = 0.5)$, and for the R-BSL-V approach, we take as our prior

$\gamma_i \sim \text{Exp}(\lambda)$ and set λ so that the prior mean is 0.3. For each procedure we consider starting values obtained from the maximum likelihood estimators of the MA(2) model, and the values of θ are obtained via a random-walk Metropolis sampler with fixed covariance matrix given by

$$\begin{pmatrix} 0.1 & 0 \\ 0 & 0.1 \end{pmatrix}.$$

We run the MCMC sampler for 50,000 iterations and discard the first 10,000 for burn-in.

Under this Monte Carlo design, the acceptance rates for R-BSL-V and R-BSL-M are 11% and 6.5%, respectively, while the acceptance rate for the standard BSL approach is just 0.58%. Indeed, the resulting posterior samples from standard BSL are unusable and deviate sharply from those obtained by the R-BSL approaches. The posterior densities for $\theta = (\theta_1, \theta_2)^\top$ across the R-BSL methods are displayed in Figure 4.¹ Both R-BSL-M and R-BSL-V display significant posterior concentration around 0. This is significant as the value that asymptotically minimizes the weighted norm $\|\mu_m(\theta) - \eta(\mathbf{y})\|_{\Sigma_m(\theta)}$, and, thus, which we would expect BSL to concentrate onto, is $\theta^* = (0, 0)^\top$.




Figure 4: Posteriors for R-BSL-M and R-BSL-V for θ_1 (left panel) and θ_2 (right panel).

To understand the poor performance of standard BSL in this example, we can examine the marginal posteriors for the different adjustment terms across R-BSL-V and R-BSL-M, which are given in Figure 5. The results demonstrate that BSL can not reliably match the first summary statistic, the sample variance, while the posteriors that correspond to the first and second-order autocorrelations do not differ from their priors (respectively given as Laplace and exponential) and signify that we can indeed match the second and third summaries. This finding is perfectly

¹To deal with excess autocorrelation in the MCMC chains, the results in Figure 4 have been thinned by taking every 10th sample. This significantly reduced the autocorrelation in the chain across each of the BSL methods and permits smoother density estimates.

in line with the above suggested theoretical behavior: to minimize $\|\mu_m(\theta) - \eta(\mathbf{y})\|$ BSL must place high posterior mass around $\theta = (0, 0)^\top$, but doing so means that we cannot adequately capture the sample variance. While the adjustment procedures can account for this issue, the BSL procedure can not and hence yields low acceptance rates and inaccurate posterior inference.²




Figure 5: Panels A, B and C given the R-BSL-M posteriors for the $\gamma_1, \gamma_2, \gamma_3$, while panels C, D and E give the R-BSL-V posteriors for the same components. The priors for the top three components are all $\text{La}(0, \lambda = 0.5)$, while the priors for the bottom three panels are $\text{Exp}(\lambda)$ where λ is chosen so that the prior mean is 0.3.

4.3 Collective Cell Spreading

4.3.1 Background

Collective cell spreading models are often used to gain insight into the biological mechanisms governing, for example, wound healing and skin cancer growth (e.g. Vo et al. (2015b,a)). Browning et al. (2018) develop a simulation-based model where cells are able to move freely in continuous space. They calibrate the model to real *in vitro* data collected from a cell proliferation assay experiment using a rejection-based ABC algorithm. Here, using our new synthetic likelihood methods, we demonstrate that the model is not compatible with the observed summary statistic and provide insight into what aspects of the data that the model is not able to recover.

The model of Browning et al. (2018) is a stochastic individual based model where cells move and interact in a two dimensional space. Here we provide only brief details of the model and refer to Browning et al. (2018) for the full description. Proliferation (cell birth) and motility (movement) for each cell evolves in continuous time according to a Poisson process. The intrinsic

²We note here that if we instead run BSL with only the first two autocorrelations as summary statistics, the poor behavior for BSL is not in evidence and we can obtain reliable inference on θ .

rates are given by p and m for proliferation and motility events, respectively. The rates of these processes are also neighbourhood-dependent, with rates decreasing as the amount of crowding around a cell increases. The closeness of cells is governed by a Gaussian kernel that depends on a fixed cell diameter, σ . When a cell proliferates, it places a new cell randomly in its neighbourhood according to an uncorrelated two dimensional Gaussian centered at the cell location with component variances of σ^2 . When a motility event occurs, the cell moves a distance of σ . The direction of the move depends on cell density, biased towards lower cell density. A parameter used to help determine the move direction, γ_b , is part of a Gaussian kernel used to measure the closeness of cells. The parameter of interest is $\theta = (p, m, \gamma_b)^\top$.

In the experiments of Browning et al. (2018), images of the cell population are taken every 12 hours starting at 0 hours with the final image taken at 36 hours. Browning et al. (2018) use the number of cells and the pair correlation computed from each image as the summary statistics, resulting in a six dimensional summary statistic here. The pair correlation is the ratio of the number of pairs of agents separated by some pre-specified distance to an expected number of cells separated by the same distance if the cells were uniformly distributed in space.

The prior distribution is set as $p \sim \mathcal{U}(0, 10)$, $m \sim \mathcal{U}(0, 0.2)$ and $\gamma_b \sim \mathcal{U}(0, 20)$ with no dependence amongst parameters, as in Browning et al. (2018). We use MCMC to sample the posterior with 50,000 iterations and no burn-in as we initialise the chain at the point estimate $\theta = (1, 0.04, 6)^\top$ reported in Browning et al. (2018). We sample over the space of a logit-type transformation of θ so that any proposal is within the prior bounds. We use a multivariate Gaussian random walk proposal on the transformed space with a covariance matrix obtained via some pilot MCMC runs. We use $m = 50$ model simulations to estimate the synthetic likelihood at each MCMC iteration. We run our methods on both simulated (using the point estimate of Browning et al. (2018)) and real data.

4.3.2 Results

For the priors for each component of γ , we use a Laplace distribution with a scale of 0.5 for the mean adjustment method and an exponential prior with mean of 0.5 for the variance inflation method.

Firstly we present results for the simulated data (model correctly specified). As shown in Figures 6 and 7, the posterior distributions on θ are similar regardless of whether BSL or R-BSL is applied. The MCMC acceptance rates for BSL and R-BSL-M are both 21% and 20%, respectively. The variance inflation seems to allow for a slightly increased acceptance rate (24%) compared to mean adjustment.

The posterior distribution for each component of γ is shown in Figure 8. It can be seen that most posteriors are similar to the prior. For both R-BSL methods, there is no indication that any of the statistics are incompatible with the model, as expected.

For the real data, the MCMC acceptance rate using BSL is only 3% as the variance of the synthetic likelihood is high generating long periods of no acceptance. Applying R-BSL-M and R-BSL-V results in an MCMC acceptance rate of roughly 12% and 18%, respectively, permitting statistical inference. Again, the variance inflation seems to produce an improved acceptance rate.

The univariate posterior distributions for γ for R-BSL-M and R-BSL-V are shown in Figures 9 and 10, respectively. Both methods identify that the model is not compatible with the pair correlation statistic at 12 and 36 hours.

In Figure 11, we show R-BSL-M posterior predictive distributions of the summary statistics




Figure 6: Contour plots of the posterior distributions based on the simulated data for the collective cell spreading example. Results are shown for BSL (solid) and R-BSL-M (dash). The true parameter values are shown as a crosses.




Figure 7: Contour plots of the posterior distributions based on the simulated data for the collective cell spreading example. Results are shown for BSL (solid) and R-BSL-V (dash). The true parameter values are shown as crosses.

without (left) and with (right) using estimated mean adjustment parameters with the observed summaries overlaid. The corresponding plots for the variance inflation is shown in Figure 12. From both figures, it is evident from the plots on the left that the model is successful in tracking the number of cells over time. However, the model underestimates the rate of decrease in the pair correlation over time. This is valuable information that might enable mathematical biologists to extend the model so that this data feature can be better captured. Both adjustment methods have allowed us to make this inference. It can be seen in the second column of Figure 11 that the mean adjustment is able to shift the predictions so that the observed statistic does not lie so far in the tails. From the second column of Figure 12, the variance adjustment expands the predictions so that the observed statistic does not lie so far in the tails.

Finally, Figure 13 compares the posterior distributions for the mean adjustment and variance inflation, together the standard synthetic likelihood results. It can be seen that the posterior




Figure 8: Posterior distributions for each component of γ (dashed lines) based on the simulated data for the collective cell spreading example for R-BSL-M (left) and R-BSL-V (right). The prior distribution of γ is shown as solid lines for both R-BSL-M (left) and R-BSL-V (right).




Figure 9: Posterior distributions (solid) for each component of γ when applying R-BSL-M to the real data of the collective cell spreading example. The prior distributions, which are Laplace distributed with scale 0.5, are also shown (dash).

distributions are broadly similar, except that the standard synthetic likelihood results suffer from substantial Monte Carlo error due to the small acceptance rate.

4.4 Toad Example

4.4.1 Background

We consider an individual-based model of a species called Fowler’s Toads (*Anaxyrus fowleri*) developed by Marchand et al. (2017), which was also analysed by An et al. (2018). Here we give very brief details, with more information in Marchand et al. (2017) and An et al. (2018).

The model assumes that a toad hides in its refuge site in the daytime and moves to a randomly




Figure 10: Posterior distributions (solid) for each component of γ when applying R-BSL-V to the real data of the collective cell spreading example. The prior distributions, which are exponential with mean 0.5, are also shown (dash).




Figure 11: Posterior predictive distributions (summarised as boxplots) of the summary statistics obtained with R-BSL-M. Shown are the posterior predictive distributions of the summaries without (left) and with (right) including the variance inflation in the predictions. The observed summary statistics are overlaid as crosses.

chosen foraging place at night. GPS location data are collected on n_t toads for n_d days, i.e. the observation matrix \mathbf{Y} is of dimension $n_d \times n_t$ ($n_t = 66$ and $n_d = 63$ here). Then \mathbf{Y} is summarised




Figure 12: Posterior predictive distributions (summarised as boxplots) of the summary statistics obtained with R-BSL-V. Shown are the posterior predictive distributions of the summaries without (left) and with (right) including the variance inflation in the predictions. The observed summary statistics are overlaid as crosses.




Figure 13: Posterior distributions for each component of θ when applying BSL (solid), R-BSL-M (dash) and R-BSL-V (dot-dash) to the real data of the collective cell spreading example. The priors used are Laplace with scale 0.5 for mean adjustment and exponential with a mean of 0.5 for variance inflation.

to 4 sets comprising the relative moving distances for time lags of 1, 2, 4, 8 days. For instance, \mathbf{y}_1 consists of the displacement information of lag 1 day, $\mathbf{y}_1 = \{|\mathbf{Y}_{i,j} - \mathbf{Y}_{i+1,j}|; 1 \leq i \leq n_d - 1, 1 \leq j \leq n_t\}$.

Simulating from the model involves two distinct processes. For each toad, we first generate an overnight displacement, Δy , then mimic the returning behaviour with a simplified model.

The overnight displacement is assumed to belong to the Lévy-alpha stable distribution family, with stability parameter α and scale parameter δ . The total returning probability is a constant p_0 , if a return occurs on day i , $1 \leq i \leq m$, then the return site is the same as the refuge site on day i , where i is selected randomly from $1, 2, \dots, m$ with equal probability. Here we consider both simulated and real datasets. For the synthetically generated data we take $\theta = (\alpha, \delta, p_0)^\top = (1.8, 45, 0.6)^\top$, which is informed by the parameter estimates obtained in Marchand et al. (2017). We use a uniform prior over $(1, 2) \times (0, 100) \times (0, 0.9)$. Marchand et al. (2017) consider three variations on the model. Here, we consider their ‘Model 2’ since there is a strong indication from their results that this model is not able to recover some of the chosen summary statistics.

As in Marchand et al. (2017), the dataset of displacements is split into two components. If the absolute value of the displacement is less than 10 metres, it is assumed the toad has returned to its starting location. For the summary statistic, we consider the number of toads that returned (Marchand et al. (2017)). For the non-returns (absolute displacement greater than 10 metres) we consider a larger collection of summaries. We calculate the log difference between adjacent p -quantiles with $p = 0, 0.1, \dots, 1$ and also the median. These statistics are computed separately for the four time lags. This results in 48 statistics in total, which is hard to handle for conventional ABC methods. An et al. (2018) demonstrate that BSL is computationally efficient enough to analyse simulated data for this application with a similar number of summary statistics.

For the γ parameters of the mean adjustment and variation inflation procedures we use the same priors as the previous example.

4.4.2 Results

We first consider the simulated dataset, where we use $n = 300$ simulations to estimate the synthetic likelihood at each MCMC iteration. Standard BSL, together with the two incompatibility extensions, produce approximate posteriors shown in Figure 14. As can be seen, the adjustments produce posteriors remarkably similar to BSL with slightly inflated variances. The MCMC acceptance rates for BSL, R-BSL-M and R-BSL-V are 11%, 9% and 22%, respectively. As consistent with previous results, the variance adjustment improves the computational efficiency, even when the model is correctly specified. The posterior distributions for γ of R-BSL are shown in Figure 15. In all cases, the posteriors are not too dissimilar to the prior.

For the real data, we required $n = 2000$ simulations for estimating the synthetic likelihood to obtain an acceptance rate of 9% for BSL. However, the chain still suffered from periods of stickiness. In contrast, with only $n = 500$, the R-BSL-M and R-BSL-V produce acceptance rates of 7% and 15%, respectively, without substantial stickiness. The variance inflation method offers a computational improvement of about one order of magnitude over BSL when accounting for both acceptance rate and number of simulations.

The posterior distributions for the components of γ for the R-BSL methods are shown in Figure 16. It is evident from the plots that our methods have identified that there are three or four statistics that the model is not compatible with. The statistic with the largest incompatibility is the number of returns for lag 1. For R-BSL-V, the 95% posterior predictive interval for this statistic is (262, 346) with an observed value of 234. Other statistics showing some incompatibility are the first quantile differences of the non-returns for lags 3 and 4. Figure 17 confirms that the observed data are not consistent with the posterior predictive distribution of the (log) non-return distances for lags 3 and 4, in that the model generally predicts larger non-return distances. The mean adjustment results are similar (not shown). Our adjustment methods




Figure 14: Univariate posterior distributions for the parameters when applying BSL (solid), R-BSL-V (dash) and R-BSL-M (dot-dash) to simulated data for the toad example. True parameter values are shown as crosses.




Figure 15: Posterior distributions for γ for R-BSL-M (right) and R-BSL-V (left) applied to the simulated data for the toad example. The thick line is the prior and the thin lines are the posteriors for the components of γ .

permit in-depth analyses such as these and may provide practitioners valuable information for improving the model.

The bivariate posterior distributions for the parameters based on the adjustment methods are shown in Figure 18. It is evident that the estimated posterior distributions are similar, with the R-BSL-M posteriors slightly more concentrated than R-BSL-V. Univariate posteriors for all BSL approaches are shown in Figure 19. It appears that the adjustments result in a significant shift of the posterior and variance inflation. The standard BSL results are adversely affected by a few long periods of no MCMC acceptance. This indicates that statistical inferences are unlikely to be robust to the model misspecification.




Figure 16: Posterior distributions for γ for R-BSL-M (right) and R-BSL-V (left) applied to the real data for the toad example. The thick line is the prior and the thin lines are the posteriors for the components of γ .




Figure 17: Posterior predictive distributions of the log non-returns for the four lags based on R-BSL-V. The thick line is the distribution for the observed data.




Figure 18: Bivariate posterior distributions visualised as contour plots for the parameters based on R-BSL-M (right) and R-BSL-V (left) applied to the real data for the toad example.




Figure 19: Univariate posterior distributions for the parameters when applying BSL (solid), R-BSL-V (dash) and R-BSL-M (dot-dash) to real data for the toad example.

5 Discussion

This paper has made two significant contributions to the literature on approximate Bayesian methods. Firstly, to our knowledge, this is the first piece of research to demonstrate that, similar to the method of approximate Bayesian computation (ABC), the method of Bayesian synthetic likelihood (BSL) can deliver unreliable inference when the assumed model is misspecified. Secondly, to circumvent the poor behavior of BSL in these settings, we have proposed a modification of BSL that is robust to model misspecification. Several Monte Carlo and empirical examples are used to illustrate the performance of this new method, with the results demonstrating the overwhelming performance gains of this robust approach relative to standard BSL when the model is misspecified.

In addition to being robust to model misspecification, this new approach also allows the user to detect precisely which summary statistics are incompatible with the assumed data generating process. Incorporating this information within subsequent rounds of model building could lead to better models that can more accurately capture the behavior exhibited by the observed summary statistics. In this sense, the robust BSL approach can be viewed as a BSL based version of the model criticism approach of Ratmann et al. (2009). In the context of ABC, Ratmann et al. (2009) propose an approach to detect aspects of the model that the summary statistics can not adequately capture. Their approach relies on treating the ABC tolerance as an unknown parameter, and augmenting the original ABC inference problem with this additional parameter. The authors argue that posterior realizations for the tolerance parameter that are “large” indicate the possibility of a mismatch between the model and the observed data.

While useful, in the case of multivariate summaries the approach of Ratmann et al. (2009) requires a tolerance parameter for each summary statistic used in the analysis, with posterior inference then required on the full set of model parameters and tolerance parameters. Therefore, even for a moderate number of summaries, this approach can exacerbate the underlying curse-of-dimensionality in ABC, as it pertains to overall number of parameters in the analysis (Frazier et al., 2018). Moreover, since summaries differ in their responsiveness to the values used to simulate the data, the scales of the individual tolerance posteriors can vary greatly and, therefore, it is not entirely clear how to accurately determine when model misspecification is in evidence. In contrast, the approach considered herein has as a direct benchmark with which to gauge the impact of misspecification on the summaries, namely the prior distribution of the adjustment components. If the corresponding posterior for the adjustment component in the robust version of BSL does not resemble the prior, this is strong evidence that this summary can not be matched by the assumed model. If one wished to put a numerical value, or conduct a formal hypothesis test, on the difference between the prior and posterior, any number of techniques could be used.

The examples illustrated a strong indication that, in particular, the variance inflation approach can significantly improve the MCMC acceptance rate, especially under model misspecification. The variance inflation approach bears some resemblance to MCMC ABC approaches that assign a distribution to the ABC tolerance to facilitate MCMC mixing by proposing a relatively large tolerance value (e.g. Bortot et al. (2007)). However, improving mixing is not our primary focus, but is simply a useful by-product. Our R-BSL approaches may also be useful for initial explorations of the parameter space when it is not known where the bulk of the posterior support is since, even for a correctly specified model, a poor parameter value will not be able to recover the observed statistic.

Lastly, we note that, given the similarities between BSL and ABC, a natural question posed during this research was whether or not the mean and variance adjustment approaches discussed in this current paper were applicable in the context of ABC. In concurrent work from the authors, preliminary investigations into a similar type of mean and variance adjusted ABC have revealed that such an approach can mitigate the poor performance of ABC under model misspecification (Frazier et al., 2017).

Acknowledgments

CD was supported by an Australian Research Council’s Discovery Early Career Researcher Award funding scheme (DE160100741). The authors are grateful to Ziwen An, who provided some useful code for this paper.

References

- An, Z., Nott, D. J., and Drovandi, C. (2018). Robust Bayesian synthetic likelihood via a semi-parametric approach. *arXiv preprint arXiv:1809.05800*.
- Andrieu, C. and Roberts, G. O. (2009). The pseudo-marginal approach for efficient Monte Carlo computations. *The Annals of Statistics*, 37(2):697–725.
- Bhattacharya, A., Pati, D., Pillai, N. S., and Dunson, D. B. (2012). Bayesian shrinkage. *arXiv preprint arXiv:1212.6088*.
- Blum, M. G. (2010). Approximate Bayesian computation: a nonparametric perspective. *Journal of the American Statistical Association*, 105(491):1178–1187.
- Bortot, P., Coles, S. G., and Sisson, S. A. (2007). Inference for stereological extremes. *Journal of the American Statistical Association*, 102(477):84–92.
- Box, G. E. (1976). Science and statistics. *Journal of the American Statistical Association*, 71(356):791–799.
- Browning, A. P., McCue, S. W., Binny, R. N., Plank, M. J., Shah, E. T., and Simpson, M. J. (2018). Inferring parameters for a lattice-free model of cell migration and proliferation using experimental data. *Journal of Theoretical Biology*, 437:251–260.
- Frazier, D. T., Martin, G. M., Robert, C. P., and Rousseau, J. (2018). Asymptotic properties of approximate bayesian computation. *Biometrika*, 105(3):593–607.
- Frazier, D. T., Robert, C. P., and Rousseau, J. (2017). Model misspecification in ABC: Consequences and diagnostics. *arXiv preprint arXiv:1708.01974*.
- Kleijn, B. and Van der Vaart, A. (2012). The Bernstein-von-Mises theorem under misspecification. *Electronic Journal of Statistics*, 6:354–381.
- Marchand, P., Boenke, M., and Green, D. M. (2017). A stochastic movement model reproduces patterns of site fidelity and long-distance dispersal in a population of Fowler’s toads (*Anaxyrus fowleri*). *Ecological Modelling*, 360:63 – 69.

- Marin, J.-M., Pillai, N. S., Robert, C. P., and Rousseau, J. (2014). Relevant statistics for Bayesian model choice. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(5):833–859.
- Marin, J.-M., Pudlo, P., Robert, C. P., and Ryder, R. J. (2012). Approximate Bayesian computational methods. *Statistics and Computing*, 22(6):1167–1180.
- Neal, R. M. (2003). Slice sampling. *The Annals of Statistics*, 31(3):705–767.
- Nott, D. J., Drovandi, C., and Kohn, R. (2019). Bayesian inference using synthetic likelihood: asymptotics and adjustments. *arXiv preprint arXiv:1902.04827*.
- Park, T. and Casella, G. (2008). The Bayesian lasso. *Journal of the American Statistical Association*, 103(482):681–686.
- Price, L. F., Drovandi, C. C., Lee, A., and Nott, D. J. (2018). Bayesian synthetic likelihood. *Journal of Computational and Graphical Statistics*, 27(1):1–11.
- Ratmann, O., Andrieu, C., Wiuf, C., and Richardson, S. (2009). Model criticism based on likelihood-free inference, with an application to protein network evolution. *Proceedings of the National Academy of Sciences*, 106(26):10576–10581.
- Sisson, S. A., Fan, Y., and Beaumont, M. (2018). *Handbook of Approximate Bayesian Computation*. Chapman and Hall/CRC.
- Vo, B. N., Drovandi, C. C., Pettitt, A. N., and Pettet, G. J. (2015a). Melanoma cell colony expansion parameters revealed by approximate Bayesian computation. *PLOS Computational Biology*, 11(12):e1004635.
- Vo, B. N., Drovandi, C. C., Pettitt, A. N., and Simpson, M. J. (2015b). Quantifying uncertainty in parameter estimates for stochastic models of collective cell spreading using approximate Bayesian computation. *Mathematical Biosciences*, 263:133–142.
- Wood, S. N. (2010). Statistical inference for noisy nonlinear ecological dynamic systems. *Nature*, 466(7310):1102.

Appendix A. Proof of Proposition 1

We first prove a result that is of independent interest. Namely, we demonstrate that under Assumptions 1-6, the standard BSL posterior concentrates all posterior mass onto the sets of the form $\{b : \|b - b_0\| \lesssim v_n^{-1}\}$. To simplify the computations, we demonstrate this result for the so-called “idealized” BSL posterior, which takes as the mean and variance the infeasible counterparts

$$b(\theta) := \mathbb{E}[\eta(\mathbf{y})|\theta] \text{ and } A_n(\theta) := \left(\mathbb{E} [(\eta(\mathbf{y}) - \mathbb{E}[\eta(\mathbf{y})|\theta])^2 | \theta] \right)^{1/2}.$$

For $\Sigma(\theta) := A_n(\theta)^\top A_n(\theta)$, the BSL “likelihood” is then given by

$$g_n(\eta|\theta) := (2\pi)^{-\frac{dn}{2}} \det(\Sigma(\theta))^{-1} \exp\left(-\frac{1}{2} \{\eta - b(\theta)\}^\top \Sigma^{-1}(\theta) \{\eta - b(\theta)\}\right).$$

Even though b and Σ can depend on n , we suppress this dependence for notational simplicity.

The following result is a modification of Corollary 1 in Marin et al. (2014), and the proof follows similarly.

Lemma 1. *Under Assumption 1-6, the idealized BSL posterior $\pi\{\cdot|\eta(\mathbf{y})\}$ concentrates at the rate $1/v_n$ onto b_0 , provided that $\alpha > \tau$.*

Proof. Take M_n to be a sequence diverging to ∞ . Define

$$m(\eta) := \int_{\Theta} g_n(\eta|\theta) \pi(\theta) d\theta,$$

and consider the set

$$T_n(M_n) := \{\theta \in \Theta : \|b(\theta) - b_0\| > M_n v_n^{-1}\}.$$

Now, consider the BSL posterior over the set $T_n(M_n)$:

$$\frac{\int_{T_n(M_n)} g_n[\eta|\theta] \pi(\theta) d\theta}{m(\eta)} \equiv \frac{\int_{T_n(M_n)} \frac{g_n[\eta|\theta]}{g_n^0(\eta)} \pi(\theta) d\theta}{m(\eta)/g_n^0(\eta)}.$$

Define

$$N_n := \int_{T_n(M_n)} \frac{g_n[\eta|\theta]}{g_n^0(\eta)} \pi(\theta) d\theta$$

and

$$D_n := m(\eta)/g_n^0(\eta).$$

By Lemma 2, we have that $D_n \gtrsim v_n^{-\tau}$. Moreover, by Lemma 3, we have that $N_n \lesssim M_n^{-\alpha} v_n^{-\alpha}$. Therefore,

$$\Pi \{T_n(M_n) | \eta\} = \frac{N_n}{D_n} = o_{P_0^n} (v_n^{-\alpha}/v_n^{-\tau}) = o_{P_0^n}(1).$$

From the above, conclude that

$$\Pi [\|b(\theta) - b_0\| > M_n v_n^{-1} | \eta] = o_P(1).$$

Applying Assumption 5, we have the stated result:

$$\Pi [\|\theta - \theta_0\| > L\{M_n v_n^{-1}\}^\kappa | \eta] = o_P(1).$$

■

Lemma 2. *Under Assumptions 1-6,*

$$\lim_{n \rightarrow \infty} P_n^0 (m(\eta) / g_n^0(\eta) \gtrsim v_n^{-\tau}) = 1.$$

Proof. Fix $\delta > 0$. By Assumption 1, there exists an M_δ such that

$$P_n^0 \{v_n \|\eta - b_0\| > M_\delta\} < \delta.$$

For all $\epsilon > 0$, by Assumption 6, there exists $U_\epsilon, \delta_\epsilon$ such that, for $\eta \in V_n$,

$$\int_{\Theta} g_n(\eta | \theta) \pi(\theta) d\theta \geq \delta_\epsilon g_n^0(\eta) \pi[\mathcal{F}_n(U_\epsilon)], \text{ for } \mathcal{F}_n(u) := \{\theta \in \Theta : \|b(\theta) - b_0\| \leq uv_n^{-1}\}.$$

Applying Assumption 4, there exists some $c(\epsilon)$ such that

$$\int_{\Theta} g_n(\eta | \theta) \pi(\theta) d\theta \geq \delta_\epsilon g_n^0(\eta) v_n^{-\tau}.$$

From the definition of V_n , it follows that, for n large enough,

$$P_n^0 \left\{ \int_{\Theta} g_n(\eta | \theta) \pi(\theta) d\theta \geq c(\epsilon) g_n^0(\eta) v_n^{-\tau} \right\} \geq 1 - \epsilon.$$

■

Lemma 3. *Under Assumptions 1-6, for the set $T_n(M_n) := \{\theta \in \Theta : \|b(\theta) - b_0\| > M_n v_n^{-1}\}$,*

$$\lim_{n \rightarrow \infty} P_n^0 \left(\int_{T_n(M_n)} \frac{g_n[\eta|\theta]}{g_n^0(\eta)} \pi(\theta) d\theta \lesssim M_n^{-\alpha} v_n^{-\alpha} \right) = 1.$$

Proof. Fix $\delta > 0$. By Assumption 1, there exists some M_δ such that, for some n large enough,

$$P_n^0 \{\|\eta - b_0\| > M_\delta / v_n\} < \delta.$$

Consider the joint probability

$$P_n^0 \left\{ \int_{T_n(M_\delta)} g_n(\eta|\theta) \pi(\theta) d\theta > g_n^0(\eta) v_n^{-\tau} \right\}$$

Applying Markov inequality and Fubini's theorem, and Assumption 4,

$$\begin{aligned} & P_n^0 \left\{ \int_{T_n(M_\delta)} g_n(\eta|\theta) \pi(\theta) d\theta > g_n^0(\eta) v_n^{-\tau} \right\} \\ & \leq P_n^0 (|\eta - b_0| > M_\delta v_n^{-1}) + v_n^\tau \int_{T_n(M_\delta)} \int_{\|\mathbf{s}_n - b(\theta)\| \geq M_\delta} \frac{1}{g_n^0(t)} g_n^0(t) g_n(t|\theta) \pi(\theta) dt d\theta \\ & \leq P_n^0 (|\eta - b_0| > M_\delta v_n^{-1}) + v_n^\tau \int_{T_n(M_\delta)} \int_{\|\eta - b(\theta)\| \geq M_\delta} g_n(t|\theta) \pi(\theta) dt d\theta \\ & \leq P_n^0 (|\eta - b_0| > M_\delta v_n^{-1}) + v_n^\tau \int_{T_n(M_\delta)} G_n[\|\eta - b(\theta)\| \geq M_\delta |\theta|] \pi(\theta) d\theta. \end{aligned}$$

It then follows that, by Assumption 3,

$$\begin{aligned} P_n^0 (|\eta - b_0| > M_\delta v_n^{-1}) + v_n^\tau \int_{T_n(M_\delta)} G_n [\|\eta - b(\theta)\| \geq M_\delta |\theta|] \pi(\theta) d\theta \\ \leq v_n^\tau (M_\delta v_n)^{-\alpha} \int_{\Theta} c(\theta) \pi(\theta) d\theta \lesssim o(v_n^{\tau-\alpha}), \end{aligned}$$

where, by Assumption 3, $\int_{\Theta} c(\theta) \pi(\theta) d\theta = O(1)$. Conclude that, for any $\delta > 0$,

$$P_n^0 \left\{ \int_{T_n(M_\delta)} g_n(\eta|\theta) \pi(\theta) d\theta > g_n^0(\eta) v_n^{-\tau} \right\} \leq \delta + o(v_n^{\tau-\alpha}) \leq 2\delta$$

for n large enough. ■

Proof of Proposition 1. We prove the result for R-BSL-M and R-BSL-V separately.

R-BSL-M: From the posterior concentration of the BSL posterior we have that, for some $M_n \rightarrow \infty$, with $M_n/v_n \rightarrow 0$,

$$\begin{aligned} \Pi [\|\Gamma\| > t|\eta(\mathbf{y})] \\ = \frac{\int_{\|\Gamma\|>t} \int_{\|b-b_0\|\leq M_n/v_n} (2\pi)^{-\frac{d_\eta}{2}} \det[\Sigma(b)]^{-1/2} e^{(-\frac{1}{2}[b-\eta-x(\Gamma)]^\top [\Sigma(b)]^{-1}[b-\eta-x(\Gamma)])} \pi(b) \pi(\Gamma) db d\Gamma}{\int \int_{\|b-b_0\|\leq M_n/v_n} (2\pi)^{-\frac{d_\eta}{2}} \det[\Sigma(b)]^{-1/2} e^{(-\frac{1}{2}[b-\eta-x]^\top [\Sigma(b)]^{-1}[b-\eta-x])} \pi(b) \pi(\Gamma) db d\Gamma} + o_P(1), \end{aligned}$$

where $x(\Gamma) := \text{diag}[\Sigma(b)]^{1/2} \Gamma$ and $\eta := \eta(\mathbf{y})$.

Define $Z := v_n(b - \eta)$ and consider the change of variables $b \mapsto v_n(b - \eta) + v_n(\eta - b_0) \equiv Z + v_n(\eta - b_0)$, which yields

$$\Pi [\|\Gamma\| > t|\eta(\mathbf{y})] = \frac{N_n}{D_n} + o_P(1),$$

where

$$N_n := \int_{\|\Gamma\|>t} \int_{\|Z\|\leq M_n} \frac{e^{(-\frac{1}{2}[Z/v_n-x(\Gamma)]^\top [\Sigma(Z/v_n+\eta)]^{-1}[Z/v_n-x(\Gamma)])}}{(2\pi)^{d/2} \det[\Sigma(Z/v_n+\eta)]^{1/2}} \pi(Z/v_n+\eta) \pi(\Gamma) dZ d\Gamma,$$

and

$$D_n := \int \int_{\|Z\|\leq M_n} \frac{e^{(-\frac{1}{2}[Z/v_n-x(\Gamma)]^\top [\Sigma(Z/v_n+\eta)]^{-1}[Z/v_n-x(\Gamma)])}}{(2\pi)^{\frac{d_\eta}{2}} \det[\Sigma(Z/v_n+\eta)]^{1/2}} \pi(Z/v_n+\eta) \pi(\Gamma) dZ d\Gamma.$$

We now analyze N_n and D_n separately.

Term D_n : By Assumption 1, $P_n^0 \{\|\eta - b_0\| \leq M_\delta/v_n\} \geq 1 - \delta$ for some $M_\delta, \delta > 0$. On this set, from the definition of M_n ,

$$\begin{aligned} \sup_{\|Z\|\leq M_n} \|\pi(Z/v_n + \eta) - \pi(b_0)\| &= o_P(1) \\ \sup_{\|Z\|\leq M_n} \|\Sigma(Z/v_n + \eta) - \Sigma(b_0)\| &= o_P(1). \end{aligned}$$

The first equation follows from continuity of $\pi(\cdot)$ and the second from continuity of $A_n(\theta) \equiv \Sigma^{1/2}(\theta)$. By the dominated convergence theorem,

$$\frac{D_n}{\pi(b_0)\det[\Sigma(b_0)]^{-1/2}} = \int \int_{\|Z\| \leq M_n} (2\pi)^{-\frac{d_n}{2}} e^{-\frac{1}{2}[Z/v_n - x(\Gamma)]^\top [\Sigma(b_0)]^{-1} [Z/v_n - x(\Gamma)]} \pi(\Gamma) dZ d\Gamma + o_P(1).$$

Define $\tilde{Z} := [\Sigma^{1/2}(b_0)v_n]^{-1} Z$ and note that, by Assumption 2, $\tilde{Z} := A(b_0)^{-1} Z + o_P(1)$, for some positive definite matrix $A(b_0)$. Defining $\tilde{\Gamma} := A^{-1}(b_0)\Gamma$, we have that

$$\frac{D_n}{\pi(b_0)\det[\Sigma(b_0)]^{-1/2}} = \int (2\pi)^{-\frac{d_n}{2}} \int_{\|\tilde{Z}\| \leq M_n} e^{-\frac{1}{2}\tilde{Z}^\top \tilde{Z} + \tilde{Z}^\top \tilde{\Gamma}} e^{-\frac{1}{2}\tilde{\Gamma}^\top \tilde{\Gamma}} \pi(\Gamma) dZ d\Gamma + o_P(1).$$

Recall the following: for $x, y \in \mathbb{R}^{d_\eta}$,

$$\int_{\mathbb{R}^{d_\eta}} e^{-\frac{1}{2}x^\top x + x^\top y} dx = (2\pi)^{d_\eta/2} e^{y^\top y/2}.$$

From the above fact and the dominated convergence theorem

$$\int_{\|\tilde{Z}\| \leq M_n} e^{-\frac{1}{2}\tilde{Z}^\top \tilde{Z} + \tilde{Z}^\top \tilde{\Gamma}} dZ \rightarrow (2\pi)^{d_\eta/2} e^{\frac{1}{2}\tilde{\Gamma}^\top \tilde{\Gamma}}.$$

Apply the above and Fubini's Theorem to deduce

$$\frac{D_n}{\pi(b_0)\det[\Sigma(b_0)]^{-1/2}} \rightarrow_p \int \pi(\Gamma) d\Gamma = 1.$$

Term N_n : Apply the same argument as for D_n , to obtain

$$\frac{N_n}{\pi(b_0)\det[\Sigma(b_0)]^{-1/2}} = \int_{\|\Gamma\| > t} (2\pi)^{-\frac{d_\eta}{2}} \int_{\|\tilde{Z}\| \leq M_n} e^{-\frac{1}{2}\tilde{Z}^\top \tilde{Z} + \tilde{Z}^\top \tilde{\Gamma}} e^{-\frac{1}{2}\Gamma^\top \Gamma} \pi(\Gamma) dZ d\Gamma \rightarrow_p \int_{\|\Gamma\| > t} \pi(\Gamma) d\Gamma.$$

Conclude that

$$\Pi[\|\Gamma\| > t | \eta(\mathbf{y})] = \frac{N_n}{D_n} + o_P(1) = \Pi[\|\Gamma\| > t] + o_P(1).$$

R-BSL-V: Define

$$V(b, \Gamma) := \Sigma(b) + \text{diag}[\Sigma^{1/2}(b)] \Gamma \Gamma^\top \text{diag}[\Sigma^{1/2}(b)]^\top$$

From posterior concentration, for some $M_n \rightarrow \infty$, with $M_n/v_n \rightarrow 0$,

$$\begin{aligned} & \Pi[\|\Gamma\| > t | \eta] \\ &= \frac{\int_{\|\Gamma\| > t} \int_{\|b-b_0\| \leq M_n/v_n} (2\pi)^{-\frac{d_\eta}{2}} \det[V(b, \Gamma)]^{-1} e^{(-\frac{1}{2}[b-\eta]^\top V(b, \Gamma)^{-1} [b-\eta])} \pi(b) \pi(\Gamma) db d\Gamma}{\int \int_{\|b-b_0\| \leq M_n/v_n} (2\pi)^{-\frac{d_\eta}{2}} \det[V(b, \Gamma)]^{-1} e^{(-\frac{1}{2}[b-\eta]^\top V(b, \Gamma)^{-1} [b-\eta])} \pi(b) \pi(\Gamma) db d\Gamma} + o_P(1), \end{aligned}$$

where $\eta := \eta(\mathbf{y})$.

Similar to the case of R-BSL-M, define $Z := v_n(b - \eta)$ and consider the change of variables $b \mapsto v_n(b - \eta) + v_n(\eta - b_0) \equiv Z + v_n(\eta - b_0)$, which yields, for some $M_n \rightarrow \infty$,

$$\Pi[\|\Gamma\| > t | \eta(\mathbf{y})] = \frac{N_n}{D_n} + o_P(1)$$

where

$$N_n := \int_{\|\Gamma\| > t} \int_{\|Z\| \leq M_n} \frac{e^{(-\frac{1}{2}[Z/v_n]^\top V^{-1}(Z/v_n + \eta, \Gamma)[Z/v_n])}}{(2\pi)^{\frac{d_n}{2}} \det[V(Z/v_n + \eta, \Gamma)]^{1/2}} \pi(Z/v_n + \eta) \pi(\Gamma) dZ d\Gamma$$

and

$$D_n := \int \int_{\|Z\| \leq M_n} \frac{e^{(-\frac{1}{2}[Z/v_n]^\top V^{-1}(Z/v_n + \eta, \Gamma)[Z/v_n])}}{(2\pi)^{\frac{d_n}{2}} \det[V(Z/v_n + \eta, \Gamma)]^{1/2}} \pi(Z/v_n + \eta) \pi(\Gamma) dZ d\Gamma.$$

Again, we analyze N_n and D_n separately.

Term D_n : Similar to the previous result, by Assumption 1, $P_n^0\{\|\eta - b_0\| \leq M_\delta/v_n\} \geq 1 - \delta$ for some $M_\delta, \delta > 0$. Similarly to the case of R-BSL-M,

$$\begin{aligned} \sup_{\|Z\| \leq M_n} \|\pi(Z/v_n + \eta) - \pi(b_0)\| &= o_P(1) \\ \sup_{\|Z\| \leq M_n} \|\Sigma(Z/v_n + \eta) - \Sigma(b_0)\| &= o_P(1). \end{aligned}$$

By Assumption 2, and the dominated convergence theorem,

$$\frac{D_n}{\pi(b_0)v_n^{-d_n}} = \int \int_{\|Z\| \leq M_n} (2\pi)^{-\frac{d_n}{2}} \det[v_n^2 V(b_0, \Gamma)]^{-1/2} e^{(-\frac{1}{2}[Z/v_n]^\top [V(b_0, \Gamma)]^{-1}[Z/v_n])} \pi(\Gamma) dZ d\Gamma + o_P(1).$$

Define

$$\tilde{Z} := \left[v_n^2 \Sigma(b_0) + \text{diag}[v_n \Sigma^{1/2}(b_0)] \Gamma \Gamma^\top \text{diag}[v_n \Sigma^{1/2}(b_0)]^\top \right]^{-1/2} Z.$$

By Assumption 2, $\tilde{Z} = A(b_0 + \Gamma)^{-1/2} Z + o_P(1)$ and we then obtain

$$\frac{D_n}{\pi(b_0)v_n^{-d_n}} = \int (2\pi)^{-\frac{d_n}{2}} \int_{\|\tilde{Z}\| \leq M_n} \det[A(b_0 + \Gamma)]^{-1/2} e^{(-\frac{1}{2}\tilde{Z}^\top \tilde{Z})} \pi(\Gamma) dZ d\Gamma + o_P(1).$$

From the dominated convergence theorem,

$$(2\pi)^{-d_n/2} \int_{\|\tilde{Z}\| \leq M_n} \det[A(b_0 + \Gamma)]^{-1/2} e^{(-\frac{1}{2}\tilde{Z}^\top \tilde{Z})} dZ \rightarrow_p 1.$$

Applying the above and Fubini's Theorem delivers

$$\frac{D_n}{\pi(b_0)v_n^{-d_n}} \rightarrow_p \int \pi(\Gamma) d\Gamma = 1.$$

Term N_n : Apply the same argument as for D_n , to deduce that

$$\frac{N_n}{\pi(b_0)v_n^{-d_n}} = \int_{\|\Gamma\| > t} (2\pi)^{-\frac{d_n}{2}} \int_{\|\tilde{Z}\| \leq M_n} \det[A(b_0 + \Gamma)]^{-1/2} e^{(-\frac{1}{2}\tilde{Z}^\top \tilde{Z})} \pi(\Gamma) dZ d\Gamma + o_P(1) \rightarrow_p \int_{\|\Gamma\| > t} \pi(\Gamma) d\Gamma.$$

Conclude that

$$\Pi[\|\Gamma\| > t | \eta(\mathbf{y})] = \frac{N_n}{D_n} + o_P(1) = \Pi[\|\Gamma\| > t] + o_P(1).$$

■