

Fast Spread in Controlled Evolutionary Dynamics

Lorenzo Zino, Giacomo Como, *Member, IEEE*, and Fabio Fagnani, *Member, IEEE*

Abstract—We study the spread of a novel state in a network, in the presence of an exogenous control. The considered controlled evolutionary dynamics is a non-homogeneous Markov process that describes the evolution of the states of all nodes in the network. Through a rigorous analysis, we estimate the performance of the system by establishing upper and lower bounds on the expected time needed for the novel state to replace the original one. Such bounds are expressed in terms of the support and intensity of the control policy (specifically, the set of nodes that can be controlled and its energy) and of the network topology and establish fundamental limitations on the system’s performance. Leveraging these results, we are able to classify network structures depending on the possibility to control the system using simple open-loop control policies. Finally, we propose a feedback control policy that, using little knowledge of the network topology and of the system’s evolution at a macroscopic level, allows for a substantial speed up of the spreading process with respect to simple open-loop control policies. All these theoretical results are presented together with explanatory examples, for which Monte Carlo simulations corroborate our analytical findings.

Index Terms—Control of network systems; Feedback control; Spreading processes; Evolutionary dynamics; Diffusion of innovation.

I. INTRODUCTION

IN the last decades, the study of spreading dynamics in network systems has significantly advanced. On the one hand, increasingly refined models have been proposed, with applications spanning from the spread of epidemic diseases and mutations to the diffusion of innovation and the adoption of social norms. On the other hand, the insight into the behavior of such systems has considerably deepened, allowing for the development of accurate strategies to effectively control and influence the spreading process. The literature on epidemics offers a paradigmatic example, where the understanding of how the network of interactions between the individuals influences the spread of a disease [2]–[6] has paved the way for the study of control policies to mitigate or stop epidemic outbreaks [7]–[11]. Also, for decision-making in social systems, the extensive analysis of the voter model [12], [13] and the study of dynamical interaction models in game

theoretic frameworks [14]–[16] have allowed for the design of techniques to optimally place a spreader in a network [17], [18] and to control networks of imitative agents [19].

In this paper, we focus on a dynamical network system that models the spread of novel states inside a population. It first appeared in the literature in [20] under the name of *evolutionary dynamics*. These dynamical systems have been used to model the evolutionary competition between a novel species and the original one in a geographic region. In evolutionary dynamics, the geographical pattern is modeled as a weighted network whose nodes represent regions, links represent proximity, and the links’ weights measure the level of interaction between individuals in adjacent regions. Each node can be in two possible states, representing the original and the novel species, respectively. The spreading process follows a probabilistic rule and takes place through pairwise interactions between adjacent nodes, which yield a competition between the species present in the two nodes to place their offsprings. This model naturally accounts for possible stochastic biases that favor one specie against the other in the pairwise competitions, thus reflecting an evolutionary advantage of one specie with respect to the other one. The key problems considered in the literature of evolutionary dynamics [20]–[24] focus on the understanding of how the network structure influences the probability that the mutant state (typically with an evolutionary advantage) diffuses in the network—that is often referred to as the fixation probability—and the duration of such a spreading process.

This model may also find valuable applications in other fields, e.g., to study the diffusion of innovation in social systems. In this setting, the two states may represent two alternative technologies, and the nodes represent users interacting in a pairwise fashion on a social network and exchanging information on the used products. As a result of these pairwise interactions, one of the two individuals may get convinced to adopt the technology used by the other one, thus leading to an imitative/contagion mechanism. As before, the model can easily incorporate an intrinsic bias towards one technology with respect to the other, thus accounting, e.g., for an intrinsic quality difference between the two products.

Despite its importance, few analytical results for evolutionary dynamics are available in the literature. To the best of our knowledge, the fixation probability has been analytically computed for very specific network topologies [21], [22], while most of the results are based on extensive Monte Carlo simulations [20], [23], [24] and no effective policy to control the spreading in the framework of evolutionary dynamics has been studied. In a preliminary work [1], we have proposed a new formalism for evolutionary dynamics, which presents two novelties with respect to the original one: i) the spreading process is modeled through a link-based (instead

Some of the results in the paper appeared in preliminary form in [1].

The authors are with the Department of Mathematical Sciences “G.L. Lagrange,” Politecnico di Torino, 10129 Torino, Italy (e-mail: {giacomo.como; fabio.fagnani}@polito.it).

Lorenzo Zino is now with the Faculty of Science and Engineering, University of Groningen, 9747 AG Groningen, The Netherlands (e-mail: lorenzo.zino@rug.nl)

Giacomo Como is also with the Department of Automatic Control, Lund University, BOX 118, 22100 Lund, Sweden, where he is a member of the excellence centre ELLIIT.

This work was partially supported by MIUR grant Dipartimenti di Eccellenza 2018–2022 [CUP: E11G18000350001], the Swedish Research Council through Project Research Grant 2015-04066, and by the Compagnia di San Paolo through a Starting Grant.

of a node-based as in [20]) activation mechanism; and ii) an exogenous control action is incorporated to model the controlled introduction of the novel species in a geographic region. This change of perspective and the explicit introduction of a control action have enabled us to gain new analytical insight concerning the expected duration of the spreading process. In [1], we introduced a first feedback control policy to speed up the spreading process, but its feasibility was limited to very specific networks and it presented two drawbacks: it was extremely sensible to small data errors and very costly from the control viewpoint. An improved feedback control policy was proposed in [25] and there tested on a real-world case study by means of Monte Carlo numerical simulations.

In this paper, we undertake a fundamental analysis of controlled evolutionary network system in great generality, encompassing both open-loop and feedback control policies. Control actions are evaluated on the basis of their cost — measured in terms of both the number of nodes where the control acts and of its total energy— and the spreading time of the induced controlled evolutionary network systems. Our main contribution is the establishment of fundamental limitations between these quantities. Specifically, Theorem 1 and Corollary 2 state general bounds on the expected spreading time in terms of the control policy implemented, its cost, and the topological network properties: notably, we highlight the key role played by the network expansiveness. From these general results, we derive an array of ready-to-use tools to estimate these quantities for specific control policies and network structures. In particular, this allows for performing a full theoretical analysis of the feedback control law presented in [25], assessing its performance.

From a technical viewpoint, our study is based on the estimation of the expected time needed for the Markov process modeling the controlled network evolutionary dynamics to reach its absorbing state. The presence of the external control makes this Markov process non-homogeneous and hinders a direct application of established results on the convergence time of homogeneous Markov processes [26], [27]. Our analysis is based on two key observations. The first one is that the considered controlled evolutionary network systems present a natural monotonicity property with respect to the initial configuration of nodes in the novel state as well with respect to the parameter measuring the evolutionary advantage of the novel state. Specifically, such monotonicity property can be formalized in terms of stochastic domination for the induced Markov process. The second observation is that, as in most epidemic models based on contact processes [9], the speed of diffusion of the novel state is proportional to the size of the boundary separating the set of nodes in the novel state from that of nodes in the original one in the network. When the network is an expansive graph, whereby the size of the boundary of subsets of nodes is proportional to their cardinality, this creates a positive feedback for the considered network evolutionary dynamics that is shown to imply fast spread (in a time that is logarithmic in the network size), even in the absence of an external control (except for the initial seeding). On the other hand, in non-expansive networks, the graph topology typically prevents this positive feedback, as the

system may reach configurations corresponding to bottlenecks where the spreading process is significantly slowed down. It is in these configurations that an external control is the most valuable in reinforcing the spreading process. In fact, the feedback control policies analyzed in Section V are based exactly on the idea of concentrating the control efforts when the system has entered configurations whose boundary is too small to guarantee a fast spread.

To better illustrate the implications of our general results, we present the rigorous analysis of three fundamental network examples: expander graphs (encompassing many models of social networks such as small-world [28] and scale-free networks [29]), stochastic block models (SBMs) [30] and ring graphs. Expander and ring graphs are presented as extreme instances of highly and poorly connected structures, respectively, and constitute benchmarks for fast and slow spread behaviors, respectively. On the other hand, SBM is an established model to represent geographic regions and real-world clustered communities [31] and it exhibits a quite interesting behavior. Our theoretical results show that, for SBMs, fast diffusion can not be reached using open-loop control policies, it can instead be achieved using the proposed feedback control law. Finally, Monte Carlo numerical simulations are offered to corroborate our theoretical results on these three examples. We believe that the capability of incorporating and analyzing control architectures in spreading dynamics and, more generally, in all types of network dynamics is a crucial step from the application viewpoint.

The rest of this paper is organized as follows. In Section II, we introduce the model. In Section III, we present the main technical results. Section IV is devoted to a detailed analysis of open-loop control policy, in particular constant control policies, for the three fundamental network examples. In Section V, we propose and analyze a feedback control policy, proving its effectiveness for SBMs. Section VI summarizes the work and outlines future research. The paper ends with an Appendix containing the proofs of the most technically involved results.

A. Notation

We gather here some notational conventions followed throughout the paper. The n -dimensional all-0 and all-1 vectors are denoted by $\mathbf{0}$ and $\mathbf{1}$, respectively, $\delta^{(i)}$ denotes a vector with a 1 in the i th entry and 0 otherwise, and, for every subset $\mathcal{S} \subseteq \{1, \dots, n\}$, $\delta^{(\mathcal{S})} = \sum_{i \in \mathcal{S}} \delta^{(i)}$. We denote by \mathbb{R}_+ the set of non-negative real numbers. Given a vector $\mathbf{x} \in \mathbb{R}^n$, \mathbf{x}^\top denotes its transpose. Given two vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, $\mathbf{x} \geq \mathbf{y}$ denotes entrywise inequality, i.e., $x_i \geq y_i$, for all $i \in \{1, \dots, n\}$. By $\mathbb{1}_A$ we denote the indicator function of set A . By convention, the value of an empty sum is equal to 0 and the ratio between a positive number and 0 is $+\infty$. Finally, the left- and right-hand limits of a function f in t_0 are denoted by

$$f(t_0^-) := \lim_{t \nearrow t_0} f(t), \quad f(t_0^+) := \lim_{t \searrow t_0} f(t).$$

respectively.

II. THE MODEL

In our model, the network is described as a weighted undirected graph whose nodes represent agents that are either in state 0 (original state) or in state 1 (novel state). A spreading mechanism and an exogenous control determine the evolution of the states of the system over time.

Weighted undirected graph. We consider a connected graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ with node set $\mathcal{V} = \{1, \dots, n\}$, undirected link set \mathcal{E} , and symmetric weight matrix $W \in \mathbb{R}_+^{n \times n}$, whose positive entries $W_{ij} = W_{ji} > 0$ correspond to the weights of the links $\{i, j\} \in \mathcal{E}$.

Spreading mechanism. Each undirected link $\{i, j\}$ of the graph \mathcal{G} is equipped with an independent rate- W_{ij} Poisson clock modeling the interactions between nodes i and j . When the clock associated with link $\{i, j\}$ ticks, if both nodes i and j are in the same state, nothing happens. Otherwise, if the current states of nodes i and j differ from one another, a conflict takes place and the winning state occupies both nodes. Conflicts are solved in a probabilistic way: the novel state 1 wins a conflict (independently of the others) with a fixed probability β , while the original state 0 wins with probability $1 - \beta$. The parameter β in $[0, 1]$ captures the *evolutionary advantage* of the original state (if $\beta < 1/2$) or of the novel state (if $\beta > 1/2$) in the considered network dynamics.

Exogenous control. We fix a locally integrable function $U : \mathbb{R}_+ \rightarrow \mathbb{R}_+^n$ whose i -th entry $U_i(t)$ represents the control rate at which the novel state is enforced at node i at time t .

The triple $(\mathcal{G}, \beta, U(t))$ shall be referred to as the *controlled evolutionary dynamics*. To it, we now associate a Markov process $X(t)$ that describes the spread of the novel state in the network [32]. The process $X(t)$ takes values in the configuration space $\{0, 1\}^n$ and has the following interpretation:

$$X_i(t) = \begin{cases} 0 & \text{if at time } t \text{ node } i \text{ is in the original state,} \\ 1 & \text{if at time } t \text{ node } i \text{ is in the novel state.} \end{cases}$$

The evolution of $X(t)$ is described as a time non-homogeneous jump Markov process where the only transitions that can possibly take place from $X(t) = \mathbf{x}$ are towards configurations that differ from \mathbf{x} in a single entry. Specifically, given a configuration \mathbf{x} and a node i , we consider the transition rates from \mathbf{x} to $\mathbf{x} + \delta^{(i)}$ (if $x_i = 0$) and to $\mathbf{x} - \delta^{(i)}$ (if $x_i = 1$) at time t formally defined as

$$\lambda_i^\pm(\mathbf{x}, t) := \lim_{h \searrow 0} \frac{1}{h} \mathbb{P}[X(t+h) = \mathbf{x} \pm \delta^{(i)} | X(t) = \mathbf{x}],$$

as illustrated in Figure 1 (see [32] for more details on Markov processes). Our model is then completely described by putting

$$\begin{aligned} \lambda_i^+(\mathbf{x}, t) &= (1 - x_i)(\beta(W\mathbf{x})_i + U_i(t)) \\ \lambda_i^-(\mathbf{x}, t) &= x_i(1 - \beta)(W(\mathbf{1} - \mathbf{x}))_i. \end{aligned} \quad (1)$$

Notice how the network topology and the exogenous control determine which transitions can actually take place and their corresponding rates. Indeed, the rate of transition with which the state of node i can jump from 0 to 1 at time t is the sum of two terms: a first term $\beta(W\mathbf{x})_i$ that is proportional to the total weight of links pointing towards neighbors of node i




Fig. 1: Transitions of the jump Markov process $X(t)$.

that are already in state 1 at time t ; and a second term $U_i(t)$ given by the control effort exerted on node i at time t . On the other hand, for the same node i , the rate of transition from 1 to 0 is determined by the term $(1 - \beta)(W(\mathbf{1} - \mathbf{x}))_i$ that is proportional to the total weight of links pointing towards neighbors of node i that are already in state 0 at time t .

Remark 1: In the absence of the external control (i.e., if $U(t) = \mathbf{0}$ for every $t \geq 0$), the model reduces to a voter model [12], which is biased if $\beta \neq 1/2$ [33]. A homogeneous version of our model without control and where clocks are associated with nodes instead of links, has been proposed in [20]. However, the analytical results on such a model are limited to the computation or the estimation of the probability that the novel state diffuses to the whole network (named fixation probability), while more in-depth analysis is limited to specific network structures [21]–[22], and the convergence time is studied only through Monte Carlo simulations.

Remark 2: An equivalent model can be obtained if we replace the exogenous controller with a fictitious stubborn node s having fixed state $X_s(t) = 1$, for $t \geq 0$ and we add time-varying links from every other node i to s having weights $W_{is} = U_i(t)/\beta$. In particular, if $U_i(t) = 0$ we assume that there is no link between i and s at time t . The dynamics driven by the spreading mechanism on this (time-varying) graph leads to exactly the same Markov process. In opinion dynamics, models with stubborn nodes have been analyzed [34]–[36]. These models typically consider the presence of multiple stubborn nodes having possibly different states but assume the graph to be constant and the main problem addressed concerns the way the various stubborn nodes influence the asymptotic behavior of the system. The problem considered in this paper concerns instead the transient behavior of the system and its analysis requires different techniques.

Throughout the paper, we make the following assumptions.

Assumption 1: The probability that in a conflict the novel state wins over the original one is $\beta > 1/2$.

Assumption 2: If $X(t) = \mathbf{0}$, then $U(t) \neq \mathbf{0}$.

The first one captures an evolutionary advantage of the novel state and the second one the presence of an exogenous control to force the evolutionary dynamics whenever the system is in the novel-free configuration $\mathbf{0}$.

From the structure of the transition rates (1), Assumption 2, and the fact that the graph \mathcal{G} is undirected and connected, it follows that the all-1 configuration $\mathbf{1}$ is the only absorbing state of the system and it is reachable from every other state. Hence, the novel state eventually spreads to the whole network almost surely in finite time [32]. From an application perspective, our interest is to shed light on the transient behavior of the system. To this aim, we introduce the following two performance

indices:

- the *spreading time*, that is,

$$T := \inf \{t \geq 0 : X(t) = \mathbf{1}\};$$

- the *control cost*, that is,

$$J = \int_0^T \mathbf{1}^\top U(t) dt.$$

In this paper, we will mostly focus on the estimation of the expected values of these two indices. Since these may depend on the initial configuration of the system $X(0)$, we shall use the following notation for the conditional expected values of the spreading time and of the control cost, respectively:

$$\begin{aligned} \mathbb{E}_{\mathbf{x}_0}[T] &:= \mathbb{E}[T | X(0) = \mathbf{x}_0], \\ \mathbb{E}_{\mathbf{x}_0}[J] &:= \mathbb{E}[J | X(0) = \mathbf{x}_0]. \end{aligned}$$

For the purpose of this paper, we will typically be interested in the scenario with $X(0) = \mathbf{0}$, that is the novel-free configuration. Hence, most of our results will be expressed as bounds on $\mathbb{E}_0[T]$ and $\mathbb{E}_0[J]$.

We will consider two main types of control policy, described in the following.

Open-loop control policies, where the control signal $U(t)$ is predetermined. A simple example of such control policies are those with $U(t) = \mathbf{u}$ constant for every $t \geq 0$. We will refer to these as *constant control policies*.

Feedback control policies, where the control signal $U(t)$ is chosen as a function of the process $X(t)$ itself. Precisely, in this case we consider a function $\nu : \{0, 1\}^n \rightarrow \mathbb{R}_+^n$ and we take $U(t) = \nu(X(t))$. The triple $(\mathcal{G}, \beta, \nu)$ is then called a *feedback controlled evolutionary dynamics*. In this case, defining the transition rates of the process $X(t)$ as in (1), we have that $X(t)$ is a time-homogeneous jump Markov process.

There are typically some constraints that we want to enforce on the admissible control policies. In general, the external control $U(t)$ is constrained to be active only at certain specified nodes, denoted by $\mathcal{U} \subseteq \mathcal{V}$. In this case, the triple $(\mathcal{G}, \beta, U(t))$ is called an \mathcal{U} -*controlled evolutionary dynamics*.

We introduce here some fundamental quantities that will be used in the following.

Definition 1: Consider an undirected weighted graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$, with $|\mathcal{V}| = n$.

- Given a subset of nodes $\mathcal{W} \subseteq \mathcal{V}$, we define its *weighted boundary* [37] as

$$\zeta(\mathcal{W}) = \sum_{i \in \mathcal{W}} \sum_{j \notin \mathcal{W}} W_{ij}. \quad (2)$$

Since W is symmetric, then $\zeta(\mathcal{W}) = \zeta(\mathcal{V} \setminus \mathcal{W})$. Since \mathcal{G} is connected, then $\zeta(\mathcal{W}) > 0$, for all $\mathcal{W} \notin \{\mathcal{V}, \emptyset\}$.

- The *minimum conductance profile* of \mathcal{G} is a function $\phi : \{1, \dots, n-1\} \rightarrow \mathbb{R}_+$, defined as

$$\phi(a) = \min_{\mathcal{W} \subset \mathcal{V}, |\mathcal{W}|=a} \zeta(\mathcal{W}). \quad (3)$$

- The *maximum expansiveness profile* is a function $\eta : \{1, \dots, n-1\} \rightarrow \mathbb{R}_+$, defined as

$$\eta(a) = \max_{\mathcal{W} \subset \mathcal{V}, |\mathcal{W}|=a} \zeta(\mathcal{W}). \quad (4)$$

We also define three one-dimensional stochastic processes as aggregate statistics of the process $X(t)$:

- the *number of nodes* with the novel state

$$A(t) := \mathbf{1}^\top X(t);$$

- the size of the *boundary* of the set of nodes that are in the novel state

$$B(t) := X(t)^\top W(\mathbf{1} - X(t)); \text{ and}$$

- the *effective control rate* in nodes that are in the original state

$$C(t) := (\mathbf{1} - X(t))^\top U(t).$$

III. GENERAL PERFORMANCE ANALYSIS

In this section, we present a series of theoretical results that allow one to estimate the performance of controlled evolutionary dynamics. First, we focus on results guaranteeing the expected spreading time to be below a certain bound that depends on the network topology and the form and intensity of the control effort. Then, we present an array of results illustrating fundamental limitations of the controlled evolutionary dynamics. They are expressed in the form of control-independent lower bounds on the expected spreading time. Leveraging these general results, ready-to-use corollaries will be derived for specific choices of the control policy.

A. Performance Guarantees

In this subsection, we present a general result estimating the performance of controlled evolutionary dynamics, which will be then applied to both open-loop and feedback control policies.

The core of the proposed analysis consists in the estimation of the expected time spent by the process $X(t)$ in the various non-absorbing configurations before reaching the all-1 absorbing configuration. To perform such an analysis, we focus on the aggregate process $A(t) = \mathbf{1}^\top X(t)$ counting the number of state-1 nodes in the current configuration, hence taking values in $\{0, 1, \dots, n\}$, and estimate the time spent by $A(t)$ before being absorbed in state n . The analysis of this process is nontrivial because, in general, $A(t)$ is not a Markov process. Indeed, notice that each transition of $X(t)$ increases or decreases $A(t)$ by 1 unit and that, by (1), in general such transitions depend on the complete vector $X(t)$ through the network weight matrix W and not just on the aggregate variable $A(t)$. From (1), we can compute the transition rates of the process $A(t)$ conditioned on $X(t)$. Precisely, when $A(t) = a$, its rate of increase (corresponding to the transition to $a+1$) conditioned on $X(t) = \mathbf{x}$ is equal to

$$\begin{aligned} \lambda_t^+(a|\mathbf{x}) &:= \lim_{h \searrow 0} \frac{1}{h} \mathbb{P}[A(t+h) = a+1 | X(t) = \mathbf{x}] \\ &= \sum_{i \in \mathcal{V}} \lambda_i^+(\mathbf{x}, t) \\ &= \beta(\mathbf{1} - \mathbf{x})^\top W \mathbf{x} + (\mathbf{1} - X(t))^\top U(t) \\ &= \beta B(t) + C(t). \end{aligned} \quad (5)$$




Fig. 2: Transition graph of the continuous-time process $A(t)$.

Notice that this quantity depends on the configuration $X(t)$ only through the two scalar aggregate statistics $B(t)$ and $C(t)$. Similarly, the rate of decrease of $A(t)$ is equal to

$$\begin{aligned} \lambda_t^-(a|\mathbf{x}) &= \frac{1}{h} \lim_{h \searrow 0} \mathbb{P}[A(t+h) = a-1 | X(t) = \mathbf{x}] \\ &= (1-\beta)B(t), \end{aligned} \quad (6)$$

and this quantity depends on the configuration $X(t)$ only through the size of the boundary $B(t)$. The transitions of the aggregate process $A(t)$ and their corresponding rates are shown in Figure 2.

In the following, it will be useful to consider the transition probabilities associated to these transition rates [32]. Precisely, for all $t \geq 0$, $a \in \{1, \dots, n-1\}$ and $\mathbf{x} \in \{0, 1\}^n$ let $p_t^+(a|\mathbf{x})$ ($p_t^-(a|\mathbf{x})$) be the conditional probability that, should the process $X(t)$ have a state transition at time t , such transition would increase the value of $A(t)$ from a to $a+1$ (respectively, decrease the value of $A(t)$ from a to $a-1$), given the current configuration $X(t^-) = \mathbf{x}$. We have

$$p_t^+(a|\mathbf{x}) = \frac{\lambda_t^+(a|\mathbf{x})}{\lambda_t^+(a|\mathbf{x}) + \lambda_t^-(a|\mathbf{x})} = \frac{\beta B(t) + C(t)}{B(t) + C(t)} \geq \beta, \quad (7)$$

$$p_t^-(a|\mathbf{x}) = \frac{\lambda_t^-(a|\mathbf{x})}{\lambda_t^+(a|\mathbf{x}) + \lambda_t^-(a|\mathbf{x})} = \frac{(1-\beta)B(t)}{B(t) + C(t)} \leq 1-\beta, \quad (8)$$

i.e., these two probabilities can be uniformly bounded by functions that are independent of the state \mathbf{x} and time t , and only depend on the evolutionary advantage β . For $a=0$, such transition probabilities reduce to $p_t^+(0|\mathbf{0}) = 1$ and $p_t^-(0|\mathbf{0}) = 0$, for every $t \geq 0$.

In our theoretical analysis, we need to bound the conditional probability that, given that the system is in a given configuration $X(t_0) = \mathbf{x}$ with $\mathbf{1}^\top \mathbf{x} = a$ state-1 nodes at time $t_0 \geq 0$, the aggregate process $A(t)$ will ever go below a at some time $t > t_0$, before being absorbed into its absorbing state n . Since the transition rates of the aggregate process $A(t)$ depend on the whole non-homogeneous Markov process $X(t)$, such probability may in general depend of the configuration of the system $X(t_0)$ and on the time t_0 . In our derivations, we will need a lower bound on such a quantity that is uniform with respect to these two variables. To this aim, we define the following lower bound on such a probability:

$$q_a := \inf_{t_0 \geq 0} \min_{\mathbf{x} : \mathbf{1}^\top \mathbf{x} = a} \mathbb{P}[A(t) \geq a, \forall t \geq t_0 | X(t_0) = \mathbf{x}].$$

Such a uniform bound is estimated using (7) in the following result. Its proof uses standard coupling arguments between stochastic processes and, for the sake of making the paper self-contained, is reported in Appendix A.

Lemma 1: For every $a \in \{1, \dots, n-1\}$, $q_a \geq (2\beta - 1)/\beta$.

Observe that Lemma 1 provides a lower bound on q_a and, ultimately, on the desired probability, depending only on the evolutionary advantage β . Using this uniform bound, we can now estimate the total amount of time that the process spends in each of its non-absorbing states, when a relation between the three aggregate statistics $A(t)$, $B(t)$, and $C(t)$ is verified. Specifically, the following is the key technical result which allows us to finally formulate our performance guarantees.

Proposition 1: Let T_a be the random variable measuring the total amount of time spent by process $A(t)$ in state a before being absorbed in n . Then, for every $f : \{0, \dots, n-1\} \rightarrow \mathbb{R}_+$ such that

$$B(t) + C(t) \geq f(A(t)), \quad (9)$$

for every $t \geq 0$ such that $X(t) \neq \mathbf{1}$, the conditional expected spreading time satisfies

$$\mathbb{E}[T_a | X(0) = \mathbf{0}] \leq \begin{cases} \frac{\beta}{2\beta-1} \frac{1}{f(0)} & \text{if } a=0, \\ \frac{1}{2\beta-1} \frac{1}{f(a)} & \text{if } a \neq 0. \end{cases} \quad (10)$$

Proof: For $s \geq 0$ and $a \in \{0, 1, \dots, n-1\}$, let $T_a(s)$ denote the time spent by the process $A(t)$ in state a from time s on. Clearly, $T_a = T_a(0)$ and, for every $s \geq 0$ and $\mathbf{x}_0 \in \{0, 1\}^n$,

$$\mathbb{E}[T_a(s) | X(s) = \mathbf{x}_0, X(0) = \mathbf{0}] = \mathbb{E}[T_a(s) | X(s) = \mathbf{x}_0],$$

because of the Markov property. For $a \in \{0, \dots, n-1\}$, put

$$\theta_a := \sup_{s \geq 0} \max_{\mathbf{x}_0 : \mathbf{1}^\top \mathbf{x}_0 \leq a} \mathbb{E}[T_a(s) | X(s) = \mathbf{x}_0]. \quad (11)$$

For $t \geq 0$, we define the random variable

$$S(t) = \inf\{h \geq 0 : X((t+h)^+) \neq X(t^+)\},$$

which models the waiting time for the first configuration change after time t . To estimate its expected value conditioned to the event $X(t^+) = \mathbf{x}$, we observe that, for $r \in [t, t+S(t)]$, the total jump rate of the process $A(t)$ can be estimated, using (5) and (6) and the relation in (9), as

$$\xi(r) = \lambda_r^+(\mathbf{1}^\top \mathbf{x} | \mathbf{x}) + \lambda_r^-(\mathbf{1}^\top \mathbf{x} | \mathbf{x}) = B(r) + C(r) \geq f(\mathbf{1}^\top \mathbf{x})$$

Standard properties of non-homogeneous Markov jump processes then yield

$$\begin{aligned} \mathbb{E}[S(t) | X(t^+) = \mathbf{x}] &= \int_0^{+\infty} e^{-\int_t^{t+s} \xi(r) dr} ds \\ &\leq \int_0^{+\infty} e^{-\int_t^{t+s} f(\mathbf{1}^\top \mathbf{x}) dr} ds \\ &= \int_0^{+\infty} e^{-sf(\mathbf{1}^\top \mathbf{x})} ds \\ &\leq \frac{1}{f(\mathbf{1}^\top \mathbf{x})}. \end{aligned}$$

We fix an initial time t_0 (not necessarily equal to 0) and an initial state $X(t_0) = \mathbf{x}_0$ such that $\mathbf{1}^\top \mathbf{x}_0 \leq a$. Let $t_1 \geq t_0$ be the time when $A(t)$ reaches the value a for the first time

after t_0 . Then, using Markov property and separating the time spent in a during its first entrance from the other contributions to $T_a(t_0)$, we write the following recursive expression:

$$\begin{aligned} & \mathbb{E}[T_a(t_0) | X(t_0) = \mathbf{x}_0, X(t_1^+) = \mathbf{x}_1] \\ &= \mathbb{E}[T_a(t_1) | X(t_1^+) = \mathbf{x}_1] \\ &= \mathbb{E}[S(t_1) | X(t_1^+) = \mathbf{x}_1] + \mathbb{E}[T_a(t_1 + S(t_1)) | X(t_1^+) = \mathbf{x}_1]. \end{aligned} \quad (12)$$

Let us define $t_2 := t_1 + S(t_1)$ and $\mathbf{x}_2 := X(t_2^+)$. Then, we estimate $\mathbb{E}[T_a(t_2) | X(t_1^+) = \mathbf{x}_1]$ depending on the direction of the jump that has occurred at time t_2 . If $A(t)$ has moved backward, then $\mathbf{1}^\top \mathbf{x}_2 = a - 1$ and, by the definition in (11),

$$\mathbb{E}[T_a(t_2) | \mathbf{1}^\top \mathbf{x}_2 = a - 1] \leq \theta_a. \quad (13)$$

If $A(t)$ instead has increased to $a + 1$, the probability that the process $A(t)$ will return to a before getting absorbed in n is bounded from the above by $(1 - q_{a+1})$, yielding

$$\mathbb{E}[T_a(t_2) | \mathbf{1}^\top \mathbf{x}_2 = a + 1] \leq (1 - q_{a+1})\theta_a. \quad (14)$$

Inserting (13) and (14) into (12), we bound

$$\begin{aligned} & \mathbb{E}[T_a(t_0) | X(t_0) = \mathbf{x}_0, X(t_1^+) = \mathbf{x}_1] \\ & \leq \frac{1}{f(a)} + p_{t_2^-}^-(a | \mathbf{x}_1)\theta_a + p_{t_2^+}^-(a | \mathbf{x}_1)(1 - q_{a+1})\theta_a. \end{aligned} \quad (15)$$

For $a = 0$, we have $p_{t_2^-}^-(0 | \mathbf{x}_1) = 0$ and $p_{t_2^+}^-(0 | \mathbf{x}_1) = 1$. Hence, using Lemma 1 we obtain

$$\mathbb{E}[T_0(t_0) | X(t_0) = \mathbf{0}] \leq \frac{1}{f(0)} + (1 - f_1)\theta_0 \leq \frac{1}{f(0)} + \frac{1 - \beta}{\beta}\theta_0.$$

This holds true for every $t_0 \geq 0$ and thus, by taking the supremum of the left-hand side with respect to t_0 , we obtain

$$\theta_0 \leq \frac{1}{f(0)} + \frac{1 - \beta}{\beta}\theta_0,$$

so that

$$\theta_0 \leq \frac{\beta}{2\beta - 1} \frac{1}{f(0)}. \quad (16)$$

For $a \in \{1, \dots, n - 1\}$, applying (7) and (8) to the right-hand side of (15), we obtain

$$\begin{aligned} & \mathbb{E}[T_a(t_0) | X(t_0) = \mathbf{x}_0, X(t_1^+) = \mathbf{x}_1] \\ & \leq \frac{1}{f(a)} + (1 - \beta)\theta_a + \beta \frac{1 - \beta}{\beta}\theta_a \\ & = \frac{1}{f(a)} + 2(1 - \beta)\theta_a. \end{aligned} \quad (17)$$

Since this holds true for every $t_0 \geq 0$ and $\mathbf{x}_0 \in \{0, 1\}^n$ such that $\mathbf{1}^\top \mathbf{x}_0 \leq a$, we immediately obtain that

$$\theta_a \leq \frac{1}{f(a)} + 2(1 - \beta)\theta_a,$$

yielding

$$\theta_a \leq \frac{1}{2\beta - 1} \frac{1}{f(a)}. \quad (18)$$

Since $\mathbb{E}[T_a | X(0) = \mathbf{0}] \leq \theta_a$, the claim now follows from (16) and (18). \blacksquare

We can now present and prove the fundamental result of this section.

Theorem 1: Let $(\mathcal{G}, \beta, U(t))$ be a controlled evolutionary dynamics with initial configuration $X(0) = \mathbf{0}$. Then, for every $f : \{0, \dots, n - 1\} \rightarrow \mathbb{R}_+$ such that

$$B(t) + C(t) \geq f(A(t)), \quad (19)$$

for $t \geq 0$, the conditional expected spreading time satisfies

$$\mathbb{E}_{\mathbf{0}}[T] \leq \frac{\beta}{(2\beta - 1)f(0)} + \frac{1}{2\beta - 1} \sum_{a=1}^{n-1} \frac{1}{f(a)}. \quad (20)$$

Proof: It follows from Proposition 1 that

$$\begin{aligned} \mathbb{E}_{\mathbf{0}}[T] &= \mathbb{E}\left[\sum_{a=0}^{n-1} T_a \mid X(0) = \mathbf{0}\right] \\ &= \mathbb{E}[T_0 | X(0) = \mathbf{0}] + \sum_{a=1}^{n-1} \mathbb{E}[T_a | X(0) = \mathbf{0}] \\ &\leq \frac{\beta}{(2\beta - 1)f(0)} + \frac{1}{2\beta - 1} \sum_{a=1}^{n-1} \frac{1}{f(a)}, \end{aligned}$$

thus proving the claim. \blacksquare

Notice how the quantity $B(t) + C(t)$ appearing in the left-hand side of (19) can be interpreted, summing the two relations in (5) and (6), as the total intensity of the process $A(t)$ (or, equivalently, of $X(t)$).

As special case of Theorem 1 for constant control policies, we have the following result.

Corollary 1: Consider a controlled evolutionary dynamics $(\mathcal{G}, \beta, \mathbf{u})$ under constant control policy $U(t) = \mathbf{u}$ with initial configuration $X(0) = \mathbf{0}$ and let ϕ be the minimum conductance profile of \mathcal{G} , defined in (3). Then, the conditional expected spreading time satisfies

$$\mathbb{E}_{\mathbf{0}}[T] \leq \frac{\beta}{(2\beta - 1)\mathbf{1}^\top \mathbf{u}} + \frac{1}{2\beta - 1} \sum_{a=1}^{n-1} \frac{1}{\phi(a)}. \quad (21)$$

Proof: Observe that, for every $a \in \{1, \dots, n\}$, for all subsets $S \subset \mathcal{V}$ with $|S| = a$, we have $\phi(a) \leq \zeta(S)$. Therefore, if $A(t) = a$, then $B(t) + C(t) \geq \phi(a)$. If $A(t) = 0$, then $C(t) = \mathbf{1}^\top \mathbf{u}$. Finally, (20) from Theorem 1 is applied with $f(a) = \phi(a)$, for $a \neq 0$, and $f(0) = \mathbf{1}^\top \mathbf{u}$. \blacksquare

B. Fundamental Performance Limitation

In this subsection we present a series of results illustrating fundamental limitations of the controlled evolutionary dynamics. We start by establishing two preliminary monotonicity properties of the process, which will be instrumental to our main results. For the sake of readability, proofs are reported in Appendices B and C, respectively.

Lemma 2: Let $(\mathcal{G}, \beta, U(t))$ and $(\mathcal{G}, \gamma, U(t))$ be two controlled evolutionary dynamics and let $X(t)$ and $Y(t)$ be, respectively, the corresponding Markov processes with initial conditions $X(0) = \mathbf{x}_0$ and $Y(0) = \mathbf{y}_0$. If $\beta \leq \gamma$ and $\mathbf{x}_0 \leq \mathbf{y}_0$, then $\mathbb{E}_{\mathbf{y}_0}[T_Y] \leq \mathbb{E}_{\mathbf{x}_0}[T_X]$ and $\mathbb{E}_{\mathbf{y}_0}[J_Y] \leq \mathbb{E}_{\mathbf{x}_0}[J_X]$, where the subscripts of T and J refer to the processes they are associated to.

Lemma 3: Let $(\mathcal{G}, 1, U(t))$ and $(\mathcal{G}, 1, 0)$ be two controlled evolutionary dynamics and let $X(t)$ and $Y(t)$ be, respectively,

the corresponding Markov processes with initial conditions $X(0) = \mathbf{x}_0$ and $Y(0) = \mathbf{y}_0$. If $\mathbf{x}_0 \leq \mathbf{y}_0$ and $(\mathbf{1} - \mathbf{y}_0)^\top U(t) = 0$, for every $t \geq 0$, i.e., if the uncontrolled process $Y(t)$ has initial condition equal to 1 in all the nodes in which the control is exerted in the controlled process $X(t)$, then, $\mathbb{E}_{\mathbf{y}_0}[T_Y] \leq \mathbb{E}_{\mathbf{x}_0}[T_X]$, where the subscripts of T and J refer to the processes they are associated to.

Remark 3: These results have straightforward consequences:

- 1) Lemma 2 implies that any lower bound on the expected spreading time obtained for the case when $\beta = 1$ will automatically yield a lower bound for every value of β .
- 2) Lemma 3 implies that, for a controlled evolutionary dynamics where the novel state always wins (i.e., when $\beta = 1$), it is always possible to establish a lower bound on the expected spreading time by considering an uncontrolled process with initial condition equal to 1 in all the nodes in which the external control is exerted. We wish to emphasize that controlled evolutionary dynamics with $\beta < 1$ do not, in general, enjoy this monotonicity property.

The previous results motivate a deeper analysis of the controlled evolutionary dynamics of type $(\mathcal{G}, 1, U(t))$. In this case, if started with an initial configuration $X(0)$ such that $\mathbf{1}^\top X(0) = k$, the Markov process $X(t)$, will always undergo exactly $n - k$ transitions before being absorbed in the all-1 configuration. Indeed, for $\beta = 1$ (1) shows that $\lambda_i^-(\mathbf{x}, t) = 0$ for every configuration \mathbf{x} , node i and time $t \geq 0$. This says that the process can only undergo transitions where the number of 1's is increasing, either driven by the spreading mechanism or by the external control.

We let $0 = T_k < T_{k+1} < \dots < T_n = T$ be the random times at which these $n - k$ jumps occur and we denote by $X_h = X(T_h^+)$ the configurations of the process after the corresponding jumps, so that $X_k = X(0)$ and $X_n = \mathbf{1}$. According to this notation, T_h is the time of the jump from X_{h-1} to X_h . Finally, we let B_h to be the corresponding values for the boundary of the process when in configuration X_h , that is $B_h = \zeta(X_h)$. This time process can be described recursively as follows. We first let σ_h to be the σ -algebra generated by T_h and by the process $X(t)$ for $t \leq T_h$. Given σ_{h-1} we consider two independent random variables t_h^s and t_h^c whose distribution functions are given, respectively, by

$$\begin{aligned} \mathbb{P}[t_h^s \geq t | \sigma_{h-1}] &= \exp(-B_{h-1}t), \\ \mathbb{P}[t_h^c \geq t | \sigma_{h-1}] &= \exp\left(-\int_{T_{h-1}}^{T_{h-1}+t} C(s)ds\right). \end{aligned} \quad (22)$$

We then put $t_h = \min\{t_h^s, t_h^c\}$. The interpretation of t_h is the following: from state X_{h-1} the system can evolve either through the spreading mechanisms or through an occurrence of the external control in a node currently in state 0; t_h^s and t_h^c are exponentially distributed random variables that model the random times at which the two phenomena would independently take place, according to their definition. According to the properties of Markov processes [32], the minimum of

these two random variables models the time to wait for the next jump, that is, $T_h = T_{h-1} + t_h$. Define

$$c_h = \int_{T_{h-1}}^{T_h} C(s)ds$$

to be the effective control rate exerted during the interval $[T_{h-1}, T_h]$. The following result shows that there is an exact algebraic relation that ties the average length of the interval $[T_{h-1}, T_h]$, the average effective control rate in this interval c_h , and the active boundary after the $(h - 1)$ -th jump B_{h-1} .

Proposition 2: For every h such that $B_{h-1} \neq 0$, the following relation holds true

$$\mathbb{E}[T_h - T_{h-1} | \sigma_{h-1}] = \frac{1 - \mathbb{E}[c_h | \sigma_{h-1}]}{B_{h-1}}. \quad (23)$$

Proof: First, observe that $T_h - T_{h-1} = t_h = \min\{t_h^s, t_h^c\}$. Using the property of the minimum of two independent exponentially distributed random variables, we note that

$$\mathbb{P}[t_h \geq t | \sigma_{h-1}] = \exp\left(-\int_{T_{h-1}}^{T_{h-1}+t} (B_{h-1} + C(s))ds\right), \quad (24)$$

which, in particular, yields

$$\frac{d}{dt}\mathbb{P}[t_h \geq t | \sigma_{h-1}] = -(B_{h-1} + C(T_{h-1} + t))\mathbb{P}[t_h \geq t | \sigma_{h-1}],$$

From this and the fact that T_{h-1} and $\{C(s) | s \in [T_{h-1}, T_h]\}$ are both σ_{h-1} -measurable, we have

$$\begin{aligned} \mathbb{E}[c_h | \sigma_{h-1}] &= \mathbb{E}\left[\int_{T_{h-1}}^{+\infty} C(s)\mathbb{1}_{[T_{h-1}, T_h]}(s)ds | \sigma_{h-1}\right] \\ &= \int_{T_{h-1}}^{+\infty} C(s)\mathbb{E}[\mathbb{1}_{[T_{h-1}, T_h]}(s) | \sigma_{h-1}] ds \\ &= \int_{T_{h-1}}^{+\infty} C(s)\mathbb{P}[t_h \geq s - T_{h-1} | \sigma_{h-1}]ds \\ &= \int_{T_{h-1}}^{+\infty} -\frac{d}{ds}\mathbb{P}[t_h \geq s - T_{h-1} | \sigma_{h-1}]ds \\ &\quad - B_{h-1} \int_{T_{h-1}}^{+\infty} \mathbb{P}[t_h \geq s - T_{h-1} | \sigma_{h-1}]ds \\ &= 1 - B_{h-1}\mathbb{E}[t_h | \sigma_{h-1}], \end{aligned} \quad (25)$$

which yields the proof. \blacksquare

If we further define

$$J_h := \int_{T_{h-1}}^{T_h} \mathbf{1}^\top U(s)ds \quad (26)$$

to be the control rate exerted during the interval $[T_{h-1}, T_h]$, we immediately observe that, by definition, $J_h \geq c_h$. (23) yields the following result.

Corollary 2: Assumethat the initial condition $X(0) = \mathbf{x}_0$ is such that $\mathbf{1}^\top \mathbf{x}_0 = k > 0$. Then,

$$\begin{aligned} \mathbb{E}_{\mathbf{x}_0}[T] &\geq \sum_{h=k+1}^n \mathbb{E} \left[\frac{1 - \mathbb{E}[J_h | \sigma_{h-1}]}{B_{h-1}} \right], \\ \mathbb{E}_{\mathbf{x}_0}[J] &= \sum_{h=k+1}^n \mathbb{E}[\mathbb{E}[J_h | \sigma_{h-1}]]. \end{aligned} \quad (27)$$

Proof: The former follows by summing and averaging (23), using that $B_h \neq 0$ for $h = k, \dots, n-1$; the latter is the definition of J , obtained by summing all the terms in (26). ■

Remark 4: In the case when $X(0) = \mathbf{0}$, the relations in (27) have to be modified as follows. The first jump is necessarily triggered by an exogenous activation, which implies

$$\mathbb{E}[t_1] = \int_0^{+\infty} \exp \left(- \int_0^t \mathbf{1}^\top U(s) ds \right) ds. \quad (28)$$

Moreover, (25) yields $\mathbb{E}[c_1] = \mathbb{E}[J_1] = 1$. Therefore, (27) is modified by setting $k = 1$ and adding the term $\mathbb{E}[t_1]$ from (28) to the right-hand side of the inequality for $\mathbb{E}_0[T]$ and 1 to the right-hand side of the equality for $\mathbb{E}_0[J]$.

The applicability of (27), in the present general form, is limited by the fact that the control efforts and the boundary evolutions can not be uncoupled when the averaging operation is taken. For \mathcal{U} -controlled evolutionary dynamics, Corollary 2 can be relaxed, yielding more explicit, even if less tight, bounds that turn to be useful in case when \mathcal{U} is sufficiently small.

Corollary 3: Let $(\mathcal{G}, \beta, U(t))$ be a \mathcal{U} -controlled evolutionary dynamics with initial condition $X(0) = \mathbf{x}_0 \leq \delta^{(\mathcal{U})}$. Then,

$$\mathbb{E}_{\mathbf{x}_0}[T] \geq \sum_{h=|\mathcal{U}|}^{n-1} \frac{1}{\eta(h)}, \quad (29)$$

where η is defined as in (4).

Proof: It follows from Lemmas 2 and 3 that it is sufficient to prove the bound for the controlled evolutionary dynamics $(\mathcal{G}, 1, 0)$ with initial condition $Y(0) = \delta^{(\mathcal{U})}$. In this case, (27) reduces to

$$\mathbb{E}_{\mathbf{x}_0}[T] \geq \mathbb{E}_{\delta^{(\mathcal{U})}}[T_Y] \geq \sum_{h=|\mathcal{U}|}^{n-1} \mathbb{E} \left[\frac{1}{B_h} \right].$$

Finally, the claim follows from the definition of $\eta(h)$ in (4). ■

For constant control policies with novel-free initial condition, we can refine the result as follows.

Corollary 4: Let $(\mathcal{G}, \beta, \mathbf{u})$ be a constant \mathcal{U} -controlled evolutionary dynamics with initial condition $X(0) = \mathbf{0}$. Then, the expected spreading time verifies

$$\mathbb{E}_0[T] \geq \frac{1}{\mathbf{1}^\top \mathbf{u}} + \sum_{h=|\mathcal{U}|}^{n-1} \frac{1}{\eta(h)}. \quad (30)$$

Proof: Consider the random time t_1 corresponding to the first jump of the process. From (28), using that $\mathbf{1}^\top \mathbf{u}$ is constant, we compute $\mathbb{E}[t_1] = 1/\mathbf{1}^\top \mathbf{u}$ and applying Corollary 3 with initial condition $X(t_1) \leq \delta^{(\mathcal{U})}$ we obtain the claim. ■

By comparing the lower bound in (30) with the corresponding upper bound in (21), we conclude that, for constant \mathcal{U} -controlled evolutionary dynamics, the control rate $\mathbf{1}^\top \mathbf{u}$ has a limited effect on the performance. In particular, $\mathbb{E}_0[T]$ remains bounded away from 0 even in the limit case as $\mathbf{1}^\top \mathbf{u} \rightarrow +\infty$, even though the expected control cost $\mathbb{E}_0[J] = (\mathbf{1}^\top \mathbf{u})\mathbb{E}_0[T]$ grows large.

Equations (29) and (30) moreover suggest that, together with the network structure, the support \mathcal{U} of the control action may play a key role in achieving suitable spreading performance. In this direction, we propose another estimation of the expected spreading time in which \mathcal{U} has a central role.

Corollary 5: Let $(\mathcal{G}, \beta, U(t))$ be a \mathcal{U} -controlled evolutionary dynamics with initial condition $X(0) = \mathbf{0}$. Then,

$$\mathbb{E}_0[T] \geq \left(\min_{\mathcal{R} \supseteq \mathcal{U}} \zeta(\mathcal{R}) \right)^{-1}.$$

Proof: Using Lemmas 2 and 3, we bound the expected spreading time by considering a controlled evolutionary dynamics $(\mathcal{G}, 1, 0)$ with initial condition $Y(0) = \delta^{(\mathcal{R})}$ where $\mathcal{R} \supseteq \mathcal{U}$. (27) yields

$$\mathbb{E}_0[T] \geq \mathbb{E}_{\delta^{(\mathcal{R})}}[T_Y] \geq \sum_{h=|\mathcal{R}|}^{n-1} \mathbb{E} \left[\frac{1}{B_h} \right] \geq \frac{1}{\zeta(\mathcal{R})},$$

where the last equality follows from the fact that $B_{|\mathcal{R}|}$ is the boundary at the initial condition $Y(0) = \delta^{(\mathcal{R})}$. Since the above inequality holds true for every $\mathcal{R} \supseteq \mathcal{U}$, we obtain the claim. ■

We conclude our analysis with a useful equivalent description of the control cost term. The following relation expresses the term $\mathbb{E}[c_h | \sigma_{h-1}]$ as the probability that the h th jump in the process is due to an exogenous event.

Proposition 3: The following relation holds true

$$\mathbb{E}[c_h | \sigma_{h-1}] = \mathbb{P}[t_h^c < t_h^s | \sigma_{h-1}].$$

Proof: It follows from (22) that

$$\begin{aligned} \mathbb{P}[t_h^c < t_h^s | \sigma_{h-1}] &= \mathbb{P}[t_h^s - t_h^c > 0 | \sigma_{h-1}] \\ &= \int_0^{+\infty} B_{h-1} e^{-B_{h-1}t} \left[1 - \exp \left(- \int_{T_{h-1}}^{T_{h-1}+t} C(s) ds \right) \right] dt \\ &= 1 - B_{h-1} \exp \left(- \int_{T_{h-1}}^{T_{h-1}+t} (B_{h-1} + C(s)) ds \right). \end{aligned}$$

The claim then follows using (24) and (25). ■

Denote now by N^c the random variable counting the total number of activations due to exerting the control action, more formally in the notation above:

$$N^c = |\{h = 1, \dots, n : t_h^c < t_h^s\}|. \quad (31)$$

Summing over h and averaging the second relation in Proposition 3, we obtain that

$$\mathbb{E}[N^c] \leq \mathbb{E}[J]. \quad (32)$$

In other terms, the expected number of activations due to the control action is a lower bound on the control cost.




Fig. 3: Examples of the topologies analyzed in this paper.

IV. THREE FUNDAMENTAL EXAMPLES

In this section, we formally introduce and analyze three key examples of graph families parameterized by their size n : expander graphs, stochastic block models (SBM), and ring graphs. For each of them, we fully analyze the behavior of the corresponding controlled evolutionary dynamics and discuss the corresponding performance behavior as a function of n .

Expander graphs and ring graphs constitute extreme cases of highly and poorly connected structures, respectively, and, as we will see, they represent benchmarks for topologies that are easy to control and hard to control, respectively. SBMs, instead, are representative of real-world geographical networks [31] and display an intermediate behavior: no fundamental limitation precludes fast diffusion, but constant control policies may fail in achieving it. In Section V we show that, for SBMs, feedback control policies play a crucial role.

For the sake of simplicity, in this section we consider graphs \mathcal{G} in which all the links in \mathcal{E} have the same weight, that is, $W_{ij} = w$, for all $\{i, j\} \in \mathcal{E}$. Also, we assume that the value of w is scaled with respect to the maximum degree Δ of the nodes in the graph, namely, we take $w = \alpha/\Delta$ where $\alpha > 0$ is a constant kept fixed as n varies. Such a choice prevents the sum of the weights insisting on each node from blowing up (which would be inconsistent with any real-world application) and guarantees a fair comparison between the topologies in the three examples.

We consider controlled evolutionary dynamics on \mathcal{G} , $(\mathcal{G}, \beta, U(t))$ always with initial condition $X(0) = \mathbf{0}$. First, we notice that the trivial bound $\eta(h) \leq \alpha h$ substituted in (29) yields, for every \mathcal{U} -controlled evolutionary dynamics, the following bound on the expected spreading time

$$\mathbb{E}_0[T] \geq \sum_{h=|\mathcal{U}|}^{n-1} \frac{1}{\alpha h} \geq \frac{1}{\alpha} \log \frac{n}{|\mathcal{U}|}. \quad (33)$$

Hence, if \mathcal{U} is assumed to be constant in n , the expected spreading time grows at least logarithmically in n .

A. Expander graphs

A family of graphs is said to be an *expander* if $\exists \gamma > 0$ such that the minimum conductance profile verifies $\phi(h) \geq \gamma \min\{h, n-h\}$, for all n and $h = 1, \dots, n-1$. Expander graphs are often used to represent well connected real-world systems. Classical models of small-world and scale-free networks, often adopted to model social networks, belong to this family.




Fig. 4: Monte Carlo estimation (200 simulations) of the expected spreading time $\mathbb{E}_0[T]$ on complete graphs for different values of n ($\alpha = 1$), with 90% confidence intervals. The two solid curves are the theoretical bounds from (33) and (34).

We now show how these networks exhibit fast spread. Precisely we demonstrate that, using constant control policies with bounded nonzero rate, the expected spreading time scales logarithmically with respect to the network size.

Let \mathcal{G} be an expander graph with n nodes. Using the expander condition on $\phi(h)$ in Corollary 1, we obtain

$$\mathbb{E}_0[T] \leq \frac{\beta}{(2\beta - 1)\mathbf{1}^\top \mathbf{u}} + \frac{2 \ln(n/2) + 2}{\gamma(2\beta - 1)}. \quad (34)$$

If we stick to control policies supported on a fixed subset \mathcal{U} , the lower bound in (33) concludes that expander graphs are a benchmark for fast-diffusive topologies. Figure 4 shows Monte Carlo estimations of the expected spreading time for complete graphs (in Figure 3(a)), together with our analytical bounds. Note that the bound in (34) is tight for the complete graph. In fact, for complete graphs the inequality (19) holds true as an equality when the function $f(h)$ coincides with the minimum conductance profile (since all the subsets with h nodes have the same weighted boundary). This suggests that the spreading time is close to the upper bound (20) in Theorem 1 when one can determine a function $f(h)$ so that the inequality (19) is close to an equality.

B. Stochastic block models

As a second example we consider *stochastic block models* (SBM), illustrated in Figure 3(b). These graphs are composed of dense (expander) subgraphs linked among each other by few connections and have been often used in the literature to represent social, economical, and geographical structures [30], [31], [38]. For the sake of simplicity, we limit our analysis to the case of two subgraphs (called communities), but our results can be easily generalized. Since SBMs are random graphs, results will be provided with high probability, i.e., with probability converging (at least polynomially) to 1 as the network size n grows. We will prove that spread is slow unless the control is exerted on all of its communities.

In this paper, we consider the following implementation of SBMs. Fixed $c \in (0, 1/2]$, we partition the nodes into two disjoint sets $\mathcal{V} = \mathcal{V}^1 \cup \mathcal{V}^2$, with $n_1 = \lfloor cn \rfloor$ and $n_2 = n - \lfloor cn \rfloor$ nodes, respectively. Without loss of generality, we assume $c \leq 1/2$. The nodes in each subset form a subgraph that is generated according to an Erdős-Rényi random

graph. In both subgraphs, each link is present with probability $p \in (0, 1]$, independent of the others [39]. On top of this, L links positioned uniformly at random connect nodes belonging to the two different subgraphs.

If the control is exerted in only one of the subgraphs, then the expected spreading time grows linearly in n . This can be shown as follows. We fix $\mathcal{U} \subseteq \mathcal{V}^1$ and consider any \mathcal{U} -controlled evolutionary dynamics $(\mathcal{G}, \beta, U(t))$ on \mathcal{G} . Considering the standing assumptions that all non-zero weights are equal to $w = \alpha/\Delta$ and that the maximal degree satisfies the inequality $\Delta \geq n_2 p \geq (1-c)np$ (w.h.p), we have that $\zeta(\mathcal{V}^1) \leq \alpha L / (1-c)np$. Hence, Corollary 5, yields

$$\mathbb{E}_0[T] \geq \frac{(1-c)np}{\alpha L}. \quad (35)$$

We demonstrate that such an estimation is asymptotically order-tight for constant control policies $U(t) = \mathbf{u}$ by leveraging Corollary 1. To this aim, we estimate the minimum conductance profile as follows. First, note that (even though SMBs are no expanders) each one of the two subgraphs is expander with $\gamma = cnpw/2$ [2]. Using the trivial bound $\Delta \leq n$, we get $\gamma \geq c\alpha p/2$. Given an integer $0 < h < n$, any subset $\mathcal{W} \subseteq \mathcal{V}$ with $|\mathcal{W}| = h$ can be written as $\mathcal{W} = \mathcal{W}^1 \cup \mathcal{W}^2$, with $\mathcal{W}^i \subseteq \mathcal{V}^i$, $|\mathcal{W}^i| = h_i$, and $h_1 + h_2 = h$. Therefore,

$$\begin{aligned} \phi(h) &\geq \frac{c\alpha p}{2} \\ &\cdot \min_{\substack{h_1, h_2 \\ h_1 + h_2 = h}} \{ \min\{h_1, n_1 - h_1\} + \min\{h_2, n_2 - h_2\} \} \\ &= \frac{c\alpha p}{2} \cdot \begin{cases} \min\{h, n_1 - h\}, & h \leq n_1, \\ \min\{h - n_1, n_2 - h\}, & n_1 < h \leq n_2, \\ \min\{h - n_2, n - h\}, & n_2 < h < n. \end{cases} \end{aligned} \quad (36)$$

For $h \in \{n_1, n_2\}$, the previous bound reduces to the trivial inequality $\phi(h) \geq 0$. However, the presence of L links between the two subgraphs ensures that

$$\phi(h) \geq \frac{L\alpha}{n}, \quad \text{if } h \in \{n_1, n_2\}. \quad (37)$$

Combining (36) and (37), using bounds on the harmonic series, and applying Corollary 1, we finally obtain

$$\begin{aligned} \mathbb{E}_0[T] &\leq \frac{1}{2\beta - 1} \left(\frac{2}{L\alpha} n + \frac{12}{c\alpha p} \left(\ln \left(\frac{n}{2} \right) + 1 \right) \right) \\ &\quad + \frac{\beta}{(2\beta - 1)\mathbf{1}^\top \mathbf{u}}. \end{aligned} \quad (38)$$

This, together with (35), demonstrates that the expected spreading time grows linearly with the network size n . In Figure 5 we show Monte Carlo estimations of the expected spreading time for a SBM (in Figure 3(b)), together with our analytical bounds.

Note that, if we allow the control nodes in both the constituent subgraphs of the SBM, we instead obtain a logarithmic growth lower bound on $\mathbb{E}_0[T]$. These considerations extend to more general SBMs composed of many dense subgraphs. They lead to the conclusion that to obtain fast spreading time (of logarithmic growth) in these graphs using constant control policies or, more generally, control policies with constant support \mathcal{U} , the set \mathcal{U} must necessarily have non-empty intersection with all the communities in the SBM. This is a drawback in




Fig. 5: Monte Carlo estimation (200 simulations) of the expected spreading time $\mathbb{E}_0[T]$ on SBMs with $\mathbf{u} = \delta^{(1)}$, $\alpha = 1$, $\beta = 0.8$, $c = 0.4$ and $L = 5$ for different values of n , with 90% confidence intervals. The two solid curves are theoretical bounds from (35) and (38).

practical applications as, not only it may require the cardinality of \mathcal{U} to be large, but also needs precise a-priori information on the network topology to suitably position the control nodes.

C. Ring graphs

Our last case study concerns ring graphs that represent an extreme scenario of poorly connected networks. We fix an undirected ring \mathcal{G} with n nodes and each link weighted $w = \alpha/2$, as represented in Figure 3(c). and we consider a controlled evolutionary dynamics $(\mathcal{G}, \beta, U(t))$ with $X(0) = \mathbf{0}$.

We bound the performance comparing our process $X(t)$ with the Markov process $Y(t)$ associated with the controlled evolutionary dynamics $(\mathcal{G}, 1, U(t))$ with $Y(0) = \mathbf{0}$. Notice that Lemma 2 ensures that the expected spreading time verifies $\mathbb{E}_0[T_X] \geq \mathbb{E}_0[J_Y]$ and the expected control cost verifies $\mathbb{E}_0[J_X] \geq \mathbb{E}_0[J_Y]$, where the subscripts denote the process.

Our aim is to bound the performance of $Y(t)$ applying Corollary 2. To this aim, we start with two simple remarks. First, for every $\mathcal{W} \subseteq \mathcal{V}$, we have $\zeta[\mathcal{W}] \leq \alpha|\mathcal{W}|$. Second, the contagion mechanism cannot increase the boundary $B(t)$: it decreases by α , if the two neighbors of the node that changes its state are both state-1 nodes, or, otherwise, it remains the same. Moreover, considering that every jump due to exerting a control action can increase the boundary by α , we have, for every h ,

$$B_h \leq \alpha N^c. \quad (39)$$

where N^c is the total number of control activations, formally defined in (31). Consider now the event $E := \{N^c \leq 2\mathbb{E}[N^c]\}$.

From (39) and Markov inequality we obtain

$$\begin{aligned} \mathbb{E} \left[\frac{1 - \mathbb{E}[J_h | \sigma_{h-1}]}{B_{h-1}} \right] &\geq \mathbb{E} \left[\frac{1 - \mathbb{E}[J_h | \sigma_{h-1}]}{\alpha N^c} \mathbb{1}_E \right] \\ &\geq \frac{\mathbb{E}[1 - \mathbb{E}[J_h | \sigma_{h-1}]] \mathbb{P}[E]}{2\alpha \mathbb{E}[N^c]} \\ &\geq \frac{1 - \mathbb{E}[\mathbb{E}[J_h | \sigma_{h-1}]]}{4\alpha \mathbb{E}[N^c]}. \end{aligned}$$

From Corollary 2, using the above computation and (32), we finally obtain

$$\mathbb{E}_0[T_X] \geq \mathbb{E}_0[T_Y] \geq \frac{n - \mathbb{E}_0[J_Y]}{4\mathbb{E}_0[J_Y]} \geq \frac{1}{4\mathbb{E}_0[J_X]} n - \frac{1}{4}.$$

This allows us to conclude that the spread is slow on large-scale ring graphs, unless adopting a control policy whose cost $\mathbb{E}_0[J]$ blows up with the network size.

V. FEEDBACK CONTROL POLICIES

Through the examples discussed in the previous section, we have unveiled different behavior of the controlled evolutionary dynamics, depending on the structure of the underlying network topologies. Specifically, we have characterized *easy-to-control* network topologies (e.g., expander graphs) where simple constant control policies can guarantee fast spread. On the other hand, we have identified *hard-to-control* network topologies (e.g., rings) where any control policy fails to achieve fast spread. There are networks, like the stochastic block models, that belong to neither of these two classes, for which the location (and possibly the number) of nodes where the control is exerted plays a crucial role in determining the performance. In such scenarios, the use of feedback control policies may allow one to achieve fast spread with reasonable control efforts, while simple open-loop control policies fail.

Our proposed feedback policies are based on the observation that, in the presence of an evolutionary advantage for the novel states (i.e., with $\beta > 1/2$), the evolutionary dynamics naturally induces a positive feedback loop enhancing the spreading process. However, the presence of bottlenecks in the graph topology (as in the example of SBMs) may slow down such a diffusion process. The objective of the proposed control policies is thus to enforce such a positive feedback whenever the process enters in a bottleneck. Technically, our policies rely on the following two considerations: (i) it is of no use to exert control on state-1 nodes: this yields $C(t) = \mathbf{1}^\top U(t)$; and (ii) in order to apply Theorem 1, the feedback control law $U(t) = \nu(X(t))$ must enforce a relation like (19). This suggests to consider feedback laws whose instantaneous rate $\nu(X(t))$ only depends on the configuration $X(t)$ through the two aggregate variables $A(t)$ and $B(t)$. Moreover, relation (19) suggests that a valuable control policy should compensate for the boundary when this becomes too small, in order to maintain the overall intensity of the process $B(t) + C(t)$ always sufficiently large.

On the basis of these considerations, we design a feedback control policy that is determined by two ingredients: a *target function* $\iota : \{0, 1\}^n \rightarrow \{1, \dots, n\}$ selecting the node in which the control has to be exerted (as a function of the current configuration); and a *rate function* $\mu : \{0, \dots, n\} \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ selecting the control rate (as a function of the number of 1's in the current configuration, that is, $A(t)$, and of the boundary $B(t)$). We assume the target function to be any function $\iota : \{0, 1\}^n \rightarrow \{1, \dots, n\}$ such that $x_{\iota(x)} = 0$ (i.e., we always choose to control a node that is currently in state 0). The rate function is assumed to have the following form. We fix a parameter $K > 0$ and we put

$$\mu(a, b) = \begin{cases} K - b & \text{if } a < n \text{ and } b < K, \\ 0 & \text{otherwise.} \end{cases} \quad (40)$$

Finally, the feedback control law $\nu(X(t))$ is given by:

$$\nu_i(X(t)) = \begin{cases} \mu(A(t), B(t)) & \text{if } i = \iota(X(t)), \\ 0 & \text{else.} \end{cases} \quad (41)$$

Briefly, in the selected node $i = \iota(X(t))$ we exert a control action with rate μ as in (40), while in all other nodes no control is exerted.

In order to analyze the expected spreading time of this feedback control policy, it is useful to consider the following *floor* version of the conductance profile.

Definition 2: For $K \geq 0$, the K -*floor conductance profile* of an undirected weighted graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ of order $|\mathcal{V}| = n$ is the function $\phi_K : \{1, \dots, n-1\} \rightarrow \mathbb{R}_+$ defined as

$$\phi_K(a) := \max\{\phi(a), K\} = \max\left\{\min_{\mathcal{W} \subset \mathcal{V}, |\mathcal{W}|=a} \zeta(\mathcal{W}), K\right\}.$$

From Theorem 1 we establish the following upper bounds on the expected spreading time and on the expected cost of the feedback control policy in (41).

Corollary 6: Consider the feedback controlled evolutionary dynamics $(\mathcal{G}, \beta, \nu)$ with initial condition $X(0) = \mathbf{0}$, where ν follows the control policy (41). Then, the expected spreading time verifies

$$\mathbb{E}_0[T] \leq \frac{\beta}{(2\beta - 1)K} + \frac{1}{2\beta - 1} \sum_{a=1}^{n-1} \frac{1}{\phi_K(a)}, \quad (42)$$

and the expected control cost is bounded as

$$\mathbb{E}_0[J] \leq \frac{\beta}{2\beta - 1} + \frac{|\{1 \leq a < n : \phi(a) < K\}|}{2\beta - 1}. \quad (43)$$

Proof: From (40) it follows that, for all t such that $A(t) < n$, $C(t) = \max\{K - B(t), 0\}$. For such values of t we thus have

$$B(t) + C(t) = \max\{B(t), K\} \geq \phi_K(A(t)).$$

The upper bound (42) follows from Theorem 1.

The estimation of the expected control cost is performed as follows. Note first that $C(t) \leq K$ for every t . Moreover, if a is such that $\phi(a) > K$, then $B(t) > K$ and $C(t) = 0$. Using the upper bounds (10) on the time spent by the process $A(t)$ in the various states, we conclude

$$\begin{aligned} \mathbb{E}_0[J] &\leq K \left(\mathbb{E}[T_0 | X(0) = \mathbf{0}] \right. \\ &\quad \left. + \sum_{a=1}^{n-1} \mathbb{E}[T_a | X(0) = \mathbf{0}] \mathbb{1}_{[0, K]}(\phi(a)) \right) \\ &= K \left(\frac{\beta}{2\beta - 1} \frac{1}{\phi_K(0)} + \frac{1}{2\beta - 1} \sum_{a=1}^{n-1} \frac{\mathbb{1}_{[0, K]}(\phi(a))}{\phi_K(a)} \right) \\ &= \frac{\beta}{2\beta - 1} + \frac{|\{1 \leq a < n : \phi(a) < K\}|}{2\beta - 1}. \end{aligned}$$

We observe that the two bounds in Corollary 6 depend on the control policy ν through the parameter K and on the network structure through the minimum conductance profile ϕ . Due to the monotonicity of ϕ_K , the upper bound in (42) is nonincreasing in K , while (43) is nondecreasing in K . This establishes a trade-off behavior between faster spread and higher cost (see, e.g., Figure 7), paving the way for the formalization of an optimization problem to choose the value K that best compromises fast spread (larger values of K) and affordable cost (lower values of K).




Fig. 6: Monte Carlo estimation (200 simulations) and 90% confidence intervals of the expected spreading time $\mathbb{E}_0[T]$ on SBMs for different n with $\alpha = 1$, $\beta = 0.8$, $c = 0.4$, and $L = 5$, under the feedback control policy in (41) with $K = 1/4$ (red squares) and constant control (blue circles), for two different values of p . The red solid curves are a numerical estimation of (29) and the theoretical bound for the feedback control from (44). The blue solid curves are theoretical bounds for the constant control from (35) and (38).

A. Fast spread on SBMs

We consider the feedback control policy in (41) on the SBMs introduced in Section IV-B, choosing $K < cap/2$. From (36), we have that

$$\phi(a) \geq K \quad \forall a \in \{1, \dots, n-1\} \setminus \{n_1, n_2\},$$

with high probability as n grows large. Therefore,

$$\begin{aligned} \phi_K(a) &= \phi(a) && \text{for } a \in \{1, \dots, n-1\} \setminus \{n_1, n_2\}, \\ \phi_K(a) &\geq K && \text{for } a \in \{n_1, n_2\}. \end{aligned}$$

The application of (42) yields the following bound on the expected spreading time

$$\mathbb{E}_0[T] \leq \frac{\beta}{(2\beta-1)K} + \frac{1}{2\beta-1} \left(\frac{2}{K} + \frac{12}{cap} (\ln(n/2) + 1) \right), \quad (44)$$

where the algebraic computations follow the ones carried on in Section IV-B. Using (43), we finally bound the expected control cost as

$$\mathbb{E}_0[J] \leq \frac{2+\beta}{2\beta-1}.$$

Hence, with a bounded control cost, we achieve an expected spreading time that grows logarithmically in n .

Figure 6 compares our feedback control policy with constant control policies, highlighting the improvement in performance of the former with respect to the latter. We also notice that, for highly connected communities (i.e., for large values of p), such an improvement increases in magnitude.

In Figure 7, we analyze the trade-off between spreading time and control cost depending on the parameter K . There is a first phase, for $K \leq cap/2$ (light blue), where, as K increases, the average spreading time shows a fast decrease, while the average control cost increases slowly. Then, for $K > cap/2$ (dark blue), the average spreading time decreases slowly against a strong growth of the control cost. In the same figure, we compare our control policy with a simpler feedback control policy that does not depend on $B(t)$, obtained using the same target function $\iota(X(t))$ but a non-negative constant




Fig. 7: Monte Carlo estimation (200 simulations) of the expected spreading time $\mathbb{E}_0[T]$ and expected cost $\mathbb{E}_0[J]$ on SBMs with $n = 800$, $\alpha = 1$, $\beta = 0.8$, $c = 0.4$, $p = 0.1$ and $L = 5$, using the feedback control policy in (41), for different values of the parameter K . Green squares are Monte Carlo estimation of the expected spreading time $\mathbb{E}_0[T]$ and expected cost $\mathbb{E}_0[J]$ under a control policy with the same position, but constant rate.

rate (different squares in green correspond to different values). This comparison supports our intuition that the use of $B(t)$ in the feedback law is key to improve in the performance of the spreading process.

Summing up, despite (i) the myopic choice of our control policy in which no optimization is done on the target node, (ii) the limited a-priori knowledge on the topology (only knowledge of ϕ is used), and (iii) the use of only two one-dimensional statistics (from an n -dimensional state variable), we have been able to design a feedback control policy whose performance is guaranteed by analytical results and whose effectiveness (already numerically tested on a case study in [25]) has been explicitly demonstrated on SBMs. We remark that the control strategy is designed without knowledge of the exact partition of nodes into communities, and even of their size. This property provides robustness in real-world situations, where limited data might be available leading to uncertainty on the exact network structure.

VI. DISCUSSION AND CONCLUSION

We have proposed a continuous-time stochastic dynamical model that allows for an analytical treatment of evolutionary dynamics and incorporates exogenous control input mechanisms. The main contributions of this paper are i) the formulation of a link-based spreading process incorporating an exogenous control and leading to a non-homogenous Markov process; ii) a rigorous analysis of the transient spreading process in terms of the network topology and of the control policy adopted; iii) the derivation of a set of ready-to-use corollaries to estimate the spreading time and the control cost for specific control policies; and iv) the design and the analysis of an effective feedback control policy, which relies on few topological data and one-dimensional variables of the system. The proposed policy provides a dramatic improvement in the speed of the evolutionary process for some network structures where simpler control policies fail to achieve fast spread.

As a consequence of our analysis, we have characterized three classes of network topologies depending on their controllability.

- a) **Topologies easy to control even with constant control policies.** Among them we mention expander graphs, which is a large family of highly connected networks including the complete graph, Erdős-Rényi random graphs, small-world networks, as well many scale-free networks such as those generated according to a preferential attachment algorithm. For these topologies, little effort is required to achieve fast spread, that is, to guarantee that the expected spreading time grows logarithmically with the network size n for every constant control policy.
- b) **Topologies easy to control only with feedback control policies.** A representative example is the stochastic block model (SBM), of which barbell graphs are well known limit cases. For these structures, no fundamental limit precludes fast spread. However, we have shown that constant control policies fail in achieving it. A major contribution of this paper then consisted in the development of a feedback control policy that remarkably improves the speed of the process, achieving fast spread.
- c) **Topologies hard to control under any control policy,** of which the ring graph is a representative element. For these topologies, we observe slow spread under any feasible control policy, where the expected spreading time shows a linear growth in the network size.

We believe that the generality of Theorem 1 and the effectiveness of the feedback control policy proposed in Section V pave the way for further research seeking for an optimal control policy for evolutionary dynamics, also extending the real-world case study in [25], in which evolutionary dynamics are proposed to model the introduction of genetically modified mosquitoes in a geographic region to substitute the species that transmit diseases [40]. The proposed feedback control policy requires limited a-priori knowledge of the network topology. When more information on the network structure is available, we believe that such information might be exploited to design and study targeted introduction policies. For instance, inspired by targeted vaccination strategies [7], the introduction of the novel states might be prioritized in nodes with high network centrality. Other avenues of future research include the use of the technical tools developed in this paper to study control policies on other dynamical processes on networks, such as opinion dynamics, diffusion of information, and epidemics.

APPENDIX

A. Proof of Lemma 1

For a given time $t_0 \in \mathbb{R}_+$ and let X_k, A_k, B_k, C_k be the discrete-time jump processes of $X(t), A(t), B(t),$ and $C(t)$, respectively, starting from time t_0^+ and onwards, with $X_0 = X(t_0) = \mathbf{x}$. Clearly,

$$\mathbb{P}[A(t) \geq a, t \geq t_0 | X(t_0) = \mathbf{x}] = \mathbb{P}[A_k \geq a, k \geq 0 | X_0 = \mathbf{x}] \quad (45)$$

It follows from (7)–(8) that the increase and decrease transition probabilities of A_k conditioned to the process X_k at time k , also satisfy

$$p_k^-(a|\mathbf{x}) \leq 1 - \beta, \quad p_k^+(a|\mathbf{x}) \geq \beta, \quad \text{for } a \notin \{0, n\}, \quad (46)$$

while $p_k^+(0|0) = 1$ for all k , and $a = n$ is an absorbing state.

Consider now a discrete-time birth-and-death chain \tilde{A}_k with state space $\{0, \dots, n\}$ and transition probabilities

$$\begin{cases} p^+(0) = 1, \\ p^+(a) = \beta, a \notin \{0, n\}, \\ p^+(n) = 0; \end{cases} \quad \begin{cases} p^-(0) = 0, \\ p^-(a) = 1 - \beta, a \notin \{0, n\}, \\ p^-(n) = 0, \end{cases}$$

with $\tilde{A}_0 = a$. A standard argument allows us to couple the two processes A_k and \tilde{A}_k in such a way that $A_k \geq \tilde{A}_k$ for every k , yielding

$$\mathbb{P}[A_k \geq a, k \geq 0 | X_0 = \mathbf{x}] \geq \mathbb{P}[\tilde{A}_k \geq a, k \geq 0]. \quad (47)$$

On the other hand, a direct computation for the birth-and-death chain \tilde{A}_k implies that

$$\mathbb{P}[\tilde{A}_k \geq a, \forall k] = \frac{1 - \left(\frac{1-\beta}{\beta}\right)}{1 - \left(\frac{1-\beta}{\beta}\right)^{n-a+1}} \geq \frac{2\beta - 1}{\beta}. \quad (48)$$

Considering that t_0 can be chosen arbitrarily, (45), (47), and (48) prove Lemma 1. ■

B. Proof of Lemma 2

First, we consider the case $\beta = \gamma$ and $\mathbf{x}_0 \leq \mathbf{y}_0$. We define the coupled process $Z(t) = (X(t), Y(t))$ on the state space $\{0, 1\}^n \times \{0, 1\}^n$, with initial condition $Z(0) = (\mathbf{x}_0, \mathbf{y}_0)$, associated with the same graph \mathcal{G} . The coupling mechanism is the following. Each link $\{i, j\}$ is equipped with an independent rate- W_{ij} Poisson clock. When the clock associated with link $\{i, j\}$ ticks, the spreading mechanism acts on that link for both $X(t)$ and $Y(t)$ as for a standard controlled evolutionary dynamics and, if a conflict occurs in both processes, then the outcome is the same. Each node i is equipped with a non-homogeneous Poisson clock with rate $U_i(t)$ associated with the external control in node i . When the clock associated with node i ticks, then both X_i and Y_i turn to 1. Each of the two marginals of $X(t)$ and $Y(t)$ coincides with the distribution of a controlled evolutionary dynamics $(\mathcal{G}, \beta, U(t))$ with initial condition $X(0) = \mathbf{x}_0$ and $Y(0) = \mathbf{y}_0$, respectively.

We now show that, under this coupling, $Y(t) \geq X(t)$, for every $t \geq 0$. At $t = 0$, this is verified by assumption. In the following, we prove that any transition of $Z(t)$ keeps the inequality preserved. We assume that a transition of the process occurs at time t . If $X(t^-) = Y(t^-)$, then, due to the coupling mechanism, $X(t^+) = Y(t^+)$. If $Y(t^-) > X(t^-)$, we analyze all the possible events which trigger a transition.

- *Spreading mechanism.* When link $\{i, j\}$ activates and a conflict occurs:

- if $X_i(t^-) = Y_i(t^-) = X_j(t^-) = 0, Y_j(t^-) = 1$ and state 1 wins, then $X_i(t^+) = X_j(t^+) = 0$ and $Y_i(t^+) = Y_j(t^+) = 1$, so that $Y(t^+) > X(t^+)$;
- if $X_i(t^-) = Y_i(t^-) = X_j(t^-) = 0, Y_j(t^-) = 1$ and state 0 wins, then $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 0$, so that $Y(t^+) \geq X(t^+)$;
- if $X_i(t^-) = Y_i(t^-) = 0, X_j(t^-) = Y_j(t^-) = 1$ and state 1 wins, then, $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 1$, so that $Y(t^+) > X(t^+)$;

- (d) if $X_i(t^-) = Y_i(t^-) = 0$, $X_j(t^-) = Y_j(t^-) = 1$, and state 0 wins, then, $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 0$, so that $Y(t^+) > X(t^+)$;
- (e) if $X_i(t^-) = 0$, $Y_i(t^-) = X_j(t^-) = Y_j(t^-) = 1$, and state 1 wins, then, $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 1$, so that $Y(t^+) \geq X(t^+)$;
- (f) if $X_i(t^-) = 0$, $Y_i(t^-) = X_j(t^-) = Y_j(t^-) = 1$, and state 0 wins, then, $X_i(t^+) = X_j(t^+) = 0$ and $Y_i(t^+) = Y_j(t^+) = 1$, so that $Y(t^+) > X(t^+)$.

• *External control.* Node i activates and it has state 0 at least in one of the two processes:

- 1) if $X_i(t^-) = Y_i(t^-) = 0$, then, $X_i(t^+) = Y_i(t^+) = 1$, so that $Y(t^+) > X(t^+)$;
- 2) if $X_i(t^-) = 0$, $Y_i(t^-) = 1$, then, $X_i(t^+) = Y_i(t^+) = 1$, so that $Y(t^+) \geq X(t^+)$.

Hence, after each transition of the process, the inequality $Y(t) \geq X(t)$ is preserved.

The proof for $\beta < \gamma$ and $\mathbf{x}_0 = \mathbf{y}_0$ follows a similar argument. We define the coupled process $Z(t) = (X(t), Y(t))$ in which each link $\{i, j\}$ is equipped with an independent Poisson clock with rate W_{ij} . When the clock associated with link $\{i, j\}$ ticks, the spreading mechanism acts on that link for both $X(t)$ and $Y(t)$. If a conflict occurs in only one of the two processes, then it is solved as in a standard controlled evolutionary dynamics with probability for the novel state to win the conflict equal to β for $X(t)$ and γ for $Y(t)$, respectively. If the conflict occurs in both processes, then with probability β the novel state wins in both $X(t)$ and $Y(t)$, with probability $\gamma - \beta$ it wins only in $Y(t)$, and with probability $1 - \gamma$ the novel state loses in both $X(t)$ and $Y(t)$. Each node i is given a non-homogeneous Poisson clock with rate $U_i(t)$. When the clock associated with node i ticks, then both X_i and Y_i turn to 1. We immediately deduce that the two marginals $X(t)$ and $Y(t)$ are controlled evolutionary dynamics $(\mathcal{G}, \beta, U(t))$ and $(\mathcal{G}, \gamma, U(t))$, respectively, with the same initial condition $X(0) = Y(0) = \mathbf{x}_0 = \mathbf{y}_0$.

Under this coupling, the inequality $Y(t) \geq X(t)$ holds true for every $t \geq 0$. In fact, at $t = 0$ it is verified, since $X(0) = Y(0)$. Then, the analysis of all possible transitions and their effect on the inequality is performed similar to above and is omitted due to space constraints. Finally, the case $\beta < \gamma$ and $\mathbf{x}_0 < \mathbf{y}_0$ is obtained by combining the two couplings above.

The coupling $Z(t) = (X(t), Y(t))$ such that $Y(t) \geq X(t)$ for every $t \geq 0$, proves the stochastic domination $Y(t) \succeq X(t)$ [3], yielding $\mathbb{E}_{\mathbf{y}_0}[T_Y] \leq \mathbb{E}_{\mathbf{x}_0}[T_X]$. Since $U(t) \geq 0$, we have

$$\begin{aligned} \mathbb{E}_{\mathbf{x}_0}[J_X] &= \mathbb{E} \left[\int_0^{T_X} U(t) dt \right] = \int_0^{\mathbb{E}_{\mathbf{x}_0}[T_X]} U(t) dt \\ &\geq \int_0^{\mathbb{E}_{\mathbf{y}_0}[T_Y]} U(t) dt = \mathbb{E}_{\mathbf{y}_0}[J_X], \end{aligned}$$

which completes the proof. ■

C. Proof of Lemma 3

We define a coupled process $Z(t) = (X(t), Y(t))$ on the state space $\{0, 1\}^n \times \{0, 1\}^n$, with initial condition $Z(0) =$

$(\mathbf{x}_0, \mathbf{y}_0)$. Here, the marginal distributions of $X(t)$ and $Y(t)$ are controlled evolutionary dynamics $(\mathcal{G}, 1, U(t))$ and $(\mathcal{G}, 1, 0)$ associated with a the same graph \mathcal{G} . The coupling mechanism is the following. Each link $\{i, j\}$ of the graph \mathcal{G} is equipped with an independent rate- W_{ij} Poisson clock. When the clock associated with link $\{i, j\}$ ticks, the spreading mechanism acts on that link for both $X(t)$ and $Y(t)$, with $\beta = 1$. Each node i is given a non-homogeneous Poisson clock with rate $U_i(t)$, associated with the external control in node i . When the clock associated with node i ticks, the state $X_i(t)$ turns to 1. We immediately deduce that the two marginals of $X(t)$ and $Y(t)$ are controlled evolutionary dynamics $(\mathcal{G}, 1, U(t))$ and $(\mathcal{G}, 1, 0)$ with the desired initial conditions, respectively.

We show now that, under this coupling, $Y(t) \geq X(t)$, for every t . At $t = 0$ this is verified by assumption. In the following, we show that, after each transition of the coupled process, the inequality is preserved.

- *Spreading mechanism.* Since $\beta = 1$, the novel state always wins. So only three transitions can occur when link $\{i, j\}$ activates and a conflict occurs:
 - (a) If $X_i(t^-) = Y_i(t^-) = X_j(t^-) = 0$ and $Y_j(t^-) = 1$, then $X_i(t^+) = X_j(t^+) = 0$, $Y_i(t^+) = Y_j(t^+) = 1$, which implies that $Y(t^+) > X(t^+)$.
 - (b) If $X_i(t^-) = Y_i(t^-) = 0$ and $X_j(t^-) = Y_j(t^-) = 1$, then $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 1$, which implies that $X(t^+) \geq Y(t^+)$.
 - (c) If $X_i(t^-) = 0$ and $Y_i(t^-) = Y_j(t^-) = X_j(t^-) = 1$, then $Y_i(t^+) = Y_j(t^+) = X_i(t^+) = X_j(t^+) = 1$, which implies that $Y(t^+) \geq X(t^+)$.
- *External control.* We observe that (i) we can only control nodes for which $Y_i(0) = 1$, and (ii) since $\beta = 1$, if $Y_i(0) = 1$ then $Y_i(t) = 1$ for every $t \geq 0$. Hence, external control preserves the ordering $Y(t) \geq X(t)$.

The stochastic domination $Y(t) \succeq X(t)$ yields the claim. ■

REFERENCES

- [1] L. Zino, G. Como, and F. Fagnani, "Fast Diffusion of a Mutant in Controlled Evolutionary Dynamics," in *Proc. 20th World Congr. Int. Fed. Autom. Control (IFAC)*, July 2017.
- [2] A. Ganesh, L. Massoulié, and D. Towsley, "The effect of network topology on the spread of epidemics," *Proc. IEEE Int. Conf. Comp. Comm. (INFOCOM)*, vol. 2, pp. 1455–1466, Mar. 2005.
- [3] M. Draief, "Epidemic processes on complex networks: The effect of topology on the spread of epidemics," *Physica A*, vol. 363, no. 1, pp. 120–131, 2006.
- [4] R. Pastor-Satorras, C. Castellano, P. Van Mieghem, and A. Vespignani, "Epidemic processes in complex networks," *Rev. Mod. Phys.*, vol. 87, no. 3, pp. 925–979, 2015.
- [5] W. Mei, S. Mohagheghi, S. Zampieri, and F. Bullo, "On the dynamics of deterministic epidemic propagation over networks," *Annu. Rev. Control*, pp. 116–128, 2017.
- [6] F. Fagnani and L. Zino, "Time to extinction for the SIS epidemic model: New bounds on the tail probabilities," *IEEE Trans. Netw. Sci. Eng.*, vol. 6, no. 1, pp. 74–81, 2019.
- [7] F. Chung, P. Horn, and A. Tsiatas, "Distributing Antidote Using PageRank Vectors," *Internet Math.*, vol. 6, no. 2, pp. 237–254, 2009.
- [8] C. Borgs, J. Chayes, A. Ganesh, and A. Saberi, "How to distribute antidote to control epidemics," *Random Struct. Algor.*, vol. 37, no. 2, pp. 204–222, 2010.
- [9] K. Drakopoulos, A. Ozdaglar, and J. N. Tsitsiklis, "An efficient curing policy for epidemics on graphs," *IEEE Trans. Netw. Sci. Eng.*, vol. 1, no. 2, pp. 67–75, 2014.

- [10] S. Trajanovski, Y. Hayel, E. Altman, H. Wang, and P. Van Mieghem, "Decentralized Protection Strategies Against SIS Epidemics in Networks," *IEEE Trans. Control Netw. Syst.*, vol. 2, no. 4, pp. 406–419, 2015.
- [11] C. Nowzari, V. M. Preciado, and G. J. Pappas, "Analysis and control of epidemics: a survey of spreading processes on complex networks," *IEEE Control Syst. Lett.*, vol. 36, no. 1, pp. 26–46, 2016.
- [12] T. M. Liggett, *Interacting particle systems*. New York, NY, USA: Springer-Verlag, 1985.
- [13] L. Frachebourg and P. L. Krapivsky, "Exact results for kinetics of catalytic reactions," *Phys. Rev. E*, vol. 53, no. 4, p. R3009, 1996.
- [14] A. Montanari and A. Saberi, "The spread of innovations in social networks," *Proc. Nat. Acad. Sci. US*, vol. 107, no. 47, pp. 20 196–20 201, 2010.
- [15] H. P. Young, "The dynamics of social innovation," *Proc. Nat. Acad. Sci. US*, vol. 108, no. 4, pp. 21 285–21 291, 2011.
- [16] W. Rossi, G. Como, and F. Fagnani, "Threshold models of cascades in large-scale networks," *IEEE Trans. Netw. Sci. Eng.*, vol. 6, no. 2, pp. 158–172, 2019.
- [17] D. Kempe, J. Kleinberg, and É. Tardos, "Maximizing the spread of influence through a social network," in *Proc. 9th ACM SIGKDD Int. Conf. Know. Discov. Data Min. (KDD)*, Aug. 2003, pp. 137–146.
- [18] S. Durand, G. Como, and F. Fagnani, "Controlling network coordination games," in *Proc. 21th World Congr. Int. Fed. Autom. Control (IFAC)*, 2020.
- [19] J. R. Riehl and M. Cao, "Towards optimal control of evolutionary games on networks," *IEEE Trans. Autom. Control*, vol. 62, no. 1, pp. 458–462, 2017.
- [20] E. Lieberman, C. Hauert, and M. A. Nowak, "Evolutionary dynamics on graphs," *Nature*, vol. 433, 2005.
- [21] J. Rychtář and B. Stadler, "Evolutionary dynamics on small-world networks," *Int. J. Comp. Math. Sci.*, vol. 2, no. 1, pp. 1–4, 2008.
- [22] H. Ohtsuki and M. A. Nowak, "Evolutionary games on cycles," *Proc. R. Soc. B*, vol. 273, no. 1598, pp. 2249–2256, 2006.
- [23] M. Broom, J. Rychtář, and B. T. Stadler, "Evolutionary dynamics on graphs - the effect of graph structure and initial placement on mutant spread," *J. Stat. Theory Pract.*, vol. 5, no. 3, pp. 369–381, 2011.
- [24] B. Allen, G. Lippner, Y.-T. Chen, B. Fotouhi, N. Momeni, S.-T. Yau, and M. A. Nowak, "Evolutionary dynamics on any population structure," *Nature*, vol. 544, no. 7649, pp. 227–230, 2017.
- [25] L. Zino, G. Como, and F. Fagnani, "Controlling Evolutionary Dynamics in Networks: A Case Study," in *7th IFAC Workshop Distrib. Estim. Control Netw. Syst. (NecSys)*, Aug. 2018.
- [26] M. Mihail, "Conductance and convergence of markov chains—a combinatorial treatment of expanders," in *Proc. 30th IEEE Annu. Symp. Found. Comput. Sci. (FOCS)*, Oct 1989, pp. 526–531.
- [27] G. Roberts and J. Rosenthal, "Quantitative bounds for convergence rates of continuous time markov processes," *Electron. J. Probab.*, vol. 1, 1996.
- [28] D. J. Watts and S. H. Strogatz, "Collective dynamics of 'small-world' networks," *Nature*, vol. 393, no. 6684, pp. 440–442, 1998.
- [29] R. Albert and A.-L. Barabási, "Statistical mechanics of complex networks," *Rev. Mod. Phys.*, vol. 74, pp. 47–97, 2002.
- [30] P. W. Holland, K. B. Laskey, and S. Leinhardt, "Stochastic blockmodels: First steps," *Soc. Netw.*, vol. 5, no. 2, pp. 109–137, 1983.
- [31] S. Fortunato and D. Hric, "Community detection in networks: A user guide," *Phys. Rep.*, vol. 659, pp. 1–44, 2016.
- [32] D. A. Levin, Y. Peres, and E. L. Wilmer, *Markov chains and mixing times*. Providence, RI, USA: American Mathematical Society, 2009.
- [33] I. Ferreira, "The probability of survival for the biased voter model in a random environment," *Stoch. Process. Their Appl.*, vol. 34, no. 1, pp. 25–38, 1990.
- [34] D. Acemoğlu, G. Como, F. Fagnani, and A. Ozdaglar, "Opinion fluctuations and disagreement in social networks," *Math. Oper. Res.*, vol. 38, no. 1, pp. 1–27, 2013.
- [35] G. Como and F. Fagnani, "From local averaging to emergent global behaviors: The fundamental role of network interconnections," *Syst. Control Lett.*, vol. 95, pp. 70–76, 2016.
- [36] A. V. Proskurnikov and R. Tempo, "A tutorial on modeling and analysis of dynamic social networks. Part I," *Annu. Rev. Control*, vol. 43, pp. 65–79, 2017.
- [37] S. Goel, R. Montenegro, and P. Tetali, "Mixing time bounds via the spectral profile," *Electron. J. Probab.*, vol. 11, pp. 1–26, 2006.
- [38] M. Barthélemy, "Spatial networks," *Phys. Rep.*, vol. 499, no. 1-3, pp. 1–101, 2011.
- [39] B. Bollobas, *Random Graphs*. Springer-Verlag, 2001.
- [40] A. Buchman, S. Gamez, M. Li, I. Antoshechkin, H.-H. Li, H.-W. Wang, C.-H. Chen, M. J. Klein, J.-B. Duchemin, J. E. Crowe, Jr., P. N. Paradkar, and O. S. Akbari, "Broad dengue neutralization in mosquitoes expressing an engineered antibody," *PLOS Pathog.*, vol. 16, no. 1, pp. 1–20, 2020.



Lorenzo Zino is a PostDoc Researcher at the Faculty of Science and Engineering, University of Groningen, Groningen, The Netherlands, since 2019. He received the Laurea degree (B.Sc.) in Applied Mathematics from Politecnico di Torino, Torino, Italy, in 2012, the Laurea Magistrale degree (M.S.) in Mathematical Engineering from Politecnico di Torino, in 2014, and the Ph.D. in Pure and Applied Mathematics from Politecnico di Torino and Università di Torino (joint doctorate program), in 2018. He was a Research Fellow at Politecnico di Torino in 2018–19 and a Visiting Research Assistant at New York University Tandon School of Engineering in 2017–18 and in 2019. His current research interests include modeling, analysis, and control of dynamical processes over networks, applied probability, network analysis, and game theory.



Giacomo Como is an Associate Professor at the Department of Mathematical Sciences, Politecnico di Torino, Italy, and at the Automatic Control Department of Lund University, Sweden. He received the B.Sc., M.S., and Ph.D. degrees in Applied Mathematics from Politecnico di Torino, in 2002, 2004, and 2008, respectively. He was a Visiting Assistant in Research at Yale University in 2006–2007 and a Postdoctoral Associate at the Laboratory for Information and Decision Systems, Massachusetts Institute of Technology, from 2008 to 2011. He currently serves as Associate Editor of the *IEEE Transactions on Network Science and Engineering* and of the *IEEE Transactions on Control of Network Systems* and as chair of the IEEE-CSS Technical Committee on Networks and Communications. He was the IPC chair of the IFAC Workshop NecSys'15 and a semiplenary speaker at the International Symposium MTNS'16. He is recipient of the 2015 George S. Axelby Outstanding Paper Award. His research interests are in dynamics, information, and control in network systems with applications to cyber-physical systems, infrastructure networks, and social and economic networks



Fabio Fagnani received the Laurea degree in Mathematics from the University of Pisa and the Scuola Normale Superiore, Pisa, Italy, in 1986. He received the PhD degree in Mathematics from the University of Groningen, Groningen, The Netherlands, in 1991. From 1991 to 1998, he was an Assistant Professor of Mathematical Analysis at the Scuola Normale Superiore. In 1997, he was a Visiting Professor at the Massachusetts Institute of Technology (MIT), Cambridge, MA. Since 1998, he has been with the Politecnico of Torino, where he is currently (since 2002) a Full Professor of Mathematical Analysis. From 2006 to 2012, he has acted as Coordinator of the PhD program in Mathematics for Engineering Sciences at Politecnico di Torino. From June 2012 to September 2019, he served as the Head of the Department of Mathematical Sciences, Politecnico di Torino. His current research topics are on cooperative algorithms and dynamical systems over graphs, inferential distributed algorithms, and opinion dynamics. He is an Associate Editor of the *IEEE Transactions on Automatic Control* and served in the same role for the *IEEE Transactions on Network Science and Engineering* and of the *IEEE Transactions on Control of Network Systems*.