

# SOBOLEV ANISOTROPIC INEQUALITIES WITH MONOMIAL WEIGHTS

F. FEO, J. MARTIN, AND M. R. POSTERARO

ABSTRACT. We derive some anisotropic Sobolev inequalities in  $\mathbb{R}^n$  with a monomial weight in the general setting of rearrangement invariant spaces. Our starting point is to obtain an integral oscillation inequality in multiplicative form.

## 1. INTRODUCTION

The study of functional and geometric inequalities with monomial weights, i.e. weights defined by

$$(1.1) \quad d\mu(x) := x^A dx = |x_1|^{A_1} \cdots |x_n|^{A_n} dx.$$

where  $A = (A_1, A_2, \dots, A_n)$  is a vector in  $\mathbb{R}^n$  with  $A_i \geq 0$  for  $i = 1, \dots, n$ , have been considered extensively recently (see for example [11], [12], [3], [9], [30] and the references quoted therein). The interest for this kind of problems appears when Cabré and Ros-Oton (motivated by an open question raised by Haim Brezis [6],[7]) studied in [13] the problem of the regularity of stable solutions to reaction-diffusion problems of double revolution in  $\mathbb{R}^2$ . A function  $u$  has symmetry of double revolution if  $u(x, y) = u(|x|, |y|)$ , with  $(x, y) \in \mathbb{R}^D = \mathbb{R}^{A_1+1} \times \mathbb{R}^{A_2+1}$  ( $A_i$  are positive integers), i.e. the function  $u$  can be seen as a suitable function in  $\mathbb{R}^2$ , and it is here where the Jacobian  $|x_1|^{A_1}|x_2|^{A_2}$  appears (see [13] for the details). In [12], the authors established a sharp isoperimetric inequality in  $(\mathbb{R}^n, \mu)$  (see also [9]) which allows them to obtain the following weighted Sobolev inequality.

**Theorem 1.1.** ([12, Theorem 1.3]) *Let  $\mu$  be defined in (1.1), let*

$$(1.2) \quad D = n + A_1 + \cdots + A_n$$

and  $1 \leq p < D$ . Then for any  $f \in C_c^1(\mathbb{R}^n)$  we have<sup>1</sup>

$$(1.3) \quad \left( \int_{\mathbb{R}^n} |f|^{p^*} d\mu \right)^{1/p^*} \leq C \left( \int_{\mathbb{R}^n} |\nabla f|^p d\mu \right)^{1/p}$$

for some positive constant  $C$ , where

$$(1.4) \quad p^* = \frac{Dp}{D-p}.$$

As in the unweighted case a scaling argument shows that the exponent  $p^*$  is optimal, in the sense that (1.4) can not hold with any other exponent. Moreover the exponent  $p^*$  is exactly the same as in the classical Sobolev inequality, but in

---

2000 *Mathematics Subject Classification.* Primary: 46E30, 26D10.

*Key words and phrases.* anisotropic inequalities, rearrangement invariant space, Sobolev embedding.

<sup>1</sup> $C_c^1(\mathbb{R}^n)$  denotes the space of  $C^1$  functions with compact support in  $\mathbb{R}^n$ .

this case the "dimension" is given by  $D$  (instead of  $n$ ). If  $A_1 = \dots = A_n = 0$ , then exponent  $p^*$  and inequality (1.3) are exactly the classical ones.

We observe that when  $p > 1$  and  $A_i < p - 1$  for all  $i = 1, \dots, n$  the weight in (1.1) belongs to the Muckenhoupt class  $A_p$ , i.e. but, in general the monomial weight does not satisfy the Muckenhoupt condition.

The main purpose of this paper is to obtain some anisotropic Sobolev inequalities on  $\mathbb{R}^n$  with monomial weight  $x^A$  in the general setting of rearrangement invariant spaces (e.g.  $L^p$ , Lorentz, Orlicz, Lorentz-Zygmund, etc...). To this end, we will use the "symmetrization by truncation principle", developed by Milman-Martín in [26] (see also [27] and [28]). This method will provide us a family of rearrangement point-wise inequalities between the special difference<sup>2</sup>  $O_\mu(f, t) := f_\mu^{**}(t) - f_\mu^*(t)$  (called the oscillation of  $f$ ) and the product of the rearrangements of the partial derivative of  $f$  (see Theorem 3.1 below) that will be the key to obtain anisotropic inequalities. More precisely we will prove that inequality (1.3) with  $p = 1$  is equivalent to the following oscillation inequality:

$$(1.5) \quad \int_0^t \left( O_\mu(f, \cdot) (\cdot)^{-\frac{1}{D}} \right)^* (s) ds \leq \int_0^t \prod_{i=1}^n \left[ \left( \frac{d}{ds} \int_{\{|f| > f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^* (\tau) \right]^{\frac{A_i+1}{D}} d\tau$$

for every  $t > 0$ . The rearrangements without subscript  $\mu$  are rearrangement with respect to Lebesgue measure on  $(0, \infty)$ ,  $f_{x_i} = \frac{\partial f}{\partial x_i}$  and symbol  $f \leq g$  means that there exists a universal constant  $c$  (independent of all parameters involved) such that  $f \leq cg$ .

Inequality (1.5) contains the basic information to obtain anisotropic Sobolev inequalities on rearrangement invariant spaces, since given a rearrangement invariant space  $X$  on  $(\mathbb{R}^n, \mu)$ , Hardy's inequality (see (2.3) below) implies<sup>3</sup>

$$(1.6) \quad \left\| O_\mu(f, t) t^{-\frac{1}{D}} \right\|_{\bar{X}} \leq \left\| \prod_{i=1}^n \left[ \left( \frac{d}{ds} \int_{\{|f| > f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^* (\tau) \right]^{\frac{A_i+1}{D}} (t) \right\|_{\bar{X}}.$$

For example, given  $p_1, \dots, p_n \geq 1$ , let  $\bar{p}$  be the weighted harmonic mean between  $p_1, \dots, p_n$ , i.e.

$$(1.7) \quad \frac{1}{\bar{p}} = \frac{1}{D} \sum_{i=1}^n \frac{A_i + 1}{p_i},$$

then (1.6) implies (see Theorem 4.3 below)

$$(1.8) \quad \left\| O_\mu(f, t) t^{-\frac{1}{D}} \right\|_{\bar{X}(\bar{p})} \leq \prod_{i=1}^n \left\| f_{x_i} \right\|_{X^{(p_i)}(\frac{A_i+1}{D})},$$

where  $X^{(p)} = \{f : |f|^p \in X\}$  endowed with the norm  $\|f\|_{X^{(p)}} = \| |f|^p \|_X^{1/p}$ .

In the particular case that  $X = L^1$  and  $\bar{p} < D$ , then (1.8) becomes (see Proposition 4.5 below)

$$\|f\|_{L_\mu^{\bar{p}^*}} \leq \prod_{i=1}^n \|f_{x_i}\|_{L_\mu^{p_i}(\frac{A_i+1}{D})} \quad \forall f \in C_c^1(\mathbb{R}^n).$$

<sup>2</sup> $f_\mu^*$  is the decreasing rearrangement of  $f$  with respect the measure  $\mu$ , and  $f_\mu^{**}(t) = \frac{1}{t} \int_0^t f_\mu^*(s) ds$  (see Subsection 2.1).

<sup>3</sup>The spaces  $\bar{X}$  are defined in Section 2.2 below.

where  $\bar{p}^* = \frac{D\bar{p}}{D-\bar{p}}$ .

In particular if  $p = p_1 = \cdots = p_n$ , then  $\bar{p} = p$ ,  $\bar{p}^* = p^*$ , and we get

$$(1.9) \quad \|f\|_{L_{\mu}^{\bar{p}^*}} \leq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{\frac{A_i+1}{p}}} \quad \forall f \in C_c^1(\mathbb{R}^n),$$

which implies (1.3).

In the unweighted case, i.e.  $A_1 = \cdots, A_n = 0$ , inequality (1.9) is well-known (see *e.g.* [34], [33] and [23]). Anisotropic inequalities involving Orlicz norm defined using an  $n$ -dimensional Young function were also studied in [14]. However, as far we know, our anisotropic inequalities involving rearrangement invariant spaces are new in this context.

The paper is organized as follows. In Section 2 we provide a brief review on the rearrangements of functions and the theory of rearrangement invariant spaces. In Section 3 we will prove our main result (Theorem 3.1 below) which establishes the equivalence between (1.3) with  $p = 1$  and (1.5). Finally in Section 4 we use the oscillation inequality (1.5) to derive anisotropic Sobolev inequalities in  $\mathbb{R}^n$  with monomial weight  $x^A$  in the general setting of rearrangement invariant spaces, with special attention in the case of Lebesgue spaces, Lorentz spaces, Lorentz-Zygmund spaces, Gamma spaces and the recent class of  $G\Gamma$  spaces.

## 2. NOTATIONS AND PRELIMINARY RESULTS

We briefly recall the basic definitions of rearrangements and of rearrangement-invariant (r.i.) spaces referring the reader to [5] and [22].

**2.1. Rearrangement of functions.** Let  $\mu$  an absolutely continuous measure with respect to Lebesgue measure on  $\mathbb{R}^n$ . For a  $\mu$ -measurable function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ , the distribution function of  $u$  is given by

$$\mu_u(s) = \mu\{x \in \mathbb{R}^n : |u(x)| > s\} \quad s \geq 0.$$

The **decreasing rearrangement**  $u_{\mu}^*$  of  $u$  is the right-continuous non-increasing function from  $[0, \infty)$  into  $[0, \infty]$  which is equimeasurable with  $u$ . Namely,

$$u_{\mu}^*(t) = \inf\{s \geq 0 : \mu_u(s) \leq t\} \quad t \geq 0.$$

We also define  $u_{\mu}^{**} : (0, \infty) \rightarrow (0, \infty)$  as

$$u_{\mu}^{**}(t) = \frac{1}{t} \int_0^t u_{\mu}^*(s) ds,$$

Note that  $u_{\mu}^{**}$  is also decreasing and  $u_{\mu}^* \leq u_{\mu}^{**}$ , moreover

$$(u + v)_{\mu}^{**}(t) \leq u_{\mu}^{**}(t) + v_{\mu}^{**}(t)$$

for  $t > 0$ .

The **oscillation** of  $u$  is defined by

$$O_{\mu}(u, t) := u_{\mu}^{**}(t) - u_{\mu}^*(t).$$

Note that

$$(2.1) \quad tO_{\mu}(u, t) = \int_{u_{\mu}^*(t)}^{\infty} \mu_u(s) ds$$

is increasing.

When rearrangements are taken with respect to the Lebesgue measure on  $(0, \infty)$ , we may omit the measure and simply write  $u^*$  and  $u^{**}$ , etc...

**2.2. Rearrangement invariant spaces.** We say that a Banach function space  $X = X(\mathbb{R}^n)$  on  $(\mathbb{R}^n, \mu)$  is **rearrangement-invariant (r.i.) space**, if  $g \in X$  implies that  $f \in X$  for all  $\mu$ -measurable functions  $f$  such that  $f_\mu^* = g_\mu^*$ , and  $\|f\|_X = \|g\|_X$ .

A basic property of rearrangements is the Hardy-Littlewood inequality which tells us that, if  $u$  and  $w$  are two  $\mu$ -measurable functions on  $\mathbb{R}^n$ , then

$$(2.2) \quad \int_{\mathbb{R}^n} |u(x)w(x)| d\mu \leq \int_0^\infty u_\mu^*(t)w_\mu^*(t) dt.$$

An important consequence (2.2) is the Hardy-Littlewood-Pólya principle stating that

$$(2.3) \quad \int_0^r f_\mu^*(s) ds \leq \int_0^r g_\mu^*(s) ds \quad \forall r > 0 \Rightarrow \|f\|_X \leq \|g\|_X \quad \text{for any r.i. space } X.$$

A r.i. space  $X(\mathbb{R}^n)$  can be represented by a r.i. space on  $(0, +\infty)$ , with Lebesgue measure,  $\bar{X} = \bar{X}(0, \infty)$ , such that

$$\|f\|_X = \|f_\mu^*\|_{\bar{X}},$$

for every  $f \in X$ . A characterization of the norm  $\|\cdot\|_{\bar{X}}$  is available (see [5, Theorem 4.10 and subsequent remarks]). The space  $\bar{X}$  is called the **representation space** of  $X$ .

If  $X$  is a r.i. space, we have

$$L_\mu^1 \cap L^\infty \subset \bar{X} \subset L_\mu^1 + L^\infty,$$

with continuous embeddings.

The **associate space**  $X'$  of  $X$  is the r.i. space of all measurable functions  $h$  for which the r.i. norm given by

$$\|h\|_{X'} = \sup_{g \neq 0} \frac{\int_{\mathbb{R}^n} |g(x)h(x)| d\mu(x)}{\|g\|_X} = \sup_{g \neq 0} \frac{\int_0^\infty h_\mu^*(s)g_\mu^*(s) ds}{\|g\|_X}.$$

In particular the following generalized Hölder inequality

$$\int_{\mathbb{R}^n} |g(x)h(x)| d\mu(x) \leq \|g\|_X \|h\|_{X'}$$

holds.

Classically conditions on r.i. spaces are given in terms of the Hardy operators defined by

$$Pf(t) = \frac{1}{t} \int_0^t f(s) ds; \quad Q_a f(t) = \frac{1}{t^a} \int_t^\infty s^a f(s) \frac{ds}{s}, \quad 0 \leq a < 1,$$

(if  $a = 0$ , we shall write  $Q$  instead of  $Q_0$ ).

The boundedness of these operators on r.i. spaces can be described in terms of the so called **Boyd indices**<sup>4</sup> defined by

$$\bar{\alpha}_X = \inf_{s>1} \frac{\ln h_X(s)}{\ln s} \quad \text{and} \quad \underline{\alpha}_X = \sup_{s<1} \frac{\ln h_X(s)}{\ln s},$$

<sup>4</sup>Introduced by D.W. Boyd in [8].

where  $h_X(s)$  denotes the norm of the compression/dilation operator  $E_s$  on  $\bar{X}$ , defined for  $s > 0$ , by  $E_s f(t) = f^*(\frac{t}{s})$ . For example if  $X = L_\mu^p$  with  $p > 1$ , then  $\bar{\alpha}_X = \underline{\alpha}_X = \frac{1}{p}$ . It is well known that

$$\begin{aligned} P \text{ is bounded on } \bar{X} &\Leftrightarrow \bar{\alpha}_X < 1, \\ Q \text{ is bounded on } \bar{X} &\Leftrightarrow \underline{\alpha}_X > a. \end{aligned}$$

The next two Lemmas will be used in Section 4.

**Lemma 2.1.** (see [22, Page 43]) *Let  $X$  be a r.i. space and let  $0 \leq \theta_i \leq 1$  such that  $\sum_{i=1}^n \theta_i = 1$ , then*

$$(2.4) \quad \left\| \prod_{i=1}^n |f_i|^{\theta_i} \right\|_X \leq \prod_{i=1}^n \|f_i\|_X^{\theta_i}.$$

**Lemma 2.2.** *Let  $g, h$  be two positive measurable functions on  $(0, \infty)$  such that*

$$g(s) \leq h^{**}(s), \text{ for all } s \in (0, \infty),$$

and

$$\int_0^t g(s) ds \leq \int_0^t h^*(s) ds, \text{ for all } t \in (0, \infty).$$

Then

$$\int_0^t g^*(s) ds \leq 4 \int_0^t h^*(s) ds \text{ for all } t \in (0, \infty).$$

Therefore for any r.i. space  $X$

$$\|g\|_X \leq 4 \|h\|_X.$$

The proof of the previous lemma is implicitly contained in the proof of Theorem 1.2 of [15], a detailed proof can be found in [27].

### 2.2.1. Examples.

**Convexifications of r.i. spaces.** A way to construct r.i. spaces is through the so-called  $p$ -convexification, which is the generalization of the procedure to construct  $L^p$  spaces,  $1 < p < \infty$ , starting from  $L^1$ . If  $X$  is a r.i. space, the  $p$ -**convexification**  $X^{(p)}$  of  $X$ , (cf. [25]) is the r.i. space defined  $X^{(p)} = \{f : |f|^p \in X\}$  endowed with the following norm

$$(2.5) \quad \|f\|_{X^{(p)}} = \| |f|^p \|_X^{1/p}.$$

The same is true for the functional  $\|\cdot\|_{X^{(p)}}$  defined as

$$(2.6) \quad \|f\|_{X^{(p)}} = \left\| \left( (|f|^p)^{**} \right)^{1/p} \right\|_X.$$

Spaces  $X^{(p)}$  have been introduced in [17] in connection with the study of Sobolev embeddings into rearrangement-invariant spaces defined by a Frostman measure.

**The generalized Lorentz spaces**  $\Lambda^{p,q}(w)$ . Given  $1 \leq p, q < \infty$ , and  $w$  a weight (a positive locally integrable function) on  $(0, \infty)$ . The generalized Lorentz spaces  $\Lambda^{p,q}(w)$  are defined by measurable functions on  $(0, \infty)$  such that

$$(2.7) \quad \|f\|_{\Lambda^{p,q}(w)} := \left( \int_0^\infty \left( t^{1/p} f^*(t) \right)^q w(t) \frac{dt}{t} \right)^{1/q} < \infty.$$

If  $p = q$ , we write  $\Lambda^p(w)$  instead  $\Lambda^{p,p}(w)$ . We denote by  $W(t) = \int_0^t w(s) ds$ . It is known (see [10]) that  $\Lambda^{p,q}(w) = \Lambda^q(W^{q/p-1}w)$ . A weight  $w$  is called a  $B_p$ -weight if there is  $C > 0$  such that

$$\int_t^\infty \frac{w(s)}{s^p} \leq \frac{C}{t^p} \int_0^t w(s) ds, \quad t > 0.$$

The  $B_p$  class satisfies that  $B_r \subset B_p$  if  $r \leq p$ . If  $w \in B_p$  then (see [32])

$$\|f\|_{\Lambda^p(w)} \simeq \left( \int_0^\infty f^{**}(t)^p w(t) dt \right)^{1/p},$$

therefore Lorentz spaces defined by  $B_p$ -weights are r.i. spaces. Moreover, since (see [29, Theorem 6.5])

$$w \in B_p \Rightarrow W^{q/p-1}w \in B_q$$

we get that  $\Lambda^{p,q}(w)$  is a r.i. space if  $w \in B_p$ .

Given  $f$  a  $\mu$ -measurable function on  $\mathbb{R}^n$  we define

$$\|f\|_{\Lambda_\mu^{p,q}(w)} = \{f : \|f_\mu^*\|_{\Lambda^{p,q}(w)} < \infty\}.$$

Typical examples of generalized Lorentz spaces are the  $L^p$ -spaces and the **Lorentz spaces**  $L^{p,q}$ , defined either for  $p = q = 1$  or  $p = q = \infty$  or  $1 < p < \infty$ ,  $1 \leq q \leq \infty$  by

$$\|f\|_{L_\mu^{p,q}} := \left\| t^{1/p-1/q} f_\mu^*(t) \right\|_{L_{[0,\infty)}^q} < +\infty,$$

and, more generally, the **Lorentz-Zygmund spaces**, defined for  $1 < p < \infty$ ,  $1 \leq q \leq \infty$  and  $\alpha \in \mathbb{R}$  by

$$\|f\|_{L_\mu^{p,q}(\log L)^\alpha} := \left\| t^{1/p-1/q} (1 + \log |f|)^{\alpha/q} f_\mu^*(t) \right\|_{L_{[0,\infty)}^q} < +\infty$$

Let us notice that, in spite of the notation, the quantities  $\|f\|_{L^{p,q}}$  and  $\|f\|_{L^{p,q}(\log L)^\alpha}$  need not be norms; however, they can be turned into equivalent norms, when  $1 < p < \infty$ , replacing  $f^*$  by  $f^{**}$ .

**The Gamma spaces**  $\Gamma^p(w)$ . Let  $1 \leq p < \infty$ . Let  $w$  be an admissible weight, i.e.

$$(2.8) \quad \int_0^t w(s) ds < \infty \quad \text{and} \quad \int_t^\infty \frac{w(s)}{s^p} ds < \infty.$$

The Gamma space  $\Gamma^p(w)$  is the r.i. space defined as the set of measurable functions such that

$$(2.9) \quad \|f\|_{\Gamma^p(w)} := \left( \int_0^\infty f^{**}(s)^p w(s) ds \right)^{1/p} < \infty.$$

Given  $f$  a  $\mu$ -measurable function on  $\mathbb{R}^n$  we define

$$\|f\|_{\Gamma_\mu^p(w)} = \{f : \|f_\mu^*\|_{\Gamma^p(w)} < \infty\}.$$

**The  $G\Gamma^p(p, m, w)$  spaces.** Let  $1 \leq p, m < \infty$  and let  $w$  be a weight satisfying that

$$(2.10) \quad \int_0^\infty \min(s, t)^{m/p} w(t) dt < \infty, \quad s > 0.$$

The  $G\Gamma(p, m, w)$ -spaces are defined by

$$(2.11) \quad G\Gamma(p, m, w) = \left\{ f : \|f\|_{G\Gamma(p, m, w)} = \left( \int_0^\infty \left( \int_0^t f^*(s)^p ds \right)^{m/p} w(t) dt \right)^{1/m} < \infty \right\}.$$

These spaces has been introduced in [19] in connection with compact Sobolev type embedding results, since then its turn out to be important and several papers devoted to the study of this spaces have been published (see e.g. [18], [20], [21] and the references quoted therein).

**2.3. Some remarks about function spaces defined by oscillations.** In this subsection we analyze functional properties of function spaces whose definition involves the oscillation of  $f$ . The principal difficulty dealing with the functional  $O_\mu(f, t)$  is its nonlinearity. Therefore function spaces whose definition involves this quantity are not linear spaces.

Consider the Hardy type operator defined on positive measurable function on  $(0, \infty)$  by

$$\bar{Q}f(t) = \int_t^\infty s^{1/D} f(s) \frac{ds}{s}.$$

**Theorem 2.3.** *Let  $X$  be a r.i. space on  $(0, \infty)$  and let us assume that  $X$  does not contain constant functions.*

*i) If  $\underline{\alpha}_X > \frac{1}{D}$ , then*

$$\|t^{-1/D} f^{**}(t)\|_X \simeq \|t^{-1/D} [f^{**}(t) - f^*(t)]\|_X.$$

*ii) If  $\bar{\alpha}_X < \frac{1}{D}$ , then*

$$\|f\|_{L^\infty} \preceq \|t^{-1/D} [f^{**}(t) - f^*(t)]\|_X + \|f\|_{L^1_\mu + L^\infty}.$$

*Proof.* *i)* Let  $f \in X$ . Since  $X$  does not contain constant functions,  $f^{**}(\infty) = 0$ . An elementary computation shows that  $(-f^{**})'(t) = \frac{f^{**}(t) - f^*(t)}{t}$ , thus by the fundamental theorem of Calculus we get

$$f^{**}(t) = \int_t^\infty (f^{**}(t) - f^*(t)) \frac{ds}{s}.$$

Therefore,

$$\begin{aligned} \|t^{-1/D} f^{**}(t)\|_X &= \left\| t^{-1/D} \int_t^\infty (f^{**}(s) - f^*(s)) \frac{ds}{s} \right\|_X \\ &= \left\| t^{-1/D} \int_t^\infty s^{1/D} s^{-1/D} (f^{**}(s) - f^*(s)) \frac{ds}{s} \right\|_X \\ &\preceq \left\| s^{-1/D} (f^{**}(s) - f^*(s)) \right\|_X \quad (\text{since } \underline{\alpha}_X > \frac{1}{D}). \end{aligned}$$

The converse inequality is obvious.

ii) Let us assume now that  $\bar{\alpha}_X < \frac{1}{D}$ . We get

$$\begin{aligned} \|f\|_{L^\infty} + \|f\|_{L_\mu^1 + L^\infty} &= f^{**}(0) - f^{**}(1) = \int_0^1 (f^{**}(s) - f^*(s)) \frac{ds}{s} \\ &= \int_0^1 s^{1/D} \left( s^{-1/D} (f^{**}(s) - f^*(s)) \right) \frac{ds}{s} \\ &\leq \|s^{-1/D} (f^{**}(s) - f^*(s))\|_X \|s^{1/D-1} \chi_{[0,1]}(s)\|_{X'}. \end{aligned}$$

It is enough to check that  $\|s^{1/D-1} \chi_{[0,1]}(s)\|_{X'} < \infty$ . Since  $\bar{\alpha}_X = 1 - \underline{\alpha}_{X'} < \frac{1}{D}$ , we can select

$$(2.12) \quad 1 - 1/D < \beta < \underline{\alpha}_{X'}.$$

By the definition of indices, there is  $c > 0$  such that

$$(2.13) \quad h_{X'}(2^{-k}) \leq c 2^{-k\beta} \quad \forall k \geq 0.$$

For any  $k \geq 0$ , write  $I_k = [2^{-k-1}, 2^{-k})$ . Since  $\chi_{[0,1]}(s) = \sum_{k=0}^{\infty} \chi_{I_k}(s)$ , we get

$$\begin{aligned} \|s^{1/D-1} \chi_{[0,1]}(s)\|_{X'} &\leq \sum_{k=0}^{\infty} \|s^{1/D-1} \chi_{I_k}(s)\|_{X'} \leq \sum_{k=0}^{\infty} 2^{k(1-1/D)} \|\chi_{I_k}(s)\|_{X'} \\ &= \sum_{k=0}^{\infty} 2^{k(1-1/D)} \|D_{2^{-k}} \chi_{[1/2,1)}(s)\|_{X'} \\ &\leq \sum_{k=0}^{\infty} 2^{k(1-1/D)} h_{X'}(2^{-k}) \|\chi_{[1/2,1)}(s)\|_{X'} \\ &= \sum_{k=0}^{\infty} 2^{k(1-1/D-\beta)} 2^{k\beta} h_{X'}(2^{-k}) \|\chi_{[1/2,1)}(s)\|_{X'} \\ &\leq c \sum_{k=0}^{\infty} 2^{k(1-1/D-\beta)} \|\chi_{[1/2,1)}(s)\|_{X'} < \infty \quad (\text{by (2.13) and (2.12)}). \end{aligned}$$

□

### 3. SELF-IMPROVEMENT OF SOBOLEV INEQUALITY

Let  $W_0^{1,1}(\mathbb{R}^n, \mu)$  be the closure of the space  $C_c^1(\mathbb{R}^n)$  under the norm

$$\|u\|_{W_0^{1,1}(\mathbb{R}^n, \mu)} = \int_{\mathbb{R}^n} (|\nabla u| + |u|) d\mu.$$

Inequality (1.3) with  $p = 1$  implies the following anisotropic Sobolev inequality

$$(3.1) \quad \|f\|_{L_\mu^{\frac{D}{D-1}}} \leq \sum_{i=1}^n \left\| \frac{\partial f}{\partial x_i} \right\|_{L_\mu^1}$$

for  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ .

**Theorem 3.1.** *Let  $\mu$  and  $D$  be defined as in (1.1) and (1.2) respectively. Let  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$  the following statements hold and are equivalent:*

i) (Poincaré inequality)

$$(3.2) \quad \|f\|_{L_\mu^{\frac{D}{D-1}}} \leq \sum_{i=1}^n \|f_{x_i}\|_{L_\mu^1}.$$



ii) (Poincaré inequality in multiplicative form)

$$(3.3) \quad \|f\|_{L_\mu^{\frac{D}{D-1}}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_\mu^1}^{\frac{A_i+1}{D}}.$$

iii) (Mazya-Talenti's inequality in multiplicative form) The function  $f_\mu^*$  is locally absolutely continuous and for all  $s > 0$  we have that

$$(3.4) \quad s^{1-1/D} (-f_\mu^*)'(s) \preceq \prod_{i=1}^n \left( \frac{d}{ds} \int_{\{|f|>f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}}.$$

iv) (Oscillation inequality in multiplicative form) For all  $1 \leq p < \infty$  and for all  $t > 0$  we get

$$(3.5) \quad \int_0^t \left( O_\mu(f, \cdot)^p (\cdot)^{-\frac{p}{D}} \right)^* (s) ds \preceq \int_0^t \prod_{i=1}^n \left[ \left( \frac{d}{ds} \int_{\{|f|>f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^* (\tau) \right]^{p \left( \frac{A_i+1}{D} \right)} d\tau,$$

where rearrangements without subscript  $\mu$  are taken with respect to Lebesgue measure on  $(0, \infty)$ .

v)

$$\|f\|_{L_\mu^{\frac{D}{D-1},1}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_\mu^1}^{\frac{A_i+1}{D}}.$$

*Proof.* i)  $\Rightarrow$  ii)

We follow a scaling argument as in [33, Lemma 7]. Indeed we apply (3.2) to function  $w(x) = f(\lambda_1 x_1, \dots, \lambda_n x_n)$  and we obtain (3.3) by choosing

$$\lambda_i = \prod_{j \neq i} \|f_{x_j}\|_{L_\mu^1}.$$

ii)  $\Rightarrow$  iii)

Since  $L_\mu^{\frac{D}{D-1}}$  is continuously embedded in  $L_\mu^{\frac{D}{D-1},\infty}$  with constant 1, inequality (3.3) implies the weaker one

$$(3.6) \quad \sup_{t>0} t |\{x \in \mathbb{R}^n : |f(x)| > t\}|^{\frac{D-1}{D}} \leq nC \prod_{i=1}^n \|f_{x_i}\|_{L_\mu^1}^{\frac{A_i+1}{D}}.$$

Let  $0 < t_1 < t_2 < \infty$ , the truncations of  $f$  are defined by

$$f_{t_1}^{t_2}(x) = \begin{cases} t_2 - t_1 & \text{if } |f(x)| > t_2, \\ |f(x)| - t_1 & \text{if } t_1 < |f(x)| \leq t_2, \\ 0 & \text{if } |f(x)| \leq t_1. \end{cases}$$

Observe that if  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$  then  $f_{t_1}^{t_2} \in W_0^{1,1}(\mathbb{R}^n, \mu)$ , therefore replacing  $f$  by  $f_{t_1}^{t_2}$  in (3.6) we obtain

$$\sup_{t>0} t |\{x \in \mathbb{R}^n : |f_{t_1}^{t_2}(x)| > t\}|^{\frac{D-1}{D}} \leq nC \prod_{i=1}^n \left( \int_{\mathbb{R}^n} \left| \frac{\partial f_{t_1}^{t_2}}{\partial x_i} \right| d\mu \right)^{\frac{A_i+1}{D}}.$$

Since

$$\sup_{t>0} t |\{x \in \mathbb{R}^n : |f_{t_1}^{t_2}(x)| > t\}|^{\frac{D-1}{D}} \geq (t_2 - t_1) |\{x \in \mathbb{R}^n : |f(x)| \geq t_2\}|^{\frac{D-1}{D}},$$

and

$$\left| \frac{\partial f_{t_1}^{t_2}}{\partial x_i} \right| = \left| \frac{\partial f}{\partial x_i} \right| \chi_{\{t_1 < |f| \leq t_2\}}$$

we get

(3.7)

$$(t_2 - t_1) |\{x \in \mathbb{R}^n : |f(x)| \geq t_2\}|^{1-1/D} \leq nC \prod_{i=1}^n \left( \int_{\{t_1 < |f| \leq t_2\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}}.$$

Using that  $|f_{x_i}| \leq |\nabla f|$ , (3.7) implies

$$(t_2 - t_1) |\{x \in \mathbb{R}^n : |f(x)| \geq t_2\}|^{1-1/D} \leq nC \int_{\{t_1 < |f| \leq t_2\}} |\nabla f| d\mu$$

and from this inequality the locally absolute continuity of  $f_\mu^*$  follows easily using the same argument as in [26, page 137].

Let  $s > 0$  and  $h > 0$ , pick  $t_1 = f_\mu^*(s+h)$ ,  $t_2 = f_\mu^*(s)$ , then by (3.7) we get

$$(f_\mu^*(s) - f_\mu^*(s+h)) s^{1-1/D} \leq nC \prod_{i=1}^n \left( \int_{\{f_\mu^*(s+h) < |f| \leq f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}}.$$

Thus,

$$\frac{(f_\mu^*(s) - f_\mu^*(s+h))}{h} s^{1-1/D} \leq nC \prod_{i=1}^n \left( \frac{1}{h} \int_{\{f_\mu^*(s+h) < |f| \leq f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}}.$$

Letting  $h \rightarrow 0$  we obtain (3.4).

*iii)  $\Rightarrow$  iv)*

Let  $1 \leq p < \infty$ . For  $0 < s < t$ , we get

$$\begin{aligned} O_\mu(f, t) &= \frac{1}{t} \int_0^t (f_\mu^*(s) - f_\mu^*(t)) ds \\ &= \frac{1}{t} \int_0^t \left( \int_s^t (-f_\mu^*)'(\tau) d\tau \right) ds \\ &= \frac{1}{t} \int_0^t s (-f_\mu^*)'(s) ds \\ &\leq \frac{t^{1/D}}{t} \int_0^t s^{1-1/D} (-f_\mu^*)'(s) ds \\ &\leq nC \frac{t^{1/D}}{t} \int_0^t \prod_{i=1}^n \left( \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}} ds \quad (\text{by (3.4)}) \\ &\leq nC \frac{t^{1/D}}{t} \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right]^{\frac{A_i+1}{D}} \right)^* (s) ds \quad (\text{by (2.2)}) \\ &\leq nC t^{1/D} \left( \frac{1}{t} \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds \right)^{1/p} \quad (\text{by Hölder}). \end{aligned}$$

Thus

$$(3.8) \quad \left( t^{-1/D} O_\mu(f, t) \right)^p \leq \frac{1}{t} \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds.$$

Notice that the previous computation shows that

$$(3.9) \quad O_\mu(f, t) = \frac{1}{t} \int_0^t s (-f_\mu^*)' (s) ds.$$

If  $p = 1$ , then using (3.9) and (3.8) we get

$$\begin{aligned} \int_0^t O_\mu(f, s) s^{-1/D} ds &= \int_0^t \frac{s^{-1/D}}{s} \left( \int_0^s z (-f_\mu^*)' (z) dz \right) ds \\ &= \int_0^t z (-f_\mu^*)' (z) \left( \int_z^t \frac{s^{-1/D}}{s} ds \right) dz \\ &\leq \int_0^t z (-f_\mu^*)' (z) \left( \int_z^\infty \frac{s^{-1/D}}{s} ds \right) dz \\ &= D \int_0^t z^{1-1/D} (-f_\mu^*)' (z) dz \\ &\leq nCD \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\cdot)\}} |f_{x_i}(x)| d\mu(x) \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds. \end{aligned}$$

If  $1 < p < \infty$ , then using (3.8), Hardy's Inequalities (see [5, page 124]) and (3.9) we get

$$\begin{aligned} \int_0^t \left( O_\mu(f, s) s^{-1/D} \right)^p ds &= \int_0^t \left( \frac{s^{-1/D}}{s} \int_0^s z (-f_\mu^*)' (z) dz \right)^p ds \\ &\leq \int_0^t \left( \frac{1}{s} \int_0^s z^{1-1/D} (-f_\mu^*)' (z) dz \right)^p ds = \\ &\leq \int_0^t \left( z^{1-1/D} (-f_\mu^*)' (z) \right)^p dz \\ &\leq \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\cdot)\}} |f_{x_i}(x)| d\mu(x) \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds. \end{aligned}$$

By Lemma 2.2 and [5, Exercise 10, page 88], we get

$$\begin{aligned} \int_0^t \left( O_\mu(f, \cdot)^p (\cdot)^{-\frac{p}{D}} \right)^* (s) ds &\leq \int_0^t \left( \prod_{i=1}^n \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\cdot)\}} |f_{x_i}(x)| d\mu(x) \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds \\ &\leq \int_0^t \prod_{i=1}^n \left( \left[ \frac{d}{d\tau} \int_{\{|f| > f_\mu^*(\cdot)\}} |f_{x_i}(x)| d\mu(x) \right]^{p \frac{A_i+1}{D}} \right)^* (s) ds \end{aligned}$$

*iv)  $\Rightarrow$  v)*

By (2.3) and (2.4), we obtain

$$(3.10) \quad \left\| O_\mu(f, s) s^{-\frac{1}{D}} \right\|_{L^1} \leq 4nCD \left\| \prod_{i=1}^n \left( \left[ \frac{d}{d\tau} \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right]^{\frac{A_i+1}{D}} \right)^* (s) \right\|_{L^1} \\ \leq 4nCD \prod_{i=1}^n \left\| \left( \frac{d}{d\tau} \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right)^* (s) \right\|_{L^1}^{\frac{A_i+1}{D}}.$$

Moreover we get

$$(3.11) \quad \left\| \left( \frac{d}{d\tau} \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right)^* (s) \right\|_{L^1} = \int_0^\infty \left( \frac{d}{d\tau} \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right)^* (s) ds \\ = \int_0^\infty \frac{d}{d\tau} \left( \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right) d\tau \\ = \int_{\mathbb{R}^n} |f_{x_i}| d\mu.$$

Taking into account that  $\frac{\partial}{\partial t} f_\mu^{**}(t) = -(f_\mu^{**}(t) - f_\mu^*(t))/t$  and  $f_\mu^{**}(\infty) = 0$  (since  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ ) by the Fundamental Theorem Calculus we have

$$f_\mu^{**}(t) = \int_t^\infty (f_\mu^{**}(\tau) - f_\mu^*(\tau)) \frac{d\tau}{\tau}.$$

Therefore

$$\|f\|_{L^{\frac{D}{D-1},1}} := \int_0^\infty f_\mu^{**}(s) s^{-\frac{1}{D}} ds = \int_0^\infty \left[ \int_s^\infty (f_\mu^{**}(\tau) - f_\mu^*(\tau)) \frac{d\tau}{\tau} \right] s^{-\frac{1}{D}} ds \\ = \int_0^\infty (f_\mu^{**}(\tau) - f_\mu^*(\tau)) \left[ \frac{1}{\tau} \int_0^\tau s^{-\frac{1}{D}} ds \right] d\tau \\ = \frac{D}{D-1} \int_0^\infty (f_\mu^{**}(\tau) - f_\mu^*(\tau)) \tau^{-\frac{1}{D}} d\tau.$$

Conclusion follows by (3.10) and (3.11).

$v) \Rightarrow i)$

It is consequence of the continuous embedding of Lorentz space  $L_\mu^{\frac{D}{D-1},1}$  into Lebesgue space  $L_\mu^{\frac{D}{D-1}}$  and the relation between weighted arithmetic and geometric mean.  $\square$

**Remark 3.2.** *As in the classical case we have that (3.1) implies a better inequality involving the Lorentz norm  $\|\cdot\|_{L^{\frac{D}{D-1},1}}$  (see e.g. [16], [1], [28] and the bibliography therein).*

**Remark 3.3.** *By (3.8) we get*

$$(3.12) \quad t^{-\frac{1}{D}} O_\mu(f, t) \leq \frac{1}{t} \int_0^t \prod_{i=1}^n \left( \frac{d}{d\tau} \int_{\{|f|>f_\mu^*(\tau)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}} ds := I(t),$$

that is a pointwise oscillation inequality in multiplicative form. Moreover by (3.12) we recover the classical oscillation inequality (see [28] and [26])

$$t^{-\frac{1}{D}} O_\mu(f, t) \leq |\nabla f|^{**}(t).$$

As matter of the fact by Hölder inequality

$$\begin{aligned} (3.13) \quad I(t) &\leq \prod_{i=1}^n \left( \frac{1}{t} \int_0^t \left( \frac{d}{ds} \int_{\{|f|>f_\mu^*(s)\}} |f_{x_i}| d\mu \right) ds \right)^{\frac{A_i+1}{D}} \\ &= \prod_{i=1}^n \left( \frac{1}{t} \int_{\{|f|>f_\mu^*(t)\}} |f_{x_i}| d\mu \right)^{\frac{A_i+1}{D}} \\ &\leq \prod_{i=1}^n \left( \frac{1}{t} \int_0^t |f_{x_i}|_\mu^*(s) ds \right)^{\frac{A_i+1}{D}} \quad (\text{by (2.2)}) \\ &\leq |\nabla f|^{**}(t). \end{aligned}$$

#### 4. ANISOTROPIC INEQUALITIES IN REARRANGEMENT INVARIANT SPACES

In this section starting from the oscillation inequality (3.5) we derive some anisotropic inequalities in  $\mathbb{R}^n$  in the general setting of rearrangement invariant spaces.

We will use throughout this Section the following notation

$$(\tilde{f}_{x_i})_\mu^*(t) := \left( \frac{d}{ds} \int_{\{|f|>f_\mu^*(s)\}} |f_{x_i}| d\mu \right)^*(t).$$

In order to prove these kind of results we need the following results.

**Lemma 4.1.** *Let  $1 < p < \infty$  and let  $v$  be a weight (a positive locally integrable function on  $(0, \infty)$ ). Let*

$$u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{v(s)^{\frac{-1}{p-1}}(s)}{s^{\frac{p}{p-1}}} ds \right)^{1-p}.$$

Then, there exists  $C > 0$  such that

$$\left( \int_0^1 f^{**}(s)^p u(s) ds \right)^{1/p} \leq C \left( \int_0^1 (O_\mu(f, s))^p v(s) ds \right)^{1/p} + \left( \int_0^1 u(s) ds \right)^{1/p} \int_0^1 f^*(t) ds.$$

*Proof.* Since

$$\begin{aligned} \left( \int_0^t u(s) ds \right)^{1/p} \left( \int_t^1 \frac{v(s)^{\frac{-1}{p-1}}}{s^{\frac{p}{p-1}}} ds \right)^{(p-1)/p} &\leq \left( 1 + \int_t^1 \frac{v(s)^{\frac{-1}{p-1}}}{s^{\frac{p}{p-1}}} ds \right)^{(1-p)/p} \left( \int_t^1 \frac{v(s)^{\frac{-1}{p-1}}}{s^{\frac{p}{p-1}}} ds \right)^{(p-1)/p} \\ &\leq 1, \end{aligned}$$

the result follows from [4, Lemma 5.4].  $\square$

**Lemma 4.2.** *(see [31] and [2]) Let  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ , then*

$$\int_0^t (\tilde{f}_{x_i})_\mu^*(\tau) d\tau \leq \int_0^t |f_{x_i}|_\mu^*(\tau) d\tau, \quad (t \geq 0)$$

therefore by (2.3) for any r.i space  $X$  on  $(\mathbb{R}^n, \mu)$  we have that

$$\left\| (\tilde{f}_{x_i})_\mu^* \right\|_{\bar{X}} \leq \left\| |f_{x_i}|_\mu^* \right\|_{\bar{X}} = \|f_{x_i}\|_X.$$

#### 4.1. Convexification of r.i. spaces.

4.1.1. *The  $X^{(q)}$  convexification.* Let  $X$  be a r.i. space on  $\mathbb{R}^n$  and  $X^{(q)}$  its  $q$ -convexification defined in (2.5). In the next theorem we state some anisotropic inequalities for functions  $f$  such that  $f_{x_i} \in X^{(p_i)}$  with  $p_i \geq 1$  for  $i = 1, \dots, n$ .

**Theorem 4.3.** *Let  $X$  be a r.i. space on  $(\mathbb{R}^n, \mu)$  and  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ . If  $p_1, \dots, p_n \geq 1$ , then*

$$(4.1) \quad \|t^{-1/D}[f_\mu^{**}(t) - f_\mu^*(t)]\|_{\bar{X}(\bar{p})} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}},$$

where  $\bar{p}$  and  $D$  are defined in (1.7) and in (1.2), respectively and the involved norms are defined as in (2.5). Moreover

i) if  $\underline{\alpha}_X > \frac{\bar{p}}{D}$ , then

$$(4.2) \quad \|t^{-1/D}f_\mu^{**}(t)\|_{\bar{X}(\bar{p})} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}};$$

ii) if  $\bar{\alpha}_X < \frac{\bar{p}}{D}$ , then

$$(4.3) \quad \|f\|_{L^\infty} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}} + \|f\|_{L_\mu^1 + L^\infty}.$$

*Proof.* Since  $\bar{X}(\bar{p})$  is a r.i space, by Theorem 3.1 part *iv*) we get

$$\begin{aligned} \left\| (f_\mu^{**}(s) - f_\mu^*(s)) s^{-\frac{1}{D}} \right\|_{\bar{X}(\bar{p})} &\leq \left\| \prod_{i=1}^n \left[ (\tilde{f}_{x_i})_\mu^* \right]^{\frac{A_i+1}{D}} \right\|_{\bar{X}(\bar{p})} \\ &= \left\| \prod_{i=1}^n \left[ (\tilde{f}_{x_i})_\mu^* \right]^{\frac{p_i}{\bar{p}} \left( \frac{\bar{p}}{p_i} \frac{A_i+1}{D} \right)} \right\|_{\bar{X}(\bar{p})} \\ &\leq \prod_{i=1}^n \left\| (\tilde{f}_{x_i})_\mu^*(t) \right\|_{\bar{X}^{(p_i)}}^{\frac{A_i+1}{D}} \quad (\text{by (2.4)}) \\ &\leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}}^{\frac{A_i+1}{D}} \quad (\text{by Lemma 4.2}). \end{aligned}$$

On the other hand, since (see [5, Proposition 5.13, Chapter 3])

$$\underline{\alpha}_{\bar{X}(\bar{p})} = \frac{\underline{\alpha}_X}{\bar{p}} \quad \text{and} \quad \bar{\alpha}_{\bar{X}(\bar{p})} = \frac{\bar{\alpha}_X}{\bar{p}}$$

(4.2) and (4.3) follows from Theorem 2.3.  $\square$

**Remark 4.4.** *We stress in the previous proof if we start from (3.13) instead of (3.5) we can not consider the case when  $p_1 = \dots = p_n = 1$ .*

In the particular case  $X = L_\mu^1$ , the previous Theorem can be detailed as in the following proposition.

**Proposition 4.5.** *Let  $p_1, \dots, p_n \geq 1$  and  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ .  
i) If  $\bar{p} < D$ , then*

$$(4.4) \quad \|f\|_{L_{\mu}^{\bar{p}^*, \bar{p}}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i}^{\frac{A_i+1}{D}}},$$

where  $\bar{p}$  is defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

ii) If  $\bar{p} = D$ , then

$$(4.5) \quad \left( \int_0^1 \left( \frac{f_{\mu}^{**}(t)}{1 + \ln \frac{1}{t}} \right)^D \frac{dt}{t} \right)^{1/D} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i}^{\frac{A_i+1}{D}}} + \|f\|_{L_{\mu}^1 + L^{\infty}}.$$

iii) If  $\bar{p} > D$ , then

$$(4.6) \quad \|f\|_{L^{\infty}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i}^{\frac{A_i+1}{D}}} + \|f\|_{L_{\mu}^1 + L^{\infty}}.$$

*Proof.* Since  $(L_{\mu}^1)^{(\bar{p})} = L_{\mu}^{\bar{p}}$  and  $\alpha_{L^{\bar{p}}} = \bar{\alpha}_{L^{\bar{p}}} = \frac{1}{\bar{p}}$ , (4.4) and (4.6) follows from Theorem 4.3. We have only to prove (4.5). If  $v(t) = \frac{1}{t}$  in Lemma 4.1, then  $u(t) = (D-1) \left( \frac{1}{1 + \ln(\frac{1}{t})} \right)^{-D} \frac{1}{s}$ . Under this choice Lemma 4.1 allows us to get

$$\left( \int_0^1 \left( \frac{f_{\mu}^{**}(t)}{1 + \ln(\frac{1}{t})} \right)^D \frac{dt}{t} \right)^{1/D} \preceq \left( \int_0^1 (f_{\mu}^{**}(t) - f_{\mu}^*(t))^D \frac{dt}{t} \right)^{1/D} + f_{\mu}^{**}(1).$$

□

**Remark 4.6.** *Our result implies Theorem 1.1. Indeed it gives an embedding in Lorentz spaces. If  $A_1 = \dots = A_n = 0$  we obtain the same results of [33] and [23] for what concerns (4.4).*

**Remark 4.7.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded domain and  $f \in C_c^1(\Omega)$ . Arguing as in the proof of Proposition 4.5, we get*

$$\left( \int_0^{\mu(\Omega)} \left( \frac{f_{\mu}^{**}(t)}{1 + \ln \frac{\mu(\Omega)}{t}} \right)^D \frac{dt}{t} \right)^{1/D} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i}^{\frac{A_i+1}{D}}} + \frac{1}{\mu(\Omega)} \int_{\Omega} |f| d\mu.$$

Since

$$\sup_{0 < t < \mu(\Omega)} \frac{f_{\mu}^{**}(t)}{\left(1 + \ln \frac{\mu(\Omega)}{t}\right)^{\frac{D-1}{D}}} \preceq \left( \int_0^{\mu(\Omega)} \left( \frac{f_{\mu}^{**}(t)}{1 + \ln \frac{\mu(\Omega)}{t}} \right)^D \frac{dt}{t} \right)^{1/D},$$

we obtain the following anisotropic Trudinger inequality

$$\sup_{0 < t < \mu(\Omega)} \frac{f_{\mu}^{**}(t)}{\left(1 + \ln \frac{\mu(\Omega)}{t}\right)^{\frac{D-1}{D}}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i}^{\frac{A_i+1}{D}}} + \frac{1}{\mu(\Omega)} \int_{\Omega} |f| d\mu.$$

4.1.2. *The  $X^{(q)}$  convexification.*

**Theorem 4.8.** *Let  $X$  be a r.i. space on  $(\mathbb{R}^n, \mu)$  and  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ . If  $p_1, \dots, p_n \geq 1$ , then*

$$\|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\bar{X}(\bar{p})} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}},$$

where  $\bar{p}$  and  $D$  are defined in (1.7) and in (1.2), respectively and the involved norms are defined as in (2.6). Moreover

i) if  $\underline{\alpha}_{\bar{X}(\bar{p})} > \frac{1}{D}$ , then

$$\|t^{-1/D}f_{\mu}^{**}(t)\|_{\bar{X}(\bar{p})} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}}^{\frac{A_i+1}{D}};$$

ii) if  $\bar{\alpha}_{\bar{X}(\bar{p})} < \frac{1}{D}$ , then

$$\|f\|_{L^\infty} \leq \prod_{i=1}^n \|f_{x_i}\|_{X^{(p_i)}}^{\frac{A_i+1}{D}} + \|f\|_{L_\mu^1 + L^\infty}.$$

*Proof.* By Theorem 3.1 part iv) with  $p = \bar{p}$ , we get

$$\begin{aligned} \frac{1}{t} \int_0^t \left( O_\mu(f, \cdot)^{\bar{p}}(\cdot)^{-\frac{\bar{p}}{D}} \right)^*(s) ds &\leq \frac{1}{t} \int_0^t \prod_{i=1}^n \left[ (\tilde{f}_{x_i})_\mu^*(\tau) \right]^{\bar{p} \left( \frac{A_i+1}{D} \right)} d\tau \\ &= \frac{1}{t} \int_0^t \prod_{i=1}^n \left[ \left( (\tilde{f}_{x_i})_\mu^*(\tau) \right)^{p_i} \right]^{\frac{\bar{p}}{p_i} \left( \frac{A_i+1}{D} \right)} d\tau \\ &\leq \prod_{i=1}^n \left( \frac{1}{t} \int_0^t \left[ \left( (\tilde{f}_{x_i})_\mu^*(\tau) \right)^{p_i} \right] d\tau \right)^{\frac{\bar{p}}{p_i} \left( \frac{A_i+1}{D} \right)} \end{aligned}$$

Let  $X$  a r.i. space, then

$$\begin{aligned} \|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\bar{X}(\bar{p})} &= \left\| \left( \frac{1}{t} \int_0^t \left( O_\mu(f, \cdot)^{\bar{p}}(\cdot)^{-\frac{\bar{p}}{D}} \right)^*(s) ds \right)^{1/\bar{p}} \right\|_{\bar{X}} \\ &\leq \left\| \prod_{i=1}^n \left( \frac{1}{t} \int_0^t \left[ \left( (\tilde{f}_{x_i})_\mu^*(\tau) \right)^{p_i} \right] d\tau \right)^{\frac{1}{p_i} \left( \frac{A_i+1}{D} \right)} \right\|_{\bar{X}} \\ &\leq \prod_{i=1}^n \left\| \left( \frac{1}{t} \int_0^t \left[ \left( (\tilde{f}_{x_i})_\mu^*(\tau) \right)^{p_i} \right] d\tau \right)^{\frac{1}{p_i} \left( \frac{A_i+1}{D} \right)} \right\|_{\bar{X}} \\ &= \prod_{i=1}^n \left\| (\tilde{f}_{x_i})_\mu^*(\tau) \right\|_{\bar{X}^{(p_i)}}^{\left( \frac{A_i+1}{D} \right)} \\ &\leq \prod_{i=1}^n \|f_{x_i}\|_{\bar{X}^{(p_i)}}^{\frac{A_i+1}{D}} \quad (\text{by Lemma 4.2}). \end{aligned}$$

The statements (4.2) and (4.3) follows from Theorem 2.3.  $\square$



**4.2. Generalized Lorentz spaces.** In this section we state some anisotropic inequalities for functions  $f$  such that the partial derivatives are in some generalized Lorentz spaces. More precisely as a consequence of Theorem 4.3 we obtain the following theorem.

**Theorem 4.9.** *Let  $w \in B_{\min(p_1, \dots, p_n)}$ ,  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ ,  $p_1, \dots, p_n \geq 1$ ,  $q_1, \dots, q_n \geq 1$  and the involved norms are defined as in (2.7).*

i) *If  $\underline{\alpha}_{\Lambda^{\bar{p}, \bar{q}}(w)} > \frac{1}{D}$ , then*

$$\|f\|_{\Lambda_{\mu}^{\bar{p}^*, \bar{q}}(w)} \lesssim \prod_{i=1}^n \|f_{x_i}\|_{\Lambda_{\mu}^{p_i, q_i}(w)}^{\frac{A_i+1}{D}},$$

where  $\bar{p}$  and  $\bar{q}$  are defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

ii) *If  $\bar{\alpha}_{\Lambda^{\bar{p}, \bar{q}}(w)} > \frac{1}{D}$ , then*

$$\|f\|_{L^\infty} \lesssim \prod_{i=1}^n \|f_{x_i}\|_{\Lambda_{\mu}^{p_i, q_i}(w)}^{\frac{A_i+1}{D}} + \|f\|_{L_{\mu}^1 + L^\infty}.$$

iii) *In the remaining cases we get*

$$\left( \int_0^1 f^{**}(s)^{\bar{q}} u(s) ds \right)^{1/\bar{q}} \lesssim \prod_{i=1}^n \|f_{x_i}\|_{\Lambda_{\mu}^{p_i, q_i}(w)}^{\frac{A_i+1}{D}} + \|f\|_{L_{\mu}^1 + L^\infty},$$

$$\text{where } u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{\left( s^{\frac{\bar{q}}{p} - \frac{\bar{q}}{D} - 1} w(s) \right)^{\frac{1}{\bar{q}-1}}}{s^{\frac{\bar{q}}{q}-1}} ds \right)^{1-\bar{q}}.$$

*Proof.* Since  $w \in B_{\min(p_1, \dots, p_n)}$  all the spaces  $\Lambda^{p_i, q_i}(w)$  are r.i spaces and from  $\min(p_1, \dots, p_n) \leq \bar{p}$  it follows that  $\Lambda^{\bar{p}, \bar{q}}(w)$  is a r.i. space. We note that

$$(4.7) \quad 1 = \frac{\bar{q}}{D} \sum_{i=1}^n \frac{A_i + 1}{q_i}.$$

By Theorem 3.1 we obtain

$$\begin{aligned} \|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Lambda^{\bar{p}, \bar{q}}(w)} &\leq \left\| \prod_{i=1}^n \tilde{f}_{x_i}(\tau)^{\frac{A_i+1}{D}} \right\|_{\Lambda^{\bar{p}, \bar{q}}(w)} \\ &= \left( \int_0^\infty \left( t^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \prod_{i=1}^n \tilde{f}_{x_i}(t)^{\frac{A_i+1}{D}} \right)^{\bar{q}} w(t) dt \right)^{\frac{1}{\bar{q}}} \\ &\leq \prod_{i=1}^n \left( \int_0^\infty \left( t^{\frac{1}{p_i} - \frac{1}{q_i}} \tilde{f}_{x_i}(t)^{\frac{A_i+1}{D}} \right)^{q_i} w(t) dt \right)^{\frac{A_i+1}{q_i D}} \quad (\text{by (4.7) and (2.4)}) \\ &= \prod_{i=1}^n \left\| \tilde{f}_{x_i} \right\|_{\Lambda^{p_i, q_i}(w)}^{\frac{A_i+1}{D}} \\ &\leq \prod_{i=1}^n \|f_{x_i}\|_{\Lambda_{\mu}^{p_i, q_i}(w)}^{\frac{A_i+1}{D}} \quad (\text{by Lemma 4.2}). \end{aligned}$$

Part *i*) follows from *i*) of Theorem 2.3, because

$$\begin{aligned} \|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Lambda^{\bar{p}, \bar{q}}(w)} &\simeq \|t^{-1/D}f_{\mu}^{**}(t)\|_{\Lambda^{\bar{p}, \bar{q}}(w)} \\ &\simeq \|f_{\mu}^{**}(t)\|_{\Lambda^{\bar{p}^*, \bar{q}}(w)} \\ &\simeq \|f\|_{\Lambda_{\mu}^{\bar{p}^*, \bar{q}}(w)}. \end{aligned}$$

Part *ii*) is consequence of *i*) of Theorem 2.3.

Let us prove now *iii*). If we denote  $I := \|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Lambda^{\bar{p}, \bar{q}}(w)}$  we get

$$\begin{aligned} I &= \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \left( t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)] \right)^*(s) \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \\ &\simeq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \frac{1}{s} \int_0^s \left( t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)] \right)^*(z) dz \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \quad (\text{since } w \in B_{\bar{p}}) \\ &\geq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \frac{1}{2s} \int_0^{2s} \left( t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)] \right)^*(z) dz \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \\ &\geq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \frac{1}{2s} \int_0^{2s} \left( z^{-1/D}[f_{\mu}^{**}(z) - f_{\mu}^*(z)] \right) dz \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \\ &\geq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \frac{1}{2s} \int_s^{2s} \left( z^{-1/D}[f_{\mu}^{**}(z) - f_{\mu}^*(z)] \right) dz \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \\ &\geq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}}} \frac{s[f_{\mu}^{**}(s) - f_{\mu}^*(s)]}{2s} \int_s^{2s} \left( z^{-1/D-1} \right) dz \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \quad (\text{by (2.1)}) \\ &\simeq \left( \int_0^{\infty} \left( s^{\frac{1}{\bar{p}} - \frac{1}{\bar{q}} - \frac{1}{D}} [f_{\mu}^{**}(s) - f_{\mu}^*(s)] \right)^{\bar{q}} w(s) ds \right)^{1/\bar{q}} \\ &= \left( \int_0^{\infty} [f_{\mu}^{**}(s) - f_{\mu}^*(s)]^{\bar{q}} s^{\frac{\bar{q}}{\bar{p}} - \frac{\bar{q}}{D} - 1} w(s) ds \right)^{1/\bar{q}}. \end{aligned}$$

Considering as weights

$$v(s) := s^{\frac{\bar{q}}{\bar{p}} - \frac{\bar{q}}{D} - 1} w(s) \quad \text{and} \quad u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{\left( s^{\frac{\bar{q}}{\bar{p}} - \frac{\bar{q}}{D} - 1} w(s) \right)^{\frac{-1}{\bar{q}-1}}}{s^{\frac{\bar{q}}{\bar{q}-1}}} ds \right)^{1-\bar{q}},$$

the result follows from Lemma 4.1.  $\square$

The following corollaries follow from the previous theorem considering  $w = 1$  and  $w(t) = (1 + |\ln t|)^{\alpha}$  with  $\alpha \in \mathbb{R}$ , respectively, and recalling that

$$\bar{\alpha}_{L^{p,q}(\log L)^{\alpha}} = \underline{\alpha}_{L^{p,q}(\log L)^{\alpha}} = \bar{\alpha}_{L^{p,q}} = \underline{\alpha}_{L^{p,q}} = \frac{1}{p}.$$

**Corollary 4.10.** *Let  $f \in C_c^1(\mathbb{R}^n)$ ,  $p_1, \dots, p_n \geq 1$  and  $q_1, \dots, q_n \geq 1$ . *i*) If  $\bar{p} < D$ , then*

$$\|f\|_{L_{\mu}^{\bar{p}, \bar{q}}} \leq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}},$$

where  $\bar{p}$  and  $\bar{q}$  are defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

ii) If  $\bar{p} > D$ , then

$$\|f\|_{L^\infty(\mathbb{R}^n)} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}}^{\frac{A_i+1}{D}} + \|f\|_{L_{\mu}^1 + L^\infty}.$$

iii) If  $\bar{p} = D$ , then

$$\left( \int_0^1 \left( \frac{f_{\mu}^{**}(s)}{1 + \ln \frac{1}{s}} \right)^{\bar{q}} \frac{ds}{s} \right)^{1/\bar{q}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}}^{\frac{A_i+1}{D}} + \|f\|_{L_{\mu}^1 + L^\infty}.$$

**Corollary 4.11.** Let  $f \in C_c^1(\mathbb{R}^n)$ ,  $p_1, \dots, p_n \geq 1$ ,  $q_1, \dots, q_n \geq 1$  and  $\alpha \in \mathbb{R}$ .

i) If  $\bar{p} < D$ , then

$$\|f\|_{L_{\mu}^{\bar{p}^*, \bar{q}}(\log L)^\alpha} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}}^{\frac{A_i+1}{D}} (\log L)^\alpha.$$

where  $\bar{p}$  and  $\bar{q}$  are defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

ii) If  $\bar{p} > D$ , then

$$\|f\|_{L^\infty(\mathbb{R}^n)} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}(\log L)^\alpha} + \|f\|_{L^1 + L^\infty}.$$

iii) If  $\bar{p} = D$ , then

$$\left( \int_0^1 \left( \frac{f_{\mu}^{**}(s)}{1 + \ln \frac{1}{s}} \right)^{\bar{q}} \left( 1 + \ln \frac{1}{s} \right)^\alpha \frac{ds}{s} \right)^{1/\bar{q}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{L_{\mu}^{p_i, q_i}(\log L)^\alpha} + \|f\|_{L_{\mu}^1 + L^\infty}.$$

When  $A_1 = \dots = A_n = 0$ , i) of Corollary 4.10 is contained in [33] and [23]. For our knowledge the other ones are new.

### 4.3. The Gamma spaces.

**Theorem 4.12.** Let  $w$  be a weight satisfying condition (2.8),  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ ,  $p_1, \dots, p_n \geq 1$  and the involved norms are defined as in (2.9).

i) If  $\underline{\alpha}_{\Gamma^{\bar{p}^*, (w)}} > \frac{1}{D}$ , then

$$\|t^{-1/D} f_{\mu}^*(t)\|_{\Gamma^{\bar{p}^*, (w)}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{\Gamma_{\mu}^{p_i}(w)},$$

where  $\bar{p}$  is defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

ii) If  $\bar{\alpha}_{\Gamma^{\bar{p}^*, (w)}} < \frac{1}{D}$ , then

$$\|f\|_{L^\infty} \preceq \prod_{i=1}^n \|f_{x_i}\|_{\Gamma_{\mu}^{p_i}(w)} + \|f\|_{L_{\mu}^1 + L^\infty}.$$

iii) In the remaining cases we get

$$\left( \int_0^1 f^{**}(s)^{\bar{p}^*} u(s) ds \right)^{1/\bar{p}^*} \preceq \prod_{i=1}^n \|f_{x_i}\|_{\Gamma_{\mu}^{p_i}(w)} + \|f\|_{L_{\mu}^1 + L^\infty},$$

$$\text{where } u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{\left( s^{-\frac{D}{\bar{p}^*}} w(s) \right)^{\frac{-1}{\bar{p}^*-1}}}{s^{\frac{D}{\bar{p}^*}-1}} ds \right)^{1-\bar{p}^*}.$$

*Proof.*

$$\begin{aligned}
\|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Gamma\bar{p}^*(w)} &\preceq \left\| \prod_{i=1}^n (\tilde{f}_{x_i})_{\mu}^*(t)^{\frac{A_i+1}{D}} \right\|_{\Gamma\bar{p}^*(w)} \\
&= \left( \int_0^{\infty} \left( \frac{1}{t} \int_0^t \prod_{i=1}^n (\tilde{f}_{x_i})_{\mu}^*(s)^{\frac{A_i+1}{D}} ds \right)^{\bar{p}^*} w(t) dt \right)^{\frac{1}{\bar{p}^*}} \\
&\leq \prod_{i=1}^n \left( \int_0^{\infty} \left( \frac{1}{t} \int_0^t (\tilde{f}_{x_i})_{\mu}^*(s) ds \right)^{p_i} w(t) dt \right)^{\frac{A_i+1}{p_i D}} \quad (\text{by (4.7) and (2.4)}) \\
&= \prod_{i=1}^n \left\| (\tilde{f}_{x_i})_{\mu}^* \right\|_{\Gamma_{\mu}^{p_i}(w)}^{\frac{A_i+1}{D}} \\
&\leq \prod_{i=1}^n \|f_{x_i}\|_{\Gamma_{\mu}^{p_i}(w)}^{\frac{A_i+1}{D}} \quad (\text{by Lemma 4.2}).
\end{aligned}$$

Part *i*) follows from *i*) of Theorem 2.3, since

$$\|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Gamma\bar{p}^*(w)} \simeq \|t^{-1/D}f_{\mu}^{**}(t)\|_{\Gamma\bar{p}^*(w)} \simeq \|t^{-1/D}f_{\mu}^*(t)\|_{\Gamma\bar{p}^*(w)}.$$

Part *ii*) is consequence of *i*) of Theorem 2.3.

Let us prove now *iii*). If we denote  $I := \|t^{-1/D}[f_{\mu}^{**}(t) - f_{\mu}^*(t)]\|_{\Gamma\bar{p}^*(w)}$  using the same argument that in part *iii*) of Theorem 4.9 we easily obtain

$$I \geq \left( \int_0^{\infty} [f_{\mu}^{**}(s) - f_{\mu}^*(s)]^{\bar{q}} s^{-\frac{\bar{p}^*}{D}} w(s) ds \right)^{1/\bar{p}^*}.$$

Considering as weights

$$v(s) := s^{-\frac{\bar{p}^*}{D}} w(s) \quad \text{and} \quad u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{\left( s^{-\frac{\bar{p}^*}{D}} w(s) \right)^{\frac{-1}{\bar{p}^*-1}}}{s^{\frac{\bar{p}^*}{\bar{p}^*-1}}} ds \right)^{1-\bar{p}^*},$$

the result follows from Lemma 4.1.  $\square$

#### 4.4. The $G\Gamma(m, p, w)$ -spaces.

**Theorem 4.13.** *Let  $w$  be a satisfying (2.10),  $f \in W_0^{1,1}(\mathbb{R}^n, \mu)$ ,  $p_1, \dots, p_n \geq 1$  and the involved norms are defined as in (2.11).*

*i*) If  $\underline{\alpha}_{G\Gamma(\bar{p}^*, m, w)} > \frac{1}{D}$ , then

$$\|t^{-1/D}f_{\mu}^*(t)\|_{G\Gamma(\bar{p}^*, m, w)} \preceq \prod_{i=1}^n \|f_{x_i}\|_{G\Gamma(p_i, m, w)}^{\frac{A_i+1}{D}},$$

where  $\bar{p}$  is defined as in (1.7) and  $\bar{p}^* = \frac{\bar{p}D}{D-\bar{p}}$ .

*ii*) If  $\bar{\alpha}_{G\Gamma(\bar{p}^*, m, w)} < \frac{1}{D}$ , then

$$\|f\|_{L^{\infty}} \preceq \prod_{i=1}^n \|f_{x_i}\|_{G\Gamma(p_i, m, w)}^{\frac{A_i+1}{D}} + \|f\|_{L_{\mu}^1 + L^{\infty}}.$$

iii) In the remaining cases we get

$$\left( \int_0^1 f^{**}(s)^m u(s) ds \right)^{1/m} \asymp \prod_{i=1}^n \|f_{x_i}\|_{G\Gamma(p_i, m, w)}^{\frac{A_i+1}{D}} + \|f\|_{L_\mu^1 + L^\infty},$$

$$\text{where } u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{(s^{-\frac{m}{D}} w(s))^{\frac{-1}{m-1}}}{s^{\frac{m}{m-1}}} ds \right)^{1-m}.$$

*Proof.* By Theorem 3.1 we get

$$\begin{aligned} \|t^{-1/D} [f_\mu^{**}(t) - f_\mu^*(t)]\|_{G\Gamma(\bar{p}^*, m, w)} &\lesssim \left\| \prod_{i=1}^n (\tilde{f}_{x_i})_\mu^*(t)^{\frac{A_i+1}{D}} \right\|_{G\Gamma(\bar{p}^*, m, w)} \\ &= \left( \int_0^\infty \left( \int_0^t \prod_{i=1}^n (\tilde{f}_{x_i})_\mu^*(s)^{\bar{p}^* \frac{A_i+1}{D}} ds \right)^{m/\bar{p}^*} w(t) dt \right)^{\frac{1}{m}} \\ &\leq \left( \int_0^\infty \prod_{i=1}^n \left( \int_0^t (\tilde{f}_{x_i})_\mu^*(s)^{p_i} ds \right)^{\frac{m}{p_i} \frac{A_i+1}{D}} w(t) dt \right)^{\frac{1}{m}} \\ &\leq \prod_{i=1}^n \left( \int_0^\infty \left( \int_0^t (\tilde{f}_{x_i})_\mu^*(s)^{p_i} ds \right)^{\frac{m}{p_i}} w(t) dt \right)^{\frac{1}{m} \frac{A_i+1}{D}} \\ &= \prod_{i=1}^n \left\| (\tilde{f}_{x_i})_\mu^* \right\|_{G\Gamma(p_i, m, w)}^{\frac{A_i+1}{D}} \\ &\leq \prod_{i=1}^n \|f_{x_i}\|_{G\Gamma(p_i, m, w)}^{\frac{A_i+1}{D}} \quad (\text{by Lemma 4.2}). \end{aligned}$$

Part *i*) follows from *i*) of Theorem 2.3, because

$$\|t^{-1/D} [f_\mu^{**}(t) - f_\mu^*(t)]\|_{G\Gamma(\bar{p}^*, m, w)} \simeq \|t^{-1/D} f_\mu^{**}(t)\|_{G\Gamma(\bar{p}^*, m, w)}$$

Part *ii*) is consequence of *i*) of Theorem 2.3.

Let us prove now *iii*). If we denote  $I := \|t^{-1/D} [f_\mu^{**}(t) - f_\mu^*(t)]\|_{G\Gamma(\bar{p}^*, m, w)}$  using the method of part *iii*) of Theorem 4.9 we get

$$I \geq \left( \int_0^\infty [f_\mu^{**}(s) - f_\mu^*(s)]^m s^{-\frac{m}{D}} w(s) ds \right)^{1/m}.$$

Considering as weights

$$v(s) := s^{-\frac{m}{D}} w(s) \quad \text{and} \quad u(t) = \frac{\partial}{\partial t} \left( 1 + \int_t^1 \frac{(s^{-\frac{m}{D}} w(s))^{\frac{-1}{m-1}}}{s^{\frac{m}{m-1}}} ds \right)^{1-m},$$

the result follows from Lemma 4.1.  $\square$

*Acknowledgments.* The first author has been partially supported by FFABR and the second one has been partially supported in part by Grants MTM2016-77635-P, MTM2016-75196-P (MINECO) and by 2017SGR358.

The second author gratefully acknowledges the generous hospitality of the Università degli Studi di Napoli “Federico II” Dipartimento di Matematica “R. Caccioppoli” where much of this work was done.

## REFERENCES

- [1] A. Alberico, A. Cianchi, L. Pick and L. Slavíková, *Sharp Sobolev type embeddings on the entire Euclidean space*. Commun. Pure Appl. Anal. **17** (2018), n. 5, 2011?-2037.
- [2] A. Alvino and G. Trombetti, *Sulle migliori costanti di maggiorazione per una classe di equazioni ellittiche degeneri*. Ricerche Mat. **27** (1978), 413–428.
- [3] D. Bakry, I. Gentil and M. Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Grundlehren der Mathematischen Wissenschaften, vol. 348. Springer, Berlin (2013).
- [4] J. Bastero, M. Milman and F.J. Ruiz. *A note on  $L(\infty, q)$ s spaces and Sobolev embeddings*. Indiana Univ. Math. J. **52** (2003), 1215–1230.
- [5] C. Bennett and R. Sharpley, *Interpolation of Operators*. Academic Press, Boston, 1988.
- [6] H. Brezis. *Is there failure of the inverse function theorem? Morse theory, Minimax theory and their applications to nonlinear differential equations*. In: Proceedings of the Workshop held at the Chinese Academy of Sciences, Beijing, 1999, 2333, New Stud. Adv. Math., 1, Int. Press, Somerville, MA, (2003)
- [7] H. Brezis and J.L. Vázquez. *Blow-up solutions of some nonlinear elliptic problems*. Rev. Mat. Univ. Complut. **10** (1997), 443–469
- [8] D.W. Boyd. *Indices of function spaces and their relationship to interpolation*. Canad. J. Math. **21** (1969), 1245–1254.
- [9] F. Brock , F. Chiacchio and A. Mercaldo. *A weighted isoperimetric inequality in an orthant*. Potential Anal. **41** (2014), 171–186.
- [10] M. J. Carro and J. Soria. *Weighted Lorentz spaces and the Hardy operator*. J. Funct. Anal. **112** (1993), 480–494.
- [11] X. Cabré. *Isoperimetric, Sobolev, and eigenvalue inequalities via the Alexandroff-Bakelman-Pucci method: a survey*. Chin. Ann. Math., **38B(1)** (2017), 201–214.
- [12] X. Cabré and X. Ros-Oton. *Sobolev and isoperimetric inequalities with monomial weights*. J. Differential Equations, **255** (2013), 4312–4336.
- [13] X. Cabré and X. Ros-Oton. *Regularity of stable solutions up to dimension 7 in domains of double revolution*. Commun. Partial Differ. Equ. **38** (2013), 135–154.
- [14] A. Cianchi. *A fully anisotropic Sobolev inequality*. Pacific J. Math. **196** (2000), n. 2, 283–295.
- [15] A. Cianchi. *Symmetrization and second-order Sobolev inequalities*. Ann.Mat. **183** (2004), 45–77.
- [16] A. Cianchi. *Optimal Orlicz-Sobolev embeddings*. Rev. Mat. Iberoamericana **20** (2004), 427–474.
- [17] A. Cianchi, L. Pick and L. Slavikova. *Sobolev embeddings. rearrangement-invariant spaces and Frostman measures*. Ann. Inst. H. Poincaré Anal. Non Linéaire (to appear).
- [18] Fiorenza, A.; Rakotoson, J. M. *Some estimates in  $G\Gamma(p, m, w)$  spaces*. J. Math. Anal. Appl. **340** (2008), 793–805.
- [19] A. Fiorenza, J.M. Rakotoson, *Compactness, interpolation inequalities for small Lebesgue-Sobolev spaces and applications*. Calc. Var. Partial Differential Equations **25** (2005) 187–203.
- [20] A. Fiorenza, M.R. Formica, A. Gogatishvili, T. Kopaliani, and J.M. Rakotoson. *Characterization of interpolation between grand, small or classical Lebesgue spaces*. Nonlinear Anal. **177** (2018),422–453.
- [21] A. Gogatishvili, M. Krepla, L. Pick and F. Soudský. *Embeddings of Lorentz-type spaces involving weighted integral means*. J. Funct. Anal. **273** (2017), 2939–2980.
- [22] S. G. Krein, Yu. I. Petunin and E. M. Semenov. *Interpolation of linear operators*, *Transl. Math. Monogr. Amer. Math. Soc.* **54**, Providence, (1982).
- [23] V. I. Kolyada and F. J. Pérez. *Estimates of difference norms for functions in anisotropic Sobolev spaces*. Mathematische Nachrichten **267(1)** (2004), 46–64.

- [24] N. Lam. *Sharp Trudinger-Moser inequalities with monomial weights*. NoDEA Nonlinear Differential Equations Appl. **24** (2017), no. 4, Art. 39, 21 pp.
- [25] J. Lindenstrauss and L. Tzafriri. *Classical Banach Spaces II. Function Spaces*, Springer-Verlag, Berlin, 1979.
- [26] J. Martin and M. Milman. *Pointwise symmetrization inequalities for Sobolev functions and applications*. Adv. Math. **225** (2010), 121–199.
- [27] J. Martin and M. Milman. *Self improving Sobolev-Poincaré inequalities, truncation and symmetrization*. Potential Anal. 29 (2008), 391–408.
- [28] J. Martin, M. Milman and E. Pustylnik. *Sobolev inequalities: Symmetrization and self-improvement via truncation*. J. Funct. Anal. **252** (2007), 677–695.
- [29] C. J. Neugebauer. *Weighted norm inequalities for averaging operators of monotone functions*. Publ. Mat. **35** (1991), 429–447.
- [30] V. H. Nguyen. *Sharp weighted Sobolev and Gagliardo-Nirenberg inequalities on half-spaces via mass transport and consequences*. Proc. Lond. Math. Soc.(3) **111** (2015), n. 1, 127–148.
- [31] J. Rakotoson. Réarrangement relatif. (French) [Relative rearrangement] Un instrument d'estimations dans les problèmes aux limites. [An estimation tool for limit problems] Mathématiques & Applications (Berlin) [Mathematics & Applications], 64. Springer, Berlin, 2008. xvi+293 pp.
- [32] E. Sawyer. *Boundedness of classical operators on classical Lorentz spaces*. Studia Math. **96** (1990), 145–158.
- [33] L. Tartar. *Imbedding theorems of Sobolev spaces into Lorentz spaces*. BUMI Serie **8 1-B** (1998), 479–500.
- [34] M. Troisi. *Teoremi di inclusione per spazi di Sobolev non isotropi*. Ricerche Mat. **18** (1969), 3–24.

DIPARTIMENTO DI INGEGNERIA, UNIVERSITÀ DEGLI STUDI DI NAPOLI “PARTHENOPÉ”, CENTRO DIREZIONALE, ISOLA C4, 80143 NAPOLI, ITALY

*E-mail address:* `filomena.feo@uniparthenope.it`

*Current address:* Universitat Autònoma de Barcelona, Department of Mathematics, Bellaterra (Barcelona), Spain

*E-mail address:* `jmartin@mat.uab.cat`

UNIVERSITÀ DEGLI STUDI DI NAPOLI “FEDERICO II”, DIPARTIMENTO DI MATEMATICA “R. CACCIOPOLI”, COMPLESSO MONTE S. ANGELO, NAPOLI, ITALY

*E-mail address:* `posterar@unina.it`