

# A FAMILY OF K3 SURFACES AND TOWERS OF ALGEBRAIC CURVES OVER FINITE FIELDS

SERGEY GALKIN AND SERGEY RYBAKOV

## 1. INTRODUCTION

Let  $\bar{k}$  be an algebraic closure of a finite field  $k$  of characteristic  $p$ , and  $\bar{Y}$  be the base change

$$\bar{Y} = Y \times_{\mathrm{Spec} k} \mathrm{Spec} \bar{k}$$

of an algebraic variety  $Y$  over  $k$ . We also fix a prime number  $\ell \neq p$ .

In the previous paper [Ry] we constructed an optimal tower of algebraic curves over  $\mathbb{F}_{p^2}$  starting from a smooth family  $f : X \rightarrow U$  of elliptic curves over an open subset  $U \subset \mathbb{P}^1$ . We used that the  $i$ -th derived étale direct image of the constant sheaf  $\mathbb{Z}/\ell^n\mathbb{Z}$  corresponds to a local system  $\mathcal{V}_n$  on  $U$ , and that one can define a fiberwise *projectivisation*  $P_n(\mathcal{V}_n)$  of this local system, which is an étale scheme  $U_n$  over  $U$  (see section 2). The challenge of this approach is to bound the genus of a connected component of  $U_n$ .

In this paper we use a special case of this construction, that lifts to characteristic zero; thus we can bound the ramification and the genus in the tower using transcendental methods. We give an example of a good tower coming from a family  $V_\lambda$  of K3 surfaces (see section 3). The family that we consider is the mirror-dual family to  $\mathbb{P}^3$ . It was studied, among others, by Dolgachev [Dolg], Batyrev [Bat], Coates–Corti–Galkin–Kasprzyk [CCGK]. Namely, Batyrev proved that  $V_\lambda$  is mirror-dual to  $\mathbb{P}^3$ , and Dolgachev computed global monodromy of  $V_\lambda$ . We also use explicit description of local monodromy operators given in [PS]. Altogether this information is enough to describe the corresponding tower.

**Theorem 1.** *There exists a tower  $C_\bullet$  over  $k = \mathbb{F}_{p^2}$  that is optimal for  $p = 3$ , and is good if  $p \equiv 3(4)$ .*

Other similar families that can be used as an input data for our construction can be found in [Gol] and [CCGK].

## 2. TOWERS OF ALGEBRAIC CURVES

In this section we give a modified version of the construction of a tower of algebraic curves from [Ry].

First we recall a generalization of the projectivization of an  $\mathbb{F}_\ell$ -vector space. Put  $\Lambda_n = \mathbb{Z}/\ell^n\mathbb{Z}$ . Let  $V$  be a finitely generated  $\Lambda_n$ -module. The set

$$V^* = \{v \in V \mid \ell^{n-1}v \neq 0\}$$

has a natural action of the group of invertible elements  $\Lambda_n^*$ . We say that the set

$$P_n(V) = V^*/\Lambda_n^*$$

is *the projectivisation of  $V$* .

---

<sup>0</sup>This work is supported by the Russian Science Foundation under grant 18-11-00121

*Key words and phrases.* optimal tower, finite field.

**Lemma 1.** [Ry, Lemma 1] *Suppose that  $V_n$  is a  $\Lambda_n$ -module, and  $V_{n-1}$  is a  $\Lambda_{n-1}$ -module. The group  $V_{n-1}$  is a  $\Lambda_n$ -module in an obvious way. Let  $\varphi : V_n \rightarrow V_{n-1}$  be a homomorphism such that  $\ell \ker \varphi = 0$ . Then there is an induced map*

$$P(\varphi) : P_n(V_n) \rightarrow P_{n-1}(V_{n-1}).$$

Let  $\mathcal{U}$  be a smooth and connected scheme over  $\mathbb{Z}$  of relative dimension one, and let  $f : \mathcal{X} \rightarrow \mathcal{U}$  be a smooth family of algebraic varieties over  $\mathcal{U}$ . Suppose that there is an action of a finite group  $H$  on  $\mathcal{X}$ , and that  $f$  is  $H$ -equivariant. According to [Mil80, VI.4.2], the  $H$ -invariants of the étale direct image

$$\mathcal{V}_n = (R_{\text{ét}}^i f_{U*}(\Lambda_n))^H,$$

is a locally constant finite sheaf of  $\Lambda_n$ -modules on  $\mathcal{U}$ .

According to [Mil80, V.1.1], the functor  $\mathcal{V}_n^*$  given by

$$T \mapsto \mathcal{V}_n(T)^*$$

is representable by a finite étale scheme over  $\mathcal{U}$  with a free action of the group  $\Lambda_n^*$ . Thus the quotient  $\tilde{\mathcal{U}}_n$  by this action represents the sheaf  $P_n(\mathcal{V}_n)$ . For each  $n$  we have a natural étale morphism

$$\pi_n : \tilde{\mathcal{U}}_n \rightarrow \tilde{\mathcal{U}}_{n-1}.$$

Let  $K$  be a field, and let  $\bar{a} \in \mathcal{U}(\bar{K})$  be a point of  $\mathcal{U}$ . The stalk  $(\mathcal{V}_n)_{\bar{a}}$  of  $\mathcal{V}_n$  is isomorphic to  $H_{\text{ét}}^i(\bar{X}_{\bar{a}}, \Lambda_n)^H$ . According to [Ry, Lemma 2] the fiber of  $\tilde{\mathcal{U}}_n$  over  $\bar{a} \in \mathcal{U}(\bar{K})$  is naturally isomorphic to  $P_n((\mathcal{V}_n)_{\bar{a}})$ . In what follows we assume that the module  $H_{\text{ét}}^i(\bar{X}_{\bar{a}}, \mathbb{Z}_\ell)^H$  is free.

Let  $\bar{a} \in \mathcal{U}(\mathbb{C})$ , and let  $\pi_1(\mathcal{U}, \bar{a})$  be the fundamental group of  $\mathcal{U}$ .

**Proposition 1.** *There is a one to one correspondence between orbits of the action of  $\pi_1(\mathcal{U}, \bar{a})$  on  $P_n(H_{\text{ét}}^i(\bar{X}_{\bar{a}}, \Lambda_n))$  and connected components of  $\tilde{\mathcal{U}}_n$ . The degree of the connected component over  $\mathcal{U}$  is equal the cardinality of the corresponding orbit.*

**Proof .** Let  $K$  be a field of functions of  $\mathcal{U}$ . A connected component  $\mathcal{L}$  of  $\tilde{\mathcal{U}}_n$  is uniquely determined by its field of functions. The group  $\pi_1(\mathcal{U}, \bar{a})$  is isomorphic to the quotient of the Galois group  $\text{Gal}(K)$ . Finite field extensions of  $K$  correspond to transitive actions of  $\text{Gal}(K)$  on finite sets (see section 2.3 of [Ry]). Finally, the degree of  $\mathcal{L}$  over  $\mathcal{U}$  is equal to the degree of the corresponding field extension.

We assume that  $U = \mathcal{U} \otimes k$  is non-empty and geometrically connected. If  $\mathcal{U}_n$  is a connected component of  $\tilde{\mathcal{U}}_n$ , then  $\mathcal{U}_{n-1} = \pi_n(\mathcal{U}_n)$  is a connected component of  $\tilde{\mathcal{U}}_{n-1}$ . Choose a sequence of connected subschemes  $\mathcal{U}_n \subset \tilde{\mathcal{U}}_n$  such that  $\mathcal{U}_{n-1} = \pi_n(\mathcal{U}_n)$ . Let  $C = C_0$  be a smooth projectivisation of  $\mathcal{U} \otimes k$ , and let  $C_n$  be a regular projectivisation of  $\mathcal{U}_n \otimes k$ . Note that every connected component  $Y$  of  $C_n$  is a smooth algebraic curve over the algebraic closure of  $k$  in  $k(Y)$ . Put  $W_n = \mathcal{U}_n \otimes \mathbb{C}$ . The smooth projectivisation  $\mathcal{C}_n$  of  $W_n$  is an algebraic curve over  $\mathbb{C}$ . We get the following diagram:

$$\begin{array}{ccccccccc} \downarrow & & \downarrow & \downarrow & \downarrow & & \downarrow & & \downarrow \\ C_n & \supset & U_n & \mathcal{U}_n & W_n & \subset & \mathcal{C}_n & & \\ \downarrow & & \downarrow & \downarrow & \downarrow & & \downarrow & & \\ \vdots & & \vdots & \vdots & \vdots & & \vdots & & \\ \downarrow & & \downarrow & \downarrow & \downarrow & & \downarrow & & \\ C_1 & \supset & U_1 & \mathcal{U}_1 & W_1 & \subset & \mathcal{C}_1 & & \\ \downarrow & & \downarrow & \downarrow & \downarrow & & \downarrow & & \\ C & \supset & U & \mathcal{U} & W & \subset & \mathcal{C} & & \end{array}$$

Let  $a \in U(k)$ . We say that a smooth variety  $Y$  over  $k$  is *strongly supersingular in degree  $i$*  if the Frobenius action on  $H_{\text{ét}}^i(\bar{Y}, \mathbb{Z}_\ell)$  is the multiplication by  $q^{i/2}$  or  $-q^{i/2}$ .

**Proposition 2.** *Assume that  $C_n$  is connected, and there exists  $a \in U(k)$  such that  $X_a$  is strongly supersingular in degree  $i$ . Then  $C_n$  is a smooth, geometrically irreducible curve over  $\mathbb{F}_p$ , and the fiber of the morphism  $C_n \rightarrow C_0$  over  $a$  is split over  $k$ .*

**Proof .** According to [Ry, Corollary 3] the statement is true after the base change to  $k$ ; thus  $C_n$  is smooth and geometrically irreducible over  $\mathbb{F}_p$  as well.

The ramification of  $C_n$  over  $\mathcal{C}$  can be computed in terms of the local monodromy of the family  $f \otimes \mathbb{C}$ . Fix a point  $\bar{a} \in W(\mathbb{C})$ . Let  $y \in \mathcal{C}(\mathbb{C}) - W(\mathbb{C})$ , and let  $G_y$  be the stabilizer of  $y$  in  $\pi_1(W, \bar{a})$ .

**Proposition 3.** [Ry, Proposition 1] *There is a bijection between orbits of the action of  $G_y$  on  $P_n((\mathcal{V}_n)_{\bar{a}})$  and points  $x \in \mathcal{C}_n(\mathbb{C})$  over  $y$ . The ramification index of  $x$  is equal to the cardinality of the corresponding orbit.*

**Remark .** Let  $k$  be an extension of odd degree of  $\mathbb{F}_p$ . Assume that the general fiber  $Y$  of our family is a K3 surface. Then, by a result of Artin [Ar, (6.8)], there are no strongly supersingular fibers in our family. This observation force us to look for a weaker supersingularity condition such that one still could prove an analog of Proposition 2 for  $k = \mathbb{F}_p$ .

### 3. A FAMILY OF K3 SURFACES

In this section we prove Theorem 1.

Projection  $\pi$  from

$$X = \{x_0^4 + x_1^4 + x_2^4 + x_3^4 + 4\lambda x_0 x_1 x_2 x_3 = 0\} \subset \mathbb{A}_\lambda^1 \times \mathbb{P}_x^3$$

to  $\mathbb{A}_\lambda^1$  is a pencil of quartic surfaces  $X_\lambda$ , and its restriction to  $\mathcal{U} = \text{Spec } \mathbb{Z}[\lambda, \frac{1}{2(\lambda^4-1)}]$  is smooth.

Let  $i$  be a primitive 4th root of unity, considered as a generator of a group scheme  $\mu_4 = \text{Spec } \mathbb{Z}[i]/(i^4 - 1)$ . Maps

$$(x_0 : x_1 : x_2 : x_3) \rightarrow (x_0 : ix_1 : x_2 : -ix_3) \text{ and } (x_0 : x_1 : x_2 : x_3) \rightarrow (ix_0 : x_1 : -ix_2 : x_3)$$

generate a non-effective action of a group scheme  $H = \mu_4 \times \mu_4$  on  $X$ . The action of  $H$  commutes with the projection  $\pi$ , hence it induces an  $H$ -action on every fiber.

**Theorem 2.** [Dolg, Theorem 7.1, Theorem 7.7, Theorem 8.2] *The general fiber of the quotient family  $(X_\lambda \otimes \mathbb{C})/H$  has 6 rational double points of type  $A_3$ . The resolution of singularities is the family  $V_\lambda$  of Kummer surfaces associated to the product  $E_1 \times E_2$ , where  $E_1$  and  $E_2$  are elliptic curves with 2-isogeny between them. The monodromy group  $G$  of this family is the modular Fricke group of level 2, i.e.,  $G$  is the subgroup of  $\text{PGL}(2, \mathbb{R})$  generated by  $\Gamma_0(2)$  and by*

$$\begin{pmatrix} 0 & -1/\sqrt{2} \\ \sqrt{2} & 0 \end{pmatrix}.$$

Exceptional lines of the resolution of 6 rational double points of type  $A_3$  generate a sublattice of rank 18 in  $\text{NS}(V_\lambda)$ . It follows that the rank of  $H$ -invariants  $H^2(X_\lambda \otimes \mathbb{C}, \mathbb{Z})^H$  is 4. Moreover, a hyperplane section gives a  $G$ -invariant sublattice of rank 1. The orthogonal complement  $T$  is naturally isomorphic to the lattice  $T$  of quadratic forms

$$\{\alpha x^2 + 2\sqrt{2}\beta xy + \gamma y^2 \mid \alpha, \beta, \gamma \in \mathbb{Z}\}$$

endowed with the quadratic form  $Q(\alpha, \beta, \gamma) = 2(2\beta^2 - \alpha\gamma)$  [Dolg, Theorem 7.1]. The action of  $G$  on  $T$  is induced from the following twisted action  $\rho$  of  $\text{PGL}(2, \mathbb{R})$  on  $T \otimes \mathbb{R}$ :

$$\rho \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1/\sqrt[4]{2} & 0 \\ 0 & \sqrt[4]{2} \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1/\sqrt[4]{2} & 0 \\ 0 & \sqrt[4]{2} \end{pmatrix}^{-1}.$$

The zero locus of the quadratic form on  $T$  is exactly the set of degenerate forms. Let  $D_n \subset P_n(T/\ell^n T)$  be the projectivisation of the cone of degenerate forms. It is straightforward to check

that the action of  $G$  on  $D_n$  is transitive for all  $n$ . It follows from Proposition 1 that the curves  $\mathcal{C}_n$  are connected; therefore  $C_n$  are connected as well. Clearly,  $D_1$  is a set of points on a conic in  $P_1(T/\ell T)$ ; thus the degree of  $\mathcal{C}_1$  over  $\mathcal{C}_0$  is equal to  $\ell + 1$ . If  $n > 1$ , then  $\pi_n$  induces a morphism  $\mathcal{C}_n \rightarrow \mathcal{C}_{n-1}$  of degree  $\ell$ .

**Lemma 2.** *For the genus  $g_n$  of  $\mathcal{C}_n$  we have*

$$g_n \leq (\ell + 1)\ell^{n-1}/2.$$

**Proof .** The family  $X_\lambda$  is mirror-dual to a family  $V_\lambda$  [Bat, Corollary 5.5.6]. In turn, the singular fibers and monodromy operators for such families are known. According to [PS, Period sequence 12] the mirror-dual family to  $\mathbb{P}^3$  has 4 singular fibers such that the monodromy operator is an involution, and one fiber with maximal unipotent monodromy. The Hurwitz formula now gives

$$2g_n - 2 = -2(\ell + 1)\ell^{n-1} + R_0 + R,$$

where  $R_0 \leq \deg \pi_n = (\ell + 1)\ell^{n-1}$  is the ramification degree of the point with maximal unipotent monodromy, and  $R$  is the ramification degree in other points. We now bound  $R$  using Proposition 3. Since the monodromy operator is an involution, there are at least  $(\ell + 1)\ell^{n-1}/2$  orbits of degree 2 on the set  $D_n$ . This gives

$$R \leq 4((\ell + 1)\ell^{n-1}/2) = 2(\ell + 1)\ell^{n-1}.$$

This proves the lemma.

By a result of Tsfasman [Ts, Example 5.9], the quartic

$$x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0$$

is strongly supersingular in degree 2 over  $k$ , when  $p \equiv 3(4)$ . According to Proposition 2, there exists at least one split fiber of  $\pi_n$  over  $k$ . We finally get

$$\lim_{n \rightarrow \infty} \frac{|C_n(k)|}{g_n} \geq 2.$$

This proves Theorem 1.

## REFERENCES

- [Ar] M. Artin, “Supersingular K3 surfaces”, *Annales scientifiques de l’École Normale Supérieure, Serie 4*, **7**:4, (1974), pp. 543-567.
- [Bat] V. Batyrev, “Dual polyhedra and mirror symmetry for Calabi–Yau hypersurfaces in toric varieties”, *J. Alg. Geom.*, **3**, (1994), 493-545.
- [CCGK] T. Coates, A. Corti, S. Galkin, and A. Kasprzyk, “Quantum periods for 3-dimensional Fano manifolds”, *Geom. Topol.*, **20**:1, (2016), 103-256.
- [PS] <http://coates.ma.ic.ac.uk/3DMinkowskiPeriodSequences.html>
- [Dolg] I. V. Dolgachev, “Mirror symmetry for lattice polarized K3 surfaces”, *Algebraic geometry 4, Itogi Nauki i Tekhniki. Ser. Sovrem. Mat. Pril. Temat. Obz.*, **33**, VINITI, Moscow, 2001, 2071; *J. Math. Sci.*, **81**:3 (1996), 2599-2630.
- [Gol] V. V. Golyshev, “Minimal Fano threefolds: Exceptional sets and vanishing cycles”, *Doklady Mathematics*, 2009, **79**:1, 1620.
- [Mil80] J. Milne, “*Étale cohomology*”, Princeton Mathematical Series 33, Princeton University Press, (1980).
- [Ry] S. Rybakov, “Families of algebraic varieties and towers of algebraic curves over finite fields”, *Math. Notes*, **104**:5 (2018), 712719.
- [Ts] M.A. Tsfasman, “Nombre de points des surfaces sur un corps fini”, *Algebraic Geometry and Coding Theory, de Gruyter, Berlin*, pp.209-224 (1996).

HSE UNIVERSITY, RUSSIAN FEDERATION

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS OF THE RUSSIAN ACADEMY OF SCIENCES, AG LABORATORY, HSE UNIVERSITY, RUSSIAN FEDERATION

*E-mail address:* rybakov.sergey@gmail.com