

Address: Professur für Mathematik ETH Zürich, HG J 16.2 Rämistrasse 101, Zürich 8092, Switzerland , E-mail: arkadij.bojko@math.ethz.ch

## NON-COMMUTATIVE COUNTING AND STABILITY

ARKADIJ BOJKO AND GEORGE DIMITROV

**ABSTRACT.** The second author and Katzarkov introduced categorical invariants based on counting of full triangulated subcategories in a given triangulated category  $\mathcal{T}$ , and they demonstrated different choices of additional properties of the subcategories being counted, in particular - an approach to make non-commutative counting in  $\mathcal{T}$  dependable on a stability condition  $\sigma \in \text{Stab}(\mathcal{T})$ . In this paper, we focus on this approach. After recalling the definitions of a *stable non-commutative curve in  $\mathcal{T}$*  and related notions, we prove a few general facts and study an example:  $\mathcal{T} = D^b(Q)$ , where  $Q$  is the acyclic triangular quiver. In previous papers, it was shown that there are two non-commutative curves of non-commutative genus 1 and infinitely many non-commutative curves of non-commutative genus 0 in  $D^b(Q)$ . Our studies here imply that for an open and dense subset in  $\text{Stab}(D^b(Q))$  the stable non-commutative curves in  $D^b(Q)$  are finite. This paper also introduces counting of semistable derived points and shows that the corresponding invariants are finite on an open dense subset of  $\text{Stab}(D^b(Q))$ .

### CONTENTS

1.	Introduction	1
2.	Notations	6
3.	Triangulated categories and Bridgeland stability conditions	7
4.	Non-commutative curve counting. Dependence on a stability condition	9
5.	Some remarks for $\sigma$ -exceptional collections and $C_{l,\sigma\sigma}(\mathcal{T})$	12
6.	The example: Stable non-commutative curves in $D^b(Q)$	15
7.	The set of stabilities, where $ C_{0,\sigma\sigma}(\mathcal{T})  < \infty$ is open and dense in $\text{Stab}(\mathcal{T})$	35
8.	Walls and chambers for $C_{1,\sigma\sigma}$	37
	Appendix A. Details on $C_{l,\sigma\sigma}(\mathcal{T})$ , $C_{l,\sigma}(\mathcal{T})$ as $\sigma$ varies in $\text{Stab}(\mathcal{T})$ and $l \geq -1$	41
	References	45

### 1. INTRODUCTION

**1.1. Overview.** Orlov’s reconstruction theorem states that for a smooth variety with an ample (anti-)canonical divisor, one can uniquely recover the variety from its bounded derived category. This motivated the point of view of non-commutative derived geometry through considering general triangulated categories as non-commutative varieties. An Example of this appeared in Kontsevich–Rosenberg [11, §3.3], where it is conjectured that the derived category of the non-commutative projective space  $N\mathbb{P}^{l-1}$  is the derived category of the wild Kronecker quiver with  $l$  arrows:

$$K(l) = 1 \begin{array}{c} \curvearrowright \\ \vdots \\ \curvearrowleft \end{array} 2$$

$l$ -times

1

Following this, the second author and Katzarkov defined non-commutative counting in [5] as taking cardinalities of certain sets of fully faithful embeddings of a fixed triangulated subcategory  $\mathcal{A}$  into a larger one  $\mathcal{T}$ . We will focus here on the cases  $\mathcal{A} = N\mathbb{P}^l$  or  $\mathcal{A} = D^b(A_1)$ . The former will be called a *non-commutative curve* and the latter a *derived point*. One can additionally introduce a dependence on a parameter  $\sigma \in \text{Stab}(\mathcal{T})$ , where  $\text{Stab}(\mathcal{T})$  is the manifold of stability conditions of Bridgeland [2]. This leads to counting of stable non-commutative curves and derived points as suggested in [5, Section 12.4].

Our main goal is to study this for non-trivial example of  $\mathcal{T} = D^b(Q)$ , where  $Q$  is the quiver

$$(1) \quad Q = \begin{array}{ccc} & & z \\ & \nearrow & \searrow \\ x & \longrightarrow & y \end{array}$$

Knowing which derived points are semistable for each stability condition, we can determine the stability on non-commutative curves. It then turns out that to obtain a complete answer, we only need to study the behavior of stable exceptional objects in detail.

We thus describe explicitly the wall and chamber structure of the space  $\text{Stab}(\mathcal{T})$ , where non-commutative curves or derived points become stable. The most important observation is that in our example, the set of  $\sigma \in \text{Stab}(\mathcal{T})$ , such that the number of semistable curves is finite is dense. This motivates Conjecture 1. As this does not apply verbatim to derived points, we propose an alternative: we count stable derived points lying outside of stable non-commutative curves. Then an analogous property is satisfied (see Conjecture 2, Theorem 9). We now give a more detailed summary of the definitions, previous work in this area and our new results.

**1.2. Summary of the results.** The idea of non-commutative counting in [5] is to count elements in a set  $C_{\mathcal{A},P}^\Gamma(\mathcal{T})$  defined as follows: given two triangulated categories  $\mathcal{A}, \mathcal{T}$ , a subgroup  $\Gamma \subset \text{Aut}(\mathcal{T})$ , and choosing some additional restrictions  $P$  of fully faithful exact functors, the set  $C_{\mathcal{A},P}^\Gamma(\mathcal{T})$  consists of equivalence classes of fully faithful exact functors from  $\mathcal{A}$  to  $\mathcal{T}$  satisfying  $P$  with two functors  $F, F'$  being equivalent iff  $F \circ \alpha \cong \beta \circ F'$  for some  $\alpha \in \text{Aut}(\mathcal{A})$ ,  $\beta \in \Gamma$  (see Definition 18). Various choices of  $\mathcal{A}, \Gamma, P, \mathcal{T}$  have been shown to result in finite sets  $C_{\mathcal{A},P}^\Gamma(\mathcal{T})$  in [5, 6, 10]. In this paper, we let  $P$  depend on  $\sigma \in \text{Stab}(\mathcal{T})$  and demonstrate an example with fixed  $\mathcal{T}$  and  $\mathcal{A}$ , in which for some points  $\sigma$  in the parameter space the set  $C_{\mathcal{A},P(\sigma)}^{\{\text{Id}\}}(\mathcal{T})$  is an infinite set which was already studied in [6]. However, for a generic parameter  $\sigma$  the set  $C_{\mathcal{A},P(\sigma)}^{\{\text{Id}\}}(\mathcal{T})$  is finite.

When the categories  $\mathcal{T}, \mathcal{A}$  are  $\mathbb{K}$ -linear, then choosing the following property  $P$  imposed on the functors: *to be  $\mathbb{K}$ -linear*, and restricting  $\Gamma$  to be a subgroup of the group of  $\mathbb{K}$ -linear auto-equivalences  $\text{Aut}_{\mathbb{K}}(\mathcal{T})$  results in the set  $C_{\mathcal{A},P}^\Gamma(\mathcal{T})$ , which is denoted by  $C_{\mathcal{A},\mathbb{K}}^\Gamma(\mathcal{T})$ .

As already mentioned, we denote  $D^b(\text{Rep}_{\mathbb{K}}(K(l+1)))$  by  $N\mathbb{P}^l$  for  $l \geq -1$  (see Section 2 for basic notations). In [5],  $N\mathbb{P}^l$  is selected as a first choice for  $\mathcal{A}$ , being non-trivial but a well studied category. For a  $\mathbb{K}$ -linear category  $\mathcal{T}$  and any  $l \geq -1$  the set  $C_{N\mathbb{P}^l,\mathbb{K}}^\Gamma(\mathcal{T})$  is denoted by  $C_l^\Gamma(\mathcal{T})$  and is referred to as *the set of non-commutative curves of non-commutative genus  $l$  in  $\mathcal{T}$  modulo  $\Gamma$* . On the other hand, in [6] the theory of intersection of the entities being counted is initiated and, in that respect, the elements in  $C_{D^b(pt),\mathbb{K}}^{\{\text{Id}\}}(\mathcal{T})$  are referred to as *derived points in  $\mathcal{T}$* .

A derived point in a proper  $\mathbb{K}$ -linear  $\mathcal{T}$  with a dg-enhancement is just a full triangulated subcategory generated by an exceptional object, whereas a non-commutative curve of non-commutative genus  $l$  is the same as a full triangulated subcategory generated by a strong exceptional pair  $(A, B)$  with  $\text{hom}(A, B) = l + 1$  (see [6, Proposition 5.5]).

For the rest of the introduction, we let  $\mathcal{T}$  be as in the previous paragraph. Let  $\Gamma \subset \text{Aut}(\mathcal{T})$  be the trivial subgroup  $\Gamma = \{\text{Id}\}$  and we will omit writing  $\{\text{Id}\}$ . We will now explain  $P(\sigma)$  (see also Definition 21 and Remark 29).

The notion of stability condition on a triangular category  $\mathcal{T}$ , introduced by T. Bridgeland in [2], has its roots in string theory and mirror symmetry. In [2], the set of so called locally finite stability conditions is equipped with a structure of a complex manifold, denoted by  $\text{Stab}(\mathcal{T})$ . Each  $\sigma \in \text{Stab}(\mathcal{T})$  distinguishes a set of non-zero objects in  $\mathcal{T}$  and labels them with real numbers, these objects are referred to as  $\sigma$ -semistable objects and the real number attached to such an object  $X$  is denoted by  $\phi_\sigma(X) \in \mathbb{R}$  and called the phase of  $X$ . We denote the set of semistable objects by  $\sigma^{ss}$ . Since the set  $\sigma^{ss}$  is invariant under the translation functor, it follows that for a derived point  $\mathcal{A} \in C_{D^b(pt),\mathbb{K}}(\mathcal{T})$  either no or all exceptional objects in  $\mathcal{A}$  are also in  $\sigma^{ss}$ , thus  $\sigma^{ss}$  naturally distinguishes those derived points satisfying the latter. The set of derived points, distinguished by  $\sigma$ , will be denoted by  $C_{D^b(pt),\sigma\sigma}(\mathcal{T}) \subset C_{D^b(pt),\mathbb{K}}(\mathcal{T})$  (see Definition 24).

We will be mainly interested in two subsets of  $C_l(\mathcal{T})$ , denoted by  $C_{l,\sigma\sigma}(\mathcal{T})$  and  $C_{l,\sigma}(\mathcal{T})$ , whose elements are referred to as  $\sigma$ -stable and, respectively  $\sigma$ -semistable non-commutative curves, as defined in [5, Section 12.4]. **A non-commutative curve  $\mathcal{A} \in C_l(\mathcal{T})$  is in  $C_{l,\sigma}(\mathcal{T})$  if and only if infinitely many derived points in it are also in  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$ , and it is in  $C_{l,\sigma\sigma}(\mathcal{T})$  if and only if all derived points in  $\mathcal{A}$  are also in  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$  (see Lemma 26).** Details on the definitions and basic properties are given in Section 4. In particular, for  $l \leq 0$  we obtain  $C_{l,\sigma}(\mathcal{T}) = \emptyset$ , whereas for  $l \geq 1$   $C_{l,\sigma\sigma}(\mathcal{T}) \subset C_{l,\sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$  (see Remark 28). For any  $\mathcal{A} \in C_l(\mathcal{T})$  the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : \mathcal{A} \in C_{l,\sigma\sigma}(\mathcal{T})\}$  is a closed subset in  $\text{Stab}(\mathcal{T})$  (see Corollary 27), which makes it the correct object to study. Because the set  $C_l(\mathcal{T})$  can often become infinite, the dependence on stability conditions is introduced to cut down to a finite number of non-commutative (semi-)stable curves and define integer invariants counting these. This is a usual approach in enumerative theories (e.g. Donaldson–Thomas invariants and Gromov–Witten invariants) where one restricts to moduli spaces of stable objects carrying well-defined virtual fundamental classes. In this paper, we also give an example, where stable  $\neq$  semistable (see Proposition 8) showing a further similarity to other enumerative theories. This naturally motivates the following question.

**Question 1.** Let  $\mathcal{T}$  be a triangulated  $\mathbb{K}$  category with a phase gap ([5, Definition 4.6]) and  $\text{rk}(K^0(\mathcal{T})) < \infty$ . Does there exist a dense open subset of  $\Sigma \subset \text{Stab}(\mathcal{T})$ , such that  $|C_{l,\sigma\sigma}^\Gamma(\mathcal{T})| < \infty$  for all  $\sigma \in \Sigma$  and  $\Gamma$  a subgroup of  $\text{Aut}(\mathcal{T})$ ?

We now formulate a conjecture for a more approachable situation. In [5, 12.1] the first author and Katzarkov define a new invariant

$$\left\{ \begin{array}{c} \text{triangulated} \\ \text{categories} \\ \text{with a phase gap} \end{array} \right\} \xrightarrow{\dim_{nc}} [0, +\infty],$$

which gives the following vanishing criterion:

$$\dim_{nc}(\mathcal{T}) \leq n \implies C_l(\mathcal{T}) = \emptyset \text{ for } l > n.$$

They show  $\dim_{nc}(D^b(Q)) = 1$  for an acyclic quiver  $Q$  if and only if it is affine. In this case, we expect the following to hold.

**Conjecture 1.** Let  $Q$  be an extended acyclic Dynking quiver and  $\mathcal{T} = D^b(Q)$ , then the set of stability conditions  $\sigma \in \text{Stab}(\mathcal{T})$  such that  $|C_{l,\sigma\sigma}| < \infty$  is open and dense.

This paper contains an example of such  $Q$ , where  $C_0(\mathcal{T})$  is infinite however the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |\cup_{l \geq -1} C_{l,\sigma\sigma}(\mathcal{T})| < \infty\}$  is open and dense subset in  $\text{Stab}(\mathcal{T})$ . Furthermore, counting the elements of  $C_{0,\sigma\sigma}(\mathcal{T})$  as  $\sigma$  varies in this open and dense subset results in all the numbers in  $\{1\} \cup 2\mathbb{Z}_{\geq 0}$ . The same approach fails when applied to derived points as shown in Proposition 7. This motivates us to introduce a new approach to counting semistable exceptional objects in terms of counting derived points. In Definition 25, we define the set  $\mathfrak{D}_\sigma \subset C_{D^b(pt),\sigma\sigma}$  of *stable derived points outside of stable curves of positive genus*. This set consists of all elements  $p \in C_{D^b(pt),\sigma\sigma}$  which are not a derived point of a curve  $c \in C_{l,\sigma\sigma}$ .

We can pose a question analogous to the Question 1. Moreover, Theorem 9 suggests the following.

**Conjecture 2.** Let  $Q$  be an extended acyclic Dynking quiver and  $\mathcal{T} = D^b(Q)$ , then the set of stability conditions  $\sigma \in \text{Stab}(\mathcal{T})$  such that  $|\mathfrak{D}_\sigma| < \infty$  is open and dense.

**1.3. Some general results and a complete example.** Before focusing on our main example, we show in Section 5 general situations ensuring non-trivial behavior of  $C_{l,\sigma\sigma}(\mathcal{T})$ ,  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$  as  $\sigma$  varies in  $\text{Stab}(\mathcal{T})$ . To that end we recall the notion of  $\sigma$ -exceptional collection, introduced in [7]. The following then summarizes multiple results.

**Proposition 3** (Cor. 34 and Prop. 36). *Let  $\mathcal{T}$  be a triangulated category with an enhancement and  $\sigma \in \text{Stab}(\mathcal{T})$ . Let  $\mathcal{E} = (E_0, \dots, E_n)$  be a full  $\sigma$ -exceptional collection such that*

- (i)  $\phi(E_{i-1}) < \phi(E_i)$  for all  $i = 1, \dots, n$ , then  $C_{D^b(pt),\sigma\sigma} = \{\langle E_0 \rangle, \langle E_1 \rangle, \dots, \langle E_n \rangle\}$ . In particular,  $C_{l,\sigma\sigma}(\mathcal{T}) = C_{l,\sigma}(\mathcal{T}) = \emptyset$  for  $l \geq 0$ .
- (ii) such that for some  $0 \leq i < n$  and  $l \geq 0$ , we have  $\text{hom}^1(E_i, E_{i+1}) = l + 1$ ,  $\text{hom}^j(E_i, E_{i+1}) = 0$  for  $j \in \mathbb{Z} \setminus \{1\}$  and  $\phi_\sigma(E_i) < \phi_\sigma(E_{i+1})$  then  $\langle E_i, E_{i+1} \rangle \in C_{l,\sigma\sigma}(\mathcal{T})$ .

In Corollary 37 this result is presented in a slightly different form. In addition to this. To study our main example, we will use in this paper more subtle tools from [7, 8] to ensure that certain elements in  $C_l(\mathcal{T})$  are in  $C_{l,\sigma\sigma}(\mathcal{T})$  or in  $C_{l,\sigma}(\mathcal{T})$  for some  $\sigma \in \text{Stab}(\mathcal{T})$ .

In [5, Proposition 12.13] the behavior of  $C_{l,\sigma}(N\mathbb{P}^k)$  and  $C_{l,\sigma\sigma}(N\mathbb{P}^k)$  was described for any  $l \geq -1$ ,  $\sigma \in \text{Stab}(N\mathbb{P}^k)$ ,  $k \geq 1$ . In this case  $C_l(N\mathbb{P}^k)$  is non-trivial only for  $l = k$ ,  $C_k(N\mathbb{P}^k)$  has only one element and it turns out that  $C_{k,\sigma}(N\mathbb{P}^k) = C_{k,\sigma\sigma}(N\mathbb{P}^k)$  for each  $\sigma$ . We see therefore that Conjecture 1 and 2 are trivially satisfied.

Here we focus on another example, namely From now on,  $\mathcal{T} = D^b(Q)$ , where  $Q$  is as in (1). Let us recall some results from [6, 7, 8]. The derived points in  $\mathcal{T}$  are ([8, Subsection 3.2, Remark 3.6]):

$$(2) \quad C_{D^b(pt),\mathbb{K}}(\mathcal{T}) = \{\langle a^m \rangle, \langle b^m \rangle : m \in \mathbb{Z}\} \cup \{\langle M \rangle, \langle M' \rangle\},$$

where  $a^m, b^m, M, M'$  are exceptional objects,  $M, M'$  is the only couple of exceptional objects in  $\text{Rep}_{\mathbb{K}}(Q)$  such that  $\text{Ext}^1(M, M') \neq 0$ ,  $\text{Ext}^1(M', M) \neq 0$  (see the beginning of [7, Subsection 2.3]).

**Proposition 4.** [6, Proposition 8.1, (34), (35), (36), (37)]. *For  $l \notin \{0, 1\}$  we have  $C_l(\mathcal{T}) = \emptyset$  and:*

$$(3) \quad C_1(\mathcal{T}) = \{\langle a^m, a^{m+1} \rangle = \langle M \rangle^\perp, \langle b^m, b^{m+1} \rangle = \langle M' \rangle^\perp\}$$

$$(4) \quad C_0(\mathcal{T}) = \{\langle M', a^m, b^{m+1} \rangle = \langle a^{m+1} \rangle^\perp : m \in \mathbb{Z}\} \cup \{\langle M, b^m, a^m \rangle = \langle b^{m+1} \rangle^\perp : m \in \mathbb{Z}\}.$$

*We will denote  $A = \langle a^m, a^{m+1} \rangle$ ,  $B = \langle b^m, b^{m+1} \rangle$ , and thus  $C_1(\mathcal{T}) = \{A, B\}$ .<sup>1</sup>*

*Denoting  $\alpha^m = \langle M', a^m, b^{m+1} \rangle$ ,  $\beta^m = \langle M, b^m, a^m \rangle$  we can write  $C_0(\mathcal{T}) = \{\alpha^m, \beta^m : m \in \mathbb{Z}\}$ .*

<sup>1</sup>here  $(a^m, a^{m+1})$ ,  $(b^m, b^{m+1})$  are exceptional pairs

In [5, Proposition 12.14] it was suggested and in [3] proved that when  $\sigma$  varies in  $\text{Stab}(\mathcal{T})$  the set  $C_{1,\sigma}(\mathcal{T})$  takes all subsets in  $C_1(\mathcal{T})$ . However, only  $C_{0,\sigma\sigma}(\mathcal{T})$  is meaningful, since we already mentioned that  $C_{0,\sigma}(\mathcal{T}) = \emptyset$  for any  $\sigma$  and any  $\mathcal{T}$ . Additionally, for any  $l \geq -1$  we expect that studying  $C_{l,\sigma\sigma}(\mathcal{T})$  is more meaningful than  $C_{l,\sigma}(\mathcal{T})$ , for example Corollary 27 does not seem to hold for  $C_{l,\sigma}(\mathcal{T})$ . These are reasons to suggest in [5, Remark 12.15] that not only  $C_{1,\sigma}(\mathcal{T})$  but also  $C_{1,\sigma\sigma}(\mathcal{T}) \subset C_{1,\sigma}(\mathcal{T})$  takes all possible subsets in  $C_1(\mathcal{T})$ .

Before giving detailed description of the functions

$$\text{Stab}(\mathcal{T}) \ni \sigma \mapsto \{C_{l,\sigma}(\mathcal{T}), C_{l,\sigma\sigma}(\mathcal{T}), C_{D^b(pt),\sigma\sigma}(\mathcal{T})\}$$

for  $l = 0, 1$  we present some features of the full picture. The following proposition captures the non-trivial behaviour of the invariants leading to wall-crossing.

**Proposition 5.** *As  $\sigma$  varies in  $\text{Stab}(\mathcal{T})$ :*

- (a) *the subset  $C_{1,\sigma\sigma}(\mathcal{T}) \subset C_1(\mathcal{T})$  takes all possible subsets of  $C_1(\mathcal{T})$ ;*
- (b) *the subset  $C_{0,\sigma\sigma}(\mathcal{T}) \subset C_0(\mathcal{T})$  is one of the following subsets and realizes each of them:  $\emptyset$ ,  $\{\alpha^i\}$ ,  $\{\beta^i\}$ ,  $\{\alpha^j, \beta^j : i \leq j \leq k\}$ ,  $\{\alpha^j, \beta^{j+1} : i \leq j \leq k\}$ ,  $\{\alpha^j, \beta^j : i \leq j\}$ ,  $\{\alpha^j, \beta^{j+1} : i \leq j\}$ ,  $\{\alpha^j, \beta^j : j \leq k\}$ ,  $\{\alpha^j, \beta^{j+1} : j \leq k\}$ , where  $i, k$  are arbitrary integers such that  $i \leq k$ . In particular  $\{|C_{0,\sigma\sigma}(\mathcal{T})| : \sigma \in \text{Stab}(\mathcal{T})\} = \{1, +\infty\} \cup 2\mathbb{Z}_{\geq 0}$ .*

Furthermore, we obtain a positive confirmation of Conjecture 1.

**Theorem 6.** *The set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| < \infty\}$  is an open and dense subset in  $\text{Stab}(\mathcal{T})$ .*

One could investigate in more details the walls separating the connected components of the open subset from Theorem 6 and to study wall-crossing. In section 8, we describe in Proposition 66 the global picture of walls and chambers for  $C_{1,\sigma\sigma}(\mathcal{T})$  by gluing the charts from tables (103), (104), (105), (106), (107) and using the results from Proposition 60.

As already mentioned in §1.1, we can recover the above from studying  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$  for each  $\sigma \in \text{Stab}(\mathcal{T})$ . However, counting elements in  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$  is not finite for generic  $\sigma$ . More precisely we have:

**Proposition 7.** *The following subsets are equal and they are open but not dense subsets in  $\text{Stab}(\mathcal{T})$ :*

$$\left\{ \sigma : \left| C_{D^b(pt),\sigma\sigma}(\mathcal{T}) \right| < \infty \right\} = \left\{ \sigma : |C_{1,\sigma}(\mathcal{T})| = 0 \right\} = \left\{ \sigma : |C_{1,\sigma\sigma}(\mathcal{T})| = 0 \right\}.$$

Furthermore,  $\left\{ \left| C_{D^b(pt),\sigma\sigma}(\mathcal{T}) \right| : \sigma \in \text{Stab}(\mathcal{T}) \right\} = \{3, 4, 5, +\infty\}$ .

For  $N\mathbb{P}^l$  we have  $C_{l,\sigma}(N\mathbb{P}^l) = C_{l,\sigma\sigma}(N\mathbb{P}^l)$  for all  $\sigma \in \text{Stab}(\mathcal{T})$  and all  $l \geq 1$ . However, for the case  $\mathcal{T} = D^b(Q)$  we observe a new phenomenon:

**Proposition 8.** *The set  $C_{1,\sigma}(\mathcal{T}) \setminus C_{1,\sigma\sigma}(\mathcal{T})$  can be non-trivial.*

Next, we obtain a positive result towards Conjecture 2.

**Theorem 9.** *The set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |\mathcal{D}_\sigma| < \infty\}$  is an open and dense subset in  $\text{Stab}(\mathcal{T})$ .*

Another result, following from Section 6, is:

**Proposition 10.** *Let  $\sigma \in \text{Stab}(\mathcal{T})$ , then  $C_{1,\sigma\sigma}(\mathcal{T}) = C_1(\mathcal{T}) \iff M, M' \in \sigma^{ss}, \phi(M) = \phi(M')$ . Furthermore,  $C_{1,\sigma\sigma}(\mathcal{T}) = C_1(\mathcal{T}) \iff C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T}) \iff C_{D^b(pt),\sigma\sigma}(\mathcal{T}) = C_{D^b(pt),\mathbb{K}}(\mathcal{T})$ .*

A complete summary of the results from which the above Theorems and Propositions were derived are collected in tables (103), (104), (105), (106), (107) and Proposition 68 in the Appendix A. We use there the notation set in §6.2. How these results glue to a global picture is described in §8.

**Acknowledgements.** The authors are grateful to Ludmil Katzarkov for his interest and encouraging remarks. The first author was supported by Clarendon Fund Scholarship at University of Oxford. The second author was supported by FWF Project P 29178-N35.

## 2. NOTATIONS

We fix a universe and assume that the set of objects and the set of morphisms of any category we consider are elements of this universe.

The shift functor in a triangulated category  $\mathcal{T}$  is designated sometimes by [1].

A *triangulated subcategory* in a triangulated category  $\mathcal{T}$  is a non-empty full subcategory  $\mathcal{B}$  in  $\mathcal{T}$ , s. t. two conditions hold:

- (a)  $Ob(\mathcal{B})[1] = Ob(\mathcal{B})$  ;
- (b) for any  $X, Y \in Ob(\mathcal{B})$  and any distinguished triangle  $X \rightarrow Z \rightarrow Y \rightarrow X[1]$  in  $\mathcal{T}$  the object  $Z$  is also in  $Ob(\mathcal{B})$ .

We write  $\langle S \rangle \subset \mathcal{T}$  for the triangulated subcategory of  $\mathcal{T}$  generated by  $S$ , when  $S \subset Ob(\mathcal{T})$ . We write  $\text{Hom}^i(X, Y)$  for  $\text{Hom}(X, Y[i])$ . In this paper  $\mathbb{K}$  denotes a field. If  $\mathcal{T}$  is  $\mathbb{K}$ -linear triangulated category we write  $\text{hom}^i(X, Y)$  for  $\dim_{\mathbb{K}}(\text{Hom}(X, Y[i]))$ , where  $X, Y \in \mathcal{T}$ .

A  $\mathbb{K}$ -linear triangulated category  $\mathcal{T}$  is called *proper* if  $\sum_{i \in \mathbb{Z}} \text{hom}^i(X, Y) < +\infty$  for any two objects  $X, Y$  in  $\mathcal{T}$ .

When we say that  $\mathcal{T}$  *has an enhancement* we mean enhancement as explained in [12].

An *exceptional object* in a  $\mathbb{K}$ -linear triangulated category is an object  $E \in \mathcal{T}$  satisfying  $\text{Hom}^i(E, E) = 0$  for  $i \neq 0$  and  $\text{Hom}(E, E) = \mathbb{K}$ . We denote by  $\mathcal{T}_{exc}$  the set of all exceptional objects in  $\mathcal{T}$ ,

We will often write *ffe functor* instead of fully faithful exact functor in the sequel.

An *exceptional collection* is a sequence  $\mathcal{E} = (E_0, E_1, \dots, E_n) \subset \mathcal{T}_{exc}$  satisfying  $\text{hom}^*(E_i, E_j) = 0$  for  $i > j$ . If in addition we have  $\langle \mathcal{E} \rangle = \mathcal{T}$ , then  $\mathcal{E}$  will be called a full exceptional collection. For a vector  $\mathbf{p} = (p_0, p_1, \dots, p_n) \in \mathbb{Z}^{n+1}$  we denote  $\mathcal{E}[\mathbf{p}] = (E_0[p_0], E_1[p_1], \dots, E_n[p_n])$ . Obviously  $\mathcal{E}[\mathbf{p}]$  is also an exceptional collection. The exceptional collections of the form  $\{\mathcal{E}[\mathbf{p}] : \mathbf{p} \in \mathbb{Z}^{n+1}\}$  will be said to be shifts of  $\mathcal{E}$ .

If an exceptional collection  $\mathcal{E} = (E_0, E_1, \dots, E_n) \subset \mathcal{T}_{exc}$  satisfies  $\text{hom}^k(E_i, E_j) = 0$  for any  $i, j$  and for  $k \neq 0$ , then it is said to be *strong exceptional collection*.

For two exceptional collections  $\mathcal{E}_1, \mathcal{E}_2$  of equal length we write  $\mathcal{E}_1 \sim \mathcal{E}_2$  if  $\mathcal{E}_2 \cong \mathcal{E}_1[\mathbf{p}]$  for some  $\mathbf{p} \in \mathbb{Z}^{n+1}$ .

An *abelian category*  $\mathcal{A}$  is said to be *hereditary*, if  $\text{Ext}^i(X, Y) = 0$  for any  $X, Y \in \mathcal{A}$  and  $i \geq 2$ , it is said to be of *finite length*, if it is *Artinian* and *Noetherian*.

By  $Q$  we denote an acyclic quiver and by  $D^b(\text{Rep}_{\mathbb{K}}(Q))$ , or just  $D^b(Q)$ , - the derived category of the category of  $\mathbb{K}$ -representations of  $Q$ . Sometimes we write  $D^b(pt)$  instead of  $D^b(A_1)$ .

For an integer  $l \geq 0$  the  $l$ -Kronecker quiver (the quiver with two vertices and  $l$  parallel arrows) will be denoted by  $K(l)$ :  $1 \begin{array}{c} \curvearrowright \\ \vdots \\ \curvearrowleft \end{array} 2$  .

For a subset  $S \subset G$  of a group  $G$  we denote by  $\langle S \rangle \subset G$  the subgroup generated by  $S$ .

The number of elements of a finite set  $X$  we denote by  $|X|$  or by  $\#(X)$ .

For integers  $a, b \in \mathbb{Z}$  we denote by  $g.c.d(a, b)$  the greatest common divisor of  $a, b$ .

For any  $a \in \mathbb{R}$  and any complex number  $z \in e^{i\pi a} \cdot (\mathbb{R} + i\mathbb{R}_{>0})$ , respectively  $z \in e^{i\pi a} \cdot (\mathbb{R}_{<0} \cup (\mathbb{R} + i\mathbb{R}_{>0}))$ , we denote by  $\arg_{(a, a+1)}(z)$ , resp.  $\arg_{(a, a+1]}(z)$ , the unique  $\phi \in (a, a+1)$ , resp.  $\phi \in (a, a+1]$ , satisfying  $z = |z| \exp(i\pi\phi)$ .

For a non-zero complex number  $v \in \mathbb{C}$  we denote the two connected components of  $\mathbb{C} \setminus \mathbb{R}v$  by:

$$(5) \quad v_+^c = v \cdot (\mathbb{R} + i\mathbb{R}_{>0}) \quad v_-^c = v \cdot (\mathbb{R} - i\mathbb{R}_{>0}) \quad v \in \mathbb{C} \setminus \{0\}.$$

**Definition 11.** Let  $\mathcal{T}$  be a triangulated category and  $\mathcal{T}_i$  for  $i = 1, \dots, n$  be its triangulated subcategories, such that  $\mathcal{T} = \langle \mathcal{T}_1, \dots, \mathcal{T}_n \rangle$ . One says that  $\mathcal{T} = \langle \mathcal{T}_1, \dots, \mathcal{T}_n \rangle$  is a semi-orthogonal decomposition of  $\mathcal{T}$  when for any  $X_i \in \text{Ob}(\mathcal{T}_i)$  and  $X_j \in \text{Ob}(\mathcal{T}_j)$  the space of morphisms  $\text{Hom}^l(X_i, X_j)$  is trivial for all  $l$  whenever  $i > j$ .

### 3. TRIANGULATED CATEGORIES AND BRIDGELAND STABILITY CONDITIONS

In this work, we use the definition of triangulated categories found for example in [1, p. 239]. We give here the form of the octahedral axiom that we will use later: If one has the following commutative diagram in a triangulated category  $\mathcal{T}$

$$(6) \quad \begin{array}{ccccc} E & \xrightarrow{\quad} & F & \xrightarrow{\quad} & G \\ & \swarrow & \downarrow & \searrow & \downarrow \\ & [1] & A & \xleftarrow{\quad} & B \\ & & & [1] & \end{array}$$

with the triangles  $E, F, A$  and  $F, B, G$  being distinguished and the others being commutative, then there exists an object  $F'$  with a diagram

$$(7) \quad \begin{array}{ccc} E & \xrightarrow{\quad} & G \\ \uparrow & \swarrow & \downarrow \\ [1] & & F' \\ \downarrow & \searrow & \downarrow \\ A & \xleftarrow{\quad} & B \\ & [1] & \end{array},$$

such that the upper and lower triangle are again distinguished and the left and right triangle are commutative. The arrows  $A \rightarrow E$ ,  $E \rightarrow G$ ,  $G \rightarrow B$  and  $B \rightarrow A$  are the same in both diagrams.

**Remark 12.** Using the axioms of a triangulated category one can also show that if one has a diagram (7), then one can construct a diagram of the form (6), where the arrows  $A \rightarrow E$  and  $G \rightarrow B$  will be the same and the arrows  $E \rightarrow G$  and  $B \rightarrow A$  will get an additional minus sign in the newly constructed diagram.

The following lemma will be used in Section 5.

**Lemma 13.** If the diagram of distinguished triangles of the form

$$(8) \quad \begin{array}{ccccc} E & \xrightarrow{\quad} & F & \xrightarrow{\quad} & G \\ & \swarrow & \downarrow & \searrow & \downarrow \\ & [1] & A & \xrightarrow{g} & B \\ & & & [1] & \end{array}$$





(3) For any non-zero  $E \in \text{Ob}(\mathcal{T})$  there exists a sequence  $\phi_1 > \phi_2 > \dots > \phi_n$  and a diagram of distinguished triangles

$$(9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & E_1 & \longrightarrow & E_2 & \longrightarrow & \dots \longrightarrow E_{n-1} & \longrightarrow & E \\ & & \swarrow & & \swarrow & & & & \swarrow \\ & & [1] & & [1] & & & & [1] \\ & & A_1 & & A_2 & & & & A_n \end{array}$$

where  $A_i \in \text{Ob}(\mathcal{P}(\phi_i))$  are non-zero.

**Remark 15.** One can prove that the diagram (9) for any object  $E$  is unique up to isomorphism. We call this diagram the Harder-Narasimhan filtration of  $E$ .

**Definition 16.** [2] Let  $\mathcal{T}$  be a triangulated category. A pair  $\sigma = (Z, \mathcal{P})$ , where  $\mathcal{P}$  is a slicing on  $\mathcal{T}$  and  $Z : K_0(\mathcal{T}) \rightarrow \mathbb{C}$  is a group homomorphism, is said to be a stability condition on  $\mathcal{T}$  when for any non-zero  $A \in \mathcal{P}(t)$  there exists  $m_\sigma(A) \in \mathbb{R}^{>0}$ , such that

$$(10) \quad Z(A) = m_\sigma(A) \exp(i\pi t).$$

The homomorphism  $Z$  is then called the central charge of  $\sigma$ .

For a stability condition  $\sigma = (Z, \mathcal{P})$  a non-zero object  $A$  in  $\mathcal{P}(t)$  for some  $t$  is called  $\sigma$ -semistable, we will write  $\sigma^{ss}$  for the set of  $\sigma$ -semistable objects. Additionally, we denote  $\phi_\sigma(A) = t$  for  $A \in \mathcal{P}(t) \setminus \{0\}$ , and  $\phi_\sigma^+(E) = \phi_\sigma(A_1)$ ,  $\phi_\sigma^-(E) = \phi_\sigma(A_n)$  for any non-zero object  $E$  with HN filtration as in (9). One is usually interested in a special case of stability conditions called *locally finite* (see [2, Definition 5.7] for details).

**Remark 17.** In [2], the set of locally finite stability conditions is equipped with a structure of a complex manifold. This complex manifold is denoted by  $\text{Stab}(\mathcal{T})$ . Furthermore, in [2] a left action by biholomorphisms of the group of exact autoequivalences  $\text{Aut}(\mathcal{T})$  on  $\text{Stab}(\mathcal{T})$  is constructed. This action is determined as follows:

$$(11) \quad \text{Aut}(\mathcal{T}) \times \text{Stab}(\mathcal{T}) \ni (\Phi, (Z, \{\mathcal{P}(t)\}_{t \in \mathbb{R}})) \mapsto \left( Z \circ [\Phi]^{-1}, \left\{ \overline{\Phi(\mathcal{P}(t))} \right\}_{t \in \mathbb{R}} \right) \in \text{Stab}(\mathcal{T}),$$

where  $[\Phi] : K_0(\mathcal{T}) \rightarrow K_0(\mathcal{T})$  is the induced isomorphism (we will often omit specifying the square brackets) and  $\overline{\Phi(\mathcal{P}(t))}$  is the full isomorphism closed subcategory containing  $\Phi(\mathcal{P}(t))$ .

For any  $\Phi \in \text{Aut}(\mathcal{T})$ ,  $\sigma \in \text{Stab}(\mathcal{T})$  let us denote  $\sigma = (\mathcal{P}_\sigma, Z_\sigma)$ ,  $\Phi \cdot \sigma = (\mathcal{P}_{\Phi \cdot \sigma}, Z_{\Phi \cdot \sigma})$ , then we have:

$$(12) \quad (\Phi \cdot \sigma)^{ss} = \overline{\Phi(\sigma^{ss})} \quad \phi_{\Phi \cdot \sigma}(X) = \phi_\sigma(\Phi^{-1}X) \quad X \in (\Phi \cdot \sigma)^{ss}$$

$$(13) \quad Z_{\Phi \cdot \sigma}(X) = Z_\sigma(\Phi^{-1}(X)) \quad X \in \mathcal{T}.$$

#### 4. NON-COMMUTATIVE CURVE COUNTING. DEPENDENCE ON A STABILITY CONDITION

In this section we present in details an approach to make non-commutative counting dependable on stability condition. The idea is sketched in [5, Section 12.4]. We make also first basic observations. The main definition is

**Definition 18.** [5, Definition 12.5] Let  $\mathcal{A}, \mathcal{T}$  be any triangulated categories. And let  $\Gamma \subset \text{Aut}(\mathcal{T})$  be a subgroup of the group of auto-equivalences. We denote

$$(14) \quad C'_{\mathcal{A}, \mathcal{P}}(\mathcal{T}) = \{ \mathcal{A} \xrightarrow{F} \mathcal{T} : F \text{ is a fully faithful exact functor satisfying properties } P \}.$$

Here we need  $P$  to be a property of fully faithful functors, such that (14) is a well defined set, examples are in Definitions 19, 21.

Next we fix an equivalence relation in  $C'_{\mathcal{A},P}(\mathcal{T})$ :

$$(15) \quad C_{\mathcal{A},P}^{\Gamma}(\mathcal{T}) = C'_{\mathcal{A},P}(\mathcal{T})/\sim \quad F \sim F' \iff F \circ \alpha \cong \beta \circ F' \text{ for some } \alpha \in \text{Aut}(\mathcal{A}), \beta \in \Gamma$$

where  $F \circ \alpha \cong \beta \circ F'$  means equivalence of exact functors between triangulated categories (this is so called graded equivalence).

In special cases, important role play the categories  $D^b(\text{Rep}_{\mathbb{K}}(K(l+1))) = N\mathbb{P}^l$ ,  $l \geq -1$ .

These special cases of Definition 18 are relevant to non-commutative curve counting and to intersection theory, are:

**Definition 19.** [5, Definition 12.8], [6, Definitions 4.3, 6.4]

Let  $\mathcal{T}$  be  $\mathbb{K}$ -linear and let  $\Gamma \subset \text{Aut}_{\mathbb{K}}(\mathcal{T})$  be a subgroup of the group of exact  $\mathbb{K}$ -linear auto-equivalences on  $\mathcal{T}$ . Let the property  $P$  in (14) be “ $F$  is  $\mathbb{K}$ -linear”.

Let  $l \geq -1$ . We denote  $C'_{N\mathbb{P}^l,P}(\mathcal{T})$  and  $C^{\Gamma}_{N\mathbb{P}^l,P}(\mathcal{T})$  by  $C'_l(\mathcal{T})$  and  $C_l^{\Gamma}(\mathcal{T})$ , respectively, and refer to the elements of  $C_l^{\Gamma}(\mathcal{T})$  as to non-commutative curves of non-commutative genus  $l$  in  $\mathcal{T}$  modulo  $\Gamma$ .

We denote  $C'_{D^b(pt),P}(\mathcal{T})$  and  $C^{\Gamma}_{D^b(pt),P}(\mathcal{T})$  by  $C'_{D^b(pt),\mathbb{K}}(\mathcal{T})$  and  $C^{\Gamma}_{D^b(pt),\mathbb{K}}(\mathcal{T})$ , respectively, and refer to the elements of  $C^{\Gamma}_{D^b(pt),\mathbb{K}}(\mathcal{T})$  as to derived points in  $\mathcal{T}$  modulo  $\Gamma$ .

Furthermore, in the special case when  $\Gamma = \{\text{Id}_{\mathcal{T}}\}$  we will omit writing  $\Gamma$  in the superscript.

The following proposition holds

**Proposition 20.** [6, Proposition 5.5]. Let  $\mathcal{T}$  be a proper  $\mathbb{K}$ -linear triangulated category which has an enhancement, e.g.  $\mathcal{T} = D^b(\text{Rep}_{\mathbb{K}}(Q))$  for an acyclic quiver  $Q$ .

Then there are bijections:

$$(16) \quad C_l(\mathcal{T}) \rightarrow \left\{ \mathcal{A} \subset \mathcal{T} : \begin{array}{l} \mathcal{A} \text{ full triangulated subcategory of } \mathcal{T} \text{ s.t. } \mathcal{A} = \langle E_0, E_1 \rangle \\ \text{for some strong exceptional pair } (E_0, E_1) \text{ with} \\ \text{hom}(E_0, E_1) = l + 1 \end{array} \right\}$$

$$(17) \quad C_{D^b(A_1),\mathbb{K}}(\mathcal{T}) \rightarrow \left\{ \mathcal{A} \subset \mathcal{T} : \begin{array}{l} \mathcal{A} \text{ is a full tr. subcategory s.t. } \mathcal{A} = \langle E \rangle \\ \text{for some exceptional object } E \end{array} \right\}$$

defined by  $[F] \mapsto \text{Im}(F)$ .

From now on, we will assume that  $\mathcal{T}$  is a proper  $\mathbb{K}$ -linear triangulated category which has an enhancement and we will use Proposition 20 to identify  $C_l(\mathcal{T})$  and  $C_{D^b(A_1),\mathbb{K}}(\mathcal{T})$  with the corresponding sets of subcategories in  $\mathcal{T}$ .

Using Definition 18, the concept of  $\sigma$ -(semi)stability of non-commutative curves in  $\mathcal{T}$  was introduced in [5]. The idea is to specify a restriction  $P$  in Definition 18 depending on a stability condition  $\sigma \in \text{Stab}(\mathcal{T})$ . The following definition is the same as [5, Definition 12.12] when  $l \geq 1$ , the cases  $l = -1, 0$  are not carefully explained in [5, Definition 12.12], which we repair here:

**Definition 21.** Let  $l \geq -1$ . Let  $\Gamma \subset \text{Aut}_{\mathbb{K}}(\mathcal{T})$  and let  $\sigma \in \mathcal{T}$ .

Let  $\mathcal{E} \subset \text{Ob}(N\mathbb{P}^l)$  be a set of exceptional objects in  $N\mathbb{P}^l$ , s.t. for each exceptional object  $X \in N\mathbb{P}^l$  there exists unique  $Y \in \mathcal{E}$ , such that  $X \cong Y[k]$  for some  $k \in \mathbb{Z}$ .

We specify the property  $P$  from Definition 18 to two different cases and denote the set  $C'_{N\mathbb{P}^l, P}(\mathcal{T})$  from (14) by  $C'_{l, \sigma}(\mathcal{T})$ ,  $C'_{l, \sigma\sigma}(\mathcal{T})$ , respectively. More precisely, (recall  $C'_l(\mathcal{T})$  from Definition 19):

$$(18) \quad C'_{l, \sigma}(\mathcal{T}) = \{F \in C'_l(\mathcal{T}) : |\{E \in \mathcal{E} : F(E) \in \sigma^{ss}\}| = \infty\}$$

$$(19) \quad C'_{l, \sigma\sigma}(\mathcal{T}) = \{F \in C'_l(\mathcal{T}) : \forall E \in \mathcal{E} \quad F(E) \in \sigma^{ss}\}.$$

The formula (15) gives two sets of equivalence classes corresponding to the two different choices of  $P$  specified above and we refer to these sets as sets of  $\sigma$ -semistable (resp.  $\sigma$ -stable) non-commutative curves of non-commutative genus  $l$  in  $\mathcal{T}$ , and modulo  $\Gamma$ , we denote them as follows:

$$C_{l, \sigma}^\Gamma(\mathcal{T}) = C'_{l, \sigma}(\mathcal{T})/\sim \quad C_{l, \sigma\sigma}^\Gamma(\mathcal{T}) = C'_{l, \sigma\sigma}(\mathcal{T})/\sim$$

Again, we will skip the superscript  $\Gamma$  when  $\Gamma$  contains only the identity.

**Remark 22.** From (18), (19) and Definition 19 we see that  $C'_{l, \sigma}(\mathcal{T})$  and  $C'_{l, \sigma\sigma}(\mathcal{T})$  are subsets of  $C'_l(\mathcal{T})$ . We show here that these subsets are unions of equivalence classes in  $C_l(\mathcal{T})$  (according to Definition 19  $C_l(\mathcal{T})$  is a set of equivalence classes, which are subsets of  $C'_l(\mathcal{T})$ ). In particular, inclusions  $C_{l, \sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$  and  $C_{l, \sigma\sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$  hold.

Indeed, let  $F \in C'_{l, \sigma}(\mathcal{T})$  and  $G \in C'_{l, \sigma}(\mathcal{T})$ , resp.  $G \in C'_{l, \sigma\sigma}(\mathcal{T})$ , be such that  $F \circ \alpha \cong G$  for some  $\alpha \in \text{Aut}(N\mathbb{P}^l)$ . Since by definition both  $F$  and  $G$  are  $\mathbb{K}$ -linear it follows from [6, Corollary 3.10] that  $\alpha$  is  $\mathbb{K}$ -linear as well, and hence  $\alpha(X)$  is exceptional iff  $X$  is exceptional for any  $X \in \text{Ob}(N\mathbb{P}^l)$ . It follows that  $F \in C'_{l, \sigma}(\mathcal{T})$ , resp.  $F \in C'_{l, \sigma\sigma}(\mathcal{T})$ , as well.

Next we will discuss here the restriction of the bijection (16) to the subsets  $C_{l, \sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$ ,  $C_{l, \sigma\sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$ . We note first that the set  $\mathcal{E}$  from Definition 21 is bijective to the set of derived points in  $N\mathbb{P}^l$  (recall Definition 19 and (17)):

**Remark 23.** The objects in a derived point  $\langle E \rangle$  in any proper  $\mathbb{K}$ -linear  $\mathcal{T}$  are exactly all the possible finite direct sums of shifts of  $E$ . Therefore the function  $\mathcal{E} \rightarrow C_{D^b(pt), \mathbb{K}}(N\mathbb{P}^l)$  defined by  $\mathcal{E} \ni E \mapsto \langle E \rangle \in C_{D^b(pt), \mathbb{K}}(N\mathbb{P}^l)$  is a bijection, where  $\mathcal{E} \subset \text{Ob}(N\mathbb{P}^l)$  is as in Definition 21. Hence for any  $F \in C'_l(\mathcal{T})$  the function  $\mathcal{E} \ni E \mapsto \langle F(E) \rangle \in C_{D^b(pt), \mathbb{K}}(\text{Im}(F))$  is also a bijection.

Taking into account this we define:

**Definition 24.** Let  $\sigma \in \text{Stab}(\mathcal{T})$ . Due to the first sentence in Remark 23 we see that if we have a derived point  $\mathcal{A} \in C_{D^b(pt), \mathbb{K}}(\mathcal{T})$  and  $\mathcal{A} = \langle A \rangle = \langle B \rangle$  with  $A, B$  exceptional objects, then  $A \in \sigma^{ss}$  if and only if  $B \in \sigma^{ss}$ . Hence the set  $C_{D^b(pt), \sigma\sigma}(\mathcal{T}) = \{\langle E \rangle \in C_{D^b(pt), \mathbb{K}}(\mathcal{T}) : E \in \sigma^{ss}\}$  is well defined.

The following is a new definition motivated by counting of derived points.

**Definition 25.** Let  $\sigma \in \text{Stab}(\mathcal{T})$ , we define

$$\mathfrak{D}_\sigma = \{\langle E \rangle \in C_{D^b(pt), \sigma\sigma} : \langle E \rangle \not\subset \mathcal{A} \quad \forall \mathcal{A} \in C_{l, \sigma\sigma}, l \geq 1\}.$$

The elements of  $\mathfrak{D}_\sigma$  are called stable derived points outside of stable curves of positive genus.

Using this terminology and Remark 23 we see that:

**Lemma 26.** Let  $l \geq -1$  and  $\sigma \in \text{Stab}(\mathcal{T})$ . In Remark 22 we explained that  $C_{l, \sigma}(\mathcal{T})$ ,  $C_{l, \sigma\sigma}(\mathcal{T})$  are subsets of  $C_l(\mathcal{T})$ . Let  $\mathcal{A} \subset \mathcal{T}$  be an element in the codomain of (16). Then

- (a)  $\mathcal{A} \in C_{l, \sigma}(\mathcal{T})$  iff  $|C_{D^b(pt), \sigma\sigma}(\mathcal{T}) \cap C_{D^b(pt), \mathbb{K}}(\mathcal{A})| = \infty$ .
- (b)  $\mathcal{A} \in C_{l, \sigma\sigma}(\mathcal{T})$  iff  $C_{D^b(pt), \mathbb{K}}(\mathcal{A}) \subset C_{D^b(pt), \sigma\sigma}(\mathcal{T})$ .

**Corollary 27.** *For any  $\mathcal{A} \in C_l(\mathcal{T})$ ,  $l \geq -1$  the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : \mathcal{A} \in C_{l,\sigma\sigma}(\mathcal{T})\}$  is a closed subset in  $\text{Stab}(\mathcal{T})$ .*

*Proof.* Let  $\mathcal{E}$  be the set of exceptional objects in  $\mathcal{A}$ . From [2, p. 342] is known that for any object  $X$  in  $\mathcal{T}$  the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : X \in \sigma^{ss}\}$  is closed. From Lemma 26 we see that  $\{\sigma \in \text{Stab}(\mathcal{T}) : \mathcal{A} \in C_{l,\sigma\sigma}(\mathcal{T})\} = \bigcap_{E \in \mathcal{E}} \{\sigma \in \text{Stab}(\mathcal{T}) : X \in \sigma^{ss}\}$ .  $\square$

**Remark 28.** *Let  $\mathcal{E}$  be a set of exceptional objects as in Definition 21.*

(a) *Let  $l \geq 1$  and let  $\{s_i\}_{i \in \mathbb{Z}}$  be a Helix in  $N\mathbb{P}^l$  (see [5, Subsection 7.1]). Then from [5, (72), (73)] we see that we can choose  $\mathcal{E} = \{s_i : i \in \mathbb{Z}\}$  and that the function  $\mathbb{Z} \ni i \mapsto s_i \in \mathcal{E}$  is bijection.*

(b) *If  $l = 0$  or  $l = -1$ , then  $N\mathbb{P}^l$ , and hence  $\mathcal{E}$ , has three or respectively two elements (see [6, Figure 1]) and it follows that for  $l = 0, l = -1$  we have always  $C_{l,\sigma}^\Gamma(\mathcal{T}) = \emptyset$ . The derived points in a genus zero nc curve cannot be ordered in a semi-orthogonal triple (follows also from [6, Figure 1]).*

(c) *From (a) it follows that for  $l \geq 1$  we have  $C_{l,\sigma\sigma}(\mathcal{T}) \subset C_{l,\sigma}(\mathcal{T}) \subset C_l(\mathcal{T})$  (recall remark 22).*

**Remark 29.** *Taking into account Remark 26, it seems reasonable to call the elements in  $C_{l,\sigma\sigma}(\mathcal{T})$  point-wise semistable and those in  $C_{l,\sigma}(\mathcal{T})$  - almost everywhere point-wise semistable. Also one can define similarly point-wise semistability for any subcategory in  $\mathcal{T}$ . Actually, provided that one has defined semi-stability for some triangulated category  $\mathcal{A}$  (could be different from derived point), then by analogy one gets in similar fashion notions of  $\mathcal{A}$ -wise semistable and of almost everywhere  $\mathcal{A}$ -wise semistable subcategories in  $\mathcal{T}$ . However, here we will use the shorter terminology fixed in Definition 21 and its equivalent description in Lemma 26.*

## 5. SOME REMARKS FOR $\sigma$ -EXCEPTIONAL COLLECTIONS AND $C_{l,\sigma\sigma}(\mathcal{T})$

*From now on, we will assume that the triangulated categories we are working with are proper.*

Firstly, we recall a Definition from [7] :

**Definition 30.** [7, Definition 3.17, Remark 3.19] *Let  $\sigma = (\mathcal{P}, Z) \in \text{Stab}(\mathcal{T})$ . An exceptional collection  $\mathcal{E} = (E_0, E_1, \dots, E_n)$  is called  $\sigma$ -exceptional collection if the following properties hold:*

- $\mathcal{E}$  is semistable w. r. to  $\sigma$  (i. e. all  $E_i$  are semistable).
- $\forall i \neq j \text{ hom}^{\leq 0}(E_i, E_j) = 0$  (i. e. this is an Ext-exceptional collection).
- There exists  $t \in \mathbb{R}$ , s. t.  $\{\phi(E_i)\}_{i=0}^n \subset (t, t+1]$  (this is equivalent to  $|\phi(E_i) - \phi(E_j)| < 1$  for  $0 \leq i < k \leq n$ ).

**Remark 31.** *Let  $\mathcal{E}$  be a full exceptional collection. The following is a well defined assignment:*

$$(20) \quad \{\sigma \in \text{Stab}(\mathcal{T}) : \mathcal{E} \subset \sigma^{ss}\} \xrightarrow{f_{\mathcal{E}}} (\{|Z(E_i)|\}_{i=0}^n, \{\phi_{\sigma}(E_i)\}_{i=0}^n) \in \mathbb{R}^{2(n+1)}$$

(a) *Let  $\mathcal{E}$  be Ext<sup>2</sup> and let  $\Theta'_{\mathcal{E}}$  denote the set of stability conditions for which  $\mathcal{E}$  is  $\sigma$ -exceptional. From [7, Corollary 3.18, Remark 3.19] we know that  $\Theta'_{\mathcal{E}}$  is an open subset in  $\text{Stab}(\mathcal{T})$  and (20) restricted to  $\Theta'_{\mathcal{E}}$  gives a homeomorphism between  $\Theta'_{\mathcal{E}}$  and the following simply connected set:*

$$\{(x_0, \dots, x_n, y_0, \dots, y_n) \in \mathbb{R}^{2(n+1)} : x_i > 0, |y_i - y_j| < 1\}.$$

(b) *Let  $\mathcal{E}$  be any full exceptional collection in a proper  $\mathcal{T}$ , then after shifting it is an Ext-exceptional collection. Let  $\Theta_{\mathcal{E}}$  denote the set of stability conditions  $\sigma \in \text{Stab}(\mathcal{T})$ , such that a shift of  $\mathcal{E}$  is  $\sigma$ -exceptional. In [8, (11)] it is shown that (20) restricted to the open subset  $\Theta_{\mathcal{E}}$  is also homeomorphism, however the image is more complicated than for  $\Theta'_{\mathcal{E}}$ .*

<sup>2</sup>satisfies the second condition in Definition 30

For a class of objects  $\mathcal{E}$  in  $\mathcal{T}$  we denote by  $\hat{\mathcal{E}}$  the extension closure in  $\mathcal{T}$  of  $\mathcal{E}$ .

**Remark 32.** Let  $E \in \mathcal{T}$  be an exceptional object. It is well known that every object  $X$  in  $\langle E \rangle$  is a direct product of finitely many objects from the class  $\{E[i]\}_{i \in \mathbb{Z}}$ , that  $\hat{E}[i]$  is a heart of a bounded  $t$ -structure in  $\langle E \rangle$ , and that  $\hat{E}[i]$  is of finite length for each  $i \in \mathbb{Z}$ . Due to  $\text{hom}^i(E, E) = 0$  whenever  $i \neq 0$ , it follows that any object  $X$  in  $\hat{E}$  is a direct product of copies of  $E$ .

**Proposition 33.** Let  $\mathcal{T}$  be a triangulated category. Let  $\sigma \in \text{Stab}(\mathcal{T})$ . If  $\mathcal{E} = (E_0, \dots, E_n)$  its full  $\sigma$ -exceptional collection with strictly increasing phases of its objects, i. e.  $\phi_\sigma(E_{i+1}) > \phi_\sigma(E_i)$  for all  $0 \leq i < n$ , then the slicing  $\mathcal{P}$  of  $\sigma$  is given in the following way.

$$(21) \quad \mathcal{P}(t) = \begin{cases} \hat{E}_i[j] & \text{when } t = \phi_\sigma(E_i) + j \\ \{0\} & \text{otherwise} \end{cases}$$

*Proof.* We note first that from the given properties of  $\mathcal{E}$  and  $\sigma$  it follows:

$$(22) \quad \phi_\sigma(E_{i_2}[j_2]) > \phi_\sigma(E_{i_1}[j_1]) \iff j_2 > j_1 \text{ or } j_2 = j_1, i_2 > i_1.$$

Let us show that  $\sigma' = (\mathcal{P}, Z)$  indeed gives a locally finite stability condition when  $\mathcal{P}$  is given by the equation (21) and  $Z$  is the central charge of the given stability condition. For that, we only need to show that  $\mathcal{P}$  is a locally finite slicing on  $\mathcal{T}$ . The first two axioms of the Definition 14 follow immediately. By a property of semi-orthogonal decomposition, for any object  $F$  there is a filtration with factors  $A_i$  that are objects of  $\langle E_i \rangle$  and as such are the direct sums of shifts of  $E_i$  by Remark 32. So  $\mathcal{T}$  is the extension closure of  $\{E_i[j]\}_{0 \leq i \leq n, j \in \mathbb{Z}}$ . However, when there is a diagram of the form

$$\begin{array}{ccccc} A & \xrightarrow{\quad} & B & \xrightarrow{\quad} & C \\ & \swarrow & \searrow & \swarrow & \searrow \\ & [1] & & [1] & \\ & & E_{i_1}[j_1] & & E_{i_2}[j_2] \end{array}$$

where  $\phi_Z(E_{i_2}[j_2]) > \phi_Z(E_{i_1}[j_1])$ , then by  $\mathcal{E}$  being Ext-exceptional and using 22 it follows that the composition  $E_{i_2}[j_2] \rightarrow B[1] \rightarrow E_{i_1}[j_1 + 1]$  is either zero morphism or isomorphism<sup>3</sup>, so we see from Lemma 13 that we can either interchange the order of the factors  $E_{i_1}[j_1]$ ,  $E_{i_2}[j_2]$  or the composition  $A \rightarrow B \rightarrow C$  is isomorphism and we can stick  $A$  and  $C$ . Doing so until any two neighboring factors  $E_{i_1}[j_1]$  and  $E_{i_2}[j_2]$  have  $\phi_Z(E_{i_1}[j_1]) \geq \phi_Z(E_{i_2}[j_2])$  and then using the octahedral axiom where two neighboring factors  $E_{i_1}[j_1]$  and  $E_{i_2}[j_2]$  have  $\phi_Z(E_{i_1}[j_1]) = \phi_Z(E_{i_2}[j_2])$ , gives us a HN-filtration of  $F$  with respect to  $\mathcal{P}$ . In this slicing there exists  $\epsilon > 0$ , such that  $\mathcal{P}(t - \epsilon, t + \epsilon)$  is either trivial or  $\hat{E}_i[j]$  for some  $i, j$ , which by Remark 32 is a locally finite abelian category, and therefore  $\mathcal{P}$  is locally finite slicing (see [2, Definition 5.7]). Therefore  $\sigma' \in \text{Stab}(\mathcal{T})$ .

Since  $\phi_{\sigma'}(E_i) = \phi_\sigma(E_i)$  for  $0 \leq i \leq n$  it follows that  $\mathcal{E}$  is also  $\sigma'$ -exceptional. Furthermore, the central charges of  $\sigma$  and  $\sigma'$  are the same. Therefore the bijection (20) has equal values on  $\sigma$  and  $\sigma'$  and therefore  $\sigma' = \sigma$ .  $\square$

**Corollary 34.** Let  $\mathcal{T}$  be a triangulated category and  $\sigma \in \text{Stab}(\mathcal{T})$ . Let  $\mathcal{E} = (E_0, \dots, E_n)$  be a full  $\sigma$ -exceptional collection with strictly increasing phases. Then the set of stable derived points, as defined in Definition 24, is  $C_{D^b(pt), \sigma\sigma} = \{\langle E_0 \rangle, \langle E_1 \rangle, \dots, \langle E_n \rangle\}$ . In particular this set is finite and  $C_{l, \sigma\sigma}(\mathcal{T}) = C_{l, \sigma}(\mathcal{T}) = \emptyset$  for  $l \geq 0$ .

<sup>3</sup>in the case  $i_1 = i_2$  and  $j_2 = j_1 + 1$  this arrow is of the form  $E_{i_1}[j_1 + 1] \rightarrow E_{i_1}[j_1 + 1]$

*Proof.* From the previous Proposition we have  $\sigma^{ss} = \{E : E \in \hat{E}_i[j] : 0 \leq i \leq n, j \in \mathbb{Z}\}$ . If  $\langle E \rangle$  is a stable derived point, then from Definition 24  $E \in \sigma^{ss}$ , hence  $E \in \hat{E}_i[j]$  for some  $0 \leq i \leq n$  and some  $j \in \mathbb{Z}$  and from Remark 32 it follows that  $\langle E \rangle = \langle E_i \rangle$ . Obviously  $\langle E_i \rangle$  is stable for each  $i$ . Therefore indeed  $C_{D^b(pt), \sigma\sigma} = \{\langle E_0 \rangle, \langle E_1 \rangle, \dots, \langle E_n \rangle\}$ , and it has  $n + 1$  elements. Now from Lemma 26 it follows that  $C_{l, \sigma}(\mathcal{T}) = C_{l, \sigma\sigma}(\mathcal{T}) = \emptyset$  for  $l \geq 1$ . On the other hand, if  $C_{0, \sigma\sigma}(\mathcal{T}) \neq \emptyset$  and  $\mathcal{A} \in C_{0, \sigma\sigma}(\mathcal{T})$ , then the three derived points in  $\mathcal{A}$  (see Remark 28 (b)) must be of the form  $\langle E_i \rangle, \langle E_j \rangle, \langle E_k \rangle$  with  $0 \leq i < j < k \leq n$ , which contradicts the last sentence in Remark 28 (b).  $\square$

**Remark 35.** *Let  $\mathcal{E} = (E_0, \dots, E_n)$  be a full Ext-exceptional collection. The homeomorphism (20) restricted to the set of stability conditions  $\sigma \in \Theta'_\mathcal{E}$  with strictly increasing phases on  $\mathcal{E}$  gives a homeomorphism between this set and the following subset of  $\mathbb{R}^{2(n+1)}$ :*

$$\left\{ (x_0, \dots, x_n, y_0, \dots, y_n) \in \mathbb{R}^{2(n+1)} \quad : \quad x_i > 0, y_0 < y_1 < \dots < y_n < y_0 + 1 \right\}.$$

**Proposition 36.** *Let  $\mathcal{T}$  have an enhancement, let  $\sigma \in \text{Stab}(\mathcal{T})$  and let  $\mathcal{E} = (E_0, \dots, E_n)$  be a full  $\sigma$ -exceptional collection, such that  $\text{hom}^1(E_i, E_{i+1}) = l + 1$  and  $\text{hom}^j(E_i, E_{i+1}) = 0$  for  $j \in \mathbb{Z} \setminus \{1\}$  for some  $0 \leq i < n$  and some  $l \geq 0$ . If  $\phi(E_i) \geq \phi(E_{i+1})$ , then  $\langle E_i, E_{i+1} \rangle \in C_{l, \sigma\sigma}(\mathcal{T})$ .*

*Proof.* From Proposition 20 we see that  $\langle E_i, E_{i+1} \rangle \in C_l(\mathcal{T})$ , in particular there is an exact equivalence

$$(23) \quad F : \langle E_i, E_{i+1} \rangle \rightarrow D^b(K(l)).$$

We will show that all exceptional objects of  $\mathcal{T}$ , which lie in  $\langle E_i, E_{i+1} \rangle$ , are in  $\sigma^{ss}$  and then from Lemma 26 will follow that  $\langle E_i, E_{i+1} \rangle \in C_{l, \sigma\sigma}(\mathcal{T})$ . Let  $\mathcal{A}$  be the extension closure of  $(E_i, E_{i+1})$  in  $\mathcal{T}$ . From [8, Proposition 2.2] we see that each exceptional object in  $\mathcal{T}$ , which is also in  $\mathcal{A}$ , is in  $\sigma^{ss}$ . To finish the proof it is enough to show that each exceptional object in  $\langle E_i, E_{i+1} \rangle$  is isomorphic to a shift of an object in  $\mathcal{A}$ . Since  $F$  in (23) is exact equivalence, it is enough to prove that

$$(24) \quad \text{Each exceptional object in } D^b(K(l+1)) \text{ is isomorphic to a shift of an object in } F(\mathcal{A}),$$

Now  $F(E_i)$  is an exceptional object in  $D^b(K(l+1))$ . Using an auto-equivalence of  $D^b(K(l+1))$  we can assume that  $F(E_i)$  is the simple representation in  $\text{Rep}_{\mathbb{K}}(K(l+1))$  with dimension vector  $(1, 0)$  (for the case  $l = 0$  we use [6, Proposition 10.9] and for the case  $l \geq 1$  we use [9, Corollary 5.3]). On the other hand  $F(E_i), F(E_{i+1})$  is an exceptional pair and it follows that  $F(E_{i+1})$  is isomorphic to a shift of the simple representation in  $\text{Rep}_{\mathbb{K}}(K(l+1))$  with dimension vector  $(0, 1)$  (this follows for example from results in [4]). However, we have also  $\text{hom}^1(F(E_i), F(E_{i+1})) = l + 1$  and it follows that this shift is trivial. Hence we see that we can assume that  $(F(E_i), F(E_{i+1}))$  is the exceptional pair of the simple representations of  $K(l)$ . It follows that each  $X \in \text{Rep}_{\mathbb{K}}(K(l+1))$  is isomorphic to  $F(A)$  for some  $A \in \mathcal{A}$ , and since  $\text{Rep}_{\mathbb{K}}(K(l+1))$  is hereditary we obtain (24).  $\square$

**Corollary 37.** *Let  $\mathcal{T}$  have an enhancement, let  $\mathcal{E} = (E_0, \dots, E_n)$  be a full exceptional collection such that for some  $0 \leq i < n$ , some  $\alpha \in \mathbb{Z}$ , and some  $l \geq 0$  holds  $\text{hom}^{1+\alpha}(E_i, E_{i+1}) = l + 1$  and  $\text{hom}^j(E_i, E_{i+1}) = 0$  for  $j \in \mathbb{Z} \setminus \{1 + \alpha\}$ . If  $\sigma \in \Theta_\mathcal{E}$  and  $\phi(E_i) \geq \phi(E_{i+1}[\alpha])$ , then  $\langle E_i, E_{i+1} \rangle \in C_{l, \sigma\sigma}(\mathcal{T})$ .*

*Proof.* By the definition of  $\Theta_\mathcal{E}$  in Remark 31, the sequence  $(E_0[p_0], \dots, E_n[p_n])$  is  $\sigma$ -exceptional for some  $(p_0, \dots, p_n) \in \mathbb{Z}^{n+1}$ . We can assume that  $p_i = 0$  and therefore  $\text{hom}^j(E_i, E_{i+1}[p_{i+1}]) = 0$  for  $j \leq 0$ ,  $|\phi(E_i) - \phi(E_{i+1}[p_{i+1}])| < 1$ . Since we are given  $\text{hom}^{1+\alpha}(E_i, E_{i+1}) = l + 1$  and  $\text{hom}^j(E_i, E_{i+1}) = 0$  for  $j \in \mathbb{Z} \setminus \{1 + \alpha\}$ , therefore  $p_{i+1} \leq \alpha$ . If  $p_{i+1} < \alpha$ , then  $p_{i+1} = \alpha - \delta$  for

some  $\delta \geq 1$  and  $\phi(E_i) - \phi(E_{i+1}[p_{i+1}]) = \phi(E_i) - \phi(E_{i+1}[\alpha]) + \delta \geq \delta \geq 1$  by the given  $\phi(E_i) - \phi(E_{i+1}[\alpha]) \geq 0$ , and this contradicts  $|\phi(E_i) - \phi(E_{i+1}[p_{i+1}])| < 1$ , Therefore  $p_{i+1} = \alpha$  and we apply the Proposition 36 to the sequence  $(E_0[p_0], \dots, E_n[p_n])$ .  $\square$

## 6. THE EXAMPLE: STABLE NON-COMMUTATIVE CURVES IN $D^b(Q)$

We fix  $\mathcal{T} = D^b(Q)$ , where  $Q$  is as in (1). Here we will study how the sets  $C_{i,\sigma\sigma}(\mathcal{T}) \subset C_{i,\sigma}(\mathcal{T})$  (defined in Definition 21) change as  $\sigma$  varies in  $\text{Stab}(\mathcal{T})$  and  $i \in \mathbb{Z}$ . Recalling that  $C_l(\mathcal{T}) = \emptyset$  for  $l \geq 2$  (see Proposition (4)) and using Remark 28 (c) we see that  $C_{l,\sigma\sigma}(\mathcal{T}) = C_{l,\sigma}(\mathcal{T}) = \emptyset$  for  $l \geq 2$ .

Our study uses to a great extent results from [6, 7, 8] regarding exceptional objects in  $\mathcal{T}$  and  $\text{Stab}(\mathcal{T})$ . We present these results in the first two subsections.

**6.1. On the exceptional objects in  $D^b(Q)$ .** Let  $\underline{\dim}(E) = (x, y, z)$  be the dimension vector of an exceptional representation in  $\text{Rep}_{\mathbb{K}}(Q)$ . Let  $\pi_{\pm}^m : \mathbb{K}^{m+1} \rightarrow \mathbb{K}^m$  and  $j_{\pm}^m : \mathbb{K}^m \rightarrow \mathbb{K}^{m+1}$  be linear maps such that

$$\begin{aligned} \pi_+^m(x_1, \dots, x_{m+1}) &= (x_1, \dots, x_m) & \pi_-^m(x_1, \dots, x_{m+1}) &= (x_2, \dots, x_{m+1}) \\ j_+^m(x_1, \dots, x_m) &= (x_1, \dots, x_m, 0) & j_-^m(x_1, \dots, x_m) &= (0, x_1, \dots, x_m). \end{aligned}$$

**Proposition 38.** [7, Proposition 2.2] *The exceptional objects of  $\mathcal{T}$  up to equivalence are:*

$$\begin{aligned} E_1^m &= \begin{array}{ccc} & \mathbb{K}^m & \\ \pi_+^m \nearrow & & \nwarrow id \\ \mathbb{K}^{m+1} & \xrightarrow{\pi_-^m} & \mathbb{K}^m \end{array} & E_2^m &= \begin{array}{ccc} & \mathbb{K}^{m+1} & \\ j_+^m \nearrow & & \nwarrow id \\ \mathbb{K}^m & \xrightarrow{j_-^m} & \mathbb{K}^{m+1} \end{array} & E_3^m &= \begin{array}{ccc} & \mathbb{K}^{m+1} & \\ j_+^m \nearrow & & \nwarrow j_-^m \\ \mathbb{K}^m & \xrightarrow{id} & \mathbb{K}^m \end{array} \\ E_4^m &= \begin{array}{ccc} & \mathbb{K}^m & \\ \pi_+^m \nearrow & & \nwarrow \pi_-^m \\ \mathbb{K}^{m+1} & \xrightarrow{id} & \mathbb{K}^{m+1} \end{array} & M &= \begin{array}{ccc} & 0 & \\ \nearrow & & \nwarrow \\ 0 & \xrightarrow{\quad} & \mathbb{K} \end{array} & M' &= \begin{array}{ccc} & \mathbb{K} & \\ \nearrow & & \nwarrow \\ \mathbb{K} & \xrightarrow{\quad} & 0 \end{array} \end{aligned}$$

where  $m$  goes over all non-negative integers.

Following [8], we denote

$$(25) \quad a^m = \begin{cases} E_1^{-m} & m \leq 0 \\ E_2^{m-1}[1] & m \geq 1 \end{cases}, \quad b^m = \begin{cases} E_4^{-m} & m \leq 0 \\ E_3^{m-1}[1] & m \geq 1 \end{cases}.$$

Due to Proposition 38 and Corollary 17 holds (2).

The following results from [7, 8] will be used often later.

**Corollary 39.** ([8, Corollary 3.7]) For each  $m \in \mathbb{Z}$  we have:

$$\begin{aligned}
(26) \quad & \text{hom}(M', a^m) \neq 0; \quad \text{hom}(M, b^m) \neq 0; \quad \text{hom}^*(a^m, M') = 0; \\
(27) \quad & \text{hom}^1(a^m, M) \neq 0; \quad \text{hom}^1(b^m, M') \neq 0; \quad \text{hom}^*(b^m, M) = 0 \\
(28) \quad & \text{hom}^*(b^m, a^{m-1}) = 0; \quad \text{hom}(b^m, a^n) \neq 0 \text{ for } m \leq n; \quad \text{hom}^1(b^m, a^n) \neq 0 \text{ for } m > n + 1; \\
(29) \quad & \text{hom}^*(a^m, b^m) = 0; \quad \text{hom}(a^m, b^{n+1}) \neq 0 \text{ for } m \leq n; \quad \text{hom}^1(a^m, b^n) \neq 0 \text{ for } m > n; \\
(30) \quad & \text{hom}^*(a^m, a^{m-1}) = 0; \quad \text{hom}(a^m, a^n) \neq 0 \text{ for } m \leq n; \quad \text{hom}^1(a^m, a^n) \neq 0 \text{ for } m > n + 1; \\
(31) \quad & \text{hom}^*(b^m, b^{m-1}) = 0; \quad \text{hom}(b^m, b^n) \neq 0 \text{ for } m \leq n; \quad \text{hom}^1(b^m, b^n) \neq 0 \text{ for } m > n + 1; \\
(32) \quad & \text{hom}^1(M, M') \neq 0 \quad \text{hom}^1(M', M) \neq 0.
\end{aligned}$$

**Corollary 40.** [7, Corollary 2.6 (b)] For any two exceptional objects  $X, Y \in D^b(Q)$  at most one element of the family  $\{\text{hom}^p(X, Y)\}_{p \in \mathbb{Z}}$  is nonzero.

In [8, Remark 3.15] is explained that for any  $p, q \in \mathbb{Z}$  there are distinguished triangles

$$(33) \quad \begin{array}{ccc} b^{p+1}[-1] & \longrightarrow & M' \\ & \searrow \text{dashed} & \swarrow \\ & a^p & \end{array} \quad \begin{array}{ccc} a^q[-1] & \longrightarrow & M \\ & \searrow \text{dashed} & \swarrow \\ & b^q & \end{array} .$$

**Remark 41.** We will denote by  $\delta \in K_0(D^b(Q))$  the element  $\delta = [E_1^0] + [M] + [E_3^0]$ . For  $\delta$  holds  $\delta = [M] + [M']$ . In particular for any stability condition  $(Z, \mathcal{P}) \in \text{Stab}(\mathcal{T})$  hold the equalities  $Z(\delta) = Z(M) + Z(M')$ . Furthermore, from (33) we see that  $Z(a^p) = Z(M') + Z(b^{p+1})$  and  $Z(b^q) = Z(M) + Z(a^q)$  for any  $p, q \in \mathbb{Z}$ . On the other hand, the latter two equalities imply that  $Z(b^p) = Z(\delta) + Z(b^{p+1})$ ,  $Z(a^p) = Z(\delta) + Z(a^{p+1})$ .

**6.2. On stability conditions on  $D^b(Q)$ .** The entire manifold  $\text{Stab}(\mathcal{T})$  was described in [8], and there it was proved that it is contractible. We recollect some material from loc cit.

For a given triple  $(A, B, C) \in \mathfrak{T}$  we will denote the open subset  $\Theta_{(A,B,C)} \subset \text{Stab}(D^b(Q))$  from Remark 31 (b) by  $(A, B, C)$ , when no confusion may arise. In [8, Proposition 2.7] it is shown that the image of the homeomorphism in Remark 31 (b) is a set of the form

$$(34) \quad \mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_0 - y_1 < 1 + \alpha \\ y_0 - y_2 < 1 + \min\{\beta, \alpha + \gamma\} \\ y_1 - y_2 < 1 + \gamma \end{array} \right\},$$

where  $\alpha, \beta, \gamma \in \mathbb{Z}$  are defined in [8, (30)]. An important role play the exceptional triples. They are:

**Proposition 42.** [8, Corollary 3.12, Remark 3.14] Define the following sets of full exceptional triples in  $\mathcal{T}$ :

- (i)  $\mathfrak{T}_a = \{(M', a^m, a^{m+1}), (a^m, b^{m+1}, a^{m+1}), (a^m, a^{m+1}, M) : m \in \mathbb{Z}\}$ ,
- (ii)  $\mathfrak{T}_b = \{(M, b^m, b^{m+1}), (b^m, a^m, b^{m+1}), (b^m, b^{m+1}, M') : m \in \mathbb{Z}\}$ .

The set of all full exceptional collections in  $\mathcal{T}$  up to shifts is:

$$(35) \quad \mathfrak{T} = \mathfrak{T}_a \cup \{(a^m, M, b^{m+1}) : m \in \mathbb{Z}\} \cup \{(b^m, M', a^m) : m \in \mathbb{Z}\} \cup \mathfrak{T}_b.$$



The stability conditions in  $(A, B, C)$  are determined by the properties that the objects  $A, B, C$  are in  $\sigma^{ss}$  and their phases satisfy the inequalities summarized in the following tables:

(36)

$\mathfrak{T}_a^{st}$	$\mathfrak{T}_b^{st}$
$\left\{ \begin{array}{l} M', a^j, a^{j+1} \in \sigma^{ss} : \phi(M') < \phi(a^j) \\ \phi(M') + 1 < \phi(a^{j+1}) \\ \phi(a^j) < \phi(a^{j+1}) \end{array} \right\}$	$\left\{ \begin{array}{l} M, b^n, b^{n+1} \in \sigma^{ss} : \phi(M) < \phi(b^n) \\ \phi(M) + 1 < \phi(b^{n+1}) \\ \phi(b^n) < \phi(b^{n+1}) \end{array} \right\}$
$\left\{ \begin{array}{l} a^p, b^{p+1}, a^{p+1} \in \sigma^{ss} : \phi(a^p) < \phi(b^{p+1}) \\ \phi(a^p) + 1 < \phi(a^{p+1}) \\ \phi(b^{p+1}) < \phi(a^{p+1}) \end{array} \right\}$	$\left\{ \begin{array}{l} b^q, a^q, b^{q+1} \in \sigma^{ss} : \phi(b^q) < \phi(a^q) \\ \phi(b^q) + 1 < \phi(b^{q+1}) \\ \phi(a^q) < \phi(b^{q+1}) \end{array} \right\}$
$\left\{ \begin{array}{l} a^m, a^{m+1}, M \in \sigma^{ss} : \phi(a^m) < \phi(a^{m+1}) \\ \phi(a^m) < \phi(M) \\ \phi(a^{m+1}) < \phi(M) + 1 \end{array} \right\}$	$\left\{ \begin{array}{l} b^i, b^{i+1}, M' \in \sigma^{ss} : \phi(b^i) < \phi(b^{i+1}) \\ \phi(b^i) < \phi(M') \\ \phi(b^{i+1}) < \phi(M') + 1 \end{array} \right\}$ .

$(a^p, M, b^{p+1})$	$(b^q, M', a^q)$
$\left\{ \begin{array}{l} a^p, M, b^{p+1} \in \sigma^{ss} : \phi(a^p) < \phi(M) + 1 \\ \phi(a^p) < \phi(b^{p+1}) \\ \phi(M) < \phi(b^{p+1}) \end{array} \right\}$	$\left\{ \begin{array}{l} b^q, M', a^q \in \sigma^{ss} : \phi(b^q) < \phi(M') + 1 \\ \phi(b^q) < \phi(a^q) \\ \phi(M') < \phi(a^q) \end{array} \right\}$ .

In [7, 8], it was shown that  $\text{Stab}(D^b(Q)) = \bigcup_{(A,B,C) \in \mathfrak{T}} (A, B, C)$  (see [8, (92)]). Following (35) we reorganize the union as follows (see [8, (95) and Section 5]):

$$(38) \quad \text{Stab}(D^b(Q)) = \mathfrak{T}_a^{st} \cup (-, M, -) \cup (-, M', -) \cup \mathfrak{T}_b^{st} \quad \mathfrak{T}_a^{st} \cap \mathfrak{T}_b^{st} = \emptyset,$$

where  $\mathfrak{T}_a^{st}$ , resp.  $\mathfrak{T}_b^{st}$ , is the union of all the subsets described in the first, resp. second, column of table (36) as  $j, p, m$ , resp.  $n, q, i$ , vary in  $\mathbb{Z}$ . The union of all the subsets described in the first, resp. second, column of table (37) as  $p$  vary in  $\mathbb{Z}$  will be denoted by  $(-, M, -)$ , resp.  $(-, M', -)$ .

More generally, the subset  $(A, B, C) \subset \text{Stab}(\mathcal{T})$  is described in [8, Proposition 2.7] for any  $\mathcal{T}$  as those stability conditions such that  $A, B, C$  are semistable and the phases  $\phi(A), \phi(B), \phi(C)$  satisfy the inequalities in (34) with  $y_0, y_1, y_2$  replaced by  $\phi(A), \phi(B), \phi(C)$ .

**6.3. First remarks on  $C_{i,\sigma}(\mathcal{T}), C_{i,\sigma\sigma}(\mathcal{T})$ .** From 2, Proposition 4, and Remark 28 we see that:

$$\begin{aligned} \alpha^m \in C_{0,\sigma\sigma}(\mathcal{T}) &\text{ iff } \{M', a^m, b^{m+1}\} \subset \sigma^{ss}; & \beta^m \in C_{0,\sigma\sigma}(\mathcal{T}) &\text{ iff } \{M, a^m, b^m\} \subset \sigma^{ss}; \\ A \in C_{1,\sigma\sigma}(\mathcal{T}) &\text{ iff } \{a^m : m \in \mathbb{Z}\} \subset \sigma^{ss}; & B \in C_{1,\sigma\sigma}(\mathcal{T}) &\text{ iff } \{b^m : m \in \mathbb{Z}\} \subset \sigma^{ss}; \\ A \in C_{1,\sigma}(\mathcal{T}) &\text{ iff } |\{m \in \mathbb{Z} : a^m \in \sigma^{ss}\}| = \infty; & B \in C_{1,\sigma}(\mathcal{T}) &\text{ iff } |\{m \in \mathbb{Z} : b^m \in \sigma^{ss}\}| = \infty. \end{aligned}$$

In particular, it follows (taking into account (2))

$$(39) \quad C_{D^b(pt),\sigma\sigma}(\mathcal{T}) = C_{D^b(pt),\mathbb{K}}(\mathcal{T}) \iff C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T}).$$

$$(40) \quad \left\{ \sigma : \left| C_{D^b(pt),\sigma\sigma}(\mathcal{T}) \right| < \infty \right\} = \left\{ \sigma : |C_{1,\sigma}(\mathcal{T})| = 0 \right\}.$$

Furthermore, if infinitely many genus zero nc curves are in  $C_{0,\sigma\sigma}(\mathcal{T})$ , then either for infinitely many  $m$  we have  $\alpha^m \in C_{0,\sigma\sigma}(\mathcal{T})$  and hence  $\{M', a^m, b^{m+1}\} \subset \sigma^{ss}$  or for infinitely many  $m$  we have  $\beta^m \in C_{0,\sigma\sigma}(\mathcal{T})$  and hence  $\{M, a^m, b^m\} \subset \sigma^{ss}$ , and it follows that  $C_{1,\sigma}(\mathcal{T}) = C_1(\mathcal{T})$ . Thus we deduce:

$$(41) \quad C_{1,\sigma}(\mathcal{T}) \neq \{A, B\} \implies |C_{0,\sigma\sigma}(\mathcal{T})| < \infty.$$

6.4. **Study of  $C_{i,\sigma}(\mathcal{T})$ ,  $C_{i,\sigma\sigma}(\mathcal{T})$ ,  $C_{D^b(pt),\sigma\sigma}(\mathcal{T})$  as  $\sigma \in \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  and  $i \geq -1$ .** Recall that  $\mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  is in (38). It will be enough to study only one of both the cases,  $\sigma \in \mathfrak{T}_a^{st}$  or  $\sigma \in \mathfrak{T}_b^{st}$ , because there is an auto-equivalence  $\zeta$  such that  $\zeta(\mathfrak{T}_a^{st}) = \mathfrak{T}_b^{st}$ . More precisely, there is  $\zeta$  such that

$$(42) \quad \zeta \in \text{Aut}_{\mathbb{K}}(\mathcal{T}) \quad \zeta(a^m) \cong b^m \quad \zeta(b^m) \cong a^{m-1} \quad \zeta(M) \cong M' \quad \zeta(M') \cong M$$

for each  $m \in \mathbb{Z}$ . Indeed from [6, Proposition 8.4] we see that there is an auto-equivalence  $\zeta \in \text{Aut}(\mathcal{T})$  such that  $\zeta(a^m) \cong b^m$ ,  $\zeta(b^m) \cong a^{m-1}$  for each  $m \in \mathbb{Z}$ . On the other hand, from the full exceptional triples  $(M', a^m, a^{m+1})$ ,  $(M, b^m, b^{m+1})$ ,  $(a^m, a^{m+1}, M)$ ,  $(b^m, b^{m+1}, M')$  it follows that  $\zeta(M') \cong M[p]$ ,  $\zeta(M) \cong M'[q]$  for some  $p, q \in \mathbb{Z}$ . However, due to Corollary 39 we have also  $0 \neq \text{hom}(M', a^m) = \text{hom}(\zeta(M'), \zeta(a^m)) = \text{hom}(M[p], b^m)$  and from Corollaries 39, 40 we see that  $p = 0$ . Also we have  $0 \neq \text{hom}(M, b^m) = \text{hom}(\zeta(M), \zeta(b^m)) = \text{hom}(M'[q], a^{m-1})$  and again Corollaries 39, 40 imply  $q = 0$ . From Proposition 4 and (42) we obtain

$$(43) \quad \zeta(\alpha^m) = \beta^m \quad \zeta(\beta^m) = \alpha^{m-1} \quad \zeta(A) = B \quad \zeta(B) = A.$$

Using (12) in Remark 17 and Subsection 6.3 we see that

**Remark 43.** For any  $\sigma \in \text{Stab}(\mathcal{T})$  and any  $m \in \mathbb{Z}$  we have

$$(44) \quad M \in \sigma^{ss} \iff M' \in (\zeta \cdot \sigma)^{ss} \Rightarrow \phi_{\zeta \cdot \sigma}(M') = \phi_{\sigma}(M)$$

$$(45) \quad M' \in \sigma^{ss} \iff M \in (\zeta \cdot \sigma)^{ss} \Rightarrow \phi_{\zeta \cdot \sigma}(M) = \phi_{\sigma}(M')$$

$$(46) \quad a^m \in \sigma^{ss} \iff b^m \in (\zeta \cdot \sigma)^{ss} \Rightarrow \phi_{\zeta \cdot \sigma}(b^m) = \phi_{\sigma}(a^m)$$

$$(47) \quad b^m \in \sigma^{ss} \iff a^{m-1} \in (\zeta \cdot \sigma)^{ss} \Rightarrow \phi_{\zeta \cdot \sigma}(a^{m-1}) = \phi_{\sigma}(b^m)$$

$$(48) \quad \alpha^m \in C_{0,\sigma\sigma}(\mathcal{T}) \iff \beta^m \in C_{0,(\zeta \cdot \sigma)(\zeta \cdot \sigma)}(\mathcal{T})$$

$$(49) \quad \beta^m \in C_{0,\sigma\sigma}(\mathcal{T}) \iff \alpha^{m-1} \in C_{0,(\zeta \cdot \sigma)(\zeta \cdot \sigma)}(\mathcal{T})$$

$$(50) \quad A \in C_{1,\sigma\sigma}(\mathcal{T}) \iff B \in C_{1,(\zeta \cdot \sigma)(\zeta \cdot \sigma)}(\mathcal{T}) \quad A \in C_{1,\sigma}(\mathcal{T}) \iff B \in C_{1,\zeta \cdot \sigma}(\mathcal{T}).$$

**Lemma 44.** Let  $p \in \mathbb{Z}$ . In any of the following cases holds  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \emptyset$ :

(a)  $a^p, a^{p+1} \in \sigma^{ss}$  and  $\phi(a^{p+1}) > \phi(a^p) + 1$ . In fact in this case  $a^j \notin \sigma^{ss}$  for  $j \notin \{p, p+1\}$ ,  $b^j \notin \sigma^{ss}$  for  $j \neq p+1$ , and it follows that  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^{p+1}\}$ ,  $\beta^{p+1} \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M, b^{p+1}\} \subset \sigma^{ss}$ ,  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M', b^{p+1}\} \subset \sigma^{ss}$ .

(b)  $b^p, b^{p+1} \in \sigma^{ss}$  and  $\phi(b^{p+1}) > \phi(b^p) + 1$ . In this case  $b^j \notin \sigma^{ss}$  for  $j \notin \{p, p+1\}$ ,  $a^j \notin \sigma^{ss}$  for  $j \neq p$ , and it follows that  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$ ,  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M, a^p\} \subset \sigma^{ss}$ ,  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M', a^p\} \subset \sigma^{ss}$ .

(c)  $a^p, b^{p+1} \in \sigma^{ss}$  and  $\phi(b^{p+1}) > \phi(a^p) + 1$ . In fact, in this case  $b^j, a^j \notin \sigma^{ss}$  for  $j \notin \{p, p+1\}$ .

*Proof.* (a) If  $b^j \in \sigma^{ss}$  for  $j \leq p$ , then from  $\text{hom}(b^j, a^p) \neq 0$  in (28) and axiom (2) in Definition 14 it follows that  $\phi(b^j) \leq \phi(a^p)$ , and therefore axiom (2) in Definition 14 implies  $\text{hom}^1(a^{p+1}, b^j) = 0$ , which contradicts (29).

If  $b^j \in \sigma^{ss}$  for  $j > p+1$ , then from  $\text{hom}(a^{p+1}, b^j) \neq 0$  in (29) and axiom (2) in Definition 14 it follows that  $\phi(a^{p+1}) \leq \phi(b^j)$ , and therefore axiom (2) in Definition 14 implies  $\text{hom}^1(b^j, a^p) = 0$ , which contradicts (28). The cases of  $a^j$  for  $j < p$ ,  $j > p+1$  are obtained by similar arguments, relying on (30).

The statement for  $C_{0,\sigma\sigma}(\mathcal{T})$  follows from the already proved part and Subsection 6.3.

(b) follows from (a) due to the auto-equivalence  $\zeta$  with (42) and (43).

(c) If  $a^j \in \sigma^{ss}$  for some  $j < p$ , then using Corollary 39 and axiom (2) in Definition 14 we deduce  $\phi(a^j) \leq \phi(a^p)$  and therefore  $\phi(a^j) + 1 < \phi(b^{p+1})$ . Hence via axiom (2) in Definition 14 we have  $\text{hom}^1(b^{p+1}, a^j) = 0$ , which contradicts (28).

If  $a^j \in \sigma^{ss}$  for some  $j > p + 1$ , then by  $\text{hom}(b^{p+1}, a^j) \neq 0$  in (28) we get  $\phi(b^{p+1}) \leq \phi(a^j)$  and therefore  $\text{hom}^1(a^j, a^p) = 0$ , which contradicts (30).

If  $b^j \in \sigma^{ss}$  for some  $j < p$ , then 28 ensures  $\phi(b^j) \leq \phi(a^p)$ , which however implies  $\text{hom}^1(b^{p+1}, b^j) = 0$ , which contradicts (31).

If  $b^j \in \sigma^{ss}$  for some  $j > p + 1$ , then by  $\text{hom}(b^{p+1}, b^j) \neq 0$  in (31) we get  $\phi(b^{p+1}) \leq \phi(b^j)$  and therefore  $\text{hom}^1(b^j, a^p) = 0$ , which contradicts (28).  $\square$

We will prove Propositions 46, 54 and 51. From these follow tables (103), (105) and then using Remark 43 follows tables (104), (106).

We will prove also Proposition 47, from which follows

**Corollary 45.** *If  $\sigma \in \mathfrak{T}_b^{st}$  and  $M, M' \in \sigma^{ss}$ , then  $\phi(M) < \phi(M')$ . If  $\sigma \in \mathfrak{T}_a^{st}$  and  $M, M' \in \sigma^{ss}$ , then  $\phi(M) > \phi(M')$ .*

Now we start proving tables (103), (105). We first determine completely the behavior of the stability and semi-stability of the curve  $B$  in  $\mathfrak{T}_b^{st}$ , and mention something about  $A$ .

**Proposition 46.** *Let  $\sigma \in \mathfrak{T}_b^{st}$ .*

*If  $\sigma \in (b^q, a^q, b^{q+1})$  for some  $q$ , then  $C_{1,\sigma}(\mathcal{T}) = C_{1,\sigma\sigma}(\mathcal{T}) = \emptyset$ .*

*If  $\sigma \in (M, b^q, b^{q+1})$  or  $\sigma \in (b^q, b^{q+1}, M')$  for some  $q$ , then for  $\phi(b^{q+1}) > \phi(b^q) + 1$  we have  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \emptyset$  and otherwise for  $\phi(b^{q+1}) \leq \phi(b^q) + 1$  holds  $B \in C_{1,\sigma\sigma}(\mathcal{T})$ .*

*Proof.* All the cases are in the second column of table (36) and therefore  $b^q, b^{q+1} \in \sigma^{ss}$ . From Lemma 44 it follows that  $A \notin C_{1,\sigma}(\mathcal{T})$  and  $B \notin C_{1,\sigma}(\mathcal{T})$ , when  $\phi(b^{q+1}) > \phi(b^q) + 1$ , furthermore for  $\sigma \in (b^q, a^q, b^{q+1})$  we see in table (36) that  $\phi(b^{q+1}) > \phi(b^q) + 1$  is automatically satisfied.

It remains the case when  $\sigma \in (M, b^q, b^{q+1})$  or  $\sigma \in (b^q, b^{q+1}, M')$  for some  $q$  and  $\phi(b^{q+1}) \leq \phi(b^q) + 1$ . In this case Corollary 37 ensure  $B \in C_{1,\sigma\sigma}(\mathcal{T})$ .  $\square$

Now we discuss the behavior of  $\phi_\sigma(M)$ ,  $\phi_\sigma(M')$  for some stability conditions.

**Proposition 47.** *Let  $p \in \mathbb{Z}$ .*

(a) *Let  $\sigma \in (b^p, a^p, b^{p+1})$ . If  $M, M' \in \sigma^{ss}$ , then  $\phi(M) < \phi(M') \leq \phi(M) + 1$ . Furthermore:*

$$(51) \quad \begin{aligned} M \in \sigma^{ss} &\iff \phi(a^p) - 1 \leq \phi(b^p) < \phi(a^p) \\ M' \in \sigma^{ss} &\iff \phi(b^{p+1}) - 1 \leq \phi(a^p) < \phi(b^{p+1}). \end{aligned}$$

*For example one can choose  $\sigma \in (b^p, a^p, b^{p+1})$  with  $\phi(b^p) = t$ ,  $\phi(a^p) = t + 1$ ,  $\phi(b^{p+1}) = t + 2$  and in this case both inequalities in (51) hold and  $\phi(M) = t$ ,  $\phi(M') = t + 1$ .*

(b) *Let  $\sigma \in (M, b^p, b^{p+1})$  or  $\sigma \in (b^p, b^{p+1}, M')$ . If  $M, M' \in \sigma^{ss}$  then  $\phi(M) < \phi(M')$ .*

*Proof.* (a) Let  $M, M' \in \sigma^{ss}$ . Combining the inequalities  $\phi(b^p) + 1 < \phi(b^{p+1})$  in (36) with the  $\phi(a^p) < \phi(b^{p+1})$

following inequalities due to (26), (27):  $\phi(b^{p+1}) - 1 \leq \phi(M') \leq \phi(a^p)$  we we obtain the following  $\phi(a^p) - 1 \leq \phi(M) \leq \phi(b^p)$

equivalent system of inequalities:  $\phi(a^p) - 1 \leq \phi(M) \leq \phi(b^p) < \phi(b^{p+1}) - 1 \leq \phi(M') \leq \phi(a^p)$ . In particular,  $\phi(M) < \phi(M') \leq \phi(M) + 1$ , if  $M, M' \in \sigma^{ss}$ .

Conversely, let  $\sigma \in (b^p, a^p, b^{p+1})$  and some of the inequalities (51) hold. The constants  $\alpha, \gamma$  defined in [8, Lemma 2.7] for the triple  $(b^p, a^p, b^{p+1})$  are  $\alpha = \gamma = -1$ , which follows from Corollary 39. From Corollary 37 we see that if the first, resp. the second, inequality in (51) holds, then the exceptional objects in the extension closures of  $(b^p, a^p[-1])$ , resp. in  $(a^p, b^{p+1}[-1])$ , are in  $\sigma^{ss}$ . Therefore  $M \in \sigma^{ss}$ , resp.  $M' \in \sigma^{ss}$  (see (33)). The last sentence follows also from (33).

(b) Let  $M, M' \in \sigma^{ss}$ . If  $\sigma \in (M, b^p, b^{p+1})$ , then from table (36) we have  $\phi(M) + 1 < \phi(b^{p+1})$ , and from (27) it follows that  $\phi(b^{p+1}) \leq \phi(M') + 1$ . Therefore  $\phi(M) < \phi(M')$ . If  $\sigma \in (b^p, b^{p+1}, M')$ , then from table (36) we have  $\phi(b^p) < \phi(M')$ , and from (26) it follows that  $\phi(M) \leq \phi(b^p)$ . Therefore  $\phi(M) < \phi(M')$ .  $\square$

**Corollary 48.** *Let  $\sigma \in (b^p, a^p, b^{p+1})$ , then  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$  and  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\phi(b^{p+1}) - 1 \leq \phi(a^p) < \phi(b^{p+1})$ ,  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\phi(a^p) - 1 \leq \phi(b^p) < \phi(a^p)$ . In particular, the first row of table (107) follows, taking into account also Proposition (46).*

*Let  $\sigma \in (a^p, b^{p+1}, a^{p+1})$ , then  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^{p+1}\}$  and  $\beta^{p+1} \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\phi(a^{p+1}) - 1 \leq \phi(b^{p+1})$ ,  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\phi(b^{p+1}) - 1 \leq \phi(a^p)$ . In particular, the second row of table (107) follows.*

*Proof.* Let  $\sigma \in (b^p, a^p, b^{p+1})$ . In table 36 we have  $b^p, a^p, b^{p+1} \in \sigma^{ss}$  and  $\phi(b^p) + 1 < \phi(b^{p+1})$ .  $\phi(b^p) < \phi(a^p)$   
 $\phi(a^p) < \phi(b^{p+1})$

From Lemma 44 (b) and Proposition 47 (a) we obtain the statement.

The second part follows from the first with the help of (42), (43) and Remark 43.  $\square$

**Corollary 49.** *Let  $\sigma \in (b^p, b^{p+1}, M')$  and  $\phi(b^{p+1}) > \phi(b^p) + 1$ . Then*

(a)  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$  and

$$(52) \quad C_{0,\sigma\sigma}(\mathcal{T}) \neq \emptyset \iff \phi(M') + 1 > \phi(b^{p+1}) \geq \phi(M') \iff \alpha^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff a^p \in \sigma^{ss}.$$

*Furthermore for  $\phi(M') + 1 > \phi(b^{p+1}) \geq \phi(M')$  we have  $\phi(a^p) = \arg_{[\phi(M'), \phi(M')+1]}(Z(b^{p+1}) + Z(M'))$  and  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff \arg_{[\phi(M'), \phi(M')+1]}(Z(b^{p+1}) + Z(M')) \leq \phi(b^p) + 1$ .*

(b)  $M \in \sigma^{ss}$  if and only if  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$ , which in turn by (a) is equivalent to  $\phi(M') + 1 > \phi(b^{p+1}) \geq \phi(M')$  and  $\arg_{[\phi(M'), \phi(M')+1]}(Z(b^{p+1}) + Z(M')) \leq \phi(b^p) + 1$ .

(c) From (a) and (b) follow the third and the second row of table (103) (taking into account Proposition 46).

*Proof.* Recall that in table (36) we have  $\phi(b^{p+1}) < \phi(M') + 1$ . From Lemma 44 (b) it follows that  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$ ,  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M, a^p\} \subset \sigma^{ss}$ ,  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M', a^p\} \subset \sigma^{ss}$ . In particular  $C_{0,\sigma\sigma}(\mathcal{T}) \neq \emptyset$  iff  $a^p \in \sigma^{ss}$ . On the other hand, if  $a^p \in \sigma^{ss}$ , then (26), (29) imply that  $\phi(M') \leq \phi(a^p) \leq \phi(b^{p+1})$  and vice versa, if  $\phi(b^{p+1}) \geq \phi(M')$ , then Corollary 37 and (27) imply that  $\alpha^p = \langle M', a^p, b^{p+1} \rangle \in C_{0,\sigma\sigma}(\mathcal{T})$  and hence  $a^p \in \sigma^{ss}$ . So we proved (52). Before continuing with analyzing the case  $\phi(b^{p+1}) \geq \phi(M')$ , we note that when  $\phi(b^{p+1}) < \phi(M')$  we have  $M \notin \sigma^{ss}$ . Indeed, the non-vanishing  $\text{hom}(M, b^p) \neq 0$  in (26) combined with inequalities  $\phi(b^{p+1}) < \phi(M')$ ,  $\phi(b^{p+1}) > \phi(b^p) + 1$ , and  $M \in \sigma^{ss}$ , recalling (2) in Definition 14, would imply  $\text{hom}^1(M', M) = 0$ , which contradicts (32). Recalling that  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M, a^p\} \subset \sigma^{ss}$  and using the already proved (52) we see that  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $M \in \sigma^{ss}$ .

Let  $\phi(b^{p+1}) \geq \phi(M')$  and  $\phi(b^{p+1}) > \phi(b^p) + 1$ . It remains to determine when  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$ , i.e. when  $M \in \sigma^{ss}$ . We already explained that  $a^p \in \sigma^{ss}$  and  $\phi(M') \leq \phi(a^p) \leq \phi(b^{p+1})$ . In table

(36) we have also  $\phi(b^p) < \phi(b^{p+1})$   
 $\phi(b^p) < \phi(M')$  . Therefore  $\phi(M') \leq \phi(a^p) \leq \phi(b^{p+1}) < \phi(M') + 1$ .  
 $\phi(b^{p+1}) < \phi(M') + 1$

Recalling that  $Z(a^p) = Z(b^{p+1}) + Z(M')$  (the first triangle in (33)) we deduce that  $\phi(a^p) = \arg_{[\phi(M'), \phi(M')+1]}(Z(b^{p+1}) + Z(M'))$ , Furthermore,  $\phi(a^p) = \phi(b^{p+1})$  iff  $\phi(b^{p+1}) = \phi(M')$  otherwise  $\phi(M') < \phi(a^p) < \phi(b^{p+1})$ . If  $\phi(b^{p+1}) = \phi(M')$ , then  $\phi(a^p) = \phi(b^{p+1})$  and therefore, taking into account  $\phi(b^p) + 1 < \phi(b^{p+1})$ , we deduce  $\phi(a^p[-1]) > \phi(b^p)$  and then the second triangle in (33) implies  $M \notin \sigma^{ss}$ . Otherwise  $\phi(b^p) < \phi(M') < \phi(a^p) < \phi(b^{p+1})$  and from table (36) it follows that  $\sigma \in (b^p, a^p, b^{p+1})$  and now applying Corollary 48 we complete the proof.  $\square$

**Corollary 50.** *Let  $\sigma \in (M, b^p, b^{p+1})$  and  $\phi(b^{p+1}) > \phi(b^p) + 1$ . Then*

(a)  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$  and

$$(53) \quad C_{0,\sigma\sigma}(\mathcal{T}) \neq \emptyset \iff \phi(b^p) > \phi(M) \geq \phi(b^p) - 1 \iff \beta^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff a^p \in \sigma^{ss}.$$

Furthermore for  $\phi(b^p) > \phi(M) \geq \phi(b^p) - 1$  we have  $\phi(a^p) = \arg_{(\phi(M), \phi(M)+1]}(Z(b^p) - Z(M))$  and  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff \phi(b^{p+1}) - 1 \leq \arg_{(\phi(M), \phi(M)+1]}(Z(b^p) - Z(M))$ .

(b)  $M' \in \sigma^{ss}$  if and only if  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$ , which in turn by (a) is equivalent to  $\phi(b^p) > \phi(M) \geq \phi(b^p) - 1$  and  $\phi(b^{p+1}) - 1 \leq \arg_{(\phi(M), \phi(M)+1]}(Z(b^p) - Z(M))$ .

(c) From (a) and (b) follow the second and the third row of table (105) (taking into account Proposition 46).

*Proof.* In table (36) we have  $\phi(M) < \phi(b^p)$ . From Lemma 44 (b) it follows that  $C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}$ ,  $\beta^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M, a^p\} \subset \sigma^{ss}$ ,  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M', a^p\} \subset \sigma^{ss}$ . In particular  $C_{0,\sigma\sigma}(\mathcal{T}) \neq \emptyset$  iff  $a^p \in \sigma^{ss}$ . On the other hand, if  $a^p \in \sigma^{ss}$ , then (27), (28) and (29) imply that  $\phi(b^p) \leq \phi(a^p) \leq \phi(M) + 1$ ,  $\phi(a^p) \leq \phi(b^{p+1})$  and vice versa, if  $\phi(M) \geq \phi(b^p) - 1$ , then Corollary 37 implies that  $\beta^p = \langle M, b^p, a^p \rangle \in C_{0,\sigma\sigma}(\mathcal{T})$  and hence  $a^p \in \sigma^{ss}$ . Thus, we proved (53).

Note that when  $\phi(M) < \phi(b^p) - 1$  we have  $M' \notin \sigma^{ss}$ . Indeed, the non-vanishing  $\text{hom}^1(b^{p+1}, M') \neq 0$  in (27) combined with inequalities  $\phi(b^p) - 1 > \phi(M)$ ,  $\phi(b^{p+1}) > \phi(b^p) + 1$ , and  $M' \in \sigma^{ss}$ , recalling (2) in Definition 14, would imply  $\text{hom}^1(M', M) = 0$ , which contradicts (32). Having (53) and  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$  iff  $\{M', a^p\} \subset \sigma^{ss}$  we see that  $M' \in \sigma^{ss} \iff \alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$ .

Let  $\phi(M) \geq \phi(b^p) - 1$  and  $\phi(b^{p+1}) > \phi(b^p) + 1$ . It remains to determine when  $\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T})$ , i.e. when  $M' \in \sigma^{ss}$ ? We already explained that  $a^p \in \sigma^{ss}$  and  $\phi(b^p) \leq \phi(a^p) \leq \phi(M) + 1$ ,

$\phi(M) < \phi(b^p)$   
 $\phi(a^p) \leq \phi(b^{p+1})$ . In table (36) we have also  $\phi(M) + 1 < \phi(b^{p+1})$  . Therefore  $\phi(M) < \phi(b^p) \leq \phi(b^p) < \phi(b^{p+1})$

$\phi(a^p) \leq \phi(M) + 1$ . Recalling that  $Z(a^p) = Z(b^p) - Z(M)$  (the second triangle in (33)) we deduce that  $\phi(a^p) = \arg_{(\phi(M), \phi(M)+1]}(Z(b^p) - Z(M))$ , Furthermore,  $\phi(a^p) = \phi(b^p)$  iff  $\phi(b^p) - 1 = \phi(M)$  otherwise  $\phi(b^p) < \phi(a^p) < \phi(M) + 1 < \phi(b^{p+1})$ . If  $\phi(b^p) - 1 = \phi(M)$ , then  $\phi(a^p) = \phi(b^p)$  and therefore, taking into account  $\phi(b^p) + 1 < \phi(b^{p+1})$ , we deduce  $\phi(b^{p+1}[-1]) > \phi(a^p)$  and then the first triangle in (33) implies  $M' \notin \sigma^{ss}$ . Otherwise  $\phi(b^p) < \phi(a^p) < \phi(M) + 1 < \phi(b^{p+1})$  and from table (36) it follows that  $\sigma \in (b^p, a^p, b^{p+1})$  and now applying Corollary 48 we complete the proof.  $\square$

In Proposition 46 and Corollaries 50, 49 we filled in some boxes of tables (103), (105). The rest boxes in these tables are proved in the next two propositions.

**Proposition 51.** *Let  $\sigma \in (b^n, b^{n+1}, M')$  for some  $n \in \mathbb{Z}$  and  $\phi(b^{n+1}) \leq \phi(b^n) + 1$ . Then  $A \notin C_{1,\sigma\sigma}(\mathcal{T})$ .*

(a) If  $\phi(b^{n+1}) < \phi(b^n) + 1$ , we denote<sup>4</sup>

$$(54) \quad t = \arg_{(\phi(b^n)-1, \phi(b^n))}(Z(b^n) - Z(b^{n+1})),$$

and then consider two subcases:

(a.1)  $\phi(M') \geq t + 1$ : in this sub-case  $a^j \notin \sigma^{ss}$  for all  $j \in \mathbb{Z}$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$  and  $A \notin C_{1,\sigma}(\mathcal{T})$ ;

(a.2)  $\phi(M') < t + 1$ : in this sub-case if we denote  $U = \arg_{(t, t+1)}(Z(M') + Z(b^{n+1}) - Z(b^n))$ , then it holds  $U > \phi(M')$ ,  $M \in \sigma^{ss}$ ,  $U = \phi(M) + 1$  and there are uniquely determined integers  $N \leq u$  such that

$$(55) \quad t < \phi(b^N) < \phi(M') \leq \phi(b^{N+1}) < t + 1$$

$$(56) \quad t < \phi(b^u) \leq U < \phi(b^{u+1}) < t + 1,$$

and for these integers  $N, u$  holds  $a^j \in \sigma^{ss} \iff N \leq j \leq u$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : N \leq j \leq u\}$ . In particular here we have  $A \notin C_{1,\sigma}(\mathcal{T})$  again.

(b) If  $\phi(b^{n+1}) = \phi(b^n) + 1$ , then we consider three sub-cases

(b.1)  $\phi(M') > \phi(b^n) + 1$ : in this case  $a^j \notin \sigma^{ss}$  for all  $j \in \mathbb{Z}$  and  $A \notin C_{1,\sigma}(\mathcal{T})$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ . Furthermore, in this case  $M \notin \sigma^{ss}$ .

(b.2)  $\phi(M') < \phi(b^n) + 1$ : in this case  $a^j \in \sigma^{ss}$  iff  $j = n$ ,  $A \notin C_{1,\sigma}(\mathcal{T})$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^n, \beta^n\}$ . Furthermore, in this case  $M \in \sigma^{ss}$  and  $\phi(M') < \phi(M) + 1$ .

(b.3)  $\phi(M') = \phi(b^{n+1}) = \phi(b^n) + 1$ : in this case  $M \in \sigma^{ss}$ ,  $\phi(M') = \phi(M) + 1$ ,  $j \geq n$  iff  $a^j \in \sigma^{ss}$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \geq n\}$ ,  $A \in C_{1,\sigma}(\mathcal{T})$ .

In particular, follows the rest of table (103) and then from Remark 42 follows table (104),

*Proof.* From Proposition 46 we see that  $\{b^j\}_{j \in \mathbb{Z}} \subset \sigma^{ss}$ . From Corollary 39 we see that  $\text{hom}(b^j, a^j) \neq 0$ ,  $\text{hom}(a^j, b^{j+1}) \neq 0$ ,  $\text{hom}(M', a^j) \neq 0$  for any  $j \in \mathbb{Z}$  and it follows

$$(57) \quad \forall j \in \mathbb{Z} \quad a^j \in \sigma^{ss} \Rightarrow \phi(b^j) \leq \phi(a^j) \leq \phi(b^{j+1}) \quad \text{and} \quad \phi(M') \leq \phi(a^j).$$

So, if  $a^j \in \sigma^{ss}$  for some  $j \in \mathbb{Z}$ , then  $\phi(M') \leq \phi(a^j)$  and from table (36) it follows that  $\phi(b^n) < \phi(a^j)$ , hence  $\text{hom}(a^j, b^n) = 0$ . Now the non-vanishing  $\text{hom}(a^p, b^{q+1}) \neq 0$  for  $p \leq q$  in corollary 39 implies that  $j \geq n$ . So we see that

$$(58) \quad j < n \Rightarrow a^j \notin \sigma^{ss} \Rightarrow A \notin C_{1,\sigma\sigma}(\mathcal{T}).$$

(a) Here we have  $\phi(b^n) < \phi(b^{n+1}) < \phi(b^n) + 1$ . Recalling that (see [8, (84)])  $Z(\delta) = Z(b^n) - Z(b^{n+1})$  we see that choosing  $t$  as in (54) and combining the facts:  $\{b^j : j \in \mathbb{Z}\} \subset \sigma^{ss}$ ,  $\text{hom}(b^i, b^j) \neq 0$  for  $i \leq j$ ,  $\text{hom}^1(b^i, b^j) \neq 0$  for  $i > j + 1$  and [8, Corollaries 3.19, 3.18] one obtains:

$$(59) \quad 0 \neq Z(\delta) = |Z(\delta)| \exp(i\pi t) \quad \lim_{j \rightarrow +\infty} \phi(b^j) = t + 1 \quad \lim_{j \rightarrow -\infty} \phi(b^j) = t \quad \forall j \in \mathbb{Z} \quad \phi(b^j) < \phi(b^{j+1}).$$

Having (59) and (57) and using again [8, Corollary 3.19] we deduce:

$$(60) \quad \{Z(a^j)\}_{j \in \mathbb{Z}} \subset Z(\delta)_+^c \quad \lim_{k \rightarrow +\infty} \arg_{(t, t+1)}(Z(a^k)) = t + 1$$

$$\forall k \in \mathbb{Z} \quad t < \arg_{(t, t+1)}(Z(a^k)) < \arg_{(t, t+1)}(Z(a^{k+1})) < t + 1.$$

(a.1) If  $\phi(M') \geq t + 1$ , then using (57) and (59) it follows that  $a^j \notin \sigma^{ss}$  for any  $j \in \mathbb{Z}$  and hence in Subsection 6.3 we deduce that  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ .

<sup>4</sup>Recall that  $Z(b^n) - Z(b^{n+1}) = Z(\delta)$  - see Remark 41.

(a.2) So, assume that  $\phi(M') < t + 1$ . In table (36) we have  $\phi(b^n) < \phi(M')$  and due to (59) we see that for some  $N \in \mathbb{Z}$  holds  $n \leq N$  and (55).

Table (36) ensure that  $\sigma \in (b^N, b^{N+1}, M')$  and recalling that  $\text{hom}^1(b^{N+1}, M') \neq 0$  in (27) we can use Corollary 37 to deduce that

$$(61) \quad \alpha^N = \langle b^{N+1}, M' \rangle = \langle a^N, b^{N+1}, M' \rangle \in C_{0, \sigma\sigma}(\mathcal{T}).$$

For  $j < N$  we use (57) and  $\phi(b^N) < \phi(M')$  to deduce that  $a^j \notin \sigma^{ss}$  for  $j < N$ . From the first triangle in (33) we have actually that for  $j > N$  holds  $Z(a^j) = Z(M') + Z(b^{j+1})$ ,  $a^j \in \mathcal{P}([\phi(M'), \phi(b^{j+1})]) \subset \mathcal{P}((t, t+1])$  hence from (55) and [8, Remark 2.1] we see that

$$(62) \quad N < j \Rightarrow \arg_{(t, t+1)}(Z(a^j)) < \arg_{(t, t+1)}(Z(b^{j+1})) = \phi(b^{j+1})$$

$$(63) \quad N < j, a^j \notin \sigma^{ss} \Rightarrow \phi_-(a^j) < \arg_{(t, t+1)}(Z(a^j)), \quad \phi(M') \leq \phi_-(a^j) \leq \phi(b^{j+1})$$

Furthermore, we will show (64) below with the help of [8, Lemma 7.2]. Let  $N < j$  and  $a^j \notin \sigma^{ss}$ . One of the five cases given there must appear. In case (a) of [8, Lemma 7.2] we have  $a^k \in \sigma^{ss}$ ,  $\phi_-(a^j) = \phi(a^k) + 1$  for some  $k < j - 1$ , however then (63) and (10) imply  $Z(a^k) \in Z(\delta)_-^c$ , which contradicts (60). In case (b) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(a^k)$  for some  $k > j$ , however then (63) (57), and (10) imply  $\arg_{(t, t+1)}(Z(a^k)) < \arg_{(t, t+1)}(Z(a^j))$  with  $k > j$ , which contradicts (60). In case (c) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(b^k) + 1$  for some  $k < j$ , which would imply  $Z(b^k) \in Z(\delta)_-^c$ , which contradicts (59). Case (d) of [8, Lemma 7.2] ensures  $\phi_-(a^j) = \phi(b^k)$  for some  $j < k$ , and then (63), (10) ensure  $\arg_{(t, t+1)}(Z(b^k)) = \phi(b^k) < \arg_{(t, t+1)}(Z(a^j))$ , which contradicts (62), (59) and  $j < k$ . So only case (e) in Lemma [8, Lemma 7.2] remains possible, hence

$$(64) \quad j > N, a^j \notin \sigma^{ss} \Rightarrow M \in \sigma^{ss}, \quad \phi_-(a^j) = \phi(M) + 1.$$

In Remark 41 we see that  $-Z(M) = Z(M') - Z(\delta)$  and (55) implies

$$(65) \quad -Z(M) \in Z(\delta)_+^c \quad \phi(M') < U = \arg_{(t, t+1)}(-Z(M)) < t + 1$$

increasing the superscript starting from  $N$  and taking into account (55) we get an integer  $u \geq N$  such that (56) holds. In (33) we see that  $Z(a^j) = Z(b^j) - Z(M)$  and from (56), (65) it follows that for  $j > u$  holds  $\arg_{(t, t+1)}(Z(a^j)) < \arg_{(t, t+1)}(Z(b^j))$ . Therefore using (57), (10) we deduce that  $a^j \in \sigma^{ss}$  for  $j > u$  implies  $\phi(a^j) < \phi(b^j)$  and  $\text{hom}(b^j, a^j) = 0$ , which contradicts (28). Hence we get (using also (64)):

$$(66) \quad j > u \Rightarrow a^j \notin \sigma^{ss} \Rightarrow M \in \sigma^{ss}, \phi_-(a^j) = \phi(M) + 1.$$

For  $N < j \leq u$  we have  $\phi(b^j) \leq U$  due to (56) and the formula  $Z(a^j) = Z(b^j) - Z(M)$  implies that  $\arg_{(t, t+1)}(Z(a^j)) \leq U$ . Using (63), (64) we see that  $N < j \leq u$ ,  $a^j \notin \sigma^{ss}$  imply  $\phi(M') = \phi(M) + 1 < U$ , which is impossible, since  $U = \arg_{(t, t+1)}(-Z(M))$  and we proved that  $M \in \sigma^{ss}$ . Therefore we proved  $a^j \in \sigma^{ss} \iff N \leq j \leq u$ . Taking into account that  $\{b^j : j \in \mathbb{Z}\} \subset \sigma^{ss}$  and  $M \in \sigma^{ss}$  we use Subsection 6.3 to deduce  $C_{0, \sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : N \leq j \leq u\}$ .

(b) Assume now  $\phi(b^{n+1}) = \phi(b^n) + 1$ . Let us denote  $\phi(b^n) = t$ , then  $\phi(b^{n+1}) = t + 1$ . Using that  $\{b^j\}_{j \in \mathbb{Z}} \subset \sigma^{ss}$ ,  $\text{hom}(b^n, b^m) \neq 0$  for  $n \leq m$  and  $\text{hom}^1(b^m, b^n) \neq 0$  for  $m > n + 1$  it follows that:

$$(67) \quad \phi(b^j) = t \text{ for } j \leq n \quad \phi(b^j) = t + 1 \text{ for } j \geq n + 1.$$

(b.1) If  $\phi(M') > t + 1$ , then in (57) we see that  $a^j \in \sigma^{ss}$  implies  $\phi(a^j) > t + 1$  and hence  $\text{hom}^1(a^j, b^k) = 0$  for small enough  $k$ , which contradicts (29). Furthermore, if  $M \in \sigma^{ss}$ , then from  $\text{hom}(M, b^n) \neq 0$  in (26) it follows that  $\phi(M) \leq \phi(b^n)$  and therefore  $\phi(M') > \phi(M) + 1$  which contradicts  $\text{hom}^1(M', M) \neq 0$  in (32).

(b.2) If  $\phi(M') < t + 1$ , due to table 36 we get  $t < \phi(M') < t + 1 = \phi(b^n) + 1$  and from Corollary 37 it follows  $\alpha^n = \langle M', a^n, b^{n+1} \rangle \in C_{0,\sigma\sigma}(\mathcal{T})$ ,  $a^n \in \sigma^{ss}$ . From  $Z(a^n) = Z(M') + Z(b^{n+1})$  in Remark 41 it follows that  $\phi(M') < \phi(a^n) < t + 1$ . Hence  $\phi(M') - 1 < \phi(a^n) - 1 < t = \phi(b^n)$ .

If  $a^j \in \sigma^{ss}$  for some  $j > n$ , then from (57) and (67), (10) we see that  $Z(b^{j+1})$ ,  $Z(a^j)$  are non-zero and collinear and from Remark 41 we know that  $Z(M') = Z(a^j) - Z(b^{j+1})$ , which contradicts  $t < \phi(M') < t + 1 = \phi(b^n) + 1$ . So for  $j > n$  we have  $a^j \notin \sigma^{ss}$ . Recalling (58) and Subsection 6.3 we deduce that  $\alpha^j, \beta^j \notin C_{0,\sigma\sigma}(\mathcal{T})$  for  $j \neq n$ . We will show that  $M \in \sigma^{ss}$  and this would imply that  $\beta^n = \langle M, a^n, b^n \rangle \in C_{0,\sigma\sigma}(\mathcal{T})$ .

Indeed, let  $j > n$ . From the first triangle in (33) we have  $a^j \in \mathcal{P}([\phi(M'), \phi(b^{j+1})]) \subset \mathcal{P}((t, t+1])$ , therefore  $\phi(M') \leq \phi_-(a^j) < \phi_+(a^j) \leq \phi(b^{j+1})$  and [8, Remark 2.1] implies

$$(68) \quad j > n \Rightarrow t < \phi_-(a^j) < \arg_{(t,t+1)}(Z(a^j)) < t + 1.$$

One of the five cases given in [8, Lemma 7.2] must appear. In case (a) of [8, Lemma 7.2] we have  $a^k \in \sigma^{ss}$ ,  $\phi_-(a^j) = \phi(a^k) + 1$  for some  $k < j - 1$ , and since we already showed that only  $j = n$  gives  $a^j \in \sigma^{ss}$  it would follow that  $\phi(a^n) + 1 = \phi_-(a^j)$ , which contradicts (68) and  $\phi(M') < \phi(a^n) < t + 1$ . In case (b) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(a^k)$  for some  $k > j$ , which contradicts the already proved instability of  $a^k$  for  $k > n$ . In cases (c) and (d) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(b^k) + 1$  or  $\phi_-(a^j) = \phi(b^k)$  for some  $k \in \mathbb{Z}$ , which contradicts (68) and the already proved fact that  $\phi(b^k) = t$  or  $\phi(b^k) = t + 1$ . So only case (e) in Lemma [8, Lemma 7.2] remains possible, hence  $M \in \sigma^{ss}$ .

Finally, from  $\text{hom}(M, b^n) \neq 0$  in (26) and  $\text{hom}^1(a^n, M) \neq 0$  in (27) we have  $\phi(a^n) - 1 \leq \phi(M) \leq t$  and then the second triangle in (33) implies that  $\phi(M') - 1 < \phi(a^n) - 1 < \phi(M) < t = \phi(b^n)$ .

(b.3) Let  $\phi(M') = t + 1$ , then for  $p \geq n$  the first distinguished triangle in (33) ensures that  $a^p \in \sigma^{ss}$  and  $\phi(a^p) = t + 1$ . In particular, we have  $\phi(a^n) = \phi(b^n) + 1$  and from the second triangle in (33) it follows that  $M \in \sigma^{ss}$ ,  $\phi(M) = \phi(b^n)$ . Recalling (58) and using Subsection 6.3 we obtain  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \geq n\}$ .  $\square$

**Lemma 52.** *Let us assume that we are in the situation of (a.1) of Proposition 51. In this case  $M \in \sigma^{ss} \iff \phi(M') = t + 1$ .*

*Proof.* First, we note that for  $\phi(M') > t + 1$  holds  $M \notin \sigma^{ss}$ . Indeed, if  $M \in \sigma^{ss}$ , then from (26) and (59) it follows that  $\phi(M) \leq t$  and  $\text{hom}^1(M', M) = 0$ , which contradicts (32).

So, it remains to consider the case  $\phi(M') = t + 1$ . Combining the triangles in (33) we get a diagram for any  $p \in \mathbb{Z}$ :

$$(69) \quad \begin{array}{ccccccc} 0 & \xrightarrow{\quad} & M'[-1] & \xrightarrow{\quad} & a^p[-1] & \xrightarrow{\quad} & M \\ & \searrow \text{dashed} & \swarrow & \searrow \text{dashed} & \swarrow & \searrow \text{dashed} & \swarrow \\ & & M'[-1] & & b^{p+1}[-1] & & b^p \end{array}$$

Therefore  $M$  is in the extension closure of  $M'[-1]$ ,  $b^{p+1}[-1]$ ,  $b^p$  for each  $p \in \mathbb{Z}$ . Since for any  $p \in \mathbb{Z}$  we have  $\phi(b^{p+1}[-1]) < t = \phi(M') - 1 < \phi(b^p)$ , it follows that  $M \in \mathcal{P}[\phi(b^{p+1}[-1]), \phi(b^p)]$  for each  $p \in \mathbb{Z}$ , hence  $\phi(b^{p+1}[-1]) \leq \phi_-(M)$  and  $\phi_+(M) \leq \phi(b^p)$  for each  $p \in \mathbb{Z}$ . Now using (59) and limiting  $p$  to  $+\infty$  we obtain  $t \leq \phi_-(M)$ , whereas limiting  $p$  to  $-\infty$  ensures  $\phi_+(M) \leq t$ , therefore  $\phi_-(M) = \phi_+(M) = t$  and indeed  $M \in \sigma^{ss}$ .  $\square$



**Proposition 53.** *Varying  $\sigma \in (b^n, b^{n+1}, M')$  in the region with  $\phi(b^{n+1}) < \phi(b^n) + 1$  and  $\phi(M') < t + 1$ , where  $t$  is as in (54), one can arrange that the pair  $(N, u)$  is  $(n, n + i)$  for any  $i \in \mathbb{Z}_{\geq 0}$ , where  $u, N$  are as in (a.2) of Proposition 51. Therefore for any  $i \in \mathbb{Z}_{\geq 0}$  we have  $\sigma$  such that  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : n \leq j \leq n + i\}$ . Hence  $|C_{0,\sigma\sigma}(\mathcal{T})|$  can be any positive even integer.*

*Proof.* From Remark 31 (b) and table (36) we know that the map (20) gives a homeomorphism from  $(b^n, b^{n+1}, M') \subset \text{Stab}(\mathcal{T})$  to  $\mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_0 - y_1 < 0 \\ y_0 - y_2 < 0 \\ y_1 - y_2 < 1 \end{array} \right\}$ . Taking into account also (10), it follows

that for any  $(r_0, r_1, r_2) \in \mathbb{R}_{>0}^3$  and any  $(y_0, y_1, y_2)$  satisfying the specified inequalities we have  $\sigma \in (b^n, b^{n+1}, M')$  such that  $Z(b^n) = r_0 \exp(i\pi y_0)$ ,  $Z(b^{n+1}) = r_1 \exp(i\pi y_1)$ ,  $Z(M') = r_2 \exp(i\pi y_2)$ . These arguments and the equalities  $Z(b^{n+1}) = Z(b^n) - Z(\delta)$ ,  $Z(\delta) = Z(M) + Z(M')$  imply that we have a stability condition  $\sigma \in (b^n, b^{n+1}, M')$ , such that  $Z(b^n)$ ,  $Z(b^{n+1})$ ,  $Z(M')$ ,  $-Z(M)$ ,  $Z(\delta)$  are as in Figure 1, and for this  $\sigma$  we have  $\phi(b^n) < \phi(M') < \phi(b^{n+1}) < \phi(b^n) + 1$ . Furthermore, choosing  $t$  as in (54), we see from Figure 1 that  $\phi(M') < t + 1$ . Hence the chosen stability condition  $\sigma$  falls in the case (a.2) of Proposition 51. In particular, denoting  $U = \arg_{(t,t+1)}(Z(M') + Z(b^{n+1}) - Z(b^n))$ , we have  $M \in \sigma^{ss}$ ,  $U = \phi(M) + 1$ , and from Figure 1 we see that not only  $\phi(b^n) < \phi(M') < \phi(b^{n+1})$  but also  $\phi(b^n) < \phi(M) + 1 = U < \phi(b^{n+1})$ . Now the defining properties of  $N, u$  in (55) and (56) imply that  $u = N = n$  for this  $\sigma$ .

The constructed  $\sigma$  corresponds to a point  $(r_0, r_1, r_2, y_0, y_1, y_2)$  via the map (20). Varying  $r_2$  to arbitrary  $s > 0$  without changing  $r_0, r_1, y_0, y_1, y_2$  corresponds, via the homeomorphism specified above, to varying the stability condition  $\sigma(s)$  in  $(b^n, b^{n+1}, M')$  and  $\sigma(s)$  falls in the case (a.2) of Proposition 51 for any  $s > 0$ . Since we change only  $r_2$ , the entities  $\phi_{\sigma(s)}(M')$ ,  $\phi_{\sigma(s)}(b^n)$ ,  $\phi_{\sigma(s)}(b^{n+1})$ ,  $Z_{\sigma(s)}(b^n)$ ,  $Z_{\sigma(s)}(b^{n+1})$  remain the same as for the initial  $\sigma = \sigma(r_2)$  for any  $s > 0$ . It follows that  $Z_{\sigma(s)}(\delta) = \arg_{(\phi_{\sigma(s)}(b^n)-1, \phi_{\sigma(s)}(b^n))}(Z_{\sigma(s)}(b^n) - Z_{\sigma(s)}(b^{n+1}))$  and  $\{Z_{\sigma(s)}(b^j)\}_{j \in \mathbb{Z}}$  remain the same as well for any  $s > 0$ . By the arguments used before (59) it follows that for any  $j \in \mathbb{Z}$  the real number  $\phi_{\sigma(s)}(b^j)$  equals  $\arg_{(t,t+1)}(Z_{\sigma(s)}(b^j))$  and therefore it does not change as  $s$  varies in  $\mathbb{R}_{>0}$ . Now from (55) we see that  $N(s)$  does not change as  $s$  varies in  $\mathbb{R}_{>0}$  and remains equal to  $n$ .

On the other hand from Figures 2 and (3) we see that  $U(s) = \phi_{\sigma(s)}(M) + 1$  increases monotone and continuously as  $s$  decreases, and that  $\lim_{s \rightarrow 0} U(s) = t + 1$ . Recalling (59) and the defining property of  $u(s)$  in (56) which hold for  $\sigma(s)$  and each  $s > 0$  we see that we can obtain an infinite sequence  $r_0 = s_0 > s_1 > s_2 > s_3 > \dots > 0$  such that  $u(s_i) = n + i$  for each  $i \in \mathbb{Z}_{\geq 0}$ . □

**Proposition 54.** *Let  $\sigma \in (M, b^n, b^{n+1})$  for some  $n \in \mathbb{Z}$  and  $\phi(b^{n+1}) \leq \phi(b^n) + 1$ . Then  $A \notin C_{1,\sigma\sigma}(\mathcal{T})$ .*



(a) *If  $\phi(b^{n+1}) < \phi(b^n) + 1$ , we denote  $t = \arg_{(\phi(b^n)-1, \phi(b^n))}(Z(b^n) - Z(b^{n+1}))$  and then must hold  $\phi(M) < t$  and consider two sub-cases:*

(a.1)  *$\phi(M) \leq t - 1$ , then in this sub-case  $a^j \notin \sigma^{ss}$  for all  $j \in \mathbb{Z}$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$  and  $A \notin C_{1,\sigma}(\mathcal{T})$ ;*

(a.2)  *$\phi(M) > t - 1$ : in this sub-case if we denote  $V = \arg_{(t,t+1)}(Z(b^n) - Z(b^{n+1}) - Z(M))$ , and then holds  $V < \phi(M) + 1$ ,  $M' \in \sigma^{ss}$ ,  $\phi(M') = V$ , and there are integers  $N \leq u$  such that*

$$(70) \quad t < \phi(b^N) < V \leq \phi(b^{N+1}) < t + 1$$

$$(71) \quad t < \phi(b^u) \leq \phi(M) + 1 < \phi(b^{u+1}) < t + 1.$$

FIGURE 1.  $n = N = u$ FIGURE 2.  $n = N < u$




FIGURE 3

and for these integers  $N, u$  holds  $a^j \in \sigma^{ss} \iff N \leq j \leq u$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : N \leq j \leq u\}$ . In particular here we have  $A \notin C_{1,\sigma}(\mathcal{T})$  again.

(b) If  $\phi(b^{n+1}) = \phi(b^n) + 1$ , then we consider three sub-cases

(b.1)  $\phi(M) < \phi(b^n) - 1$ : in this case  $a^j \notin \sigma^{ss}$  for all  $j \in \mathbb{Z}$  and  $A \notin C_{1,\sigma}(\mathcal{T})$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ . Furthermore, in this case  $M' \notin \sigma^{ss}$ .

(b.2)  $\phi(M) > \phi(b^n) - 1$ : in this case  $a^j \in \sigma^{ss}$  iff  $j = n$ ,  $A \notin C_{1,\sigma}(\mathcal{T})$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^n, \beta^n\}$ . Furthermore, in this case  $M' \in \sigma^{ss}$  and  $\phi(M') < \phi(M) + 1$ .

(b.3)  $\phi(M) = \phi(b^n) - 1$ : in this sub-case for  $j \leq n$  hold  $a^j \in \sigma^{ss}$ ,  $\phi(a^j) = \phi(b^j) = \phi(b^n)$  and for  $j > n$  hold  $a^j \notin \sigma^{ss}$ ,  $M' \in \sigma^{ss}$ ,  $\phi(M') = \phi(M) + 1$ . In particular,  $A \in C_{1,\sigma}(\mathcal{T})$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \leq n\}$ .

In particular, the rest of table (105) follows, and then from Remark 42 follows table (106),

*Proof.* From the proposition 46 we see that  $\{b^j\}_{j \in \mathbb{Z}} \subset \sigma^{ss}$ .

From Corollary 39 we see that  $\text{hom}(b^j, a^j) \neq 0$ ,  $\text{hom}(a^j, b^{j+1}) \neq 0$ ,  $\text{hom}^1(a^j, M) \neq 0$  for any  $j \in \mathbb{Z}$  and it follows

$$(72) \quad \forall j \in \mathbb{Z} \quad a^j \in \sigma^{ss} \Rightarrow \phi(b^j) \leq \phi(a^j) \leq \phi(b^{j+1}) \quad \text{and} \quad \phi(a^j) \leq \phi(M) + 1.$$

So, if  $a^j \in \sigma^{ss}$  for some  $j \in \mathbb{Z}$ , then  $\phi(a^j) \leq \phi(M) + 1$  and from table (36) it follows that  $\phi(a^j) < \phi(b^{n+1})$ , hence  $\text{hom}(b^{n+1}, a^j) = 0$ . Now the non-vanishing  $\text{hom}(b^p, a^q) \neq 0$  for  $p \leq q$  in Corollary 39 implies that  $n \geq j$ . So we see that

$$(73) \quad j > n \Rightarrow a^j \notin \sigma^{ss}$$

and therefore indeed  $A \notin C_{1,\sigma\sigma}$ .

(a) Here we have  $\phi(b^n) < \phi(b^{n+1}) < \phi(b^n) + 1$ . The arguments as in the beginning of the proof of (a) in Proposition 51 are applicable in this case and they imply the same formulas (59), (60) in this case. From (59) and the inequality  $\phi(M) + 1 < \phi(b^{n+1})$  in (36) imply that  $\phi(M) < t$ .

(a.1) If in this sub-case  $a^j \in \sigma^{ss}$  for some  $j \in \mathbb{Z}$ , (72) implies  $t < \phi(a^j)$  and  $\phi(a^j) \leq \phi(M) + 1 \leq t$ , which is a contradiction. Therefore in this case  $a^j \notin \sigma^{ss}$  for each  $j \in \mathbb{Z}$ .

(a.2) In this case  $t - 1 < \phi(M) < t$ . In table (36) we have  $\phi(M) + 1 < \phi(b^{n+1})$  and due to (59) we see that for some  $u \in \mathbb{Z}$  holds  $u \leq n$  and (71).

Table (36) ensure that  $\sigma \in (M, b^u, b^{u+1})$  and recalling that  $\text{hom}(M, b^u) \neq 0$  in (26) we can use Corollary 37 to deduce that

$$(74) \quad \beta^u = \langle M, b^u \rangle = \langle M, b^u, a^u \rangle \in C_{0, \sigma\sigma}(\mathcal{T}).$$

For  $j > u$  we use (72) and  $\phi(M) + 1 < \phi(b^{u+1})$  to deduce that  $a^j \notin \sigma^{ss}$  for  $j > u$ . From the second triangle in (33) we have for  $j < u$  holds  $Z(a^j) = Z(b^j) - Z(M)$ ,  $a^j \in \mathcal{P}([\phi(b^j), \phi(M) + 1]) \subset \mathcal{P}((t, t + 1])$  hence from (71) and [8, Remark 2.1] we see that

$$(75) \quad j < u \Rightarrow \phi(M) + 1 > \arg_{(t, t+1)}(Z(a^j)) > \arg_{(t, t+1)}(Z(b^j)) = \phi(b^j)$$

$$(76) \quad j < u, a^j \notin \sigma^{ss} \Rightarrow \phi_-(a^j) < \arg_{(t, t+1)}(Z(a^j)), \quad \phi(b^{j+1}) \leq \phi_-(a^j) \leq \phi(M) + 1$$

In Remark 41 we see that  $Z(M') = -Z(M) + Z(\delta)$  and (71) implies

$$(77) \quad Z(M') \in Z(\delta)_+^c \quad t < V = \arg_{(t, t+1)}(Z(M')) < \phi(M) + 1 < t + 1$$

decreasing the superscript starting from  $u$  and taking into account (71) we get an integer  $N \leq u$  such that (70) holds. In Remark 41 we see that  $Z(a^j) = Z(M') + Z(b^{j+1})$  and from (70) we see that

$$(78) \quad j < N \Rightarrow \arg_{(t, t+1)}(Z(a^j)) > \arg_{(t, t+1)}(Z(b^{j+1})) = \phi(b^{j+1})$$

$$(79) \quad j \geq N \Rightarrow \arg_{(t, t+1)}(Z(a^j)) \leq \arg_{(t, t+1)}(Z(b^{j+1})) = \phi(b^{j+1})$$

Therefore, using (10) and (72) we deduce that for  $j < N$  the inclusion  $a^j \in \sigma^{ss}$  implies  $\phi(a^j) > \phi(b^{j+1})$ ,  $\text{hom}(a^j, b^{j+1}) = 0$  which contradicts (29).

Now we will show that  $a^j \in \sigma^{ss}$  for  $N \leq j < u$  with the help of [8, Lemma 7.2]. Let  $N \leq j < u$  and  $a^j \notin \sigma^{ss}$ . One of the five cases given there must appear. In case (a) of [8, Lemma 7.2] we have  $a^k \in \sigma^{ss}$ ,  $\phi_-(a^j) = \phi(a^k) + 1$  for some  $k < j - 1$ , however then (76) and (10) imply  $Z(a^k) \in Z(\delta)_-^c$ , which contradicts (60). In case (b) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(a^k)$  for some  $k > j$ , however then (76) (57), and (10) imply  $\arg_{(t, t+1)}(Z(a^k)) < \arg_{(t, t+1)}(Z(a^j))$  with  $k > j$ , which contradicts (60). In case (c) of [8, Lemma 7.2] we have  $\phi_-(a^j) = \phi(b^k) + 1$  for some  $k < j$ , which would imply  $Z(b^k) \in Z(\delta)_-^c$ , which contradicts (59). Case (d) of [8, Lemma 7.2] ensures  $\phi_-(a^j) = \phi(b^k)$  for some  $j < k$ , and then (76), (10) ensure  $\arg_{(t, t+1)}(Z(b^k)) = \phi(b^k) < \arg_{(t, t+1)}(Z(a^j))$ , which contradicts (79), (59) and  $j < k$ . In the last case (e) of Lemma [8, Lemma 7.2] holds  $\phi_-(a^j) = \phi(M) + 1$ , however (75) and (76) imply that  $\phi_-(a^j) < \phi(M) + 1$  (here we take into account  $j < u$ ), which is again contradiction and we proved that  $a^j \in \sigma^{ss}$  for  $N \leq j < u$ . And in (74) we also proved that  $a^u \in \sigma^{ss}$ .

Finally, we want to prove that  $M' \in \sigma^{ss}$ . We will use [8, Lemma 2.12] with either the triple  $(a^u, M, b^{u+1}[-1])$  or the triple  $(a^{u-1}, M, b^u[-1])$  and taking into account the first triangle in (33). From Corollary 39 we see that this triple is Ext-exceptional triple. From the second triangle in (33) it follows that  $\phi(a^u) - 1 = \phi(M) = \phi(b^u) - 1$  if  $\phi(M) = \phi(b^u) - 1$  and  $\phi(b^u) - 1 < \phi(a^u) - 1 < \phi(M)$  if  $\phi(M) > \phi(b^u) - 1$ . The latter case combined with (71) and (72) imply  $\phi(a^u) - 1 < \phi(M) < \phi(b^{u+1}[-1]) < \phi(a^u)$  and indeed the second system of inequalities in the conditions of [8, Lemma 2.12] holds, and then this lemma ensures that  $M' \in \sigma^{ss}$ . Note that, if  $N = u$ , then the already proved (70), (71) and  $V < \phi(M) + 1$  imply the latter case. Thus, it remains to consider the case  $\phi(a^u) - 1 = \phi(M) = \phi(b^u) - 1$  and  $N < u$ . From (60) and since  $a^{u-1}, a^u \in \sigma^{ss}$  (recall also (72)) we see that  $\phi(a^{u-1}) < \phi(a^u) = \phi(b^u) = \phi(M) + 1$ , hence recalling

also that  $\phi(M) < t < \phi(b^{u-1}) \leq \phi(a^{u-1})$  we derive  $\phi(a^{u-1}) - 1 < \phi(b^u[-1]) = \phi(M) < \phi(a^{u-1})$ , and now [8, Lemma 2.12] applied to the triple  $(a^{u-1}, M, b^u[-1])$  ensures  $M' \in \sigma^{ss}$ . Now Subsection 6.3 shows that  $\alpha^j \in C_{0,\sigma\sigma}(\mathcal{T})$  for  $N \leq j \leq u$ .

(b) Assume now  $\phi(b^{n+1}) = \phi(b^n) + 1$ . Let us denote  $\phi(b^n) = t$ , then  $\phi(b^{n+1}) = t + 1$ . Using that  $\{b^j\}_{j \in \mathbb{Z}} \subset \sigma^{ss}$ ,  $\text{hom}(b^n, b^m) \neq 0$  for  $n \leq m$  and  $\text{hom}^1(b^m, b^n) \neq 0$  for  $m > n + 1$  it follows that:

$$(80) \quad \phi(b^j) = t \text{ for } j \leq n \quad \phi(b^j) = t + 1 \text{ for } j \geq n + 1.$$

(b.1) If  $\phi(M) < \phi(b^n) - 1$  and  $a^j \in \sigma^{ss}$  for some  $j \in \mathbb{Z}$ , then (27) imply  $\phi(a^j) < \phi(b^n)$ , hence (80) imply that  $\phi(a^j) < \phi(b^k)$ ,  $\text{hom}(b^k, a^j)$  for small enough  $k$ , which contradicts (28).

If  $\phi(M) < \phi(b^n) - 1$  and  $M' \in \sigma^{ss}$ , then from  $\text{hom}^1(b^{n+1}, M') \neq 0$  in (27) it follows that  $\phi(b^{n+1}) - 1 \leq \phi(M')$  and using that  $\phi(b^n) = \phi(b^{n+1}) - 1$  we obtain  $\phi(M) < \phi(M') - 1$  and hence  $\text{hom}^1(M', M) = 0$ , which contradicts (32).

(b.2) In this case  $t - 1 < \phi(M)$  and from table (36) we have also  $\phi(M) < t$ . Recalling that  $\text{hom}(M, b^n) \neq 0$  (see (26)) we can use Corollary 37 to deduce that  $\beta^n = \langle M, b^n, a^n \rangle \in C_{0,\sigma\sigma}(\mathcal{T})$ . Since  $Z(a^n) = Z(b^n) - Z(M)$  and  $\text{hom}(b^n, a^n) \neq 0$  (28),  $\text{hom}^1(a^n, M) \neq 0$  (27) we have  $\phi(b^{n+1}) - 1 = \phi(b^n) < \phi(a^n) < \phi(M) + 1$ .

If  $a^j \in \sigma^{ss}$  for some  $j < n$ , then from (72) and (80), (10) we see that  $Z(b^{j+1})$ ,  $Z(a^j)$ ,  $Z(b^j)$  are non-zero and collinear and from Remark 41 we know that  $Z(M) = Z(b^j) - Z(a^j)$ , which contradicts  $t - 1 < \phi(M) < t = \phi(b^n)$ . So for  $j < n$  we have  $a^j \notin \sigma^{ss}$ . Recalling (73) and Subsection 6.3 we deduce that  $\alpha^j, \beta^j \notin C_{0,\sigma\sigma}(\mathcal{T})$  for  $j \neq n$ . If we prove that  $M' \in \sigma^{ss}$ , then using again Subsection 6.3 we would obtain  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^n, \beta^n\}$ . To prove that  $M' \in \sigma^{ss}$  we use [8, Lemma 2.12] with the Ext-exceptional triple  $(a^n, M, b^{n+1}[-1])$  and the first triangle in (33). We showed above that  $\phi(b^n) < \phi(a^n) < \phi(M) + 1$ , and recalling that  $t - 1 < \phi(M) < t = \phi(b^n) = \phi(b^{n+1}) - 1$  it follows  $\phi(a^n) - 1 < \phi(M) < \phi(a^n)$ ,  $\phi(a^n) - 1 < \phi(b^{n+1}[-1]) < \phi(a^n)$ . Thus, the second system of the inequalities in the presumptions in [8, Lemma 2.12] is satisfied and this lemma ensures that  $M' \in \sigma^{ss}$  due to the first triangle in (33) with  $q = n$ .

Finally, from  $\text{hom}^1(b^{n+1}, M') \neq 0$  in (27) and  $\text{hom}(M', a^n) \neq 0$  in (26) it follows that  $\phi(b^{n+1}) - 1 \leq \phi(M') \leq \phi(a^n)$  and now from the first triangle in (33) we get  $\phi(b^{n+1}) - 1 < \phi(M') < \phi(a^n) < \phi(M) + 1$ .

(b.3) Now we have  $\phi(M) = \phi(b^n) - 1$  and hence  $\phi(M) = t - 1$ . Now for  $q \leq n$  the second distinguished triangle in (33) ensures that  $a^q \in \sigma^{ss}$  and  $\phi(a^q) = t$ . In particular, we have  $\phi(a^n) + 1 = \phi(b^{n+1})$  and from the first triangle in (33) it follows that  $M' \in \sigma^{ss}$ ,  $\phi(M') = \phi(a^n)$ . Recalling (73) and using Subsection 6.3 we obtain  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \leq n\}$ .  $\square$

**Lemma 55.** *Let us assume that we are in the situation of (a.1) of Proposition 54. In this case  $M' \in \sigma^{ss} \iff \phi(M) = t - 1$ .*

*Proof.* First, we note that for  $\phi(M) < t - 1$  holds  $M' \notin \sigma^{ss}$ . Indeed, if  $M' \in \sigma^{ss}$ , then from (27) and (59) it follows that  $\phi(M') \geq t$  and  $\text{hom}^1(M', M) = 0$ , which contradicts (32).

So, it remains to consider the case  $\phi(M) = t - 1$ . From the second triangle in (33) we have a distinguished triangle 
$$b^p \xrightarrow{\quad} a^p$$
  $\swarrow \quad \searrow$   $M[1]$  for each  $p \in \mathbb{Z}$ , and since  $\phi(b^p) > \phi(M[1]) = t$ , it follows

that this triangle is the HN filtration of  $a^p$ , hence  $a^p \in \mathcal{P}[t, \phi(b^p)]$  for each  $p \in \mathbb{Z}$ . From the first triangle in (33) we see that  $M'$  is in the extension closure of  $b^{p+1}[-1]$  and  $a^p$ , and taking into account that  $\phi(b^{p+1}[-1]) < t$ , we deduce that  $M' \in \mathcal{P}[\phi(b^{p+1}[-1]), \phi(b^p)]$  for each  $p \in \mathbb{Z}$ . Therefore

$\phi(b^{p+1}[-1]) \leq \phi_-(M')$  and  $\phi_+(M') \leq \phi(b^p)$  for each  $p \in \mathbb{Z}$ . Now using (59) and limiting  $p$  to  $+\infty$  we obtain  $t \leq \phi_-(M')$ , whereas limiting  $p$  to  $-\infty$  ensures  $\phi_+(M') \leq t$ , therefore  $\phi_-(M') = \phi_+(M') = t$  and indeed  $M' \in \sigma^{ss}$ .  $\square$

From tables (104), (106), (103), (105), (107) and subsection 6.2 follow the following two corollaries:

**Corollary 56.** *For  $\sigma \in \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  we have  $|C_{1,\sigma\sigma}(\mathcal{T})| \leq 1$ .*

**Corollary 57.** *The set  $\{\sigma \in \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st} : |C_{0,\sigma\sigma}(\mathcal{T})| = \infty\}$  is a disjoint union (any two summands are disjoint)*

$$\begin{aligned} & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) = \phi(b^p) + 1, \phi(b^p) = \phi(M) + 1\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M, a^p, a^{p+1}) : \phi(a^{p+1}) = \phi(a^p) + 1, \phi(a^p) = \phi(M') + 1\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (b^p, b^{p+1}, M') : \phi(b^{p+1}) = \phi(b^p) + 1 = \phi(M')\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (a^p, a^{p+1}, M) : \phi(a^{p+1}) = \phi(a^p) + 1 = \phi(M)\}. \end{aligned}$$

*Proof.* In the tables we see that  $|C_{0,\sigma\sigma}(\mathcal{T})| = \infty$  if and only iff  $\sigma$  is in the shown union. The fact that the union is disjoint follows from the behavior of  $C_{0,\sigma\sigma}(\mathcal{T})$ . In the tables we see that when  $\sigma$  varies in any subset, which is a summand of the given union, the set  $C_{0,\sigma\sigma}(\mathcal{T})$  does not change and that for any two different summands the corresponding sets of stable genus zero non-commutative curves are different (recall also that  $\mathfrak{T}_a^{st} \cap \mathfrak{T}_b^{st} = \emptyset$ , see (38)).  $\square$

**6.5. Study of  $C_{1,\sigma}(\mathcal{T})$ ,  $C_{1,\sigma\sigma}(\mathcal{T})$  as  $\sigma \in (-, M, -) \cup (-, M', -)$  and results for the genus zero case.** Here Remark 43 shows that the auto-equivalence  $\zeta$  transforms  $(-, M, -)$  to  $(-, M', -)$  and more precisely  $\sigma \in (a^p, M, b^{p+1})$  is mapped to  $\zeta \cdot \sigma \in (b^p, M', a^p)$  as  $\phi_{\zeta \cdot \sigma}(b^p) = \phi_\sigma(a^p)$ ,  $\phi_{\zeta \cdot \sigma}(M') = \phi_\sigma(M)$ ,  $\phi_{\zeta \cdot \sigma}(a^p) = \phi_\sigma(b^{p+1})$ . Therefore it is enough to study the behavior of  $C_{l,\sigma}(\mathcal{T})$ ,  $C_{l,\sigma\sigma}(\mathcal{T})$  in  $(a^p, M, b^{p+1})$ . To that end, we need first to strengthen [8, lemma 7.5 (e)]. So, we prove first:

**Lemma 58.** *Let  $\sigma \in (a^p, M, b^{p+1})$ , and let the following inequalities hold:*

$$(81) \quad \phi(a^p) - 1 < \phi(M) < \phi(a^p)$$

$$(82) \quad \phi(a^p) - 1 < \phi(b^{p+1}) - 1 < \phi(a^p).$$

*Then  $M' \in \sigma^{ss}$  and  $\phi(b^{p+1}) - 1 < \phi(M') = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(b^{p+1})) < \phi(a^p)$ .*

*Furthermore, if  $\phi(M) = \phi(M')$ , then  $\sigma \in (a^j, M, b^{j+1}) \cap (b^j, M', a^j)$  for all  $j \in \mathbb{Z}$ , and therefore  $C_{D^b(pt), \sigma\sigma}(\mathcal{T}) = C_{D^b(pt), \mathbb{K}}(\mathcal{T})$ . In particular  $\bigcap_{j \in \mathbb{Z}} (a^j, M, b^{j+1}) \cap (b^j, M', a^j) \neq \emptyset$ .*

*Proof.* From [8, Lemma 7.5 (a), (c)] we obtain  $b^p, M' \in \sigma^{ss}$ , and  $\phi(M) < \phi(b^p) < \phi(a^p)$ ,  $\phi(b^{p+1}) - 1 < \phi(M') = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(b^{p+1})) < \phi(a^p)$ .

Assume that  $\phi(M) = \phi(M')$  holds.

In this case we have  $\phi(a^p) - 1 < \phi(M) = \phi(M') < \phi(a^p)$ . Recalling that  $Z(\delta) = Z(M') + Z(M)$ , we see that  $t = \phi(M) = \phi(M')$  satisfies  $Z(\delta) = |Z(\delta)| \exp(i\pi t)$  and  $t < \phi(a^p) < t + 1$ . From the already obtained  $\phi(M) < \phi(b^p) < \phi(a^p)$  we get  $t < \phi(b^p) < \phi(a^p) < t + 1$ . Now we can apply [8, Corollary 3.20], which, besides  $\{Z(a^j), Z(b^j)\}_{j \in \mathbb{Z}} \subset Z(\delta)_+^c$ , gives us the formulas

$$(83) \quad \forall j \in \mathbb{Z} \quad \arg_{(t, t+1)}(Z(x^j)) < \arg_{(t, t+1)}(Z(x^{j+1}))$$

$$(84) \quad \lim_{j \rightarrow -\infty} \arg_{(t, t+1)}(Z(x^j)) = t; \quad \lim_{j \rightarrow +\infty} \arg_{(t, t+1)}(Z(x^j)) = t + 1,$$

where  $\{x^i\}_{i \in \mathbb{Z}}$  is either the sequence  $\{a^i\}_{i \in \mathbb{Z}}$  or the sequence  $\{b^i\}_{i \in \mathbb{Z}}$ . On the other hand, [8, Corollary 3.20] claims that for any three integers  $i, j, m$  we have:

$$(85) \quad j < m \leq i \Rightarrow \arg_{(t,t+1)}(Z(a^j)) < \arg_{(t,t+1)}(Z(b^m)) < \arg_{(t,t+1)}(Z(a^i)).$$

We extend the inequality  $t < \phi(b^p) < \phi(a^p) < t + 1$  to (86) as follows. We already have that  $a^p, b^p, b^{p+1} \in \sigma^{ss}$ . In (82) is given that  $\phi(a^p) < \phi(b^{p+1})$ . From  $\text{hom}^1(b^{p+1}, M') \neq 0$  (see (27)) it follows  $\phi(b^{p+1}) \leq t + 1$ , and from  $Z(b^{p+1}) \in Z(\delta)_+^c$  we get  $\phi(b^{p+1}) < t + 1 = \phi(M) + 1$ . We have also  $\phi(M) < \phi(b^{p+1})$  (due to  $\sigma \in (a^p, M, b^{p+1})$  and (37)). Therefore  $\phi(b^{p+1}) - 1 < \phi(M) < \phi(b^{p+1})$ , and from [8, Lemma 7.4 (a)] we get  $a^{p+1} \in \sigma^{ss}$  and  $\phi(b^{p+1}) - 1 < \phi(a^{p+1}) - 1 < \phi(M)$ . Thus, we derive:

$$(86) \quad \phi(a^p) - 1 < \phi(b^{p+1}) - 1 < \phi(a^{p+1}) - 1 < t < \phi(b^p) < \phi(a^p) < \phi(b^{p+1}) < \phi(a^{p+1}) < t + 1.$$

In [8, Lemma 7.5 (e)] it is shown that  $\sigma \in (a^j, M, b^{j+1})$  for each  $j \leq p$ , and therefore  $a^j, b^{j+1} \in \sigma^{ss}$  for each  $j \leq p$ .

Let  $j \geq p + 1$ . We will work now to prove that  $a^j, b^j \in \sigma^{ss}$ . From [8, Remark 3.9] we see that, if  $x^j$  is  $a^j$  or  $b^j$ , then  $x^j[-1]$  is in the extension closure of  $x^p, x^{p+1}[-1]$ , therefore we have:

$$(87) \quad j \geq p + 1 \quad x^j \in \mathcal{P}[\phi(x^{p+1}), \phi(x^p) + 1] \Rightarrow \phi_{\pm}(x^j) \in [\phi(x^{p+1}), \phi(x^p) + 1].$$

Suppose that  $a^j \notin \sigma^{ss}$  for some  $j \geq p + 1$ . Due to (86) and (87) we have  $a^j \in \mathcal{P}[\phi(a^{p+1}), \phi(a^p) + 1] \subset \mathcal{P}[\phi(b^{p+1}), \phi(b^{p+1}) + 1]$ , and [8, Remark 2.1(c)] shows that  $\arg_{(\phi(b^{p+1}), \phi(b^{p+1})+1)}(Z(a^j)) = \arg_{(t,t+1)}(Z(a^j))$ . Combining with [8, Remark 2.1 (a)], we put together:

$$(88) \quad t < \phi(a^{p+1}) \leq \phi_{-}(a^j) < \arg_{(t,t+1)}(Z(a^j)) < \phi_{+}(a^j) \leq \phi(a^p) + 1 < \phi(a^{p+1}) + 1.$$

We use [8, Lemma 7.2] to get a contradiction. One of the five cases given there must appear.

In case (a) of [8, Lemma 7.2] we have  $a^k \in \sigma^{ss}$ ,  $\phi_{-}(a^j) = \phi(a^k) + 1$  for some  $k < j - 1$ , and since we already have that  $a^q \in \sigma^{ss}$  and  $\phi(a^{q-1}) < \phi(a^q)$  for  $q \leq p$ , it follows that  $\phi(a^q) < \phi(a^k)$  for small enough  $q$ , hence  $\text{hom}^1(a^j, a^q) = 0$  for small enough  $q$ , which contradicts (28).

Case (b) of [8, Lemma 7.2] ensures  $\phi_{-}(a^j) = \phi(a^k)$  for some  $k > j$ . Using (88) and [8, Remark 2.1 (c)] we get  $\phi(a^k) = \arg_{(t,t+1)}(Z(a^k))$ , hence by (88) we get  $\arg_{(t,t+1)}(Z(a^k)) < \arg_{(t,t+1)}(Z(a^j))$ , which contradicts (83).

In case (c) of [8, Lemma 7.2] we have  $\phi_{-}(a^j) = \phi(b^k) + 1$  for some  $k < j$ , and since we already have that  $b^q \in \sigma^{ss}$  and  $\phi(b^{q-1}) < \phi(b^q)$  for  $q \leq p$ , it follows that  $\phi(b^q) < \phi(b^k)$  for small enough  $q$ , hence  $\text{hom}^1(a^j, b^q) = 0$  for small enough  $q$ , which contradicts (29).

Case (d) of [8, Lemma 7.2] ensures  $\phi_{-}(a^j) = \phi(b^k)$  for some  $j < k$ , and then (88), [8, Remark 2.1 (c)] ensure  $\arg_{(t,t+1)}(Z(b^k)) = \phi(b^k)$ , hence by (88) we get  $\arg_{(t,t+1)}(Z(b^k)) < \arg_{(t,t+1)}(Z(a^j))$ , which contradicts (85) and  $j < k$ .

In case (e) of [8, Lemma 7.2] we have  $\phi_{-}(a^j) = \phi(M) + 1 = t + 1$ , and (88), (86) imply  $t + 1 < \arg_{(t,t+1)}(Z(a^j)) < t + 2$ , which contradicts the incidence  $Z(a^j) \in Z(\delta)_+^c$ .

So far, we proved that  $a^j \in \sigma^{ss}$  for all  $j \in \mathbb{Z}$ .

Suppose that  $b^j \notin \sigma^{ss}$  for some  $j \geq p + 1$ . Due to (86) and (87) we have  $b^j \in \mathcal{P}[\phi(b^{p+1}), \phi(b^p) + 1] \subset \mathcal{P}[\phi(a^p), \phi(a^p) + 1]$ , and [8, Remark 2.1(c)] shows that  $\arg_{(\phi(a^p), \phi(a^p)+1)}(Z(b^j)) = \arg_{(t,t+1)}(Z(b^j))$ . Combining with [8, Remark 2.1 (a)] we put together:

$$(89) \quad t < \phi(b^{p+1}) \leq \phi_{-}(b^j) < \arg_{(t,t+1)}(Z(b^j)) < \phi_{+}(b^j) \leq \phi(b^p) + 1 < \phi(b^{p+1}) + 1.$$

We use [8, Lemma 7.3] to obtain a contradiction. Some of the five cases given there must appear.

Case (a) in [8, Lemma 7.3] ensures  $\phi_-(b^j) = \phi(a^k) + 1$  for some  $k < j - 1$ , and since we have already proved that  $a^q \in \sigma^{ss}$  and  $\phi(a^{q-1}) < \phi(a^q)$  for  $q \in \mathbb{Z}$ , it follows that (89) implies that  $\text{hom}^1(b^j, a^q) = 0$  for small enough  $q$ , which contradicts (28).

Case (b) in [8, Lemma 7.3] ensures  $\phi_-(b^j) = \phi(a^k)$  for some  $k \geq j$ , and then (89) and  $Z(a^k) \in Z(\delta)_+^c$  imply  $\arg_{(t,t+1)}(Z(a^k)) = \phi(a^k)$ , hence by (89) we get  $\arg_{(t,t+1)}(Z(a^k)) < \arg_{(t,t+1)}(Z(b^j))$ , which contradicts (85) and  $k \geq j$ .

Case (c) in [8, Lemma 7.3] ensures  $\phi_-(b^j) = \phi(b^k) + 1$  for some  $k < j - 1$ , and since we already have that  $b^q \in \sigma^{ss}$  and  $\phi(b^{q-1}) < \phi(b^q)$  for  $q \leq p$ , it follows that  $\text{hom}^1(b^j, b^q) = 0$  for small enough  $q$ , which contradicts (31).

In case (d) in [8, Lemma 7.3] we have  $\phi_-(b^j) = \phi(b^k)$  for some  $k > j$ . By  $Z(b^k) \in Z(\delta)_+^c$  and (89) it follows that  $\phi(b^k) = \arg_{(t,t+1)}(Z(b^k))$ , and then (89) gives  $\arg_{(t,t+1)}(Z(b^k)) < \arg_{(t,t+1)}(Z(b^j))$ , which contradicts (83) and  $k > j$ .

In case (e) using (89) we obtain  $\phi_-(b^j) = \phi(M') + 1 = t + 1 < \arg_{(t,t+1)}(Z(b^j)) < \phi_+(b^j) < t + 2$ , which contradicts the incidence  $Z(b^j) \in Z(\delta)_+^c$ .

So, we proved that  $b^j \in \sigma^{ss}$  for all  $j \in \mathbb{Z}$ .

We claim that  $\phi(M') = \phi(M) < \phi(a^j) < \phi(b^{j+1}) < \phi(a^{j+1}) < \phi(M) + 1$  for each  $j \in \mathbb{Z}$ , and then from table (37) follows that  $\sigma \in (a^j, M, b^{j+1}) \cap (b^j, M', a^j)$  for all  $j \in \mathbb{Z}$ . The desired inequality for  $j = p$  is in (86). In [8, p. 869] is shown that  $\phi(a^j) = \arg_{(t,t+1)}(Z(a^j))$  and  $\phi(b^j) = \arg_{(t,t+1)}(Z(b^j))$  for each  $j < p$ , if we show that this holds for  $j \geq p + 1$ , then the desired inequalities follow from (85). Let  $j > p$ . Since  $a^j, b^j \in \sigma^{ss}$  for each  $j$ , then by (30), (31) we get  $\phi(a^{p+1}) \leq \phi(a^j) \leq \phi(a^{p+1}) + 1$  and  $\phi(b^{p+1}) \leq \phi(b^j) \leq \phi(b^{p+1}) + 1$ , which combined with  $t < \phi(b^{p+1}) < \phi(a^{p+1}) < t + 1$  and  $Z(a^j), Z(b^j) \in Z(\delta)_+^c$  implies that  $\phi(a^j), \phi(b^j) \in (t, t + 1)$ , in particular  $\phi(a^j) = \arg_{(t,t+1)}(Z(a^j))$  and  $\phi(b^j) = \arg_{(t,t+1)}(Z(b^j))$  for each  $j \geq p + 1$ .  $\square$

**Lemma 59.** *If  $\sigma \in \cap_{j \in \mathbb{Z}}(a^j, M, b^{j+1}) \cap (b^j, M', a^j)$ , then  $\phi(M) = \phi(M')$  and (81), (82) hold for each  $p \in \mathbb{Z}$ . In particular for any  $p \in \mathbb{Z}$  holds*

$$\cap_{j \in \mathbb{Z}}(a^j, M, b^{j+1}) \cap (b^j, M', a^j) = \left\{ \sigma \in (a^p, M, b^{p+1}) : \begin{array}{l} \phi(a^p) - 1 < \phi(M) < \phi(a^p) \\ \phi(a^p) - 1 < \phi(b^{p+1}) - 1 < \phi(a^p) \\ \phi(M) = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(b^{p+1})) \end{array} \right\}$$

*Proof.* Let  $\sigma \in \cap_{j \in \mathbb{Z}}(a^j, M, b^{j+1}) \cap (b^j, M', a^j)$ . From table 37, and the non-vanishings  $\text{hom}(M, b^j) \neq 0$ ,  $\text{hom}^1(b^j, M') \neq 0$  in Corollary 39 it follows that for each  $j \in \mathbb{Z}$  we have

$$\phi(M') < \phi(a^j) < \phi(b^{j+1}) < \phi(a^{j+1}) < \phi(M') + 1 \quad \phi(M) < \phi(a^j) < \phi(b^{j+1}) < \phi(a^{j+1}) < \phi(M) + 1$$

From [8, Corollaries 3.18, Corollary 3.19] it follows that there exists  $t \in \mathbb{R}$  such that

$$(90) \quad \lim_{j \rightarrow +\infty} \phi(a^j) = t + 1, \quad \lim_{j \rightarrow -\infty} \phi(a^j) = t.$$

And therefore  $t = \phi(M) = \phi(M')$ . The last sentence now follows from Lemma 58.  $\square$

Now we pass to the behavior of  $C_{1,\sigma}(\mathcal{T})$ ,  $C_{1,\sigma\sigma}(\mathcal{T})$  in  $(a^p, M, b^{p+1})$ .

**Proposition 60.** *Let  $\sigma \in (a^p, M, b^{p+1})$  for some  $p \in \mathbb{Z}$ . Then  $a^p, M, b^{p+1} \in \sigma^{ss}$  and  $\phi(M) < \phi(b^{p+1})$ ,  $\phi(a^p) < \phi(b^{p+1})$ ,  $\phi(a^p) < \phi(M) + 1$ .*

<sup>1</sup>for some  $x \in \mathbb{Z}$



(a) If  $\phi(b^{p+1}) - 1 < \phi(M) < \phi(b^{p+1})$ , then  $a^{p+1} \in \sigma^{ss}$  and

$$(91) \quad \phi(a^{p+1}) = \arg_{(\phi(b^{p+1}), \phi(b^{p+1})+1]}(Z(b^{p+1}) - Z(M)).$$

(a.1) if  $\phi(a^p) < \phi(M)$ , then  $\sigma \in (a^p, a^{p+1}, M) \subset \mathfrak{T}_a^{st}$ .

(a.2) if  $\phi(a^p) \geq \phi(M)$ , then  $M' \in \sigma^{ss}$  and  $\phi(M') = \arg_{(\phi(b^{p+1})-1, \phi(b^{p+1}))}(Z(a^p) - Z(b^{p+1}))$  and:

$$(92) \quad \phi(b^{p+1}) - 1 < \phi(a^{p+1}) - 1 < \phi(M) \leq \phi(a^p) < \phi(b^{p+1}); \quad \phi(b^{p+1}) - 1 < \phi(M') < \phi(a^p)$$

(a.2.1) if  $\phi(M') < \phi(M)$ , then  $\sigma \in (a^j, a^{j+1}, M)$  for  $j \in \mathbb{Z}_{\leq x}^1$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \{A\}$ ,

(a.2.2) if  $\phi(M') > \phi(M)$ , then  $\sigma \in (b^j, b^{j+1}, M')$  for  $j \in \mathbb{Z}_{\leq x}^1$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \{B\}$ ,

(a.2.3) if  $\phi(M') = \phi(M)$ , then  $\sigma \in \bigcap_{j \in \mathbb{Z}} (a^j, M, b^{j+1}) \cap (b^j, M', a^j)$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \{A, B\} = C_1(\mathcal{T})$  and  $C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T})$ .

(b) If  $\phi(M) + 1 < \phi(b^{p+1})$  and  $\phi(M) < \phi(a^p)$ , then  $\sigma \in (M, b^p, b^{p+1})$  and we use table (105) with

$$(93) \quad \phi(b^p) = \arg_{(\phi(M), \phi(M)+1]}(Z(M) + Z(a^p)), \quad \phi(M) < \phi(b^p) < \phi(a^p) < \phi(M) + 1$$

(c) If  $\phi(M) + 1 < \phi(b^{p+1})$  and  $\phi(a^p) \leq \phi(M)$ , then  $C_{1,\sigma}(\mathcal{T}) = \emptyset$ ,  $M' \notin \sigma^{ss}$ , and if  $\sigma \in (A, B, C)$ , then  $(A, B, C) = (a^p, M, b^{p+1})$ . Furthermore  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^p\}$ , when  $\phi(a^p) = \phi(M)$ , whereas for  $\phi(a^p) < \phi(M)$ , we have  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$

(d) If  $\phi(M) + 1 = \phi(b^{p+1})$  and  $\phi(a^p) < \phi(M)$ , then  $C_{1,\sigma}(\mathcal{T}) = \emptyset$ ,  $M' \notin \sigma^{ss}$ , and if  $\sigma \in (A, B, C)$ , then  $(A, B, C) = (a^p, M, b^{p+1})$ . Furthermore  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{p+1}\}$ .

(e) If  $\phi(M) + 1 = \phi(b^{p+1})$  and  $\phi(a^p) > \phi(M)$ , then  $\sigma \in (b^j, b^{j+1}, M')$  for  $j \in \mathbb{Z}_{\leq x}^1$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \{B\}$ .

(f) If  $\phi(M) + 1 = \phi(b^{p+1}) = \phi(a^p) + 1$ , then  $a^j, b^j \in \sigma^{ss}$  for all  $j \in \mathbb{Z}$  and  $\phi(a^j) = \phi(b^j) = t + 1$  for each  $j \geq p + 1$ ,  $\phi(a^j) = \phi(b^j) = t$  for  $j \leq p$ . In particular  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = C_1(\mathcal{T})$ . In this case  $M' \in \sigma^{ss}$  and  $\phi(M) = \phi(M')$ , and in particular  $C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T})$ . In this case, if  $\sigma \in (A, B, C)$ , then  $(A, B, C) = (a^p, M, b^{p+1})$ .

*Proof.* Looking in table (37) we see that  $a^p, M, b^{p+1} \in \sigma^{ss}$  and the inequalities given there hold.

(a) [8, Lemma 7.4 (a)] ensures that  $a^{p+1} \in \sigma^{ss}$  and

$$(94) \quad \phi(b^{p+1}) < \phi(a^{p+1}) < \phi(M) + 1.$$

Recalling from Remark 41 that  $Z(a^{p+1}) = Z(b^{p+1}) - Z(M)$  it follows (91).

(a.1) Now  $\phi(a^p) < \phi(M)$  and [8, Lemma 7.4 (b)] ensure that  $\sigma \in (a^p, a^{p+1}, M)$  and hence  $\sigma \in \mathfrak{T}_a^{st}$ .

(a.2) Now combining (94) and (37) we see that hold the first series of inequalities in (92), and [8, Lemma 7.4 (c)] ensure that  $M' \in \sigma^{ss}$ ,  $\phi(M') = \arg_{(\phi(b^{p+1})-1, \phi(b^{p+1}))}(Z(a^p) - Z(b^{p+1}))$  and the second series of inequalities in (92).

(a.2.1) In the [8, proof of Lemma 7.4 (d)] is shown that  $\sigma \in (a^{j-1}, a^j, M)$  for small enough  $j$ . From [8, Lemma 5.2.] it follows that  $\sigma \in (a^{j-1}, a^j, M)$  and  $\sigma \in (a^{j-1}, a^j, M)$  and  $\phi(a^j) - 1 < \phi(a^{j-1}) < \phi(a^j)$  for all small enough  $j \in \mathbb{Z}$ . From Propositions (46), (51) we see that  $A$  is stable and  $B$  is not semistable.

(a.2.2) Now combining with (92) we obtain  $\phi(b^{p+1}) - 1 < \phi(a^{p+1}) - 1 < \phi(M) < \phi(M') < \phi(a^p) < \phi(b^{p+1})$  and therefore all the three conditions  $\phi(a^p) - 1 < \phi(M) < \phi(a^p)$ ,  $\phi(a^p) - 1 < \phi(b^{p+1}) - 1 < \phi(a^p)$ ,  $\phi(M) < \phi(M')$  required in [8, Lemma 7.5 (d)] are satisfied, which ensures that  $\sigma \in (b^j, b^{j+1}, M')$  for some  $j \in \mathbb{Z}$ . Furthermore, since we have  $\phi(b^{p+1}) < \phi(M) + 1$ , then in the [8, proof of Lemma 7.5 (d)] is actually shown that  $\sigma \in (b^j, b^{j+1}, M')$  for all small enough  $j \in \mathbb{Z}$ .

From [8, Lemma 5.2.] it follows that  $\sigma \in (b^{j-1}, b^j, M')$  and  $\phi(b^j) - 1 < \phi(b^{j-1}) < \phi(b^j)$  for all small enough  $j \in \mathbb{Z}$ . From Propositions (46), (51) we see that  $B$  is stable and  $A$  is not semistable.

(a.2.3) In (37) we have  $\phi(a^p) < \phi(b^{p+1})$  and (92) together with  $\phi(M) = \phi(M')$  give rise to (81), (82). Now we apply Lemma 58 and (39).

(b) The given inequalities and (37) ensure that  $\phi(a^p) - 1 < \phi(M) < \phi(a^p)$  and [8, Lemma 7.5 (a), (b)] shows that  $b^p \in \sigma^{ss}$ ,  $\phi(M) < \phi(b^p) < \phi(a^p) < \phi(M+1)$ ,  $\sigma \in (M, b^p, b^{p+1})$ . Recalling that in Remark 41 we have  $Z(b^p) = Z(M) + Z(a^p)$  and (93) follows.

(c) and (d): From the given inequalities follows  $\phi(b^{p+1}) > \phi(a^p) + 1$  and Lemma 44 we see that both  $A, B$  are not semistable. Furthermore, using [8, Lemmas 7.6, 7.7] it follows that  $\sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$ . From Lemma 44 (c) it follows that  $b^j, a^j \notin \sigma^{ss}$  for  $j \notin \{p, p+1\}$  and hence  $\sigma \notin (a^j, M, b^{j+1})$  for  $j \neq p$ . On the other hand the first triangle in (33) for  $q = p$  is a non-trivial HN filtration of  $M'$  and therefore  $M' \notin \sigma^{ss}$ , hence  $\sigma \notin (-, M', -)$ . From (38) we deduce that  $\sigma \notin \text{Stab}(\mathcal{T}) \setminus (a^p, M, b^{p+1})$ . From Subsection 6.3 it follows that  $\alpha^m \notin C_{0, \sigma\sigma}(\mathcal{T})$  for  $m \in \mathbb{Z}$ . On the other hand, since  $b^j, a^j \notin \sigma^{ss}$  for  $j \notin \{p, p+1\}$ , it follows that  $\beta^p \in C_{0, \sigma\sigma}(\mathcal{T})$  iff  $b^p \in \sigma^{ss}$  and  $\beta^{p+1} \in C_{0, \sigma\sigma}(\mathcal{T})$  iff  $a^{p+1} \in \sigma^{ss}$ . In case (d) we have  $\phi(M) = \phi(b^{p+1}) - 1$  and we can use Lemma 37 and Corollary 39 to get  $a^{p+1} \in \sigma^{ss}$ ,  $\beta^{p+1} \in C_{0, \sigma\sigma}(\mathcal{T})$ . In case (d) we have also  $\phi(a^p) < \phi(M)$ , and due to the non-vanishings  $\text{hom}(M, b^p) \neq 0$ ,  $\text{hom}(b^p, a^p) \neq 0$  in (26), (28) we see that  $b^p \notin \sigma^{ss}$ , therefore  $\beta^p \notin C_{0, \sigma\sigma}(\mathcal{T})$ . In case (c),  $a^{p+1} \notin \sigma^{ss}$ , since we have  $\text{hom}^1(a^{p+1}, M) \neq 0$ ,  $\text{hom}(b^{p+1}, a^{p+1}) \neq 0$  in (27), (28). In case (c) with  $\phi(a^p) < \phi(M)$  we have also  $b^p \notin \sigma^{ss}$  by the arguments used in case (d). Finally, in case (c) with  $\phi(a^p) = \phi(M)$  we use Corollary 37 and (27) to deduce that  $\beta^p \in C_{0, \sigma\sigma}(\mathcal{T})$ .

(e) The given inequalities and (37) give rise to  $\phi(a^p) < \phi(M) + 1 = \phi(b^{p+1}) < \phi(a^p) + 1$ . Therefore  $\phi(a^p) - 1 < \phi(M) = \phi(b^{p+1}) - 1 < \phi(a^p)$  and from [8, Lemma 7.5 (c)] we see that  $M' \in \sigma^{ss}$  and  $\phi(b^{p+1}) - 1 < \phi(M') = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(b^{p+1})) < \phi(a^p)$ . On the other hand  $\phi(M) = \phi(b^{p+1}) - 1$  implies that  $\phi(M') = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(b^{p+1})) > \phi(M)$ , hence we apply [8, Lemma 7.5 (d)]. Furthermore, we are given now that  $\phi(b^{p+1}) = \phi(M) + 1$ . Looking in the proof of [8, Lemma 7.5 (d)] we see that  $\sigma \in (b^j, b^{j+1}, M')$  for small enough  $j \in \mathbb{Z}$  and now the same arguments as in the proof of (a.2.2) imply that  $C_{1, \sigma\sigma}(\mathcal{T}) = C_{1, \sigma}(\mathcal{T}) = \{B\}$ .

(f) Let us denote  $t = \phi(M)$ . In this case we have  $t = \phi(M) = \phi(a^p) = \phi(b^{p+1}) - 1$  and by the triangles in (33) it follows that  $M', b^p, a^{p+1} \in \sigma^{ss}$  and  $\phi(b^p) = \phi(M) = \phi(M') = \phi(a^p) = t$ ,  $\phi(a^{p+1}) = t + 1$ . Using induction and (33) one sees that  $a^j, b^j \in \sigma^{ss}$ ,  $\phi(a^j) = \phi(b^j) = t + 1$  for each  $j \geq p + 1$ , and  $a^j, b^j \in \sigma^{ss}$ ,  $\phi(a^j) = \phi(b^j) = t$  for  $j \leq p$ . Due to the last sentence and table (37) it follows that  $\sigma \notin (-, M', -)$  and  $\sigma \notin (a^j, M, b^{j+1})$  for  $j \neq p$ . We showed that  $C_{1, \sigma\sigma}(\mathcal{T}) = \{A, B\}$  and from Corollary 56 it follows that  $\sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$ . From (38) we deduce that  $\sigma \notin \text{Stab}(\mathcal{T}) \setminus (a^p, M, b^{p+1})$ .  $\square$

**Corollary 61.** *For  $\sigma \in (-, M, -) \cup (-, M', -)$  we have  $C_{1, \sigma\sigma}(\mathcal{T}) = \{A, B\}$  iff  $M, M' \in \sigma^{ss}$  and  $\phi(M) = \phi(M')$ .*

*Proof.* Let  $\sigma \in (-, M, -)$ . Then  $\sigma \in (a^p, M, b^{p+1})$  for some  $p \in \mathbb{Z}$ . This  $\sigma$  must satisfy exactly one of the cases (a.1), (a.2.1), (a.2.2), (a.2.3), (b), (c), (d), (e), (f) in Proposition 60. Taking into account tables (104), (106), (105), (103) we see that  $C_{1, \sigma\sigma}(\mathcal{T}) = \{A, B\}$  if and only if  $\sigma$  is in case (a.2.3) or in case (f). On the other hand using Corollary 45 we see that  $\sigma$  satisfies case (a.2.3) or case (f) if and only if  $M, M' \in \sigma^{ss}$  and  $\phi(M) = \phi(M')$ . If  $\sigma \in (-, M', -)$  we use the already proved case and Remark 43.  $\square$

Now we can give also proof of Proposition 68. *Proof of Proposition 68* If  $\sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  due to (38) we have either  $\sigma \in (a^p, M, b^{p+1})$  or  $\sigma \in (b^p, M', a^p)$  for some  $p \in \mathbb{Z}$ . If  $\sigma \in (a^p, M, b^{p+1})$  we go through all possible cases (a.1), (a.2.1), (a.2.2), (a.2.3), (b), (c), (d), (e), (f) in Proposition 60 and from  $\sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  we must be in some of the cases (a.2.3), (c), (d), (f).

In cases (a.2.3) and (f) we have  $\{M, M'\} \subset \sigma^{ss}$ ,  $\phi(M) = \phi(M')$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = \{A, B\}$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T})$ . Furthermore using also Lemma 59 we see that  $\sigma$  satisfies case (a.2.3) if and only if  $\sigma \in \bigcap_{p \in \mathbb{Z}} (a^p, M, b^{p+1}) \cap (b^p, M', a^p)$ .

In cases (c), (d) we have  $\{M, M'\} \not\subset \sigma^{ss}$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \emptyset$ .

The case  $\sigma \in (b^p, M', a^p)$  follows from the already proved case and Remark 43.

The fact that the unions are disjoint follows from the properties  $\sigma \notin \text{Stab}(\mathcal{T}) \setminus (a^p, M, b^{p+1})$  obtained in (c), (d), (f) of Proposition 60.

The behavior of  $C_{0,\sigma\sigma}(\mathcal{T})$  specified in the first three sentences after (109) is the same as explained in (c), (d) of Proposition 60.

If  $\sigma \in (b^p, M', a^p)$  we use (43) and Remark 43.  $\square$

## 7. THE SET OF STABILITIES, WHERE $|C_{0,\sigma\sigma}(\mathcal{T})| < \infty$ IS OPEN AND DENSE IN $\text{Stab}(\mathcal{T})$

Note first that due to (41) we have that  $|C_{0,\sigma\sigma}(\mathcal{T})| = \infty$  implies  $C_{1,\sigma}(\mathcal{T}) = \{A, B\}$ . On the other hand from the already proved tables (106), (105), (104), (103), (107), Proposition 68, and (38) we see that  $C_{1,\sigma}(\mathcal{T}) = \{A, B\}$  implies  $|C_{0,\sigma\sigma}(\mathcal{T})| = \infty$ . Therefore, we see that

$$(95) \quad C_{1,\sigma}(\mathcal{T}) = \{A, B\} \iff |C_{0,\sigma\sigma}(\mathcal{T})| = \infty.$$

Note that from the results in previous sections follows Proposition 10 (see the proof in the end of Subsection A), i. e.  $C_{1,\sigma\sigma}(\mathcal{T}) = \{A, B\} \iff M, M' \in \sigma^{ss}$  and  $\phi(M) = \phi(M')$ . From (95) tables (104), (106), (103), (105), (107), Proposition 68, (38) and Subsection 6.2 it follows that the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| = \infty\}$  is union of the following three subsets in (96), (97), (98)

$$(96) \quad \{\sigma \in \text{Stab}(\mathcal{T}) : C_{1,\sigma\sigma}(\mathcal{T}) = \{A, B\}\}$$

$$(97) \quad \{\sigma \in \text{Stab}(\mathcal{T}) : B \in C_{1,\sigma\sigma}(\mathcal{T}) \text{ and } M, M' \in \sigma^{ss} \text{ and } \phi(M) + 1 = \phi(M')\} \setminus$$

$$\left( \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) < \phi(b^p) + 1\} \cup \{\sigma \in (b^p, b^{p+1}, M') : \phi(b^{p+1}) < \phi(b^p) + 1\} \right)$$

$$(98) \quad \{\sigma \in \text{Stab}(\mathcal{T}) : A \in C_{1,\sigma\sigma}(\mathcal{T}) \text{ and } M, M' \in \sigma^{ss} \text{ and } \phi(M') + 1 = \phi(M)\} \setminus$$

$$\left( \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M', a^p, a^{p+1}) : \phi(a^{p+1}) < \phi(a^p) + 1\} \cup \{\sigma \in (a^p, a^{p+1}, M) : \phi(a^{p+1}) < \phi(a^p) + 1\} \right)$$

From Corollary 27 we know that  $\{\sigma \in \text{Stab}(\mathcal{T}) : B \in C_{1,\sigma\sigma}(\mathcal{T})\}$  is a closed subset in  $\text{Stab}(\mathcal{T})$ . From [2, p. 342] is known that for any object  $X$  in  $\mathcal{T}$  the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : X \in \sigma^{ss}\}$  is closed subset and the function assigning  $\phi_\sigma(X)$  to any  $\sigma$  in this subset, is continuous in this subset. Therefore

$\{\sigma \in \text{Stab}(\mathcal{T}) : B \in C_{1,\sigma\sigma}(\mathcal{T}) \text{ and } M, M' \in \sigma^{ss} \text{ and } \phi(M) + 1 = \phi(M')\}$  is a closed subset in  $\text{Stab}(\mathcal{T})$ .

**Remark 62.** From Remark 31 (b) and table (36) we know that the map (20) gives a homeomorphism from  $(M, b^p, b^{p+1}) \subset \text{Stab}(\mathcal{T})$  to  $\mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_0 - y_1 < 0 \\ y_0 - y_2 < -1 \\ y_1 - y_2 < 0 \end{array} \right\}$ . It follows that the sub-

set  $\{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) < \phi(b^p) + 1\}$  is the inverse image of the open subset  $\mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_0 - y_1 < 0 \\ y_0 - y_2 < -1 \\ -1 < y_1 - y_2 < 0 \end{array} \right\}$  with respect to this homeomorphism, therefore  $\{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) < \phi(b^p) + 1\}$  is an open subset in  $\text{Stab}(\mathcal{T})$ .

By similar arguments one shows that  $\{\sigma \in (b^p, b^{p+1}, M') : \phi(b^{p+1}) < \phi(b^p) + 1\}$  is also an open subset in  $\text{Stab}(\mathcal{T})$ . Therefore (97) is a closed subset in  $\text{Stab}(\mathcal{T})$ . Since (98) is obtained from (97) via the auto-equivalence  $\zeta$  in (42) it follows that (98) is closed subset as well. From Corollary 27 it follows that (96) is a closed subset as well (see also Proposition 10). We proved that (96), (97), (98) are closed subset, **therefore**  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| = \infty\}$  **is a closed subset in**  $\text{Stab}(\mathcal{T})$ , **hence**  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| < \infty\}$  **is an open subset in**  $\text{Stab}(\mathcal{T})$ .

**Corollary 63.** The set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| < \infty\}$  is dense subset in  $\text{Stab}(\mathcal{T})$ .

*Proof.* We already proved that the set in question is open subset, therefore it is enough to show that for any  $\sigma \in \text{Stab}(\mathcal{T})$  such that  $|C_{0,\sigma\sigma}(\mathcal{T})| = \infty$  and for any neighborhood  $U$  of  $\sigma$  there exists  $\sigma' \in U$  such that  $|C_{0,\sigma'\sigma'}(\mathcal{T})| < \infty$ . So let  $\sigma \in \text{Stab}(\mathcal{T})$  be such that  $|C_{0,\sigma\sigma}(\mathcal{T})| = \infty$  and let  $\sigma \in U$  for some open subset  $U$ . We use Corollary 57, Proposition 68, Lemma 59 and deduce that the set  $\{\sigma \in \text{Stab}(\mathcal{T}) : |C_{0,\sigma\sigma}(\mathcal{T})| = \infty\}$  is the following disjoint union (in the last summand  $q \in \mathbb{Z}$ ):

$$\begin{aligned} & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) = \phi(b^p) + 1, \phi(b^p) = \phi(M) + 1\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (M, a^p, a^{p+1}) : \phi(a^{p+1}) = \phi(a^p) + 1, \phi(a^p) = \phi(M') + 1\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (b^p, b^{p+1}, M') : \phi(b^{p+1}) = \phi(b^p) + 1 = \phi(M')\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (a^p, a^{p+1}, M) : \phi(a^{p+1}) = \phi(a^p) + 1 = \phi(M)\} \cup \\ & \bigcup_{p \in \mathbb{Z}} \{\sigma \in (a^p, M, b^{p+1}) : \phi(M) + 1 = \phi(b^{p+1}) = \phi(a^p) + 1\} \\ & \cup \bigcup_{p \in \mathbb{Z}} \{\sigma \in (b^p, M', a^p) : \phi(M') + 1 = \phi(a^p) = \phi(b^p) + 1\} \\ & \cup \left\{ \sigma \in (a^q, M, b^{q+1}) : \begin{array}{l} \phi(a^q) - 1 < \phi(M) < \phi(a^q) \\ \phi(a^q) - 1 < \phi(b^{q+1}) - 1 < \phi(a^q) \\ \phi(M) = \arg_{(\phi(a^q)-1, \phi(a^q))}(Z(a^q) - Z(b^{q+1})) \end{array} \right\}. \end{aligned}$$

Assume that  $\sigma \in (M, b^p, b^{p+1})$  and  $\phi(b^{p+1}) = \phi(b^p) + 1, \phi(b^p) = \phi(M) + 1$  (the arguments in the other cases are analogous).

The subset  $\{\sigma \in (M, b^p, b^{p+1}) : \phi(b^{p+1}) = \phi(b^p) + 1, \phi(b^p) = \phi(M) + 1\}$  is the inverse image of the subset  $\mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_1 = y_0 + 1 \\ y_2 = y_1 + 1 \end{array} \right\}$  in  $\mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_0 - y_1 < 0 \\ y_0 - y_2 < -1 \\ y_1 - y_2 < 0 \end{array} \right\}$  with respect to the homeomorphism

discussed in Remark 62. Denote this homeomorphism by  $f$ .<sup>5</sup> In particular,  $f(\sigma) = (r_0, r_1, r_2, y_0, y_0 + 1, y_0 + 2)$  for some  $r_i > 0$  and  $y_0 \in \mathbb{R}$ . Since  $(M, b^p, b^{p+1})$  is an open subset in  $\text{Stab}(\mathcal{T})$ , it follows that  $f(U \cap (M, b^p, b^{p+1}))$  is an open subset in  $\mathbb{R}^6$  containing  $f(\sigma)$ , therefore for some  $\varepsilon > 0$  we have open subset  $V \ni \sigma$ , such that  $V \subset U \cap (M, b^p, b^{p+1})$  and such that  $f(V) = (r_0 - \varepsilon, r_0 + \varepsilon) \times (r_1 - \varepsilon, r_1 + \varepsilon) \times (r_2 - \varepsilon, r_2 + \varepsilon) \times (y_0 - \varepsilon, y_0 + \varepsilon) \times (y_0 + 1 - \varepsilon, y_0 + 1 + \varepsilon) \times (y_0 + 2 - \varepsilon, y_0 + 2 + \varepsilon)$ . Hence  $f(v) \notin \mathbb{R}_{>0}^3 \times \left\{ \begin{array}{l} y_1 = y_0 + 1 \\ y_2 = y_1 + 1 \end{array} \right\}$  for many  $v \in V$ , hence by table (105) we have  $|C_{0,vv}(\mathcal{T})| < \infty$  for many  $v \in V$ .  $\square$

### 8. WALLS AND CHAMBERS FOR $C_{1,\sigma\sigma}$

In tables (103), (104), (105), (106) and (107) and Proposition 68, we have summed up the local behaviour of  $C_{1,\sigma\sigma}$ . However, to understand how many walls there are, how they attach to one another and how the chambers with given subset of  $C_{1,\sigma\sigma}$  are separated by them, we need to combine these results with knowing how two subset  $(A, B, C), (A', B', C') \subset \text{Stab}(\mathcal{T})$  intersect or equivalently how the corresponding charts given by the homeomorphism in Remark 31 glue. We begin by labeling some of the subsets appearing in the aforementioned tables and proposition.

**Definition 64.** *The following subsets of  $\text{Stab}(\mathcal{T})$  and their connected unions will be called walls*

$$\begin{aligned} \mathcal{W}(a^m, a^{m+1}, M) &= \{\sigma \in (a^m, a^{m+1}, M) : \phi(a^{m+1}) = \phi(a^m) + 1\}, \\ \mathcal{W}(b^m, b^{m+1}, M') &= \{\sigma \in (b^m, b^{m+1}, M') : \phi(b^{m+1}) = \phi(b^m) + 1\}, \\ \mathcal{W}(M', a^m, a^{m+1}) &= \{\sigma \in (M', a^m, a^{m+1}) : \phi(a^{m+1}) = \phi(a^m) + 1\}, \\ \mathcal{W}(M, b^m, b^{m+1}) &= \{\sigma \in (M, b^m, b^{m+1}) : \phi(b^{m+1}) = \phi(b^m) + 1\}, \\ \mathcal{W}^0(A, B) &= \bigcap_{p \in \mathbb{Z}} ((a^p, M, b^{p+1}) \cap (b^p, M', a^p)), \\ \mathcal{G}(a^m, M, b^{m+1}) &= \{\sigma \in (a^m, M, b^{m+1}) : \phi(a^m) + 1 = \phi(M) + 1 = \phi(b^{m+1})\}, \\ \mathcal{G}(b^m, M', a^m) &= \{\sigma \in (b^m, M', a^m) : \phi(b^m) + 1 = \phi(M') + 1 = \phi(a^m)\}, \end{aligned}$$

for all  $m \in \mathbb{Z}$ .

*If  $X$  is a submanifold and  $Y$  a submanifold with a boundary in  $\text{Stab}(\mathcal{T})$  both of real codimension 1 and disjoint. Let  $Z \subset X$  be a submanifold of real codimension 2 contained in the boundary  $\partial Y$ , then we say that  $X$  attaches to  $Y$  in  $Z$  if  $Z = \bar{X} \cap Y$ .*

Notice that except for the last 2 walls which are of real codimension 2 submanifolds the other walls are submanifolds of real codimension 1. For  $\mathcal{W}^0(A, B)$  this follows from Lemma 59. For our purposes the following definition of a chamber will be sufficient:

**Definition 65.** *A closed chamber with  $N \subset C_1(\mathcal{T})$  stable is a closure of a non-empty maximal open connected subset  $Ch^0$  of  $\text{Stab}(\mathcal{T})$  with  $C_{1,\sigma\sigma} = N$  for all  $\sigma \in Ch^0$ .*

We now state the main result summing up the global behaviour of  $C_{1,\sigma\sigma}$ .

<sup>5</sup> $f$  can be viewed as a chart in a  $C^\infty$  manifold.

**Proposition 66.** *There is a unique closed chamber  $Ch(A)$  with  $A$  stable and a unique closed chamber  $Ch(B)$  with  $B$  stable. The complement of  $Ch(A) \cup Ch(B)$  is an open subset of  $\text{Stab}(\mathcal{T})$  with  $C_{1,\sigma\sigma} = \emptyset$  whenever  $\sigma$  lies in it. It is given by the disjoint union of open subsets  $\mathcal{O}_m$  (see (102)) going over all  $m \in \mathbb{Z}$ . In particular, the closures  $\overline{\mathcal{O}_m}$  are the closed chambers with  $\emptyset$  stable.*

There is a unique connected set with  $|C_{1,\sigma\sigma}| = 2$  given by

$$Ch(A) \cap Ch(B) = \overline{\mathcal{W}^0(A, B)} = \mathcal{W}^0(A, B) \cup \bigcup_p (\mathcal{G}(a^p, M, b^{p+1}) \cup \mathcal{G}(b^p, M', a^p)).$$

The boundary of  $Ch(A)$  is the union of the walls:

$$(99) \quad \bigcup_p (\mathcal{W}(a^p, a^{p+1}, M) \cup \mathcal{W}(M', a^p, a^{p+1})) \cup \overline{\mathcal{W}^0(A, B)},$$

where  $\mathcal{W}(a^p, a^{p+1}, M) \cup \mathcal{W}(M', a^p, a^{p+1})$  is a connected sub-manifold of real codimension 1. The union of the sets  $\mathcal{W}(a^p, a^{p+1}, M) \cup \mathcal{W}(M', a^p, a^{p+1})$  is disjoint, and moreover it is disjoint from  $\overline{\mathcal{W}^0(A, B)}$ . However, the intersection  $\mathcal{W}(a^p, a^{p+1}, M) \cap \mathcal{W}(M', a^p, a^{p+1})$  attaches to  $\overline{\mathcal{W}^0(A, B)}$  in  $\mathcal{G}(a^m, M, b^{m+1})$  and  $\mathcal{G}(b^m, M', a^m)$ . Therefore, (99) is a connected real codimension 1 wall.

The boundary of  $Ch(B)$  is the union of the walls:

$$(100) \quad \bigcup_p (\mathcal{W}(b^p, b^{p+1}, M') \cup \mathcal{W}(M, b^p, b^{p+1})) \cup \overline{\mathcal{W}^0(A, B)},$$

where  $\mathcal{W}(b^m, b^{m+1}, M') \cup \mathcal{W}(M, b^m, b^{m+1})$  is a connected sub-manifold of real codimension 1. The union of the sets  $\mathcal{W}(b^m, b^{m+1}, M') \cup \mathcal{W}(M, b^m, b^{m+1})$  is disjoint, and moreover it is disjoint from  $\overline{\mathcal{W}^0(A, B)}$ . However, the intersection  $\mathcal{W}(b^m, b^{m+1}, M') \cap \mathcal{W}(M, b^m, b^{m+1})$  attaches to  $\overline{\mathcal{W}^0(A, B)}$  in  $\mathcal{G}(a^m, M, b^{m+1})$  and  $\mathcal{G}(b^m, M', a^m)$ . Therefore, (100) is a connected real codimension 1 wall.

*Proof.* Recall that  $\mathfrak{I}_a^{\text{st}} \cap \mathfrak{I}_b^{\text{st}} = \emptyset$  and there exists an auto-equivalence  $\xi \in \text{Aut}(\text{Stab}(\mathcal{T}))$  given by (42) which identifies the two disjoint sets. We will therefore first understand the picture in  $\mathfrak{I}_a^{\text{st}}$  and then glue  $(a^p, M, b^{p+1})$  and  $(b^p, M', a^p)$  (where the last two sets are again identified using the auto-equivalence).

First, let us describe the situation in  $\bigcup_p (a^p, a^{p+1}, M)$ . Using [8, Lemma 5.4], we can describe the intersection  $(a^p, a^{p+1}, M) \cap (a^j, a^{j+1}, M)$  for  $p > j$  as

$$\{\sigma \in (a^p, a^{p+1}, M) : \phi(a^{p+1}) < \phi(a^p) + 1\}.$$

By looking at Table (104), we see that this corresponds to the interior of the unique chamber with  $A$  stable in  $(a^p, a^{p+1}, M)$ . It implies that this set is contained in the interior of the chamber with  $A$  stable in  $(a^j, a^{j+1}, M)$ . The union

$$\bigcup_p \{\sigma \in (a^p, a^{p+1}, M) : \phi(a^{p+1}) < \phi(a^p) + 1\}.$$

is connected and therefore contained in a single chamber which we denote by  $Ch(A, 1)$ . Moreover, the walls  $\mathcal{W}(a^p, a^{p+1}, M)$  and  $\mathcal{W}(a^j, a^{j+1}, M)$  are disjoint whenever  $p \neq j$ .

We do the same for  $\bigcup_p(M', a^p, a^{p+1})$ . By [8, Lemma 5.6], we see that

$$(M', a^p, a^{p+1}) \cap (M', a^j, a^{j+1}) = \{\sigma \in (M', a^p, a^{p+1}) : \phi(a^{p+1}) < \phi(a^p) + 1\}$$

for  $p < j$ . By Table (106) this coincides with the interior of the unique chamber with  $A$  stable in  $(M', a^p, a^{p+1})$ . The union

$$\bigcup_p \{\sigma \in (M', a^p, a^{p+1}) : \phi(a^{p+1}) < \phi(a^p) + 1\}.$$

is connected and thus contained in a single chamber which we denote by  $\mathcal{Ch}(A, 2)$ . The walls  $\mathcal{W}(M', a^p, a^{p+1})$  and  $\mathcal{W}(M', a^j, a^{j+1})$  are disjoint whenever  $p \neq j$ .

Let us now glue the sets  $(-, -, M)$  and  $(M', -, -)$ . The intersection  $(a^m, a^{m+1}, M) \cap (M', -, -)$  is contained by [8, Lemma 6.3] in the following subset of  $(a^m, a^{m+1}, M)$ :

$$(101) \quad \{\sigma \in (a^m, a^{m+1}, M) : \phi(a^{m+1}) < \phi(a^m) + 1 \text{ or } \phi(M) < \phi(a^{m+1})\}.$$

From the proof of [8, Lemma 6.3], we see that if the condition  $\phi(a^{m+1}) < \phi(a^m) + 1$  is true, then there exists a  $j \gg m$  with  $\sigma \in (M', a^j, a^{j+1})$ . This shows that the interior of the unique chamber with  $A$  stable in  $(a^m, a^{m+1}, M)$  intersects with the interior of the chamber  $\mathcal{Ch}(A, 2)$  non-trivially and therefore  $\mathcal{Ch}(A, 1) = \mathcal{Ch}(A, 2) =: \mathcal{Ch}(A)$ . Notice that we have already considered all subsets of  $\text{Stab}(\mathcal{T})$  with  $C_{1,\sigma\sigma} = \{A\}$  as either walls of  $\mathcal{Ch}(A)$  or as contained in its interior. We can conclude that  $\mathcal{Ch}(A)$  is the unique closed chamber with  $A$  stable.

If in (101) the condition  $\phi(M) < \phi(a^{m+1})$  is true and  $\phi(a^{m+1}) = \phi(a^m) + 1$ , then by the proof of [8, Lemma 6.3],  $\sigma$  lies in  $(M', a^m, a^{m+1})$  and in particular in  $\mathcal{W}(M', a^m, a^{m+1})$ . Therefore, we see that

$$\mathcal{W}(a^m, a^{m+1}, M) \cap \mathcal{W}(M', a^m, a^{m+1}) = \{\sigma \in \mathcal{W}(a^m, a^{m+1}, M) : \phi(M) < \phi(a^{m+1})\},$$

which is an open subset of  $\mathcal{W}(a^m, a^{m+1}, M)$ . Moreover, as we see from (101) that  $\mathcal{W}(a^m, a^{m+1}, M) \cap (M', -, -) = \mathcal{W}(a^m, a^{m+1}, M) \cap \mathcal{W}(M', a^m, a^{m+1})$ , we conclude that

$$\mathcal{W}(a^m, a^{m+1}, M) \cap \mathcal{W}(M', a^j, a^{j+1}) = \emptyset,$$

whenever  $j \neq m$ .

Let us now attach  $(a^p, b^{p+1}, a^{p+1})$  which will help us understand chambers with  $\emptyset$  stable of which they are open subsets by Table (107). From [8, Lemma 6.1 and 6.2], we know that they are pairwise disjoint, and that

$$(a^m, b^{m+1}, a^{m+1}) \cap (a^l, a^{l+1}, M) = (a^m, b^{m+1}, a^{m+1}) \cap (M', a^l, a^{l+1}) = \emptyset,$$

whenever  $m \neq l$ , while if  $m = l$  the intersections are non-trivial. Looking at the proof of [8, Lemma 6.3] we see that if  $\sigma \in (a^m, a^{m+1}, M)$  and  $\phi(M) < \phi(a^{m+1})$ ,  $\phi(a^{m+1}) > \phi(a^m) + 1$ , then  $\sigma \in (a^m, b^{m+1}, a^{m+1})$ . Therefore, we can say that

$$(a^m, a^{m+1}, M) \cap (a^m, b^{m+1}, a^{m+1}) = \{\sigma \in (a^m, a^{m+1}, M) : \phi(M) < \phi(a^{m+1}), \phi(a^{m+1}) > \phi(a^m) + 1\}.$$

Additionally, as  $(a^m, a^{m+1}, M) \cap (M', a^m, a^{m+1})$  has to contain a neighborhood of  $\mathcal{W}(a^m, a^{m+1}, M) \cap \mathcal{W}(M', a^m, a^{m+1})$ , it must intersect with  $(a^m, a^{m+1}, M) \cap (a^m, b^{m+1}, a^{m+1})$  in a nonempty open set.

If a set of the form  $(X, Y, Z) \subset \text{Stab}(\mathcal{T})$  for  $(X, Y, Z)$  full exceptional triple has a unique closed chamber with  $\emptyset$  stable, we denote by  $(X, Y, Z)_\emptyset$  its interior. We can conclude from the above that

$$(a^m, a^{m+1}, M)_\emptyset \cup (M', a^m, a^{m+1})_\emptyset \cup (a^m, b^{m+1}, a^{m+1})_\emptyset$$

are connected open subsets of  $\text{Stab}(\mathcal{T})$  for all  $m \in \mathbb{Z}$  and they are pairwise disjoint.

We now need to consider what happens in  $(a^p, M, b^{p+1})$ . This is partly handled by Proposition 60. However, we do some extra work to obtain the qualitative picture given in Figure (4), where we use  $\phi(a^m) - \phi(M)$  and  $\phi(M) + 1 - \phi(b^{m+1})$  as our coordinates under the assumption that they are the only variables that are allowed to vary. The reader should note that this picture is meant as a helpful tool for the discussion below and is not meant to depict it precisely. From now on, when we talk about a sub-case without mentioning its proposition, we mean a sub-case of Proposition 60.

In case (a.2.1), the stability condition  $\sigma$  lies in the interior of  $\mathcal{Ch}(A)$  as it lies in the intersection of multiple  $(a^j, a^{j+1}, M')$  and it connects to the case (a.1) which lies in  $(a^p, a^{p+1}, M)$ : this can be seen, because if  $\phi(a^m) - \phi(M)$  decreases while keeping  $\phi(M) + 1 - \phi(b^{m+1})$  constant, then from the formula for  $\phi(M')$  in (a.2), we see that  $\phi(M) - \phi(M')$  increases. Therefore by doing so, we eventually land in (a.2.1) on our way to  $\phi(a^p) = \phi(M)$ . Now as (a.1) connects to both (a.2.1) and (d) with  $C_{1,\sigma\sigma} = \emptyset$ , it needs to contain an open subset of  $\mathcal{W}(a^p, a^{p+1}, M)$ . We show that  $\mathcal{W}(a^p, a^{p+1}, M) \cap (a^p, M, b^{p+1})$  is contained in  $\mathcal{W}(a^p, a^{p+1}, M) \cap \mathcal{W}(M', a^p, a^{p+1})$  and that it attaches to the closure of  $\mathcal{W}^0(A, B)$  in  $(a^p, M, b^{p+1})$  through  $\mathcal{G}(a^p, M, b^{p+1})$ .

From Proposition 60, we see that in (a.1)  $\sigma \in (a^p, a^{p+1}, M)$  and  $\phi(a^{p+1}) > \phi(b^{p+1}) > \phi(M)$ . Using again the proof of [8, Lemma 6.3], we see that if additionally  $\sigma \in \mathcal{W}(M', a^p, a^{p+1})$ , then  $\sigma \in \mathcal{W}(a^p, a^{p+1}, M) \cap \mathcal{W}(M', a^p, a^{p+1})$ . Let us now assume that we are in case (a.1) and that  $\phi(M) + 1 < \phi(b^{m+1}) + \epsilon_1$  and  $\phi(a^m) > \phi(M) + \epsilon_2$  for some small  $\epsilon_1, \epsilon_2 > 0$ . This corresponds to taking a small neighborhood of  $\mathcal{G}(a^p, a^{p+1}, M)$  and intersecting it with (a.1). It follows then using Table (36) that

$$\phi(a^m) + 1 - \epsilon_1 < \phi(M) + 1 - \epsilon_1 < \phi(b^{p+1}) < \phi(a^{p+1}) < \phi(M) + 1 < \phi(a^m) + 1 + \epsilon_2.$$

and we see that we can chose  $\phi(a^{m+1}) = \phi(a^m) + 1$  in this, to obtain the intersection with  $\mathcal{W}(M', a^p, a^{p+1})$ . Therefore, each neighborhood of  $\mathcal{G}(a^p, M, b^{p+1})$  contains a point of  $\mathcal{W}(M', a^p, a^{p+1}) \cap \mathcal{W}(a^p, a^{p+1}, M)$ .

Similar arguments show that (a.2.2) connects through (e) to (b) and that the open subset of  $\mathcal{W}(M, b^p, b^{p+1})$  attaches to the closure of  $\mathcal{W}^0(A, B)$  in  $(a^p, M, b^{p+1})$  through  $\mathcal{G}(a^p, M, b^{p+1})$ . We additionally want to show that  $\mathcal{W}(M, b^p, b^{p+1}) \cap (a^p, M, b^{p+1})$  is contained in  $\mathcal{W}(b^p, b^{p+1}, M') \cap \mathcal{W}(M, b^p, b^{p+1})$ . Let us first show that in case (b) with  $\phi(b^{p+1}) = \phi(b^p) + 1$ , we have  $M' \in \sigma^{ss}$ . This follows from  $\phi(b^p) < \phi(a^p) < \phi(M) + 1$  and looking at the Table (105). To show that the inequalities for  $(b^p, b^{p+1}, M')$  from Table (36) hold, we only need to show  $\phi(M') + 1 > \phi(b^{p+1})$ , but this follows from looking at the proof of (b.2) in Proposition 54.

Let us now conclude the final results of the proposition. As the closure of  $\mathcal{W}^0(A, B)$  in  $(a^p, M, b^{p+1})$  is given by  $\mathcal{W}^0(A, B) \cup \mathcal{G}(a^p, M, b^{p+1})$ , we see that

$$\overline{\mathcal{W}^0(A, B)} = \mathcal{W}^0(A, B) \cup \bigcup_p (\mathcal{G}(a^p, M, b^{p+1}) \cup \mathcal{G}(b^p, M', a^p)).$$



Moreover, this lies in the intersection of  $\mathcal{Ch}(A)$  and  $\mathcal{Ch}(B)$ . As it is the only set, where both  $A$  and  $B$  are stable, we see that it is equal to it. Using  $\xi$ , we also know that  $\mathcal{W}(a^p, a^{p+1}, M) \cap \mathcal{W}(M', a^p, a^{p+1})$  and  $\mathcal{W}(b^p, b^{p+1}, M') \cap \mathcal{W}(M, b^p, b^{p+1})$  attach to  $\overline{\mathcal{W}^0(A, B)}$  in  $\mathcal{G}(a^p, M, b^{p+1}) \cup \mathcal{G}(b^p, M', a^p)$ .

Finally, the sets

$$(102) \quad \begin{aligned} \mathcal{O}_m = & (a^m, a^{m+1}, M)_\emptyset \cup (M', a^m, a^{m+1})_\emptyset \cup (a^m, b^{m+1}, a^{m+1})_\emptyset \cup (b^m, b^{m+1}, M')_\emptyset \\ & \cup (M', b^m, b^{m+1})_\emptyset \cup (b^m, a^m, b^{m+1})_\emptyset \cup (a^m, M, b^{m+1})_\emptyset \cup (b^m, M', a^m)_\emptyset \end{aligned}$$

are open connected subsets with  $C_{1,\sigma\sigma} = \emptyset$  and they are pairwise disjoint as the cases (c) and (d) in Proposition 60 are only contained in  $(a^p, M, b^{p+1})$ . Therefore, the set of closed chambers with  $\emptyset$  stable is given by  $\{\overline{\mathcal{O}_m}\}_{m \in \mathbb{Z}}$ . This completes the proof of the statements in the proposition.




FIGURE 4.  $(a^p, M, b^{p+1})$  with subsets as in Proposition 60 and its walls

□

**Remark 67.** Notice that to go from the chamber  $\mathcal{Ch}(A)$  into the chamber  $\mathcal{Ch}(B)$ , one needs to either pass through the wall  $\overline{\mathcal{W}^0(A, B)}$  or to go through  $\mathcal{O}_m$  for some  $m$  and pass through the walls  $\mathcal{W}(a^m, a^{m+1}, M) \cup \mathcal{W}(M', a^m, a^{m+1})$  and  $\mathcal{W}(b^m, b^{m+1}, M') \cup \mathcal{W}(M, b^m, b^{m+1})$ .

#### APPENDIX A. DETAILS ON $C_{l,\sigma\sigma}(\mathcal{T})$ , $C_{l,\sigma}(\mathcal{T})$ AS $\sigma$ VARIES IN $\text{Stab}(\mathcal{T})$ AND $l \geq -1$

In Subsection 6.4 we determine  $C_{l,\sigma}(\mathcal{T})$ ,  $C_{l,\sigma\sigma}(\mathcal{T})$  for  $\sigma \in \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  and  $l \geq -1$ . The results are contained in the tables (103), (104), (105), (106) and (107). The sets specified in the first column in each table are non-empty.

The following table collects results from Propositions 46, 51, Corollary 49, Lemma 52:

(103)

$\sigma \in (b^p, b^{p+1}, M')$ $v = \phi(b^p) + 1, q = \phi(M')$ $t = \arg_{(\phi(b^p)-1, \phi(b^p))}(Z(b^p) - Z(b^{p+1}))$	$C_{0,\sigma\sigma}(\mathcal{T}) = ?$ and $M \in \sigma^{ss}?$	$C_{1,\sigma}(\mathcal{T})$	$C_{1,\sigma\sigma}(\mathcal{T})$
$\phi(b^{p+1}) > v$ and $q > \phi(b^{p+1})$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M \notin \sigma^{ss}$	$\emptyset$	$\emptyset$
$\phi(b^{p+1}) > v$ and $q \leq \phi(b^{p+1})$	$\frac{\{\alpha^p\} \subset C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}}{\beta^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff M \in \sigma^{ss} \iff \arg_{[q, q+1]}(Z(b^{p+1}) + Z(M')) \leq v}$	$\emptyset$	$\emptyset$
$\phi(b^{p+1}) = v$ and $q > v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M \notin \sigma^{ss}$	$\{B\}$	$\{B\}$
$\phi(b^{p+1}) = v$ and $q < v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^p, \beta^p\}$ $M \in \sigma^{ss}$ and $q < \phi(M) + 1$ <sup>6</sup>	$\{B\}$	$\{B\}$
$\phi(b^{p+1}) = v$ and $q = v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \geq p\}$ $M \in \sigma^{ss}$ and $q = \phi(M) + 1$	$\{A, B\}$	$\{B\}$
$\phi(b^{p+1}) < v$ and $q - 1 \geq t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset, M \in \sigma^{ss} \iff q - 1 = t$	$\{B\}$	$\{B\}$
$\phi(b^{p+1}) < v$ and $q - 1 < t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : N \leq j \leq u\}$ , where $N \leq u$ are determined in Proposition 51 (a.2)	$\{B\}$	$\{B\}$

The last table and Remark 43 imply:

(104)

$\sigma \in (a^p, a^{p+1}, M)$ $v = \phi(a^p) + 1, q = \phi(M)$ $t = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(a^{p+1}))$	$C_{0,\sigma\sigma}(\mathcal{T}) = ?$ and $M' \in \sigma^{ss}?$	$C_{1,\sigma}(\mathcal{T})$	$C_{1,\sigma\sigma}(\mathcal{T})$
$\phi(a^{p+1}) > v$ and $q > \phi(a^{p+1})$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M' \notin \sigma^{ss}$	$\emptyset$	$\emptyset$
$\phi(a^{p+1}) > v$ and $q \leq \phi(a^{p+1})$	$\frac{\{\beta^{p+1}\} \subset C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^{p+1}\}}{\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff M' \in \sigma^{ss} \iff \arg_{[q, q+1]}(Z(a^{p+1}) + Z(M)) \leq v}$	$\emptyset$	$\emptyset$
$\phi(a^{p+1}) = v$ and $q > v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M' \notin \sigma^{ss}$	$\{A\}$	$\{A\}$
$\phi(a^{p+1}) = v$ and $q < v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{p+1}, \alpha^p\}$ $M' \in \sigma^{ss}$ and $q < \phi(M') + 1$	$\{A\}$	$\{A\}$
$\phi(a^{p+1}) = v$ and $q = v$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{j+1}, \alpha^j : j \geq p\}$ $M' \in \sigma^{ss}$ and $q = \phi(M') + 1$	$\{A, B\}$	$\{A\}$
$\phi(a^{p+1}) < v$ and $q - 1 \geq t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset, M' \in \sigma^{ss} \iff q - 1 = t$	$\{A\}$	$\{A\}$
$\phi(a^{p+1}) < v$ and $q - 1 < t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{j+1}, \alpha^j : N \leq j \leq u\}$ , where $N \leq u$ are determined in Proposition 51 (a.2) by replacing $b$ with $a$ and exchanging $M'$ and $M$	$\{A\}$	$\{A\}$

<sup>6</sup>In details, the second row of this box says that for  $\sigma \in (b^p, b^{p+1}, M')$ ,  $\phi(b^{p+1}) = \phi(b^p) + 1$ ,  $\phi(M') < \phi(b^p) + 1$  hold both  $M \in \sigma^{ss}$  and  $\phi(M') < \phi(M) + 1$ . We will use this for the proof of (97) in Section 7.

The following table collects results from Propositions 46, 54, Corollary 50, Lemma 55:

(105)

$\sigma \in (M, b^p, b^{p+1})$ $v = \phi(b^{p+1}) - 1, s = \phi(M) + 1,$ $t = \arg_{(\phi(b^p)-1, \phi(b^p))}(Z(b^p) - Z(b^{p+1}))$	$C_{0,\sigma\sigma}(\mathcal{T}) = ?$ and $M' \in \sigma^{ss}?$	$C_{1,\sigma}(\mathcal{T})$	$C_{1,\sigma\sigma}(\mathcal{T})$
$v > \phi(b^p)$ and $s < \phi(b^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M' \notin \sigma^{ss}$	$\emptyset$	$\emptyset$
$v > \phi(b^p)$ and $s \geq \phi(b^p)$	$\frac{\{\beta^p\} \subset C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^p\}}{\alpha^p \in C_{0,\sigma\sigma}(\mathcal{T}) \iff M' \in \sigma^{ss} \iff}$ $v \leq \arg_{(s-1, s]}(Z(b^p) - Z(M))$	$\emptyset$	$\emptyset$
$v = \phi(b^p)$ and $s < \phi(b^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M' \notin \sigma^{ss}$	$\{B\}$	$\{B\}$
$v = \phi(b^p)$ and $s > \phi(b^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^p, \beta^p\}$ $M' \in \sigma^{ss}$ and $\phi(M') < s$	$\{B\}$	$\{B\}$
$v = \phi(b^p)$ and $s = \phi(b^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : j \leq p\}$ $M' \in \sigma^{ss}$ and $\phi(M') = s$	$\{A, B\}$	$\{B\}$
$v < \phi(b^p)$ and $s \leq t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset, M' \in \sigma^{ss} \iff s = t$	$\{B\}$	$\{B\}$
$v < \phi(b^p)$ and $s > t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^j, \beta^j : N \leq j \leq u\},$ where $N \leq u$ are determined in Proposition 54 (a.2)	$\{B\}$	$\{B\}$

This table and Remark 43 imply:

(106)

$\sigma \in (M', a^p, a^{p+1})$ $v = \phi(a^{p+1}) - 1, s = \phi(M') + 1,$ $t = \arg_{(\phi(a^p)-1, \phi(a^p))}(Z(a^p) - Z(a^{p+1}))$	$C_{0,\sigma\sigma}(\mathcal{T}) = ?$ and $M \in \sigma^{ss}?$	$C_{1,\sigma}(\mathcal{T})$	$C_{1,\sigma\sigma}(\mathcal{T})$
$v > \phi(a^p)$ and $s < \phi(a^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M \notin \sigma^{ss}$	$\emptyset$	$\emptyset$
$v > \phi(a^p)$ and $s \geq \phi(a^p)$	$\frac{\{\alpha^p\} \subset C_{0,\sigma\sigma}(\mathcal{T}) \subset \{\alpha^p, \beta^{p+1}\}}{\beta^{p+1} \in C_{0,\sigma\sigma}(\mathcal{T}) \iff M \in \sigma^{ss}}$ $v \leq \arg_{(s-1, s]}(Z(a^p) - Z(M'))$	$\emptyset$	$\emptyset$
$v = \phi(a^p)$ and $s < \phi(a^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ and $M \notin \sigma^{ss}$	$\{A\}$	$\{A\}$
$v = \phi(a^p)$ and $s > \phi(a^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{p+1}, \alpha^p\}$ $M \in \sigma^{ss}$ and $\phi(M) < s$	$\{A\}$	$\{A\}$
$v = \phi(a^p)$ and $s = \phi(a^p)$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{j+1}, \alpha^j : j \leq p\}$ $M \in \sigma^{ss}$ and $\phi(M) = s$	$\{A, B\}$	$\{A\}$
$v < \phi(a^p)$ and $s \leq t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset, M \in \sigma^{ss} \iff s = t$	$\{A\}$	$\{A\}$
$v < \phi(a^p)$ and $s > t$	$C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{j+1}, \alpha^j : N \leq j \leq u\},$ where $N \leq u$ are determined in Proposition 54 (a.2) by replacing $b$ with $a$ and exchanging $M'$ and $M$	$\{A\}$	$\{A\}$

The following table collects results from Corollary 48 and Propositions 46, 47

(107)

	$X = C_{0,\sigma\sigma}(\mathcal{T}) = ?$ and $M, M' \in \sigma^{ss}?$	$C_{1,\sigma}(\mathcal{T})$
$\sigma \in (b^p, a^p, b^{p+1})$	$X \subset \{\alpha^p, \beta^p\}$ $M' \in \sigma^{ss} \iff \alpha^p \in X \iff \phi(b^{p+1}) - 1 \leq \phi(a^p) < \phi(b^{p+1})$ $M \in \sigma^{ss} \iff \beta^p \in X \iff \phi(a^p) - 1 \leq \phi(b^p) < \phi(a^p)$	$\emptyset$
$\sigma \in (a^p, b^{p+1}, a^{p+1})$	$X \subset \{\beta^{p+1}, \alpha^p\}$ $M \in \sigma^{ss} \iff \beta^{p+1} \in X \iff \phi(a^{p+1}) - 1 \leq \phi(b^{p+1}) < \phi(a^{p+1})$ $M' \in \sigma^{ss} \iff \alpha^p \in X \iff \phi(b^{p+1}) - 1 \leq \phi(a^p) < \phi(b^{p+1})$	$\emptyset$

In Subsection 6.5 we determine the behavior of  $C_{l,\sigma}(\mathcal{T})$ ,  $C_{l,\sigma\sigma}(\mathcal{T})$  in  $\text{Stab}(\mathcal{T}) \setminus (\mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st})$ . The results there imply Proposition 68 (the proof is given after Corollary 61):

**Proposition 68.** *Let  $\sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$ .*

*If  $\{M, M'\} \subset \sigma^{ss}$ , then  $\phi(M) = \phi(M')$  and  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \{A, B\}$ ,  $C_{0,\sigma\sigma}(\mathcal{T}) = C_0(\mathcal{T})$ .*

*If  $\{M, M'\} \not\subset \sigma^{ss}$ , then  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T}) = \emptyset$ ,  $|C_{0,\sigma\sigma}(\mathcal{T})| \leq 1$ .*

*The set  $\{\sigma : \sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st} \text{ and } \{M, M'\} \subset \sigma^{ss}\}$  is disjoint union (any two summands are non-empty and disjoint):*

$$(108) \quad \bigcup_{p \in \mathbb{Z}} \{\sigma \in (a^p, M, b^{p+1}) : \phi(M) + 1 = \phi(b^{p+1}) = \phi(a^p) + 1\} \\ \cup \bigcup_{p \in \mathbb{Z}} \{\sigma \in (b^p, M', a^p) : \phi(M') + 1 = \phi(a^p) = \phi(b^p) + 1\} \\ \cup \bigcap_{p \in \mathbb{Z}} (a^p, M, b^{p+1}) \cap (b^p, M', a^p).$$

*The set  $\{\sigma : \sigma \notin \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st} \text{ and } \{M, M'\} \not\subset \sigma^{ss}\}$  is disjoint union (any two summands are disjoint and non-empty):*

$$(109) \quad \bigcup_{p \in \mathbb{Z}} \left\{ \sigma \in (a^p, M, b^{p+1}) : \begin{array}{l} \phi(M) + 1 < \phi(b^{p+1}) \text{ and } \phi(a^p) \leq \phi(M) \\ \text{or } \phi(M) + 1 = \phi(b^{p+1}) \text{ and } \phi(a^p) < \phi(M) \end{array} \right\} \\ \cup \bigcup_{p \in \mathbb{Z}} \left\{ \sigma \in (b^p, M', a^p) : \begin{array}{l} \phi(M') + 1 < \phi(a^p) \text{ and } \phi(b^p) \leq \phi(M') \\ \text{or } \phi(M') + 1 = \phi(a^p) \text{ and } \phi(b^p) < \phi(M') \end{array} \right\}.$$

*For  $\sigma \in (a^p, M, b^{p+1})$  we have: (a)  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^p\}$ , when  $\phi(a^p) = \phi(M)$ ,  $\phi(M) + 1 < \phi(b^{p+1})$ ; (b)  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\beta^{p+1}\}$ , when  $\phi(a^p) < \phi(M)$ ,  $\phi(M) + 1 = \phi(b^{p+1})$ ; (c)  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ , when  $\phi(a^p) < \phi(M)$ ,  $\phi(M) + 1 < \phi(b^{p+1})$ .*

*For  $\sigma \in (b^p, M', a^p)$  we have: (a)  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^{p-1}\}$ , when  $\phi(b^p) = \phi(M')$ ,  $\phi(M') + 1 < \phi(a^p)$ ; (b)  $C_{0,\sigma\sigma}(\mathcal{T}) = \{\alpha^p\}$ , when  $\phi(b^p) < \phi(M')$ ,  $\phi(M') + 1 = \phi(a^p)$ ; (c)  $C_{0,\sigma\sigma}(\mathcal{T}) = \emptyset$ , when  $\phi(b^p) < \phi(M')$ ,  $\phi(M') + 1 < \phi(a^p)$ .*

Let us now discuss the proofs of the main statements which appeared in the introduction. Most of them follow straightforwardly from tables (103), (104), (105), (106), (107) and Proposition 68, however in some cases more work is required and it was carried out in the paper.

*Proof of Proposition 5.* Proposition 5 (a) is obtained by looking at all the cases in the tables and Proposition 68, recalling that all summands in the unions (108), (109) are non-empty and that stability conditions in each row of the listed tables exist. Proposition 5 (b) follows again by looking at all the cases in the tables and Proposition 68. To prove that all of the subsets in the list are taken we need to use Proposition 53 for  $\{\alpha^j, \beta^j : i \leq j \leq k\}$  and then Remark 43 for  $\{\alpha^j, \beta^{j+1} : i \leq j \leq k\}$ .  $\square$

*Proof of the first sentence in Proposition 7.* We first prove that the three sets are equal.

The first equality is in (40). If  $\sigma \in \mathfrak{T}_a^{st} \cup \mathfrak{T}_b^{st}$  we see in tables (104), (106), (103), (105), (107) that  $|C_{1,\sigma}(\mathcal{T})| = 0$  iff  $|C_{1,\sigma\sigma}(\mathcal{T})| = 0$ . Otherwise we fall in Proposition 68, where  $C_{1,\sigma\sigma}(\mathcal{T}) = C_{1,\sigma}(\mathcal{T})$ .

We prove that the set on the RHS of the proved equality is open. To that end, we note that it is the complement of  $\{\sigma \in \text{Stab}(\mathcal{T}) : A \in C_{1,\sigma\sigma}(\mathcal{T})\} \cup \{\sigma \in \text{Stab}(\mathcal{T}) : B \in C_{1,\sigma\sigma}(\mathcal{T})\}$ . And we use Corollary 27 to deduce that the latter union is a closed subset.

Looking at tables (105), (103), etc we see that there are non-empty open subsets, where  $C_{1,\sigma\sigma}(\mathcal{T}) \neq \emptyset$  (see also Remark 62), hence the set in question is not dense.  $\square$

*Proof of Proposition 8.* In tables (104), (106), (103), (105) we see that there are  $\sigma$  such that  $C_{1,\sigma}(\mathcal{T}) = \{A, B\}$ ,  $C_{1,\sigma\sigma}(\mathcal{T}) = \{A\}$  and there are  $\sigma$  such that  $C_{1,\sigma}(\mathcal{T}) = \{A, B\}$ ,  $C_{1,\sigma\sigma}(\mathcal{T}) = \{B\}$ .  $\square$

*Proof of Proposition 10.* The first equivalence follows from (38) and Corollaries 56, 45, 61. The second equivalence follows from (38), tables (104), (106), (103), (105), (107) and Proposition 68. The last equivalence is in (39).  $\square$

*Proof of Theorem 6.* This relies also on the tables and Proposition 68, however, as for the proof of Proposition 10 it requires additional work. which is carried out in Section 7.  $\square$

## REFERENCES

- [1] I. Gelfand, Y. Manin, *Methods of Homological algebra*, 2nd edition, Springer.
- [2] T. Bridgeland, *Stability conditions on triangulated categories*, *Annals of Math.* 166 no. 2, 317-345 (2007).
- [3] A. Bojko, *Stability conditions on quivers and semistable noncommutative curve counting*, Master thesis, University of Vienna. Fakultät für Mathematik, Betreuer: L. Katzarkov and G. Dimitrov, 2018. <http://othes.univie.ac.at/52820/>
- [4] W. Crawley-Boevey, *Exceptional sequences of representations of quivers*, in 'Representations of algebras', Proc. Ottawa 1992, eds V. Dlab and H. Lenzing, Canadian Math. Soc. Conf. Proc. 14 (Amer. Math. Soc., 1993), 117-124
- [5] G. Dimitrov and L. Katzarkov, *Some new categorical invariants*, *Selecta Mathematica*, doi: 10.1007/s00029-019-0493-8
- [6] G. Dimitrov and L. Katzarkov, *Non-commutative counting invariants and curve complexes*, arXiv: 1805.00294v4
- [7] G. Dimitrov and L. Katzarkov, *Non-semistable Exceptional Objects in Hereditary Categories*, *International Mathematics Research Notices*, rnv336, 85 pages. doi:10.1093/imrn/rnv336
- [8] G. Dimitrov and L. Katzarkov, *Bridgeland stability conditions on the acyclic triangular quiver*, *Advances in Mathematics* 288 (2016), 825-886
- [9] G. Dimitrov and L. Katzarkov, *Bridgeland stability conditions on wild Kronecker quivers*, *Advances in Mathematics* 352 (2019) 27-55, doi: 10.1016/j.aim.2019.05.032

- [10] G. Dimitrov and L. Katzarkov, *More finite sets coming from non-commutative counting*, arXiv:1903.00295 .
- [11] M. Kontsevich, and A. Rosenberg,; *Noncommutative smooth spaces*, The Gelfand Mathematical Seminars, 1996-1999, 85-108, Birkhäuser, Boston, 2000.
- [12] D. Orlov, Geometric realizations of quiver algebras, Proceedings of the Steklov institute of mathematics, Vol. 290, 2015

(Bojko) ETH ZÜRICH, PROFESSUR FÜR MATHEMATIK, HG J 16.2 RÄMISTRASSE 101, ZÜRICH, SWITZERLAND  
*Email address:* `arkadij.bojko@math.ethz.ch`

(Dimitrov) UNIVERSITÄT WIEN, FAKULTÄT FÜR MATHEMATIK, OSKAR-MORGENSTERN-PLATZ 1, 1090 WIEN, ÖSTERREICH,  
*Email address:* `george.dimitrov@univie.ac.at`