

Three dimensional transitive sets of C^1 generic vector fields

Manseob Lee

*Department of Mathematics, Mokwon University,
Daejeon, 302-729, Korea.*

E-mail: lmsds@mokwon.ac.kr.

Abstract

In the paper, we show that for a generic C^1 vector field X on a closed three dimensional manifold M , any isolated transitive set of X is singular hyperbolic.

1 Introduction

The transitivity is a symbol of chaotic property for differential dynamical systems. The C^1 robust transitivity for diffeomorphisms are well investigate in a series of works [2, 3, 5], and then we have a well characterization on isolated transitive sets of C^1 generic diffeomorphisms at the same time. From the main result of [1] we know that if every isolated transitive set of a C^1 generic diffeomorphism admit a nontrivial dominated splitting, then it is volume hyperbolic.

It is well known that a singularity-free flow, for an instance, a suspension of a diffeomorphism, will take similar phenomenons of diffeomorphisms. However, once the recurrent regular points can accumulates a singularity, such as the Lorenz-like systems, we will meet something new. For an instance, in [14], one have to use a new notion of singular hyperbolicity to characterize the robustly transitive sets of a 3-dimensional flow. Here the singular hyperbolicity is a generalization of hyperbolicity so that we can give the Lorenz attractor and Smale's horseshoe a unified characterization. In this article, we will show that an isolated transitive of C^1 generic vector field on 3-dimensional manifolds will be singular hyperbolic. That meas, every isolated transitive set of a C^1 generic vector field looks like a Lorenz attractor[6, 9].

Let us be precise now. Denote by M a compact $d(\geq 2)$ -dimensional smooth Riemannian manifold without boundary and by $\mathfrak{X}^1(M)$ the set of C^1 vector fields on M endowed

¹ 2010 *Mathematical Subject Classification*:37C20; 37C27; 37D50.

Key words and phrases:transitive set; generic; local star; Lyapunove stable; singular hyperbolic.

with the C^1 topology. Every $X \in \mathfrak{X}^1(M)$ generates a flow $X^t : M \times \mathbb{R} \rightarrow M$ that is a C^1 map such that $X^t : M \rightarrow M$ is a diffeomorphism for all $t \in \mathbb{R}$ and then $X^0(x) = x$ and $X^{t+s}(x) = X^t(X^s(x))$ for all $s, t \in \mathbb{R}$ and $x \in M$. An *orbit* of X corresponding a point $x \in M$ is the set $Orb(x) = \{X^t(x) : t \in \mathbb{R}\}$. Let $Sing(X)$ denotes the set of singularities of X and $Per(X)$ is the set of periodic orbits of X . Denote by $Crit(X) = Sing(X) \cup P(X)$ the set of all critical points of X .

Let $\Lambda \subset M$ be a closed X^t -invariant set. We say that Λ is a *hyperbolic set* of X if there are constants $C > 0, \lambda > 0$ and a DX^t -invariant continuous splitting $T_\Lambda M = E^s \oplus \langle X \rangle \oplus E^u$ such that

$$\|DX^t|_{E_x^s}\| \leq Ce^{-\lambda t} \quad \text{and} \quad \|DX^{-t}|_{E_x^u}\| \leq Ce^{-\lambda t}$$

for $t > 0$ and $x \in \Lambda$, where $\langle X(x) \rangle$ denotes the space spanned by $X(x)$, which is 0-dimensional if x is a singularity or 1-dimensional if x is not singularities. For any critical point $x \in Crit(X)$, if its orbit is a hyperbolic set, we denote by $index(x) = \dim E_x^s$.

Now let us recall the singular hyperbolicity firstly given by Morale, Pacifico and Pujals [14] which is an extension of hyperbolicity. We say that a compact invariant set Λ is *positively singular hyperbolic* for X if there are constants $K \geq 1$ and $\lambda > 0$, and a continuous invariant $T_\Lambda M = E^s \oplus E^c$ with respect to DX^t such that

(i) E^s is (K, λ) -dominated by E^{cu} , that is,

$$\|DX^t|_{E^s(x)}\| \cdot \|DX^{-t}|_{E^c(X^t(x))}\| \leq Ke^{-\lambda t}, \quad \forall x \in \Lambda \text{ and } t \geq 0.$$

(ii) E^s is contracting, that is,

$$\|DX^t|_{E^s(x)}\| \leq Ke^{-\lambda t}, \quad \forall x \in \Lambda \text{ and } t \geq 0.$$

(iii) E^{cu} is sectional expanding, that is, for any $x \in \Lambda$ and any 2-dimensional subspace $L \subset E^c(x)$,

$$|\det(DX^t|_L)| \geq K^{-1}e^{\lambda t}, \quad \forall t \geq 0.$$

We say that Λ is *negatively singular hyperbolic* for X if Λ is positively singular hyperbolic for $-X$, and then say that Λ is *singular hyperbolic* for X if it is either positively singular hyperbolic for X , or negatively singular hyperbolic for X . Definitely, we can see that if Λ is singular hyperbolic for X and it does not contain singularities then it is hyperbolic (see [14, Proposition 1.8] for a proof). In the paper, we consider the relation between transitivity and hyperbolicity for an isolated compact invariant set. We say that Λ is *transitive* if there is $x \in \Lambda$ such that $\omega(x) = \Lambda$, where $\omega(x)$ is the omega limit set of x . We say that a closed X^t -invariant set Λ is *isolated* (or *locally maximal*) if there exists a neighborhood U of Λ such that

$$\Lambda = \Lambda_X(U) = \bigcap_{t \in \mathbb{R}} X^t(U).$$

Here U is said to be *isolated neighborhood* of Λ .

For the 3-dimensional case, Morales, Pacifico and Pujals [14] proved that if Λ is a robustly transitive set containing singularities then it is singular hyperbolic set for X . Here we will consider C^1 generic vector fields. We say that a subset $\mathcal{G} \subset \mathfrak{X}^1(M)$ is *residual* if it contains a countable intersection of open and dense subsets of $\mathfrak{X}^1(M)$. A property is called *C^1 generic* if it holds in a residual subset of $\mathfrak{X}^1(M)$.

We give the following characterization of the isolated transitive sets of a C^1 generic vector field on 3-dimensional Riemannian manifold.

Theorem A. *For C^1 generic $X \in \mathfrak{X}^1(M)$, an isolated transitive set Λ is singular hyperbolic.*

2 Transitivity and locally Star condition

Let M be a three dimensional smooth Riemannian manifold and let $X \in \mathfrak{X}^1(M)$ be the set of C^1 vector fields on M endowed with the C^1 topology. Here we collect some known generic properties for C^1 vector fields.

Proposition 2.1 *There is a residual set $\mathcal{G}_1 \subset \mathfrak{X}^1(M)$ such that for any $X \in \mathcal{G}_1$, X satisfies the following properties:*

1. X is a Kupka-Smale system, that is, every periodic orbits and singularity of X is hyperbolic, and the corresponding invariant manifolds intersect transversely.
2. if there is a sequence of vector fields $\{X_n\}$ with critical orbit $\{P_n\}$ of X_n such that $X_n \rightarrow X$, $\text{index}(P_n) = i$ and $P_n \rightarrow_H \Lambda$ then there is a sequence of critical orbit $\{Q_n\}$ of X such that $\text{index}(Q_n) = i$ and $Q_n \rightarrow_H \Lambda$, where \rightarrow_H is the Hausdorff limit.

The item 1 is from the famous Kupka-Smale theorem (see [15]) and item 2 is from [16, Lemma 3.5]

From the item 1 of Proposition 2.1, we can see that if Λ is a trivial transitive set, that is, Λ is a periodic orbit or a singularity, then it should be hyperbolic and automatically singular hyperbolic. To prove Theorem A, we just need to consider the nontrivial case. Hereafter, we assume that Λ is a nontrivial transitive set of X . One can see that if Λ is a nontrivial transitive set, then Λ contains no hyperbolic sinks or sources.

Let U be an isolated neighborhood of Λ . Then for Y C^1 close to X , denote by

$$\Lambda_Y(U) = \bigcap_{t \in \mathbb{R}} Y^t(U)$$

the maximal invariant set of Y in U .

Lemma 2.2 *Let $\mathcal{G}_1 \subset \mathfrak{X}^1(M)$ be the residual set given in Proposition 2.1. For any $X \in \mathcal{G}_1$, if Λ is an isolated nontrivial transitive set of X , then there are a C^1 neighborhood $\mathcal{U}(X)$ of X and a neighborhood U of Λ such that for any $Y \in \mathcal{U}(X)$, we have every $\gamma \in \Lambda_Y(U) \cap \text{Per}(Y)$ is hyperbolic and $\text{index}(\gamma) = 1$.*

Proof. Let \mathcal{G}_1 be the residual set in Proposition 2.1 and let Λ be an isolated transitive set of $X \in \mathcal{G}_1$. Suppose on the contrary that for any C^1 neighborhood $\mathcal{U}(X)$ of X and any neighborhood U of Λ , there is $Y \in \mathcal{U}(X)$ such that Y has a periodic orbit Q whose index is not 1. Then we have three cases: (i) Q is not hyperbolic, (ii) Q is hyperbolic but $\text{index}(Q) = 0$ or (iii) $\text{index}(Q) = 2$. Note that if the periodic orbit Q is not hyperbolic for Y then we can take a vector field Z C^1 arbitrary close to Y such that either Q is a sink for Z or Q is a source for Z . Then we also have the case cases (ii) or (iii) happening. Thus we can take sequences $Y_n \rightarrow X$ and a periodic orbit P_n of Y_n such that $\text{index}(P_n) = 0$ or 2 and

$$\lim_{n \rightarrow \infty} P_n = \Gamma \subset \Lambda.$$

Then we can take a sequence of vector fields X_n tends to X and periodic orbits $\{Q_n\}$ of X_n with $\text{index}(Q_n) = 0$ or 2 such that

$$\lim_{n \rightarrow \infty} Q_n = \Gamma \subset \Lambda.$$

Without loss of generality, we can assume that all Q_n have the same index 0 or 2 once we take a subsequence. By the item 2 of Proposition 2.1, we know that there is a sequence P_n of periodic orbit of X with index 0 or 2 converging into Λ . Since Λ is isolated, for sufficiently large n , we have $P_n \subset \Lambda$. This is a contradiction since Λ is a nontrivial transitive set. \square

Let Λ be a closed X^t -invariant set. We say Λ is *locally star* if there are a C^1 neighborhood $\mathcal{U}(X)$ of $X \in \mathfrak{X}^1(M)$ and a neighborhood U of Λ such that for any $Y \in \mathcal{U}(X)$, every periodic orbit of Y in $\Lambda_Y(U) = \bigcap_{t \in \mathbb{R}} Y^t(U)$ is hyperbolic and indices are same.

Corollary 2.3 *There is a residual set $\mathcal{R} \subset \mathfrak{X}^1(M)$ such that for any $X \in \mathcal{R}$, if Λ is an isolated transitive set of X which is not an orbit then Λ is a local star.*

Proof. Let $X \in \mathcal{R} = \mathcal{G}_1$ and let Λ be an isolated transitive set. By Lemma 2.2, there are a C^1 neighborhood $\mathcal{U}(X)$ of X and a neighborhood U of Λ such that for any $Y \in \mathcal{U}(X)$, every periodic orbit $\gamma \in \Lambda_Y(U) \cap \text{Per}(Y)$ is hyperbolic and $\text{index}(\gamma) = 1$. Thus Λ is a local star. \square

3 Transitivity and Lyapunov stability

Suppose $\sigma \in \text{Sing}(X)$ is hyperbolic. Then we denote by

$$W^s(\sigma) = W^s(\sigma, X) = \{y \in M : d(X^t(\sigma), X^t(y)) \rightarrow 0 \text{ as } t \rightarrow \infty\}$$

$$W^u(\sigma) = W^u(\sigma, X) = \{y \in M : d(X^t(\sigma), X^t(y)) \rightarrow 0 \text{ as } t \rightarrow -\infty\},$$

where $W^s(\sigma, X)$ is said to be the *stable manifold* of σ and $W^u(\sigma, X)$ is said to be the *unstable manifold* of σ . It is known that $\text{index}(\sigma) = \dim W^s(\sigma)$.

If X is a Kupka-Smale vector field, then X contains finitely many singularities and every singularity is hyperbolic. Thus by the structural stability of hyperbolic singularity we know that there are a C^1 neighborhood $\mathcal{U}(X)$ of X and a neighborhood U of Λ such that for any $Y \in \mathcal{U}(X)$, every $\sigma \in \Lambda_Y(U) \cap \text{Sing}(Y) \subset U$ is hyperbolic.

Lemma 3.1 *Let $\mathcal{G}_1 \subset \mathfrak{X}^1(M)$ be the residual set given in Proposition 2.1. For any $X \in \mathcal{G}_1$, if Λ is an isolated nontrivial transitive set of X , then there are a C^1 neighborhood $\mathcal{U}(X)$ of X and a neighborhood U of Λ such that for any $Y \in \mathcal{U}(X)$, every singularities in $\Lambda_Y(U)$ is neither sinks nor sources.*

Proof. We prove it by contradiction. Assume the contrary of the lemma, then we can find a sequence of vector fields X_n tends to X and a sequence of singularity σ_n of X_n such that σ_n tends to a point σ such that the index of σ_n equals to 0 or 2. Without loss of generality, we assume that every σ_n has index 0, then we can see that σ is a singularity. Since $X \in \mathcal{G}_1$, we have σ is hyperbolic. By the structural stability of σ we know σ have index 0 too. This contradicts with Λ is a nontrivial transitive set. \square

Lemma 3.2 *Let Λ be a transitive set of a C^1 vector field X . If $\sigma \in \Lambda \cap \text{Sing}(X)$ is hyperbolic then $(W^s(\sigma) \setminus \{\sigma\}) \cap \Lambda \neq \emptyset$ and $(W^u(\sigma) \setminus \{\sigma\}) \cap \Lambda \neq \emptyset$.*

Proof. We consider the case of $(W^s(\sigma) \setminus \{\sigma\}) \cap \Lambda \neq \emptyset$ (Other case is similar). Since $\sigma \in \Lambda = \omega(x)$ for some $x \in \Lambda$, there is $t_n \in \mathbb{R}^+$ with $t_n \rightarrow \infty$ such that $X^{t_n}(x) \rightarrow \sigma$. Since σ is hyperbolic, we can take $\epsilon > 0$ such that

$$\{x : X^t(x) \in B_\epsilon(\sigma), \text{ for all } t > 0\} \subset W^s(\sigma).$$

Denote by $x_n = X^{t_n}(x)$. For n large enough, $x_n \in B_\epsilon(\sigma)$. Let $\tau_n = \sup\{t : X^{(-t,0)}(x_n) \subset B_\epsilon(\sigma)\}$. Then we have $X^{-\tau_n}(x_n) \in \partial B_\epsilon(\sigma)$. Let $y_n = X^{-\tau_n}(x_n)$. We can see that $\tau_n \rightarrow +\infty$ as $n \rightarrow \infty$. Take a subsequence if necessary, we can assume that $y_n \rightarrow y$ as $n \rightarrow \infty$. It is easy to see that $y \neq \sigma$. For every y_n , we have $X^{(0,\tau_n)}(y_n) \in \partial B_\epsilon(\sigma)$. By the continuity of the flow X^t , we have $X^{(0,+\infty)}(y) \subset B_\epsilon(\sigma)$, then $y \in W^s(\sigma) \setminus \{\sigma\}$. \square

The following is the connecting lemma for C^1 vector fields.

Lemma 3.3 [19] *Let $X \in \mathfrak{X}^1(M)$ and $z \in M$ be neither periodic nor singular of X . For any C^1 neighborhood $\mathcal{U}(X) \subset \mathfrak{X}^1(M)$ of X , there exist three numbers $\rho > 1, L > 1$ and $\delta_0 > 0$ such that for any $0 < \delta \leq \delta_0$ and any two points x, y outside the tube $\Delta = B_\delta(X^{[0,L]}(z))$ (or $\Delta = B_\delta(X^{[-L,0]}(z))$), if the positive X -orbit of x and the negative X -orbit of y both hit $B_{\delta/\rho}(z)$, then there exists $Y \in \mathcal{U}(X)$ with $Y = X$ outside Δ such that y is on the positive Y -orbit of x .*

Lemma 3.4 *Let Λ be a transitive set for X and $\sigma \in \Lambda \cap \text{Sing}(X)$ be hyperbolic. Then for any C^1 neighborhood $\mathcal{U}(X)$ of X , any point $y \in \Lambda$ and any neighborhood U of y , there is $Y \in \mathcal{U}(X)$ such that $W^s(\sigma, Y) \cap U \neq \emptyset$, where $W^s(\sigma, Y)$ is the stable manifold of σ with respect to Y .*

Proof. Let $\mathcal{U}(X)$ be fixed. By Lemma 3.2, there is a point $x \in (W^s(\sigma) \setminus \{\sigma\}) \cap \Lambda$. Then x is neither a singularity nor a periodic point. Let L, ρ and δ_0 be the constant given by Lemma 3.3. Take a point $X^T(x)$ with $T > L$ and $\delta > 0$ such that the tube

$$\Delta = B_\delta(X^{[0,L]}(x)) \cap X^{[T,+\infty)}(x) = \emptyset.$$

Since Λ is transitive, there is $z \in \Lambda$ such that $\omega(z) = \Lambda$. For any small neighborhood U of y , we can find $0 < s < t$ such that $X^s(z) \in U$ and $X^t(z) \in B_{\delta/\rho}(x)$. Let $q = X^T(x)$ and $p = X^s(z)$. Then by Lemma 3.3, there is $Y \in \mathcal{U}(X)$ such that $Y^t(p) = q$ for some $t > 0$. Since $q = X^T(x) \in W^s(\sigma)$, we have $p \in W^s(\sigma, Y)$. \square

From Lemma 3.1 we know that if $X \in \mathcal{G}_1$, and Λ is an isolated nontrivial transitive set of X , then every $\sigma \in \Lambda \cap \text{Sing}(X)$ has index 1 or 2.

Lemma 3.5 *There is a residual set $\mathcal{G}_2 \subset \mathfrak{X}^1(M)$ with the following property. For any $X \in \mathcal{G}_2$ and any isolated nontrivial transitive set Λ of X , if there is $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 2$ then $\Lambda \subset \overline{W^u(\sigma)}$. Symmetrically, if there is $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 1$ then $\Lambda \subset \overline{W^s(\sigma)}$.*

Proof. Let $\mathcal{O} = \{O_1, O_2, \dots, O_n, \dots\}$ be a countable basis of M . For each $m, k \in \mathbb{N}$, let

$$\begin{aligned} \mathcal{H}_{m,k} &= \{X \in \mathfrak{X}^1(M) : \text{there is a } C^1 \text{ neighborhood } \mathcal{U}(X) \text{ of } X \\ &\text{such that for any } Y \in \mathcal{U}(X), Y \text{ has a singularity } \sigma \in O_m \text{ with} \\ &\text{index}(\sigma) = 2 \text{ such that } W^u(\sigma, Y) \cap O_k \neq \emptyset\}. \end{aligned}$$

Then $\mathcal{H}_{m,k}$ is an open in $\mathfrak{X}^1(M)$. Let

$$\mathcal{N}_{m,k} = \mathfrak{X}^1(M) \setminus \overline{\mathcal{H}_{m,k}}.$$

Then $\mathcal{H}_{m,k} \cup \mathcal{N}_{m,k}$ is open and dense in $\mathfrak{X}^1(M)$. Let

$$\mathcal{G}_2 = \bigcap_{m,k \in \mathbb{N}} (\mathcal{H}_{m,k} \cup \mathcal{N}_{m,k}).$$

We will show that the residual set \mathcal{G}_2 satisfies the request of lemma. Let $X \in \mathcal{G}_2$ and Λ be an isolated transitive set and let $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 2$. Since σ is hyperbolic, we can take O_m such that O_m is an isolated neighborhood of σ . By the structural stability of hyperbolic singularity, there is a C^1 neighborhood $\mathcal{U}(X)$ of X such that for any $Y \in \mathcal{U}(X)$, Y has a unique hyperbolic singularity in O_m . For any $y \in \Lambda$ and any neighborhood U of y , we can choose $O_k \in \mathcal{O}$ such that $y \in O_k \subset U$.

Claim $X \notin \mathcal{N}_{m,k}$

Proof of Claim. For any neighborhood $\mathcal{V}(X) \subset \mathcal{U}(X)$, by Lemma 3.4, there is $Y \in \overline{\mathcal{V}(X)}$ such that Y has a singularity $\sigma \in O_m$ with $\text{index}(\sigma) = 2$ and $W^u(\sigma, Y) \cap O_k \neq \emptyset$. Note that if $W^u(\sigma, Y) \cap O_k \neq \emptyset$ then there is a C^1 neighborhood $\mathcal{U}(Y)$ of Y such that for any $Z \in \mathcal{U}(Y)$, we know that $W^u(\sigma, Z) \cap O_k \neq \emptyset$ by the continuity of the unstable manifold. Thus we have $Y \in \mathcal{H}_{m,k}$. Hence $X \in \overline{\mathcal{H}_{m,k}}$. This ends the proof of claim.

Then by claim, since $X \in \mathcal{G}_2$, we have $X \in \mathcal{H}_{m,k}$. Note that O_m is an isolated neighborhood of σ , by the definition of $\mathcal{H}_{m,k}$, we know that $W^u(\sigma) \cap O_k \neq \emptyset$. This prove that for every neighborhood U of y , we know that $W^u(\sigma) \cap U \neq \emptyset$. This means that $\Lambda \subset \overline{W^u(\sigma)}$. \square

We say that a closed X^t -invariant set Λ is *Lyapunov stable* for X if for every neighborhood U of Λ there is a neighborhood $V \subset U$ of Λ such that $X^t(V) \subset U$ for every $t \geq 0$.

Let σ be a hyperbolic singularity of X with $\dim W^u(\sigma) = 1$. Then $W^u(\sigma) \setminus \{\sigma\}$ can be divided into two connected branches Γ_1, Γ_2 , that is, $W^u(\sigma) = \{\sigma\} \cup \Gamma_1 \cup \Gamma_2$.

Lemma 3.6 *Let $X \in \mathfrak{X}^1(M)$ and Λ be a transitive set of X . Assume $\sigma \in \Lambda$ is a hyperbolic singularity of X with $\dim W^u(\sigma) = 1$. Let $\Gamma_1 = \text{Orb}(x_1)$ and $\Gamma_2 = \text{Orb}(x_2)$ be the two branches of $W^u(\sigma) \setminus \{\sigma\}$. If $x_1 \in \Lambda$, then for any neighborhood $\mathcal{U}(X)$ of X , and any neighborhood V of x_2 , there is $Y \in \mathcal{U}(X)$ such that x_1 is still in the unstable manifold of σ and the positive orbit of x_1 will cross V with respect to Y .*

Proof. We prove this lemma by a standard application of the connecting lemma. By Lemma 3.2 we know that there is a point $z \in (W^s(\sigma) \setminus \{\sigma\}) \cap \Lambda$. Then we have two triple of $\rho > 1$, $L > 1$ and δ_0 with the properties stated as in Lemma 3.3 with respect to the point x_1 and z and the neighborhood $\mathcal{U}(X)$ of X . By taking the larger ρ, L , and smaller δ_0 , we get a triple, still denoted by ρ, L and δ_0 , works both for x_1 and z .

Now we can take $\delta > 0$ small enough such that the two tubes $\Delta_1 = B_\delta(X^{[0,L]}(x_1))$ and $\Delta_2 = B_\delta(X^{[-L,0]}(z))$ are disjoint. For any neighborhood V of x_2 and any neighborhood V' of z , by the inclination lemma we know that there are a point $y \in V$ and $T > 0$ such that $X^{-T}(y) \in V'$. If $\delta > 0$ is choosing small enough, we can take y and T such that $X^{[-T,0]}(y)$ does not touch Δ_1 .

Since Λ is transitive, we can find a point $x \in \Lambda$ such that $\Lambda = \omega(x)$. Then we can find $t_1 < t_2$ such that $X^{t_1}(x) \in B_{\delta/\rho}(x_1)$ and $X^{t_2}(x) \in B_{\delta/\rho}(z)$ and a point $y \in V$ with $X^{-T}(y) \in B_{\delta/\rho}(z)$. Then apply Lemma 3.3, we can find a vector field $Y \in \mathcal{U}(X)$ differs from X at tubes Δ_1 and Δ_2 such that the negative orbit of x_1 is not changed and y is contained in the positive orbit of x_1 . It is easy to see that Y satisfies the request of lemma. \square

Lemma 3.7 *Let $\mathcal{G}_2 \subset \mathfrak{X}^1(M)$ be the residual set chosen as in Lemma 3.5. Then for any $X \in \mathcal{G}_2$ and any isolated nontrivial transitive set Λ of X , if there is a singularity $\sigma \in \Lambda$ with $\text{index}(\sigma) = 2$, then we have $\overline{W^u(\sigma)} \subset \Lambda$.*

Proof. Let $\mathcal{O} = \{O_1, O_2, \dots, O_n, \dots\}$ be a countable basis of M . Recall that for each $m, k \in \mathbb{N}$, we take

$$\mathcal{H}_{m,k} = \{X \in \mathfrak{X}^1(M) : \text{there is a } C^1 \text{ neighborhood } \mathcal{U}(X) \text{ of } X$$

such that for any $Y \in \mathcal{U}(X)$, Y has a singularity $\sigma \in O_m$ with

$$\text{index}(\sigma) = 2 \text{ such that } W^u(\sigma, Y) \cap O_k \neq \emptyset\}.$$

Then take $\mathcal{N}_{m,k} = \mathfrak{X}^1(M) \setminus \overline{\mathcal{H}_{m,k}}$ and

$$\mathcal{G}_2 = \bigcap_{m,k \in \mathbb{N}} (\mathcal{H}_{m,k} \cup \mathcal{N}_{m,k}).$$

We will see that this \mathcal{G}_2 satisfies the request of lemma.

Let $X \in \mathcal{G}_2$ and Λ be an isolated transitive set of X . Assume there is singularity $\sigma \in \Lambda$ with index 2. Let $\Gamma_1 = Orb(x_1)$ and $\Gamma_2 = Orb(x_2)$ be the two branches of $W^u(\sigma) \setminus \sigma$. By Lemma 3.2, we know that either x_1 or x_2 is contained in Λ . Without loss of generality, we assume that $x_1 \in \Lambda$. To prove $\overline{W^u(\sigma)} \subset \Lambda$, we just need to prove that x_2 is also contained in Λ . By the compactness of Λ , we just need to prove that for any neighborhood U of x_2 , one has $U \cap \Lambda \neq \emptyset$. For a given arbitrarily small neighborhood U of x , we can find k such that $O_k \subset U$. Let O_m be a isolated neighborhood of σ . Then we have

Claim $X \notin \mathcal{N}_{m,k}$

Proof of Claim : For any neighborhood $\mathcal{V}(X) \subset \mathcal{U}(X)$, by Lemma 3.6, there is $Y \in \mathcal{V}(X)$ such that Y has a singularity $\sigma \in O_m$ with $\text{index}(\sigma) = 2$ and $W^u(\sigma, Y) \cap O_k \neq \emptyset$. By the continuity of the unstable manifold we know that there is a C^1 neighborhood $\mathcal{U}(Y)$ of Y such that for any $Z \in \mathcal{U}(Y)$, $W^u(\sigma, Z) \cap O_k \neq \emptyset$. Thus we have $Y \in \mathcal{H}_{m,k}$. Hence $X \in \overline{\mathcal{H}_{m,k}}$. This ends the proof of claim.

Since $X \in \mathcal{G}_2$ and $X \notin \mathcal{N}_{m,k}$, we have $X \in \mathcal{H}_{m,k}$. Since σ is the only singularity of X in O_m , by the definition of $\mathcal{H}_{m,k}$ we can see that $W^u(\sigma) \cap O_k \neq \emptyset$. Hence for any neighborhood U of x_2 , there is a point contained in $W^u(\sigma)$. This ends the proof of Lemma 3.7. \square

The following lemma is collected from [4].

Lemma 3.8 [4, Proposition 4.1] *There is a residual set $\mathcal{G}_3 \subset \mathfrak{X}^1(M)$ such that for any $X \in \mathcal{G}_3$, $\overline{W^u(\sigma)}$ is Lyapunov stable for X and $\overline{W^s(\sigma)}$ is Lyapunov stable for $-X$ for all $\sigma \in \text{Sing}(X)$.*

Proposition 3.9 *There is a residual set $\mathcal{S} \subset \mathfrak{X}^1(M)$ such that for any $X \in \mathcal{S}$, and any isolated nontrivial transitive set Λ of X , if there is a singularity $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 2$ then Λ is Lyapunov stable for X . Symmetrically, if there is $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 1$ then Λ is Lyapunov stable for $-X$.*

Proof. Let $X \in \mathcal{S} = \mathcal{G}_2 \cap \mathcal{G}_3$ and Λ be an isolated transitive set of X . Suppose that $\sigma \in \Lambda \cap \text{Sing}(X)$ with $\text{index}(\sigma) = 2$. Then by Proposition 3.5 and Lemma 3.7, we have $\overline{W^u(\sigma)} = \Lambda$. By Lemma 3.8, Λ is Lyapunov stable for X . \square

A point $\sigma \in \text{Sing}(X)$ of X is called *Lorenz-like* if $DX(\sigma)$ has three real eigenvalues $\lambda_1, \lambda_2, \lambda_3$ such that $\lambda_2 < \lambda_3 < 0 < -\lambda_3 < \lambda_1$. Let $\sigma \in \text{Sing}(X)$ be a Lorenz-like singularity, then we use $E_\sigma^{ss}, E_\sigma^{cs}, E_\sigma^u$ to denote the eigenspaces of $DX(\sigma)$ corresponding the eigenspaces $\lambda_2, \lambda_3, \lambda_1$ respectively. Denoted by $W_X^{ss}(\sigma)$ the one-dimensional invariant manifold of X associated to the eigenvalue λ_2 . We have the following lemma was proved in [13].

Lemma 3.10 [13, Lemma A. 4] *There is a residual set $\mathcal{G}_4 \subset \mathfrak{X}^1(M)$ such that for any $X \in \mathcal{R}$, if Λ is a Lyapunov stable nontrivial transitive set of X , then every singularity $\sigma \in \Lambda$ is Lorenz-like and one has $W_X^{ss}(\sigma) \cap \Lambda = \{\sigma\}$.*

Here is the main conclusion in this section.

Proposition 3.11 *There is a residual set $\mathcal{T} \subset \mathfrak{X}^1(M)$ with the following properties. Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . If there is a singularity with index 2, then for all singularity $\sigma \in \Lambda$, one has (1) $\text{index}(\sigma) = 2$, (2) σ is Lorenz-like, and (3) $W_X^{ss}(\sigma) \cap \Lambda = \{\sigma\}$. Symmetrically, if there is singularity with index 1 then for all singularity $\sigma \in \Lambda$, one has (1) $\text{index}(\sigma) = 1$, (2) σ is Lorenz-like for $-X$, and (3) $W_X^{uu}(\sigma) \cap \Lambda = \{\sigma\}$.*

Proof. Let $X \in \mathcal{T} = \mathcal{S} \cap \mathcal{G}_4$ and Λ be an isolated transitive set of X . Suppose that there is $\eta \in \Lambda \cap \text{Sing}(X)$ such that $\text{index}(\eta) = 2$. By Proposition 3.9, Λ is Lyapunov stable for X . On the other hand, since $X \in \mathcal{G}_4$, according to Lemma 3.11, σ is Lorenz-like, and $W_X^{ss}(\sigma) \cap \Lambda = \{\sigma\}$. We directly obtained $\text{index}(\sigma) = 2$, for all $\sigma \in \Lambda \cap \text{Sing}(X)$. \square

4 Proof of Theorem A

To prove Theorem A, we prepare two techniques here. One is the extended linear Poincaré flow given by Li, Gan and Wen[10], and another one is the ergodic closing lemma given by Mañé[11, 17].

Firstly we recall the notion of linear Poincaré flow firstly given by Liao [7, 8]. For any regular point $x \in M \setminus \text{Sing}(X)$, we can put a normal space

$$N_x = \{v \in T_x M : v \perp X(x)\}.$$

Then we have a normal bundle

$$N = N(X) = \bigcup_{x \in M \setminus \text{Sing}(X)} N_x.$$

Denote by π_x the orthogonal projection from $T_x M$ to N_x for any $x \in M \setminus \text{Sing}(X)$. From the tangent flow, we can define the *linear Poincaré flow*

$$P_t^X : N(X) \rightarrow N(X)$$

$$P_t^X(v) = \pi_{X^t(x)}(DX^t(v)), \text{ for all } v \in N_x, \text{ and } x \in M \setminus \text{Sing}(X).$$

Note that the linear Poincaré flow is defined on the normal bundle over a non compact set. We consider a compactification for P_t^X as following.

Let

$$G^1 = \{L : L \text{ is a one dimensional subspaces in } T_x M, x \in M\}$$

be the Grassmannian manifold of M . Then for any $L \in G^1$, assuming $L \subset T_x M$ for some $x \in M$, we can define a normal space associated to L as in following,

$$N_L = \{v \in T_x M : v \perp L\}.$$

Now can take a normal bundle

$$N = N_{G^1} = \bigcup_{L \in G^1} N_L.$$

Note that G^1 is a compact manifold, so N_{G^1} is a bundle over a compact space.

For any $L \in G^1$ contained in $T_x M$, denoted by π_L the orthogonal projection from $T_x M$ to N_L along L . Let X be a C^1 vector field. Similar to the linear Poincaré flow, we can define the *extended linear Poincaré flow*

$$\begin{aligned} \tilde{P}_t^X : N_{G^1} &\rightarrow N_{G^1} \\ \tilde{P}_t^X(v) &= \pi_{DX^t(L)}(DX^t(v)), \end{aligned}$$

for all $L \in G^1$ and $v \in N_L$.

One can check that for any $x \in M \setminus \text{Sing}(X)$, then we have $N_x = N_{\langle X(x) \rangle}$ and $P_t^X|_{N_x} = \tilde{P}_t^X|_{N_{\langle X(x) \rangle}}$. Here, \tilde{P}_t^X is said to be the *extended linear Poincaré flow*.

For any compact invariant set Λ of the vector fields X , we use $\tilde{\Lambda}$ to denote the closure of

$$\{\langle X(x) \rangle : x \in \Lambda \setminus \text{Sing}(X)\}$$

in the space of G^1 . Let $\sigma \in \Lambda$ be a singularity, denote by

$$\tilde{\Lambda}_\sigma = \{L \in \tilde{\Lambda} : L \subset T_\sigma M\}.$$

From the facts we got from Proposition 3.11, we have the following characterization of $\tilde{\Lambda}_\sigma$.

Lemma 4.1 *Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . Suppose there is a singularity with index 2. Then for all singularity $\sigma \in \Lambda$, we have $L \subset E_\sigma^{cs} \oplus E_\sigma^u$ for all $L \in \tilde{\Lambda}_\sigma$.*

Proof. Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . Suppose on the contrary, that is, there is $L \in \tilde{\Lambda}_\sigma$ such that L is not a subspace in $E_\sigma^{cs} \oplus E_\sigma^u$. Note that $DX^t(L)$ is contained in $\tilde{\Lambda}_\sigma$ for all $t \in \mathbb{R}$ and $\tilde{\Lambda}_\sigma$ is a closed set. By taking a limit line of $DX^t(L)$ as $t \rightarrow -\infty$, we know that there is $L \in \tilde{\Lambda}_\sigma$ such that $L \subset E_\sigma^{ss}$. From now on, we assume that $L \in \tilde{\Lambda}$ and $L \subset E_\sigma^{ss}$. By the definition of $\tilde{\Lambda}$, we know that there exist $x_n \in \Lambda \setminus \text{Sing}(X)$ such that $\langle X(x_n) \rangle \rightarrow L \subset E_\sigma^{ss}$.

For the simplicity of notations, we assume everything happens in a local chart containing σ . For any $0 < \eta \leq 1$, denote by $E_\sigma^{cu} = E_\sigma^{cs} \oplus E_\sigma^u$ and

$$C_\eta^{cu}(\sigma) = \{v = v^{ss} + v^{cu} \in T_\sigma M : |v^{ss}| < \eta|v^{cu}|, v^{ss} \in E_\sigma^{ss}, v^{cu} \in E_\sigma^{cu}\}$$

the cu -cone at the singularity σ . These cones can be parallel translated to x who is close to σ . Since $E_\sigma^{ss} \oplus E_\sigma^{cu}$ is a dominated splitting for the tangent flow DX^t , there are two constants $T > 0$ and $0 < \lambda < 1$ such that

$$DX^t(C_1^{cu}(\sigma)) \subset C_\lambda^{cu}(\sigma),$$

for any $t \in [T, 2T]$. By the continuous property of the cone to a cone field in a small neighborhood U_σ of σ , for any $t \in [T, 2T]$, $X^{[0,t]}(x) \subset U_\sigma$ then we have $DX^t(C_1^{cu}(x)) \subset C_1^{cu}(X^t(x))$.

Now let $t_n = \sup\{t > 0 : X^{[-t,0]}(x_n) \subset U_\sigma\}$. We know that $t_n \rightarrow +\infty$ as $n \rightarrow \infty$ because $x_n \rightarrow \sigma$ as $n \rightarrow \infty$. Denote by $y_n = X^{-t_n}(x_n)$. Then we can take $q = \lim_{n \rightarrow \infty} y_n \in \partial U_\sigma$ by taking the subsequence if necessary. We know that for $t > 0$, $X^t(q) \in U_\sigma$ and so, $q \in W^s(\sigma)$. Since $y_n \in \Lambda$ we know $q \in \Lambda$. If $q \in W^{ss}(\sigma) \cap \Lambda$, because we have already $q \in \partial U_\sigma$, hence $q \neq \sigma$, then from the fact that $X \in \mathcal{T}_1$ and Λ is an isolated nontrivial transitive set, this is a contradiction by Proposition 3.11. Now we assume that $q \in W^s(\sigma) \setminus W^{ss}(\sigma)$. We have $\langle X(X^t(q)) \rangle \rightarrow E_\sigma^{cs}$ as $t \rightarrow +\infty$. Thus there is $T_1 > 0$ big enough such that $X(X^{T_1}(q)) \in C_1^{cu}(X^{T_1}(q))$. For n big enough we have $X(X^{T_1}(y_n)) \in C_1^{cu}(X^{T_1}(y_n))$. Since $t_n \rightarrow \infty$, we assume that $t_n - T_1 > T$. Since $X^{[T_1, t_n]}(y_n) \subset U_\sigma$, we know that

$$\begin{aligned} X(x_n) &= X(X^{t_n}(y_n)) = DX^{t_n - T_1}(X(X^{T_1}(y_n))) \\ &\in DX^{t_n - T_1}(C_1^{cu}(X^{T_1}(y_n))) \\ &\subset C_1^{cu}(X^{t_n}(y_n)) = C_1^{cu}(x_n). \end{aligned}$$

This is a contradiction with the assumption $\langle X(x_n) \rangle \rightarrow L \subset E_\sigma^{ss}$. □

It is proved in section 2 that generically, if Λ is an isolated transitive set, then it is locally star. By some well know results from the proof of stability conjecture, we have the following proposition.

Proposition 4.2 [8, 11] *Let Λ be a locally star set for $X \in \mathfrak{X}^1(M)$ and let $\mathcal{U}(X), U$ be the neighborhoods in the definition of local star. Then there are constants $0 < \lambda_0 < 1, T_0 > 0$ such that for any $Y \in \mathcal{U}(X)$ and any $p \in \Lambda_Y(U) \cap \text{Per}(Y)$, the following properties hold:*

(a) $\Delta^s \oplus \Delta^u$ is a dominated splitting with respect to the linear Poincaré flow. Precisely, for any $t \geq T_0$ and any $x \in \text{Orb}(p)$,

$$\|P_t^Y|_{\Delta^s(x)}\| \cdot \|P_{-t}^Y|_{\Delta^u(Y^t(x))}\| \leq e^{-2\lambda_0 t};$$

(b) if τ is the period of p and m is any positive integer, and if $0 = t_0 < t_1 < \dots < t_k = m\tau$ is any partition of the time interval $[0, m\tau]$ with $t_{i+1} - t_i \geq T_0$, then

$$\frac{1}{m\tau} \sum_{i=0}^{k-1} \log \|P_{t_{i+1}-t_i}^Y|_{\Delta^s(Y^{t_i}(p))}\| < -\lambda_0,$$

and

$$\frac{1}{m\tau} \sum_{i=0}^{k-1} \log \|P_{-(t_{i+1}-t_i)}^Y|_{\Delta^u(Y^{t_{i+1}}(p))}\| < -\lambda_0,$$

where $\Delta^s \oplus \Delta^u$ is the hyperbolic splitting with respect to $P_\tau^X|_{N_{\text{Orb}(p)}}$.

Now we assume that Λ is an isolated transitive set of a C^1 -generic vector field X . By the closing lemma we know that for any $x \in \Lambda \setminus \text{Sing}(X)$, one can find a sequence of periodic points p_n of X such that $p_n \rightarrow x$ as $n \rightarrow \infty$. Consequently, for any $L \in \tilde{\Lambda}$, we can find a sequence of periodic points p_n of X , such that L is the limit of $\langle X(p_n) \rangle$. Since Λ is locally star, from item (a) of Proposition 4.2 we can see that for any $L \in \tilde{\Lambda}$, we can get two one dimensional subspaces $\Delta^1(L) = \lim_{n \rightarrow \infty} \Delta^s(p_n)$ and $\Delta^2(L) = \lim_{n \rightarrow \infty} \Delta^u(p_n)$ with the property: for any $t \geq T_0$,

$$\|\tilde{P}_t^Y|_{\Delta^1(L)}\| \cdot \|\tilde{P}_{-t}^Y|_{\Delta^2(DX^t(L))}\| \leq e^{-2\lambda_0 t}.$$

This implies that there is a dominated splitting $N_{\tilde{\Lambda}} = \Delta^1 \oplus \Delta^2$ for the extended linear Poincaré flow \tilde{P}_t^X . For any $x \in \Lambda \setminus \text{Sing}(X)$, we can put $\Delta^i(x) = \Delta^i(\langle X(x) \rangle)$ for $i = 1, 2$, then we can get a dominated splitting $N_{\Lambda \setminus \text{Sing}(X)} = \Delta^1 \oplus \Delta^2$ for the linear Poincaré flow P_t^X .

If $X \in \mathcal{T}$ and Λ be an isolated transitive set of X , then we have only finitely many singularity in Λ . Without loss of generality, after a change of equivalent Riemmanian structure, we can assume that for any $\sigma \in \Lambda$ with index 2, the subspaces $E_\sigma^{ss}, E_\sigma^{cs}, E_\sigma^u$ are mutually orthogonal. From Lemma 4.1 we know that every $L \in \tilde{\Lambda}_\sigma$ is orthogonal to E_σ^{ss} , this fact derives the following lemma.

Lemma 4.3 *Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . Suppose there is a singularity with index 2. Then for all singularity $\sigma \in \Lambda$ with mutually orthogonal $E_\sigma^{ss}, E_\sigma^{cs}, E_\sigma^u$, we have $\Delta^1(L) = E_\sigma^{ss}$ and $\tilde{P}_S^X|_{\Delta^1(L)} = DX^S|_{E_\sigma^{ss}}$ for any $L \in \tilde{\Lambda}_\sigma$.*

Proof. We denote by $E_\sigma^{cu} \triangleq E_\sigma^{cs} \oplus E_\sigma^u$, for any given singularity $\sigma \in \Lambda$. For any $L \in \tilde{\Lambda}_\sigma$, we set $N^1(L) = E_\sigma^{ss}$ and $N^2(L) = E_\sigma^{cu} \cap N_L$. By the fact that L is orthogonal to E_σ^{ss} we know that $N^1(L) \subset N_L$ for any $L \in \tilde{\Lambda}_\sigma$. Now we have two subbuddles

$$N_{\tilde{\Lambda}_\sigma}^1 = \bigcup_{L \in \tilde{\Lambda}_\sigma} N^1(L), \quad N_{\tilde{\Lambda}_\sigma}^2 = \bigcup_{L \in \tilde{\Lambda}_\sigma} N^2(L).$$

These two subbuddles are \tilde{P}_t^X -invariant by the fact that $L \subset E_\sigma^{cu}$ for any $L \in \tilde{\Lambda}_\sigma$ and both E_σ^{ss} and E_σ^{cu} are DX^t -invariant.

Since $E_\sigma^{ss} \oplus E_\sigma^{cu}$ is a dominated splitting for DX^t , we know that there are constants $C > 1, \lambda > 0$ such that

$$\frac{\|DX^{-t}(u)\|}{\|DX^{-t}(v)\|} \leq Ce^{-\lambda t}$$

for any unit vectors $u \in E_\sigma^{cu}$ and $v \in E_\sigma^{ss}$ and any $t > 0$. Then for any $L \in \tilde{\Lambda}_\sigma$ and any unit vectors $u \in N^2(L), v \in N^1(L)$, we have

$$\frac{\|\tilde{P}_{-t}^X(u)\|}{\|\tilde{P}_{-t}^X(v)\|} \leq \frac{\|DX^{-t}(u)\|}{\|DX^{-t}(v)\|} \leq Ce^{-\lambda t}.$$

This says that $N_{\tilde{\Lambda}_\sigma} = N_{\tilde{\Lambda}_\sigma}^1 \oplus N_{\tilde{\Lambda}_\sigma}^2$ is a dominated splitting on $\tilde{\Lambda}_\sigma$ with respect to the extended linear Poincaré flow \tilde{P}_t^X . By the uniqueness of dominated splitting we know that $N_{\tilde{\Lambda}_\sigma}^1 = \Delta_{\tilde{\Lambda}_\sigma}^1$. Thus we have $\Delta^1(L) = E_\sigma^{ss}$ for all $L \in \tilde{\Lambda}_\sigma$. By the definition of extended linear Poincaré flow, we directly have the fact that $\tilde{P}_S^X|_{\Delta^1(L)} = DX^S|_{E_\sigma^{ss}}$ for all $L \in \tilde{\Lambda}_\sigma$. \square

Now let us recall the ergodic closing lemma. A point $x \in M \setminus \text{Sing}(X)$ is called a *well closable point* of X if for any C^1 neighborhood $\mathcal{U}(X)$ of X and any $\delta > 0$, there are $Y \in \mathcal{U}(X), z \in M, \tau > 0$ and $T > 0$ such that the following conditions are hold:

- (a) $Y^\tau(z) = z$,
- (b) $d(X^t(x), Y^t(z)) < \delta$ for any $0 \leq t \leq \tau$, and
- (c) $X = Y$ on $M \setminus B(X^{[-T, 0]}(x), \delta)$.

Denote by $\Sigma(X)$ the set of all well closable points of X . Here we will use the flow version of the ergodic closing lemma which was proved in [17].

Lemma 4.4 [17] *For any $X \in \mathfrak{X}^1(M)$, $\mu(\Sigma(X) \cup \text{Sing}(X)) = 1$ for every $T > 0$ and every X^T -invariant Borel probability measure μ .*

Assume $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . From Proposition 4.2 we have already known that there is a dominated splitting $N_{\Lambda \setminus \text{Sing}(X)} = \Delta^1 \oplus \Delta^2$ with $\dim(\Delta^1) = \dim(\Delta^2) = 1$ with respect to the linear Poincaré flow P_t^X . By applying the ergodic closing lemma, we have the following lemma.

Lemma 4.5 *Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . Suppose there is a singularity with index 2. Then there are constant $C > 1$ and $\lambda > 0$ such that*

$$\|DX^t|_{\langle X(x) \rangle}\|^{-1} \cdot \|P_t^X|_{\Delta^1(x)}\| < Ce^{-\lambda t},$$

$$\|DX^{-t}|_{\langle X(x) \rangle}\| \cdot \|P_{-t}^X|_{\Delta^2(x)}\| < Ce^{-\lambda t}$$

for all $x \in \Lambda \setminus \text{Sing}(X)$ and $t \geq 0$.

Proof. Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . Then there is a \tilde{P}_t^X invariant splitting $N_{\tilde{\Lambda}} = \Delta^1 \oplus \Delta^2$ with constant $T_0 > 0$ and $\lambda_0 > 0$ such that the followings are satisfied:

- (1) if $L = \langle X(x) \rangle$ for some $x \in \Lambda \setminus \text{Sing}(X)$, then $\Delta^i(\langle X(x) \rangle) = \Delta^i(x)$ for $i = 1, 2$,
- (2) $\|\tilde{P}_t^Y|_{\Delta^1(L)}\| \cdot \|\tilde{P}_{-t}^Y|_{\Delta^2(DX^t(L))}\| \leq e^{-2\lambda_0 t}$ for any $t > T_0$, and
- (3) $L \in \tilde{\Lambda}$.

To prove the lemma, we just need to prove that there is $C > 1$ and $\lambda > 0$ such that for any $L \in \tilde{\Lambda}$ and any $t > 0$, we have

$$\|DX^t|_L\|^{-1} \cdot \|\tilde{P}_t^X|_{\Delta^1(L)}\| < Ce^{-\lambda t},$$

$$\|DX^{-t}|_L\| \cdot \|\tilde{P}_{-t}^X|_{\Delta^2(L)}\| < Ce^{-\lambda t}.$$

Since $\tilde{\Lambda}$ is compact, we just need to show that for any $L \in \tilde{\Lambda}$, there is a $T > 0$ such that

$$\log \|\tilde{P}_T^X|_{\Delta^1(L)}\| - \log \|DX^T|_L\| < 0,$$

$$\log \|\tilde{P}_{-T}^X|_{\Delta^2(L)}\| + \log \|DX^{-T}|_L\| < 0.$$

Now let us prove these properties of $\Delta^1 \oplus \Delta^2$ by contradiction. Firstly we prove the first half part. Assume that for any $L \in \tilde{\Lambda}$ and any $t > 0$

$$\log \|\tilde{P}_t^X|_{\Delta^1(L)}\| - \log \|DX^t|_L\| \geq 0.$$

Similar to [12, Lemma I.5], by a typical application of Birkhoff ergodic theorem, for any $S > 0$ there is an ergodic DX^T -invariant measure $\tilde{\mu} \in \mathcal{M}(G^1)$ with $\text{supp}(\tilde{\mu}) \subset \tilde{\Lambda}$ such that

$$\int (\log \|\tilde{P}_S^X|_{\Delta^1(L)}\| - \log \|DX^S|_L\|) d\tilde{\mu}(L) \geq 0.$$

In the following, we will always choose S is big enough.

Claim *If S is big enough, then for any singularity $\sigma \in \Lambda \cap \text{Sing}(X)$, one has $\tilde{\mu}(\tilde{\Lambda}_\sigma) = 0$.*

Proof of Claim : According to Lemma 4.1, for every $L \in \tilde{\Lambda}_\sigma$, $L \subset E_\sigma^{cs} \oplus E_\sigma^u \triangleq E_\sigma^{cu}$. Without loss of generality, we assume that E_σ^{ss} is orthogonal to E_σ^{cu} . Then by Lemma 4.3 we have $\tilde{P}_S^X|_{\Delta^1(L)} = DX^S|_{E_\sigma^{ss}}$ for any $L \in \tilde{\Lambda}_\sigma$. Since E_σ^{ss} is dominated by E_σ^{cu} , we can take S big enough such that

$$\log \|\tilde{P}_S^X|_{\Delta^1(L)}\| - \log \|DX^S|_L\| < 0$$

for any $L \in \tilde{\Lambda}_\sigma$. If $\tilde{\mu}(\tilde{\Lambda}_\sigma) \neq 0$, then we have $\tilde{\mu}(\tilde{\Lambda}_\sigma) = 1$ by the invariant of $\tilde{\Lambda}_\sigma$ and the ergodicity of $\tilde{\mu}$, thus we have,

$$\int (\log \|\tilde{P}_S^X|_{\Delta^1(L)}\| - \log \|DX^S|_L\|) d\tilde{\mu}(L) < 0.$$

This is a contradiction. This ends the proof of claim. \square

In the following, we will take S is a multiple of T_0 which is big enough such that the above claim is satisfied. One can see S have also the properties of T_0 .

For any Borel set $A \subset \Lambda$, we denote by $\tilde{A} = \{L : L = \langle X(x) \rangle \text{ for some } x \in A\}$. Then we define $\mu(A) = \tilde{\mu}(\tilde{A})$. By the fact that $\tilde{\mu}(\tilde{\Lambda}_\sigma) = 0$ for any $\sigma \in \Lambda \cap \text{Sing}(X)$, we know that μ is an ergodic measure support in Λ with $\mu(\Lambda \setminus \text{Sing}(X)) = 1$. From the inequality

$$\int (\log \|\tilde{P}_S^X|_{\Delta^1(L)}\| - \log \|DX^S|_L\|) d\tilde{\mu}(L) \geq 0,$$

we have

$$\int_{\Lambda \setminus \text{Sing}(X)} (\log \|P_S^X|_{\Delta_x^1}\| - \log \|DX^S|_{\langle X(x) \rangle}\|) d\mu(x) \geq 0.$$

By Lemma 4.4,

$$\int_{\Lambda \cap \Sigma(X)} (\log \|P_S^X|_{\Delta^1(x)}\| - \log \|DX^S|_{\langle X(x) \rangle}\|) d\mu(x) \geq 0.$$

By the ergodic theorem of Birkhoff, there is a point $y \in \Lambda \cap \Sigma(X)$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{nS} \sum_{j=0}^{n-1} (\log \|P_S^X|_{\Delta^1(X^{jS}(y))}\| - \log \|DX^S|_{\langle X(X^{jS}(y)) \rangle}\|) \geq 0. \quad (1)$$

Claim y is not a periodic point of X .

Proof of Claim : By the fact that $\|DX^S|_{\langle X(x) \rangle}\| = \frac{|X(X^S(x))|}{|X(x)|}$, we have

$$\sum_{j=0}^{n-1} \log \|DX^S|_{\langle X(X^{jS}(y)) \rangle}\| = \sum_{j=0}^{n-1} \log \frac{|X(X^{j+1S}(y))|}{|X(X^{jS}(y))|} = \log |X(X^{nS}(y))| - \log |X(y)|.$$

If $y \in Per(X)$ then by Proposition 4.2, we have

$$\limsup_{n \rightarrow \infty} \frac{1}{nS} \sum_{j=0}^{n-1} \log \|P_S^X|_{\Delta_{X^j S(y)}^s}\| \leq -\lambda_0.$$

Since $\sup |\log(X(x))|$ is bounded for $x \in Orb(y)$, we have

$$\limsup_{n \rightarrow \infty} \frac{1}{nS} \left(\sum_{j=0}^{n-1} \log \|P_S^X|_{\Delta_{X^j S(y)}^s}\| - \log |X(X^{nS}(y))| - \log |X(y)| \right) \leq -\lambda.$$

This is contradiction by (1). Thus y is not periodic. \square

Since y is a well closable point, for any $n > 0$, there are $X_n \in \mathfrak{X}^1(M)$, $z_n \in M$, and $\tau_n > 0$ such that

- (i) $Y_n^{\tau_n}(z_n) = z_n$ and τ_n is the prime period of z_n ,
- (ii) $d(X^t(y), Y_n^t(z_n)) \leq 1/n$, for any $0 \leq t \leq \tau_n$, and
- (iii) $\|Y_n - X\| \leq 1/n$.

Since y is not a periodic point, we have $\tau_n \rightarrow +\infty$ as $n \rightarrow \infty$. We also have the following uniformly continuity for $P_t^Y|_{\Delta^1}$.

Claim For any $\epsilon > 0$ there is $\delta > 0$ and a C^1 neighborhood $\mathcal{U}(X)$ of X such that for any $x, y \in M$, if (i) $x \in \Lambda \setminus Sing(X)$, (ii) there is $Y \in \mathcal{U}(X)$ such that $y \in Per(Y)$, $Orb(y) \subset U$, and $d(x, y) < \delta$, then

$$|\log \|P_t^X|_{\Delta^1(x)}\| - \log \|P_t^Y|_{\Delta^s(y)}\|| < \epsilon, \quad (2)$$

for any $t \in [0, 2S]$. Here $\Delta^s(y)$ denotes the stable subspace of y with respect to the vector field Y .

Proof of Claim : We prove this by deriving a contradiction. Assume the contrary. Then there is $\eta > 0$ such that for any $n > 0$ there exists $t_n \in [0, 2S]$, $X_n \rightarrow X$ and two sequences $\{x_n\}, \{y_n\}$ such that (i) $x_n \in \Lambda \setminus Sing(X)$, (ii) $y_n \in Per(X_n)$ and $Orb(y_n) \subset U$, (iii) $d(x_n, y_n) < 1/n$, and

$$|\log \|P_{t_n}^X|_{\Delta^1(x_n)}\| - \log \|P_{t_n}^{X_n}|_{\Delta^s(y_n)}\|| \geq \eta,$$

Since $[0, 2S]$ and Λ are compact, we can take sequences $\{t_n\} \subset [0, 2S]$ and $\{x_n\} \subset \Lambda$ (take subsequences if necessary) such that $t_n \rightarrow t_0$ and $x_n \rightarrow x_0$. Then we have $y_n \rightarrow x_0$ by the above item (iii).

If $x_0 \notin Sing(X)$ then by the continuity of dominated splitting, we know $\Delta^1(x_n) \rightarrow \Delta^1(x_0)$ and $\Delta^s(y_n) \rightarrow \Delta^1(x_0)$ as $n \rightarrow \infty$, then we have

$$|\log \|P_{t_0}^X|_{\Delta^1(x_0)}\| - \log \|P_{t_0}^{X_n}|_{\Delta^1(x_0)}\|| \geq \eta.$$

This is a contradiction.

If $x_0 \in \text{Sing}(X)$ then we can take sequence $\{\langle X(x_n) \rangle\}, \{\langle X_n(y_n) \rangle\}$ (take subsequences if necessary) such that $\langle X(x_n) \rangle \rightarrow L \in \tilde{\Lambda}_{x_0}$ and $\langle X_n(y_n) \rangle \rightarrow L_1 \in \tilde{\Lambda}_{x_0}$. Since both $L, L_1 \in \tilde{\Lambda}_{x_0}$, we have $\tilde{P}_t^X|_{\Delta^1(L)} = \tilde{P}_t^X|_{\Delta^1(L_1)} = DX^t|_{E_{x_0}^{ss}}$ by Lemma 4.3. But on the other hand, we have

$$|\log \|\tilde{P}_t^X|_{\Delta^1(L)}\| - \log \|\tilde{P}_t^X|_{\Delta^1(L_1)}\|| \geq \eta.$$

This is also a contradiction. This ends the proof of Claim. \square

By (2), there is n_0 such that for any $k > n_0, t \in [0, 2S]$ and $t_0 \in [0, \tau_n]$, one has

$$|\log \|P_t^X|_{\Delta_{X^{t_0}(y)}^1}\| - \log \|P_t^{X_n}|_{\Delta^s(X_n^{t_0}(z_n))}\|| < S\lambda_0/3, \quad (3)$$

where λ_0 as in Proposition 4.2. Let $\tau_n = m_n S + s_n$ ($m_n \in \mathbb{Z}$ and $s_n \in [0, S]$.) Then we consider the partition

$$0 = t_0 < t_1 = S < \cdots < t_{m_n-1} = (m_n - 1)S < t_{m_n} = \tau_n,$$

According to Proposition 4.2, we know

$$\sum_{j=0}^{m_n-2} \log \|P_S^{X_n}|_{\Delta^s(X_n^{jS}(z_n))}\| + \log \|P_{S+s_n}^{X_n}|_{\Delta^s(X_n^{(m_n-1)S}(z_n))}\| \leq -\tau_n \lambda_0.$$

Then by (3) we have

$$\begin{aligned} & \sum_{j=0}^{m_n-2} \log \|P_S^X|_{\Delta^1(X^{jS}(y))}\| + \log \|P_{S+s_n}^X|_{\Delta^1(X^{(m_n-1)S}(y))}\| \\ & \leq m_n S \lambda_0 / 3 - \tau_n \lambda_0 = -2m_n S \lambda_0 / 3 - s_n \lambda_0 \leq -2m_n S \lambda_0 / 3. \end{aligned}$$

For sufficiently small $r > 0$, let $B_r(y)$ be a neighborhood of $X^{[-2S, 0]}(y)$ such that $B_r(y) \cap \text{Sing}(X) = \emptyset$.

Denote by $C = \sup\{|\log |X(x)|| : x \in B_r(y)\} + \sup\{|\log \|P_t^X|_{\Delta^s(x)}\|| : x \in B_r(y), t \in [0, 2S]\} < \infty$.

Since $d(y, z_n) < 1/n$ and $d(X^{\tau_n}(y), z_n) = d(X^{\tau_n}(y), X^{\tau_n}(z_n)) < 1/n$, we know $d(X^{\tau_n}(y), y) < 2/n$. Thus there is $n_1 > n_0$ such that for any $n > n_1$ and $t \in [0, 2S]$ we have $X^{\tau_n-t}(y) \in B_r(y)$. Since $\tau_n - (m_n - 1)S = S + s_n < 2S$, we know

$$|\log |X(X^{(m_n-1)S}(y))|| + |\log \|P_{S+s_n}^X|_{\Delta^s(P_{(m_n-1)S}^X(y)}\|| \leq C. \quad (4)$$

By (1) and $m_n \rightarrow +\infty$ as $n \rightarrow +\infty$, there is $n_2 \geq n_1$ such that for any $n > n_2$

$$\sum_{j=0}^{m_n-2} \log \|P_S^X|_{\Delta^1(X^{jS}(y))}\| - (\log |X(X^{(m_n-1)S}(y))| - \log |X(y)|) \geq -(m_n - 1)S\lambda_0/3.$$

Then by

$$\sum_{j=0}^{m_n-2} \log \|P_S^X|_{\Delta^s(X^j S(y))}\| + \log \|P_{S+s_n}^X|_{\Delta^s(X^{(m_n-1)S}(y)}\| \leq -2m_n S \lambda_0 / 3,$$

and (4), we have

$$-(m_n - 1)S \lambda_0 / 3 \leq -2m_n S \lambda_0 / 3 + C + \log |X(y)|.$$

If n is big enough then it is not happen, and so, it is a contradiction. This proves that for any $L \in \tilde{\Lambda}$, there is a $T > 0$ such that

$$\log \|\tilde{P}_T^X|_{\Delta^1(L)}\| - \log \|DX^T|_L\| < 0.$$

And then by the compactness of $\tilde{\Lambda}$, we can find $C > 1$ and $\lambda > 0$ such that for any $L \in \tilde{\Lambda}$ and any $t > 0$, we have

$$\|DX^t|_L\|^{-1} \cdot \|\tilde{P}_t^X|_{\Delta^1(L)}\| < Ce^{-\lambda t}.$$

By a similar argument we can prove that for any $L \in \tilde{\Lambda}$, there is a $T > 0$ such that

$$\log \|\tilde{P}_{-T}^X|_{\Delta^2(L)}\| + \log \|DX^{-T}|_L\| < 0,$$

and then there exist $C > 1$ and $\lambda > 0$ such that for any $L \in \tilde{\Lambda}$ and any $t > 0$, we have

$$\|DX^{-t}|_L\| \cdot \|\tilde{P}_{-t}^X|_{\Delta^2(L)}\| < Ce^{-\lambda t}.$$

This ends the proof of the lemma. \square

Theorem A is a direct corollary of Lemma 4.5 and the following lemma in [18].

Lemma 4.6 [18, Theorem A] *Assume Λ is a non-trivial transitive set such that all singularity in Λ is hyperbolic. If there is a dominated splitting $N_{\Lambda \setminus \text{Sing}(X)} = \Delta^1 \oplus \Delta^2$ on $\Lambda \setminus \text{Sing}(X)$ with respect to P_t^X and there are constant $C > 1$ and $\lambda > 0$ such that*

$$\begin{aligned} \|DX^t|_{\langle X(x) \rangle}\|^{-1} \cdot \|P_t^X|_{\Delta^1(x)}\| &< Ce^{-\lambda t}, \\ \|DX^{-t}|_{\langle X(x) \rangle}\| \cdot \|P_{-t}^X|_{\Delta^2(x)}\| &< Ce^{-\lambda t} \end{aligned}$$

for all $x \in \Lambda \setminus \text{Sing}(X)$ and $t \geq 0$, then Λ is positively singular hyperbolic.

Proof of Theorem A. Let $X \in \mathcal{T}$ and Λ be an isolated transitive set of X . If there is singularity $\sigma \in \Lambda$ with index 2, then Λ is positively singular hyperbolic by Lemma 4.5 and Lemma 4.6. If there is a singularity $\sigma \in \Lambda$ with index 1, then by reversing the vector fields, we know that Λ is negatively singular hyperbolic. This ends of the proof of Theorem A. \square

Acknowledgement. The author would like thanks to Xio Wen for very helpful comments and discussion. This work is supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Science, ICT & Future Planning (No. 2017R1A2B4001892).

References

- [1] F. Abdenur, C. Bonatti, and S. Crovisier, *Global dominated splittings and the C^1 Newhouse phenomenon*. Proc. Amer. Math. Soc. **134** (2006), no. 8, 2229–2237.
- [2] C. Bonatti and L. Díaz, *Persistent nonhyperbolic transitive diffeomorphisms*, Ann. of Math. **143** (1996), 357–396.
- [3] C. Bonatti, L. Díaz and E. Pujal, *A C^1 -generic dichotomy for diffeomorphisms: Weak forms of hyperbolicity or infinitely many sinks or sources*, Ann. of Math., **158** (2003), 355–418
- [4] C. M. Carballo, C. A. Morales and M. J. Pacifico, *Maximal transitive sets with singularities for generic C^1 vector fields*, Bol. Soc. Bras. Mat. **31** (2000), 287–303.
- [5] L. Díaz, E. Pujals, and R. Ures, *Partial hyperbolicity and robust transitivity*, Acta Math. **183** (1999), 1–43.
- [6] J. Guckenheimer, *A strange, strange attractor*, The Hopf Bifurcation Theorem and its Applications, Springer-Verlag, New York (1976).
- [7] S. Liao, *Obstruction sets (I)*, Acta Math. Sinica **23** (1980), 411–453.
- [8] S. Liao, *Obstruction sets (II)*, Acta Sci. Natur. Univ. Pekinensis **2** (1981), 1–36.
- [9] E. N. Lorenz, *Deterministic nonperiodic flow*. J. Atmosph. Sci. **20** (1963), 130–141.
- [10] M. Li, S. Gan and L. Wen, *Robustly transitive singular sets via approach of extended linear Poincare flow*, Discrete Contin. Dyn. Syst., **13** (2005), 239–269.
- [11] R. Mañé, *An erodic closing lemma*, Ann. of Math. **116** (1982), 503–540.
- [12] R. Mañé, *A proof of the C^1 stability conjecture*, Publ. Math. IHES **66** (1988), 161–210.
- [13] C. A. Morales and M. J. Pacifico, *A dichotomy for three-dimensional vector fields*, Ergod. Th. & Dynam. Syst. **23** (2003), 1575–1600.
- [14] C. A. Morales, M. J. Pacifico and E. R. Pujals, *Robust transitive singular sets for 3-flows are partially hyperbolic attractors or repellers*, Ann. of Math. **160**(2004), 375–432.
- [15] J. Palis and de Melo, *Geometric theory of dynamical system: an introduction*, Springer-Verlag, Nwe-York (1982).
- [16] L. Wen, *Generic diffeomorphisms away from homoclinic tangencies and heterodimensional cycles*, Bull. Braz. Math. Soc. New Ser. **35**(2004), 419–452.
- [17] L. Wen, *On the C^1 stability conjecture for flows*, J. Differ. Equat., **129**(1996), 334–357.

- [18] X. Wen, L. Wen and D. Yang, *A characterization of singular hyperbolicity via the linear Poincaré flow*, in preprint.
- [19] L. Wen and Z. Xia, *C^1 -connecting lemma*, Trans. Amer. Math. Soc., **352**(2000), 5213–5230.
- [20] S. Zhu, S. Gan and L. Wen, *Indices of singularities of robustly transitive sets*, Discrete Contin. Dyn. Syst., **21** (2008), 945–957.