

Review Article

Biological Value of *Centaurea damascena*: Minireview

Mohammad Jaafreh^a, Haitham Qaralleh^{b*}, Muhamad O. Al-limoun^a

^aBiology Department, Mutah University, Mutah, Karak, 61710, Jordan; ^bDepartment of Medical Laboratory Sciences, Mutah University, Mutah, Karak, 61710, Jordan

*Corresponding Author: Haitham Qaralleh, ^bDepartment of Medical Laboratory Sciences, Mutah University, Mutah, Karak, 61710, Jordan

*Corresponding author: haitham@mutah.edu.jo

Received: 4-9-2019
Revised: 7-10-2019
Published: 19-10-2019

Keywords:

Centaurea damascena
Traditional value
Antimicrobial
Antioxidant
Chemical composition

Abstract: The family Asteraceae include large number of *Centaurea* species which have been applied in folk medicine. One of the family Asteraceae members is the *Centaurea damascena* which authentically been tested for its antibacterial activity. The aim of the study was to discuss antibacterial activities of essential oil composition and methanolic extract of the same plant aerial part leaves. Thirty-seven components were characterized with 86% of oxygenated terpenes. The composition was dominated by 11.45% Fokienol, 8.8% thymol, 8.21% Alpha Terpeneol, 7.24% Chrysanthemumic acid, 7.13% Terpinen-4-ol and 6.59% Borneol with a high degree of polymorphism in the occurrence of these compounds as compared with the different species of *centaurea*.. Free radical scavenging capacity of the *C. damascna* methanol extract was calculated by DPPH and FRAP test. DPPH radicals were scavenged with an IC50 value of 17.08 µg /ml. Antioxidant capacities obtained by the FRAP was 51.9 and expressed in mg Trolox g⁻¹ dry weight. The total phenolic compounds of the methanol extracts of aerial parts, as estimated by Folin–Ciocalteu reagent method, was about 460 mg GAE/ g. The phenolic contents in the extracts highly correlate with their antioxidant activity, (R² = 0.976) confirming that the antioxidant activity of this plant extracts is considerably phenolic contents-dependent.

Cite this article as Jaafreh, M., Qaralleh, H., Al-limoun, M.O. (2019). Biological value of *Centaurea damascena*: Minireview. . Journal of basic and applied Research in Biomedicine, 5(2): 99-106



This work is licensed under a Creative Commons Attribution 4.0 License. You are free to copy, distribute and perform the work. You must attribute the work in the manner specified by the author or licensor.

INTRODUCTION

Medicinal plants continue being an important resource worldwide, to withstand critical diseases. Overwhelming of the world's population (60-80%) still relies on classical medicines for the curing of current disease (WHO, 2002; Zhang, 2004; Qaralleh et al., 2019). Plants involve various compounds that have active biological substances (Mokbel and Hashinaga, 2006). For instance, phenolic compounds and essential oil play a crucial function as vigorous natural biological factors (Cutter, 2000; Hao et al., 1998; Puupponen-Pimia et al., 2001). The routine use of antibiotic has led to the development of one or more antibiotics resistant infectious bacteria (Sarker et al., 2005; Sufferidini et al., 2004). This issue has resulted on the failure of the treatments to numerous microbial causing infectious disease. Previous investigations pointed out to a number of medicinal plant extracts by constituting a group of potent natural antimicrobial agents (Rayne and Mazza, 2007). Jordan's flora includes more than 2500 wild plant species from 700 genera, and of these, there are around 100 endemic species, 250 rare species, and 125 quite rare species (Tellawi et al., 2001). Although, most of plant species have not been explored chemically or biologically, medicinal

plants have been extensively used in traditional medication and continue being a rich source of novel therapeutic agents (Cragg et al., 1997).

Plant family Asteraceae, consists of more than 1500 genera and approximately 2500 species worldwide (Wagstaff and Breitwieser, 2002). The genus *Centaurea* comprises more than 500 species that are common worldwide, particularly the Mediterranean and western Asia area (Mabberley, 1997). Some Asteraceae species have been used in many fields, including nutrition and medicinal industries (Tepe et al., 2006; Tekeli et al., 2011). Many species of *Centaurea* have long been utilized in conventional medication to treat different diseases, like hemorrhoids, abscess and the common cold (Kargioglu et al., 2008; Kargioglu et al., 2010; Sezik et al., 2001). Thus it is worthwhile to elaborate on the antibacterial activities of different species of *Centaurea* on the case by case basis. Though most of plant species have not been explored chemically or biologically, medicinal plants have been extensively used in traditional medication and continue being a rich source of novel therapeutic agents (Cragg et al., 1997; Al Fraijat et al., 2019). In the present study, we demonstrate that methanolic extracts and essential

oil of *C. damascene* (Khleifat et al., 2019; Jaafreh et al., 2019), which are used to treat gastritis, and for flavoring hot tea on a small scale in a town south of Jordan (Bsaira). The choice of this species was based on a review of the local folk literature which points out the therapeutic properties that the genus *Centaurea* possess (Alali et al., 2007) and on a preliminary screening study done in our laboratory investigating other medicinal plants for their low cytotoxicity (Qaralleh et al., 2009; Khleifat et al., 2006a; Khleifat et al., 2010; Zeidan et al., 2013; Majali et al., 2015; Althunibat et al., 2016; Al-Asoufi et al., 2017; Qaralleh et al., 2019; ALrawashdeh et al., 2019). The aim of this study was to determine the antibacterial and antioxidant activities as well as toxicity of *Centaurea damascene* that authentically been explored in this work as one of species of *Centaurea* growing in Jordan.

Chemical Composition of *C. damascena* Essential Oil

It has been reported that the yield of essential oil of *C. damascena* (table 1). 37 components were

characterized on the basis of a typical library search and literature data with selecting only the components showing matches $\geq 90\%$, which represented about 92.97% of the essential oils from *C. damascena* (Adams, 1995; Flamini et al., 2004; Skaltsa et al., 2003, 2000; Ertugrul et al., 2003; Yayli et al., 2009; Adams, 1995). The main contents of this essential oil consist of 11.45% Fokienol, 8.8% thymol, 8.21% alpha Terpineol, 7.24% Chrysanthemumic acid, 7.13% Terpinen-4-ol and 6.59% Borneol with a high degree of polymorphism in the occurrence of these compounds between the different species of *centaurea* (Fig. 1 and table 2). For example oxygenated monoterpenes was absent or rare in the *C. wagheritzii*, *C. tossiensis* and *C. luschaniana* (Köse et al., 2007). In parallel GC analysis of *C. damascena* showed that Fokienol represent 11.45% whereas *C. wagheritzii* showed only 0.50% Fokienol (Köse et al., 2007). However, there is no previous information of investigations on the same species to make a comparison of chemical composition of other regions to show if there is a variation of the major components.




Figure 1. Variation of main constituents (of > 2%) in the essential oil of *C. damascena*.

Table 1. Yield of ethanol crude plant extracts (%)

Plants	Dry weight (g)	Weight of ethanol extract (g)	Yield of ethanol extract (%)
<i>C.damascena</i> Bioss	50	5.72	11.44%

Table 2. The chemical class distribution of the essential oil components of *C. damascena*

Class compound	Concentration (%)	Number of compounds
Hydrocarbons	0.35	1
Monoterpenes hydrocarbons	0.46	1
Oxygenated monoterpenes	46.87	14
Sesquiterpene hydrocarbons	4.41	4
Oxygenated sesquiterpenes	40.89	17

Antimicrobial Activities of Essential Oil and Methanolic Extract of *C. damascena*

The regularly used method for evaluation of antibacterial efficiency of publically used medicinal plants can be diffusion as qualitative and dilution methods of the extracts and essential oils as quantitative analyses (Köse et al., 2008). The results show that plant extracts and essential oils have antibacterial activity, while plant extracts did not have that effect as essential oils do on all the tested bacteria. The results gotten from the disc diffusion tests showed that there has been an increasing action of essential oils on the inhibition of bacterial growth on concentration dependent manner. Among the tested microorganisms, *M. lutes* was the most susceptible microorganism against lowest concentration of essential oils. The antimicrobial activity of other *Centaurea* species was focused by many authors (Hajjeh et al., 1999; Güven et al., 2005; Sarker et al., 2005; Yayli et al., 2009; Cansaran et al., 2010 and Tekeli et al., 2011). The essential oils (10 µg/ml) produced the highest inhibition zone of 24 mm that was monitored against *M. lutes*. Oils showed antibacterial activity against seven out of ten tested bacteria. Only *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* were resistant to essential oils of *C. damascene*.

For a more reliable judgment of antimicrobial action (Boğa, et al., 2016), a broth dilution assay was carried out. The MIC of essential oils of *C. damascena* ranged from and 15 ± 1000 µg/mL. The order of MIC values from lowest to highest are *M. luteus* (15µg/ml), *B. subtilis* (50µg/ml), *S. aureus* (50µg/ml), *E. aerogenes* and *S. epidermidis*(123µg/ml), *Salmonella typhi*, *P. mirabilis* and *E. coli* (370µg/ml) and *P. aeruginosa* and *Klebsiella pneumoniae* (1mg/ml). Generally speaking all MIC values for ten bacteria used in this study were in conformity with the results of disc diffusion method. Yayli et al. (2009) obtained

a moderate antibacterial activity results on Gram-positive and Gram-negative by the essential oils from *C. appendicigera* and *C. helenioides*. The essential oil of *C. aladagensis* had an antibacterial effect versus seven human pathogenic microorganisms while the essential oils of *C. nicaeensis* and *C. parlatoris* showed low activity on fourteen selected microorganisms (Kose et al., 2007). The antimicrobial characteristic of medicinal plants is really hard to explain.

The methanol extracts of the plants were active against most bacteria being tested with range of inhibition zones (0-19 mm). Among the tested bacteria, *E. aerogenes* and *S. epidermidis* were the most susceptible bacteria against methanol extracts (19 mm). The extracts exhibited antimicrobial activity against *S. aureus*, *B. subtilis* and *M. lutes* based on zone of inhibition by 18, 17, and 17 mm, respectively. An important antimicrobial effect of five *Centaurea* species (*C. ptosomipappoides*, *C. odyssei*, *C. ptosomipappa*, *C. amonicola* and *C. kurdica*) was reported by Güven et al. (2005) on *S. aureus* and *B. cereus*. Conversely, extracts of *C. appendicigera* and *C. helenioides* did not display antimicrobial activity on *E. coli* (Buruk et al., 2006).

The chloroform and ethyl alcohol extracts of *C. cariensis* subsp. *Niveotomentosa* exhibited powerful antibacterial activities on various resistant bacteria, particularly *Staphylococcus* strains (Ugur et al., 2010). The diameter zone of inhibition for tested bacteria by cephexime antibiotic ranged between 9-36 mm. However, for the plant extracts, the average zone of inhibitions observed against these pathogens ranged from 17 to 19 mm. These values fall within the range sensitive and/or intermediate sensitive when compared with control antibiotic. Although, the low values recorded for the plant extracts may be attributed to the fact that the extracts being in crude form, contain very small

amounts of bioactive compounds (Olukoya et al., 1993).

For a more reliable estimation of antimicrobial activity, a broth dilution assay was carried out. The lowest MIC value was recorded by *M. lutes* (60 $\mu\text{g mL}^{-1}$) in conformity with the result of disc diffusion method. MIC values of *C. damascena* against *E. coli* and other Gram (-) bacteria were 1100 $\mu\text{g mL}^{-1}$ with an exception of *E. aerogenes* (123 $\mu\text{g mL}^{-1}$) which they are in accordance with results of inhibition zones. The results showed that methanol extracts of the *C. damascena* exerted good antibacterial activity on all the strains being tested at different concentrations.

It is worthy of note that MBC values obtained for the extracts against tested bacteria are higher than MIC, indicating that the extracts are bacteriostatics at lower concentrations and bactericidal at higher concentrations. It noteworthy to mention that even though the MBCs are sometimes 3-fold ($\leq 3300\mu\text{g/mL}$) of that the MICs ($\leq 1100\mu\text{g/mL}$), the 3.3 mg/mL concentration is still very low as compared with the MICs and MBCs of a lot of medicinal plants in the literature (Tekeli et al., 2011). This suggests that these plant extracts, when used traditionally as antimicrobials inhibit bacterial growth through killing the bacteria and since most of the traditional preparations lack specific concentrations, this may thus account for the use of large quantity of the extracts by traditional medical practitioners for the treatment of their patients. However, the antibacterial activity of most examined *Centaurea* species were shown to be weak or solvent-dependent (Sarker et al., 2005).

However, it has no data previously been reported to discuss any possible varying degree of antibacterial activity by whole *C. damascena* against any type of microorganisms. A study on the biological activities of 12 different species of *Centaurea* was conducted by Tekeli et al., (2011). *B. cereus* was only inhibited by two of the 12 studied *Centaurea* species with 4 mg/ml concentration., *C. calolepis* displayed an action on *Salmonella enteritidis*, while *C. urvillei subsp. urvillei* was active against *Escherichia coli* and *Staphylococcus aureus*. In further study on *C. cankiriense* (Cansaran et al., 2010), the MICs of *E. coli* (Abboud et al., 2010; Tarawneh et al., 2009) and *S. aureus* against ethylacetate and methanol extracts were being as 250 and 62.5 $\mu\text{g/ml}$, respectively

The antibacterial efficacy in the methanolic extract possibly be due to the presence of tannins, flavonoids, and terpenoids (Zablotowicz et al., 1996; Hernandez et al., 2000; Mamtha et al. 2004). These medically bioactive ingredients perform

antimicrobial activity through various mechanisms. Flavonoids, which have been found to be effective antimicrobial substances against a wide array of microorganisms in vitro, are known to be synthesized in response to microbial infection by plants. They have the ability to complex with extracellular and soluble proteins and to complex with bacterial cell walls (Cowan et al., 1999; Khleifat, et al., 2006b; Khleifat, 2006; Khleifat, et al., 2008; Khleifat, et al., 2015).

The saponins have the capacity to rise leakage of metabolites from the cell (Zablotowicz et al., 1996). Tannins cause cell wall synthesis inhibition by forming irreversible complexes with prolene rich protein (Mamtha et al. 2004). Terpenoids cause dissolution of the cell wall of microorganism by undermining the membranous tissue (Hernandez et al., 2000). In this study, *C. damascena* methanol extracts are lacking steroids and this probably why Gram-negative bacteria were less susceptible to *C. damascena* extract than the Gram-positive one. The steroids are known for their antibacterial activity particularly connected with membrane lipids and cause leakage from liposomes (Epand et al., 2007).

Evaluation of the synergistic effect of antibiotics and essential oil of *C. damascena* on mainly resistant bacteria

The oil alone from *C. damascena* in the concentration of 20 $\mu\text{g/mL}$ inhibited the bacterial growth of *P. aeruginosa* and *K. pneumonia*. However, a synergetic effect against *E. coli* and *K. pneumonia* was observed when 5 $\mu\text{g/mL}$ oil was combined with some of the antibiotics tested, although some of these antibiotics did not show any activity by themselves such as gentamicin for *E. coli* and vancomycin, ampicillin and chloramphenicol for *K. pneumonia*. Even though a synergistic effect for *P. aeruginosa* with different plant extracts was previously observed (Moon et al., 2011; Nascimento et al., 2000; Rodrigues et al., 2009; Al-Asoufi et al., 2017), no synergetic effect was observed when different antibiotics were combined with 5 $\mu\text{g/mL}$ oil to inhibit the growth of *P. aeruginosa*. Probably the concentration factor is critical for synergistic effect since the concentration of 20 $\mu\text{g/mL}$ oil inhibited the bacterial growth of *P. aeruginosa* (Abboud et al., 2009).

In some cases, the mode of action of combinations vary significantly from that of the same antibiotics acting alone. The activity of the antibiotic cefixime was increased by 26% against *K. pneumonia* after combination with the oil (29 mm) since single-handedly cefixime resulted in the inhibition zone of 23 mm, showing that this oil at 5 $\mu\text{g/mL}$ influences the activity of the antibiotic and may be used as a potential adjuvant in the antibiotic cure of pneumonia caused by *K. pneumoniae*.

Antioxidant Activity of of *C. damascena* Essential Oil

DPPH is normally employed as a reagent to estimate free radical scavenging capacity of antioxidants (Zengin et al., 2010). Free radical scavenging capacity of the *C. damascena* methanol extract was calculated by DPPH test. IC₅₀ value is expressed by which the functional concentration of 50% DPPH radicals were scavenged and was determined from the graph plotting inhibition percentage versus extract concentration. IC₅₀ of methanol extract was detected by using DPPH assay and expressed as µg/ml. IC₅₀ estimate is inversely correlated to antioxidant capacity of extracts. In the present study, *C. damascena* extract exerted better antioxidant efficiency than that of previously reported *Centaurea* species extract with an IC₅₀ value of 17.08 µg /ml. The extract exhibited concentration-dependent radical scavenging activity,

As far as we know from literature survey, the *Centaurea damascena* have shown better activities than other *Centaurea* species such as *C. patula*, *C. pulchella*, and *C. tchihatchfeii* (Zengin et al., 2010), *C. mucronifera* (Tepe et al., 2006), *C. huber-morathii* (Sarker et al., 2005) and *C. centaurium* (Conforti et al., 2008) by the same DPPH assay. Antioxidant capacities obtained by the FRAP was 51.9 and expressed in mg Trolox g⁻¹ dry weight. The two methods (FRAP and DPPH) approximately showed similar results for *C. damascena* methanol extract since the two methods, the assay results were generally similar for different plant models because these methods are based on electron transfer mechanism (OZGEN et al., 2006)

The total phenolic compounds of the methanol extracts of aerial parts, as estimated by Folin–Ciocalteu reagent method, was about 460 mg GAE/g. The highest level of phenolics was found in *C. damascena* as compared with previously reported other species of *Centaurea* (Sarker et al., 2005; Tepe et al., 2006; Karamenderes et al., 2007; Alali et al., 2007; Conforti et al., 2008; Dudonné et al., 2009; Zengin et al., 2010). The results of total phenolic compounds suggested that the phenolic contents contributed significantly to the antioxidant ability of the *C. damascena* extract. It has been reported in the literature that phenolic contents have strong antioxidant capacities and these phenolics have antioxidant activity mainly as a result of their redox properties which enable them to act as reducing agents, singlet oxygen quenchers,

hydrogen donors, and metal chelator (Baratto et al., 2003; Alali et al., 2007; Zengin et al., 2010). However, it was found that methanolic plant extracts are the most effective scavenger of DPPH radical as compared with other solvents (Miliauskasa et al., 2004). Hence, it was suggested that methanol is more efficient solvent for cell walls and seeds degradation, that have nonpolar nature causing the release of polyphenols from cells. It can be observed that the phenolic contents in the extracts highly correlate with their antioxidant activity, (R² = 0.97), confirming that phenolic compounds contribute significantly to the antioxidant activity of these plant extracts (Fig. 2). The enormous disparity in *Centaurea* species antioxidant activity could result from variations in total phenolic contents (Al-Mustafa, & Al-Thunibat, 2008). Such observation agreed with several previous findings (Sarker et al., 2005; Tepe et al., 2006; Guven et al., 2005; Karamenderes et al., 2007; Alali et al., 2007; Conforti et al., 2008; Zengin et al., 2010; Karahan et al., 2016).

Moreover, phenolic compounds have various reactions to Folin-Ciocalteu assay (Sun et al., 2002). The molar response of this method is roughly proportional to the number of phenolic hydroxyl groups in a given substrate, but the reducing capacity is enhanced when two phenolic hydroxyl groups are oriented in ortho or para-position. Since these structural features of phenolic compounds are responsible for antioxidant activity (Katalinic et al., 2006). Thus, polyphenols measurements in extracts may be related to their antioxidant activities. DPPH and FRAP methods have been used by many researchers to evaluate the free radical scavenging activity of antioxidant molecules and plant extracts. DPPH does generate strongly colored solutions with methanol which is eliminated in presence of antioxidants. The data obtained from the two radical scavenging methods suggest high accuracy and constancy between them.

Using two methods, some plants showed large difference in their TEAC values, whereas others showed little differences. This may be due to variation in types of phenolic compounds, that differ significantly in their reactivity towards DPPH (Katalinic et al., 2006; Benzie and Strain, 1996). Furthermore, the affinities of different *Centaurea* species toward the two above radicals were sometimes significantly altered due to variation of DPPH solubility in aqueous medium.




Figure 2. Linear correlation between the total phenolic content and antioxidant activity TEAC for methanolic extracts: correlation coefficient $r = 0.822$, coefficient of determination $r^2 = 0.976$.

REFERENCES

- Abboud, M. M., Aljundi, I. H., Khleifat, K. M., & Dmour, S. (2010). Biodegradation kinetics and modeling of whey lactose by bacterial hemoglobin VHB-expressing *Escherichia coli* strain. *Biochemical Engineering Journal*, 48(2), 166-172.
- Abboud, M. M., Saeed, H. A., Tarawneh, K. A., Khleifat, K. M., & Al Tarawneh, A. (2009). Copper uptake by *Pseudomonas aeruginosa* isolated from infected burn patients. *Current microbiology*, 59(3), 282-287.
- Adams, R.P., 1995. Identification of Essential Oil Components by Gas Chromatography–Mass Spectroscopy. Allured Publishing, Carol Stream, IL, USA.
- Al Fraijat, B., Al-Tawarah, N. M., Khlaifat, A. M., Qaralleh, H., Khleifat, K. M., Al-Zereini, W., & Al-limotm, M. O. (2019). Urinary tract infection and non-ruptured acute appendicitis association: Uropathogens findings. *TROPICAL BIOMEDICINE*, 36(3), 620-629.
- Al-Mustafa, A. H., & Al-Thunibat, O. Y. (2008). Antioxidant activity of some Jordanian medicinal plants used traditionally for treatment of diabetes. *Pak J Biol Sci*, 11(3), 351-358.
- Al-Asoufi, A., Khlaifat, A., Tarawneh, A., Alsharafa, K., Al-Limoun, M., & Khleifat, K. (2017). Bacterial quality of urinary tract infections in diabetic and non-diabetics of the population of Ma'an Province, Jordan. *Pakistan J Biol Sci*, 20, 179-88.
- ALrawashdeh, I.N., Qaralleh, H., Al-limoun, M.O., Khleifat, K.M. (2019) Antibacterial Activity of *Asteriscus graveolens* Methanolic Extract: Synergistic Effect with Fungal Mediated Nanoparticles against Some Enteric Bacterial Human Pathogens. *Journal of basic and applied Research in Biomedicine*, 5(2): 89-98
- Althunibat, O.Y.; Qaralleh, H.; Al-Dalin, S.Y.A.; Abboud, M.; Khleifat, K.; Majali, I.S.; Aldal'in, H.K.; Rayyan, W.A.; Jaafraa, A. Effect of Thymol and Carvacrol, the Major Components of *Thymus capitatus* on the Growth of *Pseudomonas aeruginosa*. *Journal of Pure and Applied Microbiology* 2016;10:367-374
- Al-Asoufi, A., Khlaifat, A., Al Tarawneh, A., Alsharafa, K., Al-Limoun, M., & Khleifat, K. (2017). Bacterial Quality of Urinary Tract Infections in Diabetic and Non Diabetics of the Population of Ma'an Province, Jordan. *Pakistan journal of biological sciences: PJBS*, 20(4), 179-188.
- Alali, F. Q., Tawaha, K., El-Elimat, T., Syouf, M., El-Fayad, M., Abulaila, K., ... & Oberlies, N. H. (2007). Antioxidant activity and total phenolic content of aqueous and methanolic extracts of Jordanian plants: an ICBG project. *Natural Product Research*, 21(12), 1121-1131.
- Baratto, M.C., Tattini, M., Galardi, C., Pinelli, P., Romani, A., Visioli, F., Basosi, R. and Pogni, R., 2003. Antioxidant activity of galloyl quinic derivatives isolated from *P. lentiscus* leaves. *Free radical research*, 37(4), pp.405-412.
- Benzie, I. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of (antioxidant power): The FRAP assay. *Analytical Biochemistry*, 293 (1), 70-76.
- Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature*. 1958;181:1199 1200.
- Boğa, M., Alkan, H., Ertaş, A., Oral, E. V., Yılmaz, M. A., Yeşil, Y., ... & Kolak, U. (2016). Phytochemical profile and some biological activities of three *Centaurea* species from Turkey. *Tropical Journal of Pharmaceutical Research*, 15(9), 1865-1875.
- Buruk, K., Sokmen, A., Aydin, F., and M. Erturk (2006). Antimicrobial activity of some endemic plants growing in the Eastern Black Sea Region, Turkey. *Fitoterapia* 77, 388-391.
- Conforti, F., Menichini, F., Loizzo, R.M., Statti, G.A., Rapisarda, A., Menichini, F., Houghton, J.P., 2008. Antioxidant, α -amylase inhibitory and brine-shrimp toxicity studies on *Centaurea centaurium* L methanolic root extract. *Natural Product Research* 22 (16), 1457–1466

- Cansaran, A., Dogan, N.M., Oztekin, M., and G. Acar (2010). Antimicrobial activity of various extracts of *Centaurea cankiriense* A. Duran and H. Duman. *Afr. J. Microbiol. Res.* 4, 608-612.
- Cutter, C.N. (2000). Antimicrobial effect of herb extracts against *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Salmonella typhimurium* associated with beef. *Journal of Food Protect.* 63, 601-607.
- Cragg, G. M., Newman, D. J., & Snader, K. M. (1997). Natural products in drug discovery and development. *Journal of natural products*, 60(1), 52-60.
- Dudonné, S., Vitrac, X., Coutiere, P., Woillez, M., & Mérillon, J. M. (2009). Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORAC assays. *Journal of agricultural and food chemistry*, 57(5), 1768-1774.
- Epand RF, Savage PB, Epand RM. Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds (Ceragenins) *Biochimica et Biophysica Acta: Biomembranes*. 2007;1768(10):2500–2509.
- Ertugrul, K., Dural, H., Tugay, O., Flamini, G., Cioni, P.L., Morelli, I., 2003. Essential oils from flowers of *Centaurea kotschyi* var. *kotschyi* and *C. kotschyi* var. *decumbens* from Turkey. *Flav. Frag. J.* 18, 95–97.
- Flamini, G., Stoppelli, G., Morelli, I., Ertugrul, K., Dural, H., Tugay, O., & Demirelma, H. (2004). Secondary metabolites from *Centaurea isaurica* from Turkey and their chemotaxonomical significance. *Biochemical systematics and ecology*, 32(6), 553-557.
- Guven, K., Çelik, S., and I. Uysal (2005). Antimicrobial activity of *Centaurea* species. *Pharm. Biol.* 43, 67-71.
- Hajjeh, R.A., Reingold, A., Weil, A., Shutt, K., Schuchat, A., and B.A. Perkins (1999). Toxic shock syndrome in the United States: Surveillance update, 1979–1996. *Emerg. Infect. Dis.* 5, 807–810.
- Hao, Y.Y., Brackett, R.E., and M.P. Doyle (1998). Efficacy of plant extracts in inhibiting *Aeromonas hydrophila* and *Listeria monocytogenes* in refrigerated cooked poultry. *Food Microbiol.* 15, 367-378.
- Hernández, N. E., Tereschuk, M. L., & Abdala, L. R. (2000). Antimicrobial activity of flavonoids in medicinal plants from Tañi del Valle (Tucuman, Argentina). *Journal of Ethnopharmacology*, 73(1-2), 317-322.
- Hillard, N.J., Schelonka, R.L., and K.B. Waites (2003). *Bacillus cereus* bacteremia in a preterm neonate. *J. Clin. Microbiol.* 41, 3441-3444.
- Jaafreh, M., Khleifat, K.M., Qaralleh, H., Al-limoun, M.O. (2019). Antibacterial and Antioxidant Activities of *Centaurea damascena* Methanolic Extract. *Journal of basic and applied Research in Biomedicine*, 5(1): 55-63
- Kargioğlu, M., Cenkci, S., Serteser, A., Konuk, M., & Vural, G. (2010). Traditional uses of wild plants in the middle Aegean region of Turkey. *Human Ecology*, 38(3), 429-450.
- Kargioğlu, M., Cenkci, S., Serteser, A., Evliyaoglu, N., Konuk, M., Kok, M.S., and Y. Bağcı (2008). An ethnobotanical survey of inner- West Anatolia, Turkey. *Human Ecol.* 36, 763-777.
- Karahan, F., Avsar, C., Ozyigit, I. I., & Berber, I. (2016). Antimicrobial and antioxidant activities of medicinal plant *Glycyrrhiza glabra* var. *glandulifera* from different habitats. *Biotechnology & Biotechnological Equipment*, 30(4), 797-804.
- Karamenderes, C., Konyalioglu, S., Khan, S., & Khan, I. A. (2007). Total phenolic contents, free radical scavenging activities and inhibitory effects on the activation of NF-kappa B of eight *Centaurea L.* species. *Phytotherapy Research*, 21, 488–491.
- Katalinic, V., M. Milos, T. Kulisic and M. Jukic, 2006. Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chem.*, 94: 550-557.
- Khleifat, K. M., Tarawneh, K. A., Wedyan, M. A., Al-Tarawneh, A. A., & Al Sharafa, K. (2008). Growth kinetics and toxicity of *Enterobacter cloacae* grown on linear alkylbenzene sulfonate as sole carbon source. *Current microbiology*, 57(4), 364-370.
- Khleifat, K. M., Matar, S. A., Jaafreh, M., Qaralleh, H., Al-limoun, M. O., & Alsharafa, K. Y. (2019). Essential Oil of *Centaurea damascena* Aerial Parts, Antibacterial and Synergistic Effect. *Journal of Essential Oil Bearing Plants*, 22(2), 356-367.
- Khleifat, K., Abboud, M., Al-Shamayleh, W., Jiries, A., & Tarawneh, K. (2006a). Effect of chlorination treatment on gram negative bacterial composition of recycled wastewater. *Pak. J. Biol. Sci.* 9, 1660-1668.
- Khleifat, K. M., Sharaf, E. F., & Al-limoun, M. O. (2015). Biodegradation of 2-chlorobenzoic acid by *Enterobacter cloacae*: Growth kinetics and effect of growth conditions. *Bioremediation Journal*, 19(3), 207-217.
- Khleifat, K. M., Abboud, M. M., & Al-Mustafa, A. H. (2006b). Effect of *Vitreoscilla hemoglobin* gene (*vgh*) and metabolic inhibitors on cadmium uptake by the heterologous host *Enterobacter aerogenes*. *Process Biochemistry*, 41(4), 930-934.
- Khleifat, K. M., Halasah, R. A., Tarawneh, K. A., Halasah, Z., Shawabkeh, R., & Wedyan, M. A. (2010). Biodegradation of linear alkylbenzene sulfonate by *Burkholderia* sp.: Effect of some growth conditions. *Int J Agr Biol*, 12, 17-25.
- Khleifat, K. M. (2006). Correlation Between Bacterial Hemoglobin and Carbon Sources: Their Effect on Copper Uptake by Transformed *E. coli* Strain α DH5. *Current microbiology*, 52(1), 64-68.
- Köse, Y. B., İşcan, G., Demirci, B., Başer, K. H. C., & Celik, S. (2007). Antimicrobial activity of the essential oil of *Centaurea aladagensis*. *Fitoterapia*, 78(3), 253-254.
- Majali, I. S., Oran, S. A., Khaled, M. K., Qaralleh, H., Rayyan, W. A., & Althunibat, O. Y. (2015). Assessment of the antibacterial effects of *Moringa peregrina* extracts. *African Journal of microbiology research*, 9(51), 2410-2414.
- Mabberley, J. (1997). The plant book, second ed. Cambridge University Press Cambridge.
- Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical microbiology reviews*, 12(4), 564-582.
- Mamtha, B., Kavitha, K., Srinivasan, K. K., & Shivananda, P. G. (2004). An in vitro study of the effect of *Centella asiatica* [Indian pennywort] on enteric pathogens. *Indian Journal of Pharmacology*, 36(1), 41.
- Miliauskas, G., Venskutonis, P. R., & Van Beek, T. A. (2004). Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food chemistry*, 85(2), 231-237.
- Mokbel, M.S., and F. Hashinaga (2006). Evaluation of the antioxidant activity of extracts from buntan (*Citrus grandis* Osbeck) fruit tissues. *Food Chem.* 94, 529-534.

- Olukoya, D.K, Idika N.and T. Odugbemi (1993). Antimicrobial activity of some medicinal plants from Nigeria. *J. Ethnopharmacol.*39: 69-72.
- Ozgen, M., Reese, R. N., Tulio, A. Z., Scheerens, J. C., & Miller, A. R. (2006). Modified 2, 2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) method to measure antioxidant capacity of selected small fruits and comparison to ferric reducing antioxidant power (FRAP) and 2, 2'-diphenyl-1-picrylhydrazyl (DPPH) methods. *Journal of Agricultural and Food Chemistry*, 54(4), 1151-1157.
- Puupponen-Pimia, R., Nohynek, L., Meier, C., Kahkonen, M., Heinonen, M., Hopia, A., and K.M. Oksman-Caldentey (2001). Antimicrobial properties of phenolic compounds from berries. *J. Appl. Microbiol.* 90, 494-507.
- Qaralleh, H. N., Abboud, M. M., Khleifat, K. M., Tarawneh, K. A., & Althunibat, O. Y. (2009). Antibacterial activity in vitro of *Thymus capitatus* from Jordan. *Pakistan journal of pharmaceutical sciences*, 22(3).
- Qaralleh, H., Khleifat, K. M., Al-Limoun, M. O., Alzedaneen, F. Y., & Al-Tawarah, N. (2019). Antibacterial and synergistic effect of biosynthesized silver nanoparticles using the fungi *Tritirachium oryzae* W5H with essential oil of *Centaurea damascena* to enhance conventional antibiotics activity. *Advances in Natural Sciences: Nanoscience and Nanotechnology*, 10(2), 025016.
- Qaralleh, H., Al-limoun, M.O., Khleifat, K.M, Alsharafa, K.Y., Al Tarawneh, A.A. (2019). *Vitreoscilla* sp. hemoglobin (VHb) Expression in Heterologous Hosts: Some Aspects of Their Biotechnological Application. *Journal of basic and applied Research in Biomedicine*, 5(2): 81-88.
- Rayne, S., and G. Mazza (2007). Biological activities of extracts from sumac (*Rhus* spp.): a review. *Plant Foods Hum. Nutr.* 62, 165–175.
- Sarker, S.D., Kumarasamy, Y., Shoeb, M., Celik, S., Yucel, E., Middleton, M., Nahar, L., 2005. Antibacterial and antioxidant activities of three Turkish species of the genus *Centaurea*. *Oriental Pharmacy and Experimental Medicine* 5 (3), 246–250.
- Sezik, E., Yesilada, E., Honda, G., Takaishi, Y., Takeda, Y., and T. Tanaka (2001). Traditional medicine in Turkey X. Folk medicine in Central Anatolia. *J. Ethnopharmacol.* 75, 95–115.
- Skaltsa, H.D., Demetzos, C., Lazari, D., Sokovic, M., 2003. Essential oil analysis and antimicrobial activity of eight *Stachys* species from Greece. *Phytochemistry* 64, 743–752.
- Sun, J., Y.F. Chu, X. Wu and R.H. Liu, 2002. Antioxidant and antiproliferative activities of common fruits. *J. Agric. Food Chem.*, 50: 7449-7454.
- Tarawneh, K. A., Al-Tawarah, N. M., Abdel-Ghani, A. H., Al-Majali, A. M., & Khleifat, K. M. (2009). Characterization of verotoxigenic *Escherichia coli* (VTEC) isolates from faeces of small ruminants and environmental samples in Southern Jordan. *Journal of basic microbiology*, 49(3), 310-317.
- Tellawi, A. M., Prescott, M. J., Ayesb, E. M., Omari, A., Tadrus, K., Helat, N., ... & Soub, H. (2001). Conservation and sustainable use of biological diversity in Jordan. *GCEP, Amman, Jordan*, 118.
- Tekeli, Y., Zengin, G., Aktumsek, A., Sezgin, M., & Torlak, E. (2011). Antibacterial activities of extracts from twelve *Centaurea* species from Turkey. *Archives of Biological Sciences*, 63(3), 685-690.
- Tepe, B., Sokmen, M., Akpulat, H.A., Yumrutas, O., Sokmen, A., 2006. Screening of antioxidative properties of the methanolic extracts of *Pelargonium endlicherianum* Fenzl., *Verbascum wiedemannianum* Fisch. & Mey., *Sideritis libanotica* Labill. subsp. *linearis* (Benth) Borm., *Centaurea mucronifera* DC and *Hieracium cappadocicum* Freyn from Turkish flora. *Food Chemistry* 98, 9–13.
- Ugur, A., Sarac, N., Ceylan, O., and M.E. Duru (2010). Antimicrobial activity and chemical composition of endemic *Centaurea cariensis* subsp. *niveotomentosa*. *Nat. Prod. Res* 24, 861-872.
- WHO. (2002). *Traditional Medicine Strategy 2002-2005*. Geneva.
- Wagstaff, S.J., and I. Breitwieser (2002). Phylogenetic relationship of New Zealand Asteraceae inferred from ITS sequences. *Plant Syst.Evol.* 231, 203-224.
- Yayli, N., Yasar, A., Yayli, A., Albay, C., Asamaz, Y., Coskuncelebi, K., and S. Karaoglu (2009). Chemical composition and antimicrobial activity of essential oils from *Centaurea appen-dicigera* and *Centaurea helenioides*. *Pharm. Biol.* 47, 7-12.
- Zablutowicz, R. M., Hoagland, R. E., & Wagner, S. C. (1996). Effect of saponins on the growth and activity of rhizosphere bacteria. In *Saponins Used in Food and Agriculture* (pp. 83-95). Springer, Boston, MA.
- Zeidan, R.; Oran, S.; Khleifat, K.; Matar, S. Antimicrobial activity of leaf and fruit extracts of Jordanian *Rubus sanguineus* Friv.(Rosaceae). *African Journal of Microbiology Research*2013;7:5114-5118
- Zengin, G., Cakmak, Y. S., Guler, G. O., & Aktumsek, A. (2010). In vitro antioxidant capacities and fatty acid compositions of three *Centaurea* species collected from Central Anatolia region of Turkey. *Food and Chemical Toxicology*, 48(10), 2638-2641.
- Zhang, X. (2004). Traditional medicine: its importance and protection. In: Twarog S, Kapoor P (eds). *Protecting and promoting traditional knowledge: systems, national experiences and international dimensions. Part I. The role of traditional knowledge in healthcare and agriculture*. New York: USA, 3-6.