

Time-dependent approaches to open quantum systems

M. Tokieda¹ and K. Hagino^{2,*}

¹*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

²*Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

Correspondence*:

K. Hagino

hagino.kouichi.5m@kyoto-u.ac.jp

ABSTRACT

Couplings of a system to other degrees of freedom (that is, environmental degrees of freedom) lead to energy dissipation when the number of environmental degrees of freedom is large enough. Here we discuss quantal treatments for such energy dissipation. To this end, we discuss two different time-dependent methods. One is to introduce an effective time-dependent Hamiltonian, which leads to a classical equation of motion as a relationship among expectation values of quantum operators. We apply this method to a heavy-ion fusion reaction and discuss the role of dissipation on the penetrability of the Coulomb barrier. The other method is to start with a Hamiltonian with environmental degrees of freedom and derive an equation which the system degree of freedom obeys. For this, we present a new efficient method to solve coupled-channels equations, which can be easily applied even when the dimension of the coupled-channels equations is huge.

Keywords: open quantum systems, quantum friction, Caldeira-Leggett model, barrier transmission, fusion reactions

1 INTRODUCTION

Open quantum systems are ubiquitous in many branches of science. In general, a system is never isolated but couples to other degrees of freedom, which are often referred to as environment. The couplings to the environmental degrees of freedom can strongly affect the dynamics of the system. When the number of environmental degrees of freedom is huge, the couplings lead to energy dissipation. It has been demonstrated by Caldeira and Leggett that such couplings suppress the tunneling rate [1], going into a transition from quantum to classical regimes. In nuclear physics, it has been well-known that a large amount of the relative energy and angular momentum are dissipated in collisions of heavy nuclei at energies close to the Coulomb barrier, known as deep inelastic collisions [2]. In this case, the dissipation occurs due to the couplings between the relative motion of two colliding nuclei and nucleonic degrees of freedom in those nuclei. A classical Langevin equation [3] has been successfully applied to describe such collisions [2]. The Langevin approach has also been employed in order to discuss fusion reactions for syntheses of superheavy elements [4, 5, 6, 7, 8, 9].

The classical Langevin approach, by definition, is not applicable at energies around the Coulomb barrier, at which quantum effects play an important role [10, 11]. One can then ask: what is a quantum model which in the classical limit is equivalent to a classical Langevin equation? There are two approaches to address this question. One is to use a phenomenological quantum friction model, in which the expectation

values of operators obey the the classical equation of motion with friction [12, 13, 14, 15]. Recently, we solved such quantum friction Hamiltonians with a time-dependent wave packet approach in order to discuss the effect of friction on quantum tunneling [16]. The other approach to a quantum Langevin equation is to start from a system-plus-bath Hamiltonian, that is, a Hamiltonian which consists of the system and the environmental degrees of freedom, and eliminate the environmental degrees of freedom. For instance, one can employ the Caldeira-Leggett Hamiltonian [1], since the classical Langevin equation can be derived from it [2, 3]. This approach is more microscopic, thus a computation would be more involved, than the quantum friction model. It has been known that, in the Markovian limit, the time evolution for the reduced density matrix for the system degree of freedom in general takes the so called Lindblad form [17, 18].

In this paper, we discuss both of these two approaches for open quantum systems, from a point of view of time-dependent method. In the next section, we first discuss the phenomenological quantum friction models using a time-dependent wave packet approach. We apply them to heavy-ion fusion reactions around the Coulomb barrier, and discuss a role of friction in fusion dynamics. In Sec. III. we solve the Caldeira-Leggett Hamiltonian using a time-dependent coupled-channels approach. Using a quantum damped harmonic oscillator, we discuss how one can deal with a large number of degrees of freedom. A summary of the paper is then given in Sec. IV.

2 PHENOMENOLOGICAL QUANTUM FRICTION MODELS

We first consider a phenomenological approach to quantum friction. In this approach, one treats the environmental degrees of freedom implicitly and introduces a phenomenological Hamiltonian with which the classical equation of motion with a frictional force is reproduced as expectation values. For this purpose, several model Hamiltonians have been proposed so far [12, 13, 14, 15]. Among these, we focus in this paper on the one introduced by Kostin [14].

Consider a particle of mass m moving in a one dimensional space q under the influence of a potential $V(q)$. With a friction coefficient γ , the phenomenological Schrödinger equation in the Kostin model is given by [14]

$$i\hbar\frac{\partial}{\partial t}\psi(q,t) = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial q^2} + V(q) + \gamma S(q,t) \right] \psi(q,t), \quad (1)$$

where $S(q,t)$ is the phase of the wave function, $\psi(q,t) = |\psi(q,t)| \exp(iS(q,t)/\hbar)$. From this equation, it is easy to confirm that one can derive the equation of motion with a frictional force

$$\frac{d}{dt}\langle p \rangle = -\gamma\langle p \rangle - \left\langle \frac{dV}{dq} \right\rangle, \quad (2)$$

as is desired. Here, the expectation value of an operator \mathcal{O} is denoted as $\langle \mathcal{O} \rangle = \int dq \psi^*(q,t)\mathcal{O}\psi(q,t)$ and p is the momentum operator.

When one simulates an energy dissipation in nuclear collisions by means of friction, a frictional force should be active only when the colliding nuclei are close to each other. In other words, one needs to deal with a coordinate dependent friction coefficient, $\gamma = \gamma(q)$. An extension of the Kostin model along this line has been proposed in Refs. [19, 20], with which the modified Schrödinger equation is given by

$$i\hbar\frac{\partial}{\partial t}\psi(q,t) = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial q^2} + V(q) + \int^q dq_1 \gamma(q_1)\frac{\partial}{\partial q_1}S(q_1,t) \right] \psi(q,t). \quad (3)$$

To apply the phenomenological model to realistic collision problems, one further needs an extension to a three dimensional space, $\vec{q} = \vec{q}(r, \theta, \phi)$. To this end, we first expand the wave function with the Legendre polynomials $P_l(x)$ as $\psi(\vec{q}, t) = \sum_{l=0}^{\infty} u_l(r, t) P_l(\cos \theta)/r$. One can then modify the Schrödinger equation for $u_l(r, t)$ in the same way as Eq. (3) to incorporate a frictional force,

$$i\hbar \frac{\partial}{\partial t} u_l(r, t) = \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial r^2} + \frac{\hbar^2}{2m} \frac{l(l+1)}{r^2} + V(r) + \int^r dr_1 \gamma(r_1) \frac{\partial}{\partial r_1} S_l(r_1, t) \right] u_l(r, t), \quad (4)$$

where $S_l(r, t)$ is the phase of the radial wave function, $u_l(r, t) = |u_l(r, t)| \exp(iS_l(r, t)/\hbar)$. We have here assumed a spherically symmetric potential, $V(\vec{q}) = V(r)$. Notice that only the radial friction is taken into account here, while the angular momentum dissipation is neglected. In the following, we only consider the s-wave scattering, $l = 0$.

In applying Eq. (4) to scattering problems, one needs to use the time-dependent approach, since the Hamiltonian depends explicitly on time. To this end, we propagate a wave packet and observe how it bifurcates after it crosses the potential region. Since a wave packet is a superposition of various energy waves, one has to choose carefully the initial condition to get scattering quantities at certain initial energy. Notice that the energy projection approach [21] is inapplicable for our purpose, since the energy is not conserved.

In the initial condition, the width of the energy distribution must be small enough to get reasonable results. In this context, the energy means the expectation value of the asymptotic Hamiltonian, H_0 . If $V(r)$ rapidly vanishes as $r \rightarrow \infty$, one can simply take the kinetic energy operator as H_0 , that is, $H_0 = -\hbar^2/2m (\partial^2/\partial r^2)$. The minimum uncertainty wave packet in this case has been discussed in Ref. [22], which reads

$$u_0^{\min}(r, t = 0) \propto r e^{-(r-r_0)^2/4\sigma_r^2} e^{ik_0 r}, \quad (5)$$

where r_0 and σ_r are related to the mean position and the width of the wave function in the coordinate space, respectively, and k_0 is related to the mean initial energy.

In nuclear collisions, on the other hand, the potential is a sum of the nuclear potential V_N and the Coulomb potential $V_C(r) = Z_P Z_T e^2/r$ with the projectile charge Z_P and the target charge Z_T . Since the Coulomb potential is a long range potential, the asymptotic Hamiltonian H_0 has to include it, that is, $H_0 = -\hbar^2/2m (\partial^2/\partial r^2) + V_C$. Thus, the minimum uncertainty wave packet in the form of Eq. (5) would not be efficient in this case. Instead, one needs to construct a wave packet from the energy distribution, $f_C(E)$, of H_0 in the presence of the Coulomb potential. In analogy to the spherical Bessel functions, we find that this can be achieved as

$$u_0^C(r, t = 0) \propto \int_0^{\infty} dk F_0(\eta, kr) e^{ikr_0} \sqrt{k f_C(E)} \quad (6)$$

with $E = \hbar^2 k^2/2m$, where $\eta = m Z_P Z_T e^2/\hbar^2 k$ is the Sommerfeld parameter and $F_0(\eta, kr)$ is the regular Coulomb wave function.

With the initial condition given by Eq. (6), we compute the penetrability of the Coulomb barrier for the $^{16}\text{O} + ^{208}\text{Pb}$ system in the presence of friction. For the nuclear potential, we employ the optical potential in Ref. [23], that is,

$$V_N(r) = \frac{V_0}{1 + \exp((r - R_v)/a_v)} + i \frac{W_0}{1 + \exp((r - R_w)/a_w)}, \quad (7)$$




Figure 1. Energy dependence of the penetrability of the Coulomb barrier for the $^{16}\text{O}+^{208}\text{Pb}$ reaction. The dashed line shows the result without friction, while the solid line is for the result with friction.

with $V_0 = -901.4$ MeV, $R_v = 8.44$ fm, $a_v = 0.664$ fm, $W_0 = -30$ MeV, $R_w = 6.76$ fm, and $a_w = 0.4$ fm. With this potential, the Coulomb barrier height V_B is found to be 74.5 MeV.

For a friction coefficient $\gamma(r)$, we employ the surface friction model [2],

$$\gamma(r) = \frac{\gamma_0}{m} \left(V_B \frac{df_{\text{WS}}}{dr} \right)^2, \quad (8)$$

with the Woods-Saxon form factor $f_{\text{WS}} = 1/(1 + \exp((r - R_v)/a_v))$. This is a general form of the friction coefficient obtained perturbatively [24], and has successfully been applied to above barrier fusion reactions and to deep inelastic scatterings [2]. We arbitrarily set $\gamma_0 = 4.7 \times 10^{-23}$ s/MeV, which was used in the classical calculations. We compute the phase of the wave function in the same way as in Ref. [16].

For the initial energy distribution in Eq. (6), we assume the Gaussian form,

$$f_C(E) = \frac{1}{\sqrt{2\pi\sigma_E^2}} e^{-(E-E_0)^2/2\sigma_E^2}. \quad (9)$$

where E_0 and σ_E are the mean and the width of the initial energy distribution. We have confirmed that $\sigma_E = 0.5$ MeV is sufficient in the present parameter set.

Fig. 1 compares the penetrability obtained with and without friction. One finds that the penetrability with friction is shifted to higher energies around the barrier. That is, in the presence of friction, a particle needs an additional energy to penetrate the barrier, which is originated from the energy dissipation. One can also see that the penetrability does not reach unity at high energies, but it is almost saturated at around 0.9. This means that the exit channel is in a quantum superposition state of absorption and reflection even at sufficiently above barrier energies. Notice that, in classical mechanics without a fluctuation force, the penetrability can be only 0 or 1. In this sense, this is peculiar to the quantum friction model.

In low-energy fusion reactions, it has been known that low-lying collective excitations during the process play a crucial role [25]. In the calculation shown in this paper, a part of such effect is implicitly taken into account in the nuclear potential. An explicit treatment of low-lying collective excitations together with a frictional force has been carried out in Ref. [26], in which the experimental fusion cross sections for the $^{16}\text{O}+^{208}\text{Pb}$ system are well reproduced from subbarrier to above barrier energies. In that calculation, the same behavior as in Fig. 1 has been found, which may be a key to achieve a consistent description.

3 SYSTEM-PLUS-BATH MODEL

We next consider a more microscopic model for quantum friction, employing a system-plus-bath model. To be more specific, we consider the Caldeira-Leggett model [1], whose Hamiltonian is given by,

$$H_{\text{tot}} = H_S + \sum_i \hbar\omega_i a_i^\dagger a_i + h(q) \sum_i d_i (a_i^\dagger + a_i), \quad (10)$$

$$\equiv H_S + H_B + V_{\text{coup}}, \quad (11)$$

where H_S and H_B are the Hamiltonians for the system and the bath degrees of freedom, respectively, while V_{coup} is the coupling Hamiltonian between the system and the bath. Here, the bath degree of freedom is assumed to be a set of harmonic oscillators, whose creation and annihilation operators are denoted by a_i^\dagger and a_i , respectively. The coupling Hamiltonian is assumed to be separable between the system and the bath degrees of freedom. In there, d_i is the coupling strength, and $h(q)$ is the coupling form factor, where q is the coordinate of the system.

There are several ways to solve the Caldeira-Leggett Hamiltonian. In Ref. [1], the bath degrees of freedom are integrated out using the path integral in order to obtain an effective action for the system degree of freedom (see also Ref. [27]). One can also introduce the influence functional [28]. Here we discuss the coupled-channels approach [29].

In the coupled-channels approach [25], one expands the total wave function in terms of the eigen-wave functions of H_B , that is,

$$\Psi_{\text{tot}}(q, t) = \sum_{\{n_i\}} \psi_{\{n_i\}}(q, t) |\{n_i\}\rangle, \quad (12)$$

where the basis states $|\{n_i\}\rangle$ are given by,

$$|\{n_i\}\rangle = \prod_i \frac{1}{\sqrt{n_i!}} (a_i^\dagger)^{n_i} |0\rangle. \quad (13)$$

Here, $|0\rangle$ is the vacuum state defined as $a_i|0\rangle = 0$. One can derive the coupled equations for $\psi_{\{n_i\}}(q, t)$ by evaluating the equation,

$$\langle\{n_i\}| i\hbar \frac{\partial}{\partial t} |\Psi_{\text{tot}}\rangle = \langle\{n_i\}| H_{\text{tot}} |\Psi_{\text{tot}}\rangle, \quad (14)$$

that is,

$$i\hbar \frac{\partial}{\partial t} \psi_{\{n_i\}}(q, t) = \left(H_S + \sum_i n_i \hbar\omega_i \right) \psi_{\{n_i\}}(q, t) + \sum_{\{n'_i\}} \langle\{n_i\}| V_{\text{coup}} |\{n'_i\}\rangle \psi_{\{n'_i\}}(q, t). \quad (15)$$

The coupled-channels equations, Eq. (15), can be numerically solved when the number of the oscillator modes is not large [25, 30]. However, in general, the number of the oscillator modes can be huge, or the distribution of the frequency of the oscillator may even be given as a continuous function. In that situation, it is almost hopeless to solve the coupled-channels equations directly. In order to overcome this problem, we introduce a more efficient basis to expand the total wave function [29]. To this end, we first expand the

function $e^{-i\omega t}$ with a finite basis set as,

$$e^{-i\omega t} \sim \sum_{k=1}^K \eta_k(\omega) u_k(t), \quad (16)$$

where $u_k(t)$ is a known function such as a Bessel function, and $\eta_k(\omega)$ is the expansion coefficient. We then introduce a new phonon creation operator as,

$$b_k^\dagger = \sum_i \frac{d_i}{\hbar} \eta_k(\omega_i) a_i^\dagger. \quad (17)$$

Notice that the number of k is finite, k running from 1 to K , even though the number of i may be infinite. We then construct the basis states using the operators b_k^\dagger , and expand the total wave function with them. That is, instead of Eq. (12), we expand the total wave function as,

$$\Psi_{\text{tot}}(q, t) = \sum_{\{\tilde{n}_k\}} \tilde{\psi}_{\{\tilde{n}_k\}}(q, t) |\{\tilde{n}_k\}\rangle, \quad (18)$$

with

$$|\{\tilde{n}_k\}\rangle = \prod_{k=1}^K \frac{1}{\sqrt{\tilde{n}_k!}} (b_k^\dagger)^{\tilde{n}_k} |0\rangle. \quad (19)$$

One can then obtain the coupled-channels equations similar to Eq. (15), that is,

$$i\hbar \frac{\partial}{\partial t} \tilde{\psi}_{\{\tilde{n}_k\}}(q, t) = H_S \tilde{\psi}_{\{\tilde{n}_k\}}(q, t) + \sum_{\{\tilde{n}'_k\}} \langle \{\tilde{n}_k\} | H_B + V_{\text{coup}} | \{\tilde{n}'_k\} \rangle \tilde{\psi}_{\{\tilde{n}'_k\}}(q, t). \quad (20)$$

We once again emphasize that the dimension of the coupled-channels equations, Eq. (20), is much smaller than that of the original equations, Eq. (15).

The structure of the coupled-channels equations, Eq. (20), becomes simple when the basis functions $u_k(t)$ satisfy the following two conditions.

1. The matrix D defined as

$$D_{kk'} \equiv \frac{1}{\hbar^2} \sum_i d_i^2 \eta_k(\omega) \eta_{k'}^*(\omega), \quad (21)$$

is diagonal with respect k and k' . That is, $D_{kk'} = \lambda_k \delta_{k,k'}$. Notice that the matrix D can be expressed also as

$$D_{kk'} \equiv \frac{1}{\hbar} \int_{-\infty}^{\infty} d\omega J(\omega) \eta_k(\omega) \eta_{k'}^*(\omega), \quad (22)$$

with the spectral density given by,

$$J(\omega) = \frac{1}{\hbar} \sum_i d_i^2 \delta(\omega - \omega_i). \quad (23)$$

2. The basis function $u_k(t)$ is closed under differentiation, that is,

$$\frac{du_k(t)}{dt} = \sum_{k'=1}^K C_{kk'} u_{k'}(t). \tag{24}$$

Notice that Bessel functions satisfy this condition since the following relation holds,

$$\frac{d}{dx} J_k(x) = -\frac{1}{2} J_{k+1}(x) + \frac{1}{2} J_{k-1}(x), \tag{25}$$

(with $J_{-k}(x) = (-1)^k J_k(x)$ for an integer value of k).

See Eq. (31) in Ref. [29] for the explicit form of the coupled-channels equations, Eq. (20). Ref. [29] also provides an alternative derivation of the coupled-channels equations, which uses the influence functional of the path integral method. This allows one to extend the present formalism to finite temperatures.

Figures 2 and 3 show results of a time evolution for a damped harmonic oscillator [29], for which we take the Hamiltonian for the system, H_S in Eq. (10), as

$$H_S = \frac{p^2}{2M} + \frac{1}{2} M \omega_S^2 q^2 + h^2(q) \sum_i \frac{d_i^2}{\hbar \omega_i}, \tag{26}$$

where M and ω_S are the mass and the frequency of the system, respectively, and the last term represents the so called counter term. In the following, we measure the length of the system in units of the oscillator length q_S defined by $q_S \equiv \sqrt{\hbar/M\omega_S}$, and take the coupling form factor, $h(q)$, to be $h(q) = q/q_S$. We assume that the bath oscillators are distributed according to the spectral density (see Eq. (23)) as

$$J(\omega) = V_I \frac{\omega}{\Omega} \sqrt{1 - \left(\frac{\omega}{\Omega}\right)^2}. \tag{27}$$

In the numerical calculations shown below, we take $\hbar\omega_S = 2$ eV, $V_I = 1$ eV, and $\hbar\Omega = 4$ eV.

At $t = 0$, we assume that $\tilde{\psi}_{\{\tilde{n}_k\}}(q, t = 0) = 0$ for $N \equiv \sum_{k=1}^K \tilde{n}_k \neq 0$. For $N = 0$, that is, for $\tilde{n}_k = 0$ for all k , we assume that the wave function is given by,

$$\tilde{\psi}_{N=0}(q, t = 0) = \frac{1}{\sqrt[4]{2\pi\sigma_0^2}} e^{-(q-q_0)^2/4\sigma_0^2} e^{ip_0q/\hbar}, \tag{28}$$

with $q_0/q_S = -1$, $\sigma_0/q_S = 1/\sqrt{2}$, and $p_0q_S/\hbar = 0$.

The upper panel of Fig. 2 shows the norm for each phonon state N as a function of $\omega_S t$. Here, the norm is defined as,

$$\mathcal{N}_N(t) \equiv \sum_{\{\tilde{n}_k\}} \int dq |\tilde{\psi}_{\{\tilde{n}_k\}}(q, t)|^2 \delta_{\sum_k \tilde{n}_k, N}. \tag{29}$$

To draw this figure, we take Bessel functions, $J_k(\Omega t)$, for $u_k(t)$ in Eq. (16) with $K = 20$. A new basis is then constructed by diagonalizing the matrix D in Eq. (21). With this basis, we solve the coupled-channels equations by including the phonon states with $N \leq 5$. The expectation value of the norm is also shown in the lower panel. As is expected, the number of phonon in the bath gradually increases as a function of




Figure 2. (The upper panel) The norm for each phonon number, $N = \sum_{k=1}^K \tilde{n}_k$. The solid line with squares, triangles, inverted triangles, diamonds, pentagons, and circles are for $N = 0, 1, 2, 3, 4,$ and 5 , respectively. These are obtained by solving the coupled-channels equations with the phonon states up to $N_{\max} = 5$, for which the phonon operators are defined with the Bessel function basis with $K = 20$. (The Lower panel) The expectation value of the number of phonon.




Figure 3. Comparison between the present method and the exact results for the expectation values of several quantities, that is, $\xi_q = \langle q \rangle / q_S$, $\xi_p = \langle p \rangle q_S / \hbar$, $\xi_{qq} = \langle (q - \langle q \rangle)^2 \rangle / q_S^2$, and $\xi_{pp} = \langle (p - \langle p \rangle)^2 \rangle q_S^2 / \hbar^2$. The solid lines show the exact results, while the solid lines with squares, triangles, and circles are the results of the present method with $N_{\max} = 3, 4,$ and 5 , respectively.

time. Notice that the contribution of the 5-phonon states is small in the whole time range shown in the figure. This justifies the truncation at $N_{\max} = 5$ for the present parameter set. One can also see that the contribution of each phonon reaches its equilibrium at around $\omega_S t = 6$.

Fig. 3 compares the results of the present method with the exact solution for the quantum damped harmonic oscillator. To this end, we evaluate the expectation values for the following four quantities: $\xi_q \equiv \langle q \rangle / q_S$, $\xi_p \equiv \langle p \rangle q_S / \hbar$, $\xi_{qq} \equiv \langle (q - \langle q \rangle)^2 \rangle / q_S^2$, and $\xi_{pp} \equiv \langle (p - \langle p \rangle)^2 \rangle q_S^2 / \hbar^2$. We carry out the calculations with three different values of N_{\max} , that is, $N_{\max} = 3, 4,$ and 5 , and compare them with the

exact results shown by the solid lines. One can see that all of the calculations with $N_{\max} = 3, 4,$ and 5 reproduce the exact results up to $\omega_{St} \sim 5$, for which $J_{20}(\Omega t)$ is negligibly small and thus the expansion in Eq. (16) with Bessel functions up to $K = 20$ (that is, up to $J_{19}(\Omega t)$) is reasonable. The deviation from the exact results become significant for larger values of ω_{St} , especially for the second order moments, ξ_{qq} and ξ_{pp} . This is a natural consequence of the fact that the larger number of phonon states are required to describe the finer structures.

4 SUMMARY

We have discussed two time-dependent methods for quantum friction. The first method is based on an effective Hamiltonian, which is constructed so that expectation values of operators obey a classical equation of motion with friction. Such Hamiltonian is in general time-dependent, and we have solved it with a time-dependent wave packet method. The other method is to start with a total Hamiltonian with both the system and the environmental degrees of freedom and then eliminate the environmental degree of freedom to derive an equation which the system degree of freedom obeys. For this approach, we have presented a new efficient basis for coupled-channels equations. These two methods are complimentary to each other. In the first method, whereas several parameters have to be determined phenomenologically, a required computational time is much shorter than the second method. On the other hand, the second method is based on a more microscopic Hamiltonian, and thus less empirical inputs are required, even though a computational time may be large. By combining these two approaches appropriately, one may be able to achieve a quantum description of heavy-ion deep inelastic collisions as well as fusion reactions to synthesize superheavy elements.

AUTHOR CONTRIBUTIONS

The contribution of each of the authors has been significant and the manuscript is the result of an even effort of both the authors.

FUNDING

This work was supported by Tohoku University Graduate Program on Physics for the Universe (GP-PU), and JSPS KAKENHI Grant Numbers JP18J20565 and 19K03861.

ACKNOWLEDGMENTS

We would like to thank Denis Lacroix and Guillaume Hupin for useful discussions.

REFERENCES

- [1] A.O. Caldeira and A.J. Leggett, Phys. Rev. Lett. **46**, 211 (1981); Ann. Phys. **149**, 374 (1983).
- [2] P. Fröbrich and R. Lipperheide, *Theory of Nuclear Reactions*, (Oxford University Press, Oxford, 1996).
- [3] Y. Abe, S. Ayik, P-G. Reinhard, and E. Suraud, Phys. Rep. **275**, 49 (1996).
- [4] Y. Abe, D. Boilley, B.G. Giraud, and T. Wada, Phys. Rev. E **61**, 1125 (2000).
- [5] C.W. Shen, G. Kosenko, and Y. Abe, Phys. Rev. C **66**, 061602 (2002).
- [6] Y. Aritomo and M. Ohta, Nucl. Phys. A **744**, 3 (2004).

-
- [7] W.J. Swiatecki, K. Siwek-Wilczynska, and J. Wilczynski, *Acta Phys. Pol. B* **34**, 2049 (2003); *Phys. Rev. C* **71**, 014602 (2005).
- [8] V.I. Zagrebaev and W. Greiner, *Nucl. Phys. A* **944**, 257 (2015).
- [9] K. Hagino, *Phys. Rev. C* **98**, 014607 (2018).
- [10] N. Takigawa, S. Ayik, K. Washiyama, and S. Kimura, *Phys. Rev. C* **69**, 054605 (2004).
- [11] S. Ayik, B. Yilmaz, A. Gokalp, O. Yilmaz, and N. Takigawa, *Phys. Rev. C* **71**, 054611 (2005).
- [12] P. Caldirola, *Nuovo Cimento* **18**, 393 (1941).
- [13] E. Kanai, *Prog. Theor. Phys.* **3**, 440 (1948).
- [14] M.D. Kostin, *J. Chem. Phys.* **57**, 3589 (1972).
- [15] K. Albrecht, *Phys. Lett. B* **56**, 127 (1975).
- [16] M. Tokieda and K. Hagino, *Phys. Rev. C* **95**, 054604 (2017).
- [17] G. Lindblad, *Commun. Math. Phys.* **48**, 119 (1976).
- [18] P. Pearle, *Euro. J. Phys.* **33**, 805 (2012).
- [19] J.D. Immele, K.-K. Kan and J.J. Griffin, *Nucl. Phys. A* **241**, 47 (1975).
- [20] P. Bargueño and S. Miret-Artès, *Ann. Phys.* **346**, 59 (2014).
- [21] K. Yabana, *Prog. Theor. Phys.* **97**, 437 (1997).
- [22] C. Bracher, *Am. J. Phys.* **79**, 313 (2011).
- [23] M. Evers *et al.*, *Phys. Rev. C* **78**, 034614 (2008).
- [24] D.H.E. Gross and H. Kalinowski, *Phys. Rep.* **45**, 175 (1978).
- [25] K. Hagino and N. Takigawa, *Prog. Theo. Phys.* **128**, 1061 (2012).
- [26] M. Tokieda and K. Hagino, to be published in the proceeding of the International Conference on Nucleus-Nucleus collisions (NN2018).
- [27] N. Takigawa and G.F. Bertsch, *Phys. Rev. C* **29**, 2358 (1984).
- [28] A.B. Balantekin and N. Takigawa, *Ann. Phys. (N.Y.)* **160**, 441 (1985).
- [29] M. Tokieda and K. Hagino, *Ann. of Phys.* **412**, 168005 (2019); arXiv:1909.10418.
- [30] K. Hagino, N. Rowley, A.T. Kruppa, *Comput. Phys. Comm.* **123**, 143 (1999).