

# Dynamics of non cohomologically hyperbolic automorphisms of $\mathbb{C}^3$

Frédéric Protin

Received: date / Accepted: date

**Abstract** We study the dynamics of a family of non cohomologically hyperbolic automorphisms  $f$  of  $\mathbb{C}^3$ . We construct a compactification  $X$  of  $\mathbb{C}^3$  where their extensions are algebraically stable. We finally construct canonical invariant closed positive  $(1,1)$ -currents for  $f^*$ ,  $f_*$  and we study several of their properties. Moreover, we study the well defined current  $T_f \wedge T_{f^{-1}}$  and the dynamics of  $f$  on its support. Then we construct an invariant positive measure  $T_f \wedge T_{f^{-1}} \wedge \phi_\infty$ , where  $\phi_\infty$  is a function defined on the support of  $T_f \wedge T_{f^{-1}}$ . We prove that the support of this measure is compact and pluripolar. We prove also that this measure is canonical, in some sense that will be precised.

**Keywords** complex dynamics · invariant current · invariant measure

## 1 Introduction

Our purpose is to study the dynamics of a family of non cohomologically hyperbolic automorphisms (in the sense of [9]) of  $\mathbb{C}^3$ . The dynamics of such polynomial automorphisms is less understood than that of cohomologically hyperbolic automorphisms. More precisely, we study the following family:

$$f : (x, y, z) \in \mathbb{C}^3 \longmapsto (y, z, yz + by + cz + dx + e) \in \mathbb{C}^3, \quad (1)$$

where  $b, c, d, e$  are complex parameters, with  $d \neq 0$ . These are the first examples of non cohomologically hyperbolic automorphisms with a rich dynamics. This family has already been studied by Maegawa [12], but our approach is quite different. We will construct a dynamical system  $(X, f)$ , where  $X$  is a "nice compactification" of  $\mathbb{C}^3$ .

---

F. Protin  
INSA de Toulouse  
E-mail: fredprotin@yahoo.fr

In Section 1 we construct the space  $X$ , a "nice compactification" of  $\mathbb{C}^3$  useful for the study of  $f$ . We then obtain in Section 2 a canonical invariant class in  $H^{1,1}(X, \mathbb{R})$ . This is the same for  $f$  and  $f^{-1}$ . We prove that this class is Kähler. This allows us to construct, in Section 3, positive invariant currents  $T_f$  and  $T_{f^{-1}}$  for  $f$  and  $f^{-1}$ . We show that their potentials are continuous in  $\mathbb{C}^3$  and pluriharmonic on the basin of attraction of infinity. In Section 4 we study the current  $T_f \wedge T_{f^{-1}}$  and give a filtration of the dynamics of  $f$  on  $\text{supp}(T_f \wedge T_{f^{-1}})$ .

## 2 Good dynamical compactification for $f$ and $f^{-1}$

Up to conjugating  $f$  by a suitable affine automorphism, we can assume without loss of generality that  $e = 0$ . We then have

$$f^{-1}(x, y, z) = \left( \frac{z - xy - bx - cy}{d}, x, y \right). \quad (2)$$

If  $\tau(x, y, z) = (z, y, x)$ , then  $\tau^{-1} = \tau$  and  $\tau f \tau = (xy + by + cx + dz, x, y)$ , which has the same form as  $f^{-1}$ . Therefore, the general behaviour of  $f$  near infinity can be obtained from that of  $f^{-1}$  by symmetry. Note that the jacobian of  $f$  is constant and equal to  $|d|$ .

**Extension of  $f$  to  $\mathbb{P}^3$ .** Let us denote by  $[x, y, z, t]$  the homogeneous coordinates for  $\mathbb{P}^3$ , and by  $H_\infty := (t = 0)$  the hyperplane at infinity. Written in homogeneous coordinates, the extension of  $f$  to  $\mathbb{P}^3 = \mathbb{C}^3 \sqcup H_\infty$ , still denoted by  $f$ , is:

$$f = [yt : zt : yz + byt + czt + dxt : t^2]. \quad (3)$$

We denote by  $I_f$  the set of indeterminacy of  $f$ , i.e. the points on  $H_\infty$  where  $f$  does not extend continuously.

Recall that a rational transformation  $g : X \rightarrow X$  of a projective manifold  $X$  is *algebraically stable* (AS) if there is no hypersurface mapped by some iterate  $g^n$  in the set  $I^+ := \cup_n I_{g^n}$  ([6]). This is equivalent to the fact that  $g$  induces a linear action  $g^*$  on the cohomology vector space  $H^{1,1}(X, \mathbb{R})$  which is compatible with the dynamics, in the sense that  $(g^n)^* = (g^*)^n$  for all  $n \in \mathbb{N}$ . This condition is crucial in order to construct interesting invariant currents.

Note that  $f$  is not AS in  $\mathbb{P}^3$  since it contracts the hyperplane at infinity  $H_\infty = (t = 0)$  onto the point of indeterminacy  $[0, 0, 1, 0] \in I_f$ . Setting

$$L := (Y = T = 0), \quad p^+ := [1, 0, 0, 0], \quad p^- := [0, 0, 1, 0],$$

$$L' := (Z = T = 0), \quad L'' := (X = T = 0),$$

we have the following more accurate description of the behavior of  $f$  at infinity in  $\mathbb{P}^3$ .

**Lemma 1** *The indeterminacy loci of  $f$ , respectively  $f^{-1}$  in  $\mathbb{P}^3$  are*

$$I_f = L \cup L', \quad I_{f^{-1}} = L \cup L'', \quad I_{f^2} = I_f \quad \text{and} \quad I_{f^{-2}} = I_{f^{-1}}.$$

Let  $x \in L \setminus \{p^+\}$  and  $x' \in L \setminus \{p^-\}$ . Then

$$f(x) = L'' \quad \text{and} \quad f^{-1}(x') = L'.$$

Let  $y \in L' \setminus \{p^+\}$  and  $y' \in L'' \setminus \{p^-\}$ . Then

$$f(y) = L \quad \text{and} \quad f^{-1}(y') = L.$$

We have  $f(p^+) = H_\infty$  and  $f^{-1}(p^-) = H_\infty$ .

*Proof* By symmetry it is sufficient to prove the result for  $f$ . Let  $x \in L \setminus p^+$  and a sequence  $(z_n) \in \mathbb{P}^3 \setminus I_f$  such that  $z_n \rightarrow x$ . Then  $(f(z_n))$  accumulates on  $(X = T = 0)$  by (3). In a similar way we compute  $f(y) = L$  when  $y \in (Z = T = 0) \setminus p^+$ .

Now let  $(z_n) \rightarrow p^+$ ; this means that  $\frac{Y}{X} \rightarrow 0$ ,  $\frac{Z}{X} \rightarrow 0$ ,  $\frac{T}{X} \rightarrow 0$ . Moreover suppose that  $z_n$  belongs to the cubic hypersurface  $S_e$  defined by

$$\frac{z}{t} + d\frac{x}{y} = e \tag{4}$$

for  $e \in \mathbf{C}$  (note that  $p^+ \in S_e$ ). In this case, we can replace the homogeneous representation (3) by  $f = [1, \frac{Z}{Y}, b + c\frac{Z}{Y} + e, \frac{T}{Y}]$ . For all  $a', b' \in \mathbf{C}$  there exists  $e \in \mathbf{C}$  and  $(z_n) \rightarrow p^+$  such that  $(z_n) \in S$  and  $\frac{Z}{Y} \rightarrow a'$ ,  $b + c\frac{Z}{Y} + e \rightarrow b'$ .


Let us now compute the last coordinate of  $f$  in this case. We have  $\frac{T}{Y} = \frac{T}{Z} \frac{Z}{Y}$ . Since  $\frac{Z}{Y} \rightarrow a'$  and  $\frac{T}{Z} \rightarrow 0$  according to (4) (recall that  $\frac{Y}{X} \rightarrow 0$ ), we have  $\frac{T}{Y} \rightarrow 0$ . Finally, for every  $q \in H_\infty$  there exists a sequence converging to  $p^+$  that is mapped into a sequence converging to  $q$ , hence  $f(p^+) = H_\infty$ .  $\square$

We note that the hyperplane  $H_\infty \setminus I_f^+$  is contracted into the point  $p^- \in I_f$  by  $f$  and into the point  $p^+ = [1 : 0 : 0 : 0] \in I_{f^{-1}}$  by  $f^{-1}$ : thus neither  $f$  nor  $f^{-1}$  is algebraically stable on  $\mathbf{P}^3$ . Figure 1 summarizes the image by  $f$  of the points of indeterminacy;  $I_f$  is represented by straight lines and the action by  $f$  by curved arrows. The figure for  $f^{-1}$  is obtained by symmetry.

The main result of this section is the following.

**Proposition 1** *Let  $X$  denote the blow-up of  $\mathbb{P}^3$  having as smooth center the line  $L = (Y = T = 0)$ , and let  $f : X \rightarrow X$  denote the lift of  $f$ . Then  $f : X \rightarrow X$  is algebraically stable.*

More precisely, we are going to show that the exceptional divisor  $E$  of the blow-up  $\pi : X \rightarrow \mathbb{P}^3$  along  $L$  is mapped by  $f$  onto regular points of  $\tilde{H}_\infty$ , the strict transform of the hyperplane at infinity  $H_\infty$ , and that  $\tilde{H}_\infty$  is contracted by  $f$  onto a regular superattractive fixed point. Let us still denote by  $p^+$  and  $p^-$  the points where  $\tilde{H}_\infty$  is mapped by  $f^{-1}$  and  $f$  respectively, and by  $I_f, I_{f^{-1}}$  the indeterminacy locus for  $f$ , respectively  $f^{-1}$ . We denote also by  $L', L''$  the strict transform of  $L', L'' \subset \mathbb{P}^3$ .



**Fig. 1** Divisor at infinity of  $\mathbb{P}^3(\mathbb{C})$ . The indeterminacy set is represented by straight lines and the action by  $f$  by curved arrows.

*Proof Image of  $\tilde{H}_\infty$  by  $f$ .* We first express  $f$  in the chart  $Z \neq 0$  at the input and output. The origin of this chart is  $p^-$ . We have

$$\begin{aligned} f(x, y, t) &:= f[x : y : 1 : t] = [yt : t : y + byt + ct + dxt : t^2] \\ &= \left( \frac{yt}{y + byt + ct + dxt}, \frac{t}{y + byt + ct + dxt}, \frac{t^2}{y + byt + ct + dxt} \right). \end{aligned}$$

Recall that the blow-up variety  $X$  is described by:

$$\{(x, y, t, [\xi_1, \xi_2]) \in \mathbf{C}^3 \times \mathbf{P}^1, \xi_1 t = \xi_2 y\}. \quad (5)$$

The expression of  $f$  in the chart  $\xi_1 \neq 0$  at input and output is then:

$$\begin{array}{ccc} & f & \\ (\omega_1, \omega_2, \omega_3) & \longmapsto & \left( \frac{\omega_2 \omega_3}{A}, \frac{\omega_3}{A}, \omega_2 \omega_3 \right) \\ \downarrow \pi & & \downarrow \pi \\ (\omega_1, \omega_2, \omega_2 \omega_3) & \longmapsto & \left( \frac{\omega_2 \omega_3}{A}, \frac{\omega_3}{A}, \frac{\omega_2 \omega_3^2}{A} \right) \end{array} \quad (6)$$

where  $A = 1 + b\omega_2\omega_3 + c\omega_3 + d\omega_1\omega_3$ . Note that the point  $p^-$  at the origin of this chart is fixed, superattractive; we verify that  $f(\tilde{H}_\infty) = p^-$ . It remains to prove that  $E$  is never mapped by  $f^n$  onto a point of indeterminacy.

**Image of  $E$  by  $f$ .** The exceptional divisor ( $\omega_2 = 0$ ) is mapped by  $f$  (except for the points of indeterminacy) on the strict transform of the line at infinity  $L'' := (x = t = 0)$ .

Note for later use that there is no point of indeterminacy in  $E \cap \widetilde{H}_\infty$  in the chart ( $Z \neq 0, \xi_1 \neq 0$ ).

The expression of  $f$  in the charts  $\xi_2 \neq 0$  at input and  $\xi_1 \neq 0$  at output is

$$\begin{array}{ccc} & f & \\ (\omega_1, \omega_2, \omega_3) & \longmapsto & \left(\frac{\omega_2\omega_3}{B}, \frac{1}{B}, \omega_3\right) \\ \downarrow \pi & & \downarrow \pi \\ (\omega_1, \omega_2\omega_3, \omega_3) & \longmapsto & \left(\frac{\omega_2\omega_3}{B}, \frac{1}{B}, \frac{\omega_3}{B}\right) \end{array} \quad (7)$$

where  $B = \omega_2 + b\omega_2\omega_3 + c + d\omega_1$ . Here we see again that the exceptional divisor  $\omega_3 = 0$  is mapped by  $f$  on the strict transform of the line at infinity  $L'' = (t = x = 0)$ . Therefore  $E$  is contracted by  $f^2$  onto  $p^-$ .  $\square$

Let us summarize the action of  $f$  at infinity:

$$\begin{array}{ccc} \widetilde{H}_\infty \setminus I_f & \longrightarrow & p^- \\ E \setminus I_f & \longrightarrow & L'' \longrightarrow p^- \end{array}$$

The sequence of indeterminacy sets  $I_{f^n} \subset X$  of  $f^{-n}$  (resp.  $I_{f^{-n}} \subset X$  of  $f^{-n}$ ) is stationary. More precisely, we have the following


**Proposition 2** *For all  $n \geq 2$ ,  $I_{f^n} = I_{f^2} := I^+$  and  $I_{f^{-n}} = I_{f^{-2}} := I^-$ . Each of the sets  $I^+$  and  $I^-$  consists of three curves. Moreover, the set  $I^+ \cap I^-$  is finite with*

$$\#(I^+ \cap I^-) = \begin{cases} 5 & \text{if } (b-c)^2 - 4d \neq 0, \\ 4 & \text{otherwise.} \end{cases}$$

In Figure 2,  $I^+$  is represented in the divisor at infinity  $E \cup \widetilde{H}_\infty$  of  $X$ . The dynamics of the irreducible components of  $I^+$  by  $f$  is indicated by curved arrows.

**Proof Points of indeterminacy in  $E$  for  $f$  and  $f^{-1}$ .** There are points of indeterminacy for  $f$  in  $E$ . To see this, let us first consider the expression of  $f$  in the charts ( $z \neq 0$ ) at input and ( $y \neq 0$ ) at output: we have  $f(x, y, t) = \left(y, \frac{y+byt+ct+dx}{t}, t\right)$ . The expression of  $f$  in the chart ( $z \neq 0, \xi_2 \neq 0$ ) at input and the chart ( $y \neq 0$ ) at output is:

$$\begin{array}{ccc} & f & \\ (\omega_1, \omega_2, \omega_3) & \longmapsto & (\omega_2\omega_3, \omega_2 + b\omega_2\omega_3 + c + d\omega_1, \omega_3) \\ \downarrow \pi & & \downarrow \pi \\ (\omega_1, \omega_2\omega_3, \omega_3) & \longmapsto & (\omega_2\omega_3, \omega_2 + b\omega_2\omega_3 + c + d\omega_1, \omega_3) \end{array} \quad (8)$$



**Fig. 2** Divisor at infinity of  $X$ . The indeterminacy set is represented by straight lines and the action by  $f$  by curved arrows.

Thus there is no point of indeterminacy in the chart  $(Z \neq 0, \xi_2 \neq 0)$ . Moreover, we have seen that  $E \cap \tilde{H}_\infty$  does not contain any point of indeterminacy for  $f$  in the chart  $(Z \neq 0, \xi_1 \neq 0)$ . Now the local expression of  $f$  in the charts  $(X \neq 0)$  at input and  $(Z \neq 0)$  at output is

$$f = \left( \frac{yt}{yz + byt + czt + dt}, \frac{zt}{yz + byt + czt + dt}, \frac{t^2}{yz + byt + czt + dt} \right).$$

The expression of the blow-up in the chart  $\xi_2 \neq 0$  at input and output is:

$$\begin{array}{ccc} & f & \\ (\omega_1, \omega_2, \omega_3) & \longmapsto & \left( \frac{\omega_1 \omega_3}{C}, \frac{\omega_2}{\omega_3}, \frac{\omega_3}{C} \right) \\ \downarrow \pi & & \downarrow \pi \\ (\omega_1 \omega_3, \omega_2, \omega_3) & \longmapsto & \left( \frac{\omega_1 \omega_3}{C}, \frac{\omega_2}{C}, \frac{\omega_3}{C} \right) \end{array} \quad (9)$$

where  $C := \omega_1 \omega_2 + b \omega_1 \omega_3 + c \omega_2 + d$ . Therefore, there is an indeterminacy locus on  $E$ , that we call  $C'^+$ , whose equation is  $(\omega_2 = 0)$  in this chart. By symmetry,  $C'^- := I_{f^{-1}} \cap E$  is given by  $(\omega_2 = 0)$  in the chart  $(z \neq 0, \xi_2 \neq 0)$ . One sees by (9) that  $f(I_f \cap E) \subset E$  and  $f(I_{f^{-1}} \cap E) \subset E$ . Therefore

$$f(C'^+) = C'^- \quad \text{and} \quad f^{-1}(C'^-) = C'^+.$$

**Points of indeterminacy in  $E$  for  $f^2$  and  $f^{-2}$ .** Points in  $E \cap (I_{f^2} \setminus I_f)$  are mapped by  $f$  on  $[0, 1, 0, 0]$ . It follows from (8) that  $\mathcal{C}^+ := E \cap (I_{f^2} \setminus I_f)$  can be expressed in the chart  $(z \neq 0, \xi_2 \neq 0)$  by  $c + d\omega_1 = 0$ . We compute in the same way the expression of  $\mathcal{C}^- := E \cap I_{f^{-2}} \setminus I_{f^{-1}}$  in the chart  $(x \neq 0, \xi_2 \neq 0)$ , namely  $\omega_2 - \omega_1 - b - c\omega_1\omega_3 = 0$ , whereas in the chart  $(x \neq 0, \xi_1 \neq 0)$  it is given by the equation  $\omega_2\omega_3 - 1 - b\omega_3 - c\omega_1\omega_3 = 0$ .

**The set  $I^+ \cap I^-$ .** In the chart  $(z \neq 0, \xi_1 \neq 0)$ , the curve  $\mathcal{C}^+$  in  $E$  corresponds to the equation  $1 + c\omega_3 + d\omega_1\omega_3 = 0$  and the curve  $\mathcal{C}^-$  to  $\omega_3 - \omega_1 - b\omega_1\omega_3 = 0$ . A straightforward computation shows that  $\mathcal{C}^+ \cap \mathcal{C}^-$  has two points in  $E$  which coincide when  $(b - c)^2 - 4d = 0$ .

**Sequences  $I_{f^n}$  and  $I_{f^{-n}}$  are stationary.** Since neither points of  $E \setminus \{\mathcal{C}^+, \mathcal{C}^-\}$  nor points of  $\tilde{H}_\infty \setminus L'$  are mapped by  $f$  in  $I_{f^2}$ , we have  $I_{f^2} = I_{f^3} = I_{f^n}$ ,  $n \geq 2$ . Similarly  $I_{f^{-n}} = I_{f^{-2}}$  for all  $n \geq 2$ .  $\square$

*Remark 1* One can verify that  $f$  and  $f^{-1}$  become algebraically stable after blowing-up the two points  $p^+$  and  $p^-$  where the hyperplane at infinity in  $\mathbb{P}^3$  is contracted by  $f$  and  $f^{-1}$  respectively. Note however that this increases the number of charts to be considered, as well as the dimension of  $H^{1,1}(X, \mathbf{R})$ . This is one reason why we have used a slightly different compactification.

### 3 Canonical invariant classes

The action of  $f^*$  on the divisors at infinity is described by the following

**Lemma 2**  $f^*[\tilde{H}_\infty] = [\tilde{H}_\infty] + [E]$  and  $f^*[E] = [\tilde{H}_\infty]$ .

*Proof* We have  $f(\tilde{H}_\infty \setminus I_f) = p^- \in E$ , and no other hypersurface at infinity is mapped in  $E$ . So

$$f^*[E] = a[\tilde{H}_\infty], \quad a > 0.$$

Now equation (6) shows that  $a = 1$ .

Since  $\tilde{H}_\infty$  and  $E$  are both mapped into  $\tilde{H}_\infty$  by  $f$  we have

$$f^*[\tilde{H}_\infty] = b[\tilde{H}_\infty] + c[E], \quad b, c > 0.$$

The coefficients  $a = b$  and  $c$  are the order of vanishing of  $f$  along  $\tilde{H}_\infty$  and  $[E]$  respectively. By equations (6) and (7) we have  $b = c = 1$ .  $\square$

Thus the action of  $f^*$  induced on  $H^{1,1}(X, \mathbf{R})$  is represented by the matrix  $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$  in the basis  $(\{\tilde{H}_\infty\}, \{E\})$ . By symmetry, the action of  $f^{-1*}$  induced on  $H^{1,1}(X, \mathbf{R})$  is represented by the same matrix. The spectral radius of  $f^*$  (resp.  $f^{-1*}$ ) is equal to the golden mean  $\sigma := \frac{1+\sqrt{5}}{2}$ . The other eigenvalue of  $f^*$  (resp.  $f^{-1*}$ ) is  $\frac{1-\sqrt{5}}{2}$ . Recall that the *dynamical degrees* of  $f$  are given by

$$\lambda_i(f) := \lim_{n \rightarrow +\infty} \left( \int_X f^{n*}(\omega^i) \wedge \omega^{3-i} \right)^{\frac{1}{n}}, \quad i = 1, 2, 3 \quad (10)$$

for a Kähler form  $\omega$  on  $X$  (see [9]). A rational endomorphism is called *cohomologically hyperbolic* if one dynamical degree strictly dominates the others. We have thus shown the following (see also [12]):

**Proposition 3** *The dynamical degrees of  $f$  are*

$$\lambda_1(f) = \lambda_2(f) = \lambda_1(f^{-1}) = \frac{1 + \sqrt{5}}{2}, \quad \lambda_3(f) = \lambda_3(f^{-1}) = 1.$$

*Thus  $f$  is not cohomologically hyperbolic.*

There is a unique cohomology class  $\alpha \in H^{1,1}(X, \mathbf{R})$  (up to a multiplicative constant) that is invariant by  $\frac{1}{\sigma}f^*$  (resp.  $\frac{1}{\sigma}f^{-1*}$ ). The class  $\alpha$  is equal to  $\{E\} + \sigma\{\tilde{H}_\infty\}$ .

We now study the positivity properties of the invariant class  $\alpha$ .

**Proposition 4** *The invariant class  $\alpha \in H^{1,1}(X, \mathbf{R})$  is Kähler.*

*Proof* **We start by showing that  $\alpha$  is semi-positive.** We equip the line bundle  $\mathcal{O}_{\mathbb{P}^3}(H_\infty)$  with a singular positive metric ([1]) whose curvature form  $\omega'$  is smooth outside the line  $L = (Y = T = 0)$  and which has (homogeneous) Lelong number equal to 1 along this projective line. Note that  $\omega'' := \pi^*(\omega') - [E]$  is a smooth semi-positive metric on the line bundle  $\mathcal{O}_X(\tilde{H}_\infty) \simeq \pi^*\mathcal{O}_{\mathbb{P}^3}(H_\infty) \otimes \mathcal{O}_X(-E)$  which is therefore semi-positive. Thus the invariant class  $\alpha = \{\pi^*[H_\infty] + (\sigma - 1)[\tilde{H}_\infty]\}$  is semi-positive.

**The class  $\alpha$  is Kähler.** Let  $\omega_{FS}$  denote the Fubini-Study Kähler form on  $\mathbb{P}^3$ . It is cohomologous to  $[H_\infty]$ . It is classical (see e.g. [7]) that there exists a smooth closed form  $\theta \in -\{E\}$  such that  $\pi^*\omega_{FS} + \varepsilon\theta > 0$  for  $\varepsilon > 0$  sufficiently small. So

$$\alpha = (1 + \varepsilon)\pi^*\{H_\infty\} + \varepsilon\{\theta\} + (\sigma - 1 - \varepsilon)\{\tilde{H}_\infty\}$$

is Kähler.

Here is a slight reformulation: let  $\varphi : \mathbb{P}^3 \rightarrow \mathbb{P}^2$  be the projection on the hyperplane  $(Z = 0) \approx \mathbf{P}^2$  orthogonal to the line  $L = (Y = T = 0)$ ,

$$\varphi([X, Y, Z, T]) = [X, Y, 0, T].$$

Then, after blowing-up  $L$ , the meromorphic application  $\varphi$  induces a holomorphic map  $\Phi : X \rightarrow \mathbf{P}^2$ . We have

$$\begin{aligned} 0 \leq \Phi^*\mathcal{O}_{\mathbf{P}^2}(Z = T = 0) &\approx \pi^*\mathcal{O}_{\mathbf{P}^3}(T = 0) \otimes \mathcal{O}_X(-E) \\ &\approx \pi^*\mathcal{O}_{\mathbf{P}^3}(1) \otimes \mathcal{O}_X(-E) \approx \mathcal{O}_X(\tilde{H}_\infty) \end{aligned}$$

where we have denoted  $\approx$  the isomorphism between holomorphic line bundles. Therefore  $\{\tilde{H}_\infty\} = c_1(\mathcal{O}_X(\tilde{H}_\infty)) \geq 0$ . So  $\{\tilde{H}_\infty\}$  can be represented by a smooth  $(1, 1)$ -form that is positive on  $E$ . Since  $\pi^*\omega_{FS}$  is smooth and positive except on  $E$ , we conclude that  $\alpha = \pi^*\{H_\infty\} + (\sigma - 1)\{\tilde{H}_\infty\}$  is Kähler.  $\square$



*Remark 2* We can describe explicitly in  $\mathbb{C}^3$  the  $(1,1)$ -form  $\omega'$  as follows. The function  $\frac{1}{2} \log(|y|^2 + |t|^2)$  defined in  $\mathbb{C}^3 = (x, y, t)$  is in the Lelong class  $\mathcal{L}(\mathbb{C}^3)$ , i.e. we have asymptotically  $\frac{1}{2} \log(|y|^2 + |t|^2) \leq \frac{1}{2} \log(1 + |x|^2 + |y|^2 + |z|^2) + O(1)$  ([11]). The function  $g := \frac{1}{2} \log(|y|^2 + |t|^2) - \frac{1}{2} \log(1 + |x|^2 + |y|^2 + |t|^2)$  thus extends as a  $\omega_{FS}$ -plurisubharmonic function on  $\mathbb{P}^3$ . This extension (that we call  $g$  again) writes in homogeneous coordinates  $g = \frac{1}{2} \log(|Y|^2 + |T|^2) - \frac{1}{2} \log(|X|^2 + |Y|^2 + |Z|^2 + |T|^2)$  and  $\omega' = \omega_{FS} + dd^c g$ .

We now compute the volume of the class  $\alpha$ . The intermediate computations in the proof will also be useful in the sequel.

**Proposition 5** *The class  $\alpha = \{E\} + \sigma\{\tilde{H}_\infty\}$  has volume*

$$\text{Vol}(\alpha) = \alpha^3 = 1 + \frac{3}{\sigma} > 0.$$

*Proof* A potential of  $\omega \in \alpha$  in  $\mathbb{C}^3$  is  $\frac{1}{2\sigma} \log(|y|^2 + 1) + \frac{1}{2} \log(|x|^2 + |y|^2 + |z|^2 + 1)$ . Set  $u := \frac{1}{2\sigma} \log(|y|^2 + 1)$  and  $v := \frac{1}{2} \log(|x|^2 + |y|^2 + |z|^2 + 1)$ . Then

$$\text{Vol}(\alpha) = \alpha^3 = \int_{\mathbb{C}^3} (dd^c u + dd^c v)^3 = \int_{\mathbb{C}^3} (dd^c v)^3 + 3(dd^c v)^2 \wedge dd^c u = 1 + \frac{3}{\sigma}. \quad \square$$

## 4 Invariant currents

### 4.1 The Green currents $T_f$ and $T_{f^{-1}}$

Recall that  $\omega_{FS}$  is the Fubini-Study form on  $\mathbb{P}^3$  and that  $\omega''$  is a smooth  $(1,1)$ -form on  $X$  cohomologous to  $\tilde{H}_\infty$ . It follows from the previous section that  $\omega := \pi^* \omega_{FS} + (\sigma - 1)\omega''$  is a Kähler form representing the invariant class  $\alpha = \{E\} + \sigma\{\tilde{H}_\infty\}$ . Since  $f^* \alpha = \sigma \alpha$ , it follows from the  $dd^c$ -lemma that there exists an  $\omega$ -plurisubharmonic function  $\phi$  such that  $\frac{1}{\sigma} f^* \omega = \omega + dd^c \phi$ . We normalize  $\phi$  so that  $\sup_X \phi = 0$ .

We refer the reader to [11] for basic properties of  $\omega$ -plurisubharmonic functions. The sequence of positive currents

$$T_n := \frac{1}{\sigma^n} f^{n*} \omega = \omega + dd^c \left( \sum_{i=0}^{n-1} \frac{1}{\sigma^i} \phi \circ f^i \right)$$

converges towards a positive closed current  $T_f$ , since the decreasing sequence of  $\omega$ -plurisubharmonic functions  $\sum_{i=0}^{n-1} \frac{1}{\sigma^i} \phi \circ f^i$  converges in  $L^1(X)$  (it does not converge identically to  $-\infty$  by an argument due to Sibony, see e.g. the proof of Theorem 2.1 in [8]). By construction, the current  $T_f$  is invariant by  $\frac{1}{\sigma} f^*$ .

Note that the "Green function", i.e. the potential of  $T_f$  in  $\mathbf{C}^3$

$$G^+ = \lim_{n \rightarrow +\infty} \frac{1}{\sigma^n} f^{n*} \left( \frac{1}{2\sigma} \log(|y|^2 + 1) + \frac{1}{2} \log(|x|^2 + |y|^2 + |z|^2 + 1) \right) \quad (11)$$

(See Remark 2) is well defined and plurisubharmonic in  $\mathbf{C}^3$ . We construct in the same way the invariant current for  $f^{-1}$ ,

$$T_{f^{-1}} := \lim_{n \rightarrow +\infty} \frac{1}{\sigma^n} (f^{-n})^* \omega.$$

Let  $\mathcal{B}_\infty^+$  (resp.  $\mathcal{B}_\infty^-$ ) denote the basin of attraction of the superattractive point  $p^-$  (resp.  $p^+$ ) at infinity. The following result is an easy adaptation (a special case) of Theorem 2.2 in [10]:

**Proposition 6** *The potential  $G^+ \geq 0$  (resp.  $G^- \geq 0$ ) of the current  $T_f$  (resp.  $T_{f^{-1}}$ ) is pluriharmonic in  $\mathcal{B}_\infty^+$  (resp.  $\mathcal{B}_\infty^-$ ). Moreover*

$$\mathcal{B}_\infty^+ = (G^+ > 0), \quad \mathcal{B}_\infty^- = (G^- > 0)$$

and  $G^+, G^-$  are continuous in  $\mathbf{C}^3$ . We also have  $\text{supp}(T_{f^\pm}) \cap D_\infty = I^\pm$ .

#### 4.2 The invariant current $T_f \wedge T_{f^{-1}}$

The (2, 2)-current  $T_f \wedge T_{f^{-1}}$  is well defined in  $\mathbf{C}^3$  according to [2].

**Proposition 7** *The support of the current  $T_f \wedge T_{f^{-1}}$  clusters at infinity exactly along  $I^+ \cap I^-$ , where it has positive Lelong numbers.*

*Proof* We know by Proposition 6.3 in [4] that  $T_f$  (resp.  $T_{f^{-1}}$ ) has positive Lelong numbers at  $I^+$  (resp.  $I^-$ ). Thus  $T_f \wedge T_{f^{-1}}$  has positive Lelong numbers at  $I^+ \cap I^-$  by Corollary 5.10 in [2]. So  $I^+ \cap I^- \subset \text{supp}(T_f \wedge T_{f^{-1}})$ .

Now take  $x \in D_\infty \setminus I^+ \cap I^-$ . By symmetry we can assume  $x \notin I^+$ . We know by Proposition 6 that  $x \notin \text{supp}(T_f)$ , hence  $x \notin \text{supp}(T_f \wedge T_{f^{-1}})$ . Therefore  $I^+ \cap I^- = \text{supp}(T_f \wedge T_{f^{-1}}) \cap D_\infty$ .  $\square$

The current  $T_f \wedge T_{f^{-1}}$  is invariant by  $f^*$  in  $\mathbf{C}^3$  since

$$f^*(T_f \wedge T_{f^{-1}}) = f^*T_f \wedge f^*T_{f^{-1}} = \sigma T_f \wedge \frac{1}{\sigma} T_{f^{-1}} = T_f \wedge T_{f^{-1}}. \quad (12)$$

Note that this equality does not hold at infinity in  $X$ , because  $f^*(T_f \wedge T_{f^{-1}}) \geq T_f \wedge T_{f^{-1}} + a[\mathcal{C}^+]$ ,  $a > 0$ , since  $\nu(f^*S, z) \geq \nu(S, f(z))$ . Of course  $T_f \wedge T_{f^{-1}}$  is also invariant by  $(f^{-1})^*$  in  $\mathbf{C}^3$ .

Set  $J := \text{supp}(T_f \wedge T_{f^{-1}})$ . Recall that  $J \cap D_\infty$  consists in five points (Proposition 2). Using the same notations as in Proposition 2, we denote by:

- $A^+ \in J \cap E$  the only point in  $\mathcal{C}'^+$ ;

- $A^- \in J \cap E$  the only point in  $\mathcal{C}'^-$ ;
- $B, B' \in J \cap E$  the two points in  $\mathcal{C}^+ \cap \mathcal{C}^-$ ;
- $C \in J \cap H_\infty$  the point such that  $\pi(C) = [0, 1, 0, 0]$ .

We now analyse the action of  $f|_J$  at infinity.

**Proposition 8** *Setting  $f = f|_J$ , we have*

- $f(B) = f^{-1}(B) = f(B') = f^{-1}(B') = C$ ;
- $f(C) = \{A^-, B, B'\}$  and  $f^{-1}(C) = \{A^+, B, B'\}$ ;
- $f(A^+) = A^-$  and  $f^{-1}(A^-) = A^+$ ;
- $f(A^-) = C$  and  $f^{-1}(A^+) = C$ .

*Proof* Since  $f(\mathcal{C}^+) = C$ , the regular points (for  $f$  but not for  $f^2$ )  $B, B', A^- \in \mathcal{C}^+$  are mapped into  $C$  by  $f$ . We also obtain in the same way  $f^{-1}(B) = f^{-1}(B') = f^{-1}(A) = C$ . Since  $A^+ \in \mathcal{C}'^+$ , we have  $f(A^+) \in \mathcal{C}'^- \cap J$ , i.e.  $f(A^+) = A^-$ . A similar argument yields  $f^{-1}(A^-) = A^+$ .

Choose a sequence  $(p_n) \in J^\mathbb{N}$  that converges to  $B$ . Then the sequence  $z_n := f(p_n) \in \text{supp}(T_f \wedge T_{f^{-1}})$  converges to  $C$ . Therefore  $B \in f^{-1}(C)$ , since  $(z_n) \in \text{supp}(T_f \wedge T_{f^{-1}})$ . We prove in the same way that  $B', A' \in f^{-1}(C)$ , and that there are no other possibilities. Thus  $f^{-1}(C) = \{A^+, B, B'\}$ , and similarly we find  $f(C) = \{A^-, B, B'\}$ .  $\square$

For two unbounded sets  $S, S' \in \mathbb{C}^3$ , we write  $S \simeq S'$  if there exists  $R \in \mathbb{R}^+$  such that  $S \cap \{\|z\| > R\} = S' \cap \{\|z\| > R\}$ . This defines an equivalence relation, whose classes will be called **germs at infinity**. Moreover, if we have a partition  $S \cap \{\|z\| > R\} = S_1 \sqcup S_2$  for some  $R > 0$ , we will write  $S \simeq S_1 \sqcup S_2$ .

Let  $J_{A^+}, J_{A^-}, J_C, J_B$  be the germs at infinity of the intersections of  $J$  with disjoint neighborhoods of  $A^+, A^-, C, \{B, B'\}$ , respectively. It follows for instance from the previous proposition that

$$f(J_{A^+}) \simeq J_{A^-} \text{ and } f^{-1}(J_{A^-}) \simeq J_{A^+}. \quad (13)$$

Similarly, Proposition 8 yields

$$J_C \simeq f^{-1}(J_{A^+}) \sqcup f^{-1}(J_B) \simeq f(J_{A^-}) \sqcup f(J_B). \quad (14)$$

We can now describe the dynamics on  $J$  near infinity. Given two sets  $A, B \subset X$ , we write  $A \rightarrow B$  for  $f(A) \simeq B$ .


**Proposition 9** *There exists a partition of  $J|_{\mathbb{C}^3}$  near infinity into nonempty sets*

$$J_{\mathbb{C}^3} \simeq J'_{A^+} \sqcup J'_{A^-} \sqcup J'_C \sqcup J'_B \sqcup J''_{A^+} \sqcup J''_{A^-} \sqcup J''_C \sqcup J''_B,$$

where  $J'_K, J''_K \subset J_K$  for  $K \in \{A^+, A^-, C, B, B'\}$ , such that the following hold

$$J'_{A^+} \rightarrow J'_{A^-} \rightarrow J'_C \rightarrow J'_{A^+},$$

$$J''_{A^+} \rightarrow J''_{A^-} \rightarrow J''_C \rightarrow J''_B \rightarrow J''_C \rightarrow J''_{A^+}.$$



**Fig. 3** Dynamics near infinity on the support of  $T_f \wedge T_{f^{-1}}$ .

Moreover, setting  $J' := J_{A^+}' \sqcup J_{A^-}' \sqcup J_C' \sqcup J_B$  and  $J'' := J_{A^+}'' \sqcup J_{A^-}'' \sqcup J_C'' \sqcup J_B$ , the restriction of  $f$  to  $J'$  or  $J''$  extends continuously to infinity at  $A^+$ ,  $A^-$ ,  $C$ .

Thus the points in  $J$  near infinity are "weakly periodic with period 3 or 5", in the sense that each of them is mapped near itself by  $f^3$  or  $f^5$ . We summarize the above proposition in Figure 3.

*Proof* We set  $J_C''' := f(J_B)$ ,  $J_C'' := f^{-1}(J_B)$ ,  $J_C' = J_C \setminus (J_C'' \cup J_C''')$ .

We will first prove that these three sets form a partition of  $J_C$ . Note that Proposition 8 implies  $f(J_C \setminus J_C'') \subset J_{A^+}$ . It follows from (14) that

$$f(J_C \setminus J_C'') \simeq J_{A^+}. \quad (15)$$

Similarly,  $f^{-1}(J_C \setminus J_C''') \simeq J_{A^-}$ .

Now, by Proposition 8 we have  $f^2(J_{A^-}) \subset J_B \sqcup f(J_C \setminus J_C''')$ . Using (15) and the fact that  $f|_J : J \rightarrow J$  is surjective, we obtain  $f^2(J_{A^-}) \simeq J_B \sqcup J_{A^+}$ .

Rewriting this equation as  $J_{A^-} \simeq f^{-1}(J_C'') \sqcup f^{-2}(J_{A^+})$  yields

$$f^{-1}(J_C'') \subset J_{A^-}.$$

This equation, combined with  $f(J_C''') \subset J_{A^+}$  obtained in the same way, shows that  $J_C''' \cap J_C'' = \emptyset$ . So we have proved that  $J_C \simeq J_C' \sqcup J_C'' \sqcup J_C'''$ .

Set  $J_{A^+}'' := f(J_C''')$ ,  $J_{A^+}' = f(J_C')$ ,  $J_{A^-}'' := f^{-1}(J_C''')$ ,  $J_{A^-}' = f^{-1}(J_C')$ . It follows from the previous paragraph that  $J_{A^+} \simeq J_{A^+}' \sqcup J_{A^+}''$  and  $J_{A^-} \simeq$

$J'_{A^-} \sqcup J''_{A^-}$ . Indeed, the sets  $J'_{A^+}$  and  $J''_{A^+}$  are disjoint since  $J'_C$  and  $J''_C$  are, and the surjectivity of  $f|_J$  shows that  $J'_{A^+} \sqcup J''_{A^+}$  is a partition of  $J_{A^+}$ .

Let us now prove that  $f^3(J_B) \simeq f^{-2}(J_B)$  (this means informally that the previous diagram is "commutative" in a weak sense). We have

$$\begin{aligned} J_{A^+} &= (f^3(J_{A^+}) \cap J_{A^+}) \sqcup (f^3(J'_C) \cap J_{A^+}) \\ &= (f^3(J_{A^+}) \cap J_{A^+}) \sqcup (J''_{A^+} \cap J_{A^+}) \end{aligned} \quad (16)$$

Indeed, a point  $x \in J_{A^+}$  can only come from  $J_{A^+}$  or  $J'_C$  by  $f^3$ , according to Proposition 8. It follows that  $f(J''_{A^+}) \subset J''_{A^-}$ .

Similarly, one gets  $f^{-1}(J''_{A^-}) \supset J''_{A^+}$  and  $f(J'_{A^+}) \subset J'_{A^-}$ .

**The set  $J''$  is nonempty, and the set  $J'$  is nonempty for  $|b| < 1$  and  $|c|$  sufficiently small.** By Proposition 7, the set  $J = J' \sqcup J''$  reaches infinity at  $B$ . Since we have seen that  $J'$  does not reach infinity at  $B$  but  $J''$  does, the set  $J''_B$  is nonempty. Therefore, its images by  $f$  and  $f^2$  are nonempty also. Thus  $J''$  is nonempty.

For proving that  $J'$  is nonempty, it is sufficient to prove that there exists a sufficiently small neighborhood of  $A^-$  in  $X$  that is mapped by  $f^2$  into a neighborhood of  $A^+$ . It will be shown in Remark 4 below that  $J_{A^-} \cap \{(x, y, z) \in \mathbb{C}^3, |x| < \varepsilon, |y| < \varepsilon^2\} \neq \emptyset$ . If  $|c| < \varepsilon^2$  and  $|b| < 1$ , it is straightforward that  $f(x_0, y_0, z_0) \in V_{A^+}$ .  $\square$

**Corollary 1**  $f^{15}|_J : J \rightarrow J$  extends continuously at infinity. Moreover, the points at infinity are fixed by  $f^{30}|_J : J \rightarrow J$ .

*Proof* Proposition 9 shows that  $f^{15}|_J : J \rightarrow J$  extends continuously at infinity, except perhaps at  $B$  and  $B'$ . Let  $J_B \simeq J'_B \sqcup J''_B$  denote the partition of  $J_B$  into the two branches that contain the two points  $B$  and  $B'$ . We can prove by the same type of arguments as those used in the proof of the previous proposition that  $J''_C \simeq f^{-1}(J'_B) \sqcup f^{-1}(J''_B)$  and  $J'''_C \simeq f(J'_B) \sqcup f(J''_B)$ , and that  $f^5|_{J'_B}$  and  $f^5|_{J''_B}$  extend continuously at infinity.

It follows from Proposition 9 that  $f^{15}|_J : J \rightarrow J$  fixes the points  $A^+$ ,  $A^-$ ,  $C$ , and eventually permutes the points  $B$  and  $B'$ . Thus all these points are fixed by  $f^{30}|_J : J \rightarrow J$ .  $\square$

*Remark 3* We deduce from Proposition 6 that  $T_f \wedge T_f = T_{f^{-1}} \wedge T_{f^{-1}} = 0$  in  $\mathbb{C}^3$ , since  $T_f \wedge T_f = dd^c(G^+ T_f)$  and  $G^+ = 0$  on  $\text{supp}(T_f) \cap \mathbb{C}^3$ . Thus neither  $T_f \wedge T_f$  nor  $T_{f^{-1}} \wedge T_{f^{-1}}$  are of any interest.

## 5 A canonical invariant measure

We now go back to dynamics. We study the same family as before, quadratic polynomial automorphisms  $f : \mathbb{C}^3 \rightarrow \mathbb{C}^3$  which are not cohomologically hyperbolic. We would like to construct a canonical invariant probability measure for them. Recall that

$$f(x, y, z) = (y, z, yz + by + cz + dx + e), \quad b, c, d \in \mathbb{C}, |d| > 1$$

induces an algebraically stable automorphism of  $X$ , the blow-up of  $\mathbb{P}^3$  along a suitable projective line, such that  $f^{-1}$  is also algebraically stable with

$$\lambda_1(f) = \lambda_1(f^{-1}) = \frac{1 + \sqrt{5}}{2} > \lambda_3(f) = 1.$$

We have constructed the canonical invariant currents  $T_f, T_{f^{-1}}$  and  $T := \frac{\sigma}{3+\sigma} T_f \wedge T_{f^{-1}}$ . In this section we construct a continuous function  $\phi_\infty$  defined on the support of the current  $T$  with suitable invariance properties and such that it is  $T$ -plurisubharmonic. (Recall that a  $T$ -plurisubharmonic function, associated with a closed positive current  $T$ , is a function  $u : \text{supp}(T) \rightarrow \mathbb{R} \cup \{-\infty\}$ ,  $u \in L^1(|T|)$ , that is the limit on  $\text{supp}(T)$  of a sequence of plurisubharmonic functions defined on some neighborhood of  $\text{supp}(T)$ , decreasing on  $\text{supp}(T)$ .)

We then study the invariant measure  $\mu = (T_f \wedge T_{f^{-1}} \wedge dd^c \phi_\infty)|_{\mathbb{C}^3}$ . We prove that its support is compact and included in the set of points with bounded forward orbits.

We suppose in this section that  $|d| > 1$  in Equation (1). We set

$$T := \frac{\sigma}{3+\sigma} T_f \wedge T_{f^{-1}}.$$

Recall that  $J = \text{supp}(T)$ . We can decompose  $J = K \sqcup W$ , where:

$$K := \{\text{the points of } J \text{ with a bounded forward orbit}\},$$

$$W := J \setminus K.$$

Note that  $K$  is compact. Indeed, Lemma 4 below implies that  $K$  is bounded, and the same lemma implies that  $W$  is open in  $\mathbb{C}^3$  as the basin of attraction of infinity on  $J$ . The set  $W$  can be decomposed again into two "crossed" partitions:

**Lemma 3** *We can decompose  $W = W' \sqcup W''$  in  $\mathbb{C}^3$  with*

$$W' := \{z \in W : f^{3n}(z) \text{ converges towards a point at infinity when } n \rightarrow +\infty\}$$

$$W'' := \{z \in W : f^{5n}(z) \text{ converges towards a point at infinity when } n \rightarrow +\infty\}.$$

$W'$  and  $W''$  are unbounded sets invariant by  $f^3$  and  $f^5$  respectively.

We can also decompose

$$W := W_{A^+} \sqcup W_{A^-} \sqcup W_C \sqcup W_B \sqcup W_{B'},$$

where  $W_{A^+}, W_{A^-}, W_C, W_B, W_{B'}$  are the connected components of  $W$  in  $X$  that reach the points  $A^+, A^-, C, B, B'$  respectively at infinity. These five sets are invariant by  $f^{15}$ .

*Proof* A glance at Proposition 9 shows that  $W = W' \cup W''$ , with the notations of the statement. It remains to prove that  $W' \cap W'' = \emptyset$ . The set  $W' \cap W''$  is stable by  $f^3$  and  $f^5$ , therefore it is periodic with period 1, i.e. it is stable by  $f$ . This set  $W' \cap W''$  is not bounded, otherwise  $W' \cap W'' \subset K$ , which is not possible by definition of  $W$  and  $K$ . Thus  $W' \cap W''$  reaches infinity. Then Proposition 9 prevents  $W' \cap W''$  from being fixed under the action of  $f$ , a contradiction.

Now consider the decomposition  $W = W_{A^+} \cup W_{A^-} \cup W_C \cup W_B \cup W_{B'}$  given by Proposition 9. If  $x \in W_{A^+}$ , then  $f^{15n}(x) \rightarrow A^+$  when  $n \rightarrow +\infty$ . If  $W_{A^+} \cap W_C \neq \emptyset$ , then we would also have  $\lim_{n \rightarrow +\infty} f^{15n}(x) = C$ , which is a contradiction. We prove in the same way that the previous decomposition of  $W$  is a genuine partition.  $\square$

With the notations of Proposition 9, we have  $J'_{A^+}, J'_{A^-}, J'_C \subset W'''$  and  $J''_{A^+}, J''_{A^-}, J''_C \subset W''$ . Our goal is to construct a function  $\phi_\infty : J \rightarrow \mathbb{R}$  with suitable invariant properties, such that the current  $T \wedge dd^c \phi_\infty$  makes sense and yields an invariant positive measure.

### 5.1 Growth control and invariant measure

We will now construct a  $T$ -plurisubharmonic function  $\phi_\infty$  on  $J$  with invariant properties. This will allow us to construct an invariant probability measure. We set  $|T| := T \wedge \omega$  for a Kähler form  $\omega$  and denote by  $\|\cdot\|$  the max norm, and  $\|\cdot\|_+ := \max(\|\cdot\|, 1)$ .

We summarize the construction of  $\phi_\infty$  and of the invariant measure in the following theorem:

**Theorem 1** *Suppose  $|d| > 1$ . There exists a  $T$ -plurisubharmonic function  $\phi_\infty : J \rightarrow \mathbb{R}$  which satisfies the following properties:*

1. *The function  $\phi_\infty$  satisfies  $\phi_\infty \circ f^3 = \phi_\infty + \log |d|$  on  $W'$ ;*
2.  *$\phi_\infty = \log^+ \|z\| + o(1)$  as  $\|z\| \rightarrow +\infty$ ;*
3.  *$(\phi_\infty = -\infty) = K$  and  $|T|(K) = 0$ ;*
4.  *$e^{\phi_\infty}$  is continuous on  $J$ .*

*The measure  $\mu := T \wedge dd^c(\phi_\infty)|_{\mathbb{C}^3}$  is a well-defined  $f^3$ -invariant positive measure with compact support, such that  $\text{supp}(\mu) \subset (\phi_\infty = -\infty)$ .*

*Proof* Let  $V_{A^+}$  be a small neighborhood of the point of indeterminacy  $A^+$ .

**Construction of  $\phi_\infty$ .** It is sufficient to construct the function  $\phi_\infty$  on  $J_{A^+} := K \sqcup W_{A^+}$  and to check the stated properties on this set. We first construct  $\phi_\infty : J_{A^+} \rightarrow [-\infty, +\infty]$  as a limit of a decreasing sequence of functions. Let

$$\psi_n(p) := \log^+ (\|f^{3n}(p)\|) - n \log |d|.$$

Then we define a decreasing sequence on  $J_{A^+}$  by  $\psi'_n := \sup_{i \geq n}^* \psi_i$ , and its limit

$$\phi_\infty := \limsup_{n \rightarrow +\infty}^* \psi_n,$$

where we have denoted by  $\sup^*$  the usc-regularization of the sup. Note now that the limit  $\phi_\infty := \lim_{n \rightarrow +\infty} \psi'_n$  defined on  $W_{A^+} \cap V_{A^+}$  satisfies  $\phi_\infty \circ f^3 = \phi_\infty + \log |d|$ . Indeed,

$$\begin{aligned} \phi_\infty \circ f^3 &= \lim_{n \rightarrow +\infty} \sup_{i \geq n}^* (\psi_i \circ f^3) = \lim_{n \rightarrow +\infty} \sup_{i \geq n}^* (\psi_{i+1}) + \log |d| \\ &= \lim_{n \rightarrow +\infty} (\psi'_{n+1}) + \log |d| = \phi_\infty + \log |d|. \end{aligned} \quad (17)$$

**The sequence  $(\psi_n)$  is locally uniformly bounded on  $W_{A^+} \cap V_{A^+}$ .**

By Lemma 4, it suffices to verify that

$$\sigma_{n,\varepsilon} := \sum_{i=1}^n \log \left( |d| + \frac{\varepsilon}{(|d| - \varepsilon)^{i-1}} \right) - \sum_{i=1}^n \log \left( |d| - \frac{\varepsilon}{(|d| - \varepsilon)^{i-1}} \right) \quad (18)$$

does not blow up. Now

$$\sigma_{n,\varepsilon} = \sum_{i=1}^n \log \left( 1 + \frac{\varepsilon}{|d|(|d| - \varepsilon)^{i-1}} \right) - \sum_{i=1}^n \log \left( 1 - \frac{\varepsilon}{|d|(|d| - \varepsilon)^{i-1}} \right),$$

hence  $(\sigma_{n,\varepsilon})$  is a difference of uniformly convergent series, since

$$\log \left( 1 \pm \frac{\varepsilon}{|d|(|d| - \varepsilon)^{i-1}} \right) \sim \pm \frac{\varepsilon}{|d|(|d| - \varepsilon)^{i-1}}.$$

Thus  $(\psi_n)$  is locally uniformly bounded near infinity. Remark that, up to shrinking  $V_{A^+}$ , we can take  $\varepsilon$  in the previous expression as small as we want: thus  $\forall n \geq 0, \psi'_n(z) - \log^+ \|z\| = o(1)$  as  $\|z\| \rightarrow +\infty$ .

**We have  $\phi_\infty - \log^+ \|\cdot\| = o(1)$  near infinity and the sequence  $(\psi'_n)$  is locally uniformly bounded on  $J$  near infinity.** Fix a neighborhood  $V_{A^+}$  of  $A^+$  and  $\varepsilon > 0$  such that the formula given by Lemma 4 holds in  $V_{A^+} \cap J$ . Equation (18) implies that for all  $n \geq 0$  we have on  $V_{A^+} \cap J$

$$\log^+ \|z\| + c_\varepsilon \leq \psi_n(z) \leq \log^+ \|z\| + c_\varepsilon$$

for some constant  $c_\varepsilon > 0$ . We can take  $c_\varepsilon = \lim_{n \rightarrow +\infty} \sigma_{n,\varepsilon} < +\infty$ , where  $\sigma_{n,\varepsilon}$  is given by Equation (18). Then  $\lim_{\varepsilon \rightarrow 0} c_\varepsilon = 0$ . We also have on  $V_{A^+} \cap J$ , for all  $n \geq 0$ ,

$$\log^+ \|z\| + c_\varepsilon \leq \psi'_n(z) \leq \log^+ \|z\| + c_\varepsilon.$$

Letting  $n \rightarrow +\infty$ , we obtain

$$\log^+ \|z\| + c_\varepsilon \leq \phi_\infty(z) \leq \log^+ \|z\| + c_\varepsilon$$

on  $V_{A^+} \cap J$ . We conclude by shrinking  $V_{A^+}$  and letting  $\varepsilon \rightarrow 0$ .

Thus  $\phi_\infty - \log^+ \|\cdot\| = o(1)$  near infinity. In particular, the sequence  $(\psi'_n)$  is locally uniformly bounded on  $J \cap V_{A^+}$ .

**The function  $\phi_\infty$  is locally bounded on  $W_{A^+}$ .** Indeed,  $\phi_\infty$  is locally bounded near infinity since  $\phi_\infty - \log^+ \|\cdot\| = o(1)$  near infinity. Now let  $x \in W$



and  $N \geq 0$  such that  $f^{3N}(x) \in V_{A^+}$ . Thus  $\phi_\infty(x) = \phi_\infty \circ f^{3N} - N \log |d|$ . Since  $\phi_\infty - \log^+ \|\cdot\| = o(1)$  near infinity, we deduce that  $-\infty < \phi_\infty(x) < +\infty$  on  $W$ . Obviously,  $\phi_\infty|_K = -\infty$ .

**The function  $\phi_\infty$  is continuous on  $W_{A^+}$ .** Choose  $\varepsilon$  and  $V_A^+$  sufficiently small such that  $|\phi_\infty - \log^+ \|\cdot\|| < \varepsilon$  in  $W_{A^+} \cap V_A^+$ . Take  $x, y \in W$ . There exists  $N \geq 0$  such that  $x, y \in f^{-N}(W_{A^+} \cap V_{A^+})$ . Then

$$\begin{aligned} |\phi_\infty(x) - \phi_\infty(y)| &= |\phi_\infty \circ f^{3N}(x) - \phi_\infty \circ f^{3N}(y)| \\ &\leq |\log^+ \|f^{3N}(x)\| - \log^+ \|f^{3N}(y)\|| + 2\varepsilon. \end{aligned}$$

If  $x \rightarrow y$  in  $f^{-N}(W' \cap V_{A^+})$ , we obtain

$$\lim_{x \rightarrow y} |\phi_\infty(x) - \phi_\infty(y)| \leq 2\varepsilon.$$

The conclusion follows by letting  $\varepsilon \rightarrow 0$ .

**The function  $e^{\phi_\infty}$  is continuous on  $K \sqcup W_{A^+}$ .** It is sufficient to prove that  $\phi_\infty(x) \rightarrow -\infty$  when  $x \rightarrow \partial K$ . Let, for  $n \in \mathbb{N}$ ,

$$V_n := f^{-3n}(V_{A^+} \cap W) \setminus f^{-3(n-1)}(V_{A^+} \cap W)$$

be disjoint bounded sets on  $W$ . Then, by Lemma 4,  $W = \cup_{k \in \mathbb{N}} V_k$ . Fix  $k_0 > 1$  an integer. There exists a real number  $M$  such that  $\phi_\infty < M$  on  $V_{k_0}$ . Take  $k \geq k_0$ . For all  $x \in V_k$ , there exists  $N \geq 0$  such that  $f^{3N}(x) \in V_{k_0}$ . Then

$$\phi_\infty(x) = \phi_\infty \circ f^{3N}(x) - N \log |d| < M - N \log |d|.$$

Since  $|d| > 1$ , the conclusion follows by noting that  $N \rightarrow +\infty$  when  $x \rightarrow \partial K$ .

**The measure  $\mu := T \wedge dd^c \phi_\infty$  is well defined in  $\mathbb{C}^3$ .** First note that for all  $k \in \mathbb{R}$ , the function  $\phi_{\infty, k} := \max(\phi_\infty, k) : W' \rightarrow [-\infty, +\infty[$  can be extended by the value  $k$  into a  $T$ -plurisubharmonic function on the whole  $J$ . For all  $k \in \mathbb{R}$ , the current  $T \wedge dd^c \phi_{\infty, k}$  is defined by its action in  $\mathbb{C}^3$  on a test function  $\xi$ :

$$\langle T_f \wedge T_{f^{-1}} \wedge dd^c \phi_{\infty, k}, \xi \rangle := \langle \phi_{\infty, k} T_f \wedge T_{f^{-1}}, dd^c \xi \rangle.$$

Note that its support is included in  $W'$ . We want to define the current  $T \wedge dd^c \phi_\infty$  by its action in  $\mathbb{C}^3$  on a test function  $\xi$ :

$$\langle T_f \wedge T_{f^{-1}} \wedge dd^c \phi_\infty, \xi \rangle := \lim_{k \rightarrow -\infty} \langle \phi_{\infty, k} T_f \wedge T_{f^{-1}}, dd^c \xi \rangle.$$

Since the support of  $T \wedge dd^c \phi_{\infty, k}$  is included in  $W'$ , it suffices to check that  $\phi_\infty \in L_{loc}^1(|T|)$ . Note that this implies  $|T|(\phi_\infty = -\infty) = 0$ . Fix  $R > 0$ . Let  $\xi$  be a test function such that  $dd^c \xi > 0$  on the closed ball  $\overline{B}(0, R)$ . Since

$$\lim_{x \rightarrow \infty, x \in J} \phi_\infty(x) = +\infty,$$

we can suppose that  $\psi'_n \leq 0$  on  $\text{supp}(\xi)$  for some  $N > 0$  and  $n \geq N$ . Then for all  $n \geq N$

$$-\int_{\mathbb{C}^3} |\psi'_n| T \wedge dd^c \xi = \int_{\mathbb{C}^3} \xi T \wedge dd^c \psi'_n \geq c \int_{\mathbb{C}^3} T \wedge dd^c \psi'_n$$

for some  $c > 0$ . We can show that this latter integral is bounded from below by a constant independent of  $n$ .

**The support of  $\mu$  is compact in  $\mathbb{C}^3$ .** To simplify notations we write  $f$  for  $f^3$ . Note that  $f(W \cap V_{A^+}) \subset\subset W \cap V_{A^+}$  by Lemma 4. Since  $f$  is an automorphism, we have for all  $n \geq 0$

$$\int_{W \cap V_{A^+}} \mu = \int_{W \cap V_{A^+}} (f^n)_* \mu = \int_{f^n(W \cap V_{A^+})} \mu.$$

Letting  $n \rightarrow +\infty$ , we see that the previous integral is equal to  $\mu(A^+) = 0$ . For every compact  $Q \subset W$ , there exists  $N \geq 0$  such that  $Q \subset \overline{f^{-N}(W \cap V_{A^+})}$ . Thus  $\mu|_Q = f^{N*} (f_*^N \mu|_Q) = f^{N*} 0 = 0$ . Therefore  $\text{supp}(\mu) \subset \overline{K}$ . We have remarked at the beginning of this section that  $K$  is closed. Thus  $\text{supp}(\mu) \subset K$ .

**The total mass of  $\mu$  is not vanishing.** Denote  $\infty := J \cap E$ , where  $E$  is the exceptional divisor at infinity.

Now let  $A \in \mathbb{R}^- \setminus \{0\}$ . There exists  $n_0 \in \mathbb{N}$  such that  $\phi_n \leq A$  on  $\{\phi_\infty < 2A\} \subset W'$  for  $n \geq n_0$ . Let then  $\delta > 0$  sufficiently small for that  $\phi_n < A$  on  $V_\delta \cap W''$ , where  $V_\delta$  is the  $\delta$ -neighborhood of  $W'$ . Up to diminishing  $\delta$ , we can also suppose that  $U_\delta := V_\delta \cap \{\|z\| < 2\}$  does not intersect  $W''$ . Recall that  $\lim_{x \rightarrow \infty, x \in W'} \phi_\infty(x) - \log^+ \|x\| = 0$ . Then the Comparison Theorem (Corollary 2.7 in [3]) implies that

$$T \wedge dd^c \max(\phi_n, A)(U_\delta) \geq \int_{U_\delta} T \wedge dd^c \log^+ \|z\|^{\frac{1}{2}} > 0$$

for all  $n \geq n_0$ . This latter value does not depend neither on  $\delta$ ,  $A$  nor  $n$ . Thus we can diminish  $A$  towards  $-\infty$  up to diminishing  $\delta$  towards 0, in order to obtain

$$T \wedge dd^c \phi_\infty(U_\delta) = \lim_{A \rightarrow -\infty} \lim_{\delta \rightarrow 0} T \wedge dd^c \max(\phi_n, A)(U_\delta) \geq \int_{U_\delta} T \wedge dd^c \log^+ \|z\|^{\frac{1}{2}} > 0. \quad \square$$

We now come to the main technical estimate used in the proof above. Recall that  $J := \text{supp}(T_f \wedge T_{f^-})$  decomposes into three sets  $J = K \sqcup W' \sqcup W''$  invariant by  $f^{15}$ , where  $K \subset \mathbb{C}^3$  is compact and  $W', W''$  are sets that reach infinity in four or five points and are periodic under  $f$  with period 3 and 5 respectively.

**Lemma 4** *Suppose  $|d| > 1$ . For all  $\varepsilon > 0$ , there exists a neighborhood  $V_\varepsilon$  of the divisor at infinity such that for all  $p \in W' \cap V_\varepsilon$  and for all  $n \geq 1$ ,*

$$\|p\|_+ \prod_{i=1}^n \left( |d| - \frac{\varepsilon}{(|d| - \varepsilon)^{i-1}} \right) \leq \|f^{3n}(p)\|_+ \leq \|p\|_+ \prod_{i=1}^n \left( |d| + \frac{\varepsilon}{(|d| - \varepsilon)^{i-1}} \right).$$

In particular,  $W' \cap E$  is attracting for  $f^3$  on  $W'$ , in the sense that

$$f^3(W' \cap V_\varepsilon) \subset\subset (W' \cap V_\varepsilon).$$

*Proof* For  $\varepsilon \in ]0, d[$ , denote  $V_{A^+, \varepsilon} := \{|yz + by + cz| < \varepsilon|dx|\}$ ,  $V_{A^-, \varepsilon} := \{|xy + bx + cy| < \varepsilon|dz|\}$ , and  $V_{C, \frac{\varepsilon}{d}} := \{|xz + by + cz| < \varepsilon|dy|\}$ .

**The sets  $V_{A^+, \varepsilon}$  and  $V_{A^-, \varepsilon}$  are neighborhoods of  $A^+$  and  $A^-$  respectively.** The chart  $T = 0$  of  $\mathbb{P}^3(\mathbb{C})$  is parametrized by  $(\frac{y}{x}, \frac{z}{x}, \frac{1}{x})$ . Recall that  $\pi(A^+) = p^+ = [1, 0, 0, 0]$  is located at the origin of this chart. One can check that a neighborhood of  $A^+$ , centered at  $A^+$ , is parametrized by  $(y, z/x, 1/x)$ , and deduce that  $\{|y| < \varepsilon, |z/x| < \varepsilon\} \subset V_{A^+, \varepsilon}$  is a neighborhood of  $A^+$ . We prove exactly by the same arguments that  $V_{A^-, \varepsilon}$  is a neighborhood of  $A^-$ . In order to simplify notations, let us denote  $(x_0, y_0, z_0) \underset{\varepsilon}{\approx} (x_1, y_1, z_1)$  if there exist  $\delta_x, \delta_y, \delta_z \in ]-\varepsilon, \varepsilon[$  such that  $(x_0, y_0, z_0) = ((1+\delta_x)x_1, (1+\delta_y)y_1, (1+\delta_z)z_1)$ . Then it is straightforward that  $\forall (x_0, y_0, z_0) \in V_{A^+, \varepsilon}$ ,

$$f(x_0, y_0, z_0) \underset{\varepsilon}{\approx} (y_0, z_0, dx_0) \in V_{A^-, \frac{\varepsilon}{d}}. \quad (19)$$

Hence, denoting by  $\|\cdot\|$  the max norm,

$$|d - \varepsilon|\|p\|_+ \leq \|f^3(p)\| \leq |d + \varepsilon|\|p\|_+.$$

Thanks to Proposition 9, we can iterate these inequalities in order to obtain the inequalities of the statement, for  $p \in W' \cap (V_{A^+, \varepsilon} \cup V_{A^-, \varepsilon} \cup V_{C, \varepsilon})$ .  $\square$

*Remark 4* Let  $\varepsilon' \varepsilon'' \in ]0, \varepsilon[$ . By the proof of the previous lemma,  $\{|z| < \varepsilon'|dx|, |y| < \varepsilon''\}$  extends in  $X$  in a neighborhood of  $A^+$ . Then Equation (19) implies that

$$f(\{|z| < \varepsilon'|dx|, |y| < \varepsilon''\} \cap J) \subset \{|x| < \varepsilon'', |y| < \varepsilon'|dz|\} \cap J. \quad (20)$$

On the other hand

$$f(\{|z| < \varepsilon'|dx|, |y| < \varepsilon''\} \cap J) \subset \{|x| < \varepsilon|dz|, |y| < \varepsilon\} \cap J \quad (21)$$

for  $\varepsilon', \varepsilon''$  sufficiently small, since the right hand set in Equation (21) is a neighborhood of  $A^-$  in  $X$ . Then Equations (20) and (21) give

$$f(\{|z| < \varepsilon'|dx|, |y| < \varepsilon''\} \cap J) \subset \{|x| < \varepsilon'', |y| < \varepsilon\} \cap J.$$

We conclude that  $\{|x| < \varepsilon, |y| < \varepsilon'\} \cap J \neq \emptyset$ . In the same way,  $\{|z| < \varepsilon, |y| < \varepsilon''\} \cap J \neq \emptyset$ .

## 5.2 The measure $\mu$ is canonical

We push our analysis further. Let us denote by  $\log^A$  a function defined on  $J$  by  $\log^A := \max(A, \log)$  on  $W' \cup K$ , for some constant  $A \in \mathbb{R}$  sufficiently large so that  $K \subset U := \{\log^A \|z\| = A\} \cap J$ , and by  $\log^A := A$  on  $W''$ . We have

**Theorem 2** *Suppose  $|d| > 1$ . Then*

$$\mu = \lim_{n \rightarrow +\infty} T \wedge dd^c \log^A \|f^{3n}\| = \lim_{n \rightarrow +\infty} T \wedge dd^c \left( \frac{1}{n} \sum_{i=0}^{n-1} \log^A \|f^i\| \right)$$

*in the weak sense of currents.*

*Proof* Let us show the first equality.

**Preliminaries.** Define on  $J$  for all  $n \geq 0$

$$\phi'_n := \log^A \|f^{3n}\| - \sum_{k=1}^n \log \left( |d| + \frac{\varepsilon}{(|d| - \varepsilon)^{k-1}} \right). \quad (22)$$

**The sequence  $(\phi'_n)$  is decreasing near infinity.** It results from Lemma 4 that we have on  $J$  near infinity, for all  $n \geq 1$ ,

$$\|f^{3n}\| \leq \|f^{3(n-1)}\| \cdot \left( |d| + \frac{\varepsilon}{(|d| - \varepsilon)^{3(n-1)}} \right).$$

Thus for all  $n \geq 1$ ,

$$\log^A \|f^{3n}\| - \log \left( |d| + \frac{\varepsilon}{(|d| - \varepsilon)^{n-1}} \right) \leq \log^A \|f^{3(n-1)}\|.$$

Therefore, the sequence  $(\phi'_n)$  is decreasing on  $J$  near infinity. (Note that since infinity is attracting on  $W$ , for all compact  $Q$  in  $W$  there exists  $N \geq 0$  such as  $(\phi'_n)$  is decreasing on  $Q$  for  $n \geq N$ ).

**The sequence  $(\phi'_n)$  is decreasing everywhere.** The sequence  $(\phi'_n)$  is decreasing outside  $V := f^{-1}(U) \subset U$ . We claim that the sequence  $\phi'_n$  is in fact decreasing on  $W'$  (therefore on  $J$ ).

Indeed, take  $x \in W'$  and  $n \in \mathbb{N}$ . If  $f^n(x) \in V$ , then  $\log^A \|f^{n+1}(x)\| = \log^A \|f^n(x)\| = 0$ , therefore  $\phi'_{n+1} < \phi'_n$ . If  $x \notin V$ , then  $\phi'_{n+1}(x) \leq \phi'_n(x)$ .

As in the proof of Theorem 1 we set  $\psi_n := \log^+ \|f^{3n}\| - n \log |d|$ .

**The sequence  $\psi_n$  is quasi-decreasing.** We have seen in Theorem 1 that the difference  $\phi'_n - \psi_n$  on  $J$  is uniformly bounded. Thus there exists a constant  $c$  such that

$$\lim_{n \rightarrow +\infty} \phi'_n = \limsup_{n \rightarrow +\infty}^* \phi'_n = \limsup_{n \rightarrow +\infty}^* \psi_n + c = \phi_\infty + c.$$

Therefore, when  $n \rightarrow +\infty$ ,

$$T \wedge dd^c \log^+ \|f^{3n}\| = T \wedge dd^c \phi'_n \rightarrow T \wedge dd^c (\phi_\infty + c) = \mu.$$

□

**Corollary 2** *The support of the measure  $\mu$  is included in  $\overline{W} \cap K$ .*

*Proof* With the previous notations, for all  $n \geq 0$ ,

$$\text{supp} \left( T \wedge dd^c \log^A \| f^{3n} \| \right) \subset W.$$

It follows that  $\text{supp} \left( \lim_{n \rightarrow +\infty} T \wedge dd^c \log^+ \| f^{3n} \| \right) \subset \overline{W}$ . Recall finally that  $\text{supp}(\mu) \subset K$  (by Theorem 1). □

## References

1. Demailly J.-P., Singular Hermitian Metrics on Positive Line Bundles, Proceedings of the Bayreuth Conference on Complex Algebraic Varieties, April 2-6, 1990, Lecture Notes in Math., 1507, Springer (1992).
2. Demailly J.-P., Monge-Ampère operators, Lelong numbers and intersection theory, Complex analysis and geometry, pp. 115–193, Univ. Ser. Math., Plenum, New York (1993).
3. Dabbek K., Elkhadhra F. Capacité associée à un courant positif fermé, Doc. Math., 1, pp. 469–486 (2006).
4. Diller J., Favre C., Dynamics of bimeromorphic maps of surfaces. Amer. J. Math., 123, no. 6, pp. 1135–1169 (2001).
5. Dinh T.-C., Lawrence M.G., Polynomial hulls and positive currents, Ann. Fac. Sci. Toulouse Math., (6) 12, no. 3, pp. 317–334 (2003).
6. Fornæss J., Sibony N., Complex Dynamics in Higher Dimension I, Asterisque, 222, pp. 201–231 (1994).
7. Griffith P., Harris J., Principles of algebraic geometry. Reprint of the 1978 original. Wiley Classics Library, John Wiley and Sons, Inc., New York (1994).
8. Guedj V., Equidistribution towards the Green current, Bull. Soc. Math. France, 131, no. 3, pp. 359–372 (2003).
9. Guedj V., Propriétés Ergodiques des Applications Rationnelles, Panoramas et synthèses, 30, pp. 97–202 (2010).
10. Guedj V., Sibony N., Dynamics of Polynomial Automorphisms of  $\mathbb{C}^k$ , Ark. Mat., 40, pp. 207–243 (2002).
11. V. Guedj, A. Zeriahi, Intrinsic capacities on compact Kähler manifolds, J. Geom. Anal., 15, no. 4, pp. 607–639 (2005).
12. Maegawa K., Quadratic polynomial automorphisms of dynamical degree golden ratio of  $\mathbb{C}^3$ , Ergodic Theory Dynam. Systems, 21, no. 3, pp. 823–832 (2001).
13. Protin, F., Pluripotential theory on the support of closed positive currents and applications to dynamics in  $\mathbb{C}^n$ , Annali di Matematica (2019). <https://doi.org/10.1007/s10231-019-00851-y>