


Computational Science
Laboratory Technical Report
CSL-TR-19-10
December 2, 2019

Mahesh Narayanamurthi & Adrian
Sandu

*“Efficient implementation of
partitioned stiff exponential
Runge-Kutta methods”*

Computational Science Laboratory
“Compute the Future!”

Computer Science Department
Virginia Polytechnic Institute and State University
Blacksburg, VA 24060
Phone: (540)-231-2193
Fax: (540)-231-6075
Email: maheshnm@vt.edu, sandu@cs.vt.edu
Web: <http://cs1.cs.vt.edu>



Efficient implementation of partitioned stiff exponential Runge-Kutta methods

Mahesh Narayanamurthi^a, Adrian Sandu^a

^a*Computational Science Laboratory, Department of Computer Science, Virginia Tech, Blacksburg, VA 24060*

Abstract

Multiphysics systems are driven by multiple processes acting simultaneously, and their simulation leads to partitioned systems of differential equations. This paper studies the solution of partitioned systems of differential equations using exponential Runge-Kutta methods. We propose specific multiphysics implementations of exponential Runge-Kutta methods satisfying stiff order conditions that were developed in [Hochbruck et al., SISC, 1998] and [Luan and Osterman, JCAM, 2014]. We reformulate stiffly-accurate exponential Runge-Kutta methods in a way that naturally allows of the structure of multiphysics systems, and discuss their application to both component and additively partitioned systems. The resulting partitioned exponential methods only compute matrix functions of the Jacobians of individual components, rather than the Jacobian of the full, coupled system. We derive modified formulations of particular methods of order two, three and four, and apply them to solve a partitioned reaction-diffusion problem. The proposed methods retain full order for several partitionings of the discretized problem, including by components and by physical processes.

Keywords: Exponential integrators, stiff order conditions, partitioned time discretization methods.

Email addresses: maheshnm@vt.edu (Mahesh Narayanamurthi), sandu@cs.vt.edu (Adrian Sandu)

Contents

1	Introduction	4
2	Partitioned systems of differential equations	4
3	Exponential Runge-Kutta methods and W order conditions	5
4	Split exponential Runge-Kutta schemes for component partitioned systems	6
4.1	Convergence considerations for component partitioned methods	7
5	Split exponential Runge-Kutta schemes for additively partitioned systems	8
5.1	An alternative formulation of exponential Runge-Kutta methods	8
5.2	PEXPRK methods	10
5.3	Stability considerations	12
6	Particular methods	14
6.1	Second order partitioned exponential Runge-Kutta scheme	14
6.2	Third order partitioned exponential Runge-Kutta scheme	15
6.3	Fourth order partitioned exponential Runge-Kutta scheme	16
7	Computational considerations	17
8	Results	18
8.1	Gray-Scott equations	18
8.2	Component partitioning by chemical species	19
8.3	Component partitioning by splitting the spatial domain	20
8.4	Additive partitioning by physical process	21
8.5	Partitioning by physics with reaction treated explicitly, and diffusion treated exponentially	21
9	Conclusions	22

1. Introduction

Implicit methods are commonly used to solve stiff systems of ODEs since explicit methods, although cheaper per timestep, suffer significantly from stability restrictions. For large systems implicit methods become expensive as they involve the solution of (non-)linear systems of algebraic equations at each timestep. Widely used Krylov-based iterative solvers can be slow to converge in the absence of a good preconditioner [34]. For non-linear problems, obtaining a preconditioner can be challenging.

Multi-methods for time integration, such as implicit-explicit (IMEX) [6–8, 10, 43, 44, 46–49, 51] address the cost challenge by treating implicitly only the stiff components of the system, and the rest of the system explicitly. In a multi-physics system, with two or more physics based partitions, the implicit method is applied to the stiff partition and the explicit method to the non-stiff partition. IMEX methods can be inefficient when applied to systems where both partitions are stiff. Implicit-Implicit methods [4, 11, 12, 14, 36, 52] treat implicitly each component, but are less stable compared to a monolithic implicit integrator [4].

Exponential time integrators solve the linear part of a problem exactly [19, 27], are more stable than explicit methods, and can be cheaper than implicit methods per time step [22, 29]. Implicit-Exponential methods [9, 26, 32, 50] that treat one partition implicitly and the other partition exponentially have been studied in the recent years [9, 26, 32].

Our recent work [31] focuses on partitioned methods that solve each component of the system with a different exponential scheme. In [31] we developed the classical order condition theory for partitioned exponential methods constructed in the EPIRK framework [40, 41] using a GARK structure [14, 36], as well as constructed in the Rosenbrock-exponential framework [17, 20]. We found that partitioned methods satisfying the classical order conditions are amenable to certain computational optimizations that are unavailable to unpartitioned methods, can be more stable than unpartitioned methods for some experimental settings. However, as the stiffness of the coupled system increases, unpartitioned methods remain superior to partitioned methods.

In this work we study the implementation of exponential Runge–Kutta methods that satisfy stiff order conditions [18, 21, 23–25] on partitioned systems of differential equations. Special choices of the linear operator lead to partitioned exponential schemes that only compute matrix functions of the Jacobians of individual components, rather than the Jacobian of the full, coupled system. We reformulate stiffly-accurate exponential Runge–Kutta methods [18] in a way that naturally allows of the structure of multiphysics systems, and discuss their application to both component and additively partitioned systems.

The remainder of the paper is organized as follows. Section 2 discusses component and additively partitioned systems of differential equations. Section 3 reviews the exponential Runge–Kutta methods and stiff order conditions developed in [18, 21, 23–25]. Implementation of split exponential Runge–Kutta schemes for component partitioned systems is discussed in Section 4, and for additively partitioned systems in Section 5. Coefficients for particular methods satisfying the stiff order conditions are given, in the transformed and partitioned formulations, in Section 6. Considerations related to computational cost of partitioned exponential integrators are given in Section 7. Numerical experiments and results are discussed in Section 8, and conclusions are drawn in Section 9.

2. Partitioned systems of differential equations

In this paper we are concerned with solving partitioned initial value problems arising in multiphysics simulations. Additively partitioned systems

$$u' = f(u) = f^{\{1\}}(u) + f^{\{2\}}(u), \quad u(t_n) = u_n \in \mathbb{R}^N, \quad (1)$$

are characterized by a full right-hand side f that is the sum of two components $f^{\{1\}}$ and $f^{\{2\}}$, with Jacobians

$$f' = J^{\{1\}} + J^{\{2\}}, \quad J^{\{1\}} := \frac{\partial f^{\{1\}}}{\partial u}, \quad J^{\{2\}} := \frac{\partial f^{\{2\}}}{\partial u}. \quad (2)$$

Component partitioned systems split the state vector into components v and w , and have the form

$$y' = \begin{bmatrix} v' \\ w' \end{bmatrix} = \begin{bmatrix} f^{\{1\}}(v, w) \\ f^{\{2\}}(v, w) \end{bmatrix} = f(y), \quad f'(y) = \begin{bmatrix} J^{\{1\}} & \frac{\partial f^{\{1\}}}{\partial w} \\ \frac{\partial f^{\{2\}}}{\partial v} & J^{\{2\}} \end{bmatrix}, \quad J^{\{1\}} := \frac{\partial f^{\{1\}}}{\partial v}, \quad J^{\{2\}} := \frac{\partial f^{\{2\}}}{\partial w}. \quad (3)$$

Using the variable transformation $u = v + w$ [43, 44] the additively partitioned system (1) can be written in the component partitioned form (3):

$$y' = \begin{bmatrix} v' \\ w' \end{bmatrix} = \begin{bmatrix} f^{\{1\}}(v + w) \\ f^{\{2\}}(v + w) \end{bmatrix} = f(y), \quad J := f'(y) = \begin{bmatrix} J^{\{1\}} & J^{\{1\}} \\ J^{\{2\}} & J^{\{2\}} \end{bmatrix}. \quad (4)$$

We note that by adding the two coupled equations (4) we recover the original ODE (1).

3. Exponential Runge-Kutta methods and W order conditions

Application of exponential methods to solve a general ODE system,

$$y' = f(y), \quad y(t_n) = y_n \in \mathbb{R}^N, \quad (5)$$

is based on splitting the right hand side into a linear term with a fixed matrix L and a nonlinear reminder g :

$$y' = f(y) = L \cdot y + g(y), \quad L \approx f'(y), \quad g(y) := f(y) - L \cdot y, \quad (6)$$

where L captures the stiffness of the right hand side and therefore needs to approximate well the stiff directions of the Jacobian $f'(y)$.

An exponential Runge-Kutta method [18, 21, 23] applied to the semi-linearly split system (6) reads:

$$Y_i = y_n + h c_i \varphi_1(c_i h L) f(y_n) + h \sum_{j=2}^{i-1} a_{i,j}(h L) \left(g(Y_j) - g(y_n) \right), \quad i = 1, \dots, s, \quad (7)$$

$$y_{n+1} = y_n + h \varphi_1(h L) f(y_n) + h \sum_{j=2}^s b_j(h L) \left(g(Y_j) - g(y_n) \right).$$

Here, y_n is the current solution, Y_i are the internal stages of the method, s is the number of internal stages in the method, y_{n+1} is the numerical solution after one step, φ_1 is the analytic function with index $k = 1$ in the sequence

$$\varphi_0(z) = \exp(z), \quad \varphi_k(z) = \int_0^1 e^{(1-\theta)z} \cdot \frac{\theta^{k-1}}{(k-1)!} d\theta = \sum_{i=0}^{\infty} \frac{z^i}{(k+i)!}, \quad k \geq 1, \quad (8)$$

$$\varphi_{k+1}(z) = \frac{\varphi_k(z) - \varphi_k(0)}{z}, \quad \varphi_k(0) = \frac{1}{k!}.$$

The coefficients $a_{i,j}$ and b_j are analytic functions, typically linear combinations of φ_k (and scaled versions), and c_i are the abscissae corresponding to the internal stages. These coefficients can be expressed in a Butcher tableau form [5, Chapter 2] as follows:

$$\begin{array}{c|c} \mathbf{c} & \mathbf{A} \\ \hline & \mathbf{b}^T \end{array}$$

where the $a_{i,j}$ coefficients are arranged in the form of a matrix \mathbf{A} , the c_i coefficients in vector \mathbf{c} and b_j coefficients in row vector \mathbf{b}^T .

Stiff order conditions for exponential Runge–Kutta methods are given in [25, Table 1], and are repeated here in Table 1 (up to order four) for convenience. These order conditions hold for any choice of the matrix \mathbf{L} . The functions $\psi_{i,j}(h\mathbf{L})$ are defined as:

$$\psi_{i,j}(h\mathbf{L}) := \sum_{k=2}^{j-1} a_{j,k}(h\mathbf{L}) \frac{c_k^{j-1}}{(j-1)!} - c_j^i \varphi_i(c_j h\mathbf{L}). \quad (9)$$

There is one minor notational difference. In the original paper [25], conditions 3b, 4b, and 4c of Table 1 are written without any assumptions made about the Jacobian of \mathbf{g} , and therefore \mathbf{g}' is an arbitrary square matrix. Here we replace \mathbf{g}' with $\mathbf{J} - \mathbf{L}$ following equations (13), (12).

In [21, 23–25] the following assumptions are made.

Assumption 1. *There exist constants M and ω such that*

$$\|e^{t\mathbf{L}}\| \leq M e^{t\omega}, \quad t \geq 0. \quad (10)$$

As a consequence, the functions $\varphi_k(h\mathbf{L})$ (8) as well as the coefficients $a_{i,j}(h\mathbf{L})$ and $b_j(h\mathbf{L})$ are bounded operators.

Assumption 2. *The equation (6) admits a sufficiently smooth solution $y : [0, T] \rightarrow \mathbb{R}^N$. The nonlinear function \mathbf{g} is sufficiently many times (Fréchet) differentiable in a neighborhood of the solution, with uniformly bounded derivatives.*

It is shown in [21, 23–25] that, if Assumptions 1 and 2 are satisfied, then a method (7) that satisfies the order conditions of Table 1 up to order p computes a numerical solution y_n that converges to the exact solution with order p .

4. Split exponential Runge–Kutta schemes for component partitioned systems

Application of exponential methods to the component partitioned system (3) requires a linear-nonlinear split of the form (6) where $\mathbf{L} \approx \mathbf{f}'$ along the stiff subspace. The matrix \mathbf{L} has the same dimension as \mathbf{f}' , and in general has no particular structure. Evaluation of matrix functions $\varphi_k(h\mathbf{L})$ is computationally expensive.

In this paper we seek to leverage the partitioned structure of the system (3) in order to build computationally efficient implementations of exponential Runge-Kutta methods.

Specifically, we consider a linear-nonlinear splitting (6) of the component partitioned system (3) of the form

$$y' = \begin{bmatrix} v' \\ w' \end{bmatrix} = \begin{bmatrix} \mathbf{f}^{\{1\}}(v, w) \\ \mathbf{f}^{\{2\}}(v, w) \end{bmatrix} = \begin{bmatrix} \mathbf{L}^{\{1\}} \cdot v + \mathbf{g}^{\{1\}}(v, w) \\ \mathbf{L}^{\{2\}} \cdot w + \mathbf{g}^{\{2\}}(v, w) \end{bmatrix} = \mathbf{L} \cdot y + \mathbf{g}(y). \quad (11)$$

where $\mathbf{L}^{\{1\}}$ and $\mathbf{L}^{\{2\}}$ are linear operators that contain the stiffnesses of $\mathbf{f}^{\{1\}}$ and $\mathbf{f}^{\{2\}}$, respectively, $\mathbf{g}^{\{1\}}$ and $\mathbf{g}^{\{2\}}$ are the corresponding non-linear remainder functions. This splitting is of the form (6) with

$$\mathbf{L} := \begin{bmatrix} \mathbf{L}^{\{1\}} & \mathbf{0} \\ \mathbf{0} & \mathbf{L}^{\{2\}} \end{bmatrix}, \quad \mathbf{g}(v, w) := \begin{bmatrix} \mathbf{g}^{\{1\}}(v, w) \\ \mathbf{g}^{\{2\}}(v, w) \end{bmatrix} := \begin{bmatrix} \mathbf{f}^{\{1\}}(v, w) - \mathbf{L}^{\{1\}} \cdot v \\ \mathbf{f}^{\{2\}}(v, w) - \mathbf{L}^{\{2\}} \cdot w \end{bmatrix}. \quad (12)$$

The Jacobian of the non-linear remainder, \mathbf{g} , has the structure:

$$\mathbf{g}'(y) = \mathbf{f}'(y) - \mathbf{L} = \begin{bmatrix} \mathbf{J}^{\{1\}} - \mathbf{L}^{\{1\}} & \frac{\partial \mathbf{f}^{\{1\}}}{\partial w} \\ \frac{\partial \mathbf{f}^{\{2\}}}{\partial v} & \mathbf{J}^{\{2\}} - \mathbf{L}^{\{2\}} \end{bmatrix}. \quad (13)$$

Remark. Applying φ_k to the block-diagonal matrix \mathbf{L} from (12) is equivalent to applying the matrix function to each individual block:

$$\varphi_k(h\mathbf{L}) = \begin{bmatrix} \varphi_k(h\mathbf{L}^{\{1\}}) & \mathbf{0} \\ \mathbf{0} & \varphi_k(h\mathbf{L}^{\{2\}}) \end{bmatrix}. \quad (14)$$

This idea is central to obtaining computational optimizations for partitioned-exponential methods that are unavailable in the case of unpartitioned methods.

No.	Order	Order Condition
1.	1	$\sum_{j=1}^s b_j(h\mathbf{L}) = \varphi_1(h\mathbf{L}),$
2a.	2	$\sum_{j=2}^s b_j(h\mathbf{L}) c_j = \varphi_2(h\mathbf{L}),$
2b.	2	$\sum_{j=1}^s a_{i,j}(h\mathbf{L}) = c_i \varphi_1(h\mathbf{L}), \quad i = 2, \dots, s,$
3a.	3	$\sum_{j=2}^s b_j(h\mathbf{L}) \frac{c_j^2}{2} = \varphi_3(h\mathbf{L}),$
3b.	3	$\sum_{j=2}^s b_j(h\mathbf{L}) (\mathbf{J} - \mathbf{L}) \psi_{2,j}(h\mathbf{L}) = 0,$
4a.	4	$\sum_{j=2}^s b_j(h\mathbf{L}) \frac{c_j^3}{3!} = \varphi_4(h\mathbf{L}),$
4b.	4	$\sum_{j=2}^s b_j(h\mathbf{L}) (\mathbf{J} - \mathbf{L}) \psi_{3,j}(h\mathbf{L}) = 0.$
4c.	4	$\sum_{j=2}^s b_j(h\mathbf{L}) (\mathbf{J} - \mathbf{L}) \sum_{k=2}^{j-1} a_{jk}(h\mathbf{L}) (\mathbf{J} - \mathbf{L}) \psi_{2,k}(h\mathbf{L}) = 0.$
4d.	4	$\sum_{j=2}^s b_j(h\mathbf{L}) c_j \mathbf{K} \psi_{2,j}(h\mathbf{L}) = 0.$

Table 1: Stiff order conditions for exponential Runge–Kutta methods (up to order four) [25], where \mathbf{L} and \mathbf{K} are arbitrary matrices.

The exponential Runge-Kutta scheme (7) applied to (11) leads to the following efficient implementation:

$$\begin{bmatrix} V_i \\ W_i \end{bmatrix} = \begin{bmatrix} v_n + h c_i \varphi_1(c_i h \mathbf{L}^{\{1\}}) \mathbf{f}^{\{1\}}(v_n, w_n) + h \sum_{j=2}^{i-1} a_{i,j}(h \mathbf{L}^{\{1\}}) \left(\mathbf{g}^{\{1\}}(V_j, W_j) - \mathbf{g}^{\{1\}}(v_n, w_n) \right) \\ w_n + h c_i \varphi_1(c_i h \mathbf{L}^{\{2\}}) \mathbf{f}^{\{2\}}(v_n, w_n) + h \sum_{j=2}^{i-1} a_{i,j}(h \mathbf{L}^{\{2\}}) \left(\mathbf{g}^{\{2\}}(V_j, W_j) - \mathbf{g}^{\{2\}}(v_n, w_n) \right) \end{bmatrix}, \quad (15)$$

$$\begin{bmatrix} v_{n+1} \\ w_{n+1} \end{bmatrix} = \begin{bmatrix} v_n + h \varphi_1(h \mathbf{L}^{\{1\}}) \mathbf{f}^{\{1\}}(v_n, w_n) + h \sum_{j=2}^{i-1} b_j(h \mathbf{L}^{\{1\}}) \left(\mathbf{g}^{\{1\}}(V_j, W_j) - \mathbf{g}^{\{1\}}(v_n, w_n) \right) \\ w_n + h \varphi_1(h \mathbf{L}^{\{2\}}) \mathbf{f}^{\{2\}}(v_n, w_n) + h \sum_{j=2}^{i-1} b_j(h \mathbf{L}^{\{2\}}) \left(\mathbf{g}^{\{2\}}(V_j, W_j) - \mathbf{g}^{\{2\}}(v_n, w_n) \right) \end{bmatrix}.$$

Remark. In (15) the matrix functions of $\mathbf{L}^{\{1\}}$ are applied to $\mathbf{f}^{\{1\}}$, $\mathbf{g}^{\{1\}}$, and the matrix functions of $\mathbf{L}^{\{2\}}$ are applied to $\mathbf{f}^{\{2\}}$, $\mathbf{g}^{\{2\}}$.

Remark. The formulation (15) can be extended to any number of partitions.

4.1. Convergence considerations for component partitioned methods

Following [21, 23–25] we extend Assumption 1 as follows.

Assumption 3. *There exist constants $M^{\{1\}}$, $M^{\{2\}}$, $\omega^{\{1\}}$, and $\omega^{\{2\}}$, such that*

$$\|e^{t\mathbf{L}^{\{1\}}}\| \leq M^{\{1\}} e^{t\omega^{\{1\}}}, \quad \|e^{t\mathbf{L}^{\{2\}}}\| \leq M^{\{2\}} e^{t\omega^{\{2\}}}, \quad t \geq 0. \quad (16)$$

Consequently, Assumption 1 holds for the linear operator \mathbf{L} defined in (12).

We also require that Assumption 2 holds. We are particularly interested in the case where $\mathbf{L}^{\{1\}} = \mathbf{J}^{\{1\}}$ and $\mathbf{L}^{\{2\}} = \mathbf{J}^{\{2\}}$. This implies that the off-diagonal blocks of the Jacobian in equation (13) are Lipschitz continuous, with moderate size Lipschitz constants:

$$\begin{aligned} \frac{\partial \mathbf{g}^{\{1\}}(v, w)}{\partial w} &= \frac{\partial \mathbf{f}^{\{1\}}(v, w)}{\partial w}, \left\| \frac{\partial \mathbf{f}^{\{1\}}}{\partial w}(v_a, w_a) - \frac{\partial \mathbf{f}^{\{1\}}}{\partial w}(v_b, w_b) \right\| \leq \ell_1 (\|v_a - v_b\| + \|w_a - w_b\|), \\ \frac{\partial \mathbf{g}^{\{2\}}(v, w)}{\partial v} &= \frac{\partial \mathbf{f}^{\{2\}}(v, w)}{\partial v}, \left\| \frac{\partial \mathbf{f}^{\{2\}}}{\partial v}(v_a, w_a) - \frac{\partial \mathbf{f}^{\{2\}}}{\partial v}(v_b, w_b) \right\| \leq \ell_2 (\|v_a - v_b\| + \|w_a - w_b\|). \end{aligned}$$

If each component of partitioned system (11) has a dynamics that is stiff, but the coupling terms are non-stiff, then Assumption 2 holds and the convergence of the split scheme (15) follows from the general theory [21, 23–25]. Increasing the stiffness in the coupling terms leads to a decrease of the time step required for convergence.

5. Split exponential Runge-Kutta schemes for additively partitioned systems

Application of a standard exponential Runge-Kutta method (7) to the additively partitioned system (1) requires matrix operations $\varphi_k(h\mathbf{L})$ on a linear operator that approximates the sum of the individual Jacobians (2)

$$\mathbf{L} \approx \mathbf{J}^{\{1\}} + \mathbf{J}^{\{2\}}.$$

We seek to develop formulations of the scheme (7) that only perform matrix operations $\varphi_k(h\mathbf{L}^{\{1\}})$, $\varphi_k(h\mathbf{L}^{\{2\}})$ on individual Jacobian approximations $\mathbf{L}^{\{1\}} \approx \mathbf{J}^{\{1\}}$, $\mathbf{L}^{\{2\}} \approx \mathbf{J}^{\{2\}}$. In order to achieve this we first consider a reformulation of exponential Runge-Kutta methods (7).

5.1. An alternative formulation of exponential Runge-Kutta methods

We now derive an alternative formulation of (7) that will prove useful in the implementation of partitioned schemes.

Theorem 5.1. *The exponential Runge-Kutta method (7) can be rewritten as:*

$$\begin{aligned} Y_i &= y_n + h \alpha_{i,1}(h\mathbf{L}) f(y_n) + h \sum_{j=2}^{i-1} \alpha_{i,j}(h\mathbf{L}) \left(f(Y_j) - f(y_n) \right), \quad i = 1, \dots, s, \\ y_{n+1} &= y_n + h \beta_1(h\mathbf{L}) f(y_n) + h \sum_{j=2}^s \beta_j(h\mathbf{L}) \left(f(Y_j) - f(y_n) \right). \end{aligned} \tag{17}$$

We denote:

$$\mathbf{E}_{2:s,2:s}(z) = \left(\mathbf{I}_{s-1} + z \mathbf{A}_{2:s,2:s}(z) \right)^{-1}.$$

Since $\mathbf{A}_{2:s,2:s}$ is strictly lower triangular, $\mathbf{E}_{2:s,2:s}$ is lower triangular and involves only multiplications of elements of \mathbf{A} . The coefficients of (17) are defined as:

$$\begin{aligned} \alpha_{1,1:s}(z) &= \mathbf{0}, \quad \alpha_{2:s,1}(z) = \mathbf{E}_{2:s,2:s}(z) \begin{bmatrix} c_2 \varphi_1(c_2 z) \\ \vdots \\ c_s \varphi_1(c_s z) \end{bmatrix}, \quad \alpha_{2:s,2:s}(z) = \mathbf{E}_{2:s,2:s}(z) \mathbf{A}_{2:s,2:s}(z), \\ \beta_1(z) &= \varphi_1(z) - z \sum_{j=2}^s \mathbf{b}_j(z) \alpha_{j,1}(z), \quad \beta_{2:s}^T(z) = \mathbf{b}_{2:s}^T(z) \mathbf{E}_{2:s,2:s}(z). \end{aligned}$$

Proof. Using equations (6), the method (7) can be written, after a rearrangement of the internal stages, as:

$$Y_i - y_n = h c_i \varphi_1(c_i h \mathbf{L}) f(y_n) + h \sum_{j=2}^{i-1} a_{i,j}(h \mathbf{L}) \left(f(Y_j) - f(y_n) \right) - \sum_{j=2}^{i-1} a_{i,j}(h \mathbf{L}) h \mathbf{L} \left(Y_j - y_n \right),$$

$$y_{n+1} = y_n + h \varphi_1(h \mathbf{L}) f(y_n) + h \sum_{j=2}^s b_j(h \mathbf{L}) \left(f(Y_j) - f(y_n) \right) - \sum_{j=2}^s b_j(h \mathbf{L}) h \mathbf{L} \left(Y_j - y_n \right).$$

Since $Y_1 = y_n$, we have that $c_1 = 0$, and rest of the internal stages can be stacked and rearranged after multiplication by $\mathbf{E}_{2:s,2:s}$ as follows:

$$\begin{bmatrix} Y_2 - y_n \\ \vdots \\ Y_s - y_n \end{bmatrix} = h \mathbf{E}_{2:s,2:s}(h \mathbf{L}) \begin{bmatrix} c_2 \varphi_1(c_2 h \mathbf{L}) \\ \vdots \\ c_s \varphi_1(c_s h \mathbf{L}) \end{bmatrix} f(y_n) + h \mathbf{E}_{2:s,2:s}(h \mathbf{L}) \mathbf{A}_{2:s,2:s}(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}$$

$$= h \boldsymbol{\alpha}_{2:s,1}(h \mathbf{L}) f(y_n) + h \boldsymbol{\alpha}_{2:s,2:s}(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}.$$

The final one step solution reads:

$$y_{n+1} = y_n + h \varphi_1(h \mathbf{L}) f(y_n) + h \mathbf{b}_{2:s}^T(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix} - \mathbf{b}_{2:s}^T(h \mathbf{L}) h \mathbf{L} \begin{bmatrix} Y_2 - y_n \\ \vdots \\ Y_s - y_n \end{bmatrix}.$$

Substituting the expression for $[Y_2 - y_n, \dots, Y_s - y_n]^T$ from above:

$$y_{n+1} = y_n + h \varphi_1(h \mathbf{L}) f(y_n) + h \mathbf{b}_{2:s}^T(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}$$

$$- h \mathbf{b}_{2:s}^T(h \mathbf{L}) \boldsymbol{\alpha}_{2:s}(h \mathbf{L}) h \mathbf{L} f(y_n) - h \mathbf{b}_{2:s}^T(h \mathbf{L}) h \mathbf{L} \boldsymbol{\alpha}_{2:s,2:s}(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}$$

After collecting the terms,

$$y_{n+1} = y_n + h \left[\varphi_1(h \mathbf{L}) - \mathbf{b}_{2:s}^T(h \mathbf{L}) \boldsymbol{\alpha}_{2:s,1}(h \mathbf{L}) h \mathbf{L} \right] f(y_n)$$

$$+ h \mathbf{b}_{2:s}^T(h \mathbf{L}) \left[\mathbf{I} - h \mathbf{L} \boldsymbol{\alpha}_{2:s,2:s}(h \mathbf{L}) \right] \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}.$$

It can be shown that:

$$\mathbf{I} - z \boldsymbol{\alpha}_{2:s,2:s}(z) = \mathbf{I} - z \left(\mathbf{I} + z \mathbf{A}_{2:s,2:s}(z) \right)^{-1} \mathbf{A}_{2:s,2:s}(z) = \left(\mathbf{I} + z \mathbf{A}_{2:s,2:s}(z) \right)^{-1}.$$

Consequently, the final one step solution is written as:

$$y_{n+1} = y_n + h \boldsymbol{\beta}_1(h \mathbf{L}) f(y_n) + h \boldsymbol{\beta}_{2:s}^T(h \mathbf{L}) \begin{bmatrix} f(Y_2) - f(y_n) \\ \vdots \\ f(Y_s) - f(y_n) \end{bmatrix}.$$

□

Example 1. For example, for $s = 2$, we have:

$$\boldsymbol{\alpha}(z) = \begin{bmatrix} 0 & 0 \\ c_2 \varphi_1(c_2 z) & 0 \end{bmatrix}, \quad \boldsymbol{\beta}(z) = \begin{bmatrix} \varphi_1(z) - b_2(z) z c_2 \varphi_1(c_2 z) & \\ & b_2(z) \end{bmatrix}.$$

For $s = 3$, we have:

$$\boldsymbol{\alpha}(z) = \begin{bmatrix} 0 & 0 & 0 \\ c_2 \varphi_1(c_2 z) & 0 & 0 \\ c_3 \varphi_1(c_3 z) - c_2 z a_{3,2}(z) \varphi_1(c_2 z) & a_{3,2}(z) & 0 \end{bmatrix},$$

$$\boldsymbol{\beta}(z) = \begin{bmatrix} \varphi_1(z) - b_3(z) (z c_3 \varphi_1(c_3 z) - c_2 z^2 a_{3,2}(z) \varphi_1(c_2 z)) - b_2(z) z c_2 \varphi_1(c_2 z) & \\ -b_3(z) z a_{3,2}(z) + b_2(z) & \\ & b_3(z) \end{bmatrix}.$$

5.2. PEXPRK methods

We next build partitioned-exponential Runge–Kutta methods using the *transformed* exponential Runge–Kutta method formulation (17). The *transformed* scheme (17) applied to the additively partitioned system in component form (4) reads:

$$U_i := V_i + W_i, \quad Y_i := \begin{bmatrix} V_i \\ W_i \end{bmatrix},$$

$$\begin{bmatrix} V_i \\ W_i \end{bmatrix} = \begin{bmatrix} v_n \\ w_n \end{bmatrix} + h \begin{bmatrix} \boldsymbol{\alpha}_{i,1}(h\mathbb{L}^{\{1\}}) \mathbf{f}^{\{1\}}(y_n) \\ \boldsymbol{\alpha}_{i,1}(h\mathbb{L}^{\{2\}}) \mathbf{f}^{\{2\}}(y_n) \end{bmatrix} + h \sum_{j=2}^{i-1} \begin{bmatrix} \boldsymbol{\alpha}_{i,j}(h\mathbb{L}^{\{1\}}) \left(\mathbf{f}^{\{1\}}(Y_j) - \mathbf{f}^{\{1\}}(y_n) \right) \\ \boldsymbol{\alpha}_{i,j}(h\mathbb{L}^{\{2\}}) \left(\mathbf{f}^{\{2\}}(Y_j) - \mathbf{f}^{\{2\}}(y_n) \right) \end{bmatrix}, \quad i = 1, \dots, s,$$

$$\begin{bmatrix} v_{n+1} \\ w_{n+1} \end{bmatrix} = \begin{bmatrix} v_n \\ w_n \end{bmatrix} + h \begin{bmatrix} \boldsymbol{\beta}_1(h\mathbb{L}^{\{1\}}) \mathbf{f}^{\{1\}}(y_n) \\ \boldsymbol{\beta}_1(h\mathbb{L}^{\{2\}}) \mathbf{f}^{\{2\}}(y_n) \end{bmatrix} + h \sum_{j=2}^s \begin{bmatrix} \boldsymbol{\beta}_j(h\mathbb{L}^{\{1\}}) \left(\mathbf{f}^{\{1\}}(Y_j) - \mathbf{f}^{\{1\}}(y_n) \right) \\ \boldsymbol{\beta}_j(h\mathbb{L}^{\{2\}}) \left(\mathbf{f}^{\{2\}}(Y_j) - \mathbf{f}^{\{2\}}(y_n) \right) \end{bmatrix}.$$

Here V_i and W_i are the internal stages of the corresponding component ODEs, Y_i are the internal stages of the method obtained by adding the internal stages and the final one step solutions of the methods applied to the component ODEs, v_n and w_n are the current solutions of the component ODEs, y_n is the current solution of the original ODE, v_{n+1} and w_{n+1} are the final one step solutions of the methods applied to the component ODEs.

By adding the internal stages and the final solutions, the above expressions lead to the partitioned Exponential Runge–Kutta formulation.

Definition 5.1. A partitioned Exponential Runge–Kutta (PEXP RK) method reads:

$$U_i = u_n + h \boldsymbol{\alpha}_{i,1}(h\mathbb{L}^{\{1\}}) \mathbf{f}^{\{1\}}(u_n) + h \boldsymbol{\alpha}_{i,1}(h\mathbb{L}^{\{2\}}) \mathbf{f}^{\{2\}}(u_n)$$

$$+ h \sum_{j=2}^{i-1} \boldsymbol{\alpha}_{i,j}(h\mathbb{L}^{\{1\}}) \left(\mathbf{f}^{\{1\}}(U_j) - \mathbf{f}^{\{1\}}(u_n) \right) + h \sum_{j=2}^{i-1} \boldsymbol{\alpha}_{i,j}(h\mathbb{L}^{\{2\}}) \left(\mathbf{f}^{\{2\}}(U_j) - \mathbf{f}^{\{2\}}(u_n) \right), \quad i = 1, \dots, s,$$

$$u_{n+1} = u_n + h \boldsymbol{\beta}_1(h\mathbb{L}^{\{1\}}) \mathbf{f}^{\{1\}}(u_n) + h \boldsymbol{\beta}_1(h\mathbb{L}^{\{2\}}) \mathbf{f}^{\{2\}}(u_n)$$

$$+ h \sum_{j=2}^s \boldsymbol{\beta}_j(h\mathbb{L}^{\{1\}}) \left(\mathbf{f}^{\{1\}}(U_j) - \mathbf{f}^{\{1\}}(u_n) \right) + h \sum_{j=2}^s \boldsymbol{\beta}_j(h\mathbb{L}^{\{2\}}) \left(\mathbf{f}^{\{2\}}(U_j) - \mathbf{f}^{\{2\}}(u_n) \right). \tag{18}$$

Observe that method (18) applies matrix functions of $\mathbb{L}^{\{1\}}$ only to $\mathbf{f}^{\{1\}}$, and matrix functions of $\mathbb{L}^{\{2\}}$ only to $\mathbf{f}^{\{2\}}$. In this regard, it is a partitioned method.

Remark (Systems with multiple partitions). For a P -way partitioned system (1)

$$u' = f(u) = \sum_{p=1}^P f^{\{p\}}(u), \quad u(t_n) = u_n, \quad (19)$$

the PEXPRK method (18) becomes:

$$\begin{aligned} U_i &= u_n + h \sum_{p=1}^P \alpha_{i,1}(h\mathbf{L}^{\{p\}}) f^{\{p\}}(u_n) \\ &\quad + h \sum_{p=1}^P \sum_{j=2}^{i-1} \alpha_{i,j}(h\mathbf{L}^{\{p\}}) \left(f^{\{p\}}(U_j) - f^{\{p\}}(u_n) \right), \quad i = 1, \dots, s, \\ u_{n+1} &= u_n + h \sum_{p=1}^P \beta_1(h\mathbf{L}^{\{p\}}) f^{\{p\}}(u_n) + h \sum_{p=1}^P \sum_{j=2}^s \beta_j(h\mathbf{L}^{\{p\}}) \left(f^{\{p\}}(U_j) - f^{\{p\}}(u_n) \right). \end{aligned} \quad (20)$$

Theorem 5.2 (Partitioned method written in terms of residuals). *The PEXPRK method (20) can be written in terms of residual differences as follows:*

$$\begin{aligned} U_i &= u_n + h \sum_{p=1}^P \bar{\alpha}_{i,1}^{\{p\}}(h\mathbf{L}^{\{p\}}) f^{\{p\}}(u_n) \\ &\quad + h \sum_{p=1}^P \sum_{j=2}^{i-1} \bar{\alpha}_{i,j}^{\{p\}}(h\mathbf{L}^{\{p\}}) \left(g^{\{p\}}(U_j) - g^{\{p\}}(u_n) \right), \quad i = 1, \dots, s, \\ u_{n+1} &= u_n + h \sum_{p=1}^P \bar{\beta}_1^{\{p\}}(h\mathbf{L}^{\{p\}}) f^{\{p\}}(u_n) + h \sum_{p=1}^P \sum_{j=2}^s \bar{\beta}_j^{\{p\}}(h\mathbf{L}^{\{p\}}) \left(g^{\{p\}}(U_j) - g^{\{p\}}(u_n) \right). \end{aligned} \quad (21)$$

The coefficients depend on all matrices $\mathbf{L}^{\{1\}}$ to $\mathbf{L}^{\{P\}}$ and are computed as follows:

$$\begin{aligned} \bar{\mathbf{E}}(h\mathbf{L}^{\{1\}} \dots h\mathbf{L}^{\{P\}}) &= \left(\mathbf{I}_{s-1} - \sum_{p=1}^P h\mathbf{L}^{\{p\}} \alpha_{2:s,2:s}(h\mathbf{L}^{\{p\}}) \right)^{-1}, \\ \bar{\alpha}_{2:s,1:s}^{\{p\}}(h\mathbf{L}^{\{1\}} \dots h\mathbf{L}^{\{P\}}) &:= \bar{\mathbf{E}}(h\mathbf{L}^{\{1\}} \dots h\mathbf{L}^{\{P\}}) \alpha_{2:s,1:s}(h\mathbf{L}^{\{p\}}), \\ \left(\bar{\beta}_{1:s}^{\{p\}} \right)^T (h\mathbf{L}^{\{1\}} \dots h\mathbf{L}^{\{P\}}) &:= \beta_{1:s}^T(h\mathbf{L}^{\{p\}}) + \left(\sum_{q=1}^P h\mathbf{L}^{\{q\}} \beta_{2:s}^T(h\mathbf{L}^{\{q\}}) \right) \bar{\alpha}_{2:s,1:s}^{\{p\}} \end{aligned} \quad (22)$$

Since $\alpha_{2:s,2:s}$ are strictly lower triangular matrices, a formal inversion of the left hand side matrix $\bar{\mathbf{E}}$ is possible, and the inverse elements are products of the entries of the matrices $\alpha(h\mathbf{L}^{\{p\}})$.

Proof. We seek to rewrite (18) in terms of residuals. For this we first write (18) in matrix form:

$$\begin{bmatrix} U_2 - u_n \\ \vdots \\ U_s - u_n \end{bmatrix} = \sum_{p=1}^P h \alpha_{2:s,1}(h\mathbf{L}^{\{p\}}) f^{\{p\}}(u_n) + \sum_{p=1}^P h \alpha_{2:s,2:s}(h\mathbf{L}^{\{p\}}) \begin{bmatrix} f^{\{p\}}(U_2) - f^{\{p\}}(u_n) \\ \vdots \\ f^{\{p\}}(U_s) - f^{\{p\}}(u_n) \end{bmatrix}.$$

Next, we subtract back the linear terms from both sides:

$$\left(\mathbf{I}_{s-1} - \sum_{p=1}^P h\mathbf{L}^{\{p\}} \alpha_{2:s,2:s}(h\mathbf{L}^{\{p\}}) \right) \cdot \begin{bmatrix} U_2 - u_n \\ \vdots \\ U_s - u_n \end{bmatrix} = \sum_{p=1}^P h \alpha_{2:s,1:s}(h\mathbf{L}^{\{p\}}) \begin{bmatrix} f^{\{p\}}(u_n) \\ g^{\{p\}}(U_2) - g^{\{p\}}(u_n) \\ \vdots \\ g^{\{p\}}(U_s) - g^{\{p\}}(u_n) \end{bmatrix}.$$

This, together with the coefficient definition (22), gives the stage equations in (21).

From (18) we have:

$$u_{n+1} = u_n + \sum_{p=1}^P h \beta^T(hL^{\{p\}}) \begin{bmatrix} f^{\{p\}}(u_n) \\ g^{\{p\}}(U_2) - g^{\{p\}}(u_n) \\ \vdots \\ g^{\{p\}}(U_s) - g^{\{p\}}(u_n) \end{bmatrix} + \left(\sum_{p=1}^P hL^{\{p\}} \beta_{2:s}^T(hL^{\{p\}}) \right) \begin{bmatrix} U_2 - u_n \\ \vdots \\ U_s - u_n \end{bmatrix},$$

which, together with the coefficient definition (22), gives the next solution equation in (21). \square

Example 2. For example, for $s = 3$, we have:

$$\bar{\mathbf{E}} = \begin{bmatrix} \sum_{k=1}^P hL^{\{k\}} a_{3,2}(hL^{\{k\}}) & 0 \\ 1 & 1 \end{bmatrix},$$

$$\bar{\alpha}^{\{p\}} = \begin{bmatrix} 0 & 0 & 0 \\ c_2 \varphi_1(c_2 hL^{\{p\}}) & 0 & 0 \\ c_3 \varphi_1(c_3 hL^{\{p\}}) + \left(\sum_{k \neq p} hL^{\{k\}} a_{3,2}(hL^{\{k\}}) \right) c_2 \varphi_1(c_2 hL^{\{p\}}) & a_{3,2}(hL^{\{p\}}) & 0 \end{bmatrix},$$

$$\bar{\beta}^{\{p\}} = \beta_{1:s}^T(hL^{\{p\}}) + \left(\sum_{q=1}^P hL^{\{q\}} \beta_{2:s}^T(hL^{\{q\}}) \right) \bar{\alpha}_{2:s,1:s}^{\{p\}},$$

where

$$\beta(z) = \begin{bmatrix} \varphi_1(z) - b_3(z) (z c_3 \varphi_1(c_3 z) - c_2 z^2 a_{3,2}(z) \varphi_1(c_2 z)) - b_2(z) z c_2 \varphi_1(c_2 z) \\ -b_3(z) z a_{3,2}(z) + b_2(z) \\ b_3(z) \end{bmatrix}.$$

5.3. Stability considerations

The stability analysis of PEXPRK methods is more complex due to the splitting of the Jacobian $L^{\{1\}} + L^{\{2\}}$ into separate $L^{\{1\}}$ and $L^{\{2\}}$ parts. We discuss the cause of this complexity, and illustrate with the stability analysis of the PEXPRK method build using exponential Euler as the base scheme.

Consider the additive partitioned system in component form (4) and perform a linear-nonlinear splitting of the form (11):

$$y' = \begin{bmatrix} v' \\ w' \end{bmatrix} = \begin{bmatrix} f^{\{1\}}(v+w) \\ f^{\{2\}}(v+w) \end{bmatrix} = \begin{bmatrix} L^{\{1\}} \cdot v + g^{\{1\}}(v,w) \\ L^{\{2\}} \cdot w + g^{\{2\}}(v,w) \end{bmatrix} = L \cdot y + g(y), \quad (23)$$

where $L^{\{1\}}$ and $L^{\{2\}}$ are linear operators that contain the stiffnesses of $f^{\{1\}}$ and $f^{\{2\}}$, respectively, $g^{\{1\}}$ and $g^{\{2\}}$ are the corresponding non-linear remainder functions. We have

$$L := \begin{bmatrix} L^{\{1\}} & 0 \\ 0 & L^{\{2\}} \end{bmatrix}, \quad g(y) := \begin{bmatrix} g^{\{1\}}(v,w) \\ g^{\{2\}}(v,w) \end{bmatrix} := \begin{bmatrix} f^{\{1\}}(v+w) - L^{\{1\}} \cdot v \\ f^{\{2\}}(v+w) - L^{\{2\}} \cdot w \end{bmatrix}. \quad (24)$$

The standard linear-nonlinear splitting (6) of (23) using the full Jacobian evaluated at y_n as the linear operator (4) leads to

$$y' = \begin{bmatrix} v' \\ w' \end{bmatrix} = \begin{bmatrix} f^{\{1\}}(v+w) \\ f^{\{2\}}(v+w) \end{bmatrix} = \begin{bmatrix} L^{\{1\}} \cdot (v+w) + g^{\text{standard},\{1\}}(v,w) \\ L^{\{2\}} \cdot (v+w) + g^{\text{standard},\{2\}}(v,w) \end{bmatrix}. \quad (25)$$

We make the following additional assumption:

Assumption 4. The system (23) obeys Assumptions 1 and 2. In particular, there is an initial condition v_0, w_0 with $y_0 = [v_0; w_0]$ such that each component equation (25) admits a sufficiently smooth solution $v, w : [0, T] \rightarrow \mathbb{R}^N$.

The following remark shows why the PEXP stability analysis is more challenging.

Remark. Higher derivatives of our remainder (24) and of the standard remainder (25) coincide

$$\mathbf{g}^{(k)}(y) = (\mathbf{g}^{\text{standard}})^{(k)}(y), \quad k \geq 2,$$

and are therefore uniformly bounded if Assumption 4 holds. The first derivatives, however, require a closer look. The Jacobian of the standard remainder (25) is

$$(\mathbf{g}^{\text{standard}})'(y) = \begin{bmatrix} \mathbf{J}^{\{1\}}(y) - \mathbf{L}^{\{1\}} & \mathbf{J}^{\{1\}}(y) - \mathbf{L}^{\{1\}} \\ \mathbf{J}^{\{2\}}(y) - \mathbf{L}^{\{2\}} & \mathbf{J}^{\{2\}}(y) - \mathbf{L}^{\{2\}} \end{bmatrix}, \quad (26)$$

and is uniformly bounded by Assumption 4. The Jacobian of our nonlinear reminder (24)

$$\mathbf{g}'(y) = \begin{bmatrix} \mathbf{J}^{\{1\}}(y) - \mathbf{L}^{\{1\}} & \mathbf{J}^{\{1\}}(y) \\ \mathbf{J}^{\{2\}}(y) & \mathbf{J}^{\{2\}}(y) - \mathbf{L}^{\{2\}} \end{bmatrix} \quad (27)$$

is not, due to the presence of the stiff Jacobians (and not Jacobian differences) in off-diagonal positions.

While the residual Jacobian is not nicely bounded, its product with an analytical matrix function is.

Remark. The product of a function $h\varphi_k(h\mathbf{L})$ with $k \geq 1$ with the residual Jacobian reads:

$$\begin{aligned} h\varphi_k(h\mathbf{L}) \mathbf{g}'(u) &= \begin{bmatrix} h\varphi_k(h\mathbf{L}^{\{1\}}) & 0 \\ 0 & h\varphi_k(h\mathbf{L}^{\{2\}}) \end{bmatrix} \cdot \begin{bmatrix} \mathbf{J}^{\{1\}} - \mathbf{L}^{\{1\}} & \mathbf{J}^{\{1\}} \\ \mathbf{J}^{\{2\}} & \mathbf{J}^{\{2\}} - \mathbf{L}^{\{2\}} \end{bmatrix} \\ &= \begin{bmatrix} h\varphi_k(h\mathbf{L}^{\{1\}}) (\mathbf{J}^{\{1\}} - \mathbf{L}^{\{1\}}) & h\varphi_k(h\mathbf{L}^{\{1\}}) \mathbf{J}^{\{1\}} \\ h\varphi_k(h\mathbf{L}^{\{2\}}) \mathbf{J}^{\{2\}} & h\varphi_k(h\mathbf{L}^{\{2\}}) (\mathbf{J}^{\{2\}} - \mathbf{L}^{\{2\}}) \end{bmatrix}. \end{aligned} \quad (28)$$

The diagonal blocks are uniformly bounded by Assumption 4. The off-diagonal blocks are also uniformly bounded, since each term on the right of

$$\begin{aligned} h\varphi_k(h\mathbf{L}^{\{1\}}) \mathbf{J}^{\{1\}} &= h\varphi_k(h\mathbf{L}^{\{1\}}) (\mathbf{J}^{\{1\}} - \mathbf{L}^{\{1\}}) + \varphi_k(h\mathbf{L}^{\{1\}}) h\mathbf{L}^{\{1\}} \\ &= h\varphi_k(h\mathbf{L}^{\{1\}}) (\mathbf{J}^{\{1\}} - \mathbf{L}^{\{1\}}) + \left(\varphi_{k-1}(h\mathbf{L}^{\{1\}}) - \frac{1}{(k-1)!} \right), \end{aligned} \quad (29)$$

is uniformly bounded.

Consider the exponential Euler method applied to (23)–(24):

$$\begin{aligned} \begin{bmatrix} v_{n+1} \\ w_{n+1} \end{bmatrix} &= \begin{bmatrix} v_n \\ w_n \end{bmatrix} + \begin{bmatrix} h\varphi_1(h\mathbf{L}^{\{1\}}) & 0 \\ 0 & h\varphi_1(h\mathbf{L}^{\{2\}}) \end{bmatrix} \cdot \begin{bmatrix} \mathbf{f}^{\{1\}}(v_n + w_n) \\ \mathbf{f}^{\{2\}}(v_n + w_n) \end{bmatrix} \\ &= \begin{bmatrix} v_n \\ w_n \end{bmatrix} + \begin{bmatrix} h\varphi_1(h\mathbf{L}^{\{1\}}) (\mathbf{L}^{\{1\}}(v_n + w_n) + \mathbf{g}^{\text{standard}\{1\}}(v_n + w_n)) \\ h\varphi_1(h\mathbf{L}^{\{2\}}) (\mathbf{L}^{\{2\}}(v_n + w_n) + \mathbf{g}^{\text{standard}\{2\}}(v_n + w_n)) \end{bmatrix}, \end{aligned}$$

and, after adding the component equations together,

$$\begin{aligned} u_{n+1} &= u_n + h\varphi_1(h\mathbf{L}^{\{1\}}) (\mathbf{L}^{\{1\}} u_n + \mathbf{g}^{\text{standard}\{1\}}(u_n)) + h\varphi_1(h\mathbf{L}^{\{2\}}) (\mathbf{L}^{\{2\}} u_n + \mathbf{g}^{\text{standard}\{2\}}(u_n)) \\ &= (e^{h\mathbf{L}^{\{1\}}} + e^{h\mathbf{L}^{\{2\}}} - 1) u_n + h\varphi_1(h\mathbf{L}^{\{1\}}) \mathbf{g}^{\text{standard}\{1\}}(u_n) + h\varphi_1(h\mathbf{L}^{\{2\}}) \mathbf{g}^{\text{standard}\{2\}}(u_n). \end{aligned} \quad (30)$$

Assume, for simplicity, that h , $\mathbf{L}^{\{1\}}$, and $\mathbf{L}^{\{2\}}$ remain the same throughout the integration, and let

$$\mathbf{M} := e^{h\mathbf{L}^{\{1\}}} + e^{h\mathbf{L}^{\{2\}}} - 1. \quad (31)$$

The difference of two solutions of the exponential Euler scheme (30) started with initial conditions u_0 and $u_0 + \Delta u_0$ is:

$$\begin{aligned}\Delta u_{n+1} &= \mathbf{M} \Delta u_n + h\varphi_1(h\mathbf{L}^{\{1\}}) \Delta \mathbf{g}^{\text{standard}\{1\}}(u_n) + h\varphi_1(h\mathbf{L}^{\{2\}}) \Delta \mathbf{g}^{\text{standard}\{2\}}(u_n) \\ &= \mathbf{M}^n \Delta u_0 + h \sum_{k=0}^n \mathbf{M}^k \varphi_1(h\mathbf{L}^{\{1\}}) \Delta \mathbf{g}^{\text{standard}\{1\}}(u_{n-k}) \\ &\quad + h \sum_{k=0}^n \mathbf{M}^k \varphi_1(h\mathbf{L}^{\{2\}}) \Delta \mathbf{g}^{\text{standard}\{2\}}(u_{n-k}) \\ \|\Delta u_{n+1}\| &\leq \|\mathbf{M}^n\| \|\Delta u_0\| + h \left(\|\varphi_1(h\mathbf{L}^{\{1\}})\| \ell^{\{1\}} + \|\varphi_1(h\mathbf{L}^{\{2\}})\| \ell^{\{1\}} \right) \sum_{k=0}^n \|\mathbf{M}^k\| \|\Delta u_{n-k}\|,\end{aligned}$$

where $\ell^{\{i\}}$ is the (small) Lipschitz constant of the smooth function $\mathbf{g}^{\text{standard}\{i\}}$ for $i = 1, 2$. The exponential Euler scheme (30) is stable if the matrix \mathbf{M} defined in (31) is power bounded.

6. Particular methods

We now turn our attention to building particular methods of partitioned exponential Runge–Kutta type. We start with known exponential Runge–Kutta methods (7) that satisfy the stiff order conditions given in Table 1 (for up to order four). Next we use Theorem 5.1 to obtain coefficients of the *transformed* exponential Runge–Kutta method (17). Finally, we apply the *transformed* method to construct PEXPRK schemes (18). Figure 1 illustrates this procedure.




Figure 1: Procedure to build a partitioned exponential Runge–Kutta method

We build partitioned exponential Runge–Kutta schemes of orders two, three and four. For the second order scheme, we go all the way and build a partitioned method. For the third and fourth order, we only provide the coefficients of the original and the *transformed* exponential Runge–Kutta method. It is then straightforward to build a partitioned method with the coefficients of the *transformed* method.

6.1. Second order partitioned exponential Runge–Kutta scheme

We start with the second order exponential Runge–Kutta scheme (7) given by the coefficients [18, Section 5.1]:

$$\mathbf{c} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \mathbf{a}(z) = \begin{bmatrix} 0 & 0 \\ \varphi_1(z) & 0 \end{bmatrix}, \quad \mathbf{b}(z) = \begin{bmatrix} \varphi_1(z) - \varphi_2(z) \\ \varphi_2(z) \end{bmatrix},$$

that satisfy order conditions 1., 2a., and 2b. in Table 1. Here, $s = 2$, $\mathbf{A}_{2:s,2:s} = 0$ and $\mathbf{E}_{2:s,2:s} = 1$. The coefficients of the transformed method are as follows:

$$\boldsymbol{\alpha}(z) = \begin{bmatrix} 0 & 0 \\ \varphi_1(z) & 0 \end{bmatrix}, \quad \boldsymbol{\beta}(z) = \begin{bmatrix} \varphi_1(z) - z\varphi_2(z)\varphi_1(z) \\ \varphi_2(z) \end{bmatrix}.$$

The partitioned implementation (18) of the *transformed* method reads:

$$\begin{aligned}
U_1 &= u_n, \\
U_2 &= u_n + h \sum_{p=1}^2 \varphi_1(hL^{\{p\}}) f^{\{p\}}(u_n), \\
u_{n+1} &= u_n + h \sum_{p=1}^2 \left(2\varphi_1(hL^{\{p\}}) - \varphi_1^2(hL^{\{p\}}) \right) f^{\{p\}}(u_n) \\
&\quad + h \sum_{p=1}^2 \varphi_2(hL^{\{p\}}) \left(f^{\{p\}}(U_2) - f^{\{p\}}(u_n) \right).
\end{aligned} \tag{32}$$

As in Theorem 5.2, the method (32) can be written in terms of residuals as follows:

$$\begin{aligned}
u_{n+1} &= u_n + h \sum_{p=1}^2 \left(\sum_{p' \neq p} \varphi_1(hL^{\{p'\}}) \right) \varphi_1(hL^{\{p\}}) f^{\{p\}}(u_n) \\
&\quad + h \sum_{p=1}^2 \varphi_2(hL^{\{p\}}) \left(g^{\{p\}}(U_2) - g^{\{p\}}(u_n) \right).
\end{aligned} \tag{33}$$

6.2. Third order partitioned exponential Runge–Kutta scheme

Consider the third order method by Hochbruck and Ostermann [18] given by the Butcher tableau:

$$\begin{array}{c|ccc}
0 & 0 & 0 & 0 \\
c_2 & c_2\varphi_{1,2} & 0 & 0 \\
\frac{2}{3} & \frac{2}{3}\varphi_{1,3} - \frac{4}{9c_2}\varphi_{2,3} & \frac{4}{9c_2}\varphi_{2,3} & 0 \\
\hline
& \varphi_1 - \frac{3}{2}\varphi_2 & 0 & \frac{3}{2}\varphi_2
\end{array} .$$

Here, $\varphi_j \equiv \varphi_j(z)$ and $\varphi_{j,k} \equiv \varphi_j(c_k z)$. For $c_2 = 2/3$, the *transformed* method coefficients, upon applying Theorem 5.1, are given shown in Method 1.

$$\begin{aligned}
\boldsymbol{\alpha} &= \begin{bmatrix} 0 & 0 & 0 \\ \frac{2\varphi_{1,2}}{3} & 0 & 0 \\ \frac{2}{9}(3\varphi_{1,3} - 2z\varphi_{1,2}\varphi_{2,3}) & \frac{2\varphi_{2,3}}{3} & 0 \end{bmatrix}, \\
\boldsymbol{\beta} &= \begin{bmatrix} \frac{1}{3}z\varphi_2(2z\varphi_{1,2}\varphi_{2,3} - 3\varphi_{1,3}) + \varphi_1(z) \\ -z\varphi_2\varphi_{2,3} \\ \frac{3\varphi_2}{2} \end{bmatrix}.
\end{aligned}$$

Method 1: Third order transformed, unpartitioned method.

6.3. Fourth order partitioned exponential Runge–Kutta scheme

We consider next the fourth order method by Hochbruck and Ostermann [18] given by the Butcher tableau:

0	0	0	0	0	0
$\frac{1}{2}$	$\frac{1}{2}\varphi_{1,2}$	0	0	0	0
$\frac{1}{2}$	$\frac{1}{2}\varphi_{1,3} - \varphi_{2,3}$	$\varphi_{2,3}$	0	0	0
1	$\varphi_{1,4} - 2\varphi_{2,4}$	$\varphi_{2,4}$	$\varphi_{2,4}$	0	0
$\frac{1}{2}$	$\frac{1}{2}\varphi_{1,5} - 2a_{5,2} - a_{5,4}$	$a_{5,2}$	$a_{5,2}$	$\frac{1}{4}\varphi_{2,5} - a_{5,2}$	0
	$\varphi_1 - 3\varphi_2 + 4\varphi_3$	0	0	$-\varphi_2 + 4\varphi_3$	$4\varphi_2 - 8\varphi_3$

Applying Theorem 5.1 the transformed method has the coefficients given in Method 2.

$$\alpha_{2:s,1} = \begin{bmatrix} \frac{\varphi_{1,2}}{2} \\ \frac{1}{2}(\varphi_{1,3} - z\varphi_{1,2}\varphi_{2,3}) \\ \frac{1}{2}z\varphi_{2,4}(\varphi_{1,2}(z\varphi_{2,3} - 1) - \varphi_{1,3}) + \varphi_{1,4} \\ \frac{1}{8}\left(2z\varphi_{1,4}(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5}))\right. \\ \left.+ z\varphi_{1,3}(-\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) - 2\varphi_{2,5} + 4\varphi_{3,4} + 2\varphi_{3,5})\right. \\ \left.+ z\varphi_{1,2}(z\varphi_{2,3} - 1)(\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) + 2(\varphi_{2,5} - 2\varphi_{3,4} - \varphi_{3,5})) + 4\varphi_{1,5}\right) \end{bmatrix},$$

$$\alpha_{2:s,2:s} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ \varphi_{2,3} & 0 & 0 & 0 \\ (1 - z\varphi_{2,3})\varphi_{2,4} & \varphi_{2,4} & 0 & 0 \\ -\frac{1}{4}(z\varphi_{2,3} - 1)\left(2(\varphi_{2,5} - 2\varphi_{3,4} - \varphi_{3,5})\right. & \frac{1}{4}\left(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5}))\right. & \frac{1}{4}\left(-\varphi_{2,4} - \varphi_{2,5}\right. & 0 \\ \left.+ \varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1)\right) & \left.\varphi_{2,4} + \varphi_{2,4} + 2\varphi_{2,5} - 4\varphi_{3,4} - 2\varphi_{3,5}\right) & \left.+ 4\varphi_{3,4} + 2\varphi_{3,5}\right) & \end{bmatrix},$$

$$\beta_1 = \varphi_1(z) - \frac{1}{2}z\left((\varphi_2 - 2\varphi_3)(2z\varphi_{1,4}(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5}))\right. \\ \left.+ z\varphi_{1,3}(-\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) - 2\varphi_{2,5} + 4\varphi_{3,4} + 2\varphi_{3,5})\right. \\ \left.+ z\varphi_{1,2}(z\varphi_{2,3} - 1)(\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) + 2(\varphi_{2,5} - 2\varphi_{3,4} - \varphi_{3,5})) + 4\varphi_{1,5})\right. \\ \left.- (\varphi_2 - 4\varphi_3)(z\varphi_{2,4}(\varphi_{1,2}(z\varphi_{2,3} - 1) - \varphi_{1,3}) + 2\varphi_{1,4})\right),$$

$$\beta_2 = \frac{1}{4}z(z\varphi_{2,3} - 1)\left(4(\varphi_2 - 2\varphi_3)(\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) + 2(\varphi_{2,5} - 2\varphi_{3,4} - \varphi_{3,5}))\right. \\ \left.- 4(\varphi_2 - 4\varphi_3)\varphi_{2,4}\right),$$

$$\beta_{3:s} = \begin{bmatrix} z(\varphi_2 - 4\varphi_3)\varphi_{2,4} + z(\varphi_2 - 2\varphi_3)(-\varphi_{2,4}(z(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) + 1) - 2\varphi_{2,5} + 4\varphi_{3,4} + 2\varphi_{3,5}) \\ z(\varphi_2 - 2\varphi_3)(\varphi_{2,4} + \varphi_{2,5} - 2(2\varphi_{3,4} + \varphi_{3,5})) - \varphi_2 + 4\varphi_3 \\ 4(\varphi_2 - 2\varphi_3) \end{bmatrix}.$$

Method 2: Fourth order transformed, unpartitioned method.

7. Computational considerations

The most expensive computation, outside function and Jacobian evaluation, in one step of an exponential integrator is the evaluation of the φ_k function (8) times vector products, which involves evaluating the matrix exponential operator. There is a significant body of work in computing matrix exponentials as evidenced in the review article by Moler et al. [28]. A number of authors have made additional contributions to the topic [1–3, 15, 16, 33, 37, 39].

In our implementation of exponential Runge–Kutta methods, we evaluate the $\varphi_k(h\gamma J)v$ products using Krylov-subspace based approximations. If N is the number of variables in the state-space, we use the modified Arnoldi algorithm [45] to build an M -dimensional Krylov-subspace ($M \ll N$):

$$\mathcal{K}_M = \{v, Jv, J^2v, J^3v, \dots, J^{M-1}v\},$$

returning an $N \times M$ orthonormal matrix V that spans \mathcal{K}_M and an $M \times M$ upper-Hessenberg matrix, H_M . Using these, the φ_k product is approximated according to [20, 30, 35, 38, 42]:

$$\varphi_k(h\gamma J)v \approx \|v\|V_M\varphi_k(h\gamma H_M)e_1.$$

We implement an adaptive Arnoldi algorithm [35] that evaluates the error at m -th iteration, $s_m = \varphi_1(h\gamma J)v - \|v\|V_m\varphi_k(h\gamma H_m)e_1$, and stopping when the error under a chosen tolerance. Even though we only evaluate the error for φ_1 , we argue that this should be sufficient for higher order φ_k functions as they must converge faster according to their series expansion. Furthermore, since the computation of error to limit the size of the subspace (and the number of iterations of Arnoldi) in itself relies on performing a matrix exponential on the upper-Hessenberg matrix H_m , we only compute the error at some pre-determined indices [17, Section 6.4], where the cost of computing the error roughly doubles the total cost of all preceding error computations.

In the remainder of this section, we attempt to establish guidelines for when partitioned exponential methods will be computationally more efficient per timestep than traditional exponential methods applied to the full system.

Cost of unpartitioned method. The cost of an Arnoldi process, excluding the cost of error estimation code, to construct V_M and H_M is $\mathcal{O}(MN^2 + M^2N + MN)$, assuming that the matrices are dense and a matrix vector product in the full space requires $\mathcal{O}(N^2)$ operations. Since we only compute the error at indices where the cost of computing the error roughly doubles the cost of all previous error computations, the total cost of computing the error after M iterations is $\mathcal{O}(M^3)$. This cost comes from the fact that the error estimation routine requires the exponentiation of the H_M matrix, with a constant number of rows and columns augmented around it which we ignore in its cost.

The total cost per timestep of an unpartitioned exponential method that evaluates u φ matrix function vector products is $\mathcal{O}(u \times (MN^2 + M^2N + MN + M^3))$ (ignoring a constant number of vector adds). Since $N \gg M$ the cost is approximately $\sim uMN^2$.

Cost of component partitioned method. Consider a P -way component partitioned system with components of size N_i , $i = 1, \dots, P$, respectively, such that $N = N_1 + \dots + N_P$. Consider a partitioned method that performs p matrix-vector operations on each partition. Assume next that the stiff components are divided proportionately amongst the partitions, and the Arnoldi processes need the same number of M iterations to converge. The cost per step of the partitioned method is $\sim pMN_1^2 + \dots + pMN_P^2$. The partitioned method is more effective than the full method when

$$p < u \frac{\left(\sum_{i=1}^P N_i\right)^2}{\sum_{i=1}^P N_i^2}.$$

When $N_i = N/P$ the partitioned method is more effective when $p < u/P^2$.

Cost of additively partitioned method. The analytical functions are applied to matrices $hL^{\{p\}}$ of the same size. Computational savings are realized due to the simpler structure of individual matrices. First, the number of Krylov iterations for computing each individual $\varphi_k(hL^{\{p\}})v$ can be much smaller than that of the

full, more complex Jacobian $\varphi_k(hL)v$. Next, the sparsity structure of individual matrices can be much more favorable than the sparsity structure of the sum. Individual component matrices can have special structure (such as block diagonal) that can further speed up the calculations.

Further computational optimizations specific to the application of partitioned exponential methods to reaction-diffusion systems are discussed in greater detail in our earlier work [31].

8. Results

We perform fixed timestep numerical experiments using each exponential Runge–Kutta method described in Section 6 in the original formulation, transformed formulation, and partitioned formulation (for a test-specific partitioning). In the results that follow, we use the following naming convention for the methods.

Labels starting with prefix “exprks” stand for unpartitioned, exponential Runge–Kutta scheme. Likewise, labels starting with “pexprks” stand for partitioned, exponential Runge–Kutta scheme. We also use the keywords “orig” and “tran” to indicate whether the unpartitioned methods are using the original coefficients or the transformed coefficients. Following this, we specify the design order of the method with the phrase “order = design order” directly in the label itself. For example, “order = 2” in the label corresponds to the second-order method from Section 6. Lastly, we specify the structure of the Jacobian used with methods implemented using the transformed coefficients, i.e., whether it’s a full or a block Jacobian. We use the full Jacobian with the original methods but do not specify it in the label. To summarize, the nomenclature for labeling each method in the plots is as follows:


$$[\text{exprks or pexprks}]_ [\text{orig or tran}]_ [\text{order} = 2 \text{ or } 3 \text{ or } 4]_ [(\text{Full or Block Jacobian})].$$

8.1. Gray-Scott equations

The Gray-Scott equations [13]:

$$\begin{aligned} \frac{\partial a}{\partial t} &= d_a \nabla^2 a - ab^2 + f(1 - a), \\ \frac{\partial b}{\partial t} &= d_b \nabla^2 b + ab^2 - (f + k)b, \end{aligned} \tag{34}$$

model the diffusion and reaction of two species a and b over a spatial domain where the chemical reactions are:



In equation (34) a, b are the concentrations of the two chemical species, d_a and d_b are the diffusion coefficients of the species, f is the feed rate of a , and $f + k$ is drain rate of b . Here we use the parameter values $f = 0.04$, $k = 0.06$, $d_a = 2$, $d_b = 1$.

The spatial domain is $[0, 1] \times [0, 1]$ and the timespan for experiments is $[0, 0.262144]$. Initial conditions for the model are given by the expressions $v(x, y, 0) = 0.4 + 0.1(x + y) + 0.1(\sin(10 * x) \sin(20 * y))$ and $w(x, y, 0) = 0.4 + 0.1(x + y) + 0.1(\cos(10 * x) \cos(20 * y))$. The boundary conditions are periodic.

For our experiments, we use a 2D finite difference discretized Gray-Scott model with a grid resolution of 300×300 . Each method was tested for the set of stepsizes given by the MATLAB expression `0.262144 .* 2.^(-1:-1:-6)`, which roughly corresponds to the decimal range 1×10^{-1} to 4×10^{-3} . The reference solution was computed using ODE15s with absolute and relative tolerance set to 1.0×10^{-14} . Four different partitions of the discretized model were tested. For all experiments we used the additive implementation of the partitioned methods.

8.2. Component partitioning by chemical species

We first consider a component partitioning of (34), where each component is associated with a different chemical species. In equation (4) v and w variables corresponding to this splitting are

$$v = \begin{bmatrix} \bar{a} \\ \mathbf{0} \end{bmatrix}, \quad w = \begin{bmatrix} \mathbf{0} \\ \bar{b} \end{bmatrix}, \quad (35)$$

where \bar{a} and \bar{b} are the state variables corresponding to the chemical species A and B after spatial discretization. With \mathbb{L} representing the discrete Laplacian operator, and $\mathbf{1}$, a column of ones of appropriate dimension, the right-hand side functions $f^{\{1\}}$ and $f^{\{2\}}$ can be written as

$$f^{\{1\}}(v+w) = \begin{bmatrix} d_a * \mathbb{L} * \bar{a} - \bar{a} \odot \bar{b} \odot \bar{b} + f * (\mathbf{1} - \bar{a}) \\ \mathbf{0} \end{bmatrix}, \quad f^{\{2\}}(v+w) = \begin{bmatrix} \mathbf{0} \\ d_b * \mathbb{L} * \bar{b} + \bar{a} \odot \bar{b} \odot \bar{b} - (f+k) * \bar{b} \end{bmatrix}.$$

The corresponding linear operators in (18) are

$$\mathbb{L}^{\{1\}} = \begin{bmatrix} d_a * \mathbb{L} - \text{diag}(\bar{b} \odot \bar{b}) - f * \mathbb{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}, \quad \mathbb{L}^{\{2\}} = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & d_b * \mathbb{L} + 2 * \text{diag}(\bar{a} \odot \bar{b}) - (f+k) * \mathbb{I} \end{bmatrix},$$

where \mathbb{I} is the identity matrix and diag is a function that returns a diagonal matrix with arguments placed along the main diagonal.

Figure 2 shows the results of the fixed timestep experiments. All methods show the correct order of convergence. As a validation of the *transformation* Theorem 5.1, we observe that the original and transformed methods converge at the same rate as expected.




Figure 2: Gray-Scott equations (34) partitioned by chemical species (35).

Remark. For component partitioned systems, an unpartitioned exponential Runge–Kutta scheme using a block-diagonal Jacobian, \mathbb{L} , implemented as a *transformed* method is equivalent to its partitioned counterpart. If we set

$$\begin{aligned} \bar{Y} &= \begin{bmatrix} \bar{a} \\ \bar{b} \end{bmatrix}, \quad f(\bar{Y}) = \begin{bmatrix} d_a * \mathbb{L} * \bar{a} - \bar{a} \odot \bar{b} \odot \bar{b} + f * (\mathbf{1} - \bar{a}) \\ d_b * \mathbb{L} * \bar{b} + \bar{a} \odot \bar{b} \odot \bar{b} - (f+k) * \bar{b} \end{bmatrix}, \\ \mathbb{L} &= \begin{bmatrix} d_a * \mathbb{L} - \text{diag}(\bar{b} \odot \bar{b}) - f * \mathbb{I} & \mathbf{0} \\ \mathbf{0} & d_b * \mathbb{L} + 2 * \text{diag}(\bar{a} \odot \bar{b}) - (f+k) * \mathbb{I} \end{bmatrix}, \end{aligned} \quad (36)$$

the results are identical to those obtained by the partitioned methods as shown in Figure 2.

8.3. Component partitioning by splitting the spatial domain

In this experiment, we partition the variables by their spatial location into two domains as shown in Figure 3. ODE functions and linear operators are updated accordingly. In order to achieve this, we make use of permutation matrices which reorder the state variables first by grid location and then by species, such that all variables in subdomain Ω_1 are the first $N/2$ variables in the permuted state vector, and the variables in subdomain Ω_2 are the last $N/2$ ones.




Figure 3: Spatial domains when variables are partitioned by location in the numerical grid.

If \mathbf{P} is the permutation matrix which performs the rearrangement of the state variables, the variables corresponding to this splitting scheme are $v = (\mathbf{P}^\top \bar{Y})_{1:N/2}$, $w = (\mathbf{P}^\top \bar{Y})_{N/2+1:N}$, and the corresponding partition is applied to the right hand side function (36). Lastly, if \mathbf{J} is the Jacobian of the right hand side, then the linear operators are defined as $\mathbf{L}^{\{1\}} = (\mathbf{P}^\top \mathbf{J} \mathbf{P})_{1:N/2,1:N/2}$ and $\mathbf{L}^{\{2\}} = (\mathbf{P}^\top \mathbf{J} \mathbf{P})_{N/2+1:N,N/2+1:N}$.

Figure 4 shows the results of fixed timestep experiment for the scenario where the discretized model is partitioned by spatial domain. We note that all the methods show the theoretical orders of convergence.




Figure 4: Gray-Scott equations (34) partitioned by spatial domains on a 300x300 grid, as in Figure 3.

8.4. Additive partitioning by physical process

Next, we consider an additive partitioning of the Gray-Scott system. The right-hand side of the PDE (34) models diffusion of two chemical species, A and B , in a medium and the interaction between species via reaction terms. For this experiment, we split the system along these two distinct physical processes:

$$\mathbf{f}^{\{1\}}(\bar{a}, \bar{b}) = \mathbf{f}^{\text{diffusion}} = \begin{bmatrix} d_a * \mathbb{L} * \bar{a} \\ d_b * \mathbb{L} * \bar{b} \end{bmatrix}, \quad (37a)$$

and,

$$\mathbf{f}^{\{2\}}(\bar{a}, \bar{b}) = \mathbf{f}^{\text{reaction}} = \begin{bmatrix} -\bar{a} \odot \bar{b} \odot \bar{b} + f * (\mathbf{1} - \bar{a}) \\ +\bar{a} \odot \bar{b} \odot \bar{b} - (f + k) * \bar{b} \end{bmatrix}. \quad (37b)$$

The linear operators corresponding to these right-hand sides are:

$$\mathbf{L}^{\{1\}} = \mathbf{L}^{\text{diffusion}} = \begin{bmatrix} d_a * \mathbb{L} & \mathbf{0} \\ \mathbf{0} & d_b * \mathbb{L} \end{bmatrix}, \quad (37c)$$

and

$$\mathbf{L}^{\{2\}} = \mathbf{L}^{\text{reaction}} = \begin{bmatrix} -\text{diag}(\bar{b} \odot \bar{b}) - f * \mathbb{I} & -2 * \text{diag}(\bar{a} \odot \bar{b}) \\ \text{diag}(\bar{b} \odot \bar{b}) & 2 * \text{diag}(\bar{a} \odot \bar{b}) - (f + k) * \mathbb{I} \end{bmatrix}. \quad (37d)$$

Linear algebra takes advantage of the highly regular block structure of $\mathbf{L}^{\{1\}}$ and $\mathbf{L}^{\{2\}}$.

Figure 5 shows the results of fixed timestep experiment for the above scenario where the discretized model is partitioned by physics, viz., reaction and diffusion. Again, we get full convergence in all the partitioned methods.




Figure 5: Gray-Scott equations (34) partitioned by physics (37).

8.5. Partitioning by physics with reaction treated explicitly, and diffusion treated exponentially

We note that, for our settings, the reaction terms in the Gray-Scott model (34) are less stiff compared to the diffusion terms. Consequently, we test our partitioned exponential methods with an “IMEX-like” configuration [43, 44]. Specifically, we treat the stiffer diffusion partition exponentially, and the less stiff reaction partition explicitly. We make one minor modification to the previous experiment to implement this. While we define $\mathbf{f}^{\{1\}}$, $\mathbf{f}^{\{2\}}$ and $\mathbf{L}^{\{1\}}$ similarly to the previous experiment, we set $\mathbf{L}^{\{2\}} = \mathbf{0}$.

When we set the linear operator to a zero matrix, an Exponential Runge–Kutta method degenerates to the underlying Runge–Kutta method [18]. Likewise, by setting the linear operator of any one partition of a PEXPRK to a zero matrix, we obtain an explicit method in that partition.

Figure 6 shows the results of fixed timestep experiment for the scenario where the discretized model is partitioned by physics, i.e., into reaction and diffusion, with the reaction partition treated explicitly and the diffusion partition exponentially. We observe the theoretical convergence behavior in all the partitioned methods.




Figure 6: Gray-Scott equations (34) partitioned by physics with the reaction term treated explicitly, and the diffusion term treated exponentially.

9. Conclusions

In our earlier work [31] we developed classical order conditions theory for partitioned methods that solve each component of a multiphysics problem with a different exponential method. We found that the partitioned methods were amenable to certain computational optimizations that unpartitioned methods could not take advantage of, and that the partitioned methods can be more stable than unpartitioned methods for some parameter settings. However, those methods are challenged by stiff systems with a tight coupling between components.

In this work we develop an implementation of exponential Runge–Kutta methods that is specific to partitioned systems. The resulting partitioned exponential methods inherit the W stiff order conditions from the base methods [21, 23–25]. A direct application of stiff exponential Runge–Kutta methods to partitioned systems is not immediate due to the properties of the matrix exponential operator and the non-linear terms. Our approach is to introduce a transformation for exponential Runge–Kutta methods that enables the derivation of partitioned methods of that type. We apply the modified methods to component and to additively partitioned systems, using a special approximation of the Jacobian as the linear operator. This approximation allows to evaluate matrix functions on Jacobians of individual component functions rather than on the combined Jacobian. These matrix functions are combined to obtain the numerical solution. The resulting partitioned exponential methods are of exponential-exponential and exponential-explicit type.

We derive three methods of second, third and fourth order of partitioned exponential type and numerically demonstrate that they converge as expected for different types of partitions of the test problem.

Acknowledgements

This work has been supported in part by NSF through awards NSF ACI–1709727 and NSF CCF–1613905, AFOSR through the award AFOSR DDDAS 15RT1037, and by the Computational Science Laboratory at Virginia Tech.

References

References

- [1] Awad H. Al-Mohy and Nicholas J. Higham. A new scaling and squaring algorithm for the matrix exponential. *SIAM Journal on Matrix Analysis and Applications*, 31(3):970–989, jan 2010.
- [2] Awad H. Al-Mohy and Nicholas J. Higham. Computing the action of the matrix exponential, with an application to exponential integrators. *SIAM Journal on Scientific Computing*, 1(2):1–23, 2011.
- [3] Philipp Bader, Sergio Blanes, and Muaz Seydaoğlu. The scaling, splitting, and squaring Method for the exponential of perturbed matrices. *SIAM Journal on Matrix Analysis and Applications*, 36(2):594–614, jan 2015.
- [4] T. Belytschko, H.-J. Yen, and R. Mullen. Mixed methods for time integration. *Computer Methods in Applied Mechanics and Engineering*, 17-18:259–275, feb 1979.
- [5] J.C. Butcher. *Numerical Methods for Ordinary Differential Equations*. Wiley, 3 edition, 2016.
- [6] A. Cardone, Z. Jackiewicz, A. Sandu, and H. Zhang. Extrapolated IMEX Runge-Kutta methods. *Mathematical Modelling and Analysis*, 19(2):18–43, 2014.
- [7] A. Cardone, Z. Jackiewicz, A. Sandu, and H. Zhang. Extrapolation-based implicit-explicit general linear methods. *Numerical Algorithms*, 65(3):377–399, 2014.
- [8] A. Cardone, Z. Jackiewicz, A. Sandu, and H. Zhang. Construction of highly-stable implicit-explicit general linear methods. In M. de Leon, W. Feng, Z. Feng, J.L. Gomez, X. Lu, J.M. Martell, J. Parcet, D. Peralta-Salas, and W. Ruan, editors, *AIMS proceedings*, volume 0133-0189_2015_special_185 of *Dynamical Systems, Differential Equations, and Applications*, pages 185–194, Madrid, Spain, 2015.
- [9] Ching-Shan Chou, Yong-Tao Zhang, Rui Zhao, and Qing Nie. Numerical methods for stiff reaction-diffusion systems. *Discrete and Continuous Dynamical Systems - Series B*, 7(3):515–525, feb 2007.
- [10] E.M. Constantinescu and A. Sandu. Extrapolated implicit-explicit time stepping. *SIAM Journal on Scientific Computing*, 31(6):4452–4477, 2010.
- [11] Wulf G. Dettmer and Djordje Perić. A new staggered scheme for fluid-structure interaction. *International Journal for Numerical Methods in Engineering*, 93(1):1–22, jul 2012.
- [12] Charbel Farhat, K.C. Park, and Yves Dubois-Pelerin. An unconditionally stable staggered algorithm for transient finite element analysis of coupled thermoelastic problems. *Computer Methods in Applied Mechanics and Engineering*, 85(3):349–365, feb 1991.
- [13] P. Gray and S. K. Scott. Autocatalytic reactions in continuous stirred tank reactor. Technical Report 6, 1984.
- [14] M. Günther and A. Sandu. Multirate generalized additive Runge-Kutta methods. *Numerische Mathematik*, 133(3):497–524, 2016.
- [15] Nicholas J. Higham. The scaling and squaring method for the matrix exponential revisited. *SIAM Journal on Matrix Analysis and Applications*, 26(4):1179–1193, jan 2005.
- [16] Nicholas J. Higham. The scaling and squaring method for the matrix exponential revisited. *SIAM Review*, 51(4):747–764, nov 2009.
- [17] M. Hochbruck, C. Lubich, and H. Selhofer. Exponential integrators for large systems of differential equations. *SIAM Journal on Scientific Computing*, 19(5):1552–1574, 1998.
- [18] M. Hochbruck and A. Ostermann. Explicit exponential Runge–Kutta methods for semilinear parabolic problems. *SIAM Journal on Numerical Analysis*, 43:1069–1090, 2005.
- [19] M. Hochbruck and A. Ostermann. Exponential integrators. *Acta Numerica*, 19:209–286, 2012.
- [20] Marlis Hochbruck and Christian Lubich. On Krylov subspace approximations to the matrix exponential operator. *SIAM Journal on Numerical Analysis*, 34(5):1911–1925, oct 1997.
- [21] Marlis Hochbruck, Alexander Ostermann, and Julia Schweitzer. Exponential Rosenbrock-type methods. *SIAM Journal on Numerical Analysis*, 47(1):786–803, jan 2009.
- [22] J. Loffeld and M. Tokman. Comparative performance of exponential, implicit, and explicit integrators for stiff systems of ODEs. *Journal of Computational and Applied Mathematics*, 241:45–67, mar 2013.
- [23] Vu Thai Luan and Alexander Ostermann. Explicit exponential Runge–Kutta methods of high order for parabolic problems. *Journal of Computational and Applied Mathematics*, 256:168–179, jan 2014.
- [24] Vu Thai Luan and Alexander Ostermann. Exponential Rosenbrock methods of order five - Construction, analysis and numerical comparisons. *Journal of Computational and Applied Mathematics*, 255:417–431, 2014.
- [25] Vu Thai Luan and Alexander Ostermann. Stiff order conditions for exponential Runge–Kutta methods of order five. In *Modeling, Simulation and Optimization of Complex Processes - HPSC 2012*, pages 133–143. Springer International Publishing, 2014.
- [26] Vu Thai Luan, Mayya Tokman, and Greg Rainwater. Preconditioned implicit-exponential (IMEXP) time integrators for stiff differential equations. Technical report, 2016.
- [27] Borislav V Minchev and Will M Wright. A review of exponential integrators for first order semi-linear problems. 2005.
- [28] Cleve Moler and Charles Van Loan. Nineteen dubious ways to compute the exponential of a matrix, twenty-five years later. *SIAM review*, 45(1):3–49, 2003.
- [29] M. Narayanamurthi, U. Romer, and A. Sandu. Solving parameter estimation problems with discrete adjoint exponential integrators. *Optimization Methods and Software*, 33(4–6):750–770, 2018.
- [30] M. Narayanamurthi, P. Tranquilli, A. Sandu, and M. Tokman. EPIRK-W and EPIRK-K time discretization methods. *Journal of Scientific Computing*, 78(1):167–201, 2019.

- [31] Mahesh Narayanamurthi and Adrian Sandu. Partitioned exponential methods for coupled multiphysics systems. arXiv e-prints, page arXiv:1908.09434, Aug 2019.
- [32] Qing Nie, Yong-Tao Zhang, and Rui Zhao. Efficient semi-implicit schemes for stiff systems. Journal of Computational Physics, 214(2):521–537, may 2006.
- [33] Jitse Niesen and Will M. Wright. Algorithm 919: A Krylov subspace algorithm for evaluating the φ -functions appearing in exponential integrators. ACM Trans. Math. Softw., 38(3):22:1–22:19, 2012.
- [34] G. Rainwater and M. Tokman. A new approach to constructing efficient stiffly accurate EPIRK methods. Journal of Computational Physics, 323:283–309, 2016.
- [35] Y. Saad. Analysis of some Krylov subspace approximations to the matrix exponential operator. SIAM Journal on Numerical Analysis, 29(1):209–228, feb 1992.
- [36] A. Sandu and M. Günther. A generalized-structure approach to additive Runge-Kutta methods. SIAM Journal on Numerical Analysis, 53(1):17–42, 2015.
- [37] Thomas Schmelzer and Lloyd N Trefethen. Evaluating matrix functions for exponential integrators via Carathéodory-Fejér approximation and contour integrals. Electronic Transactions on Numerical Analysis, 29:1–18, 2007.
- [38] Roger B. Sidje. Expokit: a software package for computing matrix exponentials. ACM Transactions on Mathematical Software, 24(1):130–156, mar 1998.
- [39] B. Skaflestad and W.M. Wright. The scaling and modified squaring method for matrix functions related to the exponential. Applied Numerical Mathematics, 59(3-4):783–799, mar 2009.
- [40] M Tokman. A new class of exponential propagation iterative methods of Runge-Kutta type (EPIRK). Journal of Computational Physics, 230(24):8762–8778, 2011.
- [41] Mayya Tokman. Efficient integration of large stiff systems of ODEs with exponential propagation iterative (EPI) methods. Journal of Computational Physics, 213(2):748–776, apr 2006.
- [42] P. Tranquilli and A. Sandu. Exponential-Krylov methods for ordinary differential equations. Journal of Computational Physics, 278:31–46, 2014.
- [43] U.M. Ascher and S.J. Ruuth and B.T.R. Wetton. Implicit-explicit methods for time-dependent partial differential equations. SIAM Journal on Numerical Analysis, 32(3):797–823, 1995.
- [44] U.M. Ascher and S.J. Ruuth and R.J. Spiteri. Implicit-explicit Runge-Kutta methods for time-dependent partial differential equations. Applied Numerical Mathematics, 25:151–167, 1997.
- [45] Henk A. van der Vorst. Iterative Krylov methods for large linear systems. Cambridge University Press, 2003.
- [46] J. G. Verwer and B. P. Sommeijer. An implicit-explicit Runge-Kutta-Chebyshev scheme for diffusion-reaction equations. SIAM Journal on Scientific Computing, 25(5):1824–1835, 2004.
- [47] H. Zhang and A. Sandu. A second-order diagonally-implicit-explicit multi-stage integration method. In Proceedings of the International Conference on Computational Science ICCS 2012, volume 9, pages 1039–1046, April 2012.
- [48] H. Zhang, A. Sandu, and S. Blaise. Partitioned and implicit-explicit general linear methods for ordinary differential equations. Journal of Scientific Computing, 61(1):119–144, 2014.
- [49] H. Zhang, A. Sandu, and S. Blaise. High order implicit-explicit general linear methods with optimized stability regions. SIAM Journal on Scientific Computing, 38(3):A1430–A1453, 2016.
- [50] Su Zhao, Jeremy Ovadia, Xinfeng Liu, Yong-Tao Zhang, and Qing Nie. Operator splitting implicit integration factor methods for stiff reaction-diffusion-advection systems. Journal of Computational Physics, 230(15):5996–6009, jul 2011.
- [51] E. Zharovsky, A. Sandu, and H. Zhang. A class of IMEX two-step Runge-Kutta methods. SIAM Journal on Numerical Analysis, 53(1):321–341, 2015.
- [52] O. C. Zienkiewicz, D. K. Paul, and A. H. C. Chan. Unconditionally stable staggered solution procedures for soil-pore fluid interaction problems. International Journal of Rock Mechanics and Mining Sciences & Geomechanics Abstracts, 25(5):233, oct 1988.