
Distributionally Robust Deep Learning using Hardness Weighted Sampling

Lucas Fidon
BMEIS
King's College London
lucas.fidon@kcl.ac.uk

Sébastien Ourselin
BMEIS
King's College London

Tom Vercauteren
BMEIS
King's College London
tom.vercauteren@kcl.ac.uk

Abstract

Limiting failures of machine learning systems is vital for safety-critical applications. In order to improve the robustness of machine learning systems, Distributionally Robust Optimization (DRO) has been proposed as a generalization of Empirical Risk Minimization (ERM). However, its use in deep learning has been severely restricted due to the relative inefficiency of the optimizers available for DRO in comparison to the wide-spread variants of Stochastic Gradient Descent (SGD) optimizers for ERM. We propose SGD with hardness weighted sampling, a principled and efficient optimization method for DRO in machine learning that is particularly suited in the context of deep learning. Similar to a hard example mining strategy in practice, the proposed algorithm is straightforward to implement and computationally as efficient as SGD-based optimizers used for deep learning, requiring minimal overhead computation. In contrast to typical ad hoc hard mining approaches, we prove the convergence of our DRO algorithm for over-parameterized deep learning networks with ReLU activation and finite number of layers and parameters. Our experiments on brain tumor segmentation in MRI demonstrate the feasibility and the usefulness of our approach. Using our hardness weighted sampling leads to a decrease of 2% of the interquartile range of the Dice scores for the enhanced tumor and the tumor core regions. The code for the proposed hard weighted sampler will be made publicly available.

1 Introduction

Modern datasets used to train deep learning models typically aggregate data from several sources, such as different clinical imaging centers or scanner manufacturers [37]. Each source provides examples that were acquired for a specific need and using a specific protocol. As a result, there is a high risk that some subdomains are underrepresented in the aggregated training dataset.

In standard deep learning pipelines, this distributional shift is ignored and the model is trained to minimize the mean per-example loss, which corresponds to the standard Empirical Risk Minimization (ERM) problem. As a result, models trained with ERM are more likely to underperform on those examples from the underrepresented subdomains, seen as *hard examples*. This may lead to *unfair* AI systems [21]. For example, state-of-the-art deep learning models for brain tumor segmentation (currently trained using ERM) underperform for cases with confounding effects, such as low grade gliomas, despite achieving good average and median performance [5]. For safety-critical systems, such as those used in healthcare, this greatly limits their usage [14].

Distributionally Robust Optimization (DRO) is a robust generalization of ERM that has been introduced in convex machine learning to model the uncertainty in the training data distribution [10, 13, 26, 28]. Instead of minimizing the mean per-example loss on the training dataset, DRO seeks the hardest *weighted* empirical training data distribution around the (uniform) empirical

training data distribution. This suggests a link between DRO and Hard Example Mining. However, DRO as a generalization of ERM for machine learning still lacks optimization methods that are principled and computationally as efficient as SGD in the non-convex setting of deep learning. Previously proposed principled optimization methods for DRO consist in alternating between approximate maximization and minimization steps [19, 23, 28]. However, they differ from SGD methods for ERM by the introduction of additional hyperparameters for the optimizer such as a second learning rate and a ratio between the number of minimization and maximization steps. This makes DRO difficult to use as a drop-in replacement for ERM in practice.

In contrast, efficient weighted sampling methods, including Hard Example Mining [9, 24, 32] and weighted sampling [6, 18], have been empirically shown to mitigate class imbalance issues and to improve deep embedding learning [16, 35, 38]. However, even though these works typically start from an ERM formulation, it is not clear how those heuristics actually relate to ERM in theory. This suggests that bridging the gap between DRO and weighted sampling methods could lead to a principled Hard Example Mining approach, or conversely to more efficient optimization methods for DRO in deep learning.

If one could efficiently solve the inner maximization problem in DRO, DRO could be addressed by maintaining a solution of the inner maximization problem and using a minimization scheme akin to the standard ERM but over an adaptively weighted empirical distribution. However, even in the case where a closed-form solution is available for the inner maximization problem, it would require performing a forward pass over the entire training dataset at each iteration. This cannot be done efficiently for large datasets. This suggests identifying an approximate, but practically usable, solution for the inner maximization problem based on a closed-form solution.

From a theoretical perspective, analysis of previous optimization methods for non-convex DRO [19, 23, 28] made the assumption that the model is either smooth or weakly-convex, but none of those properties are true for deep neural networks with ReLU activation functions that are typically used.

In this work, we propose SGD with hardness weighted sampling, a novel, principled optimization method for training deep neural networks with DRO and inspired by Hard Example Mining, that is computationally as efficient as SGD for ERM. Compared to SGD, our method only requires introducing an additional softmax layer and maintaining a stale per-example loss vector to compute sampling probabilities over the training data.

In practice, we show that our method outperforms plain SGD in the case of class imbalance, and improves the robustness of a state-of-the-art deep learning pipeline for brain tumor segmentation by reducing the interquartile range of the Dice scores of 2% for the segmentation of the enhancing tumor and the tumor core regions. We also formally link DRO in our method with Hard Example Mining. As a result, our method can be seen as a principled Hard Example Mining approach. In this context, the robustness parameter β of the DRO problem controls the trade-off between exploitation and exploration in the Hard Example Mining process. Last but not least, we generalize recent results in the convergence theory of SGD with ERM and over-parameterized deep learning networks with ReLU activation functions [2, 1, 8, 40] to our SGD with hardness weighted sampling for DRO. This is, to the best of our knowledge, the first convergence result for deep learning networks with ReLU trained with DRO.

2 Supervised Deep Learning with Distributionally Robust Optimization

Standard training procedures in machine learning are based on Empirical Risk Minimization (ERM) [7]. For a neural network h with parameters θ , a per-example loss \mathcal{L} , and a training dataset $\{(x_i, y_i)\}_{i=1}^n$, where x_i are the inputs and y_i are the labels, the ERM problem corresponds to

$$\min_{\theta} \left\{ \mathbb{E}_{(x,y) \sim p_{\text{train}}} [\mathcal{L}(h(x; \theta), y)] = \frac{1}{n} \sum_{i=1}^n \mathcal{L}(h(x_i; \theta), y_i) \right\} \quad (1)$$

where p_{train} is the empirical uniform distribution on the training dataset. Optionally, \mathcal{L} can contain a parameter regularization term that is only a function of θ .

The ERM training formulation assumes that p_{train} is an unbiased approximation of the true data distribution. However, this is generally impossible in domains such as medical image computing.

This makes models trained with ERM at risk of underperforming on images from parts of the data distribution that are underrepresented in the training dataset.

In contrast, Distributionally Robust Optimization (DRO) is a family of generalization of ERM in which the uncertainty in the training data distribution is modelled by minimizing the worst-case expected loss over an *uncertainty set* of training data distributions [29].

In this paper, we consider training deep neural networks with DRO based on a ϕ -divergence. We denote $\Delta_n := \{(p_i)_{i=1}^n \in [0, 1]^n \mid \sum_{i=1}^n p_i = 1\}$ the set of empirical training data probabilities vectors under consideration (i.e. the uncertainty set). We use the following definition of ϕ -divergence in the remainder of the paper.

Definition 2.1 (ϕ -divergence). *Let $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}$ be two times continuously differentiable on $[0, n]$, ρ -strongly convex on $[0, n]$, i.e. $\exists \rho > 0, \forall z, z' \in [0, n], \phi(z') \geq \phi(z) + \phi'(z)(z' - z) + \frac{\rho}{2}(z - z')^2$, and satisfying $\forall z \in \mathbb{R}, \phi(z) \geq \phi(1) = 0, \phi'(1) = 0$. The ϕ -divergence D_ϕ is defined as, for all $p = (p_i)_{i=1}^n, q = (q_i)_{i=1}^n \in \Delta_n$,*

$$D_\phi(q||p) = \sum_{i=1}^n p_i \phi\left(\frac{q_i}{p_i}\right) \quad (2)$$

We refer to our example 3.1 to highlight that the KL divergence is indeed a ϕ -divergence.

The DRO problem for which we propose an optimizer for training deep neural networks can be formally defined as

$$\arg \min_{\theta} \left\{ R(\mathbf{L}(h(\theta))) := \max_{q \in \Delta_n} \left(\mathbb{E}_{(x,y) \sim q} [\mathcal{L}(h(x; \theta), y)] - \frac{1}{\beta} D_\phi(q||p_{\text{train}}) \right) \right\} \quad (3)$$

where p_{train} is the uniform empirical distribution, and $\beta > 0$ an hyperparameter. The choice of β and ϕ controls how the unknown training data distribution q is allowed to differ from p_{train} . Here and thereafter, we use the notation $\mathbf{L}(h(\theta)) := (\mathcal{L}(h(x_i; \theta), y_i))_{i=1}^n$ to refer to the vector of loss values of the n training samples for the value θ of the parameters of the neural network h .

In the remainder of the paper, we will refer to R as the *distributionally robust loss*.

3 Hardness Weighted Sampling for Distributionally Robust Deep Learning

In the case where h is a non-convex predictor (such as a deep neural network), existing optimization methods for the DRO problem (3) alternate between approximate minimization and maximization steps [19, 23, 28], requiring the introduction of additional hyperparameters compared to SGD. However, these are difficult to tune in practice and convergence has not been proven for non-smooth deep neural networks such as those with ReLU activation functions.

In this section, we present an SGD-like optimization method for training a deep learning model h with the DRO problem (3). We first highlight, in Section 3.1, mathematical properties that allow us to link DRO with stochastic gradient descent (SGD) combined with an adaptive sampling that we refer to as *hardness weighted sampling*. In Section 3.2, we present our Algorithm 1 for distributionally robust deep learning. Then, in Section 3.3, we present theoretical convergence results for our hardness weighted sampling.

3.1 A sampling approach to Distributionally Robust Optimization

The goal of this subsection is to show that a stochastic approximation of the gradient of the *distributionally robust loss* can be obtained by using a weighted sampler. This result is a first step towards our Algorithm 1 for efficient training with the *distributionally robust loss* presented in the next subsection.

Our analysis of the properties of the *distributionally robust loss* R relies on the Fenchel duality [25] and the notion of Fenchel conjugate [15].

Definition 3.1 (Fenchel Conjugate Function). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper function. The Fenchel conjugate of f is defined as $\forall \mathbf{v} \in \mathbb{R}^m, f^*(\mathbf{v}) = \max_{\mathbf{x} \in \mathbb{R}^m} \langle \mathbf{v}, \mathbf{x} \rangle - f(\mathbf{x})$.*

To reformulate R as an unconstrained optimization problem over \mathbb{R}^n (rather than Δ_n), we define

$$\forall p \in \mathbb{R}^n, \quad G(p) = \frac{1}{\beta} D_\phi(p||p_{\text{train}}) + \delta_{\Delta_n}(p) \quad (4)$$

where δ_{Δ_n} is the characteristic function of the closed convex set Δ_n , i.e.

$$\forall p \in \mathbb{R}^n, \quad \delta_{\Delta_n}(p) = \begin{cases} 0 & \text{if } p \in \Delta_n \\ +\infty & \text{otherwise} \end{cases} \quad (5)$$

The distributionally robust loss R in (3) can now be rewritten using the Fenchel conjugate function G^* of G . This allows us to obtain regularity properties for R .

Lemma 3.1 (Regularity of R). *If ϕ satisfies Definition 2.1, then G and R satisfy the following:*

$$G \text{ is } \left(\frac{n\rho}{\beta}\right)\text{-strongly convex} \quad (6)$$

$$\forall \theta, \quad R(\mathbf{L}(h(\theta))) = \max_{q \in \mathbb{R}^n} (\langle \mathbf{L}(h(\theta)), q \rangle - G(q)) = G^*(\mathbf{L}(h(\theta))) \quad (7)$$

$$R \text{ is } \left(\frac{\beta}{n\rho}\right)\text{-gradient Lipschitz continuous.} \quad (8)$$

Equation (7) follows from Definition 3.1. Proofs of (6) and (8) can be found in Appendix 11.2. According to (6), the optimization problem (7) is strictly convex and admits a unique solution in Δ_n , which we denote as

$$\bar{p}(\mathbf{L}(h(\theta))) = \arg \max_{q \in \mathbb{R}^n} (\langle \mathbf{L}(h(\theta)), q \rangle - G(q)) \quad (9)$$

Thanks to those properties, we can now show the following lemma that states that the gradient, with respect to θ , of the distributionally robust loss R can be rewritten as the expectation, with respect to the weighted empirical distribution $\bar{p}(\mathbf{L}(h(\theta)))$, of the gradient of the per-example loss. We further show that straightforward analytical formulas exist for \bar{p} , and give an example of such probability distribution for the Kullback-Leibler divergence.

Lemma 3.2 (Stochastic Gradient of the Distributionally Robust Loss). *For all θ , we have*

$$\nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta) = \mathbb{E}_{(x,y) \sim \bar{p}(\mathbf{L}(h(\theta)))} [\nabla_{\theta} \mathcal{L}(h(x; \theta), y)] \quad (10)$$

The proof is found in Appendix 11.3.

We now provide a closed-form formula for \bar{p} given $\mathcal{L}(h(\theta))$ for the KL divergence. Closed-form formulas exist for other ϕ -divergences, e.g. the Pearson χ^2 divergence.

Example 3.1. *For $\phi : z \mapsto z \log(z) - z + 1$, D_{ϕ} is the Kullback-Leibler (KL) divergence:*

$$D_{\phi}(q||p) = D_{\text{KL}}(q||p) = \sum_{i=1}^n q_i \log \left(\frac{q_i}{p_i} \right) \quad (11)$$

In this case, we have (see Appendix 11.1 for a proof)

$$\bar{p}(\mathbf{L}(h(\theta))) = \text{softmax}(\beta \mathbf{L}(h(\theta))) \quad (12)$$

3.2 Proposed Efficient Algorithm for Distributionally Robust Deep Learning

We now describe our algorithm for training deep neural networks with DRO using our hardness weighted sampling.

The equation (10) implies that $\nabla_{\theta} \mathcal{L}(h(x_i; \theta), y_i)$ is an unbiased estimator of the distributionally robust loss gradient when i is sampled with respect to $\bar{p}(\mathbf{L}(h(\theta)))$. This suggests that the distributionally robust loss can be minimized efficiently by SGD by sampling mini-batches with respect to $\bar{p}(\mathbf{L}(h(\theta)))$ at each iteration. However, even though closed-form formulas were provided in Example 3.1 for \bar{p} , evaluating exactly $\mathbf{L}(h(\theta))$, i.e. doing one forward pass on the whole training dataset at each iteration, is computationally prohibitive for large training datasets.

In practice, we propose to use a stale version of the vector of per-example loss values by maintaining an online history of the loss values of the examples seen during training $(\mathcal{L}(h(x_i; \theta^{(t_i)}), y_i))_{i=1}^n$, where for all i , t_i is the last iteration at which the per-example loss of example i has been computed. Using the Kullback-Leibler divergence as ϕ -divergence, this leads to the SGD with hardness weighted sampling algorithm proposed in Algorithm 1.

Algorithm 1 Training procedure for DRO with Hardness Weighted Sampling.

Require: $\{(x_i, y_i)\}_{i=1}^n$: training dataset with $n > 0$ the number of training samples.
Require: $b \in \{1, \dots, n\}$: batch size.
Require: \mathcal{L} : (any) smooth per-example loss function.
Require: $\beta > 0$: robustness parameter defining the distributionally robust optimization problem.
Require: θ_0 : initial parameter vector for the model h to train.
Require: L_{init} : initial stale per-example loss values vector.

- 1: $t \leftarrow 0$ (initialize the time step)
- 2: $L \leftarrow L_{init}$ (initialize the vector stale loss values)
- 3: **while** θ_t has not converged **do**
- 4: $p_t \leftarrow \text{softmax}(\beta L)$ ▷ online estimation of the *hardness weights*
- 5: $I \sim p_t$ ▷ (approximate) hardness weighted sampling
- 6: **if** not use importance sampling **then**
- 7: $\forall i \in I, w_i = 1$
- 8: **else**
- 9: $\forall i \in I, w_i \leftarrow \exp(\beta(\mathcal{L}(h(x_i; \theta), y_i) - L_i))$ ▷ importance sampling weights
- 10: $\forall i \in I, w_i \leftarrow \text{clip}(w_i, [w_{min}, w_{max}])$ ▷ clip the weights for stability
- 11: $\forall i \in I, L_i \leftarrow \mathcal{L}(h(x_i; \theta), y_i)$ ▷ update the vector of stale loss values
- 12: $g_t \leftarrow \frac{1}{b} \sum_{i \in I} w_i \nabla_{\theta} \mathcal{L}(h(x_i; \theta), y_i)$
- 13: $\theta_{t+1} \leftarrow \theta_t - \eta g_t$
- 14: **Output:** θ_t

When data augmentation is used an infinite number of training examples is virtually available. In this case, we keep one stale loss value per example irrespective of any augmentation as an approximation of the loss for this example under any augmentation.

Importance sampling is often used when sampling with respect to a desired distribution cannot be done exactly [20]. In Algorithm 1, an up-to-date estimation of the per-example losses (or equivalently the hardness weights) in a batch is only available *after* sampling and evaluation through the network. Importance sampling can be used to compensate for the difference between the initial and the updated stale losses within this batch. We propose to use importance sampling in steps 9-10 of Algorithm 1 and highlight that this is especially useful to deal with data augmentation. Indeed, in this case, the stale losses for the examples in the batch are expected to be less accurate as they were estimated under a different augmentation. For efficiency, we use the following approximation $w_i = \frac{p_i^{new}}{p_i^{old}} \approx \exp(\beta(\mathcal{L}(h(x_i; \theta), y_i) - L_i))$ where we have neglected the change in the denominator of the softmax. More details are given in Appendix 9. To tackle the typical instabilities that can arise when using importance sampling [27], the importance weights are clipped.

Compared to standard SGD-based training optimizers for the mean loss, our algorithm requires only an additional softmax operation per iteration and to store an additional vector of scalars of size n (number of training examples), thereby making it well suited for deep learning applications.

3.3 Theoretical results

In this section, we present convergence guarantees for Algorithm 1 in the framework of over-parameterized deep learning. We further demonstrate properties of our hardness weighted sampling that allow to clarify its link with Hard Example Mining.

Convergence of SGD with Hardness Weighted Sampling for Over-parameterized DNN Convergence results for over-parameterized deep learning have recently been proposed in [1]. Their work gives convergence guarantees for deep neural networks h with any activation functions (including ReLU), and with any (finite) number of layers L and parameters m , under the assumption that m is large enough. In our work, we extend the convergence theory developed by [1] for ERM and SGD to DRO using the proposed SGD with hardness weighted sampling and stale per-example loss vector (as stated in Algorithm 1). The proof in Appendix 12.4 deals with the challenges raised by the non-linearity of R with respect to the per-sample stale loss and the non-uniform dynamic sampling used in Algorithm 1.

Theorem 3.1 (Convergence of Algorithm 1 for over-parameterized neural networks with ReLU). *Let \mathcal{L} be a smooth per-example loss function, $b \in \{1, \dots, n\}$ be the batch size, and $\epsilon > 0$. If m is large enough, and the learning rate is small enough, then, with high probability over the randomness of the initialization and the mini-batches, Algorithm 1 (without importance sampling) guarantees $\|\nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta)\| \leq \epsilon$ after a finite number of iterations.*

Link between Hardness Weighted Sampling and Hard Example Mining The following result shows that using the proposed *hardness weighted sampler* those training examples with relatively high values of the loss, are sampled with higher probability.

Theorem 3.2. *Let a ϕ -divergence that satisfies Definition 2.1, and $\mathbf{L} = (L_i)_{i=1}^n \in \mathbb{R}^n$ a vector of loss values for the examples $\{x_1, \dots, x_n\}$. The proposed hardness weighted sampling probabilities vector $\bar{p}(\mathbf{L}) = (\bar{p}_i(\mathbf{L}))_{i=1}^n$ defined as in (9) verifies:*

1. For all $i \in \{1, \dots, n\}$, \bar{p}_i is an increasing function of L_i .
2. For all $i \in \{1, \dots, n\}$, \bar{p}_i is a non-increasing function of any L_j for $j \neq i$.

See Appendix 11.4 for the proof. The second part of Theorem 3.2 implies that as the loss of an example diminishes, the sampling probabilities of all the other examples increase. As a result, the proposed SGD with hardness weighted sampling balances exploitation (i.e. sampling the identified *hard examples*) and exploration (i.e. sampling any example to keep the record of *hard examples* up to date). Heuristics to enforce this trade-off are often used in Hard Example Mining methods [6, 16, 38].

4 Experiments

4.1 Toy example: MNIST with a class imbalance

The goal of this subsection is to illustrate key benefits of training a deep neural network using DRO in comparison to ERM when a part of the sample distribution is underrepresented in the training dataset. In addition, we verify the ability of our Algorithm 1 to train a neural network for DRO and illustrates the behaviour of SGD with hardness weighted sampling for different values of β .

Material: We create a bias between training and testing data distribution of MNIST [22] by keeping only 1% of the digits 3 in the training dataset, while the testing dataset remains unchanged. A Wide Residual Network [39] was used and we refer to Appendix 8.1 for the implementation details

Results: Our experiment suggests that DRO and ERM lead to different optima. Indeed, DRO for $\beta = 10$ outperforms ERM by more than 15% of accuracy on the underrepresented class, as illustrated in Appendix 8.1, Figure 1. This suggests that DRO is more robust than ERM to domain gaps between the training and the testing dataset. In addition, Appendix Figure 1 suggests that DRO with our SGD with hardness weighted sampling can converge faster than ERM with SGD.

Furthermore, the variations of learning curves with β shown in Appendix Figure 1 are consistent with our theoretical insight. As β decreases to 0, the learning curve of DRO with our Algorithm 1 converges to the learning curve of ERM with SGD.

4.2 Toward more robust deep learning methods for brain tumor segmentation

In this section, we illustrate the application of Algorithm 1 to improve the robustness of deep learning methods for brain tumor segmentation using the BraTS 2019 dataset [3, 4].

Motivation: Due to the availability of the large (for medical imaging standards) public dataset BraTS, with data coming from 19 institutions, brain tumor segmentation is often used as a benchmark for deep learning methods for medical image computing. The task consists in automatically segmenting three types of tumor components in patients with brain gliomas using four channels of Magnetic Resonance Imaging (MRI) sequences commonly used in clinical practice. An illustration of a typical case is given in Appendix 8.2, Figure 2.

Gliomas are the most common brain (primary) tumor and are associated with a poor survival rate [36]. Automatic segmentation methods can help support the training of neuroradiologists, and aid diagnosis,

Table 1: **Dice score evaluation on the BraTS 2019 online validation set (125 cases).** Metrics were computed using the BraTS online evaluation platform (<https://ipp.cbica.upenn.edu/>). ERM: Empirical Risk Minimization, DRO: Distributionally Robust Optimization, SGD: plain SGD, Nesterov: SGD with Nesterov momentum, IQR: Interquartile range. The best values are in bold.

Optim. problem	Optim. update	Enhancing Tumor			Whole Tumor			Tumor Core		
		Mean	Median	IQR	Mean	Median	IQR	Mean	Median	IQR
ERM	SGD	71.3	86.0	20.9	90.4	92.3	6.1	80.5	88.8	17.5
DRO	SGD	72.3	87.2	19.1	90.5	92.6	6.0	82.1	89.7	15.2
ERM	Nesterov	73.0	87.1	15.6	90.7	92.6	5.4	83.9	90.5	14.3
DRO	Nesterov	74.5	87.3	13.8	90.6	92.6	5.9	84.1	90.0	12.5

and disease progression assessment, in addition to assisting prognosis and radiation therapy planning. In the latest comparative study published by the organizers of the BraTS challenge [5], they remark that most deep learning methods (54/63 teams) achieve satisfactory mean and median Dice scores but that lower interquartile range (IQR) is still needed for all of them so that their robustness reach human level performance, especially for the enhanced tumor and the tumor core [5, paragraph 3.3].

Material: We split the 323 cases of the BraTS 2019 *training* dataset into 268 for training and 67 for validation. As the official testing data are not publicly available, the BraTS 2019 *validation* dataset that contains 125 cases was used for testing.

The deep learning pipeline used was based on nnUNet [18], which is a standard baseline for deep learning methods for medical image segmentation that achieve state-of-the-art performance in 49 medical imaging segmentation tasks. nnUNet is based on an intensive use of data augmentation and a set of heuristics that automatically adjust the U-net neural network architecture [30], the pre-processing and the post-processing based on the characteristics of the dataset. More details about the selected hyperparameters are given in Appendix.

Our only modification of the nnUNet pipeline is the addition of our hardness weighted sampling and the possible modification of the optimization update rule as indicated in Table 1.

Results: Table 1 summarizes the performance of training nnUNet using classical Empirical Risk Minimization (ERM) or using the proposed hardness weighted sampler for Distributionally Robust Optimization (DRO). For both ERM and DRO, the optimization update rule used was either plain SGD without momentum, or SGD with a Nesterov momentum equal to 0.99. Especially, for the latter, this implies that step 12 of Algorithm 1 is modified to use SGD with Nesterov momentum. For DRO, the results presented here are for $\beta = 100$ and using importance sampling (step 6 of Algorithm 1).

As illustrated in Table 1, DRO outperforms ERM in terms of IQR for enhanced tumor and tumor core by approximately 2% of Dice score, and in terms of mean Dice score for the enhanced tumor. The IQR is the global statistic used in [5] to measure the level of robustness of a method.

In addition, Appendix 8.2, Figure 3 shows that the lower IQR of DRO for the enhancing tumor comes specifically from a lower number of poor segmentations on cases coming from The Cancer Imaging Archive (TCIA). This suggests that DRO can deal with some of the confounding biases present in the training dataset, and lead to a model that is more fair.

The same improvements are observed independently of the optimization update rule used. This suggests that in practice Algorithm 1 still converges when a momentum is used, even if Theorem 3.1 was only demonstrated to hold for plain SGD.

Results for $\beta \in \{10, 100, 1000\}$ with and without importance sampling can be found in Appendix Table 2. The tendency described previously still holds true for β equal to 10 or 100 with or without importance sampling. However, when β becomes too large ($\beta = 1000$) a decrease of the mean and median Dice score for all regions is observed compared to ERM. For all values of β the use of importance sampling, as described in steps 6-8 of Algorithm 1, improves the interquartile range of the Dice score.

Efficiency of Algorithm 1: For our training set of $n=268$ volumes and a batch size of 2, the additional memory usage of Algorithm 1 is only 2144 bytes of memory (one float array of size n) and the additional computational time is approximately 10^{-4} sec per iteration using numpy, i.e. approximately 0.006 % of the total duration of an iteration.

5 Discussion

Group-distributionally robust deep learning: Concurrently to our work, an optimization method for group-DRO was proposed in [31]. In contrast to the formulation of DRO that we study in this paper, their method requires additional labels allowing to identify the underrepresented group in the training dataset. However, those labels may not be available or even impossible to obtain in most applications.

They show that, when associated with strong regularization of the weights of the network, their group DRO method can tackle spurious correlations that are known a priori in some classification problems. It is worth noting that, in contrast, no regularization was necessary in our experiments with MNIST.

Distributionally robust optimization in machine learning with other uncertainty set: In this work, we focus on DRO with a ϕ -divergence [12]. In this case, the data distributions that are considered in the DRO problem (3) are restricted to sharing the support of the empirical training distribution. In other words, the weights assigned to the training data can change, but the training data itself remains unchanged.

Another popular formulation is DRO with a Wasserstein distance [10, 13, 33, 34]. In contrast to ϕ -divergences, using a Wasserstein distance in DRO seeks to apply small data augmentation to the training data to make the deep learning model robust to small deformation of the data, but the sampling weights of the training data distribution typically remains unchanged. In this sense, DRO with a ϕ -divergence and DRO with a Wasserstein distance can be considered as orthogonal endeavours. While we show that DRO with ϕ -divergence can be seen as a principled Hard Example Mining method, it has been shown that DRO with a Wasserstein distance can be seen as a principled adversarial training method [33, 34].

6 Conclusion

We have shown that efficient training of deep neural networks with Distributionally Robust Optimization (DRO) with a ϕ -divergence is possible.

Our Stochastic Gradient Descent (SGD) with hardness weighted sampling is as straightforward to implement, and as computationally efficient as SGD for Empirical Risk Minimization (ERM). It can be used for deep neural networks with any activation function (including ReLU), and with any per-example loss function. We have shown that the proposed approach can formally be described as a principled Hard Example Mining strategy (Theorem 3.2). In addition, we prove the convergence of our method for over-parameterized deep neural networks (Theorem 3.1). Thereby, extending the convergence theory of deep learning of [1]. This is, to the best of our knowledge, the first convergence result for training a deep neural network based on DRO.

In practice, we have shown that our hardness weighted sampling method can be easily integrated in a state-of-the-art deep learning framework for medical image segmentation. Interestingly, the proposed algorithm remains stable when SGD with momentum is used. Our experiments on multiclass brain tumor segmentation illustrates that training deep neural networks with DRO can improve their robustness, which is one of the main limitations of current state-of-the-art deep learning methods for this application [5].

7 Statement of broader impact

Modern datasets used for training machine learning models are always subject to a selection bias, with subgroups of subjects or conditions that are underrepresented. This is in particular the case for the datasets used in medical imaging where data typically come from a limited number of clinical

imaging centers and scanner manufacturers, and do not fairly cover all possible anatomical variations associated with a disease.

Recent studies have shown limitations of deep neural networks trained on medical imaging datasets without taking into account this selection bias. State-of-the-art deep neural networks for brain tumor segmentation underperform for cases with confounding effects, such as low grade gliomas [5]. It has been shown that scans coming from 15 different studies can be correctly re-assigned to their source using a random forest classifier with 73.3% accuracy [37]. And state-of-the-art deep neural networks for the diagnosis of 14 thoracic diseases using X-ray trained on a dataset with a gender bias underperform on X-ray of female patients.

The training method proposed in this paper automatically reweights the cases with lower performance, encouraging the model to perform more consistently on all cases. In our experiments on brain tumor segmentation using MRI, we have shown that training a state-of-the-art deep neural network using our algorithm leads to a model with a higher robustness to the clinical imaging center the data is coming from. More work is needed to evaluate the impact of our method on other applications in medical imaging.

We further emphasize that the proposed method can be easily incorporated into existing machine learning methods while requiring no additional label and minimal overhead computation. This is especially important in medical imaging where the data are generally anonymized and information related to possible confounding factors, such as age and gender, are typically not available.

References

- [1] Allen-Zhu, Z., Li, Y., Song, Z.: A convergence theory for deep learning via over-parameterization. In: ICML. pp. 242–252 (2019)
- [2] Allen-Zhu, Z., Li, Y., Song, Z.: On the convergence rate of training recurrent neural networks. In: Advances in Neural Information Processing Systems 32, pp. 6676–6688. Curran Associates, Inc. (2019)
- [3] Bakas, S., Akbari, H., Sotiras, A., Bilello, M., Rozycki, M., Kirby, J.S., Freymann, J.B., Farahani, K., Davatzikos, C.: Segmentation labels and radiomic features for the pre-operative scans of the TCGA-GBM collection. The Cancer Imaging Archive (2017). <https://doi.org/10.7937/K9/TCIA.2017.KLXWJJ1Q>
- [4] Bakas, S., Akbari, H., Sotiras, A., Bilello, M., Rozycki, M., Kirby, J.S., Freymann, J.B., Farahani, K., Davatzikos, C.: Segmentation labels and radiomic features for the pre-operative scans of the TCGA-LGG collection. The Cancer Imaging Archive (2017). <https://doi.org/10.7937/K9/TCIA.2017.GJQ7R0EF>
- [5] Bakas, S., Reyes, M., Jakab, A., Bauer, S., Rempfler, M., Crimi, A., Shinohara, R.T., Berger, C., Ha, S.M., Rozycki, M., et al.: Identifying the best machine learning algorithms for brain tumor segmentation, progression assessment, and overall survival prediction in the BRATS challenge. arXiv preprint arXiv:1811.02629 (2018)
- [6] Berger, L., Eoin, H., Cardoso, M.J., Ourselin, S.: An adaptive sampling scheme to efficiently train fully convolutional networks for semantic segmentation. In: Annual Conference on Medical Image Understanding and Analysis. pp. 277–286. Springer (2018)
- [7] Bottou, L., Curtis, F.E., Nocedal, J.: Optimization methods for large-scale machine learning. *Siam Review* **60**(2), 223–311 (2018)
- [8] Cao, Y., Gu, Q.: Generalization error bounds of gradient descent for learning overparameterized deep relu networks. In: AAAI (2020)
- [9] Chang, H.S., Learned-Miller, E., McCallum, A.: Active bias: Training more accurate neural networks by emphasizing high variance samples. In: Advances in Neural Information Processing Systems. pp. 1002–1012 (2017)
- [10] Chouzenoux, E., Gérard, H., Pesquet, J.C.: General risk measures for robust machine learning. *Foundations of Data Science* **1**, 249 (2019)

- [11] Çiçek, Ö., Abdulkadir, A., Lienkamp, S.S., Brox, T., Ronneberger, O.: 3D U-Net: learning dense volumetric segmentation from sparse annotation. In: International conference on medical image computing and computer-assisted intervention. pp. 424–432. Springer (2016)
- [12] Csiszár, I., Shields, P.C., et al.: Information theory and statistics: A tutorial. *Foundations and Trends® in Communications and Information Theory* **1**(4), 417–528 (2004)
- [13] Duchi, J., Glynn, P., Namkoong, H.: Statistics of robust optimization: A generalized empirical likelihood approach. arXiv preprint arXiv:1610.03425 (2016)
- [14] European Commission: Ethics guidelines for trustworthy AI. Report, European Commission (2019)
- [15] Fenchel, W.: On conjugate convex functions. *Canadian Journal of Mathematics* **1**(1), 73–77 (1949)
- [16] Harwood, B., Kumar, B., Carneiro, G., Reid, I., Drummond, T., et al.: Smart mining for deep metric learning. In: Proceedings of the IEEE International Conference on Computer Vision. pp. 2821–2829 (2017)
- [17] Hiriart-Urruty, J.B., Lemaréchal, C.: Convex analysis and minimization algorithms I: Fundamentals, vol. 305. Springer science & business media (2013)
- [18] Isensee, F., Jäger, P.F., Kohl, S.A., Petersen, J., Maier-Hein, K.H.: Automated design of deep learning methods for biomedical image segmentation. arXiv preprint arXiv:1904.08128 (2020)
- [19] Jin, C., Netrapalli, P., Jordan, M.I.: Minmax optimization: Stable limit points of gradient descent ascent are locally optimal. arXiv preprint arXiv:1902.00618 (2019)
- [20] Kahn, H., Marshall, A.W.: Methods of reducing sample size in Monte Carlo computations. *Journal of the Operations Research Society of America* **1**(5), 263–278 (1953)
- [21] Larrazabal, A.J., Nieto, N., Peterson, V., Milone, D.H., Ferrante, E.: Gender imbalance in medical imaging datasets produces biased classifiers for computer-aided diagnosis. *Proceedings of the National Academy of Sciences* (2020)
- [22] LeCun, Y.: The MNIST database of handwritten digits. <http://yann.lecun.com/exdb/mnist/> (1998)
- [23] Lin, T., Jin, C., Jordan, M.I.: On gradient descent ascent for nonconvex-concave minimax problems. arXiv preprint arXiv:1906.00331 (2019)
- [24] Loshchilov, I., Hutter, F.: Online batch selection for faster training of neural networks. ICLR Workshop (2016)
- [25] Moreau, J.J.: Proximité et dualité dans un espace hilbertien. *Bulletin de la Société mathématique de France* **93**, 273–299 (1965)
- [26] Namkoong, H., Duchi, J.C.: Stochastic gradient methods for distributionally robust optimization with f-divergences. In: *Advances in Neural Information Processing Systems*. pp. 2208–2216 (2016)
- [27] Owen, A., Zhou, Y.: Safe and effective importance sampling. *Journal of the American Statistical Association* **95**(449), 135–143 (2000)
- [28] Rafique, H., Liu, M., Lin, Q., Yang, T.: Non-convex min-max optimization: Provable algorithms and applications in machine learning. arXiv preprint arXiv:1810.02060 (2018)
- [29] Rahimian, H., Mehrotra, S.: Distributionally robust optimization: A review. arXiv preprint arXiv:1908.05659 (2019)
- [30] Ronneberger, O., Fischer, P., Brox, T.: U-net: Convolutional networks for biomedical image segmentation. In: International Conference on Medical image computing and computer-assisted intervention. pp. 234–241. Springer (2015)

- [31] Sagawa, S., Koh, P.W., Hashimoto, T.B., Liang, P.: Distributionally robust neural networks for group shifts: On the importance of regularization for worst-case generalization. ICLR (2020)
- [32] Shrivastava, A., Gupta, A., Girshick, R.: Training region-based object detectors with online hard example mining. In: Proceedings of the IEEE conference on computer vision and pattern recognition. pp. 761–769 (2016)
- [33] Sinha, A., Namkoong, H., Duchi, J.: Certifying some distributional robustness with principled adversarial training. ICLR (2018)
- [34] Staib, M., Jegelka, S.: Distributionally robust deep learning as a generalization of adversarial training. In: NIPS workshop on Machine Learning and Computer Security (2017)
- [35] Suh, Y., Han, B., Kim, W., Lee, K.M.: Stochastic class-based hard example mining for deep metric learning. In: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition. pp. 7251–7259 (2019)
- [36] Tamimi, A.F., Juweid, M.: Epidemiology and outcome of glioblastoma. In: Glioblastoma. Codon Publications (2017)
- [37] Wachinger, C., Becker, B.G., Rieckmann, A., Pölsterl, S.: Quantifying confounding bias in neuroimaging datasets with causal inference. In: International Conference on Medical Image Computing and Computer-Assisted Intervention. pp. 484–492. Springer (2019)
- [38] Wu, C.Y., Manmatha, R., Smola, A.J., Krahenbuhl, P.: Sampling matters in deep embedding learning. In: Proceedings of the IEEE International Conference on Computer Vision. pp. 2840–2848 (2017)
- [39] Zagoruyko, S., Komodakis, N.: Wide residual networks. In: Proceedings of the British Machine Vision Conference (BMVC). pp. 87.1–87.12. BMVA Press (2016)
- [40] Zou, D., Gu, Q.: An improved analysis of training over-parameterized deep neural networks. In: Advances in Neural Information Processing Systems 32, pp. 2055–2064. Curran Associates, Inc. (2019)

— Supplementary Material —
Distributionally Robust Deep Learning
using Hardness Weighted Sampling

Contents

8 More on our experiments	13
8.1 Experiments on MNIST	13
8.1.1 Implementation details	13
8.1.2 Comment on the stability of the learning curves	14
8.2 Experiments on BraTS 2019	14
8.2.1 Implementation details	14
9 Importance sampling in Algorithm 1	16
10 Summary of the notations used in the proofs	16
10.1 Probability Theory notations	16
10.2 Machine Learning notations	16
10.3 Distributionally Robust Optimisation notations	16
10.4 Miscellaneous	17
11 Proofs	17
11.1 Proof of Example 3.1: formula of the sampling probabilities for the KL divergence	17
11.2 Proof of lemma 3.1: regularity properties of R	18
11.3 Proof of Lemma 3.2: formula of the distributionally robust loss gradient	19
11.4 Proof of Theorem 3.2: Distributionally Robust Optimization as Principled Hard Example Mining	20
12 Proof of Theorem 3.1: convergence of SGD with Hardness Weighted Sampling for Over-parameterized Deep Neural Networks with ReLU	22
12.1 Assumptions	22
12.2 Convergence theorem (restated)	23
12.3 Proofs of convergence	24
12.3.1 Proof that $R \circ L$ is one-sided gradient Lipchitz	24
12.3.2 Semi-smoothness property of the distributionally robust loss	26
12.3.3 Gradient bounds for the distributionally robust loss	27
12.3.4 Convergence of SGD with Hardness Weighted Sampling and exact per-example loss vector	28
12.3.5 Proof of technical lemma 1	32

12.4 Convergence of SGD with Hardness Weighted Sampling and stale per-example loss vector	32
12.4.1 Proof of technical lemma 2	34
12.4.2 Proof of technical lemma 3	36
12.4.3 Proof of technical lemma 4	37

8 More on our experiments

8.1 Experiments on MNIST




Figure 1: Comparison of learning curves at testing (top panels) and at training (bottom panels) for ERM with SGD (blue) and DRO with our SGD with hardness weighted sampling for different values of β ($\beta = 0.1$, $\beta = 1$, $\beta = 10$, $\beta = 100$). The models are trained on an imbalanced MNIST dataset (only 1% of the digits 3 kept for training) and evaluated on the original MNIST testing dataset.

8.1.1 Implementation details

For our experiments on MNIST, we used a Wide Residual Network (WRN) [39]. The family of WRN models has proved to be very efficient and flexible, achieving state-of-the-art accuracy on several dataset. More specifically, we used WRN-16-1 [39, section 2.3].

For the optimization we used a learning rate of 0.01. No momentum or weight decay were used. No data augmentation was used. For DRO no importance sampling was used.

We used a GPU NVIDIA GeForce GTX 1070 with 8GB of memory for the experiments on MNIST.

8.1.2 Comment on the stability of the learning curves

For large values of β (here $\beta \geq 10$), instabilities appear in the **testing learning curves**, as illustrated in the top panels of Figure 1. However, the bottom left panel of Figure 1 shows that the **training loss curves** for $\beta \geq 10$ were stable there. We also observe that during iterations where instabilities appear on the **testing set**, the standard deviation of the per-example loss on the **training set** is relatively high (i.e. the hardness weighted probability is further away from the uniform distribution). This suggests that the apparent instabilities on the **testing set** are related to differences between the distributionally robust loss and the mean loss.

8.2 Experiments on BraTS 2019




Figure 2: **Illustration of the multiclass brain tumor segmentation task on a case from the BraTS 2019 dataset.** First row: the four 3D MRI sequences available in the dataset. Second row: (from left to right) labels used for training and the groups of labels used for the evaluation.

8.2.1 Implementation details

When importance sampling is used, the minimum and maximum importance weights of step 10 of Algorithm 1 are fixed to $w_{min} = 0.1$ and $w_{max} = 10$. No other values have been investigated.

The meta-parameters used for the deep learning pipeline used were determined automatically using the heuristics developed in nnUNet [18] and the code publicly available at <https://github.com/MIC-DKFZ/nnUNet>.

The 3D CNN selected for the brain tumor data is based on 3D UNet [11] with 6 levels and 32 features after the first convolution that are multiplied by 2 at each level with a maximum fixed at 320. The 3D CNN uses leaky ReLU activation, instance normalization, max-pooling downsampling operations and linear upsampling with learnable parameters.

In addition, the network is trained using the addition of the mean Dice loss and the cross entropy, and deep supervision. The default optimization step is SGD with a momentum of 0.99 and Nesterov update, a batch size of 2, and a decreasing learning rate defined for each epoch t as

$$\lambda_t = 0.01 \times \left(1 - \frac{t}{t_{max}}\right)^{0.9}$$

where t_{max} is the maximum number of epochs fixed as 1000. Note that in nnUNet, one epoch is defined as equal to 250 batches, irrespective of the size of the training dataset.




Figure 3: **Dice scores distribution on the BraTS 2019 dataset for cases from a center of TCIA (76 cases) and cases from other centers (49 cases).** This shows that the lower interquartile range of DRO for the enhancing tumor comes specifically from a lower number of poor segmentations on cases coming from The Cancer Imaging Archive (TCIA). This suggests that DRO can deal with some of the confounding biases present in the training dataset, and lead to a model that is more fair.

Table 2: **Detailed evaluation on the BraTS 2019 online validation set (125 cases).** All the models in this table were trained using the default **SGD with Nesterov momentum** of nnUNet [18]. Dice scores were computed using the BraTS online platform for evaluation <https://ipp.cbica.upenn.edu/>. ERM: Empirical Risk Minimization, DRO: Distributionally Robust Optimization, IS: Importance Sampling is used, IQR: Interquartile range.

Optimization problem	Enhancing Tumor			Whole Tumor			Tumor Core		
	Mean	Median	IQR	Mean	Median	IQR	Mean	Median	IQR
ERM	73.0	87.1	15.6	90.7	92.6	5.4	83.9	90.5	14.3
DRO $\beta = 10$	74.6	86.8	14.1	90.8	93.0	5.9	83.4	90.7	14.5
DRO $\beta = 10$ IS	75.3	86.0	13.3	90.0	91.9	7.0	82.8	89.1	14.3
DRO $\beta = 100$	73.4	86.7	14.3	90.6	92.6	6.2	84.5	90.9	13.7
DRO $\beta = 100$ IS	74.5	87.3	13.8	90.6	92.6	5.9	84.1	90.0	12.5
DRO $\beta = 1000$	74.5	84.2	33.0	89.5	91.8	5.9	71.1	87.2	41.1
DRO $\beta = 1000$ IS	72.2	85.7	15.0	90.3	92.2	6.3	81.1	89.4	15.1

A patch size of $128 \times 192 \times 128$ was used, which is not sufficient to fit the whole brain of most of the cases. As a result, a patch-based approach is used as often in medical image segmentation applications. A large number of data augmentation methods are used: random cropping of a patch, random zoom, gamma intensity augmentation, multiplicative brightness, random rotations, random mirroring along all axes, contrast augmentation, additive Gaussian noise, Gaussian blurring and simulation of low resolution.

GPUs NVIDIA Tesla V100-SXM2 with 16GB of memory were used for the experiments on brain tumor segmentation.

9 Importance sampling in Algorithm 1

In this section, we give additional details about the approximation made in the computation of the importance weights (step 9 of Algorithm 1).

Let θ be the parameters of the neural network h , $\mathbf{L} = (L_i)_{i=1}^n$ be the stale per-example loss vector, and let i be an index in the current batch I .

We start from the definition of the importance weight w_i for example i and use the formula for the hardness weighted sampling probabilities of Example 3.1.

$$\begin{aligned} w_i &= \frac{p_i^{new}}{p_i^{old}} \\ &= \frac{\exp(\beta(\mathcal{L}(h(x_i; \theta), y_i)))}{\exp(\beta(\mathcal{L}(h(x_i; \theta), y_i))) + \sum_{j \neq i} \exp(\beta L_j)} \times \frac{\sum_{j=1}^n \exp(\beta L_j)}{\exp(\beta L_i)} \\ &\approx \exp(\beta(\mathcal{L}(h(x_i; \theta), y_i) - L_i)) \end{aligned} \quad (13)$$

where we have assumed that the two sums of exponentials are approximately equal.

10 Summary of the notations used in the proofs

For the ease of reading the proofs we first summarize our notations.

10.1 Probability Theory notations

- $\Delta_n = \{(p_i)_{i=1}^n \in [0, 1]^n, \sum_i p_i = 1\}$
- Let $\mathbf{q} = (q_i) \in \Delta_n$, and f a function, we denote $\mathbb{E}_{\mathbf{q}}[f(\mathbf{x})] := \sum_{i=1}^n q_i f(x_i)$.
- Let $\mathbf{q} \in \Delta_n$, and f a function, we denote $\mathbb{V}_{\mathbf{q}}[f(\mathbf{x})] := \sum_{i=1}^n q_i \|f(x_i) - \mathbb{E}_{\mathbf{q}}[f(\mathbf{x})]\|^2$.
- $\mathbf{p}_{\text{train}}$ is the uniform training data distribution, i.e. $\mathbf{p}_{\text{train}} = (\frac{1}{n})_{i=1}^n \in \Delta_n$

10.2 Machine Learning notations

- n is the number of training examples
- d is the dimension of the output
- \mathfrak{d} is the dimension of the input
- m is the number of nodes in each layer
- training data: $\{(x_i, y_i)\}_{i=1}^n$, where for all $i \in \{1, \dots, n\}$, $x_i \in \mathbb{R}^{\mathfrak{d}}$ and $y_i \in \mathbb{R}^d$
- $h : \mathbf{x} \mapsto \mathbf{y}$ is the predictor
- θ is the set of parameters of the predictor
- For all i , $h_i : \theta \mapsto h(x_i; \theta)$ is the output of the network for example i as a function of θ
- \mathcal{L} is the objective function
- $\mathcal{L}_i : h_i \mapsto \mathcal{L}(h_i, y_i)$ is the objective function for example i .
- We denote by \mathbf{L} the function $\mathbf{L} : (h_i)_{i=1}^n \mapsto (\mathcal{L}_i(h_i))_{i=1}^n$
- $b \in \{1, \dots, n\}$ is the batch size
- $\eta > 0$ is the learning rate
- ERM is short for Empirical Risk Minimization

10.3 Distributionally Robust Optimisation notations

- For all θ , $R(\mathbf{L}(h(\theta))) = \max_{\mathbf{q} \in \Delta_n} \mathbb{E}_{\mathbf{q}}[\mathcal{L}(h(\mathbf{x}; \theta), \mathbf{y})] - \frac{1}{\beta} D_{\phi}(\mathbf{q} \parallel \mathbf{p}_{\text{train}})$ is the **Distributionally Robust Loss** evaluated at θ , where $\beta > 0$ is the parameter that adjusts the distributionally robustness. For short, we also used the terms **distributionally robust loss** or just **robust loss** for $R(\mathbf{L}(h(\theta)))$
- DRO is short for Distributionally Robust Optimisation

10.4 Miscellaneous

By abuse of notation, and similarly to [1], we use the Bachmann-Landau notations to hide constants that do not depend on our main hyper-parameters. Let f and g be two scalars, we note:

$$\begin{cases} f \leq O(g) & \iff \exists c > 0 & \text{s.t.} & f \leq cg \\ f \geq \Omega(g) & \iff \exists c > 0 & \text{s.t.} & f \geq cg \\ f = \Theta(g) & \iff \exists c_1 > 0 \text{ and } \exists c_2 > c_1 & \text{s.t.} & c_1 g \leq f \leq c_2 g \end{cases}$$

11 Proofs

11.1 Proof of Example 3.1: formula of the sampling probabilities for the KL divergence

We give here a simple proof of the formula of the sampling probabilities for the KL divergence as ϕ -divergence (i.e. $\phi : z \mapsto z \log(z) - z + 1$)

$$\forall \boldsymbol{\theta}, \quad \bar{p}(\mathbf{L}(h(\boldsymbol{\theta}))) = \text{softmax}(\beta \mathbf{L}(h(\boldsymbol{\theta})))$$

Proof: For any $\boldsymbol{\theta}$, the distributionally robust loss for the KL divergence at $\boldsymbol{\theta}$ is given by

$$\begin{aligned} R \circ \mathbf{L} \circ h(\boldsymbol{\theta}) &= \max_{\mathbf{q} \in \Delta_n} \left(\sum_{i=1}^n q_i \mathcal{L}_i \circ h_i(\boldsymbol{\theta}) - \frac{1}{\beta} \sum_{i=1}^n q_i \log(nq_i) \right) \\ &= \max_{\mathbf{q} \in \Delta_n} \sum_{i=1}^n \left(q_i \mathcal{L}_i \circ h_i(\boldsymbol{\theta}) - \frac{1}{\beta} q_i \log(nq_i) \right) \end{aligned}$$

To simplify the notations, let us denote $\mathbf{v} = (v_i)_{i=1}^n := (\mathcal{L}_i \circ h_i(\boldsymbol{\theta}))_{i=1}^n$, and $\bar{\mathbf{p}} = (\bar{p}_i)_{i=1}^n := \bar{p}(\mathbf{L}(h(\boldsymbol{\theta})))$.

Thus $\bar{p}(\mathbf{L}(h(\boldsymbol{\theta})))$ is, by definition, solution of the optimization problem

$$\arg \max_{\mathbf{q} \in \Delta_n} \sum_{i=1}^n \left(q_i v_i - \frac{1}{\beta} q_i \log(nq_i) \right) \quad (14)$$

First, let us remark that the function $q \mapsto \sum_{i=1}^n q_i \log(nq_i)$ is strictly convex on the non empty closed convex set Δ_n as a sum of strictly convex functions. This implies that the optimization (14) has a unique solution and as a result $\bar{p}(\mathbf{L}(h(\boldsymbol{\theta})))$ is well defined.

We now reformulate the optimization problem (14) as a convex smooth constrained optimization problem by writing the condition $\mathbf{q} \in \Delta_n$ as constraints.

$$\begin{aligned} \arg \max_{\mathbf{q} \in \mathbb{R}_+^n} \sum_{i=1}^n \left(q_i v_i - \frac{1}{\beta} q_i \log(nq_i) \right) \\ \text{s.t.} \quad \sum_{i=1}^n q_i = 1 \end{aligned} \quad (15)$$

There exists a Lagrange multiplier $\lambda \in \mathbb{R}$, such that the solution \bar{p} of (15) is characterized by

$$\begin{cases} \forall i \in \{1, \dots, n\}, & v_i - \frac{1}{\beta} (\log(n\bar{p}_i) + 1) + \lambda = 0 \\ & \sum_{i=1}^n \bar{p}_i = 1 \end{cases} \quad (16)$$

Which we can rewrite as

$$\begin{cases} \forall i \in \{1, \dots, n\}, & \bar{p}_i = \frac{1}{n} \exp(\beta(v_i + \lambda) - 1) \\ & \frac{1}{n} \sum_{i=1}^n \exp(\beta(v_i + \lambda) - 1) = 1 \end{cases} \quad (17)$$

The last equality gives

$$\exp(\beta\lambda - 1) = \frac{n}{\sum_{i=1}^n \exp(\beta v_i)}$$

And by replacing in the formula of the \bar{p}_i

$$\begin{aligned} \forall i \in \{1, \dots, n\}, \quad \bar{p}_i &= \frac{1}{n} \exp(\beta v_i) \exp(\beta\lambda - 1) \\ &= \frac{\exp(\beta v_i)}{\sum_{j=1}^n \exp(\beta v_j)} \end{aligned}$$

Which corresponds exactly to

$$\bar{\mathbf{p}} = \text{softmax}(\beta \mathbf{v})$$

11.2 Proof of lemma 3.1: regularity properties of R

For the ease of reading, let us first recall that given a ϕ -divergence D_ϕ that satisfies Definition 2.1, we have defined in (3)

$$\begin{aligned} R : \mathbb{R}^n &\rightarrow \mathbb{R} \\ \mathbf{v} &\mapsto \max_{\mathbf{q} \in \Delta_n} \sum_i q_i v_i - \frac{1}{\beta} D_\phi(\mathbf{q} \| \mathbf{p}_{\text{train}}) \end{aligned} \quad (18)$$

And in (4)

$$\begin{aligned} G : \mathbb{R}^n &\rightarrow \mathbb{R} \\ \mathbf{p} &\mapsto \frac{1}{\beta} D_\phi(\mathbf{p} \| \mathbf{p}_{\text{train}}) + \delta_{\Delta_n}(\mathbf{p}) \end{aligned} \quad (19)$$

where δ_{Δ_n} is the characteristic function of the closed convex set Δ_n , i.e.

$$\forall \mathbf{p} \in \mathbb{R}^n, \quad \delta_{\Delta_n}(\mathbf{p}) = \begin{cases} 0 & \text{if } \mathbf{p} \in \Delta_n \\ +\infty & \text{otherwise} \end{cases} \quad (20)$$

We now prove Lemma 3.1 on the regularity of R .

Lemma 11.1 (Regularity of R – Restated from Lemma 3.1). *Let ϕ that satisfies Definition 2.1, G and R satisfy*

$$G \text{ is } \left(\frac{n\rho}{\beta}\right)\text{-strongly convex} \quad (21)$$

$$R(\mathbf{L}(h(\boldsymbol{\theta}))) = \max_{\mathbf{q} \in \mathbb{R}^n} (\langle \mathbf{L}(h(\boldsymbol{\theta})), \mathbf{q} \rangle - G(\mathbf{q})) = G^*(\mathbf{L}(h(\boldsymbol{\theta}))) \quad (22)$$

$$R \text{ is } \left(\frac{\beta}{n\rho}\right)\text{-gradient Lipschitz continuous.} \quad (23)$$

Proof: ϕ is ρ -strongly convex on $[0, n]$ so

$$\forall x, y \in [0, n]^2, \forall \lambda \in [0, 1], \phi(\lambda x + (1 - \lambda)y) \leq \lambda \phi(x) + (1 - \lambda)\phi(y) - \frac{\rho\lambda(1 - \lambda)}{2} |y - x|^2 \quad (24)$$

Let $\mathbf{p} = (p_i)_{i=1}^n, \mathbf{q} = (q_i)_{i=1}^n \in \Delta_n$, and $\lambda \in [0, 1]$, using (24) and the convexity of δ_{Δ_n} , we obtain:

$$\begin{aligned} G(\lambda \mathbf{p} + (1 - \lambda)\mathbf{q}) &= \frac{1}{\beta n} \sum_{i=1}^n \phi(n\lambda p_i + n(1 - \lambda)q_i) + \delta_{\Delta_n}(\lambda \mathbf{p} + (1 - \lambda)\mathbf{q}) \\ &\leq \lambda G(\mathbf{p}) + (1 - \lambda)G(\mathbf{q}) - \frac{1}{\beta n} \sum_{i=1}^n \frac{\rho\lambda(1 - \lambda)}{2} |np_i - nq_i|^2 \quad (25) \\ &\leq \lambda G(\mathbf{p}) + (1 - \lambda)G(\mathbf{q}) - \frac{n\rho}{\beta} \frac{\lambda(1 - \lambda)}{2} \|\mathbf{p} - \mathbf{q}\|^2 \end{aligned}$$

This proves that G is $\frac{n\rho}{\beta}$ -strongly convex.

Since G is convex, $R = G^*$ is also convex, and $R^* = (G^*)^* = G$ [17]. We obtain (22) using Definition 3.1.

We now show that R is Frechet differentiable on \mathbb{R}^n . Let $\mathbf{v} \in \mathbb{R}^n$.

G is strongly-convex, so in particular G is strictly convex. This implies that the following optimization problem has a unique solution that we denote $\hat{\mathbf{p}}(\mathbf{v})$.

$$\arg \max_{\mathbf{q} \in \mathbb{R}^n} (\langle \mathbf{v}, \mathbf{q} \rangle - G(\mathbf{q})) \quad (26)$$

In addition

$$\begin{aligned} \hat{\mathbf{p}} \in \Delta_n \text{ solution of (26)} &\iff 0 \in \mathbf{v} - \partial G(\hat{\mathbf{p}}) \\ &\iff \mathbf{v} \in \partial G(\hat{\mathbf{p}}) \\ &\iff \hat{\mathbf{p}} \in \partial G^*(\mathbf{v}) \\ &\iff \hat{\mathbf{p}} \in \partial R(\mathbf{v}) \end{aligned}$$

where we have used [17, Proposition 6.1.2 p.39] for the third equivalence, and (22) for the last equivalence.

As a result, $\partial R(\mathbf{v}) = \{\hat{\mathbf{p}}(\mathbf{v})\}$. this implies that R admit a gradient at \mathbf{v} , and

$$\nabla_{\mathbf{v}} R(\mathbf{v}) = \hat{\mathbf{p}}(\mathbf{v}) \quad (27)$$

Since this holds for any $\mathbf{v} \in \mathbb{R}^n$, we deduce that R is Frechet differentiable on \mathbb{R}^n .

We are now ready to show that R is $\frac{\beta}{n\rho}$ -gradient Lipchitz continuous by using the following lemma [17, Theorem 6.1.2 p.280].

Lemma 11.2. *A necessary and sufficient condition for a convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ to be c -strongly convex on a convex set C is that for all $x_1, x_2 \in C$*

$$\langle s_2 - s_1, x_2 - x_1 \rangle \geq c \|x_2 - x_1\|^2 \quad \text{for all } s_i \in \partial f(x_i), i = 1, 2.$$

Using this lemma for $f = G$, $c = \frac{n\rho}{\beta}$, and $C = \Delta_n$, we obtain:

For all $\mathbf{p}_1, \mathbf{p}_2 \in \Delta_n$, for all $\mathbf{v}_1 \in \partial G(\mathbf{p}_1)$, $\mathbf{v}_2 \in \partial G(\mathbf{p}_2)$,

$$\langle \mathbf{v}_2 - \mathbf{v}_1, \mathbf{p}_2 - \mathbf{p}_1 \rangle \geq \frac{n\rho}{\beta} \|\mathbf{p}_2 - \mathbf{p}_1\|^2$$

In addition, for $i \in \{1, 2\}$, $\mathbf{v}_i \in \partial G(\mathbf{p}_i) \iff \mathbf{p}_i \in \partial R(\mathbf{v}_i) = \{\nabla_{\mathbf{v}} R(\mathbf{v}_i)\}$.

And using Cauchy Schwarz inequality

$$\|\mathbf{v}_2 - \mathbf{v}_1\| \|\mathbf{p}_2 - \mathbf{p}_1\| \geq \langle \mathbf{v}_2 - \mathbf{v}_1, \mathbf{p}_2 - \mathbf{p}_1 \rangle$$

We conclude that

$$\frac{n\rho}{\beta} \|\nabla_{\mathbf{v}} R(\mathbf{v}_2) - \nabla_{\mathbf{v}} R(\mathbf{v}_1)\| \leq \|\mathbf{v}_2 - \mathbf{v}_1\|$$

Which implies that R is $\frac{\beta}{n\rho}$ -gradient Lipchitz continuous.

11.3 Proof of Lemma 3.2: formula of the distributionally robust loss gradient

We prove Lemma 3.2 that we restate here for the ease of reading.

Lemma 11.3 (Stochastic Gradient of the Distributionally Robust Loss – Restated from Lemma 3.2). *For all θ , we have*

$$\bar{\mathbf{p}}(\mathbf{L}(h(\theta))) = \nabla_{\mathbf{v}} R(\mathbf{L}(h(\theta))) \quad (28)$$

$$\nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta) = \mathbb{E}_{\mathbf{p}(\mathcal{L}(h(\theta)))} [\nabla_{\theta} \mathcal{L}(h(x; \theta), y)] \quad (29)$$

where $\nabla_{\mathbf{L}} R$ is the gradient of R with respect to its input.

Proof: For a given θ , equality (28) is a special case of (27) for $\mathbf{v} = \mathcal{L}(\mathbf{h}(\theta))$.

Then using the chain rule and (28),

$$\begin{aligned}\nabla_{\theta}(R \circ L \circ h)(\theta) &= \sum_{i=1}^n \frac{\partial R}{\partial v_i}(L \circ h(\theta)) \nabla_{\theta}(\mathcal{L}_i \circ h_i)(\theta) \\ &= \sum_{i=1}^n \bar{p}_i(L(h(\theta))) \nabla_{\theta}(\mathcal{L}_i \circ h_i)(\theta) \\ &= \mathbb{E}_{\bar{\mathbf{p}}(\mathcal{L}(\mathbf{h}(\theta)))} [\nabla_{\theta} \mathcal{L}(h(\mathbf{x}; \theta), y)]\end{aligned}$$

Which concludes the proof.

11.4 Proof of Theorem 3.2: Distributionally Robust Optimization as Principled Hard Example Mining

Let D_{ϕ} an ϕ -divergence satisfying Definition 2.1, and $\mathbf{v} = (v_i)_{i=1}^n \in \mathbb{R}^n$. \mathbf{v} will play the role of a generic loss vector.

ϕ is strongly convex, and Δ_n is closed and convex, so the following optimization problem has one and only one solution

$$\max_{\mathbf{p}=(p_i)_{i=1}^n \in \Delta_n} \langle \mathbf{v}, \mathbf{p} \rangle - \frac{1}{\beta n} \sum_{i=1}^n \phi(np_i) \quad (30)$$

Making the constraints associated with $p \in \Delta_n$ explicit, this can be rewritten as

$$\begin{aligned}\max_{\mathbf{p}=(p_i)_{i=1}^n \in \mathbb{R}^n} \quad & \langle \mathbf{v}, \mathbf{p} \rangle - \frac{1}{\beta n} \sum_{i=1}^n \phi(np_i) \\ \text{s.t.} \quad & \forall i \in \{1, \dots, n\}, p_i \geq 0 \\ & \sum_{i=1}^n p_i = 1\end{aligned} \quad (31)$$

There exists KKT multipliers $\lambda \in \mathbb{R}$ and $\forall i, \mu_i \geq 0$ such that the solution $\bar{\mathbf{p}} = (\bar{p}_i)_{i=1}^n$ satisfies

$$\begin{cases} \forall i \in \{1, \dots, n\}, & v_i - \frac{1}{\beta} \phi'(n\bar{p}_i) + \lambda - \mu_i = 0 \\ \forall i \in \{1, \dots, n\}, & \mu_i p_i = 0 \\ \forall i \in \{1, \dots, n\}, & p_i \geq 0 \\ & \sum_{i=1}^n \bar{p}_i = 1 \end{cases} \quad (32)$$

Since ϕ is continuously differentiable and strongly convex, we have $(\phi')^{-1} = (\phi^*)'$, where ϕ^* is the Fenchel conjugate of ϕ [see 17, Proposition 6.1.2]. As a result, (32) can be rewritten as

$$\begin{cases} \forall i \in \{1, \dots, n\}, & \bar{p}_i = \frac{1}{n} (\phi^*)'(\beta(v_i + \lambda - \mu_i)) \\ \forall i \in \{1, \dots, n\}, & \mu_i p_i = 0 \\ \forall i \in \{1, \dots, n\}, & p_i \geq 0 \\ & \frac{1}{n} \sum_{i=1}^n (\phi^*)'(\beta(v_i + \lambda - \mu_i)) = 1 \end{cases} \quad (33)$$

We now show that the KKT multipliers are uniquely defined.

The μ_i 's are uniquely defined by \mathbf{v} and λ :

Since $\forall i \in \{1, \dots, n\}$, $\mu_i p_i = 0$, $p_i \geq 0$ and $\mu_i \geq 0$, for all $\forall i \in \{1, \dots, n\}$, either $p_i = 0$ or $\mu_i = 0$.

In the case $p_i = 0$, using (33) it comes $(\phi^*)'(\beta(v_i + \lambda - \mu_i)) = 0$.

According to Definition 2.1, ϕ is strongly convex and continuously differentiable, so ϕ' and $(\phi^*)' = (\phi')^{-1}$ are continuous and strictly increasing functions. As a result, it exists a unique μ_i (dependent to \mathbf{v} and λ) such that:

$$(\phi^*)'(\beta(v_i + \lambda - \mu_i)) = 0$$

And (33) can be rewritten as

$$\left\{ \begin{array}{l} \forall i \in \{1, \dots, n\}, \quad \bar{p}_i = \text{ReLU} \left(\frac{1}{n} (\phi^*)'(\beta(v_i + \lambda)) \right) = \frac{1}{n} \text{ReLU}((\phi^*)'(\beta(v_i + \lambda))) \\ \frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v_i + \lambda))) = 1 \end{array} \right. \quad (34)$$

λ is uniquely defined by \mathbf{v} and a continuous function of \mathbf{v} :

Let $\lambda \in \mathbb{R}$ that satisfies (34).

We have $\frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v_i + \lambda))) = 1$. So there exists at least one index i_0 such that

$$\text{ReLU}((\phi^*)'(\beta(v_{i_0} + \lambda))) = (\phi^*)'(\beta(v_{i_0} + \lambda)) \geq 1$$

Since $(\phi^*)^{-1}$ is continuous and strictly increasing, $\lambda' \mapsto \text{ReLU}((\phi^*)'(\beta(v_{i_0} + \lambda')))$ is continuous and strictly increasing on a neighborhood of λ .

In addition ReLU is continuous and increasing, so for all $i \in \{1, \dots, n\}$, $\lambda' \mapsto \text{ReLU}((\phi^*)'(\beta(v_i + \lambda')))$ is a continuous and increasing function.

As a result, $\lambda' \mapsto \frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v_i + \lambda')))$ is a continuous function that is increasing on \mathbb{R} , and strictly increasing on a neighborhood of λ .

This implies that λ is uniquely defined by \mathbf{v} , and that $\mathbf{v} \mapsto \lambda(\mathbf{v})$ is continuous.

Link between Hard Weighted Sampling and Hard Example Mining:

For any pseudo loss vector $\mathbf{v} = (v_i)_{i=1}^n \in \mathbb{R}^n$, there exists a unique λ and a unique $\bar{\mathbf{p}}$ that satisfies (34), so we can define the mapping:

$$\begin{aligned} \bar{\mathbf{p}} : \mathbb{R}^n &\rightarrow \Delta_n \\ \mathbf{v} &\mapsto \bar{\mathbf{p}}(\mathbf{v}; \lambda(\mathbf{v})) \end{aligned} \quad (35)$$

where for all \mathbf{v} , $\lambda(\mathbf{v})$ is the unique $\lambda \in \mathbb{R}$ satisfying (34).

We will now demonstrate that each $\bar{p}_{i_0}(\mathbf{v})$ for $i_0 \in \{1, \dots, n\}$ is an increasing function of v_{i_0} and a decreasing function of the v_i for $i \neq i_0$. Without loss of generality we assume $i_0 = 1$.

Let $\mathbf{v} = (v_i)_{i=1}^n \in \mathbb{R}^n$, and $\epsilon > 0$.

Let us define $\mathbf{v}' = (v'_i)_{i=1}^n \in \mathbb{R}^n$, such that $v'_1 = v_1 + \epsilon$ and $\forall i \in \{2, \dots, n\}$, $v'_i = v_i$.

Similarly as in the proof of the uniqueness of λ above, we can show that there exists $\eta > 0$ such that the function

$$F : \lambda' \mapsto \frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v_i + \lambda')))$$

is continuous and strictly increasing on $[\lambda(\mathbf{v}) - \eta, \lambda(\mathbf{v}) + \eta]$, and $F(\lambda(\mathbf{v})) = 1$.

$\mathbf{v} \mapsto \lambda(\mathbf{v})$ is continuous, so for ϵ small enough $\lambda(\mathbf{v}') \in [\lambda(\mathbf{v}) - \eta, \lambda(\mathbf{v}) + \eta]$.

Let us now prove by contradiction that $\lambda(\mathbf{v}') \leq \lambda(\mathbf{v})$. Therefore, let us assume that $\lambda(\mathbf{v}') > \lambda(\mathbf{v})$. Then, as $\text{ReLU} \circ (\phi^*)'$ is an increasing function and F is strictly increasing on $[\lambda(\mathbf{v}) - \eta, \lambda(\mathbf{v}) + \eta]$,

and $\epsilon > 0$ we obtain

$$\begin{aligned}
1 &= \frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v'_i + \lambda(\mathbf{v}')))) \\
&\geq \frac{1}{n} \sum_{i=1}^n \text{ReLU}((\phi^*)'(\beta(v_i + \lambda(\mathbf{v})))) \\
&\geq F(\lambda(\mathbf{v}')) \\
&> F(\lambda(\mathbf{v})) \\
&> 1
\end{aligned}$$

which is a contradiction. As a result

$$\lambda(\mathbf{v}') \leq \lambda(\mathbf{v}) \tag{36}$$

Using equations (34) and (36), and the fact that $\text{ReLU} \circ (\phi^*)'$ is an increasing function, we obtain for all $i \in \{2, \dots, n\}$

$$\begin{aligned}
\bar{p}_i(\mathbf{v}') &= \frac{1}{n} \text{ReLU}((\phi^*)'(\beta(v'_i + \lambda(\mathbf{v}')))) \\
&= \frac{1}{n} \text{ReLU}((\phi^*)'(\beta(v_i + \lambda(\mathbf{v})))) \\
&\leq \frac{1}{n} \text{ReLU}((\phi^*)'(\beta(v_i + \lambda(\mathbf{v})))) \\
&\leq \bar{p}_i(\mathbf{v})
\end{aligned} \tag{37}$$

In addition

$$\sum_{i=1}^n \bar{p}_i(\mathbf{v}') = 1 = \sum_{i=1}^n \bar{p}_i(\mathbf{v})$$

So necessarily

$$\bar{p}_1(\mathbf{v}') \geq \bar{p}_1(\mathbf{v}) \tag{38}$$

This holds for any i_0 and any \mathbf{v} , which concludes the proof.

12 Proof of Theorem 3.1: convergence of SGD with Hardness Weighted Sampling for Over-parameterized Deep Neural Networks with ReLU

In this section, we provide the proof of Theorem 3.1. This generalizes the convergence of SGD for empirical risk minimization in [1, Theorem 2] to the convergence of SGD and our proposed hardness weighted sampler for distributionally robust optimization.

We start by describing in details the assumptions made for our convergence result in Section 12.1.

In Section 12.2, we restate Theorem 3.1 using the assumptions and notations previously introduced in Section 10.

In Section 12.3, we give the proof of the convergence theorem. We focus on providing theoretical tools that could be used to generalize any convergence result for ERM using SGD to DRO using SGD with hardness weighted sampling as described in Algorithm 1.

12.1 Assumptions

Our analysis is based on the results developed in [1] which is a simplified version of [2]. Improving on those theoretical results would automatically improve our results as well.

In the following we state our assumptions on the neural network h , and the per-example loss function \mathcal{L} .

Assumption 12.1 (Deep Neural Network). *In this section, we use the following notations and assumptions similar to [1]:*

- h is a fully connected neural network with $L + 2$ layers, ReLU as activation functions, and m nodes in each hidden layer

- For all $i \in \{1, \dots, n\}$, we denote $h_i : \theta \mapsto h_i(x_i; \theta)$ the d -dimensional output scores of h applied to example x_i of dimension \mathfrak{d} .
- For all $i \in \{1, \dots, n\}$, we denote $\mathcal{L}_i : h \mapsto \mathcal{L}(h, y_i)$ where y_i is the ground truth associated to example i .
- $\theta = (\theta_l)_{l=0}^{L+1}$ is the set of parameters of the neural network h , where θ_l is the set of weights for layer l with $\theta_0 \in \mathbb{R}^{\mathfrak{d} \times m}$, $\theta_{L+1} \in \mathbb{R}^{m \times d}$, and $\theta_l \in \mathbb{R}^{m \times m}$ for any other l .
- (Data separation) It exists $\delta > 0$ such that for all $i, j \in \{1, \dots, n\}$, if $i \neq j$, $\|x_i - x_j\| \geq \delta$.
- We assume $m \geq \Omega(d \times \text{poly}(n, L, \delta^{-1}))$ for some sufficiently large polynomial poly , and $\delta \geq O(\frac{1}{L})$. We refer the reader to [1] for details about the polynomial poly .
- The parameters $\theta = (\theta_l)_{l=0}^{L+1}$ are initialized at random such that:
 - $[\theta_0]_{i,j} \sim \mathcal{N}(0, \frac{2}{m})$ for every $(i, j) \in \{1, \dots, m\} \times \{1, \dots, \mathfrak{d}\}$
 - $[\theta_l]_{i,j} \sim \mathcal{N}(0, \frac{2}{m})$ for every $(i, j) \in \{1, \dots, m\}^2$ and $l \in \{1, \dots, L\}$
 - $[\theta_{L+1}]_{i,j} \sim \mathcal{N}(0, \frac{1}{d})$ for every $(i, j) \in \{1, \dots, d\} \times \{1, \dots, m\}$

Assumption 12.2 (Regularity of \mathcal{L}). For all i , \mathcal{L}_i is a $C(\nabla \mathcal{L})$ -gradient Lipschitz continuous, $C(\mathcal{L})$ -Lipschitz continuous, and bounded (potentially non-convex) function.

12.2 Convergence theorem (restated)

In this section, we restate the convergence Theorem 3.1 for SGD with hardness weighted sampling and stale per-example loss vector.

As an intermediate step, we will first generalize the convergence of SGD in [1, Theorem 2] to the minimization of the distributionally robust loss using SGD and an *exact* hardness weighted sampling (10), i.e. with an exact per-example loss vector.

Theorem 12.1 (Convergence of SGD with Hardness Weighted Sampling and exact per-example loss vector). Let batch size $1 \leq b \leq n$, and $\epsilon > 0$. Suppose there exists constants $C_1, C_2, C_3 > 0$ such that the number of hidden units satisfies $m \geq C_1(d\epsilon^{-1} \times \text{poly}(n, L, \delta^{-1}))$, $\delta \geq (\frac{C_2}{L})$, and the learning rate be $\eta_{exact} = C_3 \left(\min \left(1, \frac{\alpha n^2 \rho}{\beta C(\mathcal{L})^2 + 2n\rho C(\nabla \mathcal{L})} \right) \times \frac{b\delta d}{\text{poly}(n, L)m \log^2(m)} \right)$. There exists constants $C_4, C_5 > 0$ such that with probability at least $1 - \exp(-C_4 \log^2(m))$ over the randomness of the initialization and the mini-batches, SGD with hardness weighted sampling and exact per-example loss vector guarantees $\|\nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta)\| \leq \epsilon$ after $T = C_5 \left(\frac{Ln^3}{\eta_{exact} \delta \epsilon^2} \right)$ iterations.

The proof can be found in Appendix 12.3.4.

$\alpha = \min_{\theta} \min_i \bar{p}_i(\mathbf{L}(\theta))$ is a lower bound on the sampling probabilities. For the Kullback-Leibler ϕ -divergence, and for any ϕ -divergence satisfying Definition 2.1 with a robustness parameter β small enough, we have $\alpha > 0$. We refer the reader to [1, Theorem 2] for the values of the constants C_1, C_2, C_3, C_4, C_5 and the definitions of the polynomials.

Compared to [1, Theorem 2] only the learning rate differs. The $\min(1, \cdot)$ operation in the formula for η_{exact} allows us to guarantee that $\eta_{exact} \leq \eta'$ where η' is the learning rate of [1, Theorem 2].

It is worth noting that for the KL ϕ -divergence, $\rho = \frac{1}{n}$. In addition, in the limit $\beta \rightarrow 0$, which corresponds to ERM, we have $\alpha \rightarrow \frac{1}{n}$. As a result, we recover exactly Theorem 2 of [1] as extended in their Appendix A for any smooth loss function \mathcal{L} that satisfies assumption 12.2 with $C(\nabla \mathcal{L}) = 1$.

We now restate the convergence of SGD with hardness weighted sampling and a stale per-example loss vector as in Algorithm 1.

Theorem 12.2 (Convergence of SGD with Hardness Weighted Sampling and stale per-example loss vector). Let batch size $1 \leq b \leq n$, and $\epsilon > 0$. Under the conditions of Theorem 12.1, the same notations, and with the learning rate $\eta_{stale} = C_6 \min \left(1, \frac{\alpha \rho d^{3/2} \delta b \log(\frac{1}{1-\alpha})}{\beta C(\mathcal{L}) A(\nabla \mathcal{L}) L m^{3/2} n^{3/2} \log^2(m)} \right) \times \eta_{exact}$

for a constant $C_6 > 0$. With probability at least $1 - \exp(-C_4(\log^2(m)))$ over the randomness of the initialization and the mini-batches, SGD with hardness weighted sampling and stale per-example loss vector guarantees $\|\nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta)\| \leq \epsilon$ after $T = C_5 \left(\frac{Ln^3}{\eta_{stale} \delta \epsilon^2} \right)$ iterations.

The proof can be found in Appendix 12.4.

$C(\mathcal{L}) > 0$ is a constant such that \mathcal{L} is $C(\mathcal{L})$ -Lipschitz continuous, and $A(\nabla \mathcal{L}) > 0$ is a constant that bounds the gradient of \mathcal{L} with respect to its input. $C(\mathcal{L})$ and $A(\nabla \mathcal{L})$ are guaranteed to exist under assumptions 12.1.

Compared to Theorem 12.1 only the learning rate differs. Similarly to Theorem 12.1, when β tends to zero we recover Theorem 2 of [1].

It is worth noting that when β increases, $\frac{\alpha \rho d^{3/2} \delta b \log(\frac{1}{1-\alpha})}{\beta C(\mathcal{L}) A(\nabla \mathcal{L}) L m^{3/2} n^{3/2} \log^2(m)}$ decreases. This implies that η_{stale} decreases faster than η_{exact} when β increases. This was to be expected since the error that is made by using the stale per-example loss vector instead of the exact loss increases when β increases.

12.3 Proofs of convergence

In this section, we prove the results of Theorem 12.1 and 12.2.

For the ease of reading the proof, we remind here the chain rules for the distributionally robust loss that we are going to use intensively in the following proofs.

Chain rule for the derivative of $R \circ \mathbf{L}$ with respect to the network outputs h :

$$\begin{aligned} \nabla_h(R \circ \mathbf{L})(h(\theta)) &= (\nabla_{h_i}(R \circ \mathbf{L})(h(\theta)))_{i=1}^n \\ \forall i \in \{1, \dots, n\}, \quad \nabla_{h_i}(R \circ \mathbf{L})(h(\theta)) &= \sum_{j=1}^n \frac{\partial R}{\partial v_j}(\mathbf{L}(h(\theta))) \nabla_{h_i} \mathcal{L}_j(h_j(\theta)) \\ &= \bar{p}_i(\mathbf{L}(h(\theta))) \nabla_{h_i} \mathcal{L}_i(h_i(\theta)) \end{aligned} \quad (39)$$

Chain rule for the derivative of $R \circ \mathbf{L} \circ h$ with respect to the network parameters θ :

$$\begin{aligned} \nabla_{\theta}(R \circ \mathbf{L} \circ h)(\theta) &= \sum_{i=1}^n \nabla_{\theta} h_i(\theta) \nabla_{h_i}(R \circ \mathbf{L})(h(\theta)) \\ &= \sum_{i=1}^n \bar{p}_i(\mathbf{L}(h(\theta))) \nabla_{\theta} h_i(\theta) \nabla_{h_i} \mathcal{L}_i(h_i(\theta)) \\ &= \sum_{i=1}^n \bar{p}_i(\mathbf{L}(h(\theta))) \nabla_{\theta} (\mathcal{L}_i \circ h_i)(\theta) \end{aligned} \quad (40)$$

where for all $i \in \{1, \dots, n\}$, $\nabla_{\theta} h_i(\theta)$ is the transpose of the Jacobian matrix of h_i as a function of θ .

12.3.1 Proof that $R \circ \mathbf{L}$ is one-sided gradient Lipschitz

This property that $R \circ \mathbf{L}$ is one-sided gradient Lipschitz is a key element for the proof of the semi-smoothness theorem for the distributionally robust loss Theorem 12.3.

Under Definition 2.1 for the ϕ -divergence, we have shown that R is $\frac{\beta}{n\rho}$ -gradient Lipschitz continuous (Lemma 3.1). And under assumption 12.2, for all i , \mathcal{L}_i is $C(\mathcal{L})$ -Lipschitz continuous and $C(\nabla \mathcal{L})$ -gradient Lipschitz continuous.

Let $\mathbf{z} = (z_i)_{i=1}^n, \mathbf{z}' = (z'_i)_{i=1}^n \in \mathbb{R}^{dn}$.

We want to show that $R \circ \mathbf{L}$ is one-sided gradient Lipschitz, i.e. we want to prove the existence of a constant $C > 0$, independent to \mathbf{z} and \mathbf{z}' , such that:

$$\langle \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}) - \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}'), \mathbf{z} - \mathbf{z}' \rangle \leq C \|\mathbf{z} - \mathbf{z}'\|^2$$

We have

$$\begin{aligned}
& \langle \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}) - \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}'), \mathbf{z} - \mathbf{z}' \rangle \\
&= \sum_{i=1}^n \langle \nabla_{z_i}(R \circ \mathbf{L})(\mathbf{z}) - \nabla_{z_i}(R \circ \mathbf{L})(\mathbf{z}'), z_i - z'_i \rangle \\
&= \sum_{i=1}^n \langle \bar{p}_i(\mathbf{L}(\mathbf{z})) \nabla_{z_i} \mathcal{L}_i(z_i) - \bar{p}_i(\mathbf{L}(\mathbf{z}')) \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle \\
&= \sum_{i=1}^n \bar{p}_i(\mathbf{L}(\mathbf{z})) \langle \nabla_{z_i} \mathcal{L}_i(z_i) - \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle \\
&\quad + \sum_{i=1}^n (\bar{p}_i(\mathbf{L}(\mathbf{z})) - \bar{p}_i(\mathbf{L}(\mathbf{z}'))) \langle \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle
\end{aligned} \tag{41}$$

Where for all $i \in \{1, \dots, n\}$ we have used the chain rule

$$\nabla_{z_i}(R \circ \mathbf{L})(\mathbf{z}) = \sum_{j=1}^n \frac{\partial R}{\partial v_j}(\mathcal{L}(\mathbf{z})) \nabla_{z_i} \mathcal{L}_j(z_j) = \bar{p}_i(\mathbf{L}(\mathbf{z})) \nabla_{z_i} \mathcal{L}_i(z_i)$$

Let

$$A = \left| \sum_{i=1}^n \bar{p}_i(\mathbf{L}(\mathbf{z})) \langle \nabla_{z_i} \mathcal{L}_i(z_i) - \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle \right|$$

For all i , \mathcal{L}_i is $C(\nabla \mathcal{L})$ -gradient Lipschitz continuous, so using Cauchy-Schwarz inequality

$$A \leq \sum_{i=1}^n C(\nabla \mathcal{L}) \|z_i - z'_i\|^2 = C(\nabla \mathcal{L}) \|\mathbf{z} - \mathbf{z}'\|^2 \tag{42}$$

Let

$$B = \left| \sum_{i=1}^n (\bar{p}_i(\mathbf{L}(\mathbf{z})) - \bar{p}_i(\mathbf{L}(\mathbf{z}'))) \langle \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle \right|$$

Using the triangular inequality:

$$\begin{aligned}
B &\leq \left| \sum_{i=1}^n (\bar{p}_i(\mathbf{L}(\mathbf{z})) - \bar{p}_i(\mathbf{L}(\mathbf{z}'))) (\mathcal{L}_i(z_i) - \mathcal{L}_i(z'_i)) \right| \\
&\quad + \left| \sum_{i=1}^n (\bar{p}_i(\mathbf{L}(\mathbf{z})) - \bar{p}_i(\mathbf{L}(\mathbf{z}'))) (\mathcal{L}_i(z'_i) + \langle \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle - \mathcal{L}_i(z_i)) \right| \\
&\leq \langle \nabla_{\mathbf{L}} R(\mathbf{L}(\mathbf{z})) - \nabla_{\mathbf{L}} R(\mathbf{L}(\mathbf{z}')), \mathbf{L}(\mathbf{z}) - \mathbf{L}(\mathbf{z}') \rangle \\
&\quad + 2 \sum_{i=1}^n \left| \mathcal{L}_i(z'_i) + \langle \nabla_{z_i} \mathcal{L}_i(z'_i), z_i - z'_i \rangle - \mathcal{L}_i(z_i) \right| \\
&\leq \frac{\beta}{n\rho} \|\mathbf{L}(\mathbf{z}) - \mathbf{L}(\mathbf{z}')\|^2 + 2 \frac{C(\nabla \mathcal{L})}{2} \|\mathbf{z} - \mathbf{z}'\|^2 \\
&\leq \left(\frac{\beta C(\mathcal{L})^2}{n\rho} + C(\nabla \mathcal{L}) \right) \|\mathbf{z} - \mathbf{z}'\|^2
\end{aligned} \tag{43}$$

Combining equations (41), (42) and (43) we finally obtain

$$\langle \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}) - \nabla_{\mathbf{z}}(R \circ \mathbf{L})(\mathbf{z}'), \mathbf{z} - \mathbf{z}' \rangle \leq \left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \|\mathbf{z} - \mathbf{z}'\|^2 \tag{44}$$

From there, we can obtain the following inequality that will be used for the proof of the semi-smoothness property in Theorem 12.3

$$\begin{aligned}
& R(\mathbf{L}(z')) - R(\mathbf{L}(z)) - \langle \nabla_z (R \circ \mathbf{L})(z), z' - z \rangle \\
&= \int_{t=0}^1 \langle \nabla_z (R \circ \mathbf{L})(z + t(z' - z)) - \nabla_z (R \circ \mathbf{L})(z), z' - z \rangle dt \\
&\leq \frac{1}{2} \left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \|z - z'\|^2
\end{aligned} \tag{45}$$

12.3.2 Semi-smoothness property of the distributionally robust loss

We prove the following lemma which is a generalization of Theorem 4 in [1] for the distributionally robust loss.

Theorem 12.3 (Semi-smoothness of the distributionally robust loss).

Let $\omega \in \left[\Omega \left(\frac{d^{3/2}}{m^{3/2} L^{3/2} \log^{3/2}(m)} \right), O \left(\frac{1}{L^{4.5} \log^3(m)} \right) \right]$, and the $\boldsymbol{\theta}^{(0)}$ being initialized randomly as described in assumption 12.1. With probability at least $1 - \exp(-\Omega(m\omega^{3/2}L))$ over the initialization, we have for all $\boldsymbol{\theta}, \boldsymbol{\theta}' \in (\mathbb{R}^{m \times m})^L$ with $\|\boldsymbol{\theta} - \boldsymbol{\theta}^{(0)}\|_2 \leq \omega$, and $\|\boldsymbol{\theta}' - \boldsymbol{\theta}\|_2 \leq \omega$

$$\begin{aligned}
R(\mathbf{L}(h(\boldsymbol{\theta}')) &\leq R(\mathbf{L}(h(\boldsymbol{\theta}))) + \langle \nabla_{\boldsymbol{\theta}} (R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}), \boldsymbol{\theta}' - \boldsymbol{\theta} \rangle \\
&+ \|\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,1} O \left(\frac{L^2 \omega^{1/3} \sqrt{m \log(m)}}{\sqrt{d}} \right) \|\boldsymbol{\theta}' - \boldsymbol{\theta}\|_{2,\infty} \\
&+ O \left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \frac{nL^2 m}{d} \right) \|\boldsymbol{\theta}' - \boldsymbol{\theta}\|_{2,\infty}^2
\end{aligned} \tag{46}$$

where for all layer $l \in \{1, \dots, L\}$, $\boldsymbol{\theta}_l$ is the vector of parameters for layer l , and

$$\begin{aligned}
\|\boldsymbol{\theta}' - \boldsymbol{\theta}\|_{2,\infty} &= \max_l \|\boldsymbol{\theta}'_l - \boldsymbol{\theta}_l\|_2 \\
\|\boldsymbol{\theta}' - \boldsymbol{\theta}\|_{2,\infty}^2 &= \left(\max_l \|\boldsymbol{\theta}'_l - \boldsymbol{\theta}_l\|_2 \right)^2 = \max_l \|\boldsymbol{\theta}'_l - \boldsymbol{\theta}_l\|_2^2 \\
\|\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,1} &= \sum_{i=1}^n \|\nabla_{h_i} (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_2 \\
&= \sum_{i=1}^n \left\| \bar{p}_i(\mathbf{L}(h(\boldsymbol{\theta}))) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2 \quad (\text{chain rule (39)})
\end{aligned}$$

To compare this semi-smoothness result to the one in [1, Theorem 4], let us first remark that

$$\|\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,1} \leq \sqrt{n} \|\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,2}$$

As a result, our result is analogous to [1, Theorem 4], up to an additional multiplicative factor $\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right)$ in the last term of the right-hand side. It is worth noting that there is also implicitly an additional multiplicative factor $C(\nabla \mathcal{L})$ in Theorem 3 of [1] since [1] make the assumption that $C(\nabla \mathcal{L}) = 1$ [see 1, Appendix A].

Let $\boldsymbol{\theta}, \boldsymbol{\theta}' \in (\mathbb{R}^{m \times m})^L$ verifying the conditions of Theorem 12.3.

Let $A = R(\mathbf{L}(h(\boldsymbol{\theta}')) - R(\mathbf{L}(h(\boldsymbol{\theta}))) - \langle \nabla_{\boldsymbol{\theta}} (R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}), \boldsymbol{\theta}' - \boldsymbol{\theta} \rangle$, the quantity we want to bound.

Using (45) for $z = h(\boldsymbol{\theta})$ and $z' = h(\boldsymbol{\theta}')$, we obtain

$$\begin{aligned}
A &\leq \frac{1}{2} \left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \|h(\boldsymbol{\theta}') - h(\boldsymbol{\theta})\|_2^2 \\
&+ \langle \nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta})), h(\boldsymbol{\theta}') - h(\boldsymbol{\theta}) \rangle \\
&- \langle \nabla_{\boldsymbol{\theta}} (R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}), \boldsymbol{\theta}' - \boldsymbol{\theta} \rangle
\end{aligned} \tag{47}$$

Then using the chain rule (40)

$$\begin{aligned}
A &\leq \frac{1}{2} \left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \|h(\boldsymbol{\theta}') - h(\boldsymbol{\theta})\|_2^2 \\
&\quad + \sum_{i=1}^n \langle \nabla_{h_i}(R \circ \mathbf{L})(h(\boldsymbol{\theta})), h_i(\boldsymbol{\theta}') - h_i(\boldsymbol{\theta}) - (\nabla_{\boldsymbol{\theta}} h_i(\boldsymbol{\theta}))^T (\boldsymbol{\theta}' - \boldsymbol{\theta}) \rangle
\end{aligned} \tag{48}$$

For all $i \in \{1, \dots, n\}$, let us denote $\check{\text{loss}}_i := \nabla_{h_i}(R \circ \mathbf{L})(h(\boldsymbol{\theta}))$ to match the notations used in [1] for the derivative of the loss with respect to the output of the network for example i of the training set.

With this notation, we obtain exactly equation (11.3) in [1] up to the multiplicative factor $\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right)$ for the distributionally robust loss.

From there the proof of Theorem 4 in [1] being independent to the formula for $\check{\text{loss}}_i$, we can conclude the proof of our Theorem 12.3 as in [1, Appendix A].

12.3.3 Gradient bounds for the distributionally robust loss

We prove the following lemma which is a generalization of Theorem 3 in [1] for the distributionally robust loss.

Theorem 12.4 (Gradient Bounds for the Distributionally Robust Loss).

Let $\omega \in O\left(\frac{\delta^{3/2}}{n^{9/2} L^6 \log^3(m)}\right)$, and $\boldsymbol{\theta}^{(0)}$ being initialized randomly as described in assumption 12.1. With probability as least $1 - \exp(-\Omega(m\omega^{3/2}L))$ over the initialization, we have for all $\boldsymbol{\theta} \in (\mathbb{R}^{m \times m})^L$ with $\|\boldsymbol{\theta} - \boldsymbol{\theta}^{(0)}\|_2 \leq \omega$

$$\begin{aligned}
&\forall i \in \{1, \dots, n\}, \forall l \in \{1, \dots, L\}, \forall \hat{\mathbf{L}} \in \mathbb{R}^n \\
&\quad \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_i} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \leq O\left(\frac{m}{d} \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2\right) \\
&\forall l \in \{1, \dots, L\}, \forall \hat{\mathbf{L}} \in \mathbb{R}^n \\
&\quad \left\| \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_i} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \leq O\left(\frac{mn}{d} \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2\right) \\
&\quad \left\| \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_L} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \geq \Omega\left(\frac{m\delta}{dn^2} \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2\right)
\end{aligned} \tag{49}$$

It is worth noting that the loss vector $\hat{\mathbf{L}}$ used for computing the robust probabilities $\bar{p}(\hat{\mathbf{L}}) = \left(\bar{p}_i(\hat{\mathbf{L}})\right)_{i=1}^n$ does not have to be equal to $\mathbf{L}(h(\boldsymbol{\theta}))$.

We will use this for the proof of the Robust SGD with stale per-example loss vector.

The adaptation of the proof of Theorem 3 in [1] is straightforward.

Let $\boldsymbol{\theta} \in (\mathbb{R}^{m \times m})^L$ satisfying the conditions of Theorem 12.4, and $\hat{\mathbf{L}} \in \mathbb{R}^n$.

Let us denote $\mathbf{v} := \left(\bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right)_{i=1}^n$, applying the proof of Theorem 3 in [1] to our \mathbf{v} gives:

$$\begin{aligned} & \forall i \in \{1, \dots, n\}, \forall l \in \{1, \dots, L\}, \\ & \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_i} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \leq O \left(\frac{m}{d} \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \right) \\ & \forall l \in \{1, \dots, L\}, \forall \hat{\mathbf{L}} \in \mathbb{R}^n \\ & \left\| \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_i} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \leq O \left(\frac{mn}{d} \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \right) \\ & \left\| \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_L} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \geq \Omega \left(\frac{m\delta}{dn} \max_i \left(\left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \right) \right) \end{aligned}$$

In addition

$$\max_i \left(\left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \right) \geq \frac{1}{n} \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2$$

This allows us to conclude the proof of our Theorem 12.4.

12.3.4 Convergence of SGD with Hardness Weighted Sampling and exact per-example loss vector

We can now prove Theorem 12.1.

Similarly to the proof of the convergence of SGD for the mean loss (Theorem 2 in [1]), the convergence of SGD for the distributionally robust loss will mainly rely on the semi-smoothness property (Theorem 12.3) and the gradient bound (Theorem 12.4) that we have proved previously for the distributionally robust loss.

Let $\boldsymbol{\theta} \in (\mathbb{R}^{m \times m})^L$ satisfying the conditions of Theorem 12.1, and $\hat{\mathbf{L}}$ be the exact per-example loss vector at $\boldsymbol{\theta}$, i.e.

$$\hat{\mathbf{L}} = \left(\mathcal{L}_i(h_i(\boldsymbol{\theta})) \right)_{i=1}^n \quad (50)$$

For the batch size $b \in \{1, \dots, n\}$, let $S = \{i_j\}_{j=1}^b$ a batch of indices drawn from $\bar{p}(\hat{\mathbf{L}})$ without replacement, i.e.

$$\forall j \in \{1, \dots, b\}, i_j \stackrel{\text{i.i.d.}}{\sim} \bar{p}(\hat{\mathbf{L}}) \quad (51)$$

Let $\boldsymbol{\theta}' \in (\mathbb{R}^{m \times m})^L$ be the values of the parameters after a stochastic gradient descent step at $\boldsymbol{\theta}$ for the batch S , i.e.

$$\boldsymbol{\theta}' = \boldsymbol{\theta} - \eta \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \quad (52)$$

where $\eta > 0$ is the learning rate.

Assuming that $\boldsymbol{\theta}$ and $\boldsymbol{\theta}'$ satisfies the conditions of Theorem 12.3, we obtain

$$\begin{aligned} R(\mathbf{L}(h(\boldsymbol{\theta}')) \leq & R(\mathbf{L}(h(\boldsymbol{\theta}))) - \eta \langle \nabla_{\boldsymbol{\theta}} (R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}), \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \rangle \\ & + \eta \sqrt{n} \|\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,2} O \left(\frac{L^2 \omega^{1/3} \sqrt{m \log(m)}}{\sqrt{d}} \right) \left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \\ & + \eta^2 O \left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \frac{nL^2 m}{d} \right) \left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty}^2 \end{aligned} \quad (53)$$

where we refer to (40) for the form of $\nabla_{\boldsymbol{\theta}} (R \circ \mathbf{L} \circ h)(\boldsymbol{\theta})$ and to (39) for the form of $\nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}))$.

In addition, we make the assumption that for the set of values of $\boldsymbol{\theta}$ considered the hardness weighted sampling probabilities admit an upper-bound

$$\alpha = \min_{\boldsymbol{\theta}} \min_i \bar{p}_i(\mathbf{L}(\boldsymbol{\theta})) > 0 \quad (54)$$

Which is always satisfied under assumption 12.2 for Kullback-Leibler ϕ -divergence, and for any ϕ -divergence satisfying Definition 2.1 with a robustness parameter β small enough.

Let \mathbb{E}_S be the expectation with respect to S . Applying \mathbb{E}_S to (53), we obtain

$$\begin{aligned} & \mathbb{E}_S [R(\mathbf{L}(h(\boldsymbol{\theta}')))] \\ & \leq R(\mathbf{L}(h(\boldsymbol{\theta}))) - \eta \|\nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta})\|_{2,2}^2 \\ & \quad + \eta \|\nabla_h(R \circ \mathbf{L})(h(\boldsymbol{\theta}))\|_{2,2} O\left(\frac{nL^2\omega^{1/3}\sqrt{m \log(m)}}{\sqrt{d}}\right) \sqrt{\sum_{i=1}^n \max_l \|\bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_l}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta})\|^2} \\ & \quad + \eta^2 O\left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L})\right) \frac{nL^2m}{d}\right) \frac{1}{\alpha} \sum_{i=1}^n \max_l \|\bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}_l}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta})\|^2 \end{aligned} \quad (55)$$

where we have used the following results:

- For any integer $k \geq 1$, and all $(\mathbf{a}_i)_{i=1}^n \in (\mathbb{R}^k)^n$, we have (see the proof in 12.3.5)

$$\mathbb{E}_S \left[\frac{1}{b} \sum_{i \in S} \mathbf{a}_i \right] = \mathbb{E}_{\bar{p}(\hat{\mathbf{L}})} [\mathbf{a}_i] \quad (56)$$

- Using (56) for $(\mathbf{a}_i)_{i=1}^n = (\nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}))_{i=1}^n$, and the chain rule (40)

$$\mathbb{E}_S \left[\frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right] = \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) = \nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}) \quad (57)$$

- Using the triangular inequality

$$\left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \leq \frac{1}{b} \sum_{i \in S} \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \quad (58)$$

And using (56) for $(\mathbf{a}_i)_{i=1}^n = \left(\left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \right)_{i=1}^n$,

$$\begin{aligned} \mathbb{E}_S \left[\left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \right] & \leq \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty} \\ & \leq \sum_{i=1}^n \max_l \left\| \nabla_{\boldsymbol{\theta}_l}(\bar{p}_i(\hat{\mathbf{L}}) \mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2 \\ & \leq \sqrt{n} \sqrt{\sum_{i=1}^n \max_l \left\| \nabla_{\boldsymbol{\theta}_l}(\bar{p}_i(\hat{\mathbf{L}}) \mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2} \end{aligned} \quad (59)$$

where we have used Cauchy-Schwarz inequality for the last inequality.

- Using (58) and the convexity of the function $x \mapsto x^2$

$$\left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty}^2 \leq \frac{1}{b} \sum_{i \in S} \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty}^2 \quad (60)$$

And using (56) for $(a_i)_{i=1}^n = \left(\|\nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta})\|_{2,\infty}^2 \right)_{i=1}^n$,

$$\begin{aligned} \mathbb{E}_S \left[\left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty}^2 \right] &\leq \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_{2,\infty}^2 \\ &\leq \sum_{i=1}^n \frac{1}{\bar{p}_i(\hat{\mathbf{L}})} \max_l \left\| \nabla_{\boldsymbol{\theta}_l}(\bar{p}_i(\hat{\mathbf{L}}) \mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \quad (61) \\ &\leq \frac{1}{\alpha} \sum_{i=1}^n \max_l \left\| \nabla_{\boldsymbol{\theta}_l}(\bar{p}_i(\hat{\mathbf{L}}) \mathcal{L}_i \circ h_i)(\boldsymbol{\theta}) \right\|_2^2 \end{aligned}$$

Important Remark: It is worth noting the apparition of α (54) in (61). If we were using a uniform sampling as for ERM (i.e. for DRO in the limit $\beta \rightarrow 0$), we would have $\alpha = \frac{1}{n}$. So although our inequality (61) may seem brutal, it is consistent with equation (13.2) in [1] and the corresponding inequality in the case of ERM.

The rest of the proof of convergence will consist in proving that $\eta \|\nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta})\|_{2,2}^2$ dominates the two last terms in (53). As a result, we can already state that either the robustness parameter β , or the learning rate η will have to be small enough to control α .

Indeed, combining (53) with the chain rule (40), and the gradient bound Theorem 12.4 where we use our $\hat{\mathbf{L}}$ defined in (50)

$$\begin{aligned} \mathbb{E}_S [R(\mathbf{L}(h(\boldsymbol{\theta}')))] &\leq R(\mathbf{L}(h(\boldsymbol{\theta}))) - \Omega \left(\frac{\eta m \delta}{dn^2} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \\ &\quad + \eta O \left(\frac{nL^2 \omega^{1/3} \sqrt{m \log(m)}}{\sqrt{d}} \right) O \left(\sqrt{\frac{m}{d}} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \\ &\quad + \eta^2 O \left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \right) \frac{nL^2 m}{d} \right) O \left(\frac{m}{d\alpha} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \\ &\leq R(\mathbf{L}(h(\boldsymbol{\theta}))) - \Omega \left(\frac{\eta m \delta}{dn^2} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \\ &\quad + O \left(\frac{\eta n L^2 m \omega^{1/3} \sqrt{\log(m)}}{d} + K \frac{\eta^2 (n/\alpha) L^2 m^2}{d^2} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2 \quad (62) \end{aligned}$$

where we have used

$$K := \frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L}) \quad (63)$$

There are only two differences with equation (13.2) in [1]:

- in the last fraction we have n/α instead of n^2 (see remark 12.3.4 for more details), and an additional multiplicative term K . So in total, this term differs by a multiplicative factor $\frac{\alpha n}{K}$ from the analogous term in the proof of [1].
- we have $\sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta})) \right\|_2^2$ instead of $F(\mathbf{W}^{(t)})$. In fact they are analogous since in equation (13.2) in [1], $F(\mathbf{W}^{(t)})$ is the squared norm of the mean loss for the L^2 loss. We don't make such a strong assumption on the choice of \mathcal{L} (see assumption 12.2). It is worth noting that the same analogy is used in [1, Appendix A] where they extend their result to the mean loss with other objective function than the L^2 loss.

Our choice of learning rate in Theorem 12.2 can be rewritten as

$$\begin{aligned}
\eta_{exact} &= \Theta \left(\frac{\alpha n^2 \rho}{\beta C(\mathcal{L})^2 + 2n\rho C'(\nabla \mathcal{L})} \times \frac{b\delta d}{\text{poly}(n, L)m \log^2(m)} \right) \\
&= \Theta \left(\frac{\alpha n}{K} \times \frac{b\delta d}{\text{poly}(n, L)m \log^2(m)} \right) \\
&\leq \frac{\alpha n}{K} \times \eta'
\end{aligned} \tag{64}$$

And we also have

$$\eta_{exact} \leq \eta' \tag{65}$$

where η' is the learning rate chosen in the proof of Theorem 2 in [1]. We refer the reader to [1] for the details of the constant in " Θ " and the exact form of the polynomial $\text{poly}(n, L)$.

As a result, for $\eta = \eta_{exact}$, the term $\Omega \left(\frac{\eta m \delta}{dn^2} \right)$ dominates the other term of the right-hand side of inequality (62) as in the proof of Theorem 2 in [1].

This implies that the conditions of Theorem 12.4 are satisfied for all $\theta^{(t)}$, and that we have for all iteration $t > 0$

$$\mathbb{E}_{S_t} \left[R(\mathbf{L}(h(\theta^{(t+1)}))) \right] \leq R(\mathbf{L}(h(\theta^{(t)}))) - \Omega \left(\frac{\eta m \delta}{dn^2} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\theta^{(t)})) \right\|_2^2 \tag{66}$$

And using a result in Appendix A of [1], since under assumption 12.2 the distributionally robust loss is non-convex and bounded, we obtain for all $\epsilon' > 0$

$$\left\| \nabla_h (R \circ \mathbf{L})(h(\theta^{(T)})) \right\|_{2,2} \leq \epsilon' \quad \text{if } T = O \left(\frac{dn^2}{\eta \delta m \epsilon'^2} \right) \tag{67}$$

where according to (39)

$$\left\| \nabla_h (R \circ \mathbf{L})(h(\theta^{(T)})) \right\|_{2,2} = \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\theta^{(t)})) \right\|_2^2 \tag{68}$$

However, we are interested in a bound on $\left\| \nabla_{\theta} (R \circ \mathbf{L} \circ h)(\theta^{(T)}) \right\|_{2,2}$, rather than $\left\| \nabla_h (R \circ \mathbf{L})(h(\theta^{(T)})) \right\|_{2,2}$.

Using the gradient bound of Theorem 12.4 and the chain rules (40) and (39)

$$\left\| \nabla_{\theta} (R \circ \mathbf{L} \circ h)(\theta^{(T)}) \right\|_{2,2} \leq c_1 \sqrt{\frac{Lmn}{d}} \left\| \nabla_h (R \circ \mathbf{L})(h(\theta^{(T)})) \right\|_{2,2} \tag{69}$$

where $c_1 > 0$ is the constant hidden in $O \left(\sqrt{\frac{Lmn}{d}} \right)$.

So with $\epsilon' = \frac{1}{c_1} \sqrt{\frac{d}{Lmn}} \epsilon$, we finally obtain

$$\begin{aligned}
\left\| \nabla_{\theta} (R \circ \mathbf{L} \circ h)(\theta^{(T)}) \right\|_{2,2} &\leq c_1 \sqrt{\frac{Lmn}{d}} \left\| \nabla_h (R \circ \mathbf{L})(h(\theta^{(T)})) \right\|_{2,2} \\
&\leq c_1 \sqrt{\frac{Lmn}{d}} \epsilon' \\
&\leq \epsilon
\end{aligned} \tag{70}$$

If

$$T = O \left(\frac{dn^2}{\eta \delta m \epsilon'^2} \right) = O \left(\frac{dn^2}{\eta \delta m} \frac{Lmn}{d \epsilon^2} \right) = O \left(\frac{Ln^3}{\eta \delta \epsilon^2} \right) \tag{71}$$

which concludes the proof.

12.3.5 Proof of technical lemma 1

For any integer $k \geq 1$, and all $(\mathbf{a}_i)_{i=1}^n \in (\mathbb{R}^k)^n$, we have

$$\begin{aligned}
\mathbb{E}_S \left[\frac{1}{b} \sum_{i \in S} \mathbf{a}_i \right] &= \sum_{1 \leq i_1, \dots, i_b \leq n} \left[\left(\prod_{k=1}^n \bar{p}_{i_k}(\hat{\mathbf{L}}) \right) \frac{1}{b} \sum_{j=1}^b \mathbf{a}_{i_j} \right] \\
&= \frac{1}{b} \sum_{1 \leq i_1, \dots, i_b \leq n} \left[\sum_{j=1}^b \bar{p}_{i_j}(\hat{\mathbf{L}}) \mathbf{a}_{i_j} \left(\prod_{\substack{k=1 \\ k \neq j}}^n \bar{p}_{i_k}(\hat{\mathbf{L}}) \right) \right] \\
&= \frac{1}{b} \sum_{j=1}^b \left[\sum_{1 \leq i_1, \dots, i_b \leq n} \bar{p}_{i_j}(\hat{\mathbf{L}}) \mathbf{a}_{i_j} \left(\prod_{\substack{k=1 \\ k \neq j}}^n \bar{p}_{i_k}(\hat{\mathbf{L}}) \right) \right] \\
&= \frac{1}{b} \sum_{j=1}^b \left[\left(\sum_{i_j=1}^n \bar{p}_{i_j}(\hat{\mathbf{L}}) \mathbf{a}_{i_j} \right) \prod_{\substack{k=1 \\ k \neq j}}^n \left(\sum_{i_k=1}^n \bar{p}_{i_k}(\hat{\mathbf{L}}) \right) \right] \\
&= \frac{1}{b} \sum_{j=1}^b \left(\sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \mathbf{a}_i \right) \\
&= \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \mathbf{a}_i \\
&= \mathbb{E}_{\bar{p}(\hat{\mathbf{L}})} [\mathbf{a}_i]
\end{aligned} \tag{72}$$

12.4 Convergence of SGD with Hardness Weighted Sampling and stale per-example loss vector

The proof of the convergence of Algorithm 1 under the conditions of Theorem 12.2 follows the same structure as the proof of the convergence of Robust SGD with exact per-example loss vector 12.3.4. We will reuse the intermediate results of 12.3.4 when possible and focus on the differences between the two proofs due to the inexactness of the per-example loss vector.

Let t be the iteration number, and let $\boldsymbol{\theta}^{(t)} \in (\mathbb{R}^{m \times m})^L$ be the parameters of the deep neural network at iteration t . We define the stale per-example loss vector at iteration t as

$$\hat{\mathbf{L}} = \left(\mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t_i(t))})) \right)_{i=1}^n \tag{73}$$

where for all i , $t_i(t) < t$ corresponds to the latest iteration before t at which the per-example loss value for example i has been updated. Or equivalently, it corresponds to the last iteration before t when example i was drawn to be part of a mini-batch.

We also define the exact per-example loss vector that is unknown in Algorithm 1, as

$$\check{\mathbf{L}} = \left(\mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right)_{i=1}^n \tag{74}$$

Similarly to (52) we define

$$\boldsymbol{\theta}^{(t+1)} = \boldsymbol{\theta}^{(t)} - \eta \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \tag{75}$$

and using Theorem 12.3, similarly to (53), we obtain

$$\begin{aligned}
R(\mathbf{L}(h(\boldsymbol{\theta}^{t+1}))) &\leq R(\mathbf{L}(h(\boldsymbol{\theta}^{(t)}))) - \eta \langle \nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}^{(t)}), \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \rangle \\
&\quad + \eta \left\| \nabla_h(R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} O\left(\frac{L^2 \omega^{1/3} \sqrt{m \log(m)}}{\sqrt{d}}\right) \left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_{2,\infty} \\
&\quad + \eta^2 O\left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L})\right) \frac{nL^2 m}{d}\right) \left\| \frac{1}{b} \sum_{i \in S} \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_{2,\infty}^2
\end{aligned} \tag{76}$$

We can still define α as in (54)

$$\alpha = \min_{\boldsymbol{\theta}} \min_i \bar{p}_i(\mathbf{L}(\boldsymbol{\theta})) > 0 \tag{77}$$

where we are guaranteed that $\alpha > 0$ under assumptions 12.1.

Since Theorem 12.4 is independent to the choice of $\hat{\mathbf{L}}$, taking the expectation with respect to S , similarly to (62), we obtain

$$\begin{aligned}
\mathbb{E}_S \left[R(\mathbf{L}(h(\boldsymbol{\theta}^{t+1}))) \right] &\leq R(\mathbf{L}(h(\boldsymbol{\theta}^{(t)}))) - \eta \langle \nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}^{(t)}), \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \rangle \\
&\quad + \eta \left\| \nabla_h(R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} O\left(\frac{L^2 \omega^{1/3} \sqrt{nm \log(m)}}{\sqrt{d}}\right) \sqrt{\sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2^2} \\
&\quad + \eta^2 O\left(\left(\frac{\beta C(\mathcal{L})^2}{n\rho} + 2C(\nabla \mathcal{L})\right) \frac{nL^2 m}{d}\right) O\left(\frac{m}{d\alpha}\right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2^2
\end{aligned} \tag{78}$$

where the differences with respect to (62) comes from the fact that $\hat{\mathbf{L}}$ is not the exact per-example loss vector here, i.e. $\hat{\mathbf{L}} \neq \check{\mathbf{L}}$, which leads to

$$\begin{aligned}
\nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}^{(t)}) &= \sum_{i=1}^n \hat{p}_i(\check{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \\
&\neq \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)})
\end{aligned} \tag{79}$$

And

$$\begin{aligned}
\left\| \nabla_h(R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} &= \sum_{i=1}^n \left\| \hat{p}_i(\check{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2 \\
&\neq \sum_{i=1}^n \left\| \hat{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2
\end{aligned} \tag{80}$$

Let

$$K' = C(\mathcal{L}) A(\nabla \mathcal{L}) O\left(\frac{\beta L m^{3/2} \log^2(m)}{\alpha n^{1/2} \rho d^{3/2} b \log\left(\frac{1}{1-\alpha}\right)}\right) \tag{81}$$

Where $C(\mathcal{L}) > 0$ is a constant such that \mathcal{L} is $C(\mathcal{L})$ -Lipschitz continuous, and $A(\nabla \mathcal{L}) > 0$ is a constant that bound the gradient of \mathcal{L} with respect to its input.

$C(\mathcal{L})$ and $A(\nabla \mathcal{L})$ are guaranteed to exist under assumptions 12.1.

We can prove that, with probability at least $1 - \exp(-\Omega(\log^2(m)))$,

- according to lemma 12.4.1

$$\left\| \hat{p}(\hat{\mathbf{L}}) - \hat{p}(\check{\mathbf{L}}) \right\|_2 = \sqrt{\sum_{i=1}^n \left(\hat{p}_i(\hat{\mathbf{L}}) - \hat{p}_i(\check{\mathbf{L}}) \right)^2} \leq \eta \alpha K' \quad (82)$$

- according to lemma 12.4.2

$$\begin{aligned} & \left| \left\langle \nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}^{(t)}) - \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\rangle \right| \\ & \leq \eta \frac{m}{d} K' \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_2^2 \end{aligned} \quad (83)$$

- according to lemma 12.4.3

$$\left\| \nabla_h(R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} \leq (\sqrt{n} + \eta K') \sqrt{\sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_2^2} \quad (84)$$

Combining those three inequalities with (78) we obtain

$$\begin{aligned} \mathbb{E}_S \left[R(\mathbf{L}(h(\boldsymbol{\theta}^{(t+1)}))) \right] - R(\mathbf{L}(h(\boldsymbol{\theta}^{(t)}))) & \leq \\ & \eta \left[-\Omega \left(\frac{m\delta}{dn^2} \right) + O \left(\frac{nL^2m\omega^{1/3}\sqrt{\log(m)}}{d} \right) \right] \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2^2 \\ & \eta^2 O \left(K \frac{(n/\alpha)L^2m^2}{d^2} + \left(1 + \frac{m}{d} \right) K' \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2^2 \end{aligned} \quad (85)$$

One can see that compared to (62), there is only the additional term $(1 + \frac{m}{d}) K'$.

Using our choice of η ,

$$\eta = \eta_{stale} \leq O \left(\frac{\delta}{n^2 K'} \eta_{exact} \right) \quad (86)$$

where η_{exact} is the learning rate of Theorem 12.1, we have

$$\Omega \left(\frac{\eta m \delta}{dn^2} \right) \geq O \left(\eta^2 \left(1 + \frac{m}{d} \right) K' \right) \quad (87)$$

As a result, $\eta^2 \left(1 + \frac{m}{d} \right) K'$ is dominated by the term $\Omega \left(\frac{\eta m \delta}{dn^2} \right)$

In addition, since $\eta_{stale} \leq \eta_{exact}$, $\Omega \left(\frac{\eta m \delta}{dn^2} \right)$ still dominates also the other terms as in the proof of Theorem 12.1.

As a consequence, we obtain as in (66) that for any iteration $t > 0$

$$\mathbb{E}_{S_t} \left[R(\mathbf{L}(h(\boldsymbol{\theta}^{(t+1)}))) \right] \leq R(\mathbf{L}(h(\boldsymbol{\theta}^{(t)}))) - \Omega \left(\frac{\eta m \delta}{dn^2} \right) \sum_{i=1}^n \left\| \bar{p}_i(\hat{\mathbf{L}}) \nabla_{h_i} \mathcal{L}_i(h_i(\boldsymbol{\theta}^{(t)})) \right\|_2^2 \quad (88)$$

This concludes the proof using the same arguments as in the end of the proof of Theorem 12.1 starting from (66).

12.4.1 Proof of technical lemma 2

Using Lemma 3.2 and Lemma 3.1 we obtain

$$\begin{aligned} \left\| \hat{p}(\hat{\mathbf{L}}) - \hat{p}(\check{\mathbf{L}}) \right\|_2 & = \left\| \nabla_v R(\hat{\mathbf{L}}) - \nabla_v R(\check{\mathbf{L}}) \right\|_2 \\ & \leq \frac{\beta}{n\rho} \left\| \hat{\mathbf{L}} - \check{\mathbf{L}} \right\|_2 \end{aligned} \quad (89)$$

Using assumptions 12.2 and [1, Claim 11.2]

$$\begin{aligned}
\left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 &\leq \frac{\beta}{n\rho} \sqrt{\sum_{i=1}^n \left(\mathcal{L}_i \circ h_i(\boldsymbol{\theta}^{(t)}) - \mathcal{L}_i \circ h_i(\boldsymbol{\theta}^{(t_i(t))}) \right)^2} \\
&\leq \frac{\beta}{n\rho} C(\mathcal{L})C(h) \sqrt{\sum_{i=1}^n \left\| \boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t_i(t))} \right\|_{2,2}^2} \\
&\leq C(\mathcal{L})O\left(\frac{\beta L m^{1/2}}{n\rho d^{1/2}}\right) \sqrt{\sum_{i=1}^n \left\| \boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t_i(t))} \right\|_{2,2}^2}
\end{aligned} \tag{90}$$

Where $C(\mathcal{L})$ is the constant of Lipschitz continuity of the per-example loss \mathcal{L} (see assumptions 12.2) and $C(h)$ is the constant of Lipschitz continuity of the deep neural network h with respect to its parameters $\boldsymbol{\theta}$.

By developing the recurrence formula of $\boldsymbol{\theta}^{(t)}$ (75), we obtain

$$\begin{aligned}
\left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 &\leq C(\mathcal{L})O\left(\frac{\beta L m^{1/2}}{n\rho d^{1/2}}\right) \sqrt{\sum_{i=1}^n \left\| \boldsymbol{\theta}^{(t_i(t))} - \left(\sum_{\tau=t_i(t)}^{t-1} \frac{\eta}{b} \sum_{j \in S_\tau} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(\tau)}) \right) - \boldsymbol{\theta}^{(t_i(t))} \right\|_{2,2}^2} \\
&\leq \eta C(\mathcal{L})O\left(\frac{\beta L m^{1/2}}{n\rho d^{1/2}}\right) \sqrt{\sum_{i=1}^n \left\| \sum_{\tau=t_i(t)}^{t-1} \frac{1}{b} \sum_{j \in S_\tau} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(\tau)}) \right\|_{2,2}^2}
\end{aligned}$$

Let $A(\nabla \mathcal{L})$ a bound on the gradient of the per-example loss function. Using Theorem 12.4 and the chain rule

$$\forall j, \forall \tau \quad \left\| \nabla_{\boldsymbol{\theta}} (\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(\tau)}) \right\|_{2,2} \leq A(\nabla \mathcal{L})O\left(\frac{m}{d}\right) \tag{91}$$

And using the triangular inequality

$$\begin{aligned}
\left\| \sum_{\tau=t_i(t)}^{t-1} \frac{1}{b} \sum_{j \in S_\tau} \nabla_{\boldsymbol{\theta}} (\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(\tau)}) \right\|_{2,2} &\leq \sum_{\tau=t_i(t)}^{t-1} \frac{1}{b} \sum_{j \in S_\tau} \left\| \nabla_{\boldsymbol{\theta}} (\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(\tau)}) \right\|_{2,2} \\
&\leq \sum_{\tau=t_i(t)}^{t-1} A(\nabla \mathcal{L})O\left(\frac{m}{d}\right) \\
&\leq A(\nabla \mathcal{L})O\left(\frac{m}{d}\right) (t - t_i(t))
\end{aligned} \tag{92}$$

As a result, we obtain

$$\left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \leq \eta C(\mathcal{L})A(\nabla \mathcal{L})O\left(\frac{\beta L m^{3/2}}{n\rho d^{3/2}}\right) \sqrt{\sum_{i=1}^n (t - t_i(t))^2} \tag{93}$$

For all i and for any τ the probability that the sample i is not in batch S_τ is lesser than $(1 - \alpha)^b$.

Therefore, for any $k \geq 1$ and for any t ,

$$P(t - t_i(t) \geq k) \leq (1 - \alpha)^{kb} \tag{94}$$

For $k \geq \frac{1}{b}\Omega\left(\frac{\log^2(m)}{\log\left(\frac{1}{1-\alpha}\right)}\right)$, we have $(1 - \alpha)^{kb} \leq \exp(-\Omega(\log^2(m)))$, and thus with probability at least $1 - \exp(-\Omega(\log^2(m)))$,

$$\forall t, \quad t - t_i(t) \leq O\left(\frac{\log^2(m)}{b \log\left(\frac{1}{1-\alpha}\right)}\right) \tag{95}$$

As a result, we finally obtain that with probability at least $1 - \exp(-\Omega(\log^2(m)))$,

$$\begin{aligned} \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 &\leq \eta C(\mathcal{L}) A(\nabla \mathcal{L}) O\left(\frac{\beta L m^{3/2}}{n \rho d^{3/2}}\right) \sqrt{n} O\left(\frac{\log^2(m)}{b \log\left(\frac{1}{1-\alpha}\right)}\right) \\ &\leq \eta \alpha O\left(\frac{\beta L m^{3/2} \log^2(m)}{\alpha n^{1/2} \rho d^{3/2} b \log\left(\frac{1}{1-\alpha}\right)}\right) \\ &\leq \eta \alpha K' \end{aligned} \quad (96)$$

12.4.2 Proof of technical lemma 3

Let us first denote

$$\begin{aligned} A &= \left| \langle \nabla_{\boldsymbol{\theta}}(R \circ \mathbf{L} \circ h)(\boldsymbol{\theta}^{(t)}) - \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \rangle \right| \\ &= \left| \langle \sum_{i=1}^n (\bar{p}_i(\check{\mathbf{L}}) - \bar{p}_i(\hat{\mathbf{L}})) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{i=1}^n \bar{p}_i(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \rangle \right| \end{aligned} \quad (97)$$

Using Cauchy-Schwarz inequality

$$\begin{aligned} A &= \left| \sum_{i=1}^n (\bar{p}_i(\check{\mathbf{L}}) - \bar{p}_i(\hat{\mathbf{L}})) \langle \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \rangle \right| \\ &\leq \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \sqrt{\sum_{i=1}^n \left(\langle \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \rangle \right)^2} \end{aligned} \quad (98)$$

Let

$$B = \langle \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}), \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \rangle \quad (99)$$

Using again Cauchy-Schwarz inequality

$$B \leq \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2} \left\| \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2} \quad (100)$$

As a result, A becomes

$$\begin{aligned} A &\leq \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \left\| \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2} \sqrt{\sum_{i=1}^n \left\| \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2}^2} \\ &\leq \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \left\| \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2} \sqrt{\sum_{i=1}^n \frac{1}{\alpha^2} \left\| \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_i \circ h_i)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2}^2} \\ &\leq \frac{1}{\alpha} \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \left\| \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \nabla_{\boldsymbol{\theta}}(\mathcal{L}_j \circ h_j)(\boldsymbol{\theta}^{(t)}) \right\|_{2,2}^2 \end{aligned} \quad (101)$$

Using the triangular inequality, Theorem 12.4, and Lemma 12.4.1, we finally obtain

$$\begin{aligned}
A &\leq \frac{m}{\alpha d} \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \sum_{j=1}^n \left\| \bar{p}_j(\hat{\mathbf{L}}) \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2}^2 \\
&\leq \eta \frac{m}{d} K' \sum_{j=1}^n \left\| \bar{p}_j(\hat{\mathbf{L}}) \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2}^2
\end{aligned} \tag{102}$$

12.4.3 Proof of technical lemma 4

We have

$$\begin{aligned}
\left\| \nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} &= \sum_{j=1}^n \bar{p}_j(\check{\mathbf{L}}) \left\| \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2} \\
&= \sum_{j=1}^n \bar{p}_j(\hat{\mathbf{L}}) \left\| \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2} \\
&\quad + \sum_{j=1}^n \left(\frac{\bar{p}_j(\check{\mathbf{L}}) - \bar{p}_j(\hat{\mathbf{L}})}{\bar{p}_j(\hat{\mathbf{L}})} \right) \bar{p}_j(\hat{\mathbf{L}}) \left\| \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2}
\end{aligned} \tag{103}$$

Using Cauchy-Schwarz inequality

$$\left\| \nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} \leq \left(\sqrt{n} + \sqrt{\sum_{j=1}^n \left(\frac{\bar{p}_j(\check{\mathbf{L}}) - \bar{p}_j(\hat{\mathbf{L}})}{\bar{p}_j(\hat{\mathbf{L}})} \right)^2} \right) \sqrt{\sum_{j=1}^n \left\| \bar{p}_j(\hat{\mathbf{L}}) \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2}^2} \tag{104}$$

Using Lemma 12.4.1

$$\begin{aligned}
\sum_{j=1}^n \left(\frac{\bar{p}_j(\check{\mathbf{L}}) - \bar{p}_j(\hat{\mathbf{L}})}{\bar{p}_j(\hat{\mathbf{L}})} \right)^2 &\leq \frac{1}{\alpha} \left\| \hat{\mathbf{p}}(\hat{\mathbf{L}}) - \hat{\mathbf{p}}(\check{\mathbf{L}}) \right\|_2 \\
&\leq \eta K'
\end{aligned} \tag{105}$$

Therefore, we finally obtain

$$\left\| \nabla_h (R \circ \mathbf{L})(h(\boldsymbol{\theta}^{(t)})) \right\|_{1,2} \leq (\sqrt{n} + \eta K') \sqrt{\sum_{j=1}^n \left\| \bar{p}_j(\hat{\mathbf{L}}) \nabla_{h_j} \mathcal{L}(h_j(\boldsymbol{\theta}^{(t)})) \right\|_{2,2}^2} \tag{106}$$