

Supporting Information for:

Changing the Phosphorus Allotrope from a Square Columnar Structure to a Planar Zigzag Nanoribbon by Increasing the Diameter of Carbon Nanotube Nanoreactors

Jinying Zhang,^{*,†} Chengcheng Fu,[†] Shixin Song,[‡] Hongchu Du,[§] Dan Zhao,[†] Hongyang Huang,[†] Lihui Zhang,[†] Jie Guan,^{*,‡} Yifan Zhang,[⊥] Xinluo Zhao,[⊥] Chuansheng Ma,^{//} Chun-Lin Jia,^{*,§, //} David Tománek[#]

[†]State Key Laboratory of Electrical Insulation and Power Equipment, Center of Nanomaterials for Renewable Energy, School of Electrical Engineering, Xi'an Jiaotong University, Xi'an, Shaanxi, P. R. China, 710049

[‡]The School of Physics, Southeast University, Nanjing, Jiangsu, P. R. China, 211189

[§]Ernst Ruska Centre for Microscopy and Spectroscopy with Electrons, Forschungszentrum Jülich, 52425 Jülich, Germany

[⊥]Department of Physics, Shanghai University, Shanghai, P. R. China, 200444

// The School of Microelectronics and State Key Laboratory for Mechanical Behaviour of Materials, Xi'an Jiaotong University, Xi'an, Shaanxi, P. R. China, 710049

#Physics and Astronomy Department, Michigan State University, East Lansing, Michigan 48824-2320, USA




Figure S1. Structural models of (a) AR-P@SWCNT and (b) AR-P@SWCNT.




Figure S2. Raman spectra of pristine and phosphorus-filled (a) SWCNTs and (b) MWCNTs. Spectra of pristine nanotubes are shown in black and those of P-filled nanotubes in red.




Figure S3. (a, b) structural model and (c, d) vibrational phonon spectra of unstable straight (left) and stable twisted (right) SC-P structures.




Figure S4. Electronic band structure and density of states (DOS) for the twisted SC-P allotrope based on DFT-PBE calculations. An indirect band gap is indicated by the red arrow.