

Quantum computation of molecular response properties

Xiaoxia Cai,¹ Wei-Hai Fang,¹ Heng Fan,² and Zhendong Li^{1,*}

¹Key Laboratory of Theoretical and Computational Photochemistry, Ministry of Education, College of Chemistry, Beijing Normal University, Beijing 100875, China

²Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

Accurately predicting response properties of molecules such as the dynamic polarizability and hyperpolarizability using quantum mechanics has been a long-standing challenge with widespread applications in material and drug design. Classical simulation techniques in quantum chemistry are hampered by the exponential growth of the many-electron Hilbert space as the system size increases. Building upon the quantum linear system algorithm [Harrow et al., Phys. Rev. Lett. 103, 150502 (2009)], we describe an algorithm for computing molecular response properties on quantum computers, which scales polynomially in the system size instead of the dimension of the exponentially large Hilbert space, and hence achieves an exponential speedup over existing classical algorithms. Moreover, the variational hybrid quantum-classical variant of the proposed algorithm can be readily applied on near-term quantum devices.

Introduction.—How molecules response upon the action of external fields determines the properties of materials. For weak external fields, the response is fully characterized by the linear and nonlinear response functions[1, 2], such as the polarizability tensor $\alpha_{ij}(\omega)$ and hyperpolarizability $\beta_{ijk}(\omega_1, \omega_2)$ ($i, j, k \in \{x, y, z\}$). The dynamic polarizability $\alpha_{ij}(\omega)$ describes how the dipole moment of a molecule responses to an oscillating electric field to the leading order, and can be linked to the photoabsorption cross section, while the first-order hyperpolarizability describes nonlinear response processes such as second-harmonic generation in nonlinear optical materials. Besides, these response functions are also the key to understand intermolecular interactions. Notably, the van der Waals C_6 coefficients, which are of paramount importance in quantifying the dispersion interaction between drug molecules and proteins in drug design, can be computed from dynamic polarizabilities at imaginary frequencies via the Casimir-Polder integral[3].

Developing reliable quantum mechanical methods for accurately predicting molecular response properties has been one of the major challenges in quantum chemistry[2, 4, 5]. The full configuration interaction (FCI)[6–8], also known as the exact diagonalization method, represents the most accurate method within a molecular orbital basis set, however, is limited to small molecular systems due to the exponential growth of the many-electron Hilbert space as the system size increases. Over the past decades, a plethora of approximate methods along with efficient algorithms have been developed[4, 5]. Unfortunately, approximations adopted in these methods in order to describe the correlation among electrons efficiently, such as the mean-field approximation[6] or approximate exchange-correlation functionals in density functional theory[9], can sometimes fail miserably. In particular, the strong electron correlation[10], which is the root for many fascinating phenomena in materials such as high-temperature superconductivity, cannot be accurately accounted for by these approximate methods. A satisfactory classi-

cal simulation method for predicting molecular response properties, which works in all regime of electron correlation, is lacking.

Initially advocated by Feynman[11], quantum computation shows a great promise for solving interacting fermion problems in physics and chemistry[12–19]. The quantum phase estimation (QPE) algorithm[20] was applied to obtain the ground state energies of molecules with an exponential speedup over the classical FCI[21]. It also allows to compute molecular static properties via energy derivatives[22, 23]. While QPE has only been realized for two-electron systems[24–27] due to the requirement of long circuit depth, the variational quantum eigensolver (VQE)[28, 29] is more suitable for the noisy intermediate-scale quantum (NISQ)[30] devices. Unlike QPE, its advantage over classical simulation techniques in quantum chemistry is still an open question and being actively explored. Nevertheless, VQE has been experimentally demonstrated on various platforms for small molecules such as H_2 , HeH^+ , LiH and BeH_2 [27, 28, 31–33]. In view of such progresses on the ground state problem, it is a natural question to ask whether computing molecular response properties, as the next logical step after computing the ground state, will also benefit from quantum computation.

In this Letter, we propose an algorithm for computing molecular response properties on quantum computers. While dynamical properties can alternatively be obtained by Fourier transform of the corresponding correlation functions in the time domain[16, 34, 35] determined from real-time Hamiltonian simulations, analogous to the classical computation side[2, 4, 5] it is highly desirable to have a quantum algorithm for computing a target response property such as $\alpha_{ij}(\omega)$ or $\beta_{ijk}(\omega_1, \omega_2)$ at given frequencies directly. Because in many molecular applications[2, 4], only a small range of frequencies is of interest, including the simulations of (hyper)polarizabilities at specific frequencies of applied electromagnetic fields[8, 36], absorption spectra in an interested visible/ultraviolet/X-ray region[5], and multi-dimensional spectroscopies for studying couplings

between selected modes[37]. By reformulating the target property into a symmetric expression with the help of a set of auxiliary quantum states, we convert the most demanding part of computations into linear systems of equations for determining these states, which can be solved on quantum computers using quantum algorithms for linear systems of equations[38–42] or variational hybrid quantum-classical algorithms[43, 44]. Depending on the subroutine employed for determining auxiliary states, the resulting variant of the proposed algorithm can be considered as the analog of QPE or VQE for molecular response properties. Furthermore, we show that in combination with the quantum linear system algorithm invented by Harrow, Hassidim, and Lloyd (HHL)[38], the runtime complexity of the proposed algorithm scales polynomially in the molecular system size, instead of the dimension of the exponentially large many-electron Hilbert space. Thus, an exponential speedup is achieved compared with the classical FCI-based approach[7, 8] for computing molecular response properties, which has a direct impact on designing new materials, drugs and catalysts.

Theory.—For concreteness, we consider the calculation of the polarizability $\alpha_{zz}(\omega)$ for a molecule under a monochromatic electric field with optical frequency ω in the z -direction. The static polarizability will be obtained as a special case where $\omega = 0$. Extensions to off-diagonal components of the polarizability tensor as well as nonlinear response properties are straightforward and will be discussed later.

Suppose initially without external fields, a molecule with N electrons is in the ground state $|\Psi_0\rangle$ of the second quantized electronic Hamiltonian \hat{H}_0 , expressed in an orthonormal molecular spin-orbital basis $\{\psi_p\}_{p=1}^K$ (K is proportional to the system size N) as

$$\hat{H}_0 = \sum_{p,q=1}^K h_{pq} a_p^\dagger a_q + \frac{1}{2} \sum_{p,q,r,s=1}^K h_{pqrs} a_p^\dagger a_q^\dagger a_s a_r, \quad (1)$$

where $a_p^{(\dagger)}$ represents the fermionic annihilation (creation) operator, and h_{pq} (h_{pqrs}) represent the one-electron (two-electron) integrals. The dynamic electric field in the dipole approximation is associated with the perturbation operator

$$\hat{z} = \sum_{p,q=1}^K z_{pq} a_p^\dagger a_q, \quad (2)$$

where $z_{pq} \triangleq \langle \psi_p | z | \psi_q \rangle$ represent the dipole-moment integrals. By the time-dependent perturbation theory, the frequency-dependent polarizability $\alpha_{zz}(\omega)$ can be expressed in a sum-over-state (SOS) form[1, 2]

$$\alpha_{zz}(\omega) = \sum_{n>0} \left[\frac{\langle \Psi_0 | \hat{z} | \Psi_n \rangle \langle \Psi_n | \hat{z} | \Psi_0 \rangle}{\omega_{n0} - (\omega + i\gamma)} + \frac{\langle \Psi_0 | \hat{z} | \Psi_n \rangle \langle \Psi_n | \hat{z} | \Psi_0 \rangle}{\omega_{n0} + (\omega + i\gamma)} \right], \quad (3)$$

with $\gamma > 0$ being a phenomenological damping parameter, which physically is associated with the inverse lifetime of excited states. Computing $\alpha_{zz}(\omega)$ allows to access important information of molecules such as the transition dipole moments $\langle \Psi_0 | \hat{z} | \Psi_n \rangle$ between the ground state $|\Psi_0\rangle$ and the n -th excited state $|\Psi_n\rangle$, as well as the associated excitation energy $\omega_{n0} \triangleq E_n - E_0$. Moreover, the imaginary part of $\alpha_{zz}(\omega)$ is related with the photoabsorption cross section $\sigma(\omega) \propto \omega \Im \alpha(\omega)$ for visible/ultraviolet/X-ray absorption spectra, which is one of the central quantities considered in designing functional materials.

In the standard FCI-based approach[7, 8] for computing $\alpha_{zz}(\omega)$, Eq. (3) is reformulated as

$$\alpha_{zz}(\omega) = \langle \Psi_0 | \hat{z} | \Psi(\omega) \rangle + \langle \Psi_0 | \hat{z} | \Psi(-\omega) \rangle, \quad (4)$$

where the frequency-dependent response wavefunctions $|\Psi(\pm\omega)\rangle$ are obtained by solving the response equations

$$\hat{Q}[\hat{H}_0 - E_0 \mp (\omega + i\gamma)]\hat{Q}|\Psi(\pm\omega)\rangle = \hat{Q}\hat{z}|\Psi_0\rangle, \quad (5)$$

with the projector $\hat{Q} = 1 - |\Psi_0\rangle\langle\Psi_0|$, in the full N -electron Hilbert space, and hence avoids the need for determining all excited states in the SOS form (3). The computational complexity of solving Eq. (5) using the best classical iterative algorithm[45] for linear systems of equations scales linearly in the dimension of the N -electron Hilbert space D . For the molecular problem with \hat{H}_0 (1), D is exponential in N , e.g., $D = \binom{K}{N}$ with $K = 2N$ for the half-filling case. Therefore, like solving the ground-state eigenvalue problem, viz., $\hat{H}_0|\Psi_0\rangle = E_0|\Psi_0\rangle$, this FCI-based approach scales exponentially in the system size N , and in practical is limited to very small molecules (ca. $N \lesssim 16$ assuming $K = 2N$) in routine quantum chemistry applications[4, 5, 8].

Just as QPE and VQE have been applied to the ground state problem, we attempt to utilize the advantage of quantum algorithms for linear systems of equations[38–44] in computing molecular response properties. However, while QPE can be applied readily to the ground state problem, both Eqs. (4) and (5) are not in a form that is amenable to compute on quantum computers directly, due to the asymmetric form of each term in Eq. (4) and the involvement of the projector \hat{Q} . To resolve these two problems, we introduce the notation

$$\hat{A}(\pm\omega) \triangleq \hat{H}_0 - E_0 \mp (\omega + i\gamma) \quad (6)$$

for brevity and rewrite the first part of $\alpha_{zz}(\omega)$ (4) as

$$\begin{aligned} \langle \Psi_0 | \hat{z} | \Psi(\omega) \rangle &= \langle \Psi_0 | \hat{z} \hat{Q} [\hat{Q} \hat{A}(\omega) \hat{Q}]^{-1} \hat{Q} \hat{z} | \Psi_0 \rangle \\ &= \langle \Psi_0 | \hat{z} \hat{Q} \hat{A}^{-1}(\omega) \hat{z} | \Psi_0 \rangle \\ &= \langle \Psi_0 | \hat{z} [\hat{A}^\dagger(\omega)]^{-1} \hat{A}^\dagger(\omega) \hat{Q} \hat{A}^{-1}(\omega) \hat{z} | \Psi_0 \rangle, \end{aligned} \quad (7)$$

where the second equality follows from the spectral decompositions $\hat{A}(\omega) = \sum_{n>0} |\Psi_n\rangle[\omega_{n0} - (\omega + i\gamma)]\langle\Psi_n|$ and $\hat{Q} = \sum_{n>0} |\Psi_n\rangle\langle\Psi_n|$, which immediately imply that $\hat{A}(\omega)$ is invertible for $\gamma > 0$ regardless of ω , and

$\hat{Q}[\hat{Q}\hat{A}(\omega)\hat{Q}]^{-1}\hat{Q} = \sum_{n>0} |\Psi_n\rangle[\omega_{n0} - (\omega + i\gamma)]^{-1}\langle\Psi_n| = \hat{Q}\hat{A}^{-1}(\omega)\hat{Q} = \hat{Q}\hat{A}^{-1}(\omega)$. To reach a symmetric expression, the identity $[\hat{A}^\dagger(\omega)]^{-1}\hat{A}^\dagger(\omega) = 1$ has been inserted in the last line of Eq. (7), which suggests to introduce an auxiliary state $|Z(\omega)\rangle$ satisfying an equation similar to Eq. (5) but without the projector \hat{Q}

$$\hat{A}(\omega)|Z(\omega)\rangle = \hat{z}|\Psi_0\rangle. \quad (8)$$

Consequently, Eq. (7) can be recast into a symmetric form

$$\begin{aligned} \langle\Psi_0|\hat{z}|\Psi(\omega)\rangle &= \langle Z(\omega)|\hat{A}^\dagger(\omega)|Z(\omega)\rangle \\ &+ (\omega - i\gamma)|\langle Z(\omega)|\Psi_0\rangle|^2. \end{aligned} \quad (9)$$

Now the explicit dependence on the projector \hat{Q} , which makes the design of a quantum algorithm difficult, has been removed from both the response equation (8) and the expression for the polarizability (9). Its effect is only reflected in the second term of Eq. (9). Likewise, the second part of $\alpha_{zz}(\omega)$ (4) can be expressed in a similar symmetric form

$$\begin{aligned} \langle\Psi_0|\hat{z}|\Psi(-\omega)\rangle &= \langle Z(-\omega)|\hat{A}^\dagger(-\omega)|Z(-\omega)\rangle \\ &- (\omega - i\gamma)|\langle Z(-\omega)|\Psi_0\rangle|^2. \end{aligned} \quad (10)$$

More explicitly, we can separate $\alpha_{zz}(\omega)$ into real and imaginary parts

$$\begin{aligned} \alpha_{zz}(\omega) &= \Re\alpha_{zz}(\omega) + i\Im\alpha_{zz}(\omega), \\ \Re\alpha_{zz}(\omega) &= \langle Z(\omega)|\hat{H}_0 - E_0|Z(\omega)\rangle, \\ &+ \langle Z(-\omega)|\hat{H}_0 - E_0|Z(-\omega)\rangle - \frac{\omega}{\gamma}\Im\alpha_{zz}(\omega), \\ \Im\alpha_{zz}(\omega) &= \gamma(\langle Z(\omega)|Z(\omega)\rangle - \langle Z(-\omega)|Z(-\omega)\rangle) \\ &- |\langle Z(\omega)|\Psi_0\rangle|^2 + |\langle Z(-\omega)|\Psi_0\rangle|^2, \end{aligned} \quad (11)$$

where the expected symmetry relations $\Re\alpha_{zz}(-\omega) = \Re\alpha_{zz}(\omega)$ and $\Im\alpha_{zz}(-\omega) = -\Im\alpha_{zz}(\omega)$ are obvious. In fact, from Eq. (8) one can further find $\langle\Psi_0|Z(\pm\omega)\rangle = \mp \frac{\langle\Psi_0|\hat{z}|\Psi_0\rangle}{\omega + i\eta}$, such that the second terms in Eqs. (9) and (10) will cancel each other in $\alpha_{zz}(\omega)$. Building upon the reformulation of the standard response theory, Eqs. (8)-(11), we are ready to present a quantum algorithm for computing $\alpha_{zz}(\omega)$.

Algorithm. — Our algorithm assumes that the ground state problem has been solved, namely, the ground-state wavefunction $|\Psi_0\rangle$ and its associated energy E_0 are available either by QPE or VQE. The central step is to solve the response equation (8), when expressed in the full many-electron Hilbert space as a linear system of equation with dimension D , on quantum computers. One choice suitable for NISQ devices is to use the variational hybrid quantum-classical algorithms[43, 44] for solving (8). In this case, an appropriate variational wavefunction ansatz for the normalized state $|x\rangle = |Z(\omega)\rangle/\sqrt{\langle Z(\omega)|Z(\omega)\rangle}$ can be designed and optimized by minimizing the following cost function

$$\mathcal{C}[|x\rangle] = 1 - \frac{|\langle\Psi_0|\hat{z}\hat{A}(\omega)|x\rangle|^2}{\langle\Psi_0|\hat{z}\hat{z}|\Psi_0\rangle\langle x|\hat{A}^\dagger(\omega)\hat{A}(\omega)|x\rangle}. \quad (12)$$

The solution of Eq. (8) corresponding to $\mathcal{C}[|x\rangle] = 0$ can be readily used to compute $\alpha_{zz}(\omega)$ (11) by efficient measurements (vide post). Together with VQE for the ground state $|\Psi_0\rangle$, this approach provides a feasible variational hybrid quantum-classical algorithm for computing molecular response properties on near-term devices. The other choice for solving (8) is to use the quantum algorithms for linear systems of equations[38–42], which are in principle exact and have the potential for quantum speedup. It is also noteworthy that the HHL algorithm[38] has recently been experimentally realized on small scale problems[46–49]. In the following context, we will focus on the demonstration of quantum advantage for computing molecular response properties using the HHL algorithm as a subroutine.

Since $\hat{A}(\omega)$ (6) is non-Hermitian for $\gamma > 0$, $|Z(\omega)\rangle$ can be determined using the HHL algorithm[38] either by

$$\begin{bmatrix} 0 & \hat{A}(\omega) \\ \hat{A}^\dagger(\omega) & 0 \end{bmatrix} \begin{bmatrix} 0 \\ |Z(\omega)\rangle \end{bmatrix} = \begin{bmatrix} \hat{z}|\Psi_0\rangle \\ 0 \end{bmatrix}, \quad (13)$$

as suggested in the original work[38] or by the following equivalent equation

$$\hat{A}^\dagger(\omega)\hat{A}(\omega)|Z(\omega)\rangle = \hat{A}^\dagger(\omega)\hat{z}|\Psi_0\rangle, \quad (14)$$

with a Hermitian matrix on the left hand side (LHS) of the same dimension as Eq. (8) at the cost of increasing the condition number. For a linear system of equations $\mathbf{A}\mathbf{x} = \mathbf{b}$, where \mathbf{A} is a Hermitian matrix of dimension D with an eigendecomposition $\mathbf{A} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^\dagger$, the HHL algorithm[38] essentially prepares a solution following the sequence $\mathbf{x} = \mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\dagger\mathbf{b}$. The transformation to the eigenbasis of \mathbf{A} and the backtransformation are executed by QPE subroutines[20], which require the implementation of the controlled time evolution $e^{i\mathbf{A}t}$, while the realization of the nonunitary operation $\mathbf{\Lambda}^{-1}$ is through controlled rotations also with the help of ancilla qubits. The runtime complexity of the HHL algorithm is $O(\log(D)s^2\kappa^2/\epsilon)$ [38], where ϵ is the desired precision, s is the sparsity parameter specifying the maximal number of nonzero entries per row in \mathbf{A} , and κ is the condition number of \mathbf{A} , i.e., $\kappa = |\lambda_{\max}|/|\lambda_{\min}|$, which is the ratio between the maximal and minimal eigenvalues by moduli of \mathbf{A} . The real advantage of the HHL algorithm over classical algorithms crucially depends on the efficiency of four major steps[38, 50]: (1) preparation of \mathbf{b} on quantum computers, (2) Hamiltonian simulation $e^{i\mathbf{A}t}$, (3) dependence of κ on D , and (4) readout of the output quantum state $|x\rangle = A^{-1}|b\rangle/\|A^{-1}|b\rangle\|$ encoding the solution \mathbf{x} . Any slowdown in one of the steps could kill the exponential speedup promised by the HHL algorithm. Here, we show how an exponential speedup can be achieved for computing molecular response properties step-by-step by utilizing the specialities of the problem.

Step 1: Provided $|\Psi_0\rangle$ is available, the state $\hat{z}|\Psi_0\rangle$ on the right hand side (RHS) of Eq. (13) or $\hat{A}^\dagger(\omega)\hat{z}|\Psi_0\rangle$ in Eq. (14) can be prepared with a cost of poly(N) using

the linear combination of unitaries (LCU) algorithm[51, 52]. This is because both the one-body perturbation \hat{z} (2) and the Hamiltonian \hat{H}_0 (1) in $\hat{A}(\omega)$ can be expressed as a sum of poly(N) Pauli matrices, e.g.,

$$\hat{z} = \sum_{\mu} z_{\mu} P_{\mu}, \quad P_{\mu} = \sigma_{\mu_1} \otimes \sigma_{\mu_2} \otimes \cdots \otimes \sigma_{\mu_K}, \quad (15)$$

where $\sigma_{\mu_k} \in \{I_2, \sigma_x, \sigma_y, \sigma_z\}$ and the number of terms is quadratic in N for \hat{z} (2), through a fermion-to-qubit mapping such as the Jordan-Wigner transformation[53] or the Bravyi-Kitaev transformation[54, 55].

Step 2: Given the RHS of Eq. (13) (or Eq. (14)) prepared on quantum computers, the HHL algorithm is applied to prepare a normalized solution state $|x\rangle = |Z(\omega)\rangle / \sqrt{\langle Z(\omega)|Z(\omega)\rangle}$ for Eq. (8). For molecular systems with \hat{H}_0 (1), it is known that the Hamiltonian simulation can be accomplished efficiently in poly(N), just as in applying QPE to the molecular ground state problem[13, 14, 21]. Because \hat{H}_0 involves at most two-body Coulomb interactions, the sparsity parameter s is only quartic in N . Thus, the most crucial part for the runtime complexity of the HHL algorithm is the condition number κ . On one hand, since \hat{H}_0 can be written as a sum over $O(N^4)$ Pauli terms $\hat{H}_0 = \sum_{\mu} h_{\mu} P_{\mu}$, as for \hat{z} in Eq. (15), $|\lambda_{\max}|$ of \hat{H}_0 (and $\hat{A}(\omega)$) is bounded by a system-dependent constant $\max_{\mu} |h_{\mu}|$ times N^4 . Assuming we consider the scaling with respect to the variation of the system size for systems of the same kind, such as water clusters of different sizes in a given atomic orbital basis set, then $\max_{\mu} |h_{\mu}|$ is independent of N , such that $|\lambda_{\max}|$ is of poly(N). On the other hand, the operator $\hat{H}_0 - E_0 - \omega$ becomes singular whenever the frequency ω matches the excitation energy ω_{n0} , such that in the worst case $|\lambda_{\min}|$ of $\hat{A}^{\dagger}(\omega)\hat{A}(\omega)$ in Eq. (14) is γ^2 , and likewise for Eq. (13) $|\lambda_{\min}|$ equals γ . In practice, the parameter γ is a fixed input parameter for spectral resolution, which determines the half width at half maximum (HWHM) of peaks in $\Im\alpha(\omega)$ (11). Thus, the condition numbers κ for the coefficient matrices in Eqs. (13) and (14) are polynomial in the system size N instead of the dimension of the Hilbert space D , which is exponential in N . This concludes that the runtime complexity of the HHL algorithm for preparing the normalized solution state $|x\rangle$ from either Eq. (13) or (14) is poly(N).

Step 3: After applying the HHL algorithm to Eq. (13) or (14), the first part of $\alpha_{zz}(\omega)$ (4) can be computed from $|x\rangle$ using Eq. (9) in poly(N), without accessing its individual entry. This is achieved by first noting that the norm of $|Z(\omega)\rangle$ required in Eq. (9) can be computed using Eq. (8) as

$$\langle Z(\omega)|Z(\omega)\rangle = \langle \Psi_0|\hat{z}\hat{z}|\Psi_0\rangle / \langle x|\hat{A}^{\dagger}(\omega)\hat{A}(\omega)|x\rangle, \quad (16)$$

which only requires the measurements of $\langle \Psi_0|\hat{z}\hat{z}|\Psi_0\rangle$ and $\langle x|\hat{A}^{\dagger}(\omega)\hat{A}(\omega)|x\rangle$. Then, the first term in Eq. (9) involving $\langle Z(\omega)|\hat{H}_0|Z(\omega)\rangle$ can be obtained from the measurement $\langle x|\hat{H}_0|x\rangle = \sum_{\mu} h_{\mu} \langle x|P_{\mu}|x\rangle$ in the same way as

obtaining the energy in VQE[28, 29], while the second term $|\langle Z(\omega)|\Psi_0\rangle|^2$ can be computed from $|\langle x|\Psi_0\rangle|^2$ by the SWAP test[56, 57] or simply from $\frac{|\langle \Psi_0|\hat{z}|\Psi_0\rangle|^2}{\omega^2 + \gamma^2}$ following Eq. (8). Therefore, the necessary information for computing $\alpha_{zz}(\omega)$ from the output state $|x\rangle$ of the HHL algorithm (and its counterpart for $|Z(-\omega)\rangle$) can be accessed through $\langle x|\hat{H}_0^2|x\rangle$, $\langle x|\hat{H}_0|x\rangle$, and $|\langle x|\Psi_0\rangle|^2$ with a cost of poly(N).

Using the above procedure, we show that the dynamic polarizability tensor $\alpha_{zz}(\omega)$ of molecules can be computed on quantum computers with poly(N) runtime complexity, achieving an exponential speedup compared with the classical FCI-based approach[7, 8]. This becomes possible due to the specialities of the molecular response problem, such that all the limitations of the HHL algorithm can be overcome in this application: the RHS of Eq. (8) can always be efficiently prepared given $|\Psi_0\rangle$, $e^{i\hat{H}_0 t}$ can be efficiently simulated due to the sparse structure of \hat{H}_0 (1), the condition numbers κ for matrices in Eqs. (13) and (14) are polynomial in N , and finally only partial information of the solution is required for computing $\alpha_{zz}(\omega)$. Therefore, the molecular response problem is an ideal application of the HHL algorithm[38], and the same conclusion is also generalizable to its improved variants[39–41].

Discussion.— The above quantum algorithm can be readily generalized to compute different linear and nonlinear response properties, using the same idea of first deriving a symmetric expression for the target property by introducing appropriate auxiliary states, and then determining these states by solving response equations with appropriate quantum or hybrid algorithms. For off-diagonal components of $\alpha_{ij}(\omega)$, e.g., $\alpha_{xz}(\omega)$ containing a form of $\langle \Psi_0|\hat{x}\hat{Q}[\hat{Q}\hat{A}(\omega)\hat{Q}]^{-1}\hat{Q}\hat{z}|\Psi_0\rangle$, a symmetric expression can be derived by applying the polarization identity, which involves a linear combination of four symmetric terms $\langle \Psi_0|(\hat{x} \pm (i)\hat{z})^{\dagger}\hat{Q}[\hat{Q}\hat{A}(\omega)\hat{Q}]^{-1}\hat{Q}(\hat{x} \pm (i)\hat{z})|\Psi_0\rangle$ and each of them can be computed using the same algorithm for $\alpha_{zz}(\omega)$. An increasingly important nonlinear response functions for probing elementary excitations in complex correlated electron systems is the resonant inelastic X-ray scattering (RIXS) amplitudes[5, 58], given by the Kramers-Heisenberg formula[59] as

$$\mathcal{F}_{zz}^{f0}(\omega) = \sum_n \left[\frac{\langle \Psi_f|\hat{z}|\Psi_n\rangle \langle \Psi_n|\hat{z}|\Psi_0\rangle}{\omega_{n0} - (\omega + i\gamma)} + \frac{\langle \Psi_f|\hat{z}|\Psi_n\rangle \langle \Psi_n|\hat{z}|\Psi_0\rangle}{\omega_{n0} + (\omega' + i\gamma)} \right], \quad (17)$$

where $\omega' \triangleq \omega - \omega_{f0}$ and $|\Psi_f\rangle$ represents the final state of interest involved in the inelastic scattering process. Eq. (17) takes a similar form as Eq. (3) for $\alpha_{zz}(\omega)$, and hence a similar strategy can be designed to compute $|\mathcal{F}_{zz}^{f0}(\omega)|^2$ for the scattering cross section[58]. In particular, within the rotating wave approximation, where the second term of Eq. (17) is neglected, the scattering cross section is simply given by $|\mathcal{F}_{zz}^{f0}(\omega)|^2 = |\langle \Psi_f|\hat{z}|Z(\omega)\rangle|^2$

with the same $|Z(\omega)\rangle$ in Eq. (8), which can be computed by modifications of the SWAP test[56]

In summary, we presented a general algorithm for computing molecular response properties on quantum computers. The most demanding step involves a set of linear systems of equations for auxiliary quantum states, which can be solved either by quantum algorithms[38–42] or variational hybrid quantum-classical algorithms[43, 44]. The resulting two variants enable the computation of molecular response properties for interested frequencies directly. While the later variational hybrid variant is more suitable for near-term applications, it is demonstrated that the former with the HHL algorithm[38] as a subroutine has a provable quantum speedup over existing classical FCI-based approach[7, 8]. Our work provides a new evidence that quantum chemistry is among the first fields that will benefit from quantum computations. Enabling accurate and efficient predictions of molecular response properties on quantum computers will potentially open up many new applications of quantum computations in material science and drug discovery in the near future.

Acknowledgements The author (Z.L.) acknowledges Garnet Kin-Lic Chan, Jiajun Ren, Man-Hong Yung, and Zheng Li for critically reading the manuscript and helpful comments. We thank Zi-Yong Ge, Zhengan Wang, and Kai Xu for discussions. This work was supported by the National Natural Science Foundation of China (Grants No. 21973003) and the Beijing Normal University Startup Package.

* zhendongli@bnu.edu.cn

- [1] R. W. Boyd, *Nonlinear optics* (Elsevier, 2003).
- [2] P. Norman, K. Ruud, and T. Saue, *Principles and practices of molecular properties: Theory, modeling, and simulations* (John Wiley & Sons, 2018).
- [3] H. B. Casimir and D. Polder, *Physical Review* **73**, 360 (1948).
- [4] T. Helgaker, S. Coriani, P. Jørgensen, K. Kristensen, J. Olsen, and K. Ruud, *Chemical Reviews* **112**, 543 (2012).
- [5] P. Norman and A. Dreuw, *Chemical Reviews* **118**, 7208 (2018).
- [6] A. Szabo and N. S. Ostlund, *Modern quantum chemistry: introduction to advanced electronic structure theory* (Courier Corporation, 2012).
- [7] J. Olsen and P. Jørgensen, *The Journal of Chemical Physics* **82**, 3235 (1985).
- [8] H. Koch and R. J. Harrison, *The Journal of Chemical Physics* **95**, 7479 (1991).
- [9] A. J. Cohen, P. Mori-Sánchez, and W. Yang, *Science* **321**, 792 (2008).
- [10] P. R. Kent and G. Kotliar, *Science* **361**, 348 (2018).
- [11] R. P. Feynman, *International Journal of Theoretical Physics* **21**, 467 (1982).
- [12] S. Lloyd, *Science* **273**, 1073 (1996).
- [13] D. S. Abrams and S. Lloyd, *Physical Review Letters* **79**, 2586 (1997).
- [14] D. S. Abrams and S. Lloyd, *Physical Review Letters* **83**, 5162 (1999).
- [15] G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, *Physical Review A* **64**, 022319 (2001).
- [16] R. Somma, G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, *Physical Review A* **65**, 042323 (2002).
- [17] I. M. Georgescu, S. Ashhab, and F. Nori, *Reviews of Modern Physics* **86**, 153 (2014).
- [18] Y. Cao, J. Romero, J. P. Olson, M. Degroote, P. D. Johnson, M. Kieferová, I. D. Kivlichan, T. Menke, B. Peropadre, N. P. Sawaya, *et al.*, *Chemical Reviews* **119**, 10856 (2019).
- [19] S. McArdle, S. Endo, A. Aspuru-Guzik, S. C. Benjamin, and X. Yuan, *Reviews of Modern Physics* **92**, 015003 (2020).
- [20] A. Y. Kitaev, arXiv preprint quant-ph/9511026 (1995).
- [21] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, *Science* **309**, 1704 (2005).
- [22] I. Kassal and A. Aspuru-Guzik, *The Journal of Chemical Physics* **131**, 224102 (2009).
- [23] T. E. O’Brien, B. Senjean, R. Sagastizabal, X. Bonet-Monroig, A. Dutkiewicz, F. Buda, L. DiCarlo, and L. Visscher, *npj Quantum Information* **5**, 1 (2019).
- [24] B. P. Lanyon, J. D. Whitfield, G. G. Gillett, M. E. Goggin, M. P. Almeida, I. Kassal, J. D. Biamonte, M. Mohseni, B. J. Powell, M. Barbieri, *et al.*, *Nature Chemistry* **2**, 106 (2010).
- [25] J. Du, N. Xu, X. Peng, P. Wang, S. Wu, and D. Lu, *Physical Review Letters* **104**, 030502 (2010).
- [26] Y. Wang, F. Dolde, J. Biamonte, R. Babbush, V. Bergholm, S. Yang, I. Jakobi, P. Neumann, A. Aspuru-Guzik, J. D. Whitfield, *et al.*, *ACS Nano* **9**, 7769 (2015).
- [27] P. J. O’Malley, R. Babbush, I. D. Kivlichan, J. Romero, J. R. McClean, R. Barends, J. Kelly, P. Roushan, A. Tranter, N. Ding, *et al.*, *Physical Review X* **6**, 031007 (2016).
- [28] A. Peruzzo, J. McClean, P. Shadbolt, M.-H. Yung, X.-Q. Zhou, P. J. Love, A. Aspuru-Guzik, and J. L. O’Brien, *Nature Communications* **5**, 4213 (2014).
- [29] J. R. McClean, J. Romero, R. Babbush, and A. Aspuru-Guzik, *New Journal of Physics* **18**, 023023 (2016).
- [30] J. Preskill, *Quantum* **2**, 79 (2018).
- [31] Y. Shen, X. Zhang, S. Zhang, J.-N. Zhang, M.-H. Yung, and K. Kim, *Physical Review A* **95**, 020501 (2017).
- [32] A. Kandala, A. Mezzacapo, K. Temme, M. Takita, M. Brink, J. M. Chow, and J. M. Gambetta, *Nature* **549**, 242 (2017).
- [33] C. Hempel, C. Maier, J. Romero, J. McClean, T. Monz, H. Shen, P. Jurcevic, B. P. Lanyon, P. Love, R. Babbush, *et al.*, *Physical Review X* **8**, 031022 (2018).
- [34] A. Chiesa, F. Tacchino, M. Grossi, P. Santini, I. Tavernelli, D. Gerace, and S. Carretta, *Nature Physics* **15**, 455 (2019).
- [35] A. Francis, J. Freericks, and A. Kemper, *Physical Review B* **101**, 014411 (2020).
- [36] P. Norman, D. M. Bishop, H. J. A. Jensen, and J. Oddershede, *The Journal of Chemical Physics* **123**, 194103 (2005).
- [37] S. Mukamel, *Annual Review of Physical Chemistry* **51**, 691 (2000).
- [38] A. W. Harrow, A. Hassidim, and S. Lloyd, *Physical Review Letters* **103**, 150502 (2009).
- [39] A. Ambainis, arXiv:1010.4458 (2010).
- [40] B. D. Clader, B. C. Jacobs, and C. R. Sprouse, *Physical*

- Review Letters **110**, 250504 (2013).
- [41] A. M. Childs, R. Kothari, and R. D. Somma, *SIAM Journal on Computing* **46**, 1920 (2017).
- [42] Y. Subaşı, R. D. Somma, and D. Orsucci, *Physical review letters* **122**, 060504 (2019).
- [43] X. Xu, J. Sun, S. Endo, Y. Li, S. C. Benjamin, and X. Yuan, arXiv preprint arXiv:1909.03898 (2019).
- [44] C. Bravo-Prieto, R. LaRose, M. Cerezo, Y. Subasi, L. Cincio, and P. J. Coles, arXiv preprint arXiv:1909.05820 (2019).
- [45] Y. Saad, *Iterative methods for sparse linear systems*, Vol. 82 (Society for Industrial and Applied Mathematics, 2003).
- [46] X.-D. Cai, C. Weedbrook, Z.-E. Su, M.-C. Chen, M. Gu, M.-J. Zhu, L. Li, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, *Physical Review Letters* **110**, 230501 (2013).
- [47] J. Pan, Y. Cao, X. Yao, Z. Li, C. Ju, H. Chen, X. Peng, S. Kais, and J. Du, *Physical Review A* **89**, 022313 (2014).
- [48] S. Barz, I. Kassal, M. Ringbauer, Y. O. Lipp, B. Dakić, A. Aspuru-Guzik, and P. Walther, *Scientific Reports* **4**, 6115 (2014).
- [49] Y. Zheng, C. Song, M.-C. Chen, B. Xia, W. Liu, Q. Guo, L. Zhang, D. Xu, H. Deng, K. Huang, *et al.*, *Physical Review Letters* **118**, 210504 (2017).
- [50] S. Aaronson, *Nature Physics* **11**, 291 (2015).
- [51] A. M. Childs and N. Wiebe, *Quantum Info. Comput.* **12**, 901 (2012).
- [52] D. W. Berry, A. M. Childs, R. Cleve, R. Kothari, and R. D. Somma, *Physical Review Letters* **114**, 090502 (2015).
- [53] P. Jordan and E. Wigner, *Z. Phys* **47**, 14 (1928).
- [54] S. B. Bravyi and A. Y. Kitaev, *Annals of Physics* **298**, 210 (2002).
- [55] J. T. Seeley, M. J. Richard, and P. J. Love, *The Journal of Chemical Physics* **137**, 224109 (2012).
- [56] H. Buhrman, R. Cleve, J. Watrous, and R. De Wolf, *Physical Review Letters* **87**, 167902 (2001).
- [57] J. C. Garcia-Escartin and P. Chamorro-Posada, *Physical Review A* **87**, 052330 (2013).
- [58] L. J. Ament, M. Van Veenendaal, T. P. Devereaux, J. P. Hill, and J. Van Den Brink, *Reviews of Modern Physics* **83**, 705 (2011).
- [59] H. A. Kramers and W. Heisenberg, *Z. Phys.* **31**, 681 (1925).