

Autocorrelation of a class of quaternary sequences of period $2p^m$

Qiuyan Wang, Weigang Kong, Yang Yan, Chenhuang Wu and Minghui Yang

Abstract

Sequences with good randomness properties are quite important for stream ciphers. In this paper, a new class of quaternary sequences is constructed by using generalized cyclotomic classes of \mathbb{Z}_{2p^m} ($m \geq 1$). The exact values of autocorrelation of these sequences are determined based on cyclotomic numbers of order 2 with respect to p^m . Results show that the presented sequences have the autocorrelations with at most 4 values.

keywords: autocorrelation, quaternary sequences, generalized cyclotomic numbers

I. INTRODUCTION

pseudorandom sequences with good cryptography properties have wide applications in code-division multiple-access systems, global positioning systems and stream ciphers. From the viewpoint of implementation, binary and quaternary sequences are preferred. Recently, quaternary sequences over \mathbb{Z}_4 have received a lot of study. The mainly methods for constructing quaternary sequences are the inverse Gray mapping and the generalized cyclotomy. In [6], [7], [8], [9], quaternary sequences over \mathbb{Z}_4 with good autocorrelation have been constructed by using the inverse Gray mapping. In [10], [11], [12], the authors used the generalized cyclotomy to present quaternary sequences with different periods.

The periodic autocorrelation function, as an important merit for sequence design, measures the level of similarity of a periodic sequence and a shift of this sequence. In [13], Ding et al. gave several families of binary sequences of period $2p$ with optimal three-level autocorrelation. In [14], Kim, Jin and Song determined the autocorrelation and the linear complexity of prime cube sequences. In [15], Edemskiy made use of generalized cyclotomic classes of \mathbb{Z}_{2p} to construct quaternary sequences with low autocorrelation. In [16], some quaternary cyclotomic sequences of period $2p$ were given and the autocorrelation function of these sequences were computed. Now it is interesting to see if it is possible to construct quaternary sequences of length $2p^m$, where m is a positive integer.

We first recall the definition of generalized cyclotomic classes of \mathbb{Z}_{2p^m} , where p is an odd prime. Assume that g is a primitive root of p^m . Without loss of generality, we assume that g is an odd integer. By Chinese Remainder Theorem [2], g is also a primitive root of $2p^m$. Furthermore, g is a common primitive root of p^i and $2p^i$ for any integer $1 \leq i \leq m$. Define

$$D_0^{(p^m)} = \langle g^2 \rangle = \{g^{2t} \bmod p^m : 0 \leq t \leq p^{m-1}(p-1)/2 - 1\}, D_1^{(p^m)} = gD_0^{(p^m)},$$

The work was supported by National Science Foundation of China No. 61602342, Natural Science Foundation of Tianjin under grant No. 18JCQNJC70300, NFSC 11701553, the Science and Technology Development Fund of Tianjin Education Commission for Higher Education No. 2018KJ215, KYQD1817, and the China Scholarship Council (No. 201809345010 and No. 201907760008), Key Laboratory of Applied Mathematics of Fujian Province University (Putian University) (No. SX201904 and SX201804), NSFT No.16JCYBJC42300, NFSC No. 61872359, 61972456 and 61802281, the Science and Technology Development Fund of Tianjin Education Commission for Higher Education No. 2017KJ213.

Q. Wang is with the School of Computer Science and Technology, Tiangong University, Tianjin, 300387, China, and with the Provincial Key Laboratory of Applied Mathematics, Putian University, Putian, Fujian 351100, China. Email: wangyan198801@163.com

W. Kong is with the School of Computer Science and Technology, Tiangong University, Tianjin, 300387, China. Email: 15002233496@163.com

Y. Yan is with the School of information and Technology Engineering, Tianjin University of Technology and Education, Tianjin, 300, China. Email: youngucas@126.com

C. Wu is with the Provincial Key Laboratory of Applied Mathematics, Putian University, Putian, Fujian 351100, China. Email: ptuwch@163.com

M. Yang is with the State Key Laboratory of Information Security, Institute of Information Engineering, Chinese Academy of Sciences, Beijing 100195, China. Email: yangminghui@iie.ac.cn.

$$D_0^{(2p^m)} = \langle g^2 \rangle = \{g^{2t} \bmod 2p^m : 0 \leq t \leq p^{m-1}(p-1)/2 - 1\}, D_1^{(2p^m)} = gD_0^{(2p^m)}.$$

Then $D_i^{(p^m)}$ and $D_i^{(2p^m)}$ for $0 \leq i \leq 1$ are called the generalized cyclotomic classes of order 2 of \mathbb{Z}_{p^m} and \mathbb{Z}_{2p^m} , respectively. For $1 \leq i \leq m$, it can be easily checked that

$$\mathbb{Z}_{p^i}^* = D_0^{(p^i)} \cup D_1^{(p^i)}, \quad \mathbb{Z}_{2p^i}^* = D_0^{(2p^i)} \cup D_1^{(2p^i)},$$

and

$$\begin{aligned} \mathbb{Z}_{2p^m} &= \bigcup_{i=1}^m p^{m-i} \left(\mathbb{Z}_{2p^i}^* \cup 2\mathbb{Z}_{p^i}^* \right) \cup \{0, p^m\} \\ &= \bigcup_{i=1}^m \left(p^{m-i} D_0^{(2p^i)} \cup p^{m-i} D_1^{(2p^i)} \cup 2p^{m-i} D_0^{(p^i)} \cup 2p^{m-i} D_1^{(p^i)} \right) \cup \{0, p^m\}. \end{aligned}$$

The quaternary sequence $\{s(n)\}$ of period $2p^m$ is defined as

$$s(n) = \begin{cases} 0, & \text{if } n \equiv 0 \pmod{2p}, \\ 2, & \text{if } n \equiv p \pmod{2p}, \\ 0, & \text{if } n \in D_0^{(2p^m)}, \\ 1, & \text{if } n \in D_1^{(2p^m)}, \\ 2, & \text{if } n \in 2D_0^{(p^m)}, \\ 3, & \text{if } n \in 2D_1^{(p^m)}. \end{cases} \quad (1)$$

A quaternary sequence $\{s(n)\}$ of period N over \mathbb{Z}_4 is called balanced if

$$\max_{j \in \mathbb{Z}_4} |\{i : 0 \leq i \leq N-1, s(i) = j\}| - \min_{j \in \mathbb{Z}_4} |\{i : 0 \leq i \leq N-1, s(i) = j\}| \leq 1.$$

Thus, the sequence $\{s(n)\}$ given by (1) is balanced for $m = 1$.

The rest of this paper is organized as follows. Basic definitions and related properties are introduced in Section II. Section III determines the autocorrelation of the quaternary sequence $\{s(n)\}$ defined by (1). Finally, we summarize our study in Section V.

II. PRELIMINARIES

In this section, we present some notations and preliminaries that will be used in the sequel. We begin with the definition of cross-correlation functions of periodic sequences.

Definition 1. *The corss-correlation function of quaternary sequences $\{u(n)\}$ and $\{v(n)\}$ of period N is defined by*

$$C_{u,v}(\tau) = \sum_{n=0}^{N-1} \omega^{u(n+\tau)-v(n)},$$

where ω is a complex 4-th root of unity.

When $\{u(n)\} = \{v(n)\}$, we write $C_u = C_{u,u}$ for short, and C_u is called the auto-correlation function of $\{u(t)\}$.

Lemma 1 ([3], [4]). $2 \in D_i^{(p)}$ if and only if $2 \in D_i^{(p^k)}$ for $k \geq 1$ and $i = 0, 1$.

Lemma 2 ([1], [2]). $2 \in D_0^{(p)}$ if $p \equiv \pm 1 \pmod{8}$, and $2 \in D_1^{(p)}$ if $p \equiv \pm 3 \pmod{8}$.

Lemma 3 (Lemma 5, [17]). *Assume that t is an odd integer, then for $i = 0, 1$ and $j = 1, 2, \dots, m$, we have*

$$1) \quad 2D_i^{(p^j)} + pt \pmod{2p^j} = \begin{cases} D_i^{(2p^j)}, & \text{if } 2 \in D_0^{(p)}, \\ D_{i+1}^{(2p^j)}, & \text{if } 2 \in D_1^{(p)}. \end{cases}$$

$$2) D_i^{(2p^j)} + 2pt \pmod{2p^j} = D_i^{(2p^j)}.$$

Lemma 4 (Lemma 1, [17]). *For $1 \leq k \leq m$ and $i = 0, 1$, we have*

- 1) $D_i^{(p^k)} = \{x + py : x \in D_i^{(p)}, y \in \mathbb{Z}_{p^{k-1}}\}$.
- 2) $D_i^{(2p^k)} = \{x + py + \delta_{x,y} : x \in D_i^{(p)}, y \in \mathbb{Z}_{p^{k-1}}\}$, where

$$\delta_{x,y} = \begin{cases} 0, & \text{if } x + py \text{ is odd,} \\ p^k, & \text{otherwise.} \end{cases}$$

Lemma 5. *Let symbols be the same as before. Then*

$$-1 \pmod{p^k} \in D_0^{(p^k)} \text{ if and only if } -1 \pmod{2p^k} \in D_0^{(2p^k)} \text{ for } k \geq 1.$$

Proof. It can be easily proved by Lemma 4. □

Let $(i, j)_{p^m}$ denote the cyclotomic numbers of order 2 with respect to p^m , i.e.,

$$(i, j)_{p^m} = \left| (D_i^{(p^m)} + 1) \cap D_j^{(p^m)} \right|.$$

The following lemma is well-known.

Lemma 6 (Theorem 1, [3]). *If $p \equiv 1 \pmod{4}$, then*

$$(1, 0)_{p^m} = (1, 1)_{p^m} = (0, 1)_{p^m} = \frac{(p-1)p^{m-1}}{4}, \quad (0, 0)_{p^m} = \frac{p^{m-1}(p-5)}{4}.$$

If $p \equiv 3 \pmod{4}$, then

$$(1, 0)_{p^m} = (0, 0)_{p^m} = (1, 1)_{p^m} = \frac{(p-3)p^{m-1}}{4}, \quad (0, 1)_{p^m} = \frac{(p+1)p^{m-1}}{4}.$$

Definition 2. *Let p be an odd prime. The two sequences $\{u(n)\}$ and $\{v(n)\}$ of period p^m are defined as*

$$u(n) = \begin{cases} 0, & \text{if } n \pmod{p^m} \in C_0^{(p^m)} = D_0^{(p^m)}, \\ 1, & \text{if } n \pmod{p^m} \in C_1^{(p^m)} = D_1^{(p^m)} \cup \{0\} \cup A; \end{cases}$$

$$v(n) = \begin{cases} 0, & \text{if } (2n+1) \pmod{2p^m} \in C_0^{(2p^m)} = D_0^{(2p^m)}, \\ 1, & \text{if } (2n+1) \pmod{2p^m} \in C_1^{(2p^m)} = B \cup D_1^{(2p^m)}, \end{cases}$$

where $A = p\mathbb{Z}_{p^{m-1}}$ and $B = \{p, 3p, \dots, (2p^{m-1})p\}$.

For $\tau \in \mathbb{Z}_{p^m}$, and $i, j = 0, 1$, define

$$\begin{aligned} d_u(i, j; \tau) &= |C_i^{(p^m)} \cap (C_j^{(p^m)} + \tau)|, \\ d_v(i, j; \tau) &= |C_i^{(2p^m)} \cap (C_j^{(2p^m)} + 2\tau)|, \\ d_{u,v}(i, j; \tau) &= |2C_i^{(p^m)} \cap (C_j^{(2p^m)} + 2\tau - 1)|, \\ d_{v,u}(i, j; \tau) &= |C_i^{(2p^m)} \cap (2C_j^{(p^m)} + 2\tau - 1)|. \end{aligned}$$

Lemma 7. *Let symbols be the same as before. Then we have*

$$d_u(1, 0; \tau) = \begin{cases} \frac{p^{m-1}(p+3)}{4}, & \text{if } \tau \in D_0^{(p^m)}, p \equiv 1 \pmod{4}, \\ \frac{p^{m-1}(p-1)}{4}, & \text{if } \tau \in D_1^{(p^m)}, p \equiv 1 \pmod{4}, \\ \frac{p^{m-1}(p+1)}{4}, & \text{if } p \equiv 3 \pmod{4}, \\ 0, & \text{if } \tau \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}. \end{cases}$$

Proof. It can be proved by Lemma 6. □

Lemma 8. *Let symbols be the same as before. Then we have*

$$d_v(1, 0; \tau) = \begin{cases} \frac{p^{m-1}(p-1)}{4}, & \text{if } \tau \in D_1^{(p^m)}, p \equiv 1 \pmod{8}, \\ & \text{or } \tau \in D_0^{(p^m)}, p \equiv 5 \pmod{8}, \\ \frac{p^{m-1}(p+3)}{4}, & \text{if } \tau \in D_0^{(p^m)}, p \equiv 1 \pmod{8}, \\ & \text{or } \tau \in D_1^{(p^m)}, p \equiv 5 \pmod{8}, \\ \frac{p^{m-1}(p+1)}{4}, & \text{if } \tau \in \mathbb{Z}_{p^m}^*, p \equiv 3 \text{ or } 7 \pmod{8}, \\ 0, & \text{if } \tau \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}. \end{cases}$$

Proof. Note that

$$\begin{aligned} d_v(1, 0; \tau) &= |C_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)| \\ &= |B \cap (C_0^{(2p^m)} + 2\tau)| + |D_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)|, \end{aligned}$$

where $B = \{p, 2p, \dots, (2p^{m-1} - 1)p\}$.

If $p \mid \tau$, by Lemma 3 we obtain $C_0^{(2p^m)} + 2\tau = C_0^{(2p^m)}$ and then $d_v(1, 0; \tau) = 0$.

In the sequel, we consider $\tau \in \mathbb{Z}_{p^m}^*$. In this case, we first compute $|B \cap (C_0^{(2p^m)} + 2\tau)|$. If $-2\tau \in D_0^{(p)}$, by Lemma 3

$$\begin{aligned} C_0^{(2p^m)} + 2\tau &= \{py + \delta_y : y \in \mathbb{Z}_{p^{m-1}}\}, \text{ where} \\ \delta_y &= \begin{cases} 0, & \text{if } y \text{ is odd,} \\ p^{m-1}, & \text{otherwise.} \end{cases} \end{aligned}$$

According to Lemmas 2 and 5, we have

1) when $p \equiv 1$ or $3 \pmod{8}$,

$$|B \cap (C_0^{(2p^m)} + 2\tau)| = \begin{cases} p^{m-1}, & \text{if } \tau \in D_0^{(p^m)}, \\ 0, & \text{if } \tau \in D_1^{(p^m)}; \end{cases}$$

2) when $p \equiv 5$ or $7 \pmod{8}$,

$$|B \cap (C_0^{(2p^m)} + 2\tau)| = \begin{cases} 0, & \text{if } \tau \in D_0^{(p^m)}, \\ p^{m-1}, & \text{if } \tau \in D_1^{(p^m)}. \end{cases}$$

Next, we determine $|D_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)|$. Since

$$\begin{aligned} |D_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)| &= |D_1^{(p^m)} \cap (D_0^{(p^m)} + 2\tau) \pmod{p^m}| \\ &= |2^{-1}\tau^{-1}D_1^{(p^m)} \cap (2^{-1}\tau^{-1}D_0^{(p^m)} + 1)|, \end{aligned}$$

by Lemma 2 we have

1) when $p \equiv \pm 1 \pmod{8}$,

$$|D_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)| = \begin{cases} (0, 1)_{p^m}, & \text{if } \tau \in D_0^{(p^m)}, \\ (1, 0)_{p^m}, & \text{if } \tau \in D_1^{(p^m)}; \end{cases}$$

2) when $p \equiv \pm 1 \pmod{8}$,

$$|D_1^{(2p^m)} \cap (C_0^{(2p^m)} + 2\tau)| = \begin{cases} (1, 0)_{p^m}, & \text{if } \tau \in D_0^{(p^m)}, \\ (0, 1)_{p^m}, & \text{if } \tau \in D_1^{(p^m)}; \end{cases}$$

Combining the above computations and Lemma 6, we can get the conclusions of this lemma. \square

Lemma 9. *Let symbols be the same as before. Then we have*

$$d_{u,v}(1, 0; \tau) = \begin{cases} \frac{p^{m-1}(p+3)}{4}, & \text{if } p \equiv 1 \pmod{8}, 2\tau - 1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(p-1)}{4}, & \text{if } p \equiv 1 \pmod{8}, 2\tau - 1 \in D_1^{(2p^m)}, \\ & \text{or } p \equiv 5 \pmod{8}, 2\tau - 1 \in \mathbb{Z}_{2p^m}^*, \\ \frac{p^{m-1}(p+1)}{4}, & \text{if } p \equiv 3 \pmod{8}, 2\tau - 1 \in D_1^{(2p^m)}, \\ & \text{or } p \equiv 7 \pmod{8}, 2\tau - 1 \in \mathbb{Z}_{2p^m}^*, \\ \frac{p^{m-1}(p-3)}{4}, & \text{if } p \equiv 3 \pmod{8}, 2\tau - 2 \in D_0^{(2p^m)}, \\ p^m, & \text{if } p \equiv \pm 1 \pmod{8}, 2\tau - 1 \equiv p \pmod{2p}, \\ p^{m-1}, & \text{if } p \equiv \pm 3 \pmod{8}, 2\tau - 1 \equiv p \pmod{2p}. \end{cases}$$

Proof. Note that

$$\begin{aligned} d_{u,v}(1, 0; \tau) &= |2C_1^{(p^m)} \cap (C_0^{(2p^m)} + 2\tau - 1)| \\ &= |2p\mathbb{Z}_{p^{m-1}} \cap (D_0^{(2p^m)} + 2\tau - 1)| + |2D_1^{(p^m)} \cap (D_0^{(2p^m)} + 2\tau - 1)|. \end{aligned} \quad (2)$$

It follows from Lemma 4 that

$$|2p\mathbb{Z}_{p^{m-1}} \cap (C_0^{(2p^m)} + 2\tau - 1)| = \begin{cases} p^{m-1}, & \text{if } -(2\tau - 1) \pmod{p} \in D_0^{(p)}, \\ 0, & \text{otherwise.} \end{cases} \quad (3)$$

Since $|2D_1^{(p^m)} \cap (D_0^{(2p^m)} + 2\tau - 1)| = |2D_1^{(p^m)} \cap (D_0^{(p^m)} + 2\tau - 1)|$, according to Lemma 2, Lemma 3 and Lemma 5 we have

1) if $p \equiv \pm 1 \pmod{8}$, then $2 \in D_0^{(p)}$ and

$$|2D_1^{(p^m)} \cap (D_0^{(2p^m)} + 2\tau - 1)| = \begin{cases} 0, & \text{if } 2\tau - 1 \equiv p \pmod{2p}, \\ (0, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ (1, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}. \end{cases} \quad (4)$$

2) if $p \equiv \pm 3 \pmod{8}$, then $2 \in D_1^{(p)}$ and

$$|2D_1^{(p^m)} \cap (D_0^{(2p^m)} + 2\tau - 1)| = \begin{cases} \frac{p^{m-1}(p-1)}{2}, & \text{if } 2\tau - 1 \equiv p \pmod{2p}, \\ (0, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ (1, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}. \end{cases} \quad (5)$$

Then the desirable results follow from Lemma 6, (2), (3) and (4). \square

Lemma 10. *Let symbols be the same as before. Then we have*

$$d_{v,u}(1, 0; \tau) = \begin{cases} \frac{p^{m-1}(p+3)}{4}, & \text{if } p \equiv 1 \pmod{8}, 2\tau - 1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(p-3)}{4}, & \text{if } p \equiv 3 \pmod{8}, 2\tau - 1 \in D_1^{(2p^m)}, \\ \frac{p^{m-1}(p-1)}{4}, & \text{if } p \equiv 5 \pmod{8}, 2\tau - 1 \in \mathbb{Z}_{2p^m}^*, \\ & \text{or } p \equiv 1 \pmod{8}, 2\tau - 1 \in D_1^{(2p^m)}, \\ \frac{p^{m-1}(p+1)}{4}, & \text{if } p \equiv 7 \pmod{8}, 2\tau - 1 \in \mathbb{Z}_{2p^m}^*, \\ & \text{or } p \equiv 3 \pmod{8}, 2\tau - 1 \in D_0^{(2p^m)}, \\ p^m, & \text{if } p \equiv \pm 1 \pmod{8}, 2\tau - 1 \equiv p \pmod{2p}, \\ p^{m-1}, & \text{if } p \equiv \pm 3 \pmod{8}, 2\tau - 1 \equiv p \pmod{2p}. \end{cases}$$

Proof. Note that

$$d_{v,u}(1, 0; \tau) = |B \cap (2C_0^{(2p^m)} + 2\tau - 1)| + |D_1^{(2p^m)} \cap (2C_0^{(p^m)} + 2\tau - 1)|, \quad (6)$$

where $B = \{p, 2p, \dots, (2p^{m-1} - 1)p\}$.

The computation on $d_{v,u}(1, 0; \tau)$ is divided into the following two cases.

Case (i). $2\tau - 1 \equiv p \pmod{2p}$. By Lemma 3, we know $|B \cap (2C_0^{(2p^m)} + 2\tau - 1)| = 0$ and

$$2C_0^{(p^m)} + 2\tau - 1 = \begin{cases} D_0^{(2p^m)}, & \text{if } 2 \in D_0^{(p)}, \\ D_1^{(2p^m)}, & \text{if } 2 \in D_1^{(p)}, \end{cases}$$

$$|D_1^{(2p^m)} \cap (2C_0^{(p^m)} + 2\tau - 1)| = \begin{cases} 0, & \text{if } p \equiv \pm 1 \pmod{8}, \\ \frac{p^{m-1}(p-1)}{2}, & \text{if } p \equiv \pm 3 \pmod{8}. \end{cases}$$

Hence, we obtain

$$d_{v,u}(1, 0; \tau) = \begin{cases} 0, & \text{if } p \equiv \pm 1 \pmod{8}, \\ \frac{p^{m-1}(p-1)}{2}, & \text{if } p \equiv \pm 3 \pmod{8}. \end{cases} \quad (7)$$

Case (ii). $2\tau - 1 \in \mathbb{Z}_{2p^m}^*$. Simple computation shows that

1) if $p \equiv \pm 1 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = |B \cap (C_0^{(2p^m)} + 2\tau - 1)| + |D_1^{(2p^m)} \cap (D_0^{(p^m)} + 2\tau - 1)|.$$

2) if $p \equiv \pm 3 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = |B \cap (D_1^{(2p^m)} + 2\tau - 1)| + |D_1^{(2p^m)} \cap (D_1^{(p^m)} + 2\tau - 1)|.$$

Since $|D_1^{(2p^m)} \cap (D_0^{(p^m)} + 2\tau - 1)| = |D_1^{(p^m)} \cap (D_0^{(p^m)} + 2\tau - 1)|$, by Lemma 4 we obtain

1) if $p \equiv 1 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = \begin{cases} p^{m-1} + (0, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ (1, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}; \end{cases} \quad (8)$$

2) if $p \equiv 3 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = \begin{cases} p^{m-1} + (1, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ (0, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}; \end{cases} \quad (9)$$

3) if $p \equiv 5 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = \begin{cases} (1, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ p^{m-1} + (0, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}; \end{cases} \quad (10)$$

4) if $p \equiv 7 \pmod{8}$, then

$$d_{v,u}(1, 0; \tau) = \begin{cases} (0, 1)_{p^m}, & \text{if } 2\tau - 1 \in D_0^{(2p^m)}, \\ p^{m-1} + (1, 0)_{p^m}, & \text{if } 2\tau - 1 \in D_1^{(2p^m)}. \end{cases} \quad (11)$$

The results of this lemma can be derived from (6)-(11) and Lemma 6.

□

III. AUTOCORRELATION OF QUATERNARY CYCLOTOMIC SEQUENCES WITH PERIOD $2p^m$

In order to determine the autocorrelation of $\{s(n)\}$ defined by (1), we introduce two kinds of quaternary cyclotomic sequences of period p^m .

Let

$$s^{(1)}(n) = \begin{cases} 0, & \text{if } n \equiv 0 \pmod{p}, \\ 2, & \text{if } n \in D_0^{(p^m)}, \\ 3, & \text{if } n \in D_1^{(p^m)}; \end{cases} \quad (12)$$

$$s^{(2)}(n) = \begin{cases} 2, & \text{if } 2n+1 \equiv p \pmod{2p}, \\ 0, & \text{if } 2n+1 \in D_0^{(2p^m)}, \\ 1, & \text{if } 2n+1 \in D_1^{(2p^m)}. \end{cases} \quad (13)$$

With the help of $\{s^{(1)}(n)\}$ and $\{s^{(2)}(n)\}$, the autocorrelation of the quaternary sequence $\{s(n)\}$ can be expressed as follows.

Lemma 11. *Let symbols be the same as before. Then we have*

$$C_s(\tau) = \begin{cases} C_{s^{(1)}}(k) + C_{s^{(2)}}(k), & \text{if } \tau \equiv 2k \pmod{2p^m}, 1 \leq k \leq p^m - 1, \\ C_{s^{(2)}s^{(1)}}(k-1) + C_{s^{(1)}s^{(2)}}(k), & \text{if } \tau \equiv 2k-1 \pmod{2p^m}, 1 \leq k \leq p^m. \end{cases}$$

Proof. The proof of this lemma can be directly obtained from Def. 1. \square

Lemma 12. *The autocorrelation of the quaternary sequence $\{s^{(1)}(n)\}$ of period p^m can be given by:*

1) *when $p \equiv 1 \pmod{4}$, we have*

$$C_{s^{(1)}}(k) = \begin{cases} \frac{p^{m-1}(p-7)}{2}, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-3)}{2}, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}; \end{cases}$$

2) *when $p \equiv 3 \pmod{4}$, we have*

$$C_{s^{(1)}}(k) = \begin{cases} \frac{p^{m-1}(p-5)}{2} + 2\omega, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-5)}{2} - 2\omega, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}. \end{cases}$$

Proof. The autocorrelation of $\{s^{(1)}(n)\}$ can be given as follows:

$$\begin{aligned} C_{s^{(1)}}(k) &= \sum_{i=0}^{p^m-1} \omega^{s^{(1)}(i+k)-s^{(1)}(i)} \\ &= \sum_{i \in E} \omega^{s^{(1)}(i+k)-s^{(1)}(i)} + \sum_{i \in D} \omega^{s^{(1)}(k+i)-s^{(1)}(i)} \\ &\quad + \sum_{\substack{i \in \mathbb{Z}_{p^m} \\ i \notin D \cup E}} \omega^{s^{(1)}(i+k)-s^{(1)}(i)}, \end{aligned}$$

where $D = p\mathbb{Z}_{p^{m-1}} - k$ and $E = p\mathbb{Z}_{p^{m-1}}$.

Note that

$$s^{(1)}(n) = \begin{cases} u(n) + 2, & \text{if } n \in \mathbb{Z}_{p^m}^*, \\ 0, & \text{if } n \in p\mathbb{Z}_{p^m}. \end{cases}$$

Hence, we **obtain**

$$\begin{aligned} C_{s(1)}(k) &= \sum_{i \in E} \omega^{u(i+k)+2} + \sum_{i \in D} \omega^{-2-u(i)} \\ &\quad + \sum_{i=0}^{p^m-1} \omega^{u(i+k)-u(i)} - \sum_{i \in D \cup E} \omega^{u(i+k)-u(i)}, \end{aligned} \quad (14)$$

It can be easily checked that

$$\begin{aligned} \sum_{i=0}^{p^m-1} \omega^{u(i+k)-u(i)} &= d_u(1, 0; k)\omega + d_u(0, 0; k)\omega^0 + d_u(0, 1; k)\omega^{-1} + d_u(1, 1; k)\omega^0 \\ &= p^m - 2d_u(1, 0; k), \end{aligned}$$

where the last identity was due to $d_u(1, 0; k) = d_u(0, 1; k)$ and

$$d_u(1, 0; k) + d_u(0, 0; k) + d_u(0, 1; k) + d_u(1, 1; k) = p^m.$$

Based on (14), we get

$$C_{s(1)}(k) = p^m - 2d_u(1, 0; k) \text{ for } k \in \mathbb{Z}_{p^m} \setminus \{0\}, \quad (15)$$

and

$$\begin{aligned} C_{s(1)}(k) &= \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{u(i+k)+2} + \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{-u(-k+i)-2} \\ &\quad + p^m - 2d_u(1, 0; k) - \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{u(i+k)-u(i)} - \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{u(i)-u(-k+i)} \end{aligned} \quad (16)$$

for $k \in \mathbb{Z}_{p^m}^*$.

When $p \equiv 1 \pmod{4}$, by Lemma 5 $k \in D_i^{(p^m)} \Leftrightarrow -k \in D_i^{(p^m)}$. Hence, $k \in D_j^{(p^m)}$ implies $u(i+k) = u(-k+i) = j$ for $j = 0, 1$. Similarly, when $p \equiv 3 \pmod{4}$, by Lemma 5 $k \in D_i^{(p^m)} \Leftrightarrow -k \in D_{(i+1) \bmod 2}^{(p^m)}$. Hence, $k \in D_j^{(p^m)}$ implies $u(i+k) = j$ and $u(-k+i) = (j+1) \bmod 2$ for $j = 0, 1$.

By Lemmas 2, 5 and 7 and the above analysis, we complete the proof of this lemma. \square

Lemma 13. *The autocorrelation of the quaternary sequence $\{s^{(2)}(n)\}$ of period p^m can be given by:*

1) when $p \equiv 1 \pmod{8}$, we have

$$C_{s(2)}(k) = \begin{cases} \frac{p^{m-1}(p-7)}{2}, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-3)}{2}, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}; \end{cases}$$

2) when $p \equiv 3 \pmod{8}$, we have

$$C_{s(2)}(k) = \begin{cases} \frac{p^{m-1}(p-5)}{2} - 2p^{m-1}\omega, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-5)}{2} + 2p^{m-1}\omega, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}; \end{cases}$$

3) when $p \equiv 5 \pmod{8}$, we have

$$C_{s(2)}(k) = \begin{cases} \frac{p^{m-1}(p-3)}{2}, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-7)}{2}, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}; \end{cases}$$

4) when $p \equiv 7 \pmod{8}$, we have

$$C_{s^{(2)}}(k) = \begin{cases} \frac{p^{m-1}(p-5)}{2} + 2p^{m-1}\omega, & \text{if } k \in D_0^{(p^m)}, \\ \frac{p^{m-1}(p-5)}{2} - 2p^{m-1}\omega, & \text{if } k \in D_1^{(p^m)}, \\ p^m, & \text{if } k \in p\mathbb{Z}_{p^{m-1}} \setminus \{0\}. \end{cases}$$

Proof. Note that

$$s^{(2)}(n) = \begin{cases} v(n), & \text{if } 2n+1 \in \mathbb{Z}_{p^m}^*, \\ 2, & \text{if } n \in C, \end{cases}$$

where

$$C = \{(p-1)/2, (3p-1)/2, \dots, ((2p^{m-1}-1)p-1)/2\}. \quad (17)$$

Similar to the procedure in the case of $\{s^{(1)}(n)\}$, the autocorrelation of $\{s^{(2)}(n)\}$ satisfies:

1) if $p \mid k$ and $k \neq 0$, then we have

$$\begin{aligned} C_{s^{(2)}}(k) &= p^m - 2d_v(1, 0; k) \\ &= p^m, \end{aligned} \quad (18)$$

where the last equality holds due to Lemma 8.

2) if $k \in \mathbb{Z}_{p^m}^*$, then we have

$$\begin{aligned} C_{s^{(2)}}(k) &= \sum_{i \in C} \omega^{2-v(-k+i)} + \sum_{i \in C} \omega^{v(i+k)-2} \\ &\quad - \sum_{i \in C} \omega^{1-v(-k+i)} - \sum_{i \in C} \omega^{v(i+k)-1} + p^m - 2d_v(1, 0; k). \end{aligned} \quad (19)$$

By the assumption, we know $2(-k+t) \in D_j^{(2p^m)}$ for $t \in C$ means that $-2k \in D_j^{(p^m)}$, and $2(k+t) \in D_j^{(2p^m)}$ for $t \in C$ means that $2k \in D_j^{(p^m)}$. And the inverse also holds. Therefore, we get

1) if $p \equiv 1 \pmod{8}$, then $k \in D_j^{(p^m)}$ if and only if $2k \in D_j^{(p^m)}$ if and only if $-2k \in D_j^{(p^m)}$ for $j = 0, 1$. Hence, $k \in D_j^{(p^m)}$ implies

$$v(i+k) = v(i-k) = j \text{ for } i \in C;$$

2) if $p \equiv 3 \pmod{8}$, then $k \in D_j^{(p^m)}$ if and only if $2k \in D_{(j+1) \bmod 2}^{(p^m)}$ if and only if $-2k \in D_j^{(p^m)}$ for $j = 0, 1$. Hence, $k \in D_j^{(p^m)}$ implies

$$v(i+k) = j+1 \pmod{2} \text{ and } v(i-k) = j \text{ for } i \in C;$$

3) if $p \equiv 5 \pmod{8}$, then $k \in D_j^{(p^m)}$ if and only if $2k \in D_{(j+1) \bmod 2}^{(p^m)}$ if and only if $-2k \in D_j^{(p^m)}$ for $j = 0, 1$. Hence, $k \in D_j^{(p^m)}$ implies

$$v(i+k) = v(i-k) = (j+1) \bmod 2 \text{ for } i \in C;$$

4) if $p \equiv 7 \pmod{8}$, then $k \in D_j^{(p^m)}$ if and only if $2k \in D_j^{(p^m)}$ if and only if $-2k \in D_{(j+1) \bmod 2}^{(p^m)}$ for $j = 0, 1$. Hence, $k \in D_j^{(p^m)}$ implies

$$v(i+k) = j \text{ and } v(i-k) = (j+1) \bmod 2 \text{ for } i \in C.$$

Combining the above analysis and Lemma 8, we get the conclusions of this lemma. \square

Lemma 14. For $1 \leq k \leq p^m$, the cross-correlation of the two quaternary sequences $\{s^{(1)}(n)\}$ and $\{s^{(2)}(n)\}$ of period p^m can be given by:

1) when $p \equiv 1 \pmod{8}$, we have

$$C_{s^{(1)}s^{(2)}}(k) = \begin{cases} -p^m, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(7-p)}{2}, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(3-p)}{2}, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

2) when $p \equiv 3 \pmod{8}$, we have

$$C_{s^{(1)}s^{(2)}}(k) = \begin{cases} -p^{m-1}, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(1-p)}{2}, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(5-p)}{2}, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

3) when $p \equiv 5 \pmod{8}$, we have

$$C_{s^{(1)}s^{(2)}}(k) = \begin{cases} -p^{m-1}, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(3-p)}{2} + 2p^{m-1}\omega, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(3-p)}{2} - 2p^{m-1}\omega, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

4) when $p \equiv 7 \pmod{8}$, we have

$$C_{s^{(1)}s^{(2)}}(k) = \begin{cases} -p^m, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(5-p)}{2} - 2p^{m-1}\omega, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(5-p)}{2} + 2p^{m-1}\omega, & \text{if } 2k-1 \in D_1^{(2p^m)}. \end{cases}$$

Proof. Let $\{s^{(1)}(n)\}$ and $\{s^{(2)}(n)\}$ be the two quaternary sequences defined by (12) and (13), respectively. We distinguish between the following two cases to calculate the cross-correlation of $\{s^{(1)}(n)\}$ and $\{s^{(2)}(n)\}$:

$$\begin{aligned} C_{s^{(1)}s^{(2)}}(k) &= \sum_{i=0}^{p^m-1} \omega^{s^{(1)}(i+k)-s^{(2)}(i)} \\ &= \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{-s^{(2)}(i-k)} + \sum_{i \in C} \omega^{s^{(1)}(i+k)+2} + \sum_{i=0}^{p^m-1} \omega^{u(i+k)+2-v(i)} \\ &\quad - \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{u(i)+2-v(i-k)} - \sum_{i \in C} \omega^{u(i+k)+2-v(i)}, \end{aligned} \tag{20}$$

where C is defined by (17).

Case (i): $2k-1 \equiv p \pmod{2p}$. In this case $s^{(2)}(-k+i) = 2$ and $v(-k+i) = 1$, $u(i+k) = 1$ and $s^{(1)}(i+k) = 0$ for $i \in C$. It follows from (20) that

$$C_{s^{(1)}s^{(2)}}(k) = - \sum_{i=0}^{p^m-1} \omega^{u(i+k)-v(i)}.$$

Case (ii): $2k-1 \in \mathbb{Z}_{2p^m}^*$. By (20), we have

$$\begin{aligned} C_{s^{(1)}s^{(2)}}(k) &= \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{-v(-k+i)} + \sum_{i \in C} \omega^{u(i+k)} - \sum_{i=0}^{p^m-1} \omega^{u(i+k)-v(i)} \\ &\quad + \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{-v(-k+i)} - \sum_{i \in C} \omega^{u(i+k)+1}. \end{aligned}$$

It is easy to verify that

$$d_{u,v}(0, 1; k) = d_{u,v}(1, 0; k),$$

$$\sum_{j=0}^1 \sum_{i=0}^1 d_{u,v}(i, j; k) = p^m.$$

Hence, we deduce

$$\sum_{i=0}^{p^m-1} \omega^{u(i+k)-v(i)} = p^m - 2d_{u,v}(1, 0; k).$$

According to Lemma 2 and Lemma 5, we have

- 1) if $p \equiv 1 \pmod{8}$, then $v(-k+i) = u(i+k) = j$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in C$;
- 2) if $p \equiv 3 \pmod{8}$, then $v(-k+i) = u(i+k) = j+1 \pmod{2}$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in C$;
- 3) if $p \equiv 5 \pmod{8}$, then $v(-k+i) = j$ and $u(i+k) = (j+1) \pmod{2}$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in C$;
- 4) if $p \equiv 7 \pmod{8}$, then $v(-k+i) = j+1 \pmod{2}$ and $u(i+k) = j$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in C$.

The desirable results follow from the above discussions together with Lemma 9. \square

Lemma 15. For $1 \leq k \leq p^m$, the cross-correlation of the two quaternary sequences $\{s^{(2)}(n)\}$ and $\{s^{(1)}(n)\}$ of period p^m can be given by:

- 1) when $p \equiv 1 \pmod{8}$, we have

$$C_{s^{(2)}s^{(1)}}(k) = \begin{cases} -p^m, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(7-p)}{2}, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(3-p)}{2}, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

- 2) when $p \equiv 3 \pmod{8}$, we have

$$C_{s^{(2)}s^{(1)}}(k) = \begin{cases} -p^{m-1}, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(5-p)}{2}, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(1-p)}{2}, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

- 3) when $p \equiv 5 \pmod{8}$, we have

$$C_{s^{(2)}s^{(1)}}(k) = \begin{cases} -p^{m-1}, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(3-p)}{2} - 2p^{m-1}\omega, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(3-p)}{2} + 2p^{m-1}\omega, & \text{if } 2k-1 \in D_1^{(2p^m)}; \end{cases}$$

- 4) when $p \equiv 7 \pmod{8}$, we have

$$C_{s^{(2)}s^{(1)}}(k) = \begin{cases} -p^m, & \text{if } 2k-1 \equiv p \pmod{2p}, \\ \frac{p^{m-1}(5-p)}{2} - 2p^{m-1}\omega, & \text{if } 2k-1 \in D_0^{(2p^m)}, \\ \frac{p^{m-1}(5-p)}{2} + 2p^{m-1}\omega, & \text{if } 2k-1 \in D_1^{(2p^m)}. \end{cases}$$

Proof. To begin with, we consider $C_{s^{(2)}s^{(1)}}(k)$.

When $2k-1 \equiv p \pmod{2p}$, it is easy to verify that

$$C_{s^{(2)}s^{(1)}}(k) = - \sum_{i=0}^{p^m-1} \omega^{v(k-1+i)-u(i)}.$$

When $2k-1 \in \mathbb{Z}_{2p^m}^*$, on the other hand, we can deduce that

$$\begin{aligned} C_{s^{(2)}s^{(1)}}(k) &= \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{v(k-1+i)} + \sum_{i \in C} \omega^{-u(-k+1+i)} - \sum_{i=0}^{p^m-1} \omega^{v(k-1+i)-u(i)} \\ &\quad + \sum_{i \in p\mathbb{Z}_{p^{m-1}}} \omega^{v(k-1+i)-1} - \sum_{i \in C} \omega^{-u(-k+1+i)-1}. \end{aligned}$$

Hence, we need to determine $\sum_{i=0}^{p^m-1} \omega^{v(k-1+i)-u(i)}$. Based on

$$\sum_{j=0}^1 \sum_{i=0}^1 d_{v,u}(i, j; k) = p^m$$

and

$$d_{v,u}(1, 0; k) = d_{v,u}(0, 1; k),$$

we obtain

$$\sum_{i=0}^{p^m-1} \omega^{v(k-1+i)-u(i)} = p^m - 2d_{v,u}(1, 0; k).$$

It follows from Lemma 2 and Lemma 5 that

- 1) if $p \equiv 1$ or $3 \pmod{8}$, then $v(-k+1+i) = u(k-1+i) = j$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in A$;
- 2) if $p \equiv 5$ or $7 \pmod{8}$, then $v(-k+1+i) = j$ and $u(k-1+i) = j+1 \pmod{2}$ for $2k-1 \in D_j^{(2p^m)}$ and $i \in A$.

Combining the above analysis and Lemma 10, we complete the proof of this lemma. \square

Theorem 16. *The autocorrelation of the constructed quaternary sequence $\{s(n)\}$ of period $2p^m$ can be given by:*

- 1) if $p \equiv 1 \pmod{8}$, we have

$$C_s(\tau) = \begin{cases} 2p^m, & \text{if } \tau \equiv 0 \pmod{2p}, \\ -2p^m, & \text{if } \tau \equiv p \pmod{2p}, \\ p^{m-1}(p-7), & \text{if } \tau \in 2D_0^{(p^m)}, \\ p^{m-1}(p-3), & \text{if } \tau \in 2D_1^{(p^m)}, \\ p^{m-1}(7-p), & \text{if } \tau \in D_0^{(2p^m)}, \\ p^{m-1}(3-p), & \text{if } \tau \in D_1^{(2p^m)}; \end{cases}$$

- 2) if $p \equiv 3$ or $5 \pmod{8}$, we have

$$C_s(\tau) = \begin{cases} 2p^m, & \text{if } \tau \equiv 0 \pmod{2p}, \\ -2p^{m-1}, & \text{if } \tau \equiv p \pmod{2p}, \\ p^{m-1}(p-5), & \text{if } \tau \in 2\mathbb{Z}_{p^m}^*, \\ p^{m-1}(3-p), & \text{if } \tau \in \mathbb{Z}_{2p^m}^*; \end{cases}$$

- 3) if $p \equiv 7 \pmod{8}$, we have

$$C_s(\tau) = \begin{cases} 2p^m, & \text{if } \tau \equiv 0 \pmod{2p}, \\ -2p^m, & \text{if } \tau \equiv p \pmod{2p}, \\ p^{m-1}(p-5) + 4p^{m-1}\omega, & \text{if } \tau \in 2D_0^{(p^m)}, \\ p^{m-1}(p-5) - 4p^{m-1}\omega, & \text{if } \tau \in 2D_1^{(p^m)}, \\ p^{m-1}(5-p) - 4p^{m-1}\omega, & \text{if } \tau \in D_0^{(2p^m)}, \\ p^{m-1}(5-p) + 4p^{m-1}\omega, & \text{if } \tau \in D_1^{(2p^m)}. \end{cases}$$

Proof. The desired conclusions on the auto-correlation of $\{s(n)\}$ follow from Lemma 11, Lemma 12, Lemma 13, Lemma 14, and Lemma 15. \square

Example 1. Let $m = 2$ and $p = 3$. Then

$$\begin{aligned} D_0^{(18)} &= \{1, 7, 13\}, & D_1^{(18)} &= \{5, 11, 17\}, \\ 2D_0^{(9)} &= \{2, 8, 14\}, & 2D_1^{(9)} &= \{4, 10, 16\}. \end{aligned}$$

It can be checked by Magma that

$$C_s(\tau) = \begin{cases} 0, & \text{if } \tau \in \mathbb{Z}_{18}^*, \\ 18, & \text{if } \tau \equiv 0 \pmod{6}, \\ -6 & \text{if } \tau \equiv 3 \pmod{6} \text{ or } \tau \in 2\mathbb{Z}_9^*, \end{cases}$$

which accords with Theorem 16.

IV. CONCLUDING REMARKS

In this paper, we present a class of quaternary cyclotomic sequences of period $2p^m$, where p is an odd prime. The auto-correlation of these quaternary sequences is determined and results show that they have at most 4 values of autocorrelation in terms of magnitude.

DATA AVAILABILITY

No data were used to support this study.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

- [1] D. M. Burton, Elementary Number Theory, 4th edn. McGraw-Hill International Editions, New York, 1998.
- [2] T. Cusick, C. Ding, A. Renvall, Stream ciphers and number theory. N.-Holl. Math. Libr. 55, 198-212, 1998.
- [3] Ding C, Helleseht T. New generalized cyclotomy and its applications, Finite fields and their applications, 4(2), 140-166, 1998.
- [4] C. Ding, Linear complexity of some generalized cyclotomic sequences, Int. J. Algebra Comput., 8(4), 431-442, 1998.
- [5] C. Ding, T. Helleseht, Generalized cyclotomy and its applications, Finite Fields Appl., 4, 467-474, 1999.
- [6] Y. -S. Kim, J. -W. Jang, S. -H. Kim, and J. -S. No, New construction of quaternary sequences with ideal autocorrelation from Legendre sequences, Proc. ISIT2009, 282-285, Seoul, Korea, 2009.
- [7] J. -W. Jang and D. -W. Lim, Large low probability of intercept properties of the quaternary sequence with optimal correlation property constructed by legendre sequences, Cryptogr. Commun., 8(4), 593-604, 2016.
- [8] V. Edemskiy and A. Ivanov, The linear complexity of balanced quaternary sequences with optimal autocorrelation value, Cryptogr. Commun., 7(4), 485-496, 2015.
- [9] X. Tang and C. Ding, New classes of balanced quaternary and almost balanced binary sequences with optimal autocorrelation value, IEEE Trans. Inf. Theory, 56(12), 6398-6405, 2010.
- [10] P. Ke, Z. Yang, and J. Zhang, On the autocorrelation and linear complexity of some $2p$ periodic quaternary cyclotomic sequences over \mathbb{F}_4 , IEICE Trans. Fundamentals, E94-A(11), 2472-2477, 2011.
- [11] X. Shen, Y. Jia, J. Wang, and L. Zhang, New families of balanced quaternary sequences of even period with three-level optimal autocorrelation, IEEE Commun. Lett., 21(10), 2150-2153, 2017.
- [12] D. Li, Q. Wen, J. Zhang, and Z. Chang, Linear complexity of generalized cyclotomic quaternary sequences with period pq , IEICE Trans. Fundamentals, E97-A(5), 1153-1158, 2014.
- [13] C. Ding, T. Helleseht, H. M. Martinsen, New families of binary sequences with optimal three-level autocorrelation, IEEE Trans. Inf. Theory, 47, 428C433, 2001.
- [14] Y. -J. Kim, S. -Y. Jin, H. -Y. Song, Linear complexity and autocorrelation of prime cube sequences. In Applied Algebra, Algebraic Algorithms, and Error-Correcting Codes; Lecture Notes in Computer Science, Springer: Berlin, Germany, 4851, 188-197, 2007.
- [15] V. Edemskiy and A. Ivanov, Autocorrelation and linear complexity of quaternary sequences of period $2p$ based on cyclotomic classes of order four, 2013 IEEE International Symposium on Information Theory, Istanbul, Turkey, 3120-3124, 2013.
- [16] Y. J. Kim, Y. P. Hong, H. Y. Song, Autocorrelation of some quaternary cyclotomic sequences of length $2p$, IEICE Transactions on Fundamentals of Electronics, Communications and Computer Sciences, 91(12), 3679-3684, 2008.
- [17] P. Ke, J. Zhang, S. Zhang, On the linear complexity and the autocorrelation of generalized cyclotomic binary sequences of length $2p^m$, Designs, codes and cryptography, 67(3), 325-339, 2013.