

Quantitative measure equivalence between finitely generated groups

Thiebaut Delabie*, Juhani Koivisto, François Le Maître† and Romain Tessera

June 2, 2021

Abstract

We initiate a quantitative study of measure equivalence (and orbit equivalence) between finitely generated groups, which extends the classical setting of L^p measure equivalence. We focus on two classes of groups: amenable groups and hyperbolic groups. The first one is motivated by the classical theorem of Ornstein and Weiss, and the second by lattices in rank one simple Lie groups. Our contribution consists in both rigidity and flexibility results.

On the rigidity side for amenable groups, we prove a general monotonicity property satisfied by the isoperimetric profile, which implies in particular its invariance under L^1 measure equivalence. This yields explicit “lower bounds” on how integrable a measure coupling between two amenable groups can be. This result also has an unexpected application to geometric group theory: the isoperimetric profile turns out to be monotonous under coarse embedding between amenable groups. This has various applications, among which the existence of an uncountable family of 3-solvable groups which pairwise do not coarsely embed into one another.

On the flexibility side, we construct explicit orbit equivalences between amenable groups with prescribed integrability conditions. Our main tool is a (new) notion of Følner tiling sequences. We show in a number of instances that the bounds given by the isoperimetric profile are sharp up to a logarithmic error. We also deduce from this study that two important quasi-isometric invariants are *not* preserved under L^1 orbit equivalence: the asymptotic dimension and finite presentability.

For hyperbolic groups, Shalom had proved that lattices in $SO(n, 1)$ are L^p measure equivalent for all $p < n - 1$. His proof yields a stronger statement, which we prove to be optimal in some sense. This is obtained via a general result saying that hyperbolicity is preserved under measure equivalence satisfying certain quantitative conditions.

Contents

1	Introduction	2
1.1	Monotonicity of the isoperimetric profile (and applications)	4
1.2	Construction of measure couplings between amenable groups	7
1.3	Rigidity of hyperbolicity	9
1.4	A sort of “asymmetric distance” between finitely generated groups	12
1.5	Plan of the paper	13
2	Variations on measure equivalence	13
2.1	Smooth actions and fundamental domains	13
2.2	Asymmetric couplings	15
2.3	Composition of asymmetric measure couplings	16

*supported by grant P2NEP2 181564 of the Swiss National Science Foundation

†Research partially supported by Projet ANR-17-CE40-0026 AGRUME, Projet ANR-19-CE40-0008 AODynG and IdEx Université de Paris, ANR-18-IDEX-0001.

2.4	Integrability conditions	18
2.5	Composition of φ -integrable couplings	20
2.6	Variations on orbit equivalence	24
3	Revisiting Lewis Bowen’s monotonicity theorem for the volume growth	26
4	Monotonicity of the isoperimetric profile	28
4.1	Overview of the proof	30
4.2	Λ -gradients on Λ -spaces	30
4.3	The induction technique	32
4.4	Monotonicity under at most m -to-one measure sub-quotients	34
4.5	Monotonicity under measure quotients	35
4.6	Can there be a quantitative version of the Ornstein-Weiss theorem?	38
5	L^∞ measure subgroups	39
5.1	L^∞ measure subgroups and regular maps	39
5.2	A continuum of 3-solvable groups	41
5.3	Stability of Liouville property for lamplighter groups	43
6	Følner tilings	44
6.1	Følner tiling sequences and orbit equivalence	44
6.2	Applications to groups with polynomial growth	46
6.3	A coupling between \mathbb{Z} and the lamplighter groups	49
7	Almost optimal couplings between \mathbb{Z} and iterated wreath products	51
7.1	Wreath products of measure-preserving equivalence relations	51
7.2	Wreath products and quantitative orbit equivalence	52
8	Unstable properties under exponential couplings	53
8.1	Finite presentability is unstable	53
8.2	Finite asymptotic dimension is unstable	55
9	Hyperbolicity	56
9.1	Geometric preliminaries	57
9.2	Rigidity of hyperbolicity	63

1 Introduction

Gromov introduced measure equivalence between countable groups as a measured analogue of quasi-isometry. A classical instance of a pair of measure equivalent groups is given by lattices in a common locally compact group. Another source of examples is given by orbit equivalent groups, i.e. groups which admit free measure-preserving actions on a standard probability space which share the same orbits.

The notion of measure equivalence has been extensively studied over the past 20 years, and we refer the reader to [Gab05, Sec. 2] for an overview of its main properties as well as its tight connections with invariants such as cost or ℓ^2 Betti numbers. Various rigidity phenomena have also been uncovered. A famous example is Furman’s superrigidity results for lattices in higher rank semi-simple Lie groups [Fur99], which implies for instance that any countable group which is measure-equivalent to a lattice in $\mathrm{PSL}_3(\mathbb{R})$ is commensurable up to finite kernel to another lattice in $\mathrm{PSL}_3(\mathbb{R})$. Another example is provided by Kida’s work on mapping class groups of surfaces: he showed that most surfaces can be reconstructed from the measure equivalence class of their mapping class group [Kid08], and that every

group which is measure equivalent to a mapping class group must actually be commensurable up to finite kernel to it [Kid10].

In the opposite direction of flexibility, a celebrated result of Ornstein and Weiss implies that all infinite countable amenable groups are orbit equivalent and hence measure equivalent [OW80]. So most coarse geometric invariants (such as volume growth) are not preserved under orbit equivalence. Also, it is known that the class of groups measure equivalent to lattices in $\mathrm{PSL}_2(\mathbb{R})$ is very diverse and contains groups that are not virtually isomorphic to lattices of the latter (for instance, all free products of infinite amenable groups belong to this class). But as we will now see, measure equivalence admits natural refinements which capture meaningful coarse geometric invariants.

Assume for simplicity that we are given an *orbit equivalence coupling* of two finitely generated groups $\Gamma = \langle S_\Gamma \rangle$ and $\Lambda = \langle S_\Lambda \rangle$ over a probability space (X, μ) , i.e. two measure-preserving free actions of Γ and Λ on (X, μ) which share the same orbits. Then we can equip the space (X, μ) with the Schreier graph metrics d_{S_Γ} and d_{S_Λ} , and consider for each $\gamma \in \Gamma$ and each $\lambda \in \Lambda$ the following *distance maps*

$$x \mapsto d_{S_\Lambda}(x, \gamma \cdot x) \text{ and } x \mapsto d_{S_\Gamma}(x, \lambda \cdot x).$$

The fact that the two actions share the same orbits means that these functions do not take the value $+\infty$, in other words they belong to the space $L^0(X, \mu, \mathbb{R})$ of measurable maps $X \rightarrow \mathbb{R}$. We then say that Γ and Λ are L^p orbit equivalent when all these functions are in $L^p(X, \mu; \mathbb{R})$. A similar definition can be given for measure equivalence, yielding the notion of L^p measure equivalence (see Section 2.4).

In the past ten years, L^1 measure equivalence has been intensely investigated. In the context of non-amenable groups, Bader, Furman and Sauer showed that any group that is L^1 measure equivalent to a lattice in $\mathrm{SO}(n, 1)$ for a given $n \geq 2$ must be virtually isomorphic to another such lattice [BFS13]. In particular, there is a L^1 -measure equivalence rigidity phenomenon for lattices in $\mathrm{PSL}_2(\mathbb{R})$, as opposed to the pure measure equivalence case. Somewhat more surprising is a result of Austin in the context of amenable groups: he showed that L^1 measure equivalent virtually nilpotent groups have bi-Lipschitz asymptotic cones [Aus16]. Another important result is due to Bowen, who showed in the appendix of Austin's aforementioned paper that volume growth is invariant under L^1 measure equivalence.

This paper is in the direct continuation of such results, but we also aim at a deeper understanding by going in two new directions: finer integrability notions and asymmetric versions of measure equivalence. For this introduction, let us however only give the definitions in the context of orbit equivalence (see Sec. 2.4 for the measure equivalence versions and its asymmetric counterparts).

Assume again that we are given two orbit equivalent actions of two finitely generated groups $\Gamma = \langle S_\Gamma \rangle$ and $\Lambda = \langle S_\Lambda \rangle$. Given any two unbounded increasing positive function $\varphi, \psi: (0, \infty) \rightarrow (0, \infty)$, we say that that we have a *(φ, ψ) -integrable orbit equivalence coupling from Γ to Λ* if for each $\gamma \in \Gamma$ and each $\lambda \in \Lambda$, there are constants $c_\gamma, c_\lambda > 0$ such that the associated distance functions satisfy the following conditions:

$$\int_X \varphi \left(\frac{d_{S_\Lambda}(x, \gamma \cdot x)}{c_\gamma} \right) d\mu(x) < \infty \text{ and } \int_X \psi \left(\frac{d_{S_\Gamma}(x, \lambda \cdot x)}{c_\lambda} \right) d\mu(x) < \infty.$$

Two remarks are in order: since X has finite measure, the (φ, ψ) -integrability of the distance functions is only sensitive to the speed at which φ and ψ tend to infinity. The constants $c_\gamma, c_\lambda > 0$ are partly motivated by the fact that we want our notion of (φ, ψ) -integrability to be independent of the choice of generating subset. We address these technical points in Section 2.4.

For consistency with the literature, we write L^p to replace the map $t \mapsto t^p$ for $p \in (0, \infty)$ and L^∞ if the distance maps are essentially bounded. If no assumption is made on the distance maps we shall write L^0 . Also we won't always use the adjective integrable for the sake of brevity.

Saying that two groups are L^p orbit equivalent for some $p \geq 1$ means in our terminology that there exists an (L^p, L^p) orbit equivalence coupling between them (the same is true for measure equivalence). Note that for $p < 1$, what one would like to call L^p orbit equivalence fails to be a transitive relation, so we will refrain from using such a terminology. Nevertheless, such a notion actually defines a very

natural pseudo-distance on the space of all finitely generated groups (see Sec. 1.4), and we will see for instance that the Heisenberg group and \mathbb{Z}^4 are at distance zero, while the distance between \mathbb{Z}^n and \mathbb{Z}^m is equal to $|\log n - \log m|$.

The rest of this introduction contains a summarized description of our main results, and is organized in four subsections. The first two ones are about amenable groups: §1.1 contains a general rigidity result related to the isoperimetric profile, while §1.2 has a flexibility flavor as it deals with constructions of concrete orbit equivalences. In §1.3, we state a rigidity result for hyperbolic groups. Finally, in 1.4 we briefly discuss a point of view that emerges from the notion of (φ, ψ) -integrable coupling (this can be seen as an introduction to the results of Section 2).

1.1 Monotonicity of the isoperimetric profile (and applications)

The isoperimetric profile. Bowen’s aforementioned result allows to distinguish groups with different growth functions up to L^1 measure equivalence. However this fails to address the case of solvable groups that are not virtually nilpotent, as these have exponential growth. By contrast the isoperimetric profile offers a fine invariant of amenable groups with exponential growth. Its asymptotic behavior has notably been computed for a wide class of solvable groups. Let us start recalling its definition.

Let Γ be a finitely generated group, and let S_Γ be a finite generating subset. Given a function $f: \Gamma \rightarrow \mathbb{R}$ and $p > 0$ we define the L^p -norm of its left gradient by the equation

$$\|\nabla_{S_\Gamma}^l f\|_p^p = \sum_{g \in \Gamma, s \in S} |f(sg) - f(g)|^p.$$

Given a function $f: \Gamma \rightarrow \mathbb{R}$, its support is the set $\text{supp } f = \{g \in \Gamma: f(g) \neq 0\}$. Following [Cou00, Now07, Tes08], we define for every $0 < p \leq \infty$, the ℓ^p -isoperimetric profile¹ of Γ as the non-decreasing function

$$j_{p,\Gamma}(n) = \sup_{|\text{supp } f| \leq n} \frac{\|f\|_p}{\|\nabla_{S_\Gamma}^l f\|_p}.$$

Of course this quantity depends on a choice of generating subset, but we will be interested in the asymptotic behavior of $j_{p,\Gamma}$, which does not.

Given two monotonous real-valued functions f and g we say that f is *asymptotically less* than g and write $f \preceq g$ if there exists a positive constant C such that $f(n) = O(g(Cn))$ as $n \rightarrow \infty$. We say that f and g are *asymptotically equivalent* and write $f \approx g$ if $f \preceq g$ and $g \preceq f$. The asymptotic behavior of f is its equivalence class modulo \approx .

Recall that for $p = 1$, the ℓ^1 isoperimetric profile is simply called the isoperimetric profile because of the following geometric interpretation [Cou00]:

$$j_{1,\Gamma}(n) \approx \sup_{|A| \leq n} \frac{|A|}{|\partial A|},$$

where $\partial A = S_\Gamma A \Delta A$. Certain authors prefer to work with the Følner function, defined for all $k \geq 1$ as follows

$$\text{Føl}_\Gamma(k) = \inf \left\{ |A|: \frac{|\partial A|}{|A|} \leq 1/k \right\}.$$

Note that the isoperimetric profile and the Følner function are generalized inverses of one another.

For $p = 2$, the asymptotic behavior of $j_{2,\Gamma}$ is intimately related to the probability of return of the simple random walks on the Cayley graph (Γ, S_Γ) as described in [CS93].

Note that a group is amenable if and only if its isoperimetric profile is unbounded. So the isoperimetric profile can be interpreted as a measurement of amenability: the faster it tends to infinity, the “more amenable” the group is. This is well illustrated by the following examples. For all $p \geq 1$ we have²:

¹Note that in [BZ21], the isoperimetric profile is defined as $I_{p,\Gamma} = \frac{1}{j_{p,\Gamma}}$.

²The notion of wreath product of groups (denoted by $G \wr H$) is recalled in §6.3.

- $j_{p,\Gamma}(n) \approx n^{1/d}$ for a group of polynomial growth of degree d (e.g. for $\Gamma = \mathbb{Z}^d$) (see [Cou00]);
- $j_{p,\Gamma}(n) \asymp \log n$ for all amenable groups with exponential growth [CS93];
- $j_{p,\Gamma}(n) \approx \log n$ for a wide class of solvable groups with exponential growth including those that are polycyclic, or the so-called “lamplighter” groups, i.e. group of the form $F \wr \mathbb{Z}$, where F is a finite non-trivial group [CGP01, Pit95, Pit00, Tes08];
- $j_{1,\Gamma}(n) \approx (\log n)^{1/d}$ for a group of the form $F \wr \Sigma$, where F is a non-trivial finite group and Σ has polynomial growth of degree d [Ers03].

A general monotonicity result. Our first main result is the following theorem.

Theorem 1. *Assume that there exists (φ, L^0) measure equivalence coupling from a finitely generated group Γ to a finitely generated group Λ . Then*

- if $\varphi(t) = t^p$ for some $p \geq 1$, then

$$j_{p,\Gamma} \succcurlyeq j_{p,\Lambda};$$

- if φ and $t \mapsto t/\varphi(t)$ are increasing then

$$j_{1,\Gamma} \succcurlyeq \varphi \circ j_{1,\Lambda}.$$

In particular, the ℓ^p -isoperimetric profile is stable under L^p measure equivalence for all $p \geq 1$. As a concrete application we deduce for instance that $\mathbb{Z} \wr \mathbb{Z}$ is not L^1 measure equivalence to $\mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}$, nor to any polycyclic group. Indeed, for these groups, the isoperimetric profile is $\asymp \log n$, while for $\mathbb{Z} \wr \mathbb{Z}$ it is $\approx \log n / \log \log n$ [Ers03].

Remark 1.1 (Isoperimetric profile versus Følner function). In the special case $p = 1$, we deduce that the asymptotic behavior of the Følner function is invariant under L^1 measure equivalence and the second statement translates into the following inequality: $F_{\text{øl}} \succcurlyeq F_{\text{øl}_\Lambda} \circ \varphi^{-1}$. This strengthens the following unpublished result obtained by Bowen in 2013: assuming Γ and Λ are L^1 orbit equivalent, there exists $r > 1$ such that $F_{\text{øl}_\Gamma}(k) \leq F_{\text{øl}_\Lambda}(r^k)$.

A special case of interest is when $\Gamma = \mathbb{Z}$: since $j_{1,\mathbb{Z}}(n) \approx n$, the second statement of Theorem 1 says in this case that $\varphi(j_{1,\Gamma}(n)) \succcurlyeq n$. Translated in terms of Følner functions, this means that $\varphi \succcurlyeq F_{\text{øl}_\Gamma}$. This triggers the following question asking for some kind of converse.

Question 1.2. Given an amenable finitely generated group Γ , does there exist a $(F_{\text{øl}_\Gamma}, L^0)$ orbit equivalence coupling from Γ to \mathbb{Z} ?

We shall see in the next subsection that this holds up to a logarithmic error in various examples.

Application to regular maps. Theorem 1 turns out to be a special case of a more general monotonicity property of the ℓ^p -isoperimetric profile (Theorems 4.2 and 4.3) which encompasses its well-known monotonicity under subgroups and quotients [Tes08]. Writing down this statement motivated us to introduce and study natural “measured generalizations” of the notions of subgroup, quotient, and subgroup of a quotient (see Sec. 2.2).

Let us mention here a striking application of this monotonicity result. Recall that a regular map $f: \Gamma \rightarrow \Lambda$ between two finitely generated groups is a Lipschitz map whose pre-images have uniformly bounded cardinality. Particular cases of regular maps are Lipschitz injective embeddings and coarse embeddings. We recall that a coarse embedding is a Lipschitz map $f: \Gamma \rightarrow \Lambda$ such that there exists a proper increasing function $\rho: [0, \infty) \rightarrow [0, \infty)$ satisfying

$$d_\Lambda(f(\gamma), f(\gamma')) \geq \rho(d_\Gamma(\gamma, \gamma')),$$

for all $\gamma, \gamma' \in \Gamma$.

A trick essentially due to Shalom allows us to interpret the existence of a regular map from an amenable group Γ to a group Λ as “ Γ being an L^∞ measure subgroup of Λ ” (see §2.4). This enable us to deduce from Theorem 4.2 the following rather unexpected fact.

Theorem 2 (see Cor. 5.6). *For every $1 \leq p \leq \infty$, the ℓ^p -isoperimetric profile is monotonous under regular map between amenable groups.*

The mere monotonicity under coarse embedding is already very surprising and actually false if the embedded group is not assumed to be amenable: for instance the lamplighter group $\mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}$ admits a quasi-isometrically embedded 4-regular tree. Hence, the free group on 2 generators quasi-isometrically (and hence coarsely) embeds into the lamplighter group (see for instance [Woe05]). But the free group is non-amenable and therefore its isoperimetric profile is bounded below, while the isoperimetric profile of the lamplighter, which is amenable, tends to zero.

Only very few geometric invariants are known to be monotonous under coarse embedding: these are the volume growth, the asymptotic dimension, the separation profile [BST12], and more recently the Poincaré profiles [HMT20]. All of these turn out to be monotonous under regular map as well (see [BST12] for the case of asymptotic dimension). For solvable groups of exponential growth, the asymptotic dimension is generally infinite, and the Poincaré profiles have only been computed in very specific examples. By contrast, our theorem provides us with a powerful obstruction. As a concrete application, it prevents the existence of a regular map from $\mathbb{Z} \wr \mathbb{Z}^a$ to $\mathbb{Z} \wr \mathbb{Z}^b$ as soon as $a > b$ (note that these groups have infinite asymptotic dimension). Combining it with a construction of Erschler and Zheng [EZ17, Cor. 3.3] (see also [BZ21, Thm. 1.1]) we obtain the following result.

Theorem 3 (see Cor. 5.8). *There exists an uncountable family of 3-step solvable groups Γ_i that do not pairwise regularly embed into one another, and that are pairwise non L^1 measure equivalent. Moreover, one can assume that these groups all have asymptotic dimension one.*

We note that finding an uncountable family of groups that do not pairwise regularly embed into one another can be done by other means. Indeed, one can produce an uncountable family of groups of intermediate growth whose growth functions are pairwise incomparable (this can be done exploiting [Gri85, Theorem 7.1], or more directly invoking [Bri14, Theorem 7.2]). Moreover by Lewis Bowen’s theorem (see Theorem 3.1), these groups are pairwise non L^1 measure equivalent. Other examples of uncountable families of groups that pairwise do not coarsely embed into one another are due to Hume [Hum17, Thm. 1.2]: they involve (non-amenable) groups that contain isometrically embedded families of expanders. However none of these other methods can yield families of solvable (nor even elementary amenable) groups.

No quantitative version of Orstein-Weiss. In a recent groundbreaking work, Brioussel and Zheng managed to construct amenable groups with prescribed isoperimetric profiles [BZ21]. In particular, they show that for every increasing unbounded function φ , there exists an amenable group whose isoperimetric profile does not dominate φ . Combining this with Theorem 1, we deduce that Orstein-Weiss’ orbit equivalence coupling between amenable groups can be as poorly integrable as possible. More precisely, we obtain the following corollary.

Corollary 4 (see §4.6). *For every amenable finitely generated group Λ and every increasing unbounded function φ , there exists an amenable finitely generated group Γ such that there are no (φ, L^0) measure equivalence coupling from Γ to Λ .*

Stability of Liouville property among lamplighter groups. Let G be a countable infinite group and let μ be a probability measure on G whose support generates G (as a group). A function $f : G \rightarrow \mathbb{R}$ is called harmonic if for all $g \in G$,

$$f(g) = \sum_{g' \in G} \mu(gg').$$

If all bounded harmonic functions are constant, then we say that (G, μ) has the Liouville property. In what follows, to simplify the discussion, we shall say that a measure μ is admissible if it is symmetric, and its support is finite and generates G . We refer to [KV83] for an introduction to this topic.

A well-known open question due to Itai Benjamini is whether Liouville property is “invariant under quasi-isometry”. More precisely does there exist (G_1, μ_1) and (G_2, μ_2) such that the measures μ_i are admissible, the groups G_i are quasi-isometric, and (G_1, μ_1) is Liouville while (G_2, μ_2) is not? Intuitively, being Liouville is a property of “small groups”, so we might even expect it to be stable under coarse (or even regular) embeddings. However, this is false in general as shown by the following counterexample: being non-amenable, the free group on two generators is non-Liouville (with respect to any μ), while $(\mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}, \mu)$ is Liouville for any admissible measure μ (see [KV83]). But as noticed earlier, the free group coarsely embeds into $\mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}$.

By contrast, Theorem 4.2 can be used to prove that being Liouville is stable under regular embedding (and under L^1 measure equivalence) among lamplighter groups³. More precisely, we obtain:

Theorem 1.3 (See Theorem 5.12). *For $i = 1, 2$, let F_i be a non-trivial finite group, and let H_i be a finitely generated group. Let μ_i be an admissible probability measure on $\Gamma_i = F_i \wr H_i$. Assume either that*

- Γ_1 regularly embeds into Γ_2 ,
- or there exists a (L^1, L^0) measure equivalence coupling from Γ_1 to Γ_2 .

Then if (Γ_2, μ_2) is Liouville, then so is (Γ_1, μ_1) .

1.2 Construction of measure couplings between amenable groups

Følner tilings. Most of the known examples of L^p measure equivalence couplings come from lattices in semi-simple Lie groups. Here we introduce a general method for constructing orbit equivalent couplings with prescribed integrability conditions between amenable groups. Our main tool is a notion of Følner tiling, developed in §6.1. A Følner tiling sequence in a group Γ is a sequence of finite subsets T_n satisfying three conditions:

- (i) (T_n) is a left Følner sequence for Γ ,
- (ii) each T_n is obtained as a union of right translates of T_{n-1} .

We note that Følner sequences T_n such that Γ can be obtained as a disjoint union of right translates of T_n do exist in all elementary amenable groups and all residually finite amenable groups thanks to the work of Weiss [Wei01]. Our second condition appears to be new, and is crucial in our construction. An easy example of a Følner tiling sequence is provided by $T_n = \{1, \dots, 2^n\}$ in \mathbb{Z} .

Starting from a Følner tiling sequence for a group Γ , we construct a free measure preserving action of Γ on the infinite product space $\prod_n (F_n, \nu_n)$, where F_n is a sequence of finite subsets of Γ such that $T_n = \bigsqcup_{\gamma \in F_n} T_{n-1}\gamma$, and where ν_n is the renormalized counting measure on F_n . We show that the equivalence relation generated by the action is the co-finite equivalence relation (in the above example the action is given by the dyadic odometer): two sequences are equivalent if they coincide except possibly on finitely many indices. It follows from our construction that if another group Λ admits a Følner tiling sequence whose basic tiles T'_n satisfy $|T_n| = |T'_n|$, then the corresponding action of Λ is orbit equivalent to that of Γ . Moreover, the degree of integrability of this orbit equivalence is controlled by the properties of the two Følner sequences (T_n) and (T'_n) (see Proposition 6.7).

Theorem 1 provided us with an obstruction for finding φ -integrable couplings with certain functions φ between two amenable groups. We will now show that in many cases of interest (and especially in the case of couplings from certain amenable groups to \mathbb{Z}) this obstruction is close to being optimal.

³The term lamplighter group is commonly used for groups of the form $F \wr H$, where F is a non-trivial finite group.

Let us start with the case of groups with polynomial growth. We deduce from a straightforward extension of Bowen’s theorem on volume growth (see Corollary 3.4), or alternatively from the monotonicity of the isoperimetric profile (Corollary 4.6), that \mathbb{Z}^m and \mathbb{Z}^n do not admit L^p measure equivalence couplings for $p > m/n$ if $n < m$. We show that this threshold is sharp:

Theorem 5 (see Thm. 6.10). *For every pair of positive integers $n < m$, there exists an orbit equivalence coupling between \mathbb{Z}^m and \mathbb{Z}^n that is L^p for every $p < n/m$.*

Let us mention that these couplings between \mathbb{Z}^m and \mathbb{Z}^n were independently discovered by Matthieu Joseph.

Note that \mathbb{Z}^n being a “smaller group” than \mathbb{Z}^m , we expect the coupling to be less integrable from \mathbb{Z}^m to \mathbb{Z}^n . Indeed, Theorem 6.10 yields an orbit equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n which is $(L^p, L^{p'})$ for all $p < n/m$ and $p' < m/n$, while Corollary 3.4 implies that there is no (L^p, L^0) measure equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n for $p > n/m$.

Theorem 5 leaves open the following question (see also Question 6.12).

Question 1.4. Let $n < m$, are \mathbb{Z}^m and \mathbb{Z}^n $L^{n/m}$ measure equivalent?

We may also consider the following asymmetric version of this question: is there a $(L^{n/m}, L^0)$ measure coupling from \mathbb{Z}^m to \mathbb{Z}^n ?

Regarding groups with same degree of growth, Austin proved that L^1 measure equivalent groups of polynomial growth have bi-Lipschitz asymptotic cones [Aus16]. Combined with a famous theorem of Pansu [Pan89, Thm. 3], this implies for instance that a non-virtually abelian nilpotent group cannot be L^1 measure equivalent to an abelian group. By contrast we show the following surprising converse for the Heisenberg group.

Theorem 6 (see Thm. 6.14). *There exists an orbit equivalence coupling between \mathbb{Z}^4 and $\mathbb{H}(\mathbb{Z})$ that is L^p for every $p < 1$.*

In all cases considered in this paper, the actions constructed by means of Følner tiling sequences are profinite actions but it seems likely that non profinite actions can be built in this manner as well. To the best of our knowledge, this way of producing orbit equivalences between amenable groups is new, and quite different from the much more abstract quasi-tiling machinery of Ornstein and Weiss [OW87]. Note that latter allows them to prove that *all* free measure-preserving ergodic actions of *all* amenable groups are orbit equivalent. This brings us to the complementary problem of studying quantitatively orbit equivalence between actions of a fixed amenable group. This is obviously a vast program, so we shall simply mention the following natural question, which is already interesting for the group \mathbb{Z} .

Question 1.5. Let Γ be an amenable finitely generated group. Does there exist a single unbounded increasing function φ such that any two free measure-preserving ergodic actions of Γ are φ orbit equivalent?

Orbit equivalence and wreath products. In §7.2, we provide a natural notion of “wreath product of orbit equivalence couplings”. This construction allows us to obtain orbit equivalence couplings between wreath product of groups with good integrability conditions (see Corollary 7.3 for a precise statement). In this paragraph, we will focus on some concrete applications of this general result.

Theorem 7. *For every pair of positive integers $a < b$, and every non-trivial finite group F there exists an $(L^p, L^{1/p})$ orbit equivalence coupling from $F \wr \mathbb{Z}^b$ to $F \wr \mathbb{Z}^a$ for every $p < a/b$.*

On the other hand, we deduce from the monotonicity of the isoperimetric profile that there exists no (L^p, L^0) measure equivalence coupling from $F \wr \mathbb{Z}^a$ to $F \wr \mathbb{Z}^b$ for $p > 1 - 1/a + 1/b$. Note that this is sharp for $a = 1$.

We now focus on the existence of couplings with \mathbb{Z} . We first consider couplings between \mathbb{Z} with the lamplighter group. Since the isoperimetric profile of the lamplighter group decays like $1/\log n$, we know that a (φ, L^0) coupling from the latter to \mathbb{Z} must satisfy $\varphi(t) \preccurlyeq \log t$. This turns out to be sharp up to a logarithmic error.

Theorem 8 (see Prop. 6.18). *For every finite group F , there exists an orbit equivalence coupling from $F \wr \mathbb{Z}$ to \mathbb{Z} which is $(\varphi_\varepsilon, \text{exp})$ -integrable for every $\varepsilon > 0$, where $\varphi_\varepsilon(x) = \frac{\log(x)}{\log(\log(x))^{1+\varepsilon}}$.*

This result is obtained by means of Følner tilings. Once again, the asymmetry of the condition reflects the intuition that \mathbb{Z} is a “small” group compared to $F \wr \mathbb{Z}$.

More generally, one can consider iterated wreath products as follows. Let $G_0 = \mathbb{Z}$, and for all $n \geq 1$, $G_n = F \wr G_{n-1}$. By [Ers03], we know that the isoperimetric profile of G_k is asymptotically equivalent to \log^{*n} , namely the n -times iteration of \log with itself. Similarly we show that this is almost sharp: see Corollary 7.5.

Exponentially integrable couplings. The lamplighter group $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ and the Baumslag-Solitar group $\text{BS}(1, k)$ are known to have very similar geometries. Indeed the first one can be seen as a horocyclic product of two $(k+1)$ -regular trees [Woe05], while the second is quasi-isometric to a horocyclic product of a $(k+1)$ -regular tree with a hyperbolic plane [FM98]. A consequence of this description is that they have isometric asymptotic cones (this follows from the analysis in [dC08, §9]). They are also known to have same isoperimetric profile [CGP01]. However they are not quasi-isometric, as $\text{BS}(1, k)$ is finitely presented, while $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ is not.

In §8.1, we construct a very natural orbit coupling between these groups, which satisfies a very strong integrability condition.

Theorem 9 (see Thm. 8.1). *For every $k \geq 2$, there exists an (L^∞, exp) orbit equivalence coupling from $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ to $\text{BS}(1, k)$.*

The asymmetry is not an artifact of the proof: indeed we could not have an (L^∞, exp) -orbit equivalence coupling from $\text{BS}(1, k)$ to $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ as the asymptotic dimension is monotonous under (L^∞, L^0) orbit equivalence (Corollary 5.5), and $\text{BS}(1, k)$ has asymptotic dimension 2, while $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ has asymptotic dimension 1. Moreover, this shows that the asymptotic dimension is not preserved under L^p orbit equivalence for all $p < \infty$. Considering slightly more sophisticated examples, we similarly show that the finiteness of the asymptotic dimension is not preserved either.

1.3 Rigidity of hyperbolicity

The main motivation to study quantitative versions of measure equivalence is to distinguish groups that are measure equivalent but have different geometric properties. Beyond the case of amenable groups, another source of examples are lattices in a same locally compact group. Recall⁴ that uniform lattices in rank one simple Lie groups are hyperbolic but non-uniform ones are not hyperbolic, except for $\text{SL}(2, \mathbb{R})$ where all lattices are hyperbolic. Hence these are interesting instances of groups that are measure equivalent, yet with different geometric properties. With this main application in mind, we will look for general integrability conditions ensuring that hyperbolicity is preserved under measure equivalence. We will not state our main results in this introduction as these are quite technical (see Theorem 9.12 and Theorem 9.11). Instead we shall focus on two specific corollaries.

Uniform versus non-uniform lattices in rank one simple Lie groups. Shalom proved that any two lattices in $\text{SO}(n, 1)$ are L^p measure equivalent for all $p < n - 1$. When one of the two lattices is uniform, one can strengthen this statement as follows. Let $n \geq 2$, and let Γ (resp. Λ) be a uniform (resp. non-uniform) lattice in $\text{SO}(n, 1)$. We consider the coupling associated to the action of Λ and Γ respectively by left and right-translations on the measure space $\text{SO}(n, 1)$ equipped with an invariant Haar measure. Shalom showed that for a suitable fundamental domain X_Λ for Λ and any relatively compact fundamental domain X_Γ for Γ , the resulting coupling is an (L^p, L^∞) measure equivalence coupling from Γ to Λ for all $p < n - 1$. Exploiting the relative compactness of X_Γ , one can check that this coupling satisfies the following additional property: there is a finite subset $F \subset \Lambda$ such that

⁴This is classical: see for instance [DuB, Example 22.1 and Proof of Theorem 22.32].

$X_\Gamma \subset FX_\Lambda$. In what follows, we shall refer to this property that as *coboundedness* of the coupling. We summarize this as follows.

Theorem 10 (Shalom). *Let Γ and Λ be two lattices in $SO(n, 1)$ such that Γ is uniform. Then there exists a cobounded coupling from Γ to Λ that is (L^p, L^∞) -integrable for all $p < n - 1$.*

By contrast we prove the following rigidity result.

Theorem 11 (see Cor. 9.13). *Let Γ be a finitely generated hyperbolic group. There exists $p > 0$ such that if there exists a cobounded (L^p, L^∞) measure equivalence coupling from Γ to a finitely generated group Λ , then Λ is also hyperbolic.*

Remark 1.6. The value of p for which the conclusion holds is explicit: assuming that Γ admits a Cayley graph that is δ -hyperbolic and has volume entropy at most α , one can take $p = 75\delta\alpha + 2$. For the definition of volume entropy, see the paragraph which precedes Theorem 9.11.

We immediately deduce the following converse of Shalom's result.

Corollary 12. *Assume that Γ is a uniform lattice in a center-free, real rank 1 simple Lie group G and Λ is another lattice of G . There exists p only depending on G such that if there exists a cobounded (L^p, L^∞) coupling from Γ to Λ , then Λ must be uniform as well.*

This raises the following question.

Question 1.7. What is the infimum over all p such that the previous result holds? Does it match Shalom's value $n - 1$ for lattices in $SO(n, 1)$?

Note that the L^∞ condition from Λ to Γ is the strongest possible. It can be relaxed to an L^p -type condition, but at the cost of imposing a stretched exponential integrability condition in the other direction. More precisely, we obtain:

Theorem 13 (see Cor. 9.14). *Let Γ be a finitely generated hyperbolic group. For every $p > q > 0$, if there is a cobounded (φ, ψ) -integrable measure equivalence coupling from Γ to any finitely generated group Λ where $\varphi(t) = \exp(t^p)$ and $\psi(t) = t^{1+1/q}$, then Λ is also hyperbolic.*

Once again we deduce the following corollary for lattices in rank 1 simple Lie groups.

Corollary 14. *Assume that Γ is a uniform lattice in a center-free, real rank 1 simple Lie group G and Λ is another lattice of G . For every $p > q > 0$, if there is a cobounded (φ, ψ) -integrable measure equivalence coupling from Γ to Λ where $\varphi(t) = \exp(t^p)$ and $\psi(t) = t^{1+1/q}$, then Λ is uniform as well.*

As observed by Mikael de la Salle, Corollaries 12 and 14 are in sharp contrast with what happens for lattices in higher rank simple Lie groups: indeed if Γ and Λ are lattices in a simple Lie group Γ of rank ≥ 2 , then if X_Λ and X_Γ are Dirichlet fundamental domains for Λ and Γ , the resulting measure equivalence coupling is exponentially integrable [de 19, Lemme 5.6]. Hence we obtain:

Theorem 15 (de la Salle). *Let Γ and Λ be two lattices in a simple Lie group of rank at least 2, such that Γ is uniform. Then there exists a cobounded (φ, L^∞) measure equivalence coupling from Γ to Λ , where $\varphi(t) = \exp(ct)$ for some $c > 0$.*

Remark 1.8. In Corollaries 12 and 14, the case of $SL(2, \mathbb{R})$ was already known and actually a much stronger conclusion holds in that case: Bader, Furman and Sauer have proved that non-uniform lattices and uniform ones are not L^1 measure equivalent.

Remark 1.9. Theorems 11 and 13 should be compared with a theorem of Bowen saying that if there exists an (L^1, L^0) orbit equivalence coupling from a finitely generated accessible group Λ to a virtually free group, then Λ is virtually free.

Stability of hyperbolicity under quantitative orbit equivalence The connection between measure equivalence and orbit equivalence is recalled in §2.6. Roughly, from an orbit equivalence coupling, a measure equivalence coupling can be constructed in such a way that both groups share a common fundamental domain. In particular, such a coupling is automatically cobounded. Hence Theorem 11 and Theorem 13 have the following immediate corollaries.

Corollary 16. *Let Γ be a finitely generated hyperbolic group. There is $p > 0$ such that if there exists a (L^p, L^∞) orbit equivalence coupling from Γ to a finitely generated group Λ , then Λ is also hyperbolic.*

Corollary 17. *Let Γ be a finitely generated hyperbolic group. For every $p > q > 0$, if there is a (φ, ψ) -integrable orbit equivalence coupling from Γ to any finitely generated group Λ where $\varphi(t) = \exp(t^p)$ and $\psi(t) = t^{1+1/q}$, then Λ is also hyperbolic.*

Hyperbolicity and embedded cycles. The proofs of Theorem 11 and Theorem 13 are based on the following seemingly new characterization of *non*-hyperbolic spaces which might be useful for future applications. We denote by C_n the standard Cayley graph of the cyclic group $\mathbb{Z}/n\mathbb{Z}$.

Theorem 18 (see Cor. 9.10). *Let (X, d) be a geodesic space that is not hyperbolic. Then there are arbitrarily large integers n and embeddings $\varphi_n: C_n \rightarrow (X, d)$ such that*

$$\frac{1}{17820}d_{C_n}(x, y) \leq d(\varphi_n(x), \varphi_n(y)) \leq d_{C_n}(x, y),$$

for every $x, y \in C_n$.

The strategy of proof consists in confronting this result with the following one (a very close statement is proved in [VS14] for the real hyperbolic space).

Theorem 19 (see Corollary 9.3). *Let $a \geq 0$, $b \geq 1$, and $\delta \geq 1$. There is an integer $n_0 = n_0(a, b) \geq 2$ such that the following holds. For all δ -hyperbolic geodesic space X , for all $n \geq n_0$, if there is a map $\varphi: C_n \rightarrow X$ such that for all $x, y \in C_n$*

$$ad_{C_n}(x, y) \leq d(\varphi(x), \varphi(y)) \leq bd_{C_n}(x, y)$$

then we have

$$a < 12\delta \cdot \frac{\log n}{n}. \tag{1}$$

Sketch of proof. We prove Theorem 11 and Theorem 13 by contradiction. Let us briefly sketch the argument for an orbit equivalence coupling. We assume that Λ is non hyperbolic and consider the map $\varphi_n: C_n \rightarrow \Lambda$ provided for some large integer n by Theorem 18. Identifying the orbits of Λ and Γ , we obtain for a.e. $x \in X$, a map $\psi_{n,x}: C_n \rightarrow \Gamma$. By exploiting the integrability condition from Λ to Γ , we obtain a bound on the Lipschitz constant of $\psi_{n,x}$, which is satisfied on a subset of X of sufficiently large measure. Observe that this step is trivial under the hypotheses of Theorem 11, as the L^∞ -condition ensures that $\psi_{n,x}$ is Lipschitz, uniformly with respect to $x \in X$. The more subtle part of the argument consists in estimating the Lipschitz constant of the inverse of $\psi_{n,x}$, in order to obtain a contradiction with Theorem 19.

In case of a measure equivalence coupling, a difficulty arises in the second step of the proof: in order to estimate the Lipschitz constant of the inverse of $\psi_{n,x}$, we need to go back to Λ and exploit the lower bound $\frac{1}{17820}$ on the Lipschitz constant of the inverse of φ_n . This where the coboundedness condition comes in, allowing us to relate the fundamental domains of the two groups.

Further remarks and questions. As commented above, the coboundedness assumption plays an important (though technical) role in the proofs. It would be interesting to know whether it can be avoided. In particular, this raises the following question.

Question 1.10. Assume $n \geq 3$, and let Γ (resp. Λ) be a uniform (resp. non-uniform) lattice in $SO(n, 1)$. Are Γ and Λ L^p measure equivalent for all $p < \infty$? Or at the opposite are they not L^p -measure equivalent as soon as $p > n - 1$ (or maybe even $p = n - 1$)?

Coboundedness was also considered by Sauer in his PhD thesis [Sau02]. He proved that two amenable finitely generated groups are quasi-isometric if and only if they admit an L^∞ coupling which is cobounded in both directions: namely there exist finite subsets $F_\Lambda \subset \Lambda$ and $F_\Gamma \subset \Gamma$ such that $X_\Gamma \subset F_\Lambda X_\Lambda$ and $X_\Lambda \subset F_\Gamma X_\Gamma$. These considerations lead us to the following maybe surprisingly open question.

Question 1.11. Is hyperbolicity invariant under L^∞ measure equivalence?

1.4 A sort of “asymmetric distance” between finitely generated groups

We may view the functions φ and ψ for which there exists a (φ, ψ) measurable coupling from a group Γ to a group Λ as an asymmetric way of quantifying how geometrically close these groups are from one another. This is of course only interesting within a class of groups that are measure equivalent, such as lattices (uniform and non-uniform ones) in a given locally compact group, or among amenable groups.

Let us illustrate this point of view in the following special case: given two finitely generated groups Γ and Λ , let $\alpha(\Gamma, \Lambda)$ be the infimum of $-\log p$ over all $p \leq 1$ such that there exists an (L^p, L^p) integrable measure equivalence coupling between Γ and Λ . It follows from Proposition 2.26 that α satisfies the triangle inequality. Therefore, it defines a pseudo-distance between (isomorphism classes of) finitely generated groups. Note that Theorem 6 yields $\alpha(\mathbb{Z}^4, \text{Heis}(\mathbb{Z})) = 0$, while Theorem 5 and the remark that precedes imply $\alpha(\mathbb{Z}^n, \mathbb{Z}^m) = |\log n - \log m|$.

It is interesting to note that for $p \geq 1$ (and more generally for convex functions φ), things behave differently, as admitting an (L^p, L^p) integrable measure equivalence coupling is an equivalence relation. This means that balls of radius 0 for α can be equipped with an ultrametric distance $\beta(\Gamma, \Lambda)$ defined for instance by the infimum of $1/p$ over all $p \geq 1$ such that the groups are L^p measure equivalent. In other words, we can distinguish two scales: a “large” scale measured by the pseudo distance α , and a finer scale, measured by the ultrametric pseudo-distance β .

More generally, we will see in §2.5 that if the functions φ and ψ are concave, then (φ, ψ) -integrable measurable equivalence couplings satisfy a natural composition rule, which we may view as analogous to a “triangle inequality”. Moreover, as for the case of L^p integrability, we shall see that for functions that grow faster than linear, e.g. $t \rightarrow t^p$ for $p \geq 1$, or $t \rightarrow \exp(t^a)$ for $a > 0$, one has to adopt a different point of view. Indeed, a stronger version of the integrability condition is required (which is automatic for L^p) in order for them to be well-behaved under composition. Actually under this stronger condition, we also get a stronger conclusion, which is analogous to an ultrametric property.

As a matter of fact, these composition rules are better stated in an even more asymmetric situation, where the measure coupling is no longer a measure *equivalence* coupling. If we drop on one side the finiteness of the measure of the fundamental domain, we obtain a natural notion of measured subgroup, while if we drop the freeness for one of the actions, we obtain a measurable notion of quotient. Combining them, we get a measurable notion of sub-quotient. In these situations, an integrability condition only makes sense in one direction, so that we obtain notions of φ -integrable measurable subgroups, quotients or sub-quotients, which we develop in the next section. To see how these notions can be useful, note that Theorem 1 is a corollary of the monotonicity of the isoperimetric profile under L^1 measure quotient, while Theorem 2 involves the notion of L^∞ measure subgroup.

1.5 Plan of the paper

In Section 2, we settle the general framework of this paper. More precisely, in §2.1 we introduce new notions of measured subgroups, quotients and sub-quotients that are defined via asymmetric couplings where the actions of both groups are smooth but where the freeness or the finite covolume assumption are dropped for one of the actions. Such notions satisfy natural composition rules that are gathered in Proposition 2.9. In §2.4, we make these notions quantitative by adding integrability conditions. The behavior of these integrability conditions under composition is studied in §2.5. Finally, §2.6 is dedicated to similar notions in the orbit equivalence setting.

In Section 3, we prove a general monotonicity result of the volume growth under measured sub-quotients satisfying certain integrability conditions. The proof is largely inspired (although different) from Bowen’s original proof of the invariance of volume growth under L^1 measure equivalence.

Section 4 focuses on two monotonicity results regarding the isoperimetric profile, namely Theorems 4.2 and 4.3, which generalize Theorem 1. In §4.1 a quick overview of their proofs is given. Sections 4.2 to 4.5 are dedicated to the actual proofs of these theorems. This section ends with §4.6 where we prove Corollary 4.

In Section 5, we show that the existence of a regular map from an amenable finitely generated group Γ to a finitely generated group Λ can be used to construct an L^∞ subgroup coupling from Γ to Λ . This allows us to deduce Theorem 2 from Theorem 4.2. We then exploit this result in §5.2 to establish Theorem 3.

In Section 6 we show how Følner tilings can be used to produce orbit equivalent pmp actions of groups with a control on the integrability properties of the corresponding cocycles. In §6.2 we give applications to groups with polynomial growth, and in §6.3 we treat the case of \mathbb{Z} and the Lamplighter group.

In Section 7.2, we study the behavior of orbit equivalence under wreath products, and we show that the integrability conditions pass through. This enables us to construct couplings between \mathbb{Z} and iterated wreath products with nearly sharp integrability conditions.

Section 8 deals with properties that are “very” unstable. In §8.1, we construct a very explicit exponential orbit equivalence coupling between $BS(1, n)$ and $\mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}$, proving that finite presentability is not preserved. §8.2, we construct another explicit coupling between two solvable groups, one with finite asymptotic dimension and the other not.

Finally Section 9 is dedicated to our rigidity results for hyperbolic groups. Theorem 18 and Theorem 19 are first proved in §9.1. We then exploit these geometric statements in §9.11 to prove our main result, namely Theorem 9.11 from which we deduce the theorems announced in the introduction.

Despite the fact that the whole paper relies on the definitions introduced in Section 2, each other section can be read independently. We therefore advise the reader not to refrain from going directly to the section that mostly interests her or him, keeping in mind that the basic notions and notations can be found in Section 2.

Acknowledgments. We thank Matthieu Joseph for many helpful conversations around this project. We also thank Yves Cornuier and Amandine Escalier for their useful remarks on a preliminary version of this work. We thank Lewis Bowen for sharing with us his unpublished note on the behavior of the Følner function under L^1 orbit equivalence. Finally we are grateful to the anonymous referee for his numerous remarks and corrections.

2 Variations on measure equivalence

Convention. Throughout the paper, we allow metrics and pseudo-metrics to take the value $+\infty$.

2.1 Smooth actions and fundamental domains

A **standard Borel space** $(\Omega, \mathcal{B}(\Omega))$ is a measurable space whose σ -algebra $\mathcal{B}(\Omega)$ consists of the Borel subsets coming from some Polish (separable and completely metrizable) topology on Ω . The

elements of $\mathcal{B}(\Omega)$ are called **Borel**, and so are called measurable maps between standard Borel spaces. A **standard measure space** is a standard Borel space $(\Omega, \mathcal{B}(\Omega))$ equipped with a nonzero σ -finite measure μ on $\mathcal{B}(\Omega)$. We simply denote such a space by (Ω, μ) . A Borel subset $\Omega_0 \subseteq \Omega$ is said to be **conull** or of **full measure** if $\mu(\Omega \setminus \Omega_0) = 0$ and a property that holds for all $\omega \in \Omega_0$ is said to hold **almost surely** or **for almost every** $\omega \in \Omega$. The space (Ω, μ) is said to be a standard probability space if $\mu(\Omega) = 1$.

A **measure-preserving action** of a discrete countable group Γ on (Ω, μ) , for short $\Gamma \curvearrowright (\Omega, \mu)$, is a Γ -action on Ω such that the action map $(\gamma, x) \mapsto \gamma \cdot x$ is Borel and that $\mu(\gamma \cdot E) = \mu(E)$ for all $\gamma \in \Gamma$ and all Borel $E \subseteq \Omega$. A measure-preserving action on a standard probability space is said to be probability measure-preserving.

Since the groups that we are dealing with are countable, if we are given $\Gamma \curvearrowright (\Omega, \mu)$, then every full measure Borel subset $A \subseteq \Omega$ contains a Γ -invariant full measure Borel subset, namely the set $\bigcap_{\gamma \in \Gamma} \gamma \cdot A$. Moreover, since any full measure Borel subset of a standard measure space is a standard measure space, there will be no harm in considering that some properties that hold almost everywhere actually hold everywhere, so we will often do so without explicitly restricting to a full measure Borel subset. For instance, one says that a measure-preserving Γ -action on (Ω, μ) is *free* if for all $\gamma \in \Gamma \setminus \{e_\Gamma\}$, we have $\gamma \cdot x \neq x$ for almost every $x \in \Omega$. We may as well assume that the latter holds for *every* $x \in \Omega$ since this becomes true once we restrict the action to a full measure Γ -invariant Borel set.

Finally, given $\Gamma \curvearrowright (\Omega, \mu)$, the **full pseudo-group** of the action $\Gamma \curvearrowright \Omega$ is the set of all partially defined Borel bijections $\varphi: A \rightarrow B$, where A and B are Borel subsets of Ω , such that for all $x \in A$, we have $\varphi(x) \in \Gamma \cdot x$. Every such element is measure-preserving, see [KM04, Prop. 2.1].

Definition 2.1. A **fundamental domain** for $\Gamma \curvearrowright (\Omega, \mu)$ is a Borel set $X_\Gamma \subseteq \Omega$ which intersects almost every Γ -orbit at exactly one point: there is a full measure Γ -invariant Borel set $\Omega_0 \subseteq \Omega$ such that for all $x \in \Omega_0$ we have that the intersection of the Γ -orbit of x with X_Γ is a singleton

Equivalently, a Borel set $X_\Gamma \subseteq \Omega$ is a fundamental domain if and only if the quotient map $\pi: X_\Gamma \rightarrow \Omega/\Gamma$ is a bijection. Note that since a fundamental domain intersects almost every orbit, the union of its translates has full measure, so every fundamental domain must have positive measure. Also note that since a fundamental domain for $\Gamma \curvearrowright (\Omega, \mu)$ intersects almost every orbit, the union of its translates has full measure, so every fundamental domain must have strictly positive measure whenever this is the case with Ω .

Moreover, after possibly passing to a Γ -invariant Borel subset of full measure, the existence of a Borel set $X_\Gamma \subseteq \Omega$ that intersects every Γ -orbit exactly once is equivalent to the fact that Ω/Γ is standard Borel, or in other words that the quotient map $\pi: X_\Gamma \rightarrow \Omega/\Gamma$ is a Borel bijection between standard Borel spaces. In this case, the orbit equivalence relation is said to be smooth, and we make the following definition.

Definition 2.2. A measure-preserving action of a countable group Γ on a standard measured space (Ω, μ) is **smooth** if it admits a fundamental domain.

Given a smooth action $\Gamma \curvearrowright (\Omega, \mu)$, if X is a fundamental domain for the Γ -action, we denote by π_X the map which takes (almost) every $\omega \in \Omega$ to the unique element of the Γ -orbit of ω which belongs to X . Observe that by definition, if X_1 and X_2 are two measure-fundamental domains, then the restriction of π_{X_1} to X_2 is an element of the full pseudo-group of the Γ -action whose inverse is the restriction of π_{X_2} to X_1 .

In particular, X_1 and X_2 have the same measure, and so given any smooth action $\Gamma \curvearrowright (\Omega, \mu)$, we can unambiguously endow the quotient space Ω/Γ with the measure obtained by identifying it with one of the fundamental domains. We will still denote this measure by μ .

Finally, given a fundamental domain X , we denote by ι_X the inverse of the projection map $X \rightarrow \Omega/\Gamma$.

Convention. We shall use the notation “ $\gamma * x$ ” instead of “ $\gamma \cdot x$ ” for smooth Γ -actions. This distinction will prove useful later on since we will also have induced actions on fundamental domains, which we will denote by \cdot .

2.2 Asymmetric couplings

We start by recalling the notion of measure equivalence coupling, introduced by Gromov.

Definition 2.3. Let Γ and Λ be countable groups, a **measure equivalence coupling** from Γ to Λ is a quadruple $(\Omega, X_\Gamma, X_\Lambda, \mu)$, where (Ω, μ) is a standard Borel measure space equipped with commuting measure-preserving smooth Γ and Λ actions such that

- (i) both actions are free;
- (ii) X_Λ (resp. X_Γ) is a fixed fundamental domain for the Λ -action (resp. for the Γ -action);
- (iii) X_Λ and X_Γ have finite measures.

When there exists such a coupling, we say that Γ and Λ are **measure equivalent**.

The most basic example of a measure equivalence coupling is the following. Given a countable group Λ viewed as a standard measured space equipped with the counting measure c , we take $\Gamma = \Lambda$ and we have a measure equivalence coupling between Γ and Λ by making Γ act on Λ by left translation, and Λ act on itself by right translation. A finite measure fundamental domain for both actions is given by $X_\Gamma = X_\Lambda = \{e\}$, so $(\Gamma, \{e_\Lambda\}, \{e_\Lambda\}, c)$ is a measure equivalence coupling between Γ and Λ . Motivated by this example, we see that if Γ was only an *infinite index subgroup* of Λ , then it cannot have a finite measure fundamental domain, although it has an infinite measure fundamental domain. We thus first relax the notion of measure equivalence coupling as the following asymmetric version.

Definition 2.4. Let Γ and Λ be countable groups. A **measure subgroup coupling** from Γ to Λ is a triple (Ω, X_Λ, μ) , where (Ω, μ) is a standard Borel measure space equipped with commuting measure-preserving smooth Γ and Λ actions such that

- (i) the Γ and Λ -actions are free;
- (ii) X_Λ is a fixed fundamental domain for the Λ -action;
- (iii) X_Λ has finite measure.

When there exists such a coupling, we say that Γ is a **measure subgroup** of Λ .

As explained above, the basic example of a measure subgroup coupling is obtained when $\Gamma \leq \Lambda$: it is the triple $(\Lambda, \{e_\Lambda\}, c)$ making Γ act by left-translation on Λ and Λ by right-translation on itself.

Remark 2.5. Being symmetric in Γ and Λ , measure equivalence couplings require the data of two fundamental domains X_Γ and X_Λ . By contrast, for the above asymmetric notion and its latter variants, only an explicit fundamental domain for the Λ -action is required. The latter will be crucial when making these notions quantitative (see §2.4).

Let us now tackle the case when Γ is a quotient of Λ . Here we can make Γ act on itself by left translation, and Λ act on Γ by right translation via the quotient map. Compared to measure subgroup couplings, we have lost the freeness of the Λ -action, but we gained that Γ has a finite measure fundamental domain, arriving at the following other asymmetric definition.

Definition 2.6. Let Γ and Λ be countable groups. A **measure quotient coupling** from Γ to Λ is a triple (Ω, X_Λ, μ) , where (Ω, μ) is a standard Borel measure space equipped with commuting measure-preserving smooth Γ and Λ actions such that

- (i) the Γ -action is free and admits a finite measure fundamental domain;
- (ii) X_Λ is a fixed fundamental domain for the Λ -action;
- (iii) X_Λ has finite measure.

When there exists such a coupling, we say that Γ is a **measure quotient** of Λ .

Finally, one can find a common denominator to measure subgroup couplings and measure quotient couplings as follows.

Definition 2.7. Let Γ and Λ be countable groups. A **measure sub-quotient coupling** from Γ to Λ is a triple (Ω, X_Λ, μ) , where (Ω, μ) is a standard Borel measure space equipped with commuting measure-preserving smooth Γ and Λ actions such that

- (i) the Γ action is free;
- (ii) X_Λ is a fixed fundamental domain for the Λ -action;
- (iii) X_Λ has finite measure.

When there exists such a coupling, we say that Γ is a **measure sub-quotient** of Λ .

It is a good exercise for the reader to describe how such a coupling arises when Γ is a subgroup of a quotient of Λ .

Remark 2.8. Every measure equivalence coupling yields two measure sub-quotient couplings (one from Γ to Λ , the other one from Λ to Γ) which are both subgroup and quotient couplings.

To lighten the presentation, and since there will be no ambiguity, we shall frequently remove the word “measure” in front of the word “coupling” from there on. Let us end this section by pointing a few nontrivial examples of couplings:

- An important instance of subgroup coupling arises via regular embeddings, using amenability. It will be described in §5.
- Other examples arising from the notion of orbit equivalence will be given in §2.6.
- Another way to produce more sophisticated examples of couplings is to compose them, as we will see now.

2.3 Composition of asymmetric measure couplings

Our goal here is to extend the (classical) notion of composition of measure equivalence couplings to our asymmetric setting (see for instance [BFS13] for compositions of measure equivalence couplings). Starting from a triplet of groups Γ , Λ and Σ , a coupling from Γ to Λ and a coupling from Λ to Σ , we want to define a coupling from Γ to Σ . Of course we want such a composition rule should be coherent with those holding for usual notions of subgroups, quotients... For instance we will get that a measure subgroup of a measure subgroup is again a measure subgroup. Similarly, we will obtain that a measure subgroup of a measure quotient is a measure sub-quotient (see Proposition 2.9). However, a difficulty arises when considering the composition of two measure quotients: it can only be defined at the cost of losing the smoothness of the action of the target group. On the other hand, the condition that both actions are smooth plays a crucial role in our proof of Theorem 1, which is the reason why we kept it in the definition.

Composition of couplings. Let Γ , Λ and Σ be three countable groups let $(\Omega_1, X_{1,\Lambda}, \mu_1)$ be a subgroup coupling from Γ to Λ and let $(\Omega_2, X_{2,\Sigma}, \mu_2)$ be a sub-quotient coupling from Λ to Σ . The **composition** of these two couplings $(\Omega_3, X_{3,\Sigma}, \mu_3)$ is the sub-quotient coupling from Γ to Σ obtained as follows.

- We consider the diagonal action $\Lambda \curvearrowright (\Omega_1 \times \Omega_2, \mu_1 \otimes \mu_2)$ which is smooth and commutes with the Γ -action on the first coordinate and the Λ -action on the second coordinate.

- Then the measured space of the coupling is $\Omega_3 := (\Omega_1 \times \Omega_2)/\Lambda$ equipped with the induced Γ and Λ actions.
- Identifying Ω_3 with a Λ -fundamental domain of $\Omega_1 \times \Omega_2$, we define the measure μ_3 as the restriction of the product measure.
- We let $X_{3,\Sigma} = \pi_{\Omega_3}(X_{1,\Lambda} \times X_{2,\Sigma})$, where π_{Ω_3} is the projection $\Omega_1 \times \Omega_2 \rightarrow \Omega_3$.
- Furthermore, letting $X_{1,\Gamma}$ (resp. $X_{2,\Lambda}$) be fundamental domains for the action of Γ on Ω_1 (resp. for the action of Λ on Ω_2), the subset $X_{3,\Gamma} = \pi_{\Omega_3}(X_{1,\Gamma} \times X_{2,\Lambda})$ of Ω_3 defines a fundamental domain for the action of Γ .

Proposition 2.9. *Let Γ , Λ and Σ be three countable groups, let $(\Omega_1, X_{1,\Lambda}, \mu_1)$ be a subgroup coupling from Γ to Λ and let $(\Omega_2, X_{2,\Sigma}, \mu_2)$ be a sub-quotient coupling from Λ to Σ . The composition of these two couplings is a sub-quotient coupling from Γ to Σ .*

If both couplings are subgroup (resp. equivalence) couplings, then their composition is also a subgroup (resp. equivalence) coupling.

Finally if the coupling from Γ to Λ is a measure equivalence coupling and the coupling from Λ to Σ is a quotient coupling, then their composition is a quotient coupling.

Before proceeding to the proof, we need to introduce some notation. Given a sub-quotient coupling (Ω, X_Λ, μ) from Γ to Λ , we denote their two commuting actions by $*$. We also have a natural action of Γ on Ω/Λ which we call the **induced action** and denote by \cdot . Through the natural identification of Ω/Λ to X_Λ , this induced action can be described as follows: for all $x \in X_\Lambda$, $\gamma \in \Gamma$,

$$\{\gamma \cdot x\} = \Lambda * \gamma * x \cap X_\Lambda.$$

Note that the induced Γ -action defines elements of the pseudo full group of the Λ -action on Ω , so it is a measure-preserving action.

Proof of Proposition 2.9. First, the diagonal Λ -action on $\Omega_1 \times \Omega_2$ is smooth because $X_{1,\Lambda} \times \Omega_2$ is a fundamental domain for it. We then define $\Omega_3 := (\Omega_1 \times \Omega_2)/\Lambda$, and we denote by μ_3 the measure induced on Ω_3 from the product measure on $\Omega_1 \times \Omega_2$. Finally, we denote by \star the induced Γ -action on Ω_3 . Through the identification of Ω_3 with the fundamental domain $X_{1,\Lambda} \times \Omega_2$ we see that the induced Σ -action is given by:

$$\sigma \star (x, \omega) = (x, \sigma * \omega),$$

for all $(x, \omega) \in X_{1,\Lambda} \times \Omega_2$ and all $\sigma \in \Sigma$. In particular, we see that $X_{3,\Sigma} := \pi_{\Omega_3}(X_{1,\Lambda} \times X_{2,\Sigma})$ is a fundamental domain for it. If the Σ -action on Ω_2 is free then so is the Σ -action on Ω_3 . Note that $X_{3,\Sigma}$ has finite measure because $X_{2,\Sigma}$ does. Finally, if we fix a fundamental domain $X_{2,\Lambda}$ for the Λ -action on Ω_2 , then we also have a natural identification of Ω_3 with $\Omega_1 \times X_{2,\Lambda}$. This shows that the Γ -action on Ω_3 is free.

All the properties stated in the proposition can now directly be inferred from the two previous paragraphs. \square

Remark 2.10. Keeping the notation of the proof, we can also describe the Γ -action on Ω_3 when the latter is identified with $X_{1,\Lambda} \times \Omega_2$ as follows:

$$\gamma \star (x, \omega) = (\gamma \cdot x, \alpha(\gamma, x) * \omega),$$

where \cdot denotes the induced Γ -action on $X_{1,\Lambda}$, and where $\alpha: \Gamma \times X_{1,\Lambda} \rightarrow \Lambda$ is the cocycle given by the equation $\alpha(\gamma, x) * \gamma * x = \gamma \cdot x$ (we shall come back to this important notion in the next section). Note that the induced Γ -action (denoted by \bullet) on the fundamental domain $X_{3,\Sigma}$, identified to $X_{1,\Lambda} \times X_{2,\Sigma}$, is then given by:

$$\gamma \bullet (x, y) = (\gamma \cdot x, \alpha(\gamma, x) \cdot y).$$

This point of view will be useful when we explore how integrability conditions behave under composition of couplings.

2.4 Integrability conditions

We now come to the central notion studied in this paper. Our goal is to add quantitative constraints on a coupling that extend the well-known L^p -integrability condition for measure equivalence couplings. Classically (for measure equivalence couplings), this is done via the notion of cocycle. Given a subgroup coupling (Ω, X_Λ, μ) from Γ to Λ , we can define the cocycle $\alpha: \Gamma \times X_\Lambda \rightarrow \Lambda$ via by the equation

$$\alpha(\gamma, x) * \gamma * x = \gamma \cdot x.$$

It is non-ambiguously defined for a.e. $x \in X_\Lambda$, as we assume that the Λ -action is free.

Definition 2.11. Given any non-decreasing map $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and a finite generating subset $S_\Lambda \subset \Lambda$, we say that the subgroup coupling (Ω, X_Λ, μ) is φ -integrable if for all $\gamma \in \Gamma$ there is $c_\gamma > 0$ such that

$$\int_{X_\Lambda} \varphi \left(\frac{|\alpha(\gamma, x)|_{S_\Lambda}}{c_\gamma} \right) d\mu(x) < +\infty.$$

This definition is very convenient but only makes sense when the Λ action is free. So we need to find a substitute for measure quotients or measure sub-quotients. To that purpose, it will be useful to work with Schreier graph metrics.

Definition 2.12. Let Λ be a finitely generated group acting smoothly on a standard measured space (Ω, μ) . We denote by d_{S_Λ} the **Schreier graph metric** on the Λ -orbits, namely for $y \in \Lambda * x$, we let

$$d_{S_\Lambda}(x, y) = \min\{n \in \mathbb{N} : \exists s_1, \dots, s_n \in S_\Lambda, y = s_1 \cdots s_n * x\}.$$

Observe that if we are given $\Lambda \curvearrowright (\Omega, \mu)$ and two finite generating sets S_1 and S_2 for Λ , then there is $C > 0$ such that for all $x \in \Omega$ and all $y \in \Lambda * x$,

$$\frac{1}{C} d_{S_1}(x, y) \leq d_{S_2}(x, y) \leq C d_{S_1}(x, y). \quad (2)$$

We recall that given a smooth action $\Lambda \curvearrowright (\Omega, \mu)$ and a measure fundamental domain X , the map ι_X is the inverse of the (bijective) projection $\pi_{\Omega/\Lambda}: X \rightarrow \Omega/\Lambda$.

Definition 2.13. Given any non-decreasing map $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and two fundamental domains X_1 and X_2 of a smooth Λ -action on (Ω, μ) , we say that they are **φ -equivalent** if there is $c > 0$ such that

$$\int_{\Omega/\Lambda} \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))}{c} \right) d\mu(x) < +\infty.$$

Note that this does not depend on the choice of the symmetric generating set S_Λ by virtue of inequality (2).

Remark 2.14. If φ satisfies that for every $c > 0$, there is a constant $C > 0$ such that for all $x \geq 0$, $\varphi(cx) \leq C\varphi(x)$, then X_1 and X_2 are φ -equivalent if and only if

$$\int_{\Omega/\Lambda} \varphi(d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))) d\mu(x) < +\infty,$$

which is then also equivalent to: for every $c > 0$ we have

$$\int_{\Omega/\Lambda} \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))}{c} \right) d\mu(x) < +\infty.$$

This is the case if $\varphi(x) = x^p$ for some $p > 0$, or more generally if φ is subadditive. An example where this is not true is when φ is the exponential map.

In order to show that φ -equivalence is indeed an equivalence relation, we introduce the following quantity: given a smooth Λ -action and two fundamental domains X_1 and X_2 , we let

$$c_{\varphi, S_\Lambda}(X_1, X_2) = \inf \left\{ c > 0: \int_{\Omega/\Lambda} \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))}{c} \right) d\mu(x) < +\infty \right\}.$$

Proposition 2.15. *The map c_{φ, S_Λ} is a pseudo-metric on the set of fundamental domains (recall that pseudo-metrics are allowed to take the value $+\infty$).*

The proof relies on the following elementary observation.

Lemma 2.16. *Let $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a non-decreasing function, let $a, b \geq 0$ and $c, d > 0$. Then*

$$\varphi \left(\frac{a+b}{c+d} \right) \leq \varphi \left(\frac{a}{c} \right) + \varphi \left(\frac{b}{d} \right).$$

Proof. By symmetry, we may as well assume that $\frac{a}{c} \geq \frac{b}{d}$, in which case $\frac{a}{c} \geq \frac{a+b}{c+d}$, so we have $\varphi \left(\frac{a+b}{c+d} \right) \leq \varphi \left(\frac{a}{c} \right) \leq \varphi \left(\frac{a}{c} \right) + \varphi \left(\frac{b}{d} \right)$. \square

Proof of Proposition 2.15. The map c_{φ, S_Λ} is clearly symmetric and satisfies $c_{\varphi, S_\Lambda}(X, X) = 0$ for every fundamental domain X , so we only need to check that it satisfies the triangle inequality.

To this end, let X_1, X_2 and X_3 be fundamental domains, let $c_1 > c_{\varphi, S_\Lambda}(X_1, X_2)$ and $c_2 > c_{\varphi, S_\Lambda}(X_2, X_3)$. We have for every $x \in \Omega/\Lambda$ that

$$\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_3}(x))}{c_1 + c_2} \leq \frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x)) + d_{S_\Lambda}(\iota_{X_2}(x), \iota_{X_3}(x))}{c_1 + c_2}.$$

By the previous lemma, we thus have

$$\varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_3}(x))}{c_1 + c_2} \right) \leq \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))}{c_1} \right) + \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_2}(x), \iota_{X_3}(x))}{c_2} \right).$$

By integrating and using our assumptions on c_1 and c_2 , we then deduce that

$$\int_{\Omega/\Lambda} \varphi \left(\frac{d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_3}(x))}{c_1 + c_2} \right) d\mu(x) < +\infty,$$

and so $c_{\varphi, S_\Lambda}(X_1, X_3) \leq c_{\varphi, S_\Lambda}(X_1, X_2) + c_{\varphi, S_\Lambda}(X_2, X_3)$ as wanted. \square

Corollary 2.17. *The notion of φ -equivalence is an equivalence relation between fundamental domains.*

We are now ready to define our notion of φ -integrability for sub-quotient couplings.

Definition 2.18. Let $\varphi, \psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be non-decreasing maps.

- A sub-quotient (resp. subgroup, quotient) coupling (Ω, X_Λ, μ) from Γ to Λ is called **φ -integrable** if for every $\gamma \in \Gamma$ we have that X_Λ and $\gamma * X_\Lambda$ are φ -equivalent as fundamental domains of the Λ -action. In that case, we call Γ a **φ -integrable measure sub-quotient** (resp. **subgroup or quotient**) of Λ .
- A measure equivalence coupling $(\Omega, X_\Gamma, X_\Lambda, \mu)$ from Γ to Λ is called **(φ, ψ) -integrable** when the coupling (Ω, X_Λ, μ) from Γ to Λ is φ -integrable and the coupling (Ω, X_Γ, μ) from Λ to Γ is ψ -integrable. We then say that Γ and Λ are **(φ, ψ) -integrably measure equivalent**, or simply that they are (φ, ψ) -measure equivalent.

Remark 2.19. By spelling out what φ -integrability means, we see that a measure sub-quotient coupling (Ω, X_Λ, μ) from Γ to Λ is φ -integrable if and only if for every $\gamma \in \Gamma$, there is $c_\gamma > 0$ such that

$$\int_{X_\Lambda} \varphi \left(\frac{d_{S_\Lambda}(\gamma \cdot x, \gamma * x)}{c_\gamma} \right) d\mu(x) < +\infty,$$

where $*$ denotes the Γ -action on Ω and \cdot denotes the induced Γ -action on X_Λ . We deduce that a measure subgroup coupling from Γ to Λ is a φ -coupling if and only if the cocycle $\alpha: \Gamma \times X_\Lambda \rightarrow \Lambda$ is φ -integrable in the sense of Definition 2.11.

Proposition 2.20. *A sub-quotient coupling (Ω, X_Λ, μ) from $\Gamma = \langle S_\Gamma \rangle$ to Λ is φ -integrable if and only if for every $s \in S_\Gamma$ we have that X_Λ and $s * X_\Lambda$ are φ -equivalent.*

Proof. Since the Γ -action commutes with the Λ -action, it must preserve the equivalence relation of φ -equivalence, so for every $\gamma \in \Gamma$ and every $s \in S_\Gamma$ we have that $\gamma * X_\Lambda$ and $\gamma s * X_\Lambda$ are φ -equivalent. From there, the statement follows by induction from Proposition 2.15. \square

Remark 2.21. It is not hard to see that being a φ -integrable sub-quotient coupling only depends on the asymptotic behavior of φ . More precisely, if $\varphi \preceq \psi$, then every ψ -integrable sub-quotient coupling is also a φ -integrable.

For convenience and consistency with the literature, for $p \in (0, \infty)$ we talk about L^p couplings instead of $x \mapsto x^p$ -integrable couplings. We also say that we have an **L^∞ sub-quotient coupling** from Γ to Λ when the Γ -action satisfies for every $\gamma \in \Gamma$ that the map $\Omega/\Lambda \rightarrow \Lambda: x \mapsto d_{S_\Lambda}(\gamma \cdot x, \gamma * x)$ is essentially bounded. Note that every L^∞ sub-quotient coupling is φ -integrable for any increasing map $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$.

The following definition plays an important role in our study of hyperbolicity.

Definition 2.22. A measure equivalence coupling $(\Omega, X_\Gamma, X_\Lambda, \mu)$ from Γ to Λ is called **cobounded** when there is a finite subset $F \subseteq \Lambda$ such that $X_\Gamma \subseteq FX_\Lambda$.

Example 2.23. We will see in Section 2.6 that every orbit equivalence coupling is cobounded.

Let us now explain how various established notions fit into our asymmetric framework.

- Two finitely generated groups Γ and Λ are *L^p measure equivalent* in the sense of [BFS13] when there is an (L^p, L^p) measure equivalence coupling from Γ to Λ .
- Two finitely generated groups Γ and Λ are *uniform measure equivalent* in the sense of [Sha04] when there is an (L^∞, L^∞) measure equivalence coupling between Γ and Λ .
- Two finitely generated groups Γ and Λ are *bounded measure equivalent* in the sense of [Sau02] when there is an (L^∞, L^∞) measure equivalence coupling from Γ to Λ which is cobounded in both directions.

2.5 Composition of φ -integrable couplings

We study how integrability conditions behave under composition of couplings. We first consider the case where $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is subadditive, e.g. when φ is concave. Our arguments will follow closely those from [BFS13, Sec. A.2].

Lemma 2.24. *Let Γ and Λ be two finitely generated groups, let $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a non-decreasing subadditive map and let (Ω, X_Λ, μ) be a φ -integrable sub-quotient coupling from Γ to Λ . Then there is a constant $C > 0$ such that for every $\gamma \in \Gamma$*

$$\int_{X_\Lambda} \varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x) \leq C |\gamma|_{S_\Gamma}.$$

Proof. Assume (Ω, X_Λ, μ) is a φ -integrable sub-quotient coupling from Γ to Λ . Given two fundamental domains X_1 and X_2 for the Λ -action, we define their φ -distance $d_{\varphi, S_\Lambda}(X_1, X_2)$ by

$$d_{\varphi, S_\Lambda}(X_1, X_2) = \int_{\Omega/\Lambda} \varphi(d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x))) d\mu(x)$$

Note that this distance is symmetric and satisfies the triangle inequality, so it is a pseudometric as soon as $\varphi(0) = 0$. Moreover, using Remark 2.14, we have for every $\gamma \in \Gamma$ that

$$d_{\varphi, S_\Lambda}(X_\Lambda, \gamma * X_\Lambda) < +\infty,$$

while by Remark 2.19 we have $d_{\varphi, S_\Lambda}(X_\Lambda, \gamma * X_\Lambda) = \int_{X_\Lambda} \varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x)$.

Let $C = \max_{\gamma \in S_\Gamma} d_{\varphi, S_\Lambda}(X_\Lambda, \gamma * X_\Lambda)$. The Γ -action on the set of Λ -fundamental domains preserves d_{φ, S_Λ} because it commutes with the Λ -action, so for every $s_1, \dots, s_n \in S_\Gamma$ we have by the triangle inequality

$$\begin{aligned} d_{\varphi, S_\Lambda}(s_1 \cdots s_n * X_\Lambda, X_\Lambda) &\leq \sum_{i=1}^n d_{\varphi, S_\Lambda}(s_1 \cdots s_i * X_\Lambda, s_1 \cdots s_{i-1} * X_\Lambda) \\ &\leq \sum_{i=1}^n d_{\varphi, S_\Lambda}(s_i * X_\Lambda, X_\Lambda) \\ &\leq Cn, \end{aligned}$$

which yields the desired result. \square

Remark 2.25. Note that when φ is subadditive and non-decreasing, we always have $\varphi(x) \leq x\varphi(1) + \varphi(1)$, in particular every L^1 sub-quotient coupling is φ -integrable.

Let us now study how couplings compose in the subadditive regime.

Proposition 2.26. *Let $\varphi, \psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be non-decreasing subadditive maps with φ moreover concave and let Γ, Λ and Σ be three finitely generated groups. Let $(\Omega_1, X_{1,\Lambda}, \mu_1)$ be a φ -integrable subgroup coupling from Γ to Λ and let $(\Omega_2, X_{2,\Sigma}, \mu_2)$ be a ψ -integrable sub-quotient coupling from Λ to Σ . Then the composition of these two couplings is a $\varphi \circ \psi$ -integrable sub-quotient coupling from Γ to Σ .*

Proof. Thanks to Lemma 2.24 we find $C > 0$ such that

$$\int_{X_{2,\Sigma}} \psi(d_{S_\Sigma}(\lambda \cdot x, \lambda * x)) d\mu(x) \leq C|\lambda|_{S_\Lambda} \text{ for every } \lambda \in \Lambda. \quad (3)$$

By scaling the measure μ_2 we may assume that $\mu_2(X_{2,\Sigma}) = 1$. Denote by $\alpha: \Gamma \times X_{1,\Lambda} \rightarrow \Lambda$ the cocycle defined by the equation $\alpha(\gamma, x) * \gamma * x = \gamma \cdot x$. By Remark 2.10 and the definition of the composition of our two couplings, we need to show that the following quantity is finite:

$$\int_{X_{1,\Lambda}} \int_{X_{2,\Sigma}} \varphi \circ \psi(d_{S_\Sigma}(\alpha(\gamma, x) \cdot y, \alpha(\gamma, x) * y)) d\mu_2(y) d\mu_1(x).$$

Now by Jensen's inequality, this is at most

$$\int_{X_{1,\Lambda}} \varphi \left(\int_{X_{2,\Sigma}} \psi(d_{S_\Sigma}(\alpha(\gamma, x) \cdot y, \alpha(\gamma, x) * y)) d\mu_2(y) \right) d\mu_1(x),$$

which by inequality (3) is bounded above by $\int_{X_{1,\Lambda}} \varphi(C|\alpha(\gamma, x)|_{S_\Lambda}) d\mu_1(x)$. The latter is indeed finite by our assumption on the first coupling and Remark 2.14. \square

The above result can be combined with Proposition 2.9 to obtain a $\varphi \circ \psi$ -integrable subgroup coupling or a $\varphi \circ \psi$ -integrable quotient coupling by composition.

Remark 2.27. Given two finitely generated groups Γ and Λ , one could define

$$\alpha(\Gamma, \Lambda) = -\log(\sup\{p \leq 1 : \Gamma \text{ and } \Lambda \text{ are } L^p \text{ measure equivalent}\})$$

The previous proposition implies that this is a pseudo-metric on the space of isomorphism classes of finitely generated groups. It would be interesting to understand this pseudo-metric further. For instance, Theorem 6.10 implies that $d(\mathbb{Z}^n, \mathbb{Z}^m) = |\log n - \log m|$.

For non subadditive maps φ , we need a stronger notion of φ -integrability so that it behaves well with respect to composition.

Definition 2.28. Let $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be an increasing map. We say that a coupling (Ω, X_Λ, μ) from Γ to Λ is **strongly φ -integrable** or φ^\diamond -integrable if for every $\varepsilon > 0$ there are $\delta > 0$ and $C > 0$ such that for every $\gamma \in \Gamma$,

$$\int_{X_\Lambda} \varphi(\delta d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x) \leq C\varphi(\varepsilon|\gamma|_{S_\Gamma})$$

Note again that thanks to inequality (2) strong integrability does not depend on the choice of the finite generating set S_Γ . However, the above condition has to be checked on *every* element of Γ .

Proposition 2.29. Let $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be an increasing map and let Γ, Λ and Σ be three finitely generated groups. Let $(\Omega_1, X_{1,\Lambda}, \mu_1)$ be a strongly φ -integrable subgroup coupling from Γ to Λ and let $(\Omega_2, X_{2,\Sigma}, \mu_2)$ be a strongly φ -integrable sub-quotient coupling from Λ to Σ . Then the composition of these two couplings is a strongly φ -integrable sub-quotient coupling from Γ to Σ .

Proof. Let $\varepsilon > 0$. As $(\Omega_1, X_{1,\Lambda}, \mu_1)$ is strongly φ -integrable, there are $\delta_\Lambda > 0$ and $C_\Gamma > 0$ such that

$$\int_{X_{1,\Lambda}} \varphi(\delta_\Lambda d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu_1(x) \leq C_\Gamma \varphi(\varepsilon|\gamma|_{S_\Gamma}) \text{ for every } \gamma \in \Gamma. \quad (4)$$

and as $(\Omega_2, X_{2,\Sigma}, \mu)$ is strongly φ -integrable, there exist $\delta_\Sigma > 0$ and $C_\Lambda > 0$ such that

$$\int_{X_{2,\Sigma}} \varphi(\delta_\Sigma d_{S_\Sigma}(\lambda \cdot x, \lambda * x)) d\mu_2(x) \leq C_\Lambda \varphi(\delta_\Lambda |\lambda|) \text{ for every } \lambda \in \Lambda. \quad (5)$$

By Remark 2.10 and the definition of the composition of our two couplings, we need to estimate the following quantity:

$$\int_{X_{1,\Lambda}} \int_{X_{2,\Sigma}} \varphi(\delta_\Sigma d_{S_\Sigma}(\alpha(\gamma, x) \cdot y, \alpha(\gamma, x) * y)) d\mu_2(y) d\mu_1(x).$$

By inequality (5), this is bounded above by

$$\int_{X_{1,\Lambda}} C_\Lambda \varphi(\delta_\Lambda |\alpha(\gamma, x)|_{S_\Lambda}) \leq C_\Lambda C_\Gamma \varphi(\varepsilon|\gamma|_{S_\Gamma})$$

as wanted, where the last inequality is a consequence of inequality (4), and the fact that by definition $|\alpha(\gamma, x)|_{S_\Lambda} = d_{S_\Lambda}(\gamma \cdot x, \gamma * x)$. \square

For some maps φ we can weaken the strong integrability condition. Most notably for L^p couplings, where $p \geq 1$.

Proposition 2.30. Let $p \geq 1$. Every L^p sub-quotient coupling from Γ to Λ is actually a strongly L^p sub-quotient coupling.

Proof. We follow an approach similar to that of Lemma 2.24. For two fundamental domains X_1 and X_2 for the Λ -action, we define their L^p distance by

$$d_{L^p, S_\Lambda}(X_1, X_2) = \left(\int_{\Omega/\Lambda} (d_{S_\Lambda}(\iota_{X_1}(x), \iota_{X_2}(x)))^p d\mu(x) \right)^{1/p}$$

It is not hard to check that d_{L^p, S_Λ} is a metric, and since the Γ -action commutes with the Λ -action, the group Γ acts on the set of Λ -fundamental domains by isometries. Let X_Λ be one of them. As in the proof of Lemma 2.24, we find $C_\Gamma > 0$ such that for every $\gamma \in \Gamma$, $d_{L^p, S_\Lambda}(X_\Lambda, \gamma * X_\Lambda) \leq C_\Gamma |\gamma|_{S_\Gamma}$. This means that for all $\gamma \in \Gamma$, we have

$$\int_{X_\Lambda} d_{S_\Lambda}(\gamma \cdot x, \gamma * x)^p d\mu(x) \leq C_\Gamma^p |\gamma|_{S_\Gamma}^p,$$

from which the result easily follows: given $\varepsilon > 0$, we take $\delta = \varepsilon$, $C = C_\Gamma^p$ and note that

$$\int_{X_\Lambda} (\delta d_{S_\Lambda}(\gamma \cdot x, \gamma * x))^p d\mu(x) = \delta^p \int_{X_\Lambda} d_{S_\Lambda}(\gamma \cdot x, \gamma * x)^p d\mu(x) \leq C_\Gamma^p (\varepsilon |\gamma|_{S_\Gamma})^p$$

as wanted. \square

Say for $p \geq 1$ that two finitely generated groups are L^p **measure equivalent** when there is an (L^p, L^p) measure equivalence coupling from Γ to Λ . We deduce from the two previous results that L^p measure equivalence is an equivalence relation between finitely generated groups, as proven by Bader, Furman and Sauer in [BFS13]. Note however that this is not true anymore for $p < 1$; counter-examples are provided by Theorem 6.10.

For exponentially integrable couplings, the next proposition states that it is enough to find a single ε so as to witness strong exponential integrability.

Proposition 2.31. *Let Γ and Λ be two finitely generated groups and let (Ω, X_Λ, μ) be a sub-quotient coupling from Γ to Λ . If there are $\varepsilon' > 0$, $\delta' > 0$ and $C' > 0$ such that*

$$\int_{X_\Lambda} \exp(\delta' d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x) \leq C' \exp(\varepsilon' |\gamma|_{S_\Gamma}) \text{ for every } \gamma \in \Gamma,$$

then the coupling is strongly exp-integrable.

Proof. Let $\varepsilon > 0$. If $\varepsilon \geq \varepsilon'$, we take $C = C'$ and $\delta = \delta'$ and check that Definition 2.28 holds with these constants. If $\varepsilon < \varepsilon'$, then take $C = C' \frac{\varepsilon}{\varepsilon'}$ and $\delta = \frac{\delta' \varepsilon}{\varepsilon'}$. Then we have for every $\gamma \in \Gamma$ that

$$\begin{aligned} \int_{X_\Lambda} \exp(\delta d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x) &= \int_{X_\Lambda} \exp(\delta' d_{S_\Lambda}(\gamma \cdot x, \gamma * x))^{\frac{\varepsilon}{\varepsilon'}} d\mu(x) \\ &\leq \left(\int_{X_\Lambda} \exp(\delta' d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) d\mu(x) \right)^{\frac{\varepsilon}{\varepsilon'}} \\ &\leq (C' \exp(\varepsilon' |\gamma|))^{\frac{\varepsilon}{\varepsilon'}} \\ &= C \exp(\varepsilon |\gamma|). \end{aligned}$$

Thus, the coupling is strongly exp-integrable as wanted. \square

Interesting examples of strongly (exp, exp)-integrable measure equivalence couplings will be provided in Section 8.

2.6 Variations on orbit equivalence

We now turn our attention to orbit equivalence of groups, which implies measure equivalence. To make the connection, we need to introduce measure-preserving equivalence relations (see [KM04] for details).

Definition 2.32. Given a measure-preserving action of a countable group Λ on a standard probability space (X, μ) , we associate to it a **measure-preserving equivalence relation** \mathcal{R}_Λ defined by

$$\mathcal{R}_\Lambda := \{(x, \lambda \cdot x) : x \in X, \lambda \in \Lambda\}.$$

A key property of measure-preserving equivalence relations is that they can be endowed with a natural σ -finite measure M . Namely, for \mathcal{R} a measure-preserving equivalence relation, and A a Borel subset of \mathcal{R} , we let

$$M(A) = \int_X |A_x| d\mu(x),$$

where $A_x = \{y \in X : (x, y) \in A\}$. Such a measure is invariant under the flip map $(x, y) \mapsto (y, x)$ (see [KM04, p. 34]), which means that

$$M(A) = \int_X |A^y| d\mu(y),$$

where $A^y = \{x \in X : (x, y) \in A\}$.

Definition 2.33. Let Γ and Λ be two finitely generated groups.

- An **orbit sub-quotient coupling** from Γ to Λ is a triple (X, Y, μ) where
 - (i) (Y, μ) is a standard σ -finite space equipped with a measure-preserving Λ -action,
 - (ii) X is a Borel subset of Y of measure 1 equipped with a free measure-preserving Γ -action;
 - (iii) and finally for almost every $x \in X$ we have that $\Gamma \cdot x \subseteq \Lambda \cdot x$.
- If the Λ -action is also free, we say that (X, Y, μ) is an **orbit subgroup coupling** of Γ with Λ . If $X = Y$ and for almost every $x \in X$ we have that $\Gamma \cdot x = \Lambda \cdot x$, we say that (X, Y, μ) is an **orbit quotient coupling**, which we then simply write as (X, μ) . Finally, (X, μ) is an **orbit equivalence coupling** of Γ with Λ if it is both an orbit quotient and an orbit subgroup coupling.

Remark 2.34. To every orbit sub-quotient coupling is naturally associated a measure sub-quotient coupling. We will see later that this statement admits some kind of converse (see Proposition 2.43).

- The coupling space is $\Omega := \mathcal{R}_\Lambda \cap (X \times Y)$ equipped with the measure induced by M .
- The commuting actions are defined as follows: for every $\gamma \in \Gamma$, $\lambda \in \Lambda$ and every $(x, y) \in \mathcal{R}_\Lambda$,

$$\gamma * (x, y) = (\gamma \cdot x, y) \text{ and } \lambda * (x, y) = (x, \lambda \cdot y).$$

- The chosen Λ -fundamental domain is the diagonal: $X_\Lambda = \{(x, x) : x \in X\}$.
- The Γ -action is smooth. Indeed, a Borel fundamental domain can be obtained as the intersection with $X \times Y$ of a disjoint union of graphs of *Borel choice functions* for the subequivalence relation $(\mathcal{R}_\Gamma \cup \Delta_Y) \leq \mathcal{R}_\Lambda$ (see [IKT09, Sec. 2.(A)]).

Note furthermore that (X_Λ, M) is naturally isomorphic to (X, μ) via $x \mapsto (x, x)$, and that the induced action on X_Λ is conjugate via the inverse of this map to the original action on (X, μ) .

Definition 2.35. An orbit sub-quotient coupling from Γ to Λ is φ -integrable when it is φ -integrable as a measure sub-quotient coupling.

Note that φ -integrability for an orbit sub-quotient coupling as above means that for all $\gamma \in \Gamma$, there is $c_\gamma > 0$ such that

$$\int_X \varphi \left(\frac{d_{S_\Lambda}(x, \gamma \cdot x)}{c_\gamma} \right) d\mu(x) < +\infty.$$

Remark 2.36. When $\varphi(x) = x^p$ for some $p \geq 1$, this means that Γ is contained in the L^p full group of the Λ -action, as defined in [LM18].

Remark 2.37. We note that admitting an orbit equivalence coupling agrees with the definitions of orbit equivalence between countable groups in the literature.

Remark 2.38. An orbit sub-quotient coupling (X, Y, μ) from Γ to Λ such that for almost every $x \in X$ we have that $\Gamma \cdot x = \Lambda \cdot x$ is an orbit quotient coupling.

Remark 2.39. Every orbit equivalence (resp. subgroup, quotient) coupling is a measure equivalence (resp. subgroup, quotient) coupling. We can thus define (φ, ψ) -integrable orbit equivalence couplings as well as φ -integrable sub-quotient (resp. subgroup, quotient) couplings. Finally, we can also define strong φ -integrability conditions for orbit couplings.

It is well known that orbit equivalence couplings are exactly those of the form $(\Omega, X_\Gamma, X_\Lambda, \mu)$ with $X_\Gamma = X_\Lambda := X$ of measure 1 and both induced actions on X are free. The freeness hypothesis is not a restriction, as we will see now.

Proposition 2.40. *Let (Ω, X_Λ, μ) be a measure sub-quotient coupling from Γ to Λ , let (Y, ν) be a standard probability space equipped with a free Γ -action. Then $(\Omega \times Y, X_\Lambda \times Y, \mu \otimes \lambda)$ is a measure sub-quotient coupling from Γ to Λ , where Γ acts diagonally and Λ acts on the first coordinate. Moreover, the induced Γ -action on $X_\Lambda \times Y$ is free, the coupling $(\Omega \times Y, X_\Lambda \times Y, \mu \otimes \lambda)$ is φ -integrable if and only if (Ω, X_Λ, μ) was, and if X_Γ was a fundamental domain for $\Gamma \curvearrowright \Omega$, then $X_\Gamma \times Y$ is a fundamental domain for $\Gamma \curvearrowright \Omega \times Y$.*

Proof. It is clear that $X_\Lambda \times Y$ is a fundamental domain for the Λ -action. Since the Γ -action on Ω is free, $X_\Gamma \times Y$ is a fundamental domain for the new diagonal action. The induced Γ -action on $X_\Lambda \times Y$ is the diagonal action obtained from its induced action on X_Λ and its action on Y . We deduce that this action is free. Finally, the statement about φ -integrability follows directly from the fact that for every $y \in Y$ and every $\omega, \omega' \in \Omega$, we have $d_{S_\Lambda}((\omega, y), (\omega', y)) = d_{S_\Lambda}(\omega, \omega')$. \square

Remark 2.41. Note that the above lemma can be applied to any countable group Γ : if Γ is infinite, one can take (Y, ν) as a Bernoulli shift of Γ , and if Γ is finite, one can take $Y = \Gamma$ acted upon by left translation, equipped with the normalized counting measure.

Proposition 2.42. *Let Γ and Λ be countable groups. If there is a (φ, ψ) measure equivalence (resp. measure quotient) coupling from Γ to Λ where the two fundamental domains coincide, then there is a (φ, ψ) orbit equivalence (resp. orbit quotient) coupling from Γ to Λ .*

Proof. Let $(\Omega, X_\Gamma, X_\Lambda, \mu)$ be a (φ, ψ) -integrable measure equivalence coupling from Γ to Λ , and denote by $X = X_\Gamma = X_\Lambda$ the common fundamental domain. Up to rescaling the measure (which does not impact the integrability conditions), we may as well assume that X has measure 1. Using the previous remark, we can apply the above proposition twice and see that without loss of generality, we can also assume that the induced Γ - and Λ -actions on X are free. We denote them by \cdot .

Now observe that for every $\gamma \in \Gamma$ and every $x \in X$, we have $\gamma \cdot x \in \Lambda * \gamma * x$ so there is $\lambda \in \Lambda$ such that $\gamma \cdot x = \lambda * \gamma * x = \gamma * \lambda * x$. In particular $\lambda \cdot x = \gamma \cdot x$, so we conclude that $\Gamma \cdot x \subseteq \Lambda \cdot x$. By the symmetry, we also have $\Lambda \cdot x \subseteq \Gamma \cdot x$, so we conclude that (X, μ) is an orbit equivalence coupling. Finally, the map $(\gamma \cdot x, x) \mapsto \gamma * x$ is a $\Gamma \times \Lambda$ -equivariant bijection from $\mathcal{R}_\Gamma = \mathcal{R}_\Lambda$ to Ω which takes $\{(x, x) : x \in X\}$ to X , and thus the (φ, ψ) -integrability of the orbit coupling (X, μ) follows from that of $(\Omega, X_\Gamma, X_\Lambda, \mu)$.

The statement for orbit quotient couplings follows from the same argument, except we apply the above proposition only once so as to make the induced Γ -action free. \square

Similar comparisons can be made for orbit subgroup and sub-quotient couplings. Observe that in this situation there is no canonical choice for a Γ fundamental domain X_Γ , but X_Λ intersects every Γ -orbit at most once.

Proposition 2.43. *Let Γ and Λ be two finitely generated groups. If there is a φ -integrable measure sub-quotient (resp. measure subgroup) coupling (Ω, X_Λ, μ) from Γ to Λ such that X_Λ intersects every Γ -orbit at most once, then there is a φ -integrable orbit sub-quotient (resp. orbit subgroup) coupling from Γ to Λ .*

Proof. We first assume that (Ω, X_Λ, μ) is a φ -integrable measure sub-quotient coupling from Γ to Λ such that X_Λ intersects every Γ -orbit at most once. Up to scaling the measure and using Proposition 2.40, we may as well assume that $\mu(X_\Lambda) = 1$ and that the induced Γ -action on X_Λ is free.

We define $Y := \Omega/\Gamma$ and $X := \Omega/\Lambda$, equipped with their respective actions of Λ and Γ . By assumption the restriction of $\pi_{\Omega/\Gamma}$ to X_Λ is injective, so identifying X with $\pi(X_\Lambda)$ we may as well view X as a subset of Y . To prove that this defines an orbit sub-quotient coupling, it remains to show that $\Gamma \cdot x \subseteq \Lambda \cdot x$ for every $x \in X$. For every $\gamma \in \Gamma$ and $x \in X_\Lambda$ there is $\lambda \in \Lambda$ such that $\gamma \cdot x = \lambda * \gamma * x \in X_\Lambda$. Then $\gamma * (\lambda * x) \in X_\Lambda$, so $\lambda \cdot x = \gamma * (\lambda * x) = \gamma \cdot x$. Thus, $\Gamma \cdot x \subseteq \Lambda \cdot x$, and (X, Y, μ) is indeed an orbit sub-quotient coupling.

We now check that the map $\lambda * x \mapsto (x, \lambda \cdot x)$ from Ω to $\mathcal{R}_\Lambda \cap X \times Y$ is well-defined and bijective. First, if $\lambda * x = \lambda' * x'$ for some $x, x' \in X_\Lambda$ and $\lambda, \lambda' \in \Lambda$, we have $x = x'$ since X_Λ is a fundamental domain, and then by definition of the induced action we must have $\lambda \cdot x = \lambda' \cdot x'$: our map is well-defined. In order to show it is injective, assume that $\lambda \cdot x = \lambda' \cdot x$, then $\lambda^{-1}\lambda' \cdot x = x$, so there is $\gamma \in \Gamma$ such that $\lambda^{-1}\lambda' * x = \gamma * x$, so $\gamma \cdot x = x$. By freeness of the induced Γ -action, we deduce that $\gamma = e_\Gamma$, and so $\lambda * x = \lambda' * x$ as wanted. Surjectivity is clear from the definition. Note that this map can be more formally defined as $(\pi_{\Omega/\Lambda}, \pi_{\Omega/\Gamma})$. Its $\Gamma \times \Lambda$ -equivariance is obvious from that description. We conclude that the orbit sub-quotient coupling that we obtained is φ -integrable if and only if our original coupling was.

For the orbit subgroup case we proceed as before, except we first make sure that both induced actions are free by using Proposition 2.40 twice. \square

Remark 2.44. The two previous propositions show that orbit couplings can be composed as in Proposition 2.9. This is shown by combining Proposition 2.9 with Proposition 2.42 and Proposition 2.43. The integrability of this composition coupling satisfies the same result as in Proposition 2.26 and Proposition 2.29.

3 Revisiting Lewis Bowen’s monotonicity theorem for the volume growth

We let Γ be a finitely generated group equipped with a finite generating subset S_Γ . By a slight abuse of notation, we denote $B_\Gamma(\gamma, n)$ the ball of radius n centered at $\gamma \in \Gamma$ for the left-invariant word metric d_{S_Γ} (therefore omitting the mention of S_Γ). We also denote $V_\Gamma(n) = |B_\Gamma(\gamma, n)|$. The volume growth of Γ is the asymptotic behavior of V_Γ (which does not depend on a specific choice of S_Γ). In [Aus16, Theorem B2], Lewis Bowen proves that the volume growth is invariant under L^1 measure equivalence⁵. We strengthen his result as follows.

Theorem 3.1. *Let φ be an increasing, subadditive function such that $\varphi(0) = 0$ and let Γ and Λ be finitely generated groups. If Γ is a φ -integrable measure sub-quotient of Λ , then*

$$V_\Gamma(n) \preceq V_\Lambda(\varphi^{-1}(n)),$$

where φ^{-1} denotes the inverse function of φ .

⁵Bowen actually proves a more general statement: he shows that the volume growth is monotonous under “integrable-embedding”. His notion seems strongly related to that of L^1 -measure subgroup, although we did not investigate this further.

The following key lemma is stated in [Aus16] for free actions, but the proof works as well for non-free actions.

Lemma 3.2. [Aus16, Lemma B.11] *Let Γ be a finitely generated group. Let $\Gamma \curvearrowright (X, \mu)$ be a measure-preserving action on a standard probability space, and let $X_0 \subseteq X$. For $x \in X_0$ let $R_{X_0}(x) := \{\gamma \in \Gamma : \gamma \cdot x \in X_0\}$ be its associated **return time set**. Then, for every $n \in \mathbb{N}$ we have*

$$\int_{X_0} \frac{|R_{X_0}(x) \cap B_\Gamma(e_\Gamma, n)|}{V_\Gamma(n)} d\mu(x) \geq 2\mu(X_0) - 1.$$

Proof of Theorem 3.1. Let (Ω, X_Λ, μ) be a sub-quotient coupling from Γ to Λ such that for all $s \in S_\Gamma$,

$$\int_{X_\Lambda} \varphi(d_{S_\Lambda}(s \cdot x, s * x)) d\mu(x) < \infty.$$

The set X_Λ has finite measure, so by replacing μ by $\frac{\mu}{\mu(X_\Lambda)}$, we may as well assume that $\mu(X_\Lambda) = 1$. We let X_Γ be a fundamental domain for the action of Γ . Since $\Gamma * X_\Gamma = \Omega$ there exists $R > 0$ such that the subset $X_0 = X_\Lambda \cap B_\Gamma(e_\Gamma, R) * X_\Gamma$ satisfies $\mu(X_0) \geq \frac{9}{10}$.

By Lemma 2.24 there exists $C > 0$ such that for every $\gamma \in \Gamma$,

$$\int_{X_\Lambda} \varphi(d_{S_\Gamma}(\gamma \cdot x, \gamma * x)) d\mu(x) \leq C|\gamma|. \quad (6)$$

We then fix $n \in \mathbb{N}$ and define

$$X_1 = \left\{ x \in X_\Lambda : \sum_{\gamma \in B_\Gamma(e_\Gamma, n)} \frac{\varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x))}{|\gamma|} \leq 10CV_\Gamma(n) \right\}.$$

Note that by applying Markov's inequality and then inequality (6), we have

$$\mu(X_\Lambda \setminus X_1) \leq \frac{1}{10CV_\Gamma(n)} \int_{X_\Lambda} \sum_{\gamma \in B_\Gamma(e_\Gamma, n)} \frac{\varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x))}{|\gamma|} d\mu(x) \leq \frac{1}{10}.$$

For every $x \in X_\Lambda$, we let

$$\Gamma_x = \{\gamma \in B_\Gamma(e_\Gamma, n) : \varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) \leq 60C|\gamma|\}.$$

Applying Markov's inequality to the finite probability space $B_\Gamma(e_\Gamma, n)$ equipped with the normalized counting measure and the definition of X_1 , we then have for all $x \in X_1$

$$|\Gamma_x| \geq \frac{5}{6}V_\Gamma(n). \quad (7)$$

Defining $X_2 = X_1 \cap X_0$, we have

$$\mu(X_2) \geq \frac{8}{10}. \quad (8)$$

We deduce from Lemma 3.2 that

$$\int_{X_2} \frac{|R_{X_2}(x) \cap B_\Gamma(e_\Gamma, n)|}{V_\Gamma(n)} \geq 2\mu(X_2) - 1 \geq \frac{6}{10}. \quad (9)$$

As $\Gamma_x \subseteq B_\Gamma(e_\Gamma, n)$ we have that

$$\begin{aligned} \int_{X_2} |R_{X_2}(x) \cap \Gamma_x| d\mu(x) &\geq \int_{X_2} (|R_{X_2}(x) \cap B_\Gamma(e_\Gamma, n)| + |\Gamma_x| - V_\Gamma(n)) d\mu(x) \\ &\geq V_\Gamma(n) \left(\frac{6}{10} + \frac{5}{6}\mu(X_2) - \mu(X_2) \right) \quad \text{by (7) and (9)} \\ &\geq \frac{1}{10}V_\Gamma(n). \end{aligned}$$

On the other hand

$$\begin{aligned}
\int_{X_2} |R_{X_2}(x) \cap \Gamma_x| d\mu(x) &\leq \int_{X_2} |\{\gamma \in R_{X_2}(x) : \varphi(d_{S_\Lambda}(\gamma \cdot x, \gamma * x)) \leq 60Cn\}| d\mu(x) \\
&\leq \sum_{\lambda \in B_\Lambda(e_\Lambda, \varphi^{-1}(60Cn))} \int_{X_2} |\{\gamma \in R_{X_2}(x) : \lambda * \gamma * x \in X_\Lambda\}| d\mu(x) \\
&\leq V_\Lambda(\varphi^{-1}(60Cn)) V_\Gamma(R) \mu(X_2),
\end{aligned}$$

where the last inequality results from the inclusion $X_2 \subset X_0$ together with the following claim.

Claim 3.3. Given $\lambda \in \Lambda$ and $x \in X_\Lambda$, the cardinality of $\{\gamma \in R_{X_0}(x) : \lambda * \gamma * x \in X_\Lambda\}$ is at most $V_\Gamma(R)$.

Proof. Indeed, let γ be in the above set. We have $\lambda * \gamma * x = \gamma \cdot x \in X_0$. In particular $\lambda * \gamma * x \in B_\Gamma(e_\Gamma, R) * X_\Gamma$, from which we deduce that $\gamma \in B_\Gamma(e_\Gamma, R) * \lambda^{-1} * X_\Gamma$. The conclusion follows by freeness of the Γ -action and the fact that $\lambda^{-1} * X_\Gamma$ is a fundamental domain for it. \square

Exploiting our upper and lower bounds for $\int_{X_2} |R_{X_2}(x) \cap \Gamma_x| d\mu(x)$ we deduce that for all $n \in \mathbb{N}$,

$$\frac{1}{10} V_\Gamma(n) \leq V_\Lambda(\varphi^{-1}(60Cn)) V_\Gamma(R),$$

which yields the conclusion of the theorem. \square

Corollary 3.4. Assume that two groups Γ and Λ satisfying $V_\Gamma(r) \approx r^a$ and $V_\Lambda(r) \approx r^b$, with $a > b$. Then Γ is not an L^p measured sub-quotient of Λ if $p > a/b$.

Proof. Since $a/b < 1$, and L^p -integrability gets stronger as p increases, it is enough to check it for $p \leq 1$. In this case, Theorem 3.1 applies as $x \mapsto x^p$ is subadditive. \square

4 Monotonicity of the isoperimetric profile

In this section we state and prove a general monotonicity result satisfied by the isoperimetric profile. We obtain two different conclusions depending on whether the coupling is L^p for some $p \geq 1$, or if it is φ -integrable for a sublinear function φ . However the first statement for $p = 1$ and the second one for $\varphi(t) = t$ have the same content. Roughly speaking the following theorem says that the isoperimetric profile behaves well under measure quotient (and so in particular under measure equivalence). In order to obtain a similar statement for measure sub-quotients, and in particular for measure subgroups, we will need the following technical assumption.

Definition 4.1. Given $m \in \mathbb{N}$, we say that a sub-quotient (resp. subgroup, resp quotient) coupling (Ω, X_Λ, μ) from Γ to Λ is **at most m -to-one** if for every $x \in X_\Lambda$ the map $\gamma \mapsto \gamma^{-1} * (\gamma \cdot x) \in \Lambda * x$ has pre-images of size at most m .

Theorem 4.2. Let $p \geq 1$, let Γ and Λ be finitely generated groups. Assume that we are in one of the following situations.

- (i) Γ is an L^p measure quotient of Λ ;
- (ii) Γ is an at most m -to-one L^p measure sub-quotient of Λ .

Then their L^p isoperimetric profiles satisfy the following inequality:

$$j_{p,\Gamma} \succcurlyeq j_{p,\Lambda}.$$

Theorem 4.3. Let $\varphi: (0, \infty) \rightarrow (0, \infty)$ be a function such that φ and $t \mapsto t/\varphi(t)$ are non-decreasing, let Γ and Λ be finitely generated groups. Assume that we are in one of the following situations.

- (i) Γ is a φ -integrable measure quotient of Λ ;
- (ii) Γ is an at most m -to-one φ -integrable measure sub-quotient of Λ

Then their isoperimetric profiles satisfy the following inequality:

$$j_{1,\Gamma} \succcurlyeq \varphi \circ j_{1,\Lambda}.$$

Remark 4.4. Formulated with the Følner function instead of the isoperimetric profile, we obtain $F_{\emptyset|\Gamma} \succcurlyeq F_{\emptyset|\Lambda} \circ \varphi^{-1}$.

Remark 4.5. We do not know whether the condition of being “at most m -to-one” is necessary. Fortunately, we shall see in §5 that it is satisfied in a case of interest: when Γ admits a regularly map to Λ , then Γ turns out to be an at most m -to-one L^∞ measure subgroup of Λ .

We illustrate this by the following corollary.

Corollary 4.6. Assume that Γ and Λ have polynomial growth of degree b and a respectively, with $b > a$, and let F and K be non-trivial finite groups.

- then Γ is not an L^p measured quotient (nor an at most m -to-one L^p measured sub-quotient) of Λ if $p > a/b$.
- $F \wr \Gamma$ is not an L^p measured quotient (nor an at most m -to-one L^p measured sub-quotient) of $K \wr \Lambda$ if $p > 1 - 1/a + 1/b$.

Proof. We recall the following estimates.

- $j_{1,\Gamma}(n) \approx n^{1/d}$ for a group of polynomial growth of degree d (e.g. for $\Gamma = \mathbb{Z}^d$) (see [Cou00]);
- $j_{1,\Gamma}(n) \approx (\log n)^{1/d}$ for a group of the form $F \wr \Sigma$, where F is a non-trivial finite group, and Σ has polynomial growth of degree d [Ers03].

We obtain both statements by confronting Theorem 4.2(ii) to these estimates. \square

Remark 4.7. The first item is slightly weaker than the conclusion of Corollary 3.4 obtained by comparing the volume growths (as the latter does not require the at most m -to-one assumption).

We end this section discussing some important situations where the condition of being at most m -to-one is satisfied.

Proposition 4.8. A measure sub-quotient coupling (Ω, X_Λ, μ) from Γ to Λ is at most 1-to-one if and only if X_Λ intersects every Γ -orbit at most once.

Proof. Suppose first that X_Λ intersects each Γ -orbit at most once. Let $x \in X_\Lambda$, suppose that $\gamma_1^{-1} * \gamma_1 \cdot x = \gamma_2^{-1} * \gamma_2 \cdot x$. Then $\gamma_1 \cdot x$ and $\gamma_2 \cdot x$ are two elements of the same Γ -orbit which belong to X_Λ . So by our assumption they are equal. Since the Γ -action on Ω is free, we conclude that $\gamma_1 = \gamma_2$ as wanted.

Conversely, suppose that the coupling is at most 1-to-one. Let $x \in X_\Lambda$, suppose that $\gamma * x \in X_\Lambda$ for some $\gamma \in \Gamma$. Then $\gamma \cdot x = \gamma * x$, so $\gamma^{-1} * \gamma \cdot x = x$, and so by injectivity $\gamma = e_\Gamma$. We conclude that $\gamma * x = x$, so X_Λ intersects each Γ -orbit at most once as announced. \square

An important example of at most 1-to-one coupling is provided by the notion of *orbit sub-quotient couplings* (see Section 2.6). In fact, Proposition 2.43 implies that every at most 1-to-one measure sub-quotient coupling can be turned into such a coupling after making the induced action free. In particular, we have

Corollary 4.9. Let $\varphi: (0, \infty) \rightarrow (0, \infty)$ be a function such that φ and $t \mapsto t/\varphi(t)$ are non-decreasing. Assume that Γ is a φ -integrable orbit sub-quotient of Λ . Then $j_{1,\Gamma} \succcurlyeq \varphi \circ j_{1,\Lambda}$. \square

4.1 Overview of the proof

Let us start describing the main steps of the proof when the Λ -action is free.

The first step is to start from a finitely supported function f on Λ with small *right* gradient and extend it to a function \tilde{f} on the coupling space Ω through its identification with $X_\Lambda \times \Lambda$. In other words, we identify Ω to a field copies of Λ which are pointed at X_Λ . The key observation is that the function \tilde{f} has small gradient with respect to the Γ -action. This is because each element $\gamma \in \Gamma$ induces an isomorphism of Λ -spaces, only shifting the basepoint by an amount which is precisely controlled by our integrability condition: γ takes the field of copies of Λ pointed at X_Λ to the field of copies of Λ pointed at $\gamma * X_\Lambda$. The precise statement which allows to control the difference between $\gamma \cdot \tilde{f}$ and \tilde{f} is provided by Lemma 4.12.

Once this is established, we identify Ω with $\Gamma \times X_\Gamma$ and view \tilde{f} as a family (f^x) of finitely supported functions on Γ . By the previous fact, the gradient of these functions is well behaved on a large measure subset of X_Γ . We need to simultaneously control from below the ℓ^p -norm of the function, and control from above the size of the support of these functions. Let us describe the simpler situation where the coupling is assumed to be at most m -to-one. This condition immediately implies that the size of the support of f^x is (almost surely) at most m times the size of the support of f . On the other hand, a pigeonhole argument allows us to find a set of positive measure of X_Γ on which the lower bound on the norm is satisfied simultaneously with the upper bound on the gradient. This is enough to conclude for the existence of a subset of positive measure for which f^x satisfies all three conditions simultaneously and therefore for the proof of the theorem.

If we do not assume that the coupling is at most m -to-one, a difficulty arises that is overcome in §4.5. What remains possible using pigeonhole arguments is the following: we can find a subset of X_Γ with measure –say at least $1/2$ – where an upper bound on both the size of the support of f^x and the norm of its gradient hold simultaneously. However, the issue comes from the lower bound on the norm of f^x , which may be only valid on a subset of arbitrarily small measure. The solution that we find is rather subtle and consists in slightly modifying f in such a way that the lower bound on the norm of f^x is actually satisfied on a subset of X_Γ of relatively large measure (see Remark 4.21 for a more detailed overview of the argument).

Finally, when the action of Λ is not free, we still find a way to define a function \tilde{f} on Ω which in some sense extends f . The rest of proof is identical.

4.2 Λ -gradients on Λ -spaces

It will be convenient to define a notion of gradient on functions defined on measure spaces equipped with a measure-preserving group action.

Definition 4.10. Let Λ be a finitely generated group equipped with a finite symmetric generating set S_Λ . Given any standard measured Λ -space (Ω, μ) and any function $f \in L^p(\Omega, \mu)$, define its gradient to be the function $\nabla_{S_\Lambda} f: S_\Lambda \times \Omega \rightarrow \mathbb{R}$ defined by

$$\nabla_{S_\Lambda} f(s, x) = f(x) - f(s^{-1} * x).$$

Remark 4.11. We have

$$\|\nabla_{S_\Lambda} f\|_p^p = \int_\Omega \sum_{s \in S_\Lambda} |f(\omega) - f(s^{-1} * \omega)|^p d\mu(\omega) = \sum_{s \in S_\Lambda} \|f - s * f\|_p^p,$$

where $s * f(x) = f(s^{-1} * x)$. Note that by symmetry of S_Λ , we also have

$$\|\nabla_{S_\Lambda} f\|_p^p = \int_\Omega \sum_{s \in S_\Lambda} |f(\omega) - f(s * \omega)|^p d\mu(\omega)$$

There are two natural ways to view Λ as a Λ -space (action by left or right multiplication), so we define the *left* S_Λ -gradient f as the function $\nabla_{S_\Lambda}^l f: S_\Lambda \times \Lambda \rightarrow \mathbb{R}$ by the formula

$$\nabla_{S_\Lambda}^l f(s, \lambda) = f(\lambda) - f(s^{-1}\lambda);$$

and the *right* S_Λ -gradient f by

$$\nabla_{S_\Lambda}^r f(s, \lambda) = f(\lambda) - f(\lambda s).$$

The isoperimetric profile can be alternatively defined using the left or right gradient. This clearly does not change its value as the inverse map intertwines the actions by left and right multiplication. We will exploit this remark in our proof as follows: we will start with a function on Λ with small *right* gradient and induce on Γ a function of comparable support with small *left* gradient.

In the following lemma, we consider a Γ -space X . Starting with a function defined on Λ , we define a functions on X depending on some basepoint x . The main objective of the lemma is to control the L^p -norm of the difference between functions associated to different base points. It is reminiscent of the fact that right Reiter functions on a group Λ can be pushed to Reiter functions on any equivalence relation induced by a Λ -action (see e.g. [KM04, Prop. 9.2]).

Lemma 4.12. *Let X be a transitive Λ -set and let $x_0 \in X$. Let $p \geq 1$. The map $\mathbb{R}^\Lambda \rightarrow \mathbb{R}^X$ which associates to every $f \in \mathbb{R}^\Lambda$ the function f_{x_0} given by*

$$f_{x_0}(y) = \left(\sum_{\lambda: \gamma * x_0 = y} |f(\lambda)|^p \right)^{1/p}$$

satisfies $\|f_{x_0}\|_p = \|f\|_p$. Moreover, if S_Λ is a finite generating set for Λ and if $f \in \ell^p(\Lambda)$ then for all $x_1 \in X$,

$$\|f_{x_0} - f_{x_1}\|_p \leq d_{S_\Lambda}(x_0, x_1) \|\nabla_{S_\Lambda}^r f\|_p. \quad (10)$$

Proof. Now assume that $p \geq 1$. The verification that $\|f_{x_0}\|_p = \|f\|_p$ is immediate. To obtain the inequality (10), note that for each $s \in S_\Gamma$ and $x_0 \in X$, we have by the triangle difference inequality

$$\begin{aligned} \|f_{x_0} - f_{s \cdot x_0}\|_p^p &= \sum_{y \in X} \left| \left(\sum_{\lambda: \lambda * x_0 = y} |f(\lambda)|^p \right)^{1/p} - \left(\sum_{\lambda: \lambda s * x_0 = y} |f(\lambda)|^p \right)^{1/p} \right|^p \\ &= \sum_{y \in X} \left| \left(\sum_{\lambda: \lambda * x_0 = y} |f(\lambda)|^p \right)^{1/p} - \left(\sum_{\lambda: \lambda * x_0 = y} |f(\lambda s^{-1})|^p \right)^{1/p} \right|^p \\ &\leq \sum_{y \in X} \sum_{\lambda: \lambda * x_0 = y} |f(\lambda) - f(\lambda s^{-1})|^p \\ &\leq \sum_{\lambda \in \Lambda} |f(\lambda) - f(\lambda s^{-1})|^p, \end{aligned}$$

so by the definition of the gradient we have $\|f_{x_0} - f_{s \cdot x_0}\|_p \leq \|\nabla_{S_\Lambda}^r f\|_p$. The conclusion now follows using the triangle inequality for the L^p -norm and the fact that Γ is acting by isometries on $\ell^p(X)$. \square

We also need the following variant of Lemma 4.12.

Lemma 4.13. *Let $\varphi: (0, \infty) \rightarrow (0, \infty)$ be a function such that both φ and $t \mapsto t/\varphi(t)$ are non-decreasing. Given a transitive Λ -set X and a point $x_0 \in X$, the map $\mathbb{R}^\Lambda \rightarrow \mathbb{R}^X$ which associates to every $f \in \mathbb{R}^\Lambda$ the function f_{x_0} given by*

$$f_{x_0}(y) = \sum_{\lambda: \lambda * x_0 = y} |f(\lambda)|$$

satisfies $\|f_{x_0}\|_1 = \|f\|_1$. Moreover, if S_Λ is a finite generating set for Λ , $x_0, x_1 \in X$ and f is a finitely supported function on Λ such that $\|\nabla_{S_\Lambda}^r f\|_1 = 1$, then

$$\frac{\|f\|_1}{\|f_{x_0} - f_{x_1}\|_1} \geq \frac{\varphi(\|f\|_1)}{2\varphi(d_{S_\Lambda}(x_0, x_1))}. \quad (11)$$

Proof. By triangle inequality, we obtain

$$\|f_{x_0} - f_{x_1}\|_1 \leq 2 \max_{i=0,1} \|f_{x_i}\|_1 = 2\|f\|_1.$$

Using the monotonicity of $t/\varphi(t)$ we have

$$\|f_{x_0} - f_{x_1}\|_1 = \varphi(\|f_{x_0} - f_{x_1}\|_1) \frac{\|f_{x_0} - f_{x_1}\|_1}{\varphi(\|f_{x_0} - f_{x_1}\|_1)} \leq \varphi(\|f_{x_0} - f_{x_1}\|_1) \frac{2\|f\|_1}{\varphi(2\|f\|_1)}.$$

Now, since $\|\nabla f\|_1 = 1$, by applying Lemma 4.12 for $p = 1$ and using the monotonicity of φ we get

$$\|f_{x_0} - f_{x_1}\|_1 \leq 2\varphi(d_{S_\Gamma}(x_0, x_1)) \frac{\|f\|_1}{\varphi(\|f\|_1)}$$

So the lemma follows. \square

4.3 The induction technique

In this section, we shall prove that if $p \geq 1$ and Γ is an L^p measure *sub-quotient* of Λ , then every ℓ^p function on Λ induces a function on Ω whose S_Γ -gradient is well behaved (and a similar statement for φ measure sub-quotient). To deal with sub-quotients, we need a non-free analogue of the fact that if Λ is acting freely on Ω and X_Λ is a fundamental domain, the map $(x, \lambda) \mapsto \lambda * x$ is a measure-preserving bijection between $X_\Lambda \times \Lambda$ and Ω .

Lemma 4.14. *Suppose X_Λ is a fundamental domain for a measure-preserving Λ -action on a standard measured space (Ω, μ) . Then for every Borel $A \subseteq \Omega$, we have*

$$\mu(A) = \int_{X_\Lambda} |(\Lambda * x) \cap A| d\mu(x).$$

In particular, for every measurable function $f: \Omega \rightarrow \mathbb{R}$, we have

$$\int_{\Omega} f d\mu = \int_{X_\Lambda} \sum_{y \in \Lambda * x} f(y) d\mu(x).$$

Proof. Since every subset of Ω can be written as a countable disjoint union of Λ -translates of Borel subsets of X_Λ , it suffices to show that the right term defines a Borel measure on Ω which is Λ -invariant and coincides with μ when restricted to X_Λ . The fact that the formula

$$m(A) = \int_{X_\Lambda} |(\Lambda * x) \cap A| d\mu(x)$$

defines a Borel measure follows from the fact that the map $x \mapsto |(\Lambda * x) \cap A|$ is Borel. Λ -invariance is clear, and the fact that the two measures coincide when restricted on X_Λ is also straightforward to check. \square

Proposition 4.15 (Monotonicity of the ℓ^p -gradient under L^p measured sub-quotient). *Let $p \geq 1$. If Γ is an L^p measure sub-quotient of Λ via a coupling (Ω, X_Λ, μ) , and if $f \in \ell^p(\Lambda)$, then the induced function \tilde{f} on Ω defined by $\tilde{f}(\omega) = (\sum_{\lambda: \lambda^{-1} * \omega \in X_\Lambda} |f(\lambda)|^p)^{1/p}$ satisfies*

$$\left\| \tilde{f} \right\|_p^p = \|f\|_p^p \mu(X_\Lambda),$$

and

$$\left\| \nabla_{S_\Gamma} \tilde{f} \right\|_p^p \leq C \left\| \nabla_{S_\Lambda}^r f \right\|_p^p,$$

where $C = |S_\Gamma| \max_{s \in S_\Gamma} \int_{X_\Lambda} d_{S_\Lambda}(s \cdot x, s * x)^p d\mu(x)$.

Proof. The first equality is obvious. For all $(x, y) \in \Omega^2$ such that $y \in \Lambda * x$, we let

$$f_x(y) = \left(\sum_{\lambda: \lambda * x = y} |f(\lambda)|^p \right)^{1/p}.$$

Identifying Ω with the set of pairs (x, y) such that $x \in X_\Lambda$ and $y \in \Lambda * x$, we have

$$\tilde{f}(x, y) = f_x(y).$$

For every $\gamma \in \Gamma$,

$$\tilde{f}(\gamma * (x, y)) = f_{\gamma \cdot x}(\gamma * y).$$

Note that since the actions of Γ and Λ commute, we have for all $\gamma \in \Gamma$

$$f_{\gamma * x}(\gamma * y) = f_x(y).$$

Hence, denoting $\gamma^{-1} * \tilde{f}: \omega \mapsto \tilde{f}(\gamma * \omega)$,

$$\gamma^{-1} * \tilde{f}(x, y) - \tilde{f}(x, y) = f_{\gamma \cdot x}(\gamma * y) - f_{\gamma * x}(\gamma * y).$$

We can now invoke Lemma 4.12: letting $s \in S_\Gamma$, we have

$$\begin{aligned} \sum_{y \in \Lambda * x} \left| s^{-1} * \tilde{f}(x, y) - \tilde{f}(x, y) \right|^p &= \sum_{y \in \Lambda * x} |f_{s \cdot x}(s * y) - f_{s * x}(s * y)|^p \\ &= \sum_{y \in \Lambda * x} |f_{s \cdot x}(y) - f_{s * x}(y)|^p \\ &\leq d_{S_\Lambda}(s \cdot x, s * x)^p \left\| \nabla_{S_\Lambda}^r f \right\|_p^p \end{aligned}$$

We can now use Lemma 4.14 to compute for all $s \in S_\Gamma$

$$\begin{aligned} \left\| s^{-1} * \tilde{f} - \tilde{f} \right\|_p^p &= \int_{X_\Lambda} \sum_{y \in \Lambda * x} \left| s^{-1} * \tilde{f}(x, y) - \tilde{f}(x, y) \right|^p \\ &\leq \int_{X_\Lambda} d_{S_\Lambda}(s \cdot x, s * x)^p \left\| \nabla_{S_\Lambda}^r f \right\|_p^p \end{aligned}$$

The desired inequality now follows by the definition of $\nabla_{S_\Gamma} \tilde{f}$ and the symmetry of S_Γ . \square

Proposition 4.16 (Monotonicity of the ℓ^1 -gradient under φ measured sub-quotient). *Let $\varphi: (0, \infty) \rightarrow (0, \infty)$ be a function such that φ and $t/\varphi(t)$ are non-decreasing. If Γ is an φ measure sub-quotient of Λ via a coupling (Ω, X_Λ, μ) , and if f is a finitely supported function on Λ such that $\left\| \nabla_{S_\Lambda}^r f \right\|_1 = 1$, then the induced function \tilde{f} on Ω defined by $\tilde{f}(\omega) = \sum_{h: h^{-1} * \omega \in X_\Lambda} f(h)$ satisfies*

$$\frac{\left\| \nabla_{S_\Gamma} \tilde{f} \right\|_1}{\left\| \tilde{f} \right\|_1} \leq \frac{2C}{\varphi(\|f\|_1)},$$

where $C = \frac{|S_\Gamma|}{\mu(X_\Lambda)} \max_{s \in S_\Gamma} \int_{X_\Lambda} \varphi(d_{S_\Lambda}(s \cdot x, s * x)) d\mu(x)$.

Proof. The proof is similar to that of Proposition 4.15: using Lemma 4.13 applied to $f_x(y) := \sum_{\lambda: \lambda * x = y} |f(\lambda)|$, we obtain

$$\frac{\sum_{y \in \Lambda * x} \left| s^{-1} * \tilde{f}(x, y) - \tilde{f}(x, y) \right|}{\|f\|_1} \leq \frac{2\varphi(d_{S_\Lambda}(s \cdot x, s * x))}{\varphi(\|f\|_1)},$$

which together with Lemma 4.14 and the fact that $\|\tilde{f}\|_1 = \mu(X_\Lambda) \|f\|_1$ yields the desired inequality. \square

Remark 4.17. Using Fubini's theorem and the natural identification of $X_\Gamma \times \Gamma$ with Ω given by $(x, g) \mapsto g * x$, we may rewrite $\left\| \nabla_{S_\Gamma} \tilde{f} \right\|_p^p$ as

$$\left\| \nabla_{S_\Gamma} \tilde{f} \right\|_p^p = \int_{X_\Gamma} \sum_{s \in S_\Gamma} \sum_{\gamma \in \Gamma} \left| \tilde{f}(\gamma * x) - \tilde{f}(s\gamma * x) \right|^p d\mu(x).$$

Now for each $x \in X_\Gamma$, the function \tilde{f} defines a function f^x on Γ given by $f^x(\gamma) = \tilde{f}(\gamma * x)$, and the previous equality may thus be rewritten as

$$\left\| \nabla_{S_\Gamma} \tilde{f} \right\|_p^p = \int_{X_\Gamma} \left\| \nabla_{S_\Gamma}^l f^x \right\|_p^p d\mu(x).$$

So the conclusion of the two previous propositions is that under the right assumption, any function f on Λ of small *right* ℓ^p -gradient induces functions f^x on Γ which have on average a small *left* ℓ^p gradient. From there on, as we will see in the next section, it is not hard to conclude the proof that the isoperimetric profile goes down for $p \geq 1$ under the assumption that the coupling is finite-to-one.

But without the finite-to-one assumption, we lose the uniform control on the size of the support of the functions f^x . We will circumvent this by simultaneously controlling the size of the support of f^x and bounding its norm from below on a large portion of the fundamental domain of Γ . This will be done in Section 4.5.

4.4 Monotonicity under at most m -to-one measure sub-quotients

In this section we shall prove the second items of Theorem 4.2 and Theorem 4.3.

The common feature between these statements is that the coupling is supposed to be at most m -to-one. This has the following consequence.

Lemma 4.18. *Let (Ω, X_Λ, μ) be an at most m -to-one measure sub-quotient coupling between Γ and Λ . Let X_Γ be a fundamental domain for the Γ -action. Let f be a function on Λ whose support has cardinality at most K , and let $(f^x)_{x \in X_\Gamma}$ be the family of functions on Γ defined by*

$$f^x(\gamma) = \left(\sum_{\lambda: \lambda * (\gamma * x) \in X_\Lambda} |f(\lambda)|^p \right)^{1/p},$$

for some $p > 0$. Then for each $x \in X_\Gamma$, the function f^x has support of cardinality at most mK .

Proof. Let $x \in X_\Gamma$, let $\lambda_0 \in \Lambda$ such that $\lambda_0 * x \in X_\Lambda$. By definition for every $\gamma \in \Gamma$ and $\lambda \in \Lambda$ we have $\lambda * (\gamma * x) \in X_\Lambda$ if and only if $\lambda * (\gamma * x) = \gamma \cdot (\lambda_0 * x)$, which since the Γ and Λ -actions on Ω commute is in turn equivalent to $\lambda * x = \gamma^{-1} * (\gamma \cdot (\lambda_0 * x))$. So we have

$$f^x(\gamma) = \sum_{\lambda: \lambda * x = \gamma^{-1} * (\gamma \cdot (\lambda_0 * x))} f(\lambda).$$

The conclusion now follows from our assumption that the map $\gamma \mapsto \gamma^{-1} * (\gamma \cdot (\lambda_0 * x))$ is at most m -to-one. \square

Proof of Theorem 4.2(ii). We start with a function f that realize the L^p -isoperimetric profile of Λ , and we consider the function \tilde{f} on Ω defined in Proposition 4.15. By Proposition 4.15, there exists C' only depending on the coupling such that

$$\frac{\|\nabla_{S_\Gamma} \tilde{f}\|_p^p}{\|\tilde{f}\|_p^p} \leq C' \frac{\|\nabla_{S_\Lambda}^r f\|_p^p}{\|f\|_p^p}. \quad (12)$$

This implies that on a set of positive measure, the function f^x on Γ satisfies

$$\frac{\|\nabla_{S_\Gamma}^l f^x\|_p^p}{\|f^x\|_p^p} < 2C' \frac{\|\nabla_{S_\Lambda}^r f\|_p^p}{\|f\|_p^p}. \quad (13)$$

Indeed, assume by contradiction that the reverse inequality holds on a subset Z of full measure. This means that for all $x \in Z$,

$$\|\nabla_{S_\Gamma}^l f^x\|_p^p \geq 2C' \frac{\|\nabla_{S_\Lambda}^r f\|_p^p}{\|f\|_p^p} \|f^x\|_p^p.$$

But integrating over $x \in Z$, we get a contradiction with (12). On the other hand, by Lemma 4.18, the support of f^x has size at most $m|\text{supp}(f)|$. Hence we deduce that $j_{p,\Gamma}(mn) \geq \frac{j_{p,\Lambda}(n)}{2C'}$, so we are done. \square

Remark 4.19. Note that the previous argument does not provide any information on the measure on which inequality (13) holds.

Proof of Theorem 4.3(ii). As before we start with a function f that realizes the L^1 -isoperimetric profile of Λ . We then normalize f such that $\|\nabla_{S_\Lambda}^r f\|_1 = 1$ and consider the function \tilde{f} on Ω defined in Proposition 4.15. We deduce from Proposition 4.16 that

$$\frac{\|\nabla_{S_\Gamma} \tilde{f}\|_1}{\|\tilde{f}\|_1} \leq \frac{C'}{\varphi(\|f\|_1)},$$

and the rest of the proof is identical. \square

4.5 Monotonicity under measure quotients

In this section we prove the first items of Theorem 4.2 and Theorem 4.3. On rescaling the measure, we may assume that X_Γ has measure 1. Note that since X_Λ intersects every Λ -orbit, we can then find a finite subset W of Λ so that $Z := (W * X_\Lambda) \cap X_\Gamma$ has measure at least $3/4$. Let us fix once and for all such a set W .

Given $n \geq 1$, we let f_1 be a function that realizes the (right) isoperimetric profile of Λ at n . We start modifying it as follows. We define a function f_2 on Λ by

$$f_2(\lambda) = \left(\sum_{w \in W} |f_1(w^{-1}\lambda)|^p \right)^{1/p}.$$

The important fact is that f_2 “almost” realizes the isoperimetric profile of Λ : the support of f_2 has size at most $|W|$ times the support of f_1 , and we have $\|f_2\|_p \geq \|f_1\|_p$. Using the same inequalities as in the proof of Lemma 4.12 we obtain

$$\|\nabla_{S_\Gamma}^r f_2\|_p^p \leq |W| \|\nabla_{S_\Gamma}^r f_1\|_p^p. \quad (14)$$

For all $p > 0$, we set

$$\tilde{f}_2(\omega) = \left(\sum_{\lambda: \lambda * \omega \in X_\Lambda} f_2(\lambda)^p \right)^{1/p}.$$

As in the previous section, this provides a random function f_2^x on Γ for $x \in X_\Gamma$ given by

$$f_2^x(\gamma) = \tilde{f}_2(\gamma * x) = \left(\sum_{\lambda: \lambda * \gamma * x \in X_\Lambda} f_2(\lambda)^p \right)^{1/p}.$$

Therefore, under the assumption that $p \geq 1$ (in order to prove Theorem 4.2.(i)), Proposition 4.15 and Remark 4.17 imply that for all $\varepsilon > 0$, we find two subsets V and V' of X_Γ of measure $1 - \varepsilon$, such that for all $x \in V$,

$$\|\nabla_{S_\Gamma}^l f_2^x\|_p^p \leq \frac{C}{\varepsilon} \|\nabla_{S_\Gamma}^r f_2\|_p^p,$$

and for all $x \in V'$

$$|\text{supp}(f_2^x)| \leq \frac{1}{\varepsilon} \mu(X_\Lambda) |\text{supp}(f_2)|.$$

For the proof of Theorem 4.3, we normalize f_2 such that $\|\nabla_{S_\Gamma} f_2\|_1 = 1$. Using Proposition 4.16 instead of Proposition 4.15, we find two subsets V and V' of X_Γ of measure $1 - \varepsilon$, such that for all $x \in V$,

$$\frac{\|\nabla_{S_\Gamma}^l f_2^x\|_1}{\|f_2\|_1} \leq \frac{C}{\varepsilon \varphi(\|f_2\|_1)},$$

and for all $x \in V'$

$$|\text{supp}(f_2^x)| \leq \frac{1}{\varepsilon} \mu(X_\Lambda) |\text{supp}(f_2)|.$$

In what follows, it will be sufficient to take $\varepsilon = 1/5$. The proof of Items (i) of Theorem 4.2 and Theorem 4.3 will be finished if we can find a subset $Y \subset X_\Gamma$ of measure $> 2/5$ on which the ℓ^p -norm of f_2^x satisfies a uniform lower bound. Indeed, this ensures that for all x in the non-empty set $V \cap V' \cap Y$, the function f_2^x gives the required bound on the (left) isoperimetric profile of Γ . The existence of such a subset is given by the following lemma, whose proof occupies the rest of the subsection. This is where we use our initial assumption that the set $Z = (W * X_\Lambda) \cap X_\Gamma$ has measure at least $3/4$.

Lemma 4.20. *There exists a measurable subset $Y \subset X_\Gamma$ of measure at least $1/2$ such that for all $y \in Y$,*

$$\|f_2^y\|_p^p \geq \frac{1}{4} \|f_1\|_p^p.$$

Remark 4.21. Before proceeding to the proof of the lemma, let us briefly explain its main idea. The issue that prevents us from obtaining such a lower bound directly from the lower bound on the norm of \tilde{f}_2 comes from the fact that we cannot rule out the possibility that in the integral

$$\left\| \tilde{f}_2 \right\|_p^p = \int_{X_\Gamma} \|f_2^x\|_p^p d\mu(x),$$

most of the contribution comes from a very tiny portion of X_Γ where $\|f_2^x\|_p$ is much bigger than $\|f_2\|_p$. If we could combine this lower bound with an uniform upper bound $\|f_2^x\|_p^p \leq K \|f_2\|_p^p$, then this problem would not arise. The reason why such an upper bound is a priori not available is due to the fact that the Γ -orbit of $x \in X_\Gamma$ could potentially meet many times a same translate of X_Λ . In other words, when writing $\|f_2^x\|_p^p = \sum_\gamma \sum_{\lambda: \lambda * \gamma * x \in X_\Lambda} f_2(\lambda)^p = \sum_\lambda n_x(\lambda) f_2(\lambda)^p$, the ‘‘multiplicities’’ $n_x(\lambda) \in \mathbb{N}$ might be very large. Therefore, our strategy consists in replacing this expression by a sum where all the multiplicities are at most one. To be more precise, we will replace it by $\sum_{\lambda, \lambda \cdot x \in Z} |f_1(\lambda)|^p$.

We first have to make sure that this new expression is controlled from above by $\left\| \tilde{f}_2 \right\|_p^p$ (Inequation 15)

and check that after integrating over X_Γ , this new expression still captures a reasonable fraction of $\|f_1\|_p^p$ (Inequation 16). The conclusion will then follow by confronting this lower bound on the integral with the pointwise trivial upper bound $\sum_{\lambda, \lambda \cdot x \in Z} |f_1(\lambda)|^p \leq \|f_1\|_p^p$ (Claim 4.22).

Proof. We start by establishing the following upper bound.

$$\sum_{\lambda, \lambda \cdot x \in Z} |f_1(\lambda)|^p \leq \|f_2^x\|_p^p. \quad (15)$$

First note that for all $x \in X_\Gamma$ and $\lambda \in \Lambda$ such that $\lambda \cdot x \in Z$, there is $\gamma \in \Gamma$ such that $\lambda * \gamma * x \in W * X_\Lambda$. Indeed, this follows by inspection of the definitions: $\lambda \cdot x \in Z$ means that there exists $\gamma \in \Gamma$ such that $\lambda * \gamma * x \in Z$, and therefore that $\lambda * \gamma * x \in W * X_\Lambda$.

We deduce the following crude inequality, which we then rewrite by exchanging orders of summation:

$$\begin{aligned} \sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p &\leq \sum_{\lambda} \sum_{\gamma: \lambda * \gamma * x \in W * X_\Lambda} |f_1(\lambda)|^p \\ &\leq \sum_{\gamma} \sum_{\lambda: \lambda * \gamma * x \in W * X_\Lambda} |f_1(\lambda)|^p. \end{aligned}$$

We can bound the above sum by

$$\begin{aligned} \sum_{\gamma} \sum_{w \in W} \sum_{\lambda: w \lambda * \gamma * x \in X_\Lambda} |f_1(\lambda)|^p &= \sum_{\gamma} \sum_{w \in W} \sum_{\lambda: \lambda * \gamma * x \in X_\Lambda} |f_1(w^{-1}\lambda)|^p \\ &= \sum_{\gamma} \sum_{\lambda: \lambda * \gamma * x \in X_\Lambda} f_2(\lambda)^p \\ &= \|f_2^x\|_p^p. \end{aligned}$$

Putting this together with the previous inequality, we obtain (15).

We then have the following inequality, using our initial assumption that the set $Z = (W * X_\Lambda) \cap X_\Gamma$ has measure at least $3/4$:

$$\int_{X_\Gamma} \sum_{\lambda, \lambda \cdot x \in Z} |f_1(\lambda)|^p d\mu(x) \geq \frac{3}{4} \|f_1\|_p^p. \quad (16)$$

Indeed, the following chain of inequalities hold:

$$\begin{aligned} \int_{X_\Gamma} \sum_{\lambda, \lambda \cdot x \in Z} |f_1(\lambda)|^p d\mu(x) &\geq \int_{X_\Gamma} \sum_{\lambda \in \Lambda} |f_1(\lambda)|^p 1_{\lambda^{-1} \cdot Z}(x) d\mu(x) \\ &\geq \sum_{\lambda \in \Lambda} |f_1(\lambda)|^p \int_{X_\Gamma} 1_{\lambda^{-1} \cdot Z}(x) d\mu(x) \\ &= \|f_1\|_p^p \mu(Z) \geq \frac{3}{4} \|f_1\|_p^p, \end{aligned}$$

so (16) is proved. We now use this inequality in our final claim.

Claim 4.22. There is a subset $Y \subset X_\Gamma$ of measure at least $1/2$ such that for all $x \in Y$,

$$\sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p \geq \frac{1}{4} \|f_1\|_p^p.$$

Proof. Indeed, assume by contradiction that

$$\sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p < \frac{1}{4} \|f_1\|_p^p$$

for all x in a set $U \subset X_\Gamma$ of measure strictly larger than $1/2$. Then by Inequation 16, we have

$$\begin{aligned} \frac{3}{4} \|f_1\|_p^p &\leq \int_U \sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p d\mu(x) + \int_{U^c} \sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p d\mu(x) \\ &< \frac{1}{4} \|f_1\|_p^p + \frac{1}{2} \|f_1\|_p^p \\ &< \frac{3}{4} \|f_1\|_p^p, \end{aligned}$$

which is a contradiction. \square

Putting the above claim together with inequality (15) we deduce that for all $x \in Y$,

$$\sum_{\gamma \in \Gamma} |f_2^x(\gamma)|^p \geq \sum_{\lambda: \lambda \cdot x \in Z} |f_1(\lambda)|^p \geq \frac{1}{4} \|f_1\|_p^p$$

as wanted. \square

4.6 Can there be a quantitative version of the Ornstein-Weiss theorem?

In this section, we first observe that any two amenable groups admit a (φ, φ) -integrable orbit equivalence coupling for some φ which grows slower than the logarithm, and then that there cannot be a universal such φ , thus proving Corollary 4 from the introduction.

Proposition 4.23. *For every orbit equivalence coupling between two finitely generated groups Γ and Λ , there exists a concave increasing unbounded function φ with $\varphi(0) = 0$ and a (φ, φ) orbit equivalence coupling between them. Moreover one can assume that $\varphi(t)/\log t$ is non-increasing on $[1, \infty)$.*

Proof. Let

$$k(t) = \mu \left(\left\{ x \in X; \max_{s \in S_\Gamma} |\alpha(s, x)|_{S_\Lambda} \geq t \right\} \right) + \mu \left(\left\{ x \in X; \max_{s \in S_\Lambda} |\beta(s, x)|_{S_\Gamma} \geq t \right\} \right).$$

Since k tends to zero at infinity, there exists an increasing sequence of positive integers (a_n) such that $a_0 = 0$, and $k(a_n) \leq 2^{-n}$. Up to taking a subsequence, we can assume that $a_{n+1} - a_n$ and $\frac{\log a_n}{n}$ are non-decreasing. Let φ be the continuous piecewise linear function with breakpoints a_n satisfying $\varphi(a_n) = n$. Note that φ is increasing, concave and such that $\varphi(t)/\log t$ is non-increasing on $[1, \infty)$. Moreover we have for all $s \in S_\Gamma$,

$$\begin{aligned} \int \varphi(|\alpha(s, x)|_{S_\Lambda}) d\mu(x) &\leq \sum_{n \geq 0} \varphi(a_n) \mu(\{x \in X; |\alpha(s, x)|_{S_\Lambda} \geq a_n\}) \\ &\leq \sum_{n \geq 0} nk(a_n) \\ &\leq \sum_{n \geq 0} 2^{-n} n < \infty, \end{aligned}$$

The same computation shows that for all $s \in S_\Lambda$,

$$\int \varphi(|\beta(s, x)|_{S_\Gamma}) d\mu(x) < \infty.$$

So we are done. \square

We deduce the following quantitative version of Ornstein and Weiss's theorem.

Corollary 4.24. *Let Γ and Λ be infinite finitely generated amenable groups. There exists a concave increasing unbounded function φ satisfying $\varphi(0) = 0$ and $\varphi(t)/\log t$ is non-increasing on $[1, \infty)$ such that there is a (φ, φ) orbit equivalence coupling between them.*

Recall the following theorem from [BZ21].

Theorem 4.25. *Let F be an increasing function such that $F(t)/\log t$ is non-increasing on $[1, \infty)$. Then there exists a finitely generated group Γ whose isoperimetric profile satisfies $j_{1,\Gamma} \approx F$.*

We deduce the following corollary, which is in sharp contrast with the previous corollary.

Corollary 4.26. *For every concave increasing unbounded function ψ such that $\psi(t)/\log t$ is non-increasing on $[1, \infty)$ there exists a finitely generated amenable group Γ with the following property: for every concave function φ such that $\varphi(0) = 0$, if Γ is a φ -integrable measure quotient (or an at most m -to-one measure sub-quotient) of \mathbb{Z} , then $\varphi \preceq \psi$.*

Proof. Consider the function $F(t) = \psi(t)$, then $F(t)/\log t = \psi(t)/\log t$, so it is non-increasing on $[1, +\infty)$. We can then apply Theorem 4.25 and find a finitely generated group Γ whose isoperimetric profile satisfies $j_{1,\Gamma} \approx F$.

Now let φ be a concave non-decreasing function such that $\varphi(0) = 0$, then by concavity we have that $t/\varphi(t)$ is non-decreasing. If there is a φ -integrable measure quotient coupling from Γ to \mathbb{Z} , then we can apply Theorem 4.3, and since $j_{1,\mathbb{Z}}(n) \approx n$ and $j_{1,\Gamma} \approx \psi$ we get that $\varphi \preceq \psi$ as wanted. \square

Proof of Corollary 4. Let Γ and Λ be finitely generated amenable groups, and let ϕ be a positive, unbounded increasing function. Assume that there exists a (ϕ, L^0) measure equivalence coupling from Γ to Λ . Reasoning as in the proof of Corollary 4.24, we can always find $\phi' \leq \phi$ that is increasing, unbounded, concave and such that $\phi'(0) = 0$. Hence we can assume without loss of generality that $\phi = \phi'$. By Corollary 4.24, there exist another concave increasing function θ and a (θ, L^0) measure equivalence coupling from Λ to \mathbb{Z} . Composing these coupling, we obtain by Proposition 2.26 a $(\phi \circ \theta, L^0)$ measure equivalence coupling from Γ to \mathbb{Z} . Note that the two conditions: f is increasing and $t/f(t)$ is non-decreasing are stable under composition. Therefore, to prove Corollary 4 it suffices to apply Corollary 4.26 to the function $\psi = \log \circ \phi \circ \theta$, as it is strictly slower than $\phi \circ \theta$. \square

5 L^∞ measure subgroups

In this section we discuss the notion of at most m -to-one L^∞ -measure subgroups. We shall see that for amenable groups this notion is equivalent to that of regular map, which will allow us to deduce various results stated in the introduction.

5.1 L^∞ measure subgroups and regular maps

The following proposition can be extracted from the proof of [Sha04, Thm. 2.1.2], and provides a useful way of building measured subgroup couplings from Borel sub-quotient couplings.

Proposition 5.1. *Let Ω be a standard Borel space, suppose that we have a $\Gamma \times \Lambda$ -action on Ω with the following properties:*

- *the Λ -action is free and admits a Borel fundamental domain X_Λ ,*
- *there exists a Γ -invariant probability measure μ for the induced action on X_Λ .*

Then μ has a unique Λ -invariant extension to a σ -finite invariant measure m on Ω which is Γ -invariant as well. In particular, if the Γ -action on Ω is free and has a Borel fundamental domain, then Γ is a measure subgroup of Λ .

Proof. Since every Borel subset of Ω is a countable disjoint union of Λ -translates of Borel subsets of X_Λ , the Λ -invariant extension of μ is unique. We build it by letting

$$m(A) = \sum_{\lambda \in \Lambda} \mu((\lambda * A) \cap X_\Lambda).$$

By definition, the measure m is Λ -invariant, let us show that it is Γ -invariant as well. Let $\gamma \in \Gamma$, if A is a Borel subset of X_Λ , by the definition of the induced action on X_Λ we have

$$\gamma \cdot A = \bigsqcup_{\lambda \in \Lambda} (\lambda * \gamma * A) \cap X_\Lambda$$

and so $m(\gamma * A) = \mu(\gamma \cdot A) = \mu(A) = m(A)$. Now for an arbitrary $\lambda \in \Lambda$ and a Borel subset $A \subseteq X_\Lambda$, we have

$$m(\gamma * (\lambda * A)) = m(\lambda * (\gamma * A)) = m(\gamma * A) = m(A) = m(\lambda * A),$$

and since every Borel subset of Ω is a countable disjoint union of Λ -translates of Borel subsets of X_Λ , we conclude that the Γ -action on Ω preserves m . \square

Recall the following definition from [BST12].

Definition 5.2. Let Γ and Λ be two countable finitely generated groups, a map $f: \Gamma \rightarrow \Lambda$ is a **regular map** if it is Lipschitz and there exists $m \in \mathbb{N}$ such that the preimages of f have size at most m for some $m \in \mathbb{N}$. When there exists such a map, we say that Γ regularly embeds in Λ .

Remark 5.3. Coarse embeddings are special cases of regular maps.

The second part of the following theorem is a slightly generalized version of [Sha04, Thm. 2.1.2] (which was proved for coarse embeddings).

Theorem 5.4. *Let Γ and Λ be finitely generated groups.*

- *Assume that Γ is an at most m -to-one L^∞ measure subgroup of Λ . Then Γ regularly embeds into Λ .*
- *Conversely, if Γ regularly embeds into Λ and Γ is amenable, then there exists $m \in \mathbb{N}$ such that Γ is an at most m -to-one L^∞ measure subgroup of Λ .*

We deduce from the first statement of Theorem 5.4 and the fact that asymptotic dimension is monotonous under regular map (see [BST12]) the following corollary.

Corollary 5.5. *The asymptotic dimension is monotonous under taking at most m -to-one L^∞ measure subgroup.* \square

And from the second part of the theorem, we deduce the following result, which we announced in the introduction.

Corollary 5.6. *For every $1 \leq p \leq \infty$, the isoperimetric profile is monotonous under regular map between amenable groups.*

Proof. Let Γ and Λ be finitely generated amenable groups. By the previous theorem, the existence of a regular map from Γ to Λ implies that of an at most m -to-one L^∞ measure subgroup coupling from Γ to Λ . Such a coupling is in particular L^1 , so the result follows from the second item in Theorem 4.2. \square

The rest of this subsection is dedicated to the proof of the theorem.

Proof of Theorem 5.4. To prove the first statement, we let (Ω, X_Λ, μ) be an m -to-one L^∞ subgroup coupling from Γ to Λ , and consider the associated cocycle $c: \Gamma \times X_\Lambda \rightarrow \Lambda$. Let $K \geq 0$ be such that $|c(s, x)|_{S_\Lambda} \leq K$ for all $s \in S_\Gamma$ and a.e. $x \in X$. The assumptions imply that for a.e. $x \in X_\Lambda$, the map $c(\cdot, x): \Gamma \rightarrow \Lambda$ is at most m -to-one and K -Lipschitz, hence is a regular map.

We now prove the second statement of the theorem. We let $K \geq 1$ and $m \in \mathbb{N}$ such that there is an m -to-one K -Lipschitz maps $\Gamma \rightarrow \Lambda$. We then let F be a finite set of cardinality m , and denote by $\pi_\Lambda: \Lambda \times F \rightarrow \Lambda$ the projection on the first coordinate. Our coupling space is

$$\Omega := \{f: \Gamma \rightarrow \Lambda \times F: f \text{ is injective and } \pi_\Lambda \circ f \text{ is } K\text{-Lipschitz}\}$$

Note that since F has cardinality m , every at most m -to-one map $\Gamma \rightarrow \Lambda$ can be lifted to an injective map $\Gamma \rightarrow \Lambda \times F$, so Ω is not empty. The Lipschitz condition and the fact that balls in Λ are finite ensures that Ω is locally compact for the product topology. As before we have a $\Gamma \times \Lambda$ -action given by $(\gamma, \lambda) \cdot f(g) = \lambda f(\gamma^{-1}g)$, and a compact fundamental for the Λ -action is given by $X_\Lambda = \{f \in \Omega: \pi_\Lambda(f(1_\Gamma)) = 1_\Lambda\}$. A Borel fundamental domain X_Γ for the Γ -action can be obtain as follows: we fix a well-order $<$ on Λ , and let X_Γ be the set of functions f which attain their $<$ -minimum at 1_Γ .

The cocycle $c: \Gamma \times X_\Lambda \rightarrow \Lambda$ is given by $c(\gamma, f) = \pi_\Lambda(f(\gamma^{-1}))^{-1}$, and the injectivity condition implies that Γ acts freely. Finally for each $\gamma \in \Gamma$, the Lipschitz condition implies that $c(\gamma, \cdot)$ is bounded, and since Γ is amenable we may find a Γ -invariant measure on X_Λ which we extend via Proposition 5.1 in order to get the desired L^∞ subgroup coupling. \square

5.2 A continuum of 3-solvable groups

In this section we prove the following result, announced in the introduction.

Theorem 5.7. *There exists a uncountable family of groups Γ_i , such that*

- (i) $\Gamma_i = N_i \rtimes \mathbb{Z}$, where N_i is locally finite, and 2-step nilpotent;
- (ii) for any $i \neq j$ and any $m \geq 1$, Γ_i is not an at most m -to-one L^1 -measure sub-quotient (nor an L^1 -measure quotient) of Γ_j .

Note that (i) implies in that Γ_i is 3-step solvable and has asymptotic dimension 1. We deduce the result announced in the introduction.

Corollary 5.8. *There exists a uncountable family of groups Γ_i , such that*

- (i) $\Gamma_i = N_i \rtimes \mathbb{Z}$, where N_i is locally finite, and 2-step nilpotent;
- (ii) for any $i \neq j$ and any $m \geq 1$, Γ_i does not regularly embed into Γ_j .

Proof of Theorem 5.7. To prove the theorem, we shall use [EZ17, Corollary 3.3] which is phrased in terms of Følner function. Since the Følner function is equal to to the generalized inverse of the L^1 -isoperimetric profile, i.e. $F\phi_\Gamma(k) = \inf \{n: j_{1,\Gamma}(n) \geq k\}$, Corollary 5.6 implies that it is monotonous under regular map between amenable groups.

Now by [EZ17, Corollary 3.3], for any non-decreasing function $\tau: [1, \infty) \rightarrow [1, \infty)$ such that $\tau(n+1) - \tau(n) \leq n$ and $\tau(n) \geq n$, there exists a group as in (i) whose Følner function is asymptotically equivalent to $\exp(\tau(n))$. Let us now explain how to get uncountably many asymptotically incomparable such functions.

Claim 5.9. For every $1 < a < b < 2$, there is a non-decreasing function $\tau_{a,b}: \mathbb{N} \rightarrow \mathbb{N}$ such that for all $n \in \mathbb{N}$ we have

$$\tau_{a,b}(n) \geq n \quad \text{and} \quad \tau_{a,b}(n+1) - \tau_{a,b}(n) \leq n, \tag{*}$$

and the following three conditions are met:

- (1) for all but finitely many $n \in \mathbb{N}$ we have $n^a \leq \tau_{a,b}(n) \leq n^b$;
- (2) there are infinitely many $n \in \mathbb{N}$ such that $\tau_{a,b}(n) = n^b$;
- (3) the set of $n \in \mathbb{N}$ such that $\tau_{a,b}(n) = n^a$ contains arbitrarily large intervals.

Moreover, for every $1 < a' < a < b < b' < 2$, the functions $\exp(\tau_{a,b})$ and $\exp(\tau_{a',b'})$ are asymptotically incomparable.

Proof of the claim. Let us start by proving the existence of $\tau_{a,b}$ as above for $1 < a < b < 2$. First, we fix $N \in \mathbb{N}$ such that for all $n \geq N$, we have $(n+1)^b - n^b \leq n$, so in particular $(n+1)^a - n^a \leq n$. Since $b < 2$, we may and do fix $N_0 > N$ such that

$$\sum_{n=0}^{N_0-1} n \geq N_0^b.$$

We then define $\tau_{a,b}$ by induction on the interval $[0, N_0]$ by $\tau_{a,b}(0) = 0$ and for all $n < N_0$,

$$\tau_{a,b}(n+1) = \min(\tau_{a,b}(n) + n, N_0^b).$$

Observe that by the previous inequality $\tau_{a,b}(N_0) = N_0^b$.

Suppose now by induction $\tau_{a,b}$ has been defined on an interval $[0, N_k]$, satisfies (*) and satisfies $\tau_{a,b}(N_k) = N_k^b$, then it suffices to explain how to extend its definition to a bigger interval $[0, N_{k+1}]$ so that it still satisfies (*) and

- (1') for all $n \in (N_k, N_{k+1}]$ we have $n^a \leq \tau_{a,b}(n) \leq n^b$;
- (2') $\tau_{a,b}(N_{k+1}) = N_{k+1}^b$;
- (3') there is an interval I_k of size k such that for all $n \in I_k$, $\tau_{a,b}(n) = n^a$.

In order to do so, let M_k be the first integer such that $(M_k)^a \geq N_k^b$. We then let $\tau_{a,b}(n) = N_k^b$ for all $n \in (N_k, M_k)$, and then for all $n \in [M_k, M_k + k]$ we let $\tau_{a,b}(n) = n^a$, which takes care of condition (3').

Then, since $b < 2$, we may and do define N_{k+1} as the least integer such that

$$\tau_{a,b}(M_k + k) + \sum_{n=M_k+k}^{N_{k+1}-1} n \geq N_{k+1}^b.$$

We now define by induction on $n \in [M_k + k, N_{k+1})$

$$\tau_{a,b}(n+1) = \min(n + \tau_{a,b}(n), n^b),$$

which by the previous inequality guarantees that $\tau_{a,b}(N_{k+1}) = N_{k+1}^b$, so (2') is satisfied. Finally, since for all $n \geq N_k$, we have $(n+1)^a - n^a \leq n$, an inspection of the definition shows that condition (*) is still satisfied, and (1') is clearly satisfied. This finishes the construction of the desired $\tau_{a,b}$.

We now prove the incomparability statement. First note that we cannot have $\exp(\tau_{a',b'})$ asymptotically bounded by $\exp(\tau_{a,b})$ since we have $\tau_{a',b'}(n) = n^{b'}$ for infinitely many $n \in \mathbb{N}$ but $\tau_{a,b}(n) \leq n^b$ for all but finitely many $n \in \mathbb{N}$ and $b < b'$.

Conversely, suppose that there is $C \in \mathbb{N}$ such that for all $n \in \mathbb{N}$, we have $\exp(\tau_{a,b}(n)) \leq C \exp(\tau_{a',b'}(Cn))$. Note that the set $\{Cn : n \in \mathbb{N}\}$ has to intersect infinitely many times any set which contains arbitrarily large intervals. Since $\tau_{a',b'}$ satisfies condition (3) we thus find infinitely many $n \in \mathbb{N}$ such that $\tau_{a',b'}(Cn) = (Cn)^{a'}$. On the other hand for all but finitely many $n \in \mathbb{N}$ we have $\tau_{a,b}(n) \geq n^a$. We conclude that there are infinitely many $n \in \mathbb{N}$ such that

$$\exp(n^a) \leq C \exp((Cn)^{a'}),$$

a contradiction. □_{claim}

We now fix for every $1 < a < b$ a function $\tau_{a,b}$ as above. Then it satisfies the assumption of [EZ17, Corollary 3.3], so there is a group $\Gamma_{a,b}$ satisfying (i) whose Følner function is asymptotically equivalent to $\exp(\tau_{a,b})$.

Now, as explained right before the previous claim, the Følner function is monotonous under regular embeddings. So by the claim for all $1 < a' < a < b < b' < 2$, the groups $\Gamma_{a,b}$ and $\Gamma_{a',b'}$ are not regularly embeddable one another. Therefore the family $(\Gamma_{1+\varepsilon, 2-\varepsilon})_{0 < \varepsilon < 1/2}$ is the uncountable family we were looking for. \square

5.3 Stability of Liouville property for lamplighter groups

The Liouville property is now completely understood for lamplighter groups.

Theorem 5.10. *Let F be a non-trivial finite group, and let H be a finitely generated group. Let μ be a probability measure on $\Gamma = F \wr H$ whose support is finite and generates Γ , and let $\bar{\mu}$ be the projected probability on H . Then (Γ, μ) is Liouville if and only if $(H, \bar{\mu})$ is recurrent.*

This result is due to Kaimanovich [V.A85] when F is abelian, and to Lyons and Peres in the general case [LY15]. By a celebrated result of Varopoulos, if μ is an admissible probability measure on G , (G, μ) is recurrent if and only if G has at most quadratic volume growth. Therefore, we deduce the following corollary.

Corollary 5.11. *Let F be a non-trivial finite group, and let H be a finitely generated group. Let μ be an admissible probability measure on $\Gamma = F \wr H$. Then (Γ, μ) is Liouville if and only if H has at most quadratic volume growth.*

We immediately deduce that for lamplighter groups, the Liouville property is independent on the (admissible) measure μ . We now state the main result of this subsection.

Theorem 5.12. *For $i = 1, 2$, let F_i be a non-trivial finite group, and let H_i be a finitely generated group. Let μ_i be a probability measure on $\Gamma_i = F_i \wr H_i$ whose support is finite and generates Γ_i . Assume that Γ_1 is either an at most m -to-one L^1 measure sub-quotient, or an L^1 measure quotient of Γ_2 . Then if (Γ_2, μ_2) is Liouville, then so is (Γ_1, μ_1) .*

A well-known open question asked by Benjamini in [Ben12] is whether Liouville property is “invariant under quasi-isometry”. We suggest the following variant.

Question 5.13. Let G_1 and G_2 be finitely generated groups equipped with admissible probability measures μ_1 and μ_2 . Assume that (G_2, μ_2) is Liouville and that G_1 regularly embeds into G_2 . Must (G_1, μ_1) be also Liouville?

This question is motivated by the following immediate corollary of Theorem 5.12 and Theorem 5.4.

Corollary 5.14. *For $i = 1, 2$, let F_i be a non-trivial finite group, and let H_i be a finitely generated group. Let μ_i be a probability measure on $\Gamma_i = F_i \wr H_i$ whose support is finite and generates Γ_i . Assume that Γ_1 regularly embeds into Γ_2 . Then if (Γ_2, μ_2) is Liouville, then so is (Γ_1, μ_1) .*

Proof of Theorem 5.12. By [Ers03], the Følner function of Γ_i satisfies

$$\text{Føl}_{\Gamma_i}(n) \approx e^{\text{Føl}_{H_i}(n)}.$$

On the other hand, by Remark 4.4, we have

$$\text{Føl}_{\Gamma_1} \preceq \text{Føl}_{\Gamma_2},$$

from which we deduce that

$$\text{Føl}_{H_1} \preceq \text{Føl}_{H_2}.$$

By Corollary 5.11, we know that Γ_2 has at most quadratic growth. Hence by [Cou00], we have $\text{Føl}_{H_2} \preceq n^2$, and therefore $\text{Føl}_{H_1} \preceq n^2$. Hence by [Cou00], H_1 has at most quadratic growth and so is recurrent. So we conclude thanks to Corollary 5.11. \square

6 Følner tilings

6.1 Følner tiling sequences and orbit equivalence

Definition 6.1. Let Γ be an amenable group and (F_k) be a sequence of finite subsets of Γ . We call (F_k) a (left) **Følner tiling sequence** if the sequence of *tiles* (T_k) defined inductively by $T_0 = F_0$ and $T_{k+1} = T_k F_{k+1}$ satisfies the following conditions:

- (i) (tiling condition) for all $k \in \mathbb{N}$, we can write T_{k+1} as a *disjoint* union:

$$T_{k+1} = \bigsqcup_{\gamma \in F_{k+1}} T_k \gamma;$$

- (ii) (Følner condition) (T_k) is a left Følner sequence: for all $\gamma \in \Gamma$, $\lim_{k \rightarrow +\infty} \frac{|\gamma T_k \setminus T_k|}{|T_k|} = 0$.

If in addition there exists a decreasing sequence of finite index subgroups Γ_k such that the F_k 's are left coset representatives of Γ_{k-1} modulo Γ_k , then we call (F_k) a **profinite Følner tiling sequence** associated to (Γ_k) .

Remark 6.2. The first condition amounts to saying that every element of T_k can uniquely be written as $f_0 \cdots f_k$ where each f_i belongs to F_i .

Remark 6.3. In some examples it will be more convenient to consider *right* Følner tiling sequences (F_k) , i.e. sequences such that (F_k^{-1}) is a left Følner tiling sequence. Nevertheless, every Følner tiling sequence will be a *left* Følner tiling sequence unless specified otherwise.

To every Følner tiling sequence (F_k) , we associate a measure-preserving Γ -action constructed as follows. We consider the standard Borel probability space $(X = \prod_k F_k, \mu)$, where each factor is equipped with the normalized counting measure, and μ is the product measure. Each element $x = (x_k)_{k \in \mathbb{N}}$ of X defines a sequence $(g_k(x))_{k \in \mathbb{N}}$ of elements of Γ given by

$$g_k(x) = x_0 \cdots x_k \in T_k$$

Observe that by condition (i), each g_k is an equidistributed random element of T_k . In other words, the map g_n defines an isomorphism from the finite product $(X_n = \prod_{k=0}^n F_k, \mu_n)$ to T_n equipped with the renormalized counting measure.

Since (T_k) is a left Følner sequence, we deduce that for every $\gamma \in \Gamma$ and almost every $x \in X$, there is n such that $\gamma g_n(x) \in T_n$. We can then write uniquely $\gamma g_n(x) = x'_0 \cdots x'_n$ where $x'_i \in F_i$, and we then define

$$\gamma \cdot (x_k)_{k \in \mathbb{N}} = (x'_0, \dots, x'_n, x_{n+1}, x_{n+2}, \dots).$$

Note that this does not depend on the choice of n because we have $\gamma g_{n+1}(x) = \gamma g_n(x) x_{n+1} \in T_{n+1}$.

We claim that up to measure zero, this group action induces the equivalence relation of equality up to a finite set of indices, also called the **cofinite equivalence relation** E_{cof} . In particular the action is measure preserving: indeed the cofinite equivalence relation can be realized through the natural action of the direct sum $\bigoplus_k \mathfrak{S}(F_k)$, which is obviously measure preserving.

Indeed, we have just seen that for almost every $x \in X$ and all $\gamma \in \Gamma$, x and $\gamma \cdot x$ are equal up to a finite set of indices. Conversely, if x and x' are such that $x_j = x'_j$ for all $j \geq k+1$, the element $\gamma = g_k(x) g_k(x')^{-1} \in G$ satisfies $\gamma g_k(x) = g_k(x') \in T_k$, and hence $\gamma \cdot x = x'$.

For instance, if $\Gamma = \mathbb{Z}$, $F_k = \{0, 2^k\}$, the tiles are $T_k = \{0, \dots, 2^{k+1} - 1\}$ and for this example we get the usual odometer, up to renaming each F_k as $\{0, 1\}$. More generally, when the tiles T_k are left coset representatives for finite index subgroups Γ_k of Γ , and if $F_k \subset \Gamma_{k-1}$, then we get the profinite action $\Gamma \curvearrowright \text{proj lim } \Gamma/\Gamma_k$. We deduce from this discussion the following proposition.

Proposition 6.4. *Assume that (F_k) is a Følner tiling sequence for Γ . Then Γ has a measure-preserving action on the infinite product probability space $(X = \prod_k F_k, \mu)$, which almost surely generates the co-finite equivalence relation on this product. Moreover if (F_k) is a profinite Følner tiling sequence associated to (Γ_k) , then this action is isomorphic to the profinite action of Γ on $\varprojlim \Gamma/\Gamma_k$.*

Consider the following (possibly infinite) measurable distance on X given by

$$\rho(x, x') = \inf\{n \in \mathbb{N} : \forall k \geq n, x_k = x'_k\}$$

Observe that x and x' are equal up to a finite set of indices if and only if $\rho(x, x') < +\infty$. Also, by the definition of our action, for every $\gamma \in \Gamma$ and almost every $x \in X$ we have that $\rho(\gamma \cdot x, x) > k$ if and only if $\gamma g_k(x) \notin T_k$. In particular,

$$\mu(\{x, \rho(\gamma \cdot x, x) > k\}) = \frac{|T_k \setminus \gamma^{-1}T_k|}{|T_k|} = \frac{|\gamma T_k \Delta T_k|}{2|T_k|}. \quad (17)$$

In order to obtain quantitative statements, we introduce the following parameters.

Definition 6.5. Let (ε_k) be a sequence of strictly positive numbers tending to 0, and (R_k) be a sequence of positive reals. Say that a Følner tiling sequence (F_k) of a finitely generated group Γ equipped with a finite generating set S_Γ is an (ε_k, R_k) -**Følner tiling sequence** if

- (i) each tile T_k has d_{S_Γ} -diameter at most R_k ,
- (ii) every $s \in S_\Gamma$ satisfies $|T_k \setminus sT_k| \leq \varepsilon_k |T_k|$.

We shall need the following key observation.

Lemma 6.6. *Let (F_k) be (ε_k, R_k) -Følner tiling sequence of a finitely generated group Γ equipped with a finite generating set S_Γ . Then*

- (i) for all $s \in S_\Gamma$, we have for all $k \geq 0$ that $\mu(\{x \in X : \rho(s \cdot x, x) > k\}) \leq \varepsilon_k$;
- (ii) for almost every $x \in X$, if $|\gamma|_{S_\Gamma} > 2R_k$, then $\rho(\gamma \cdot x, x) > k$.

Proof. The first item follows from equation (17), and the second item by its first part since if $|\gamma|_{S_\Gamma} > 2R_k$, then $\gamma T_k \cap T_k = \emptyset$ as $\text{diam}(T_k) \leq R_k$. \square

Observe that if two amenable groups Γ and Γ' admit respective Følner tiling sequences (F_k) and (F'_k) such that $|F_k| = |F'_k|$, then there is a natural measure-preserving bijection Ψ between the corresponding product spaces (X, μ) and (X', μ') that preserves the cofinite equivalence relation, and Proposition 6.4 now provides us with an explicit orbit equivalence between them. In what follows, we identify X and X' via Ψ . The following proposition relates the parameters of the Følner tiling sequences with the integrability of the cocycles.

Proposition 6.7. *Suppose that (F_k) is an (ε_k, R_k) Følner tiling sequences for Γ and (F'_k) is an (ε'_k, R'_k) Følner tiling sequence for Γ' . Then for all $s \in S_\Gamma$ and all $k \geq 0$,*

$$\mu(\{x \in X : d_{S_{\Gamma'}}(x, s \cdot x) > 2R'_k\}) \leq \varepsilon_k$$

In particular, the groups are orbit equivalent, and if $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a non-decreasing function such that the sequence $(\varphi(2R'_k)(\varepsilon_{k-1} - \varepsilon_k))_{k \in \mathbb{N}}$ is summable, then for all $s \in S_\Gamma$

$$\int_X \varphi(d_{S_{\Gamma'}}(x, s \cdot x)) d\mu(x) < \infty.$$

Proof. The first part follows from Lemma 6.6 as $|F_k| = |F'_k|$ for all $k \in \mathbb{N}$ and

$$\mu(\{x \in X : d_{S_\Gamma}(x, s \cdot x) > 2R'_k\}) \leq \mu(\{x \in X : \rho(s \cdot x, x) > k\}) \leq \varepsilon_k.$$

To deduce the second part, we exploit the fact that ϕ is increasing: for $s \in S_\Gamma$ we have that

$$\begin{aligned} \int_X \varphi(d_{S_\Gamma}(x, s \cdot x)) d\mu(x) &\leq \varphi(2R'_0) + \sum_{k=1}^{\infty} \varphi(2R'_k) \mu(\{x \in X : 2R'_{k-1} < d_{S_\Gamma}(x, s \cdot x) \leq 2R'_k\}) \\ &\leq \varphi(R'_0) + \sum_{k=1}^{\infty} \varphi(2R'_k) (\varepsilon_{k-1} - \varepsilon_k), \end{aligned}$$

which is finite by assumption. \square

6.2 Applications to groups with polynomial growth

We start applying Proposition 6.7 to torsion-free abelian groups.

Proposition 6.8. *Let n be a positive integer. The group \mathbb{Z}^n (equipped with its standard generating set) admits a profinite (ε_k, R_k) -Følner tiling sequence (F_k) , with $|F_k| = 2^n$, $R_k = n2^{k+1}$ and $\varepsilon_k = 2^{-(k+1)}$ for any $k \geq 0$.*

Proof. We let $F_k = \{0, 2^k\}^n$ for any $k \geq 0$. One can check that $T_k = \{0, 1, \dots, 2^{k+1} - 1\}^n$, which is a coset representative for the finite index subgroup $\Gamma_k = (2^{k+1}\mathbb{Z})^n$. The diameter of T_k is bounded by $n2^{k+1}$ and its size equals $2^{n(k+1)}$. Finally take s a generator of \mathbb{Z}^n . Without loss of generality, we can assume that s is the first basis vector in \mathbb{Z}^n . Then, we have

$$T_k \setminus ((1, 0, \dots, 0) + T_k) = \{0\} \times \{0, 1, \dots, 2^{k+1} - 1\}^{n-1},$$

whose cardinality is 2^{k+1} smaller than that of T_k , so we are done. \square

Corollary 6.9. *Let n and m be positive integers. The group \mathbb{Z}^n (equipped with its standard generating set) admits a (ε_k, R_k) -Følner tiling sequence (F'_k) , with $|F'_k| = 2^{nm}$, $R_k = n2^{m(k+1)}$ and $\varepsilon_k = 2^{-m(k+1)}$ for any $k \geq 0$.*

Proof. Let $(F_k)_k$ be the Følner tiling sequence given in Proposition 6.8 and for any $k \geq 0$ let $F'_k = F_{mk}F_{m(k+1)} \dots F_{m(k+m-1)}$. Note that $F'_k = \{0, 2^{mk}, 2 \cdot 2^{mk}, \dots, (2^m - 1)2^{mk}\}^n$ and $T'_k = \{0, 1, \dots, 2^{mk+m} - 1\}^n$.

As T'_k is the set $T_{m(k+m-1)}$ from Proposition 6.8 we have that the diameter of T'_k is at most $n2^{mk+m}$ and the set $T'_k \setminus (s + T'_k)$ has cardinality at most $2^{-mk-m}|T'_k|$ for any standard generator s of \mathbb{Z}^n . \square

The following theorem is almost sharp since by our extensions of Bowen's theorem (Theorem 3.1), if $n < m$ and $p > \frac{n}{m}$, then there cannot be an L^p measure sub-quotient coupling from \mathbb{Z}^m to \mathbb{Z}^n .

Theorem 6.10. *For every n and m , there exists an orbit equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n which is for every $\varepsilon > 0$ a $(\varphi_\varepsilon, \psi_\varepsilon)$ -integrable coupling, where*

$$\varphi_\varepsilon(x) = \frac{x^{n/m}}{\log(x)^{1+\varepsilon}} \text{ and } \psi_\varepsilon(x) = \frac{x^{m/n}}{\log(x)^{1+\varepsilon}}.$$

In particular if $n < m$, then there is an $(L^p, L^{p'})$ orbit equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n for all $p < \frac{n}{m}$ and $p' < \frac{m}{n}$.

Proof. Applying Corollary 6.9, we find a (ε_k, R_k) -Følner tiling sequence $(F_k)_k$ for \mathbb{Z}^m with $|F_k| = 2^{nm}$, $R_k = m2^{n(k+1)}$ and $\varepsilon_k = 2^{-n(k+1)+1}$. Similarly, we find a (ε'_k, R'_k) -Følner tiling sequence $(F'_k)_k$ for \mathbb{Z}^n with $|F'_k| = 2^{nm}$, $R'_k = n2^{m(k+1)}$ and $\varepsilon'_k = 2^{-m(k+1)+1}$. Note that $\varphi_\varepsilon(R'_k)\varepsilon_{k-1} = O(k^{-(1+\varepsilon)})$ and $\psi_\varepsilon(R_k)\varepsilon'_{k-1} = O(k^{-(1+\varepsilon)})$. Thus, by Proposition 6.7, we obtain an orbit equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n which is $(\varphi_\varepsilon, \psi_\varepsilon)$ -integrable. Finally if $n < m$ then for all $p < n/m$ we both have that $x^p = o(\varphi_\varepsilon(x))$ and $x^{p'} = o(\psi_\varepsilon(x))$ as $x \rightarrow +\infty$ so our coupling is also $(L^p, L^{p'})$. \square

Remark 6.11. The expert reader will recognize in the above proof an explicit orbit equivalence between the dyadic \mathbb{Z}^n and \mathbb{Z}^m odometers. Moreover, using this point of view it can be shown that this coupling is not ψ_0 -integrable, so we ask the following refinements of question 1.4.

Question 6.12. Let $n < m$, is there a (φ, L^∞) measure equivalence coupling from \mathbb{Z}^m to \mathbb{Z}^n , where $\varphi(x) = \frac{x^{n/m}}{\log(x)}$? What about a (φ, L^0) measure equivalence coupling?

Recall that the Heisenberg group is the 2-step torsion-free nilpotent group that can be defined as the group of triples $(x, y, z) \in \mathbb{Z}^3$ equipped with the group operation

$$(x, y, z) \cdot (x', y', z') = (x + x', y + y', z + z' + yx'),$$

which comes from its identification with the group of matrices of the form $\begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ z & y & 1 \end{bmatrix}$. We equip it with the standard generating set $S = \{E_1^{\pm 1}, E_2^{\pm 1}\}$, where $E_1 = (1, 0, 0)$ and $E_2 = (0, 1, 0)$.

Proposition 6.13. *The Heisenberg group admits a profinite (ε_k, R_k) -Følner tiling (F_k) such that $|F_k| = 16$, $R_k = 10 \cdot 2^{k+2}$, and $\varepsilon_k = 2^{-k}$ for any $k \geq 0$.*

Proof. For every $k \geq 0$ let

$$F_k = \{(2^k x, 2^k y, 4^k z) : x, y \in \{0, 1\}, z \in \{0, 1, 2, 3\}\}.$$

We claim that (F_k) is the profinite left Følner tiling we are looking for.

First note that our sequence is *profinite* because the set F_k is a right coset representative of Γ_{k-1} modulo Γ_k , where Γ_{k-1} is the (non normal) finite index subgroup

$$\Gamma_{k-1} = \{(x, y, z) : x \equiv y \equiv 0 \pmod{2^k}, z \equiv 0 \pmod{4^k}\}.$$

We then check that (F_k) satisfies the *tiling condition*. Let $A = (x_A, y_A, z_A) \in T_k$. By our definition of T_k , we can write A as

$$A = (x_0, y_0, z_0)(2x_1, 2y_1, 4z_1) \cdots (2^k x_k, 2^k y_k, 4^k z_k) \quad (18)$$

with $x_i, y_i \in \{0, 1\}$, and $z_i \in \{0, 1, 2, 3\}$, which yields

$$x_A = \sum_{i=0}^k 2^i x_i, \quad y_A = \sum_{i=0}^k 2^i y_i, \quad \text{and} \quad z_A = \sum_{i=0}^k 4^i z_i + \sum_{i=1}^k 2^i x_i \sum_{j=0}^{i-1} 2^j y_j \quad (19)$$

We thus see that we can recover the finite sequences $(x_i)_{i=0}^k$ and $(y_i)_{i=0}^k$ from the coefficients (x_A, y_A, z_A) of A as the binary expansions of x_A and y_A respectively, and then similarly $((z_i)_{i=0}^k)$ is also completely determined by (x_A, y_A, z_A) . Therefore the above decomposition of A as an element of $F_0 \cdots F_k$ is unique for every element in T_k .

We now check that the *diameters of tiles* are as claimed. Let $B \in F_k$, we have $B = (2^k x, 2^k y, 4^k z)$ for some $x, y \in \{0, 1\}$, $z \in \{0, 1, 2, 3\}$ and a straightforward computation shows $B = E_1^{2^k x} E_2^{2^k y} [E_1^{2^k}, E_2^{2^k z}]$. We thus have that F_k is contained in the ball of radius $10 \cdot 2^k$ around the identity. Hence T_k is contained in a ball of radius $10 \cdot 2^{k+1}$, and so has diameter at most $10 \cdot 2^{k+2}$ as claimed.

We finally check the *quantitative Følner condition*. We first estimate $|T_k \setminus E_1 T_k|$. To do so, let us consider an elements $A \in T_k$ such that $E_1 A \in T_k$. We decompose A according to (18):

$$A = (x_0, y_0, z_0)(2x_1, 2y_1, 4z_1) \cdots (2^k x_k, 2^k y_k, 4^k z_k) = (x_A, y_A, z_A).$$

We have $E_1A = (x_A + 1, y_A, z_A)$, so if the latter belongs to T_k , then by equation (19) one of the x_i 's must vanish. We let m be the least i such that $x_i = 0$ and we will now count the number of possible A 's for a given $m \in \{0, \dots, k\}$.

First note that there are 2^{k-m} possibilities for x_0, \dots, x_k since we must have $x_0 = \dots = x_{m-1} = 1$ and $x_m = 0$. There are 2^{k+1} possibilities for y_0, \dots, y_k , and we now have to understand precisely what are the conditions on the z_i 's: we will bound from below the number of possible parameters $z_0, \dots, z_k \in \{0, 1, 2, 3\}$ such that the element

$$A = (x_0, y_0, z_0)(2x_1, 2y_1, 4z_1) \cdots (2^k x_k, 2^k y_k, 4^k z_k) = (x_A, y_A, z_A)$$

satisfies $E_1A \in T_k$. We have $E_1A = (x_A + 1, y_A, z_A)$, so if the latter belongs to T_k , then by equation (19) its decomposition

$$E_1A = (x'_0, y'_0, z'_0)(2x'_1, 2y'_1, 4z'_1) \cdots (2^k x'_k, 2^k y'_k, 4^k z'_k)$$

is given by $x'_i = 0$ for $i < m$, $x'_m = 1$ and $x'_i = x_i$ for $i > m$, $y'_i = y_i$ for all $i \in \{0, \dots, k\}$, and finally the z'_i 's are subject to the equation

$$\sum_{i=0}^k 4^i z'_i + \sum_{i=1}^k 2^i x'_i \sum_{j=0}^{i-1} 2^j y_j = \sum_{i=0}^k 4^i z_i + \sum_{i=1}^k 2^i x_i \sum_{j=0}^{i-1} 2^j y_j.$$

We thus have:

$$\sum_{i=0}^k 4^i z'_i = \sum_{i=0}^k 4^i z_i + \sum_{i=1}^k 2^i (x_i - x'_i) \sum_{j=0}^{i-1} 2^j y_j, \quad (20)$$

Now recall that $x'_i = 0$ for $i < m$, $x'_m = 1$ and $x'_i = x_i$ for $i > m$, so we can rewrite the last term as

$$\sum_{i=1}^k 2^i (x_i - x'_i) \sum_{j=0}^{i-1} 2^j y_j = \sum_{i=1}^m 2^i \sum_{j=0}^{i-1} 2^j y_j - 2^m \sum_{j=0}^{m-1} 2^j y_j$$

We deduce that $\sum_{i=1}^k 2^i (x_i - x'_i) \sum_{j=0}^{i-1} 2^j y_j$ is negative, and its absolute value is strictly less than 4^m . Hence (20) has a solution, and thus $E_1A \in T_k$, as soon as $\sum_{i=0}^k 4^i z_i \geq 4^m$. Taking the contrapositive, we see that there are at most 4^m possibilities for the sequence $(z_i)_{i=0}^k$ so that $E_1A \notin T_k$.

We conclude that for a fixed m , we have 2^{k-m} possible x_A 's, 2^{k+1} possible y_A 's and at most 4^m possible z_A 's such that $E_1A \notin T_k$, which yields a total of $2^{k-m} 2^{k+1} 4^m = 2^{2k+1} 2^m$ possibilities. When $x_i = 1$ for every i , we never have $E_1A \in T_k$, and then this adds $2^{k+1} 4^{k+1} = 2^{3k+3}$ possibilities.

We conclude that there are at most $2^{3k+3} + \sum_{m=0}^k 2^{2k+1} 2^m < 2^{3k+4}$ choices of A such that $E_1A \notin T_k$,

which means that $|E_1T_k \setminus T_k| < 2^{3k+4}$.

To estimate the cardinality of $E_2T_k \setminus T_k$, we proceed similarly. We consider $A \in T_k$ such that $E_2A \in T_k$. We let m be such that $y_0 = \dots = y_{m-1} = 1$, $y_m = 0$. Similarly as before, if $E_2A \in T_k$ we may write it as

$$E_2A = (x_A, y_A + 1, x_A + z_A) = (x'_0, y'_0, z'_0)(2x'_1, 2y'_1, 4z'_1) \cdots (2^k x'_k, 2^k y'_k, 4^k z'_k).$$

So by equation (19) for all $i \in \{0, \dots, k\}$ we have $x'_i = x_i$, $y'_0 = \dots = y'_{m-1} = 0$, $y'_m = 1$ and $y'_i = y_i$ for all $i > m$. Finally the z'_i 's satisfy

$$\sum_{i=0}^k 4^i z'_i + \sum_{i=1}^k 2^i x_i \sum_{j=0}^{i-1} 2^j y'_j = \sum_{i=0}^k 2^i x_i + \sum_{i=0}^k 4^i z_i + \sum_{i=1}^k 2^i x_i \sum_{j=0}^{i-1} 2^j y_j.$$

We may rewrite our previous equation as

$$\sum_{i=0}^k 4^i z'_i - \sum_{i=0}^k 4^i z_i = \sum_{i=1}^k 2^i x_i \sum_{j=0}^{i-1} 2^j (y_j - y'_j) + \sum_{i=0}^k 2^i x_i$$

We then decompose the first sum in the right term, noting that for all $i > m$, $\sum_{j=0}^{i-1} 2^j (y_j - y'_j) = -1$:

$$\begin{aligned} \sum_{i=0}^k 4^i z'_i - \sum_{i=0}^k 4^i z_i &= \sum_{i=1}^m 2^i x_i \sum_{j=0}^{i-1} 2^j - \sum_{i=m+1}^k 2^i x_i + \sum_{i=0}^k 2^i x_i \\ &= \sum_{i=1}^m 2^i x_i (2^i - 1) + \sum_{i=0}^m 2^i x_i \\ &= \sum_{i=1}^m 4^i x_i + x_0 \\ &< 2^{2m+1} \end{aligned}$$

So for these x_A and y_A there exists at most 2^{2m+1} values of z_A such that $E_2 A \notin T_k$ (namely those such that $\sum_{i=0}^k 4^i z_i + 2^{2m+1} \geq 4^{k+1}$) and as before we conclude that there exists at most

$$2^{k+1} 4^{k+1} + \sum_{m=0}^k 2^{k+1} 2^{k-m} 2^{2m+1} < 2^{3k+4}$$

choices of A such that $E_2 A \notin T_k$. Thus, altogether we have that $|E_i T_k \setminus T_k| \leq 2^{3k+4}$ for $i \in \{1, 2\}$, and since $|T_k| = 2^{4k+4}$ we conclude that

$$|E_i T_k \setminus T_k| \leq 2^{-k} |T_k|,$$

so that we can indeed pick $\varepsilon_k = 2^{-k}$. □

Theorem 6.14. *There exists an orbit equivalence coupling between \mathbb{Z}^4 and $\text{Heis}(\mathbb{Z})$ which is $(\varphi_\varepsilon, \varphi_\varepsilon)$ -integrable for all $\varepsilon > 0$, where*

$$\varphi_\varepsilon(x) = \frac{x}{\log(x)^{1+\varepsilon}}.$$

In particular the coupling is (L^p, L^p) for all $p < 1$.

Proof. This follows from Proposition 6.7, using the Følner tilings provided by Proposition 6.8 and Proposition 6.13. □

Remark 6.15. As explained in the introduction, there cannot be an L^1 measure equivalence coupling between \mathbb{Z}^4 and $\text{Heis}(\mathbb{Z})$. We thus ask the following question.

Question 6.16. Is there a (φ_0, φ_0) -integrable measure equivalence coupling between \mathbb{Z}^4 and $\text{Heis}(\mathbb{Z})$, where $\varphi_0(x) = \frac{x}{\log(x)}$?

6.3 A coupling between \mathbb{Z} and the lamplighter groups

Given two countable groups Λ, Γ and a function $f: \Gamma \rightarrow \Lambda$, the **support** of f is given by $\text{supp } f = \{\gamma \in \Gamma: f(\gamma) \neq e_\Lambda\}$. The space Λ^Γ is a group for pointwise multiplication. We then define $\bigoplus_{\gamma \in \Gamma} \Lambda$ as the subgroup of Λ^Γ consisting of all functions which have finite support. Finally, we the **wreath product** $\Lambda \wr \Gamma$ is by definition the semi-direct product

$$\Lambda \wr \Gamma := \bigoplus_{\gamma \in \Gamma} \Lambda \rtimes \Gamma,$$

where the group multiplication is given by $(f, \gamma) \cdot (f', \gamma') = (f\gamma \cdot f', \gamma\gamma')$ and $\gamma \cdot f(g) = f(\gamma^{-1}g)$. When the group Λ is finite and non-trivial, such groups are sometimes referred to as lamplighter groups. Here we shall consider the special case $\mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z}$, for which a standard generating set is provided by the pair $\{(0, 1), (\delta_0, 0)\}$ where $\delta_0(n) = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{otherwise.} \end{cases}$

The lamplighter point of view consists in viewing each element (f, n) of the group as a pair where f is a configuration of lamps (off at position i if $f(i) = 0$), and where n is the position of the ‘‘lamplighter’’. Multiplying (f, n) on the right by the first generator amounts to moving the lamplighter from position n to $n + 1$. Multiplying it by the second generator amounts to switching the light at position n .

Proposition 6.17. *Let $m \geq 2$ be a positive integer. The group $\mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z}$ (equipped with the generating set $S = \{(0, 1), (\delta_0, 0)\}$) admits a (ε_k, R_k) -Følner tiling sequence $(F_k)_k$, with $|F_0| = 2m^2$, and $|F_k| = 2 \cdot m^{2^k}$, $R_k = (m + 1)2^{k+1}$ and $\varepsilon_k = 2^{-(k+1)}$ for $k \geq 1$.*

Proof. Given the convention used to define the wreath product, it will be more convenient to construct a right Følner tiling sequence (F_k) . Recall that (F_k^{-1}) will then be a left Følner tiling sequence.

We define $F_0 = \{(f, n) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq \{0, 1\}, n \in \{0, 1\}\}$ and

$$F_k = \{(f, 0) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [2^k, 2^{k+1} - 1]\} \\ \cup \{(f, 2^k) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^k - 1]\}.$$

By induction, we show that $T_k = \{(f, n) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^{k+1} - 1], n \in [0, 2^{k+1} - 1]\}$. Indeed, we have that

$$T_{k+1} = F_{k+1}T_k \\ = \{(f, n) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^{k+2} - 1], n \in [0, 2^{k+1} - 1]\} \\ \cup \{(f, n) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^{k+2} - 1], n \in [2^{k+1}, 2^{k+2} - 1]\} \\ = \{(f, n) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^{k+2} - 1], n \in [0, 2^{k+2} - 1]\}.$$

A straightforward cardinality computation then shows that the F_{k+1} -translates of T_k must be pairwise disjoint, so the tiling condition is satisfied.

To bound the diameter of T_k , observe that to join two elements (f, n) and (f', n') in T_n , the lamplighter may travel from position n to n' , passing through the whole interval $[0, 2^{k+1} - 1]$, while possibly switching all the lamps (at most once) along the way. Since switching each lamp uses the generator $(\delta_0, 0)$ at most $\lfloor \frac{m}{2} \rfloor$ times, such a path has length at most $\lfloor \frac{m}{2} \rfloor 2^{k+1} + 2(2^{k+1} - 1)$. Hence the diameter of T_k is at most $\lfloor \frac{m}{2} \rfloor 2^{k+1} + 2(2^{k+1} - 1)$, which is less than $(m + 1)2^{k+1}$.

Finally, in order to prove the quantitative Følner condition, it suffices to show that $|T_k s \setminus T_k| \leq 2^{-(k+1)} |T_k|$ for all $s \in S$. If $s = (\delta_0, 0)$, then $T_k s \setminus T_k = \emptyset$. If $s = (0, 1)$, then

$$T_k s \setminus T_k = \{(f, 2^{k+1}) \in \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z} : \text{supp}(f) \subseteq [0, 2^{k+1} - 1]\}.$$

So either way, $|T_k s \setminus T_k| \leq m^{2^{k+1}} = 2^{-(k+1)} |T_k|$, which finishes the proof of the proposition. \square

Proposition 6.18. *For any integer $m \geq 2$, there exists an orbit equivalence coupling from \mathbb{Z} to $\mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z}$ which is $(\exp, \varphi_\varepsilon)$ -integrable for all $\varepsilon > 0$, where $\varphi_\varepsilon(x) = \frac{\log(x)}{\log(\log(x))^{1+\varepsilon}}$.*

Proof. Let $(F_k)_k$ be the Følner tiling sequence of $\mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z}$ constructed in Proposition 6.17. And let $(F'_k)_k$ the Følner tiling sequence of \mathbb{Z} defined by $F'_0 = [0, 2m^2 - 1]$ and

$$F'_k = |T_{k-1}| [0, |F_k| - 1] = \{0, m^{2^k} 2^k, 2m^{2^k} 2^k, \dots, (2m^{2^k} - 1)m^{2^k} 2^k\}.$$

Note that $T'_k = [0, |T_k| - 1]$. So $(F'_k)_k$ is a (ε'_k, R'_k) -Følner tiling sequence with $R'_k = |T'_k| - 1 \leq m^{2^{k+1}} 2^{k+1}$ and $\varepsilon'_k = \frac{\log}{|T'_k|} = \frac{1}{m^{2^{k+1}} 2^k}$. Proposition 6.7 yields an explicit orbit equivalence coupling (X, μ) between

the groups $\Gamma' = \mathbb{Z}$ and $\Gamma = \mathbb{Z}/m\mathbb{Z} \wr \mathbb{Z}$. Moreover, defining $z > 1$ as the solution of the equation $z^{2^3(m+1)}/m = 1$, we have

$$z^{2R_k} \varepsilon'_{k-1} \leq \frac{z^{2(m+1)2^{k+2}}}{m^{2^k} 2^{k-1}} = (z^{2^3(m+1)}/m)^{2^k} \frac{1}{2^{k-1}} = \frac{1}{2^{k-1}},$$

so we deduce from Proposition 6.7 that for every $s \in S_\Gamma$,

$$\int_X z^{d_{S_{\Gamma'}}(x, s \cdot x)} d\mu(x) < \infty.$$

On the other hand, we have

$$\varphi_\varepsilon(2R'_k) \varepsilon_{k-1} \leq \frac{2^{k+1} \log m + (k+2) \log 2}{\log(2^{k+1} \log m + (k+2) \log 2)^{1+\varepsilon}} \frac{1}{2^k} = O\left(\frac{1}{k^{1+\varepsilon}}\right).$$

Hence we deduce from Proposition 6.7 that that for every $s \in S_{\Gamma'}$,

$$\int_X \varphi(d_{S_\Gamma}(x, s \cdot x)) d\mu(x) < \infty.$$

This concludes the proof of the proposition. \square

7 Almost optimal couplings between \mathbb{Z} and iterated wreath products

In this section, we show that taking wreath products is well-behaved with respect to φ -integrable orbit equivalence, and use this to find almost optimal orbit equivalence couplings between \mathbb{Z} and iterated wreath products.

7.1 Wreath products of measure-preserving equivalence relations

In this section, we develop the concept of wreath products of measure-preserving equivalence relations in order to obtain a natural proof of the following statement: if Γ_1, Γ_2 are orbit equivalent, and Λ_1, Λ_2 are orbit equivalent, then $\Lambda_1 \wr \Gamma_1$ is orbit equivalent to $\Lambda_2 \wr \Gamma_2$. The measure-preserving equivalence relation that witnesses that they are orbit equivalent will be the wreath product $\mathcal{S} \wr \mathcal{R}$ of the equivalence relation \mathcal{R} witnessing the orbit equivalence between Γ_1 and Γ_2 with the equivalence relation \mathcal{S} witnessing the orbit equivalence between Λ_1 and Λ_2 .

Let \mathcal{R} be a measure-preserving equivalence relation on (X, μ) , let \mathcal{S} be a measure-preserving equivalence relation on (L, ν) . Consider the space $L^{\mathcal{R}}$ which consists of couples $(x, (l_y)_{y \in [x]_{\mathcal{R}}})$ where $l_y \in L$ for every $y \in [x]_{\mathcal{R}}$. This space can be equipped with a natural standard Borel space structure which we don't make explicit for now since in our concrete case it will be easy to describe. Moreover, one can endow it with the following probability measure η given by

$$\eta = \int_X \nu^{\otimes [x]_{\mathcal{R}}} d\mu(x).$$

We then equip the space $(L^{\mathcal{R}}, \eta)$ with the **wreath product** $\mathcal{S} \wr \mathcal{R}$ of \mathcal{S} by \mathcal{R} defined by saying two couples $(x, (l_y)_{y \in [x]_{\mathcal{R}}})$ and $(x', (l'_y)_{y \in [x]_{\mathcal{R}}})$ are $\mathcal{S} \wr \mathcal{R}$ -equivalent as soon as $(x, x') \in \mathcal{R}$ and the following two conditions are satisfied:

- for all but finitely many $y \in [x]_{\mathcal{R}}$, we have $l_y = l'_y$;
- for all $y \in [x]_{\mathcal{R}}$, we have $(l_y, l'_y) \in \mathcal{S}$.

One can check that $\mathcal{S} \wr \mathcal{R}$ is indeed a measure-preserving equivalence relation.

Let us make all this completely explicit in the case of interest to us: we now suppose that \mathcal{R} comes from a fixed *free* Γ -action. We then have a cocycle map $c: \mathcal{R} \rightarrow \Gamma$ which takes $(x, y) \in \mathcal{R}$ to the unique $\gamma \in \Gamma$ satisfying $\gamma \cdot x = y$. The standard Borel space structure on $L^{\mathcal{R}}$ then comes from its natural identification to the standard Borel space $X \times L^{\Gamma}$ given by the bijection $\Phi: X \times L^{\Gamma} \rightarrow L^{\mathcal{R}}$ defined by

$$\Phi(x, (l_{\gamma})_{\gamma \in \Gamma}) = (x, (l_{c(x,y)})_{y \in [x]_{\mathcal{R}}}).$$

The inverse map is given by $\Phi^{-1}(x, (l_y)_{y \in [x]_{\mathcal{R}}}) = (x, (l_{\gamma \cdot x})_{\gamma \in \Gamma})$. Note that $\Phi_*(\mu \otimes \nu^{\otimes \Gamma}) = \eta$. We have a natural Γ -action on $L^{\mathcal{R}}$ given by $\gamma \cdot (x, l) = (\gamma \cdot x, l)$. The corresponding action on $X \times L^{\Gamma}$ is given by

$$\gamma \cdot (x, (l_g)_{g \in \Gamma}) = (\gamma \cdot x, (l_{g\gamma})_{g \in \Gamma}).$$

Finally, suppose that the equivalence relation \mathcal{S} comes from a free Λ -action on L . We then also have a natural $\bigoplus_{\gamma \in \Gamma} \Lambda$ -action on $X \times L^{\Gamma}$ given by: for all $f \in \bigoplus_{\gamma \in \Gamma} \Lambda$,

$$f \cdot (x, (l_{\gamma})_{\gamma \in \Gamma}) = (x, (f(g^{-1}) \cdot l_g)_{g \in \Gamma}),$$

or from the $L^{\mathcal{R}}$ viewpoint, $f \cdot (x, (l_y)_{y \in [x]_{\mathcal{R}}}) = (x, (f(c(y, x)) \cdot l_y)_{y \in [x]_{\mathcal{R}}})$.

One can check that the Γ and $\bigoplus_{\gamma \in \Gamma} \Lambda$ -actions actually extend to a free action of $\Lambda \wr \Gamma$ on $L^{\mathcal{R}}$ given by

$$(f, \gamma) \cdot \left(x, (l_y)_{y \in [x]_{\mathcal{R}}} \right) = \left(\gamma \cdot x, (f(c(y, \gamma \cdot x)) \cdot l_y)_{y \in [x]_{\mathcal{R}}} \right).$$

One can check that the equivalence relation induced by this action is equal to $\mathcal{S} \wr \mathcal{R}$. Since the action is measure-preserving, this proves that $\mathcal{S} \wr \mathcal{R}$ is measure-preserving. We have shown the following result.

Proposition 7.1. *Let $\Gamma \curvearrowright (X, \mu)$ freely, let $\Lambda \curvearrowright (L, \nu)$ freely, where (X, μ) and (L, ν) are two standard probability spaces and the actions are measure-preserving. Denote by \mathcal{R} and \mathcal{S} the respective associated equivalence relations, and by $c: \mathcal{R} \rightarrow \Gamma$ the cocycle defined by $c(x, \gamma \cdot x) = \gamma$. Then the measure-preserving free $\Gamma \wr \Lambda$ -action on $L^{\mathcal{R}}$ given by*

$$(f, \gamma) \cdot \left(x, (l_y)_{y \in [x]_{\mathcal{R}}} \right) = \left(\gamma \cdot x, (f(c(y, \gamma \cdot x)) \cdot l_y)_{y \in [x]_{\mathcal{R}}} \right)$$

induces the equivalence relation $\mathcal{S} \wr \mathcal{R}$. □

7.2 Wreath products and quantitative orbit equivalence

It follows directly from the previous proposition that when Γ_1, Γ_2 are orbit equivalent, and Λ_1, Λ_2 are orbit equivalent, then $\Lambda_1 \wr \Gamma_1$ is orbit equivalent to $\Lambda_2 \wr \Gamma_2$. We now give a quantitative version of this fact.

For this, it is useful to identify Γ and Λ to subgroups of $\Gamma \wr \Lambda$ as follows: first we embed Λ in $\bigoplus_{\gamma \in \Gamma} \Lambda$ by associating to every $\lambda \in \Lambda$ the function

$$\iota(\lambda): \gamma \mapsto \begin{cases} \lambda & \text{if } \gamma = e_{\Gamma} \\ e_{\Lambda} & \text{otherwise.} \end{cases}$$

Then we embed Λ in $\Lambda \wr \Gamma$ via $\lambda \mapsto (e_{\Gamma}, \iota(\lambda))$, and we embed Γ into $\Lambda \wr \Gamma$ via $\gamma \mapsto (\gamma, \iota(e_{\Lambda}))$. It is well known that if S_{Γ} is a generating set for Γ and S_{Λ} is a generating set for Λ , then through the above identification $S_{\Lambda} \cup S_{\Gamma}$ is a generating set for the wreath product $\Lambda \wr \Gamma$.

Proposition 7.2. *Let (X, μ) be an orbit equivalence coupling between finitely generated groups $\Gamma_1 = \langle S_{\Gamma_1} \rangle$ and $\Gamma_2 = \langle S_{\Gamma_2} \rangle$, and let (L, ν) be an orbit equivalence coupling between $\Lambda_1 = \langle S_{\Lambda_1} \rangle$ and $\Lambda_2 = \langle S_{\Lambda_2} \rangle$. Denote by \mathcal{R} the equivalence relation generated by the Γ_i -action.*

Then the orbit coupling $L^{\mathcal{R}}$ satisfies that for every $(x, l) \in L^{\mathcal{R}}$, every $\gamma_1 \in \Gamma_1$ and every $\lambda_1 \in \Lambda_1$,

$$\begin{aligned} d_{S_{\Lambda_2} \cup S_{\Gamma_2}}((x, l), \gamma_1 \cdot (x, l)) &= d_{S_{\Gamma_2}}(x, \gamma_1 \cdot x) \\ d_{S_{\Lambda_2} \cup S_{\Gamma_2}}((x, l), \lambda_1 \cdot (x, l)) &= d_{S_{\Lambda_2}}(l_{e_{\Gamma}}, \lambda_1 \cdot l_{e_{\Gamma}}) \end{aligned}$$

Proof. By definition for all $i \in \{1, 2\}$, all $\gamma_i \in \Gamma_i$, and all $(x, l) \in L^{\mathcal{R}}$, we have $\gamma_i \cdot (x, l) = (\gamma_i \cdot x, l)$, so the first equation is clear. The second equation follows similarly by noting that if $\lambda_i \in \Lambda_i$, then their action on (x, l) only changes the value at e_{Γ} of l according to the λ_i -action on L . \square

Corollary 7.3. *If Γ_1 and Γ_2 admit a (φ, ψ) -integrable orbit equivalence coupling, and if Λ_1 and Λ_2 admit a (φ, ψ) -integrable orbit equivalence coupling, then the wreath products $\Lambda_1 \wr \Gamma_2$ and $\Lambda_2 \wr \Gamma_2$ also admit a (φ, ψ) -integrable orbit equivalence couplings.* \square

In combination with Theorem 6.10 we find the following:

Corollary 7.4. *Let $n, m \in \mathbb{N}$ with $n < m$ and let Δ be any finitely generated group, then there is an $(L^p, L^{1/p})$ -orbit equivalence coupling from $\Delta \wr \mathbb{Z}^m$ to $\Delta \wr \mathbb{Z}^n$ for every $p < \frac{n}{m}$.* \square

We also obtain the following corollary by using compositions of couplings. Recall that we denote by \log^{*n} the function \log composed with itself n times. We denote $L^{<\infty}$ the intersection of L^p for all $1 \leq p < \infty$.

Corollary 7.5. *Let $m > 0$, and let G_n be defined inductively by $G_0 = \mathbb{Z}$, and $G_{n+1} = \mathbb{Z}/m\mathbb{Z} \wr G_n$. For any integer $n \geq 1$, there exists an orbit equivalence coupling from \mathbb{Z} to G_n which is $(L^{<\infty}, \varphi_{n,\varepsilon})$ -integrable for every $\varepsilon > 0$, where $\varphi_{n,\varepsilon} = \log^{*n} / (\log^{*(n+1)})^{1+\varepsilon}$.*

Proof. We first claim that for all $n \geq 2$, there exists an orbit equivalence coupling from G_{n-1} to G_n which is $(L^{<\infty}, \varphi_{1,\varepsilon})$ -integrable for all $\varepsilon > 0$: this follows by induction using Proposition 7.2, the case $n = 2$ resulting from Proposition 6.18.

We now pass to the proof of the corollary, which we also do by induction on n . The base case $n = 1$ follows from the claim, so we take $n \geq 2$ and assume that we have a orbit equivalence coupling from \mathbb{Z} to G_{n-1} which is $(L^{<\infty}, \varphi_{n-1,\varepsilon})$ -integrable for all $\varepsilon > 0$. Using Proposition 2.26, we compose this coupling with the coupling from the claim between G_{n-1} and G_n , and obtain a $(\varphi_{1,\varepsilon} \circ \varphi_{n-1,\varepsilon}, L^{<\infty})$ -coupling from \mathbb{Z} to G_n . Finally one easily shows that $\varphi_{1,\varepsilon} \circ \varphi_{n-1,\varepsilon} \succcurlyeq \varphi_{n,\varepsilon}$, so we are done. \square

8 Unstable properties under exponential couplings

8.1 Finite presentability is unstable

In this section we prove that being finitely presented is not preserved by strongly exponentially integrable orbit equivalence couplings (see Definition 2.28). We do this by constructing an explicit strongly exponentially integrable orbit coupling between the wreath product $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ (also known as a lamplighter group) and the Baumslag-Solitar group $\text{BS}(1, k)$, for every $k \geq 2$.

Recall from the previous section that $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z} = \bigoplus_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z} \rtimes \mathbb{Z}$ where \mathbb{Z} acts by a shift on $\bigoplus_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$, that is

$$((x_i)_{i \in \mathbb{Z}}, m_x)((y_i)_{i \in \mathbb{Z}}, m_y) = ((x_i + y_{i-m_x})_{i \in \mathbb{Z}}, m_x + m_y).$$

Also recall that $\text{BS}(1, k) = \mathbb{Z}[1/k] \rtimes \mathbb{Z}$, where $\mathbb{Z}[1/k] = \{\frac{x}{k^m} : x, m \in \mathbb{Z}\}$ and \mathbb{Z} acts on $\mathbb{Z}[1/k]$ by multiplication by $1/k$, that is

$$\left(\frac{x}{k^{m_x}}, n_x\right) \left(\frac{y}{k^{m_y}}, n_y\right) = \left(\frac{x}{k^{m_x}} + \frac{y}{k^{m_y+n_x}}, n_x + n_y\right).$$

It is well known that the second group is finitely presented, while the first one is not by a result of Baumslag [Bau61]. In particular these groups are not quasi-isometric.

We will consider actions of these two groups on the standard Borel space $X = \prod_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ equipped with the measure μ defined as the infinite product of the normalized counting measure. The action of the lamplighter group $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ is the standard one, that is

$$((x_i)_{i \in \mathbb{Z}}, m) \cdot (y_i)_{i \in \mathbb{Z}} = ((x_i + y_{i-m})_{i \in \mathbb{Z}}, m).$$

For the Baumslag-Solitar group $\text{BS}(1, k)$ we first define the action of the subgroup $\mathbb{Z}[1/k] = \langle (k^m, 0) : m \in \mathbb{Z} \rangle$ as follows: for all $m \in \mathbb{Z}$, we decompose the space X as

$$X = \prod_{i < m} \mathbb{Z}/k\mathbb{Z} \times \prod_{i \geq m} \mathbb{Z}/k\mathbb{Z},$$

and then $(k^m, 0)$ acts trivially on the first factor, and as the k -adic odometer on the second factor. To be completely explicit, $(k^m, 0) \cdot (x_i)_{i \in \mathbb{Z}} = (y_i)_{i \in \mathbb{Z}}$, where $x_i = y_i$ for every $i < m$, $y_N = x_N + 1$ where $N \geq m$ the smallest such that $x_N \neq k - 1$, $y_i = 0$ for $m \geq i < N$ and $y_i = x_i$ for $i > N$. If $x_i = k - 1$ for every $i \geq m$, then we take $y_i = 0$ for every $i \geq m$. Note that this action is realized by letting $\mathbb{Z}[1/k]$ act on \mathbb{Q}_k by addition and extending this action to the bi-infinite product $X = \prod_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$. Also note that the equivalence relation generated by the action of $\mathbb{Z}[1/k]$ is the cofinite relation up to measure zero.

As with the lamplighter group we then let \mathbb{Z} act by $(0, 1) \cdot (x_i)_{i \in \mathbb{Z}} = (x_{i-1})_{i \in \mathbb{Z}}$. This defines a free pmp action of the group $\mathbb{Z}[1/k] \rtimes \mathbb{Z}$ on X . Since the natural action of $\bigoplus_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ is the cofinite one, we deduce that the two actions we have build yield an orbit equivalence coupling (X, μ) from the lamplighter group $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ to the Baumslag-Solitar group $\text{BS}(1, k)$.

Theorem 8.1. *The orbit equivalence coupling (X, μ) we just constructed is an $(L^\infty, \text{exp}^\diamond)$ orbit equivalence coupling from $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ to $\text{BS}(1, k)$.*

Proof. We equip $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ with the generating subset $S = \{\delta_0, 1_{\mathbb{Z}}\}$ and $\text{BS}(1, k)$ with the generating subset $T = \{1_{\mathbb{Z}[1/k]}, 1_{\mathbb{Z}}\}$. We denote by $S^\pm = S \cup S^{-1}$ and $T^\pm = T \cup T^{-1}$ the corresponding symmetric generating sets.

Note that the generator of the \mathbb{Z} copy in the two groups acts exactly the same. Moreover, the two subgroups $\bigoplus_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ and $\mathbb{Z}[1/k]$ have the same orbits.

Let us start by showing that our orbit equivalence coupling is L^∞ as a coupling from the lamplighter group to the Baumslag-Solitar group. For this, we only need to check that $d_{T^\pm}(s \cdot x, x)$ is uniformly bounded for all $s \in S$. Note that the generator of the \mathbb{Z} copy in the two groups acts exactly the same. So for $s = 1_{\mathbb{Z}}$, we have $d_{T^\pm}(s \cdot x, x) = 1$. Then observe that the generator δ_0 changes only x_0 , which is achieved by the action of s^m for some $m \in \{-k + 1, \dots, k - 1\}$, where $s = 1_{\mathbb{Z}[1/k]}$. Hence $d_{T^\pm}(\delta_0 \cdot x, x) \leq k - 1$, which proves that our coupling is L^∞ in one direction.

To prove that the coupling is strongly exponential in the other direction, we need to obtain estimates for every $g \in \text{BS}(1, k)$. So let $g \in \text{BS}(1, k)$, we will show that g satisfies the following estimate: for all $M \geq 0$,

$$\mu\left(\{x \in X : d_{S^\pm}(g \cdot x, x) \geq (k + 1)(2|g|_T + 2M + 3)\}\right) \leq k^{-M+1} \quad (21)$$

Let $n = |g|_{T^\pm}$, write $g = (z, j)$, with $|j| \leq n$ and $z \in \mathbb{Z}[1/k]$. First note that by the triangle inequality and the fact that $d_{S^\pm}((0, j) \cdot x, x) = j$ for all $x \in X$, it suffices to show that

$$\mu\left(\{x \in X : d_{S^\pm}(z \cdot x, x) \geq k(2|g|_T + 2M + 3)\}\right) \leq k^{-M+1} \quad (22)$$

By symmetry we may assume that $z > 0$. A straightforward induction on n then shows that we can always write z as $z = \sum_{i=-n}^n a_i k^i$, where $a_i \in \{0, \dots, k - 1\}$.

Let $x \in X$. Observe that the coefficients of $z \cdot x$ that differ from those of x are contained in the interval $[-n, \infty)$. Moreover, consider the event $(z \cdot x)_{n+m+2} \neq x_{n+m+2}$ for some $m \geq M$, and observe that its occurrence forces the coefficients of x_{n+i} for $i = 1, 2, \dots, m + 1$ to be equal to $k - 1$. Therefore this event has probability at most $\sum_{m \geq M} k^{-m} \leq k^{-M+1}$.

Now note that $|f|_{S^\pm} \leq k(2n + M + 2)$ for every $f \in \bigoplus_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ with $\text{supp}(f) \subseteq [-n, n + M + 1]$, so the estimate (22) holds, and hence so does (21).

Let us finally check that this estimate yields the desired result. For a given $\delta > 0$ and $g \in \text{BS}(1, k)$,

by the integration by slices inequality and (21) we have

$$\begin{aligned} \int_X \exp(\delta d_{S^\pm}(x, g \cdot x)) d\mu(x) &\leq \sum_{M \geq 0} k^{-M+1} \exp(\delta k(2|g|_T + 2M + 3)) \\ &= \sum_{M \geq 0} k \exp(\delta k(2|g|_T + M(2 - \log k/\delta) + 3)) \end{aligned}$$

So given $\varepsilon > 0$, we take δ such that $2\delta k < \varepsilon$ and $2 - \log k/\delta < 0$, and then we will have a positive constant $C > 0$ such that for every $g \in \text{BS}(1, k)$,

$$\int_X \exp(\delta d_{S^\pm}(x, g \cdot x)) d\mu(x) < C \exp(\varepsilon |g|_T),$$

which finishes the proof that the coupling is strongly exp-integrable from $\text{BS}(1, k)$ to $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$. \square

Corollary 8.2. *Finite presentability is unstable under the equivalence relation of strongly exponential orbit equivalence. In particular, it is unstable under $L^{<\infty}$ orbit equivalence.*

8.2 Finite asymptotic dimension is unstable

We now consider the following group:

$$\Gamma_1 = \bigoplus_{\mathbb{Z}} \mathbb{Z}[1/k] \rtimes \mathbb{Z}^2,$$

where the first coordinate of \mathbb{Z}^2 acts by shift on the direct sum $\bigoplus_{\mathbb{Z}} \mathbb{Z}[1/k]$, and the second coordinate multiplies each factor by the corresponding power of k . Recall the action of $\mathbb{Z}[1/k]$ on $X = \prod_{\mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ as defined in section 8.1 induces the cofinite equivalence relation up to measure zero. For every $n \in \mathbb{Z}$, we let X_n be a copy of X . We deduce an action of the direct sum $\bigoplus_{n \in \mathbb{Z}} \mathbb{Z}[1/k]$ on $Y = \prod_{n \in \mathbb{Z}} X_n = \prod_{\mathbb{Z}^2} \mathbb{Z}/k\mathbb{Z}$ which induces the cofinite equivalence relation up to measure zero. The cofinite equivalence relation is also induced by the natural action of the group $\bigoplus_{\mathbb{Z}^2} \mathbb{Z}/k\mathbb{Z}$ on Y .

Combining these actions with the Bernoulli action of \mathbb{Z}^2 produces an orbit equivalence coupling between Γ_1 and the wreath product

$$\Gamma_2 = \mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}^2.$$

We equip Γ_1 with the generating subset $S_1 = \{\pm 1_{\mathbb{Z}[1/k]_0}\} \cup S_{\mathbb{Z}^2}$, and Γ_2 with $S_2 = \{\delta_0\} \cup S_{\mathbb{Z}^2}$, where $S_{\mathbb{Z}^2}$ is the canonical generating set of \mathbb{Z}^2 .

Theorem 8.3. *The coupling we just defined is an (L^∞, exp) -integrable orbit equivalence coupling from Γ_2 to Γ_1 . Actually the exponential integrability from Γ_1 to Γ_2 is uniform in the following sense: there exists a constant $c > 0$ such that for all $g \in \Gamma_1$*

$$\int_X \exp(cd_{S_2}(x, g \cdot x)) d\mu(x) < \infty.$$

Proof. The proof that the coupling is L^∞ from Γ_2 to Γ_1 is exactly the same as in the proof of Theorem 8.1, so we don't write it down. Also similar to the proof of Theorem 8.1, we observe that the two respective copies of \mathbb{Z}^2 act the same. As a warm up, let us prove the exponential integrability from Γ_1 to Γ_2 . Thanks to Proposition 2.20 only needs to consider the action of the generators $\pm 1_{\mathbb{Z}[1/k]_0}$. Observe that the coupling induces on every X_n an orbit-equivalence coupling between $\mathbb{Z}[1/k]_0$ and $\bigoplus_{\mathbb{Z} \times \{0\}} \mathbb{Z}/k\mathbb{Z}$ that coincides with that of section 8.1. Hence the result follows from Theorem 8.1.

The uniform exponential integrability requires a bit more work. It will result from the following statement: given $g \in \Gamma_1$ and $M \geq 0$ we have that

$$\mu(\{y \in Y : d_{S_2}(y, g \cdot y) \geq (k+1)(2M + (2|g|+5)(2|g|+1)) + |g|\}) \leq \int_{t \geq M} t^{2|g|+1} k^{-t} dt, \quad (23)$$

where we write $|g|$ instead of $|g|_{S_1}$ in order to simplify notation. Indeed, this implies that

$$\mu(\{y \in Y : d_{S_2}(y, g \cdot y) \geq (k+1)(2M + (2|g| + 5)(2|g| + 1)) + 3|g| + 1\}) \leq \frac{k^{-M}}{\log k}.$$

and therefore

$$\int_X \exp(cd_{S_2}(x, g \cdot x)) d\mu(x) < \infty$$

for all $c < \log k$.

We will use the fact that the n^{th} copy $\mathbb{Z}[1/k]$ and $\oplus_{\{n\} \times \mathbb{Z}} \mathbb{Z}/k\mathbb{Z}$ have the same orbits. Moreover they only act non-trivially on the n^{th} factor X_n , and their actions are those defined in section 8.1. Let $y = (y_{i,j})$ and $g = ((x_i), (m, n))$. Note that $|n| + |m| \leq |g|$. Hence

$$d_{S_2}(y, g \cdot y) \leq |g| + \sum_{i=-|g|}^{|g|} d_T((y_{i-n,j-m})_{j \in \mathbb{Z}}, x_i \cdot (y_{i-n,j-m})_{j \in \mathbb{Z}}),$$

where T is the generating subset of $\mathbb{Z}/k\mathbb{Z} \wr \mathbb{Z}$ from section 8.1. Now for every $-|g| \leq i \leq |g|$, let $M_i \in \mathbb{R}$ be such that

$$d_T((y_{i-n,j-m})_{j \in \mathbb{Z}}, x_i \cdot (y_{i-n,j-m})_{j \in \mathbb{Z}}) = (k+1)(2|g| + 2M_i + 5).$$

We deduce that $\sum_{i=-|g|}^{|g|} M_i \geq M$. Note that $|g| \geq |(x_i)_{i \in \mathbb{Z}}|_S$, where S is the generating subset of $\text{BS}(1, k)$ from section 8.1. On the other hand, by (21) we have that for all $(t_i) \in \mathbb{R}_+^{2|g|+1}$,

$$\mu(\{y \in Y : d_T((y_{i-n,j-m})_{j \in \mathbb{Z}}, x_i \cdot (y_{i-n,j-m})_{j \in \mathbb{Z}}) \geq (k+1)(2|g|_S + 2t_i + 5)\}) \leq k^{-t_i}.$$

For every $t \geq 0$ and $m \in \mathbb{N}$, denote by $v_m(t) = \text{Vol}\{(t_i) \in \mathbb{R}_+^m, \sum_i t_i = t\}$, and observe that $v_m(t) \leq t^m$. Finally, integrating over all values for $t_i = \max\{M_i, 0\}$ for which $t = \sum_i t_i \geq M$, we obtain

$$\begin{aligned} \mu(\{y \in Y : d_{S_2}(y, g \cdot y) \geq (k+1)(2M + (2|g| + 5)(2|g| + 1)) + |g|\}) &\leq \int_{t \geq M} k^{-t} v_{2|g|+1}(t) dt \\ &\leq \int_{t \geq M} t^{2|g|+1} k^{-t} dt \end{aligned}$$

which ends the proof of (23) and therefore of the theorem. \square

Observing that Γ_1 has infinite asymptotic dimension, while Γ_2 has asymptotic dimension 2 (e.g. as a consequence of the results from [DS06, Sec. 3]), we deduce the following corollary.

Corollary 8.4. *There exist a group with asymptotic dimension 2, and a group of infinite asymptotic dimension that admit an exponential-integrable orbit equivalence measure coupling.*

9 Hyperbolicity

In this section we prove the results of §1.3. As explained in the introduction, our main tool will be to characterize Gromov hyperbolicity in terms of distortion of embedded cycles. More precisely we shall confront the facts that an n -cycle in a hyperbolic graph must have distortion in $\Theta(n/\log n)$ (see Proposition 9.2), while any non-hyperbolic graph admits arbitrary large cycles with universally bounded distortion (see Proposition 9.8). A version of the first statement for real hyperbolic spaces is proved in [VS14, Lemma 3.3] and a weaker version of our second statement is obtained in [VS14, Theorem 4.1].

9.1 Geometric preliminaries

Let X be a graph, a **discrete path** in X of length $l \geq 1$ is a map $\alpha: \{0, \dots, l\} \rightarrow X$ such that for all $i \in \{0, \dots, l-1\}$, we have that $\alpha(i)$ and $\alpha(i+1)$ are connected by an edge. We will also say that α is a path from $\alpha(0)$ to $\alpha(l)$, and we will often identify a path to its range.

Every connected graph X is viewed as a metric space (X, d) equipped with the **discrete path metric**, defined by setting $d(x, y)$ as the minimum length of a path from x to y . Any discrete path which realizes the discrete path metric between two points is called a **discrete geodesic**, and it is then an isometric embedding from $\{0, \dots, d(x, y)\}$ to (X, d) .

Another important metric space that we can get out of a connected graph X is given by the (continuous) **path metric** which we define as in [Gro07, 1.15₊]. We first identify each edge to the interval $[0, 1]$ isometrically, thus obtaining a *length structure* on our graph. The metric associated to this length structure is denoted by d_l , and it is by definition the continuous path metric on X . It agrees with the discrete path metric on the vertices of X , and it is geodesic. Every geodesic between vertices defines a discrete geodesic, and every discrete geodesic can be lifted to a geodesic between vertices.

A (geodesic) **triangle** in a metric space (X, d) with **vertices** $a_1, a_2, a_3 \in X$ is the set $[a_1, a_2, a_3] \subseteq X$ obtained by taking the union of a choice of geodesics $[a_1, a_2]$, $[a_2, a_3]$, and $[a_3, a_1]$ between its vertices. In the same way, we define a (geodesic) **n -gon** with vertices $a_1, \dots, a_n \in X$, and denote it by $[a_1, \dots, a_n]$. Given an n -gon where $n \geq 3$, we will frequently call any of its defining geodesics a **side**.


Now, recall that a geodesic space (X, d) is **δ -hyperbolic** in the sense of Rips if there exists a $\delta \geq 0$ such that for every geodesic triangle $[a_1, a_2, a_3]$ and for every $x \in [a_1, a_2]$, there exists an element y in either $[a_1, a_3]$ or $[a_2, a_3]$ such that $d(x, y) \leq \delta$; or equivalently, that the side $[a_1, a_2]$ is contained in the δ -neighborhood of $[a_1, a_3] \cup [a_2, a_3]$. Moreover, we say that a geodesic space (X, d) is **hyperbolic** if it is δ -hyperbolic for some δ , and that a finitely generated group Γ with generating set S is hyperbolic whenever its Cayley graph is hyperbolic when equipped with the continuous path metric.

We shall need the following well-known lemma.

Lemma 9.1. *Let X be a δ -hyperbolic geodesic space, let α be a path of length $\ell \geq 1$ between two points x_1 and x_2 , and let y belong to a geodesic from x_1 to x_2 . Then*

$$d(y, \alpha) \leq \delta \log_2(\ell) + 1.$$

Proof. Let us prove it by induction on $[\ell]$. The case $[\ell] = 1$ is clear. So assume $n \geq 2$ and suppose that the lemma is true for all paths of length $< n$. Let α be a path of length $\ell \in [n, n+1[$ from x_1 to x_2 , represented as a gray path in the following figure. For every $x \in \alpha$, using that the geodesic triangle $[x_1, x, x_2]$ is δ -thin, then either $d(y, [x_1, x]) \leq \delta$ or $d(y, [x, x_2]) \leq \delta$. By connectedness of α , there exists x_3 such that both conditions are satisfied. By exchanging x_1 and x_2 if necessary, we can assume that the portion α_1 of α from x_1 to x_3 has length $\ell_1 \leq \ell/2$.



Hence there exists a point $y' \in [x_1, x_3]$ such that $d(y, y') \leq \delta$. Now applying the induction hypothesis to the path α_1 and the point $y' \in [x_1, x_3]$, we obtain

$$d(y', \alpha_1) \leq \delta \log_2(\ell_1) \leq \delta \log_2 \ell + 1 - \delta.$$

We deduce by the triangle inequality

$$d(y, \alpha) \leq d(y, y') + d(y', \alpha_1) \leq \delta \log_2(\ell) + 1.$$

So we are done. \square

First we need an alternative definition of hyperbolicity in terms of embedded cycles. In what follows, the **cycle** C_n of length $n \geq 2$ is the Cayley graph of $\mathbb{Z}/n\mathbb{Z}$ with respect to the generating set containing only the element $1 \bmod n$, which we view as a *discrete* metric space denoted by (C_n, d_{C_n}) .

Proposition 9.2. *Let (X, d) be a δ -hyperbolic geodesic space, let n be a positive integer. Then for every $a \geq 0$ and every $b \geq 1$, if there is a map $\varphi: C_{2n} \rightarrow X$ such that for every $x, y \in C_{2n}$,*

$$ad_{C_{2n}}(x, y) \leq d(\varphi(x), \varphi(y)) \leq bd_{C_{2n}}(x, y)$$

then we have

$$a \leq \frac{4\delta \log_2(bn) + 4 + 2b}{n}$$

Before proving the above proposition, let us note the following straightforward corollary, using the estimate $\frac{1}{\log 2} < 3/2$.

Corollary 9.3. *Let $a \geq 0$, $b \geq 1$. There is an integer $n_0 = n_0(b) \geq 2$ such that the following holds. For all $n \geq n_0$ and all $\delta \geq 1$, if X is a δ -hyperbolic geodesic space and if there is a map $\varphi: C_n \rightarrow X$ such that for all $x, y \in C_n$*

$$ad_{C_n}(x, y) \leq d(\varphi(x), \varphi(y)) \leq bd_{C_n}(x, y)$$

then we have

$$a < 12\delta \cdot \frac{\log n}{n}. \tag{24}$$

In order to prove the proposition, we need the following additional notion. Given a discrete path β in a graph Y and a map $\varphi: Y \rightarrow (X, d)$ where (X, d) is geodesic, we say that a continuous path α in X is a φ -**direct image** of β if it is obtained by concatenating geodesics between $\varphi(\beta(i))$ and $\varphi(\beta(i+1))$ where i ranges from 0 to $\ell(\beta) - 1$.


Proof of Proposition 9.2. Let $a_1, a_2 \in C_{2n}$ be such that $d_{C_{2n}}(a_1, a_2) = n$ and define $x_1 = \varphi(a_1)$, $x_2 = \varphi(a_2)$. Consider a geodesic $[x_1, x_2]$ from x_1 to x_2 . In C_{2n} there are two discrete geodesic paths from a_1 to a_2 , both with length n . Denote by α and α' some respective φ -direct images of those paths in X , which by assumption have length at most bn . Let $y \in [x_1, x_2]$, we deduce from Lemma 9.1 that

$$d(y, \alpha) \leq \delta \log_2(\ell(\alpha)) + 1 \leq \delta \log_2(bn) + 1,$$

where $\ell(\alpha)$ is the length of α , and by the same argument $d(y, \alpha') \leq \delta \log_2(bn) + 1$. If we then pick z_y and z'_y points in C_{2n} such that their φ -images are in α and α' respectively and minimize the distance to y , we have

$$\max(d(y, \varphi(z_y)), d(y, \varphi(z'_y))) \leq \delta \log_2(bn) + 1 + \frac{b}{2}. \tag{25}$$

Note that for any $y \in [x_1, x_2]$, any geodesic from z_y to z'_y must pass through x_1 or through x_2 . Moreover there are some $y \in [x_1, x_2]$ for which the first case occurs, and some for which the second case occurs. For all $\varepsilon > 0$, we may thus find $y_1, y_2 \in [x_1, x_2]$ such that $d(y_1, y_2) \leq \varepsilon$, the geodesic from z_{y_1} to z'_{y_1} passes through x_1 and the geodesic from z_{y_2} to z'_{y_2} passes through x_2 .



Then we have that

$$d(z_{y_1}, z_{y_2}) + d(z_{y_2}, z'_{y_2}) + d(z'_{y_2}, z'_{y_1}) + d(z'_{y_1}, z_{y_1}) \geq 2n.$$

Hence one of these four distances is at least $\frac{n}{2}$. However, a straightforward case analysis using the bound (25) on the lengths of the dotted geodesics, the triangle inequality and the fact that $d(y_1, y_2) \leq \varepsilon$ shows that the distance between the corresponding images of the points in the previous inequality is always at most $2(\delta \log_2(bn) + 1 + \frac{b}{2}) + \varepsilon$. Using our assumption on φ , we thus have the following inequality: for all $\varepsilon > 0$,

$$\frac{an}{2} \leq 2\delta \log_2(bn) + 2 + b + \varepsilon.$$

So the proposition follows. \square

In what follows, given a constant $a \geq 1$, we will say that a function $f: (X, d_X) \rightarrow (Y, d_Y)$ between two metric spaces **contracts distances by at most the factor a** if for every $x_1, x_2 \in X$ we have $d_X(x_1, x_2) \leq a d_Y(f(x_1), f(x_2))$.

Lemma 9.4. *Let $n \geq 4$ and $D \geq 1$, let (X, d) be a geodesic metric space containing an n -gon $[a_1, \dots, a_n]$ such that the distance between points on non-adjacent sides is at least D . Let $R \geq D$ be the length of its longest side. Then there is a cycle C of length at least $\frac{Dn}{2}$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ which contracts distances by at most a factor $\frac{3nR}{D}$.*

Proof. Our cycle will be obtained via the following claim.

Claim 9.5. There is a $2n$ -gon $C = [b_1, c_1, b_2, \dots, b_n, c_n]$ such that for every $i \in \{1, \dots, n\}$,

- $b_i \in [a_{i-1}, a_i]$ and $c_i \in [a_i, a_{i+1}]$, where we let $a_0 = a_n$;
- $d(b_i, c_i) = \frac{D}{8R}(d(b_i, a_i) + d(a_i, c_i))$;
- for each pair of points $x \in [c_{i-1}, b_i]$ and $y \in [c_i, b_{i+1}]$, we have $d(x, y) \geq \frac{D}{8R}(d(x, a_i) + d(a_i, y))$.

Proof of the claim. We start with the existence of b_1 and c_1 . Note that for all points $u \in [a_n, a_1]$ and $v \in [a_1, a_2]$, we have $d(u, a_1) + d(a_1, v) \leq 2R$ and so we have $\frac{D}{8R}(d(u, a_1) + d(a_1, v)) \leq \frac{D}{4}$.

Taking $u = a_1 = v$, we see that there are $u \in [a_n, a_1]$ and $v \in [a_1, a_2]$ such that $d(u, v) = \frac{D}{8R}(d(u, a_1) + d(a_1, v))$. Also, for $u = a_n$ and $v = a_2$, we have $d(u, v) \geq D$ and so $d(u, v) > \frac{D}{8R}(d(u, a_1) + d(a_1, v))$.

Now by continuity we can take $u \in [a_n, a_1]$ and $v \in [a_1, a_2]$ such that $d(u, v) = \frac{D}{8R}(d(u, a_1) + d(a_1, v))$, but with $d(u, a_1) + d(a_1, v)$ maximal. Then for all $(u', v') \in [a_n, u] \times [v, a_2]$ such that $(u, v) \neq (u', v')$, we have $d(u', v') > \frac{D}{8R}(d(u', a_1) + d(a_1, v'))$. Indeed if not, by continuity since $d(a_n, a_2) > \frac{D}{8R}(d(a_n, a_1) + d(a_1, a_2))$ we could find $(u'', v'') \in [a_n, u'] \times [v', a_2]$ such that $d(u'', v'') = \frac{D}{8R}(d(u'', a_1) + d(a_1, v''))$,

contradicting the maximality of $d(u, a_1) + d(a_1, v)$. Taking $b_1 = u$ and $c_1 = v$, we do have the desired properties for $i = 1$.

Similarly, for every $i \in \{2, \dots, n-1\}$ one can find $b_i \in [c_{i-1}, a_i]$ and $c_i \in [a_i, a_{i+1}]$ such that $d(b_i, c_i) = \frac{D}{8R}(d(b_i, a_i) + d(a_i, c_i))$, and such that $d(b_i, a_i) + d(a_i, c_i)$ is maximal in the sense that for every $x \in [a_{i-1}, b_i]$ and $y \in [c_i, a_{i+1}]$ we have that $d(x, y) \geq \frac{D}{8R}(d(x, a_i) + d(a_i, y))$. This can be done repeating the previous argument, and based on the fact that

$$d(c_{i-1}, a_{i+1}) \geq d(b_{i-1}, a_{i+1}) - d(b_{i-1}, c_{i-1}) \geq \frac{3D}{4}.$$

Finally, we take $b_n \in [c_{n-1}, a_n]$ and $c_n \in [a_n, b_1]$ as previously, completing the trip around $[a_1, \dots, a_n]$ (this time, this relies on the fact that $d(c_{n-1}, b_1) \geq D/2$). \square_{claim}

Now note that the length ℓ of C , which is the sum of its edge lengths, lies between $\frac{Dn}{2} \geq 2$ and nR since each side $[c_i, b_{i+1}]$ is of length at least $D - D/4 - D/4 = \frac{D}{2}$ and the sides of $[a_1, \dots, a_n]$ have length at most R . Denote the path metric on C by d_C . Then (C, d_C) is isometric to the cycle of real length ℓ , i.e. to $\mathbb{R}/\ell\mathbb{Z}$ equipped with the quotient metric (for $\ell \in \mathbb{N}$, this is isometric to the continuous path metric on the graph C_ℓ). So we can find a 1-Lipschitz map $(C_{\lceil \ell \rceil}, d_{C_{\lceil \ell \rceil}}) \rightarrow (C, d_C)$ which contracts distances by at most the factor $\frac{\lceil \ell \rceil}{\ell} \leq \frac{3}{2}$. In order to prove the theorem, we will now show that the inclusion (C, d_C) in (X, d) contracts distances by at most the factor $\frac{2nR}{D}$, since by composing this will yield a 1-Lipschitz map $C_{\lceil \ell \rceil} \rightarrow (X, d)$ which contracts distances by at most the factor $\frac{3nR}{D}$ and the length of the cycle is at least $\frac{Dn}{2}$.

For each pair of points $x \in [c_{i-1}, b_i]$ and $y \in [c_i, b_{i+1}]$, we have $d(x, y) \geq \frac{D}{8R}(d(x, a_i) + d(a_i, y))$. Since $d_C(x, y) \leq d(x, a_i) + d(a_i, y)$, we deduce that

$$d_C(x, y) \leq \frac{8R}{D}d(x, y). \quad (26)$$

We now give a similar bound for two points on adjacent sides of C . Towards this, let $x \in [b_i, c_i]$ and $y \in [c_i, b_{i+1}]$. By Claim 9.5, we have $d(y, b_i) \leq \frac{D}{8R}(d(y, a_i) + d(a_i, b_i))$, $d(y, a_i) = d(y, c_i) + d(c_i, a_i)$, and $d(b_i, c_i) = \frac{D}{8R}(d(c_i, a_i) + d(a_i, b_i))$. Using successively these facts and the triangle inequality, we deduce:

$$\begin{aligned} d(x, y) &\geq d(y, b_i) - d(x, b_i) \\ &\geq \frac{D}{8R}(d(y, a_i) + d(a_i, b_i)) - d(b_i, c_i) + d(x, c_i) \\ &= \frac{D}{8R}(d(y, c_i) + d(c_i, a_i) + d(a_i, b_i)) - d(b_i, c_i) + d(x, c_i) \\ &= \frac{D}{8R}d(y, c_i) + d(x, c_i) \\ &\geq \frac{D}{8R}(d(y, c_i) + d(x, c_i)) \\ &= \frac{D}{8R}d_C(x, y). \end{aligned}$$

Hence (26) is also valid for such pairs (x, y) . The case where $x \in [b_i, c_i]$ and $y \in [c_i, b_{i+1}]$ is analogous. Finally, for $x, y \in C$ not in any of the previous cases, there exist two non-adjacent sides S_x and S_y in the n -gon $[a_1, \dots, a_n]$ such that $d(x, S_x) \leq \frac{D}{4}$ and $d(y, S_y) \leq \frac{D}{4}$, and hence $d(x, y) \geq \frac{D}{2}$. As the perimeter of C is at most nR , it follows that

$$d_C(x, y) \leq \frac{2nR}{D}d(x, y). \quad (27)$$

The lemma now results from the combination of (26) and (27). \square

Lemma 9.6. *Let (X, d) be a geodesic space and let $r > 0$. Assume that X contains a quadrilateral $[a_1, a_2, a_3, a_4]$ with the following properties:*

- $d(x, y) \geq \frac{r}{2}$ for every $x \in [a_2, a_3]$ and $y \in [a_1, a_4]$;
- $d(a_1, a_2) \leq r$;
- $d(a_2, a_3) \geq 4r$;
- $d(a_3, a_4) \leq 2r$.

Then there exist a cycle C of length at least $\lceil r \rceil$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances by at most a factor of 360 in X .

Proof. First note that by the triangle inequality, $d(x, y) \geq r$ for all $x \in [a_1, a_2]$ and $y \in [a_3, a_4]$. We deduce that the distance between pairs of opposite sides are at least $r/2$.

Let us first consider the case $d(a_2, a_3) \leq 12r$, in which case we deduce that the longest side has length at most $15r$. By Lemma 9.4, it now follows that there exist a cycle C of length at least $\lceil r \rceil$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances at most by a factor of $(3 \cdot 4 \cdot 15r)/(r/2) = 360$ in X .

Let $m \in \mathbb{N}$ be such that $2mr \leq d(a_2, a_3) < 2(m+1)r$. We will prove the lemma by induction on $m \geq 0$, noting that the previous paragraph actually covers the steps $m = 0, \dots, 5$.

Let $m \geq 5$, and assume that the lemma is true for all $r > 0$, and all quadrilaterals $[a_1, a_2, a_3, a_4]$ with $d(a_2, a_3) < 2mr$. Let $r > 0$, and consider a quadrilateral $[a_1, a_2, a_3, a_4]$ such that $d(a_2, a_3) < 2(m+1)r$. We may as well assume $d(a_2, a_3) \geq 12r$ because otherwise our previous analysis applies.

Case 1. Assume that there exists $b \in [a_2, a_3]$ such that $d(a_2, b) \geq 2r$ and $d(a_3, b) \geq 2r$ that satisfies $d(b, [a_1, a_4]) < r$. Let $c \in [a_1, a_4]$ be such that $d(b, c) < r$. Now, we have that $d(a_2, b) \leq 2mr$ and $d(b, a_3) \leq 2mr$, and at least of them must be more than $4r$ since $d(a_2, a_3) \geq 12r$. From this it follows that either $[a_1, a_2, b, c]$ or $[c, b, a_3, a_4]$ satisfies the induction hypothesis.

Case 2. Else, for every $b \in [a_2, a_3]$, if $d(b, [a_1, a_4]) < r$, then b must be $2r$ -close to either a_2 or a_3 . We let $b_2 \in [a_2, a_3]$ be the furthest point from a_2 such that $d(a_2, b_2) \leq 2r$ and $d(b_2, [a_1, a_4]) \leq r$. Similarly, we let $b_3 \in [a_2, a_3]$ be the furthest point from a_3 which satisfies $d(b_2, a_3) \leq 2r$, and $d(b_2, [a_1, a_4]) \leq r$. Then for every $x \in [b_1, b_2]$, we have $d(x, [a_1, a_4]) \geq r$. Recall that since $d(a_2, a_3) \geq 12r$, we have $d(b_2, b_3) \geq 8r$. We then find $b_1 \in [a_1, a_4]$ such that $d(b_1, b_2) \leq r$, and $b_4 \in [a_1, a_4]$ such that $d(b_4, b_3) \leq r$. In particular, we have :

- $d(x, y) \geq r$ for every $x \in [b_2, b_3]$ and $y \in [b_1, b_4]$;
- $d(b_1, b_2) \leq 2r$;
- $d(b_2, b_3) \geq 8r$;
- $d(b_3, b_4) \leq 4r$.

This means that we can apply our induction hypothesis to $r' = 2r$, noting that $d(b_2, b_3) < 2(m+1)r < 2mr'$. So, there exist a cycle C of length at least $\lceil 4r \rceil$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances at most by a factor of 360 in X , which concludes the proof by induction. \square

Corollary 9.7. *Let (X, d) be a geodesic space and let $R \geq r > 0$. Assume that X contains a quadrilateral $[a_1, a_2, a_3, a_4]$ with the following properties:*

- $d(x, y) \geq \frac{r}{2}$ for every $x \in [a_2, a_3]$ and $y \in [a_1, a_4]$;
- $d(a_1, a_2) \leq r$;

- $d(a_2, a_3) \geq \max\{12(R - r), 4r\}$;
- $d(a_3, a_4) \leq R$.

Then there exist a cycle C of length at least $\lceil r \rceil$ and a 1-Lipschitz-embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances by at most a factor of 360 in X .

Proof. By Lemma 9.6, the corollary is true for $R \leq 2r$, so suppose that $R \geq 2r$ and let $m \in \mathbb{N}$ be such that $2^m r < R \leq 2^{m+1} r$. We now prove the claim by induction on m . If $m = 0$, then $R \leq 2r$ and the corollary is true. Next, suppose the corollary is true for $R \leq 2^m r$. We claim that it is then also true for $R \leq 2^{m+1} r$, which concludes the proof by induction.

Now we take $b \in [a_2, a_3]$ such that $d(b, [a_1, a_4]) = r$ and $d(a_2, b)$ is maximal. Also take $c \in [a_1, a_4]$ such that $d(b, c) = r$. If $d(a_2, b) \geq 4r$, then by Lemma 9.6 the quadrilateral $[a_1, a_2, b, c]$ satisfies the claim, which concludes the proof by induction. On the other hand, if $d(a_2, b) < 4r$, then $d(b, a_3) \geq d(a_2, a_3) - d(b, a_2) > 12(R - r) - 4r = 12(R - 2r) + 8r \geq \max\{12(R - 2r), 8r\}$, and we can apply the induction hypothesis to the quadrilateral $[c, b, a_3, a_4]$ as $d(c, b) \leq 2r$, $d(a_3, a_4) \leq R$ and $d(x, [a_1, a_4]) \geq r$ for every $x \in [a_2, a_3]$, from which it follows that there exist a cycle C of length at least $\lceil r \rceil$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances at most by a factor 360 in X , which concludes the proof by induction. \square

Proposition 9.8. *Let (X, d) be a geodesic space that is not δ -hyperbolic. Then, there exist a cycle C of length at least $\lceil \frac{\delta}{15} \rceil$ and a 1-Lipschitz embedding $\varphi: (C, d_C) \rightarrow (X, d)$ that contracts distances by at most a factor of 17820 in X .*

Proof. As (X, d) is not δ -hyperbolic, there exists a geodesic triangle $[a, b, c]$ such that the minimum $D > 0$ such that $[a, b]$ is contained in the D -neighborhood of the union of $[a, c]$ and $[b, c]$ is $> \delta$. Let $x \in [a, b]$ be such that $d(x, [a, c] \cup [b, c]) = D$.

As the map $y \mapsto d(y, [a, c] \cup [b, c])$ is continuous, there exists $x_a \in [a, x]$ such that $d(x_a, [a, c] \cup [b, c]) = \frac{D}{15}$. Now, take $y_a \in [a, c] \cup [b, c]$ such that $d(x_a, y_a) = \frac{D}{15}$. Similarly, let $x_b \in [x, b]$ and $y_b \in [a, c] \cup [b, c]$ such that $d(x_b, [a, c] \cup [b, c]) = d(x_b, y_b) = \frac{D}{15}$. Moreover, take x_a and x_b above such that $d(x_a, x)$ and $d(x_b, x)$ are minimal.

Case 1. Suppose first that y_a and y_b are on the same side $[a, c]$ or $[b, c]$. To begin, note that $d(x_a, x_b) = d(x_a, x) + d(x, x_b) \geq \frac{28D}{15}$ since $D \leq d(x, y_a) \leq d(x, x_a) + \frac{D}{15}$ and $D \leq d(x, y_b) \leq d(x, x_b) + \frac{D}{15}$. Thus, we have for $[y_a, x_a, x_b, y_b]$ that $d(y_a, x_a) = \frac{D}{15} \leq \frac{2D}{15}$, $d(x_a, x_b) \geq \frac{28D}{15}$, $d(x_b, y_b) = \frac{D}{15} \leq \frac{2D}{15}$, and $d(x', [y_a, y_b]) \geq \frac{1}{2} \frac{2D}{15}$ for all $x' \in [x_a, x_b]$. Now, by Corollary 9.7 it follows by taking $r = \frac{2D}{15}$ and $R = r$ that there exist a cycle C of length at least $\lceil \frac{2D}{15} \rceil$ and a 1-Lipschitz embedding of C into X that contracts distances by at most a factor of 360 in X .

Case 2. Next, suppose that y_a and y_b are not on the same geodesic $[a, c]$ or $[b, c]$, but that $d(y_a, [y_b, c]) \leq \frac{3D}{15}$. Let $u \in [y_b, c]$ be such that $d(y_a, u) \leq \frac{3D}{15}$. Thus, we have for $[y_b, x_b, x_a, u]$ that $d(y_b, x_b) = \frac{D}{15} \leq \frac{2D}{15}$, $d(x_b, x_a) \geq \frac{28D}{15} \geq \max\{12(\frac{4D}{15} - \frac{2D}{15}), 4\frac{2D}{15}\}$, $d(x_a, u) \leq d(x_a, y_a) + d(y_a, u) \leq \frac{4D}{15}$, and $d(x', [y_b, u]) \geq \frac{1}{2} \frac{2D}{15}$ for every $x' \in [x_b, x_a]$. Thus, by Corollary 9.7 it follows by taking $r = \frac{2D}{15}$ and $R = 2r$ that there exist a cycle C of length at least $\lceil \frac{2D}{15} \rceil$ and a 1-Lipschitz embedding of C into X that contracts distances by at most by a factor of 360 in X .

Case 3. Finally, suppose that y_a and y_b are not on the same geodesic $[a, c]$ or $[b, c]$, and that $d(y_a, [y_b, c]) > \frac{3D}{15}$ and $d(y_b, [y_a, c]) > \frac{3D}{15}$. Take $z_a \in [y_a, c]$ and $z_b \in [y_b, c]$ such that $d(z_a, z_b) = \frac{D}{15}$ and $d(z_a, y_a)$ is minimal. Now consider the hexagon $[x_a, x_b, y_b, z_b, z_a, y_a]$. Recall that the sides $[x_a, x_b]$, $[y_a, z_a]$ and $[y_b, z_b]$ are $\frac{D}{15}$ -disjoint and have length of at least $\frac{2D}{15}$, and that the length of the other sides is $\frac{D}{15}$.

Claim 9.9. We claim that all non-adjacent sides are $\frac{D}{45}$ -disjoint.

Proof of the claim. Suppose w and w' are on non-adjacent sides. Recalling that the sides $[x_a, x_b]$, $[y_a, z_a]$ and $[y_b, z_b]$ are $\frac{D}{15}$ -disjoint, we may assume without loss of generality that w is not on one of these sides, say $w \in [z_a, z_b]$. If $w' \in [x_a, x_b]$, then either $d(w, w') \geq d(w', [y_a, z_a]) - d(w, [y_a, z_a]) \geq \frac{D}{15} - \frac{D}{30} = \frac{D}{30}$ or $d(w, w') \geq d(w', [y_b, z_b]) - d(w, [y_b, z_b]) \geq \frac{D}{15} - \frac{D}{30} = \frac{D}{30}$ as it can not be that both $d(w, [y_a, z_a])$ and $d(w, [y_b, z_b])$ are strictly greater than $\frac{D}{30}$ because $d(z_a, z_b) = \frac{D}{15}$. So, suppose w' is either in $[x_a, y_a]$ or in $[x_b, y_b]$. Without loss of generality, we can assume that $w' \in [x_a, y_a]$. Now, if $d(x_a, w') + d(z_b, w) \leq \frac{2D}{45}$, then $d(w, w') \geq d(z_b, x_a) - d(x_a, w') - d(w, z_b) \geq d(z_b, x_a) - \frac{2D}{45} \geq \frac{D}{45}$ since $d(z_b, x_a) \geq \frac{2D}{15}$, and we are done. On the other hand, if $d(x_a, w') + d(z_b, w) \geq \frac{2D}{45}$ then $d(y_a, w') + d(z_a, w) \leq \frac{4D}{45}$ since the side length of $[x_a, y_a]$ and $[z_a, z_b]$ is $\frac{D}{15}$, and thus $d(w, w') \geq d(w, y_a) - d(y_a, w') \geq d(z_a, y_a) - (d(w, z_a) + d(y_a, w')) \geq d(z_a, y_a) - \frac{2D}{45} = \frac{2D}{15} - \frac{2D}{45} = \frac{2D}{45}$, and we are done. In other words, all non-adjacent sides are $\frac{D}{45}$ -disjoint, which completes the proof. \square_{claim}

We can now finish the proof of the third case. First, if every side has length at most $22D$, then by Lemma 9.4 there exist a cycle C of length at least $\lceil \frac{6D}{2 \cdot 45} \rceil = \lceil \frac{D}{15} \rceil$ and a 1-Lipschitz embedding of C into X that contracts distances by at most a factor of $(3 \cdot 6 \cdot 22D)/(D/45) = 17820$ in X . On the other hand, if at least one of the sides has length more than $22D$, take u as the midpoint of that side, and $v \in [a, b, c]$ such that $d(u, v) \leq D$ where u and v are not on the same side of the triangle. Then, we find two adjacent corners w_u and w_v in $[x_a, x_b, y_b, z_b, z_a, y_a]$ such that $[u, v]$ splits this hexagon into the quadrilateral $[w_v, w_u, u, v]$ and a new hexagon, where $d(w_v, w_u) \leq \frac{2D}{15}$, $d(w_u, u) > \frac{22D}{2} = 11D \geq \max\{12(D - \frac{2D}{15}), 4\frac{2D}{15}\}$, $d(u, v) \leq D$ and $d(x', [w_v, v]) \geq \frac{1}{2} \frac{2D}{15}$ for each $x' \in [w_u, u]$. Thus, once again by Corollary 9.7 it follows by taking $r = \frac{2D}{15}$ and $R = D$ that there exist a cycle C of length at least $\lceil \frac{2D}{15} \rceil$ and a 1-Lipschitz embedding of C into X that contracts distances by at most by a factor of 360 in X . \square

We can finally prove Theorem 18, which we restate as a corollary of the above proposition.

Corollary 9.10. *Let (X, d) be a geodesic space that is not hyperbolic. Then, there exist arbitrarily large integers n and 1-Lipshitz embeddings $\varphi_n: C_n \rightarrow (X, d)$ which contract distances by at most a factor of 17820 in X .*

Proof. Consider a sequence δ_n tending to infinity and apply Proposition 9.8 for every n to the non δ_n -hyperbolic space (X, d) . \square

9.2 Rigidity of hyperbolicity

We now prove rigidity results, saying that hyperbolicity is preserved under cobounded measure equivalence couplings satisfying certain integrability conditions. To simplify the exposition, we choose to state two results, but the second one should be seen as a degenerate version of the first one (where one of the conditions becomes an L^∞ -condition). Anyway the two results share basically the same proof, which consists in confronting Proposition 9.2 with Theorem 18. Since these results are quite technical, we shall first deduce two more appealing corollaries (namely Theorem 11 and Theorem 13 from the introduction).

Let us start with some notation: given a group Γ equipped with a finitely generated subset S_Γ , we denote the growth function $\text{Vol}_{S_\Gamma}(n) = |S_\Gamma^n|$ and define the associated **volume entropy** as $\text{Ent}(S_\Gamma) = \limsup_{r \rightarrow \infty} \frac{\log(\text{Vol}_{S_\Gamma}(r))}{r}$.

Theorem 9.11. *Let Γ and Λ be two finitely generated groups such that Γ is δ -hyperbolic. We let $L \geq 1$ and φ, ψ and r be increasing unbounded functions. Assume that the following conditions are satisfied:*

$$\lim_{n \rightarrow \infty} \frac{n^2 r(n) \text{Vol}_{S_\Gamma}(r(n))}{\varphi(n/r(n))} \rightarrow 0; \quad (28)$$

and⁶ for all large enough n ,

$$\frac{r(n)}{17820} \geq 8(\delta + 1) \log_2 n + 3\psi^{-1}(3Ln). \quad (29)$$

Assume that (Ω, μ) is a cobounded measure equivalence coupling from Γ to Λ , normalized so that $\mu(X_\Gamma) = 1$, and such that the associated cocycles $\alpha: \Gamma \times X_\Lambda \rightarrow \Lambda$ and $\beta: \Lambda \times X_\Gamma \rightarrow \Gamma$ satisfy the following properties.

(i) for all $s \in S_\Gamma$,

$$\int_{X_\Lambda} \varphi(|\alpha(s, x)|_{S_\Lambda}) d\mu(x) < \infty;$$

(ii) for all $t \in S_\Lambda$

$$\int_{X_\Gamma} \psi(|\beta(t, x)|_{S_\Gamma}) d\mu(x) \leq L.$$

Then Λ is hyperbolic.

Assuming that β is bounded, we have the following variant.

Theorem 9.12. *Let Γ and Λ be two finitely generated groups such that Γ is δ -hyperbolic. We let φ, ψ and r be increasing unbounded functions. Assume that (28) holds, and that for all large enough n ,*

$$\frac{r(n)}{17820} \geq 13(\delta + 1) \log n. \quad (30)$$

Assume that (Ω, μ) is a cobounded measure equivalence coupling from Γ to Λ , normalized so that $\mu(X_\Gamma) = 1$, and such that the associated cocycles $\alpha: \Gamma \times X_\Lambda \rightarrow \Lambda$ and $\beta: \Lambda \times X_\Gamma \rightarrow \Gamma$ satisfy the following properties.

(i) for all $s \in S_\Gamma$,

$$\int_{X_\Lambda} \varphi(|\alpha(s, x)|_{S_\Lambda}) d\mu(x) < \infty;$$

(ii) for all $t \in S_\Lambda$

$$|\beta(t, \cdot)|_{S_\Gamma} \in L^\infty(X_\Gamma).$$

Then Λ is hyperbolic.

Theorem 11 is an immediate consequence of the following corollary of Theorem 9.12.

Corollary 9.13. *Let Γ be a finitely generated δ -hyperbolic group and let $p > 75\delta \text{Ent}(S_\Gamma) + 2$. Assume that there exists a cobounded (L^p, L^∞) -integrable measure equivalence coupling from Γ to any finitely generated group Λ . Then Λ is hyperbolic.*

Proof. We apply Theorem 9.12 with $p = 75\delta(\text{Ent}(S_\Gamma) + \varepsilon) + 2$ for some $\varepsilon > 0$. We let $r(n) = 75\delta \log n$. By definition of $\text{Ent}(S_\Gamma)$, there exists have $\text{Vol}_{S_\Gamma}(r(n)) = o(e^{(\text{Ent}(S_\Gamma) + \varepsilon/2)r(n)})$. Hence we deduce that

$$\text{Vol}_{S_\Gamma}(r(n)) = o(n^{75\delta(\text{Ent}(S_\Gamma) + \varepsilon/2)}),$$

which combined with the fact that $\phi(t) = t^{75\delta(\text{Ent}(S_\Gamma) + \varepsilon) + 2}$ implies that (28) is satisfied. Thus, Λ is hyperbolic. \square

⁶The constant 17820 in (29) comes from the use of Theorem 18, while the right-hand term comes from the use of Proposition 9.2.

Corollary 9.14. *Let Γ be a finitely generated hyperbolic group. For every $p > q > 0$ such that if there is a cobounded (φ, ψ) -integrable measure equivalence coupling from Γ to any finitely generated group Λ where $\varphi(t) = \exp(t^p)$ and $\psi(t) = t^{1+1/q}$, then Λ is also hyperbolic.*

Proof. First we take the cobounded (φ, ψ) -integrable measure equivalence coupling $(\Omega, X_\Gamma, X_\Lambda, \mu)$ and normalize such that $\mu(X_\Gamma) = 1$. We pick η strictly between q and p , we let $r(n) = n^{\frac{\eta}{1+\eta}}$, and we define

$$L = \max_{s \in S_\Lambda} \int_{X_\Gamma} \psi(|\beta(s, x)|_{S_\Gamma}) d\mu(x).$$

Note that $\psi^{-1}(t) = t^{\frac{q}{1+q}}$. So (29) follows from the fact that $\eta > q$. Finally, take $\varepsilon > 0$ such that

$$\max_{s \in S_\Gamma} \int_{X_\Lambda} \varphi_\varepsilon(|\alpha(s, x)|) d\mu(x) < \infty,$$

where $\varphi_\varepsilon(t) = \varphi(\varepsilon t)$. Note that $n/r(n) = n^{\frac{1}{1+\eta}}$. Hence $\varphi_\varepsilon(n/r(n)) = \exp(\varepsilon^p n^{\frac{p}{1+\eta}})$, while

$$\text{Vol}_{S_\Gamma}(r(n)) \leq |S_\Gamma| n^{\frac{\eta}{1+\eta}}$$

Hence since $\eta < p$, we have

$$\text{Vol}_{S_\Gamma}(r(n))/\varphi_\varepsilon(n/r(n)) = O(n^{-k})$$

for any $k > 0$. So (28) is satisfied and we conclude by Theorem 9.11 that Λ is hyperbolic. \square

Proof of Theorems 9.11 and 9.12. Let us start by strengthening the coboundedness condition.

Claim 9.15. On replacing (29) and (30) respectively by the slightly weaker conditions

$$\frac{r(n)}{17820} \geq 4\delta \log_2 n + 3\psi^{-1}(3Ln) - 2, \quad (31)$$

and

$$\frac{r(n)}{17820} \geq 13\delta \log n, \quad (32)$$

it is enough to prove Theorem 9.11 and Theorem 9.12 under the assumption that $X_\Gamma \subseteq X_\Lambda$.

Proof of the claim. Assuming we have a coupling satisfying the conditions of Theorem 9.11 (resp. Theorem 9.12), we build a new coupling satisfying (31) (resp. (32)) with $X_\Gamma \subseteq X_\Lambda$.

Since our initial coupling is cobounded, there exists a finite subset F of Λ such that $X_\Gamma \subseteq F * X_\Lambda$. Consider the new coupling space $\tilde{\Omega} := \Omega \times F$, let K be a finite group which acts simply transitively on F , and let $\tilde{\Gamma} = \Gamma \times K$ act on $\tilde{\Omega}$ by $(\gamma, k) * (\omega, f) = (\gamma * \omega, k \cdot f)$. This action is smooth, and we take as a fundamental domain the set

$$\tilde{X}_{\tilde{\Gamma}} := \bigsqcup_{f \in F} (X_\Gamma \cap f * X_\Lambda) \times \{f\}$$

The Λ -action on $\tilde{\Omega}$ is the action on the first coordinate; a fundamental domain is provided by

$$\tilde{X}_\Lambda = \bigsqcup_{f \in F} (f * X_\Lambda) \times \{f\}.$$

Viewing both Γ and K as subgroups of $\tilde{\Gamma}$, the latter has $S_{\tilde{\Gamma}} = S_\Gamma \cup K$ as a finite generating set. In fact, with this generating set $\tilde{\Gamma}$ is $\tilde{\delta}$ -hyperbolic, with $\tilde{\delta} = \delta + 1$. Observe that the volume growth of $\tilde{\Gamma}$ is at least $|K|$ times that of Γ .

Next note that condition (i) is still met by this new generating set. Indeed \tilde{X}_Λ is L^∞ -equivalent to the fundamental domain $X_\Lambda \times F$, and for all $\gamma \in \Gamma$, $x \in X_\Lambda$ and $f \in F$ we have $d_{S_\Lambda}(\gamma * (x, f), \gamma \cdot (x, f)) =$

$d_{S_\Lambda}(\gamma * x, \gamma \cdot x)$, so for all $\tilde{\gamma} \in S_{\tilde{\Gamma}}$ we have that $\tilde{\gamma} * X_\Lambda \times F$ is φ -equivalent to $X_\Lambda \times F$, so the same is true of \tilde{X}_Λ .

For condition (ii), we have, by construction, for all $x \in X_\Gamma$ and all $f, f' \in F$ that the distance $d_{S_{\tilde{\Gamma}}}((x, f), (x, f')) \leq 1$, so for every $\lambda \in \Lambda$ we have

$$d_{S_{\tilde{\Gamma}}}(\lambda \cdot (x, f), \lambda * (x, f)) \leq 1 + d_{S_\Gamma}(\lambda \cdot x, \lambda * x) = 1 + |\beta(\lambda, x)|,$$

hence the new coupling satisfies the same conditions on replacing $\psi(t)$ by $\tilde{\psi}(t) = \psi(\max\{t-1, 0\})$ in Theorem 9.11. Note that for n large enough, $\psi^{-1}(3Ln) \geq 1$ and $\tilde{\psi}(t) = \psi(t-1)$ for $t \geq 1$, from which we easily deduce that for large enough n ,

$$\frac{r(n)}{17820} \geq 4\tilde{\delta} \log_2 n + 3\tilde{\psi}^{-1}(3Ln) - 3.$$

So the claim is proved. □_{claim}

From now on, we assume that $X_\Gamma \subseteq X_\Lambda$ and we normalize the measure so that $\mu(X_\Gamma) = 1$. Suppose by contradiction that Λ is not hyperbolic. By Theorem 18 we can take a cycle C_n of arbitrary large length n , and a bi-Lipschitz embedding $\iota: C_n \rightarrow \Lambda$ that are 1-Lipschitz and contracts distances at most by a factor 17820. In what follows we consider C_n as a subset of Λ . Let K be such that $\int_{X_\Lambda} \varphi(|\alpha(s, x)|_{S_\Lambda}) d\mu(x) \leq K$ for all $s \in S_\Gamma$.

For every $x \in X_\Gamma$ we denote by $b_x: \Lambda \rightarrow \Gamma$ the map defined by $b_x(\lambda) = \beta(\lambda^{-1}, x)^{-1}$ for every $\lambda \in \Lambda$. We will use throughout the following immediate consequence of the cocycle relation: let $u, v \in \Lambda$, we have

$$b_x(v)^{-1}b_x(u) = \beta(u^{-1}v, u^{-1} \cdot x). \quad (33)$$

Upper estimates for the restriction of b_x to C_n . In the case of Theorem 9.12, we trivially have that b_x is a.e. L -Lipschitz for some constant L .

Under the assumption of Theorem 9.11, we claim that with probability at least $2/3$, the restriction of b_x to C_n is $\psi^{-1}(3Ln)$ -Lipschitz. Here we use the integrability condition for β . For every u and v adjacent in C_n there exists an $s_{u,v} \in S_\Lambda$ such that $v = us_{u,v}$. By (33), we have:

$$d_{S_\Gamma}(b_x(v), b_x(u)) = |b_x(v)^{-1}b_x(u)|_{S_\Gamma} = |\beta(s_{u,v}, u^{-1} \cdot x)|_{S_\Gamma}.$$

Next consider for any $u \in C_n$ the set of all $x \in X_\Gamma$ such that $\psi(|\beta(s_{u,v}, u^{-1} \cdot x)|_{S_\Gamma}) \geq 3Ln$. By Markov's inequality, these sets have measure at most $\frac{1}{3n}$ and therefore the set of all $x \in X_\Gamma$ such that b_x is $\psi^{-1}(3Ln)$ -Lipschitz in restriction to C_n has measure at least $1 - n \cdot \frac{1}{3n} = \frac{2}{3}$. So our claim follows.

Lower estimates for the restriction of b_x to C_n . Providing lower estimates on the quasi-isometric embedding constants is more involved as this requires to apply the cocycle α to C_n . We shall use the inverse relation between α and β and the inclusion $X_\Gamma \subseteq X_\Lambda$: for all $x \in X_\Gamma$ and $\lambda \in \Lambda$

$$\alpha(\beta(\lambda, x), x) = \lambda \quad (34)$$

We claim that we have the following key inequality.

Claim 9.16. For every $R > 0$, and u and v in Λ , we have

$$\mu\left(\left\{x \in X_\Gamma : d_{S_\Gamma}(b_x(v), b_x(u)) \leq R\right\}\right) \leq KR \frac{\text{Vol}_{S_\Gamma}(R)}{\varphi\left(\frac{d_{S_\Lambda}(u,v)}{R}\right)}.$$

Proof of the claim. For any $\gamma \in \Gamma$, we define the set

$$A_\gamma = \{x \in X_\Gamma : b_x(v)^{-1}b_x(u) = \gamma\}.$$

By (34) and (33), we have that for every $x \in A_\gamma$,

$$\alpha(\gamma, u^{-1} \cdot x) = \alpha(\beta(u^{-1}v, u^{-1} \cdot x), u^{-1} \cdot x) = u^{-1}v,$$

from which we deduce that $|\alpha(\gamma, u^{-1} \cdot x)|_{S_\Lambda} = d_{S_\Lambda}(u, v)$.

Let us start giving an upper bound of $\mu(A_\gamma)$ as a function of $|\gamma|$. Write $\gamma = s_1 \dots s_{|\gamma|_{S_\Gamma}}$ with $s_i \in S_\Gamma$. By the cocycle relation and the triangular inequality, there exists an i such that the set

$$\left\{ x \in A_\gamma : \left| \alpha \left(s_i, s_{i+1} \dots s_{|\gamma|_{S_\Gamma}} \cdot (u^{-1} \cdot x) \right) \right|_{S_\Lambda} \geq \frac{d_{S_\Lambda}(u, v)}{|\gamma|_{S_\Gamma}} \right\}$$

has measure at least $\frac{\mu(A_\gamma)}{|\gamma|_{S_\Gamma}}$. Letting $s = s_i$, we have that

$$\mu \left(\left\{ y \in X_\Lambda : |\alpha(s, y)|_{S_\Lambda} \geq \frac{d_{S_\Lambda}(u, v)}{|\gamma|_{S_\Gamma}} \right\} \right) \geq \frac{\mu(A_\gamma)}{|\gamma|_{S_\Gamma}},$$

from which we deduce the following upper bound on the measure of A_γ :

$$\mu(A_\gamma) \leq |\gamma|_{S_\Gamma} \mu \left(\left\{ y \in X_\Lambda : |\alpha(s, y)|_{S_\Lambda} \geq \frac{d_{S_\Lambda}(u, v)}{|\gamma|_{S_\Gamma}} \right\} \right)$$

By Markov's inequality, we deduce

$$\mu(A_\gamma) \leq \frac{K|\gamma|_{S_\Gamma}}{\varphi \left(\frac{d_{S_\Lambda}(u, v)}{|\gamma|_{S_\Gamma}} \right)}.$$

Using that φ is non-decreasing, we get that for all $R > 0$,

$$\begin{aligned} \mu(\{x \in X_\Gamma : |b_x(v)^{-1}b_x(u)|_{S_\Gamma} \leq R\}) &= \sum_{\gamma \in B_\Gamma(e_\Gamma, R)} \mu(A_\gamma) \\ &\leq \sum_{\gamma \in B_\Gamma(e_\Gamma, R)} \frac{K|\gamma|_{S_\Gamma}}{\varphi \left(\frac{d_{S_\Lambda}(u, v)}{R} \right)} \\ &\leq KR \frac{\text{Vol}_{S_\Gamma}(R)}{\varphi \left(\frac{d_{S_\Lambda}(u, v)}{R} \right)}. \end{aligned}$$

So the claim is proved. □_{claim}

Applying Claim 9.16 with $R = \frac{r(n)}{n} d_{S_\Lambda}(u, v)$, $u, v \in C_n$, observing that $d_{S_\Lambda}(u, v) \leq n$, we obtain

$$\mu \left(\left\{ x \in X_\Gamma : d_{S_\Gamma}(b_x(v), b_x(u)) \leq \frac{r(n)}{n} d_{S_\Lambda}(u, v) \right\} \right) \leq \frac{Kr(n) \text{Vol}_{S_\Gamma}(r(n))}{\varphi(n/r(n))}.$$

As there are at most n^2 pairs (u, v) in C_n , we deduce that

$$\mu \left(\left\{ x \in X_\Gamma : \exists u, v \in C_n : d_{S_\Gamma}(b_x(v), b_x(u)) \leq \frac{r(n)}{n} d_{S_\Lambda}(u, v) \right\} \right) \leq \frac{Kn^2 r(n) \text{Vol}_{S_\Gamma}(r(n))}{\varphi(n/r(n))}.$$

By (28), there exists n_0 such that for $n \geq n_0$, there exists a subset B of X_Γ of measure at least $2/3$ on which for all $u, v \in C_n$,

$$d_{S_\Gamma}(b_x(v), b_x(u)) \geq \frac{r(n)}{n} d_{S_\Lambda}(u, v).$$

Finally, for all x in the subset $A \cap B$ which has positive measure, we deduce for every $u, v \in C_n$ that

$$a_n d_{C_n}(u, v) \leq d_{S_\Gamma}(b_x(v), b_x(u)) \leq b_n d_{C_n}(u, v),$$

where in the case of Theorem 9.11, $a_n = \frac{r(n)}{17820n}$ and $b_n = \psi^{-1}(3Ln)$; and in the case of 9.12, $a_n = \frac{r(n)}{17820n}$ and $b_n = L$. In the first case, assuming (31), we have

$$\frac{r(n)}{17820} \geq 4\delta \log_2 n + 3\psi^{-1}(3Ln),$$

from which we deduce that for n large enough (as $b_n \rightarrow \infty$),

$$na_n \geq 4\delta \log_2 n + 3b_n > 4\delta \log_2(b_n n) + 4 + 2b_n,$$

which contradicts Proposition 9.2. Similarly, in the second case, we check that (32) contradicts Corollary 9.3. \square

References

- [Aus16] Tim Austin. Integrable measure equivalence for groups of polynomial growth. *Groups, Geometry, and Dynamics*, 10(1):117–154, February 2016.
- [Bau61] Gilbert Baumslag. Wreath products and finitely presented groups. *Mathematische Zeitschrift*, 75(1):22–28, December 1961.
- [Ben12] Itai Benjamini. Instability of the liouville property for quasi-isometric graphs and manifolds of polynomial volume growth. *Journal of Theoretical Probability*, 4:631–637, 2012.
- [BFS13] Uri Bader, Alex Furman, and Roman Sauer. Integrable measure equivalence and rigidity of hyperbolic lattices. *Inventiones mathematicae*, 194(2):313–379, November 2013.
- [Bri14] Jérémie Brieussel. Growth behaviors in the range $\alpha \in [1, 2]$. *Afrika Matematika*, 25(4):1143–1163, December 2014.
- [BST12] Itai Benjamini, Oded Schramm, and Ádám Timár. On the separation profile of infinite graphs. *Groups, Geometry, and Dynamics*, 6(4):639–658, November 2012.
- [BZ21] Brieussel and Zheng. Speed of random walks, isoperimetry and compression of finitely generated groups. *Annals of Mathematics*, 193(1):1, 2021.
- [CGP01] Thierry Coulhon, Alexander Grigor’yan, and Christophe Pittet. A geometric approach to on-diagonal heat kernel lower bounds on groups. *Annales de l’Institut Fourier*, 51(6):1763–1827, 2001.
- [Cou00] Thierry Coulhon. Random walks and geometry on infinite graphs. In *Lectures Notes on Analysis in Metric Spaces*, pages 5–30. Luigi Ambrosio, Francesco Serra Cassano, eds., Scuola Normale Superiore di Pisa, 2000.
- [CS93] Thierry Coulhon and Laurent Saloff-Coste. Isopérimétrie pour les groupes et les variétés. *Revista Matemática Iberoamericana*, 9(2):293–314, August 1993.
- [dC08] Yves de Cornulier. Dimension of asymptotic cones of Lie groups. *J. Topol.*, 1(2):342–361, 2008.
- [de 19] Mikael de la Salle. Strong property (T) for higher-rank lattices. *Acta Mathematica*, 223(1):151–193, September 2019.
- [DS06] A. Dranishnikov and J. Smith. Asymptotic dimension of discrete groups. *Fundamenta Mathematicae*, 189:27–34, 2006.
- [DuB] Kapovitch Michael Dru, u Badea, Cornelia. *Quasi-isometry rigidity of groups*.

- [Ers03] Anna Erschler. On Isoperimetric Profiles of Finitely Generated Groups. *Geometriae Dedicata*, 100(1):157–171, August 2003.
- [EZ17] Anna Erschler and Tianyi Zheng. Isoperimetric inequalities, shapes of Følner sets and groups with Shalom’s property H_{FD} . *arXiv:1708.04730 [math]*, August 2017.
- [FM98] B. Farb and L. Mosher. A rigidity theorem for the solvable Baumslag-Solitar groups. *Invent. Math.*, 131(2):419–451, 1998. With an appendix by Daryl Cooper.
- [Fur99] Alex Furman. Gromov’s Measure Equivalence and Rigidity of Higher Rank Lattices. *Annals of Mathematics*, 150(3):1059–1081, 1999.
- [Gab05] D. Gaboriau. Examples of groups that are measure equivalent to the free group. *Ergodic Theory and Dynamical Systems*, 25(6):1809–1827, December 2005.
- [Gri85] R. I. Grigorchuk. Degrees of growth of finitely generated groups and the theory of invariant means. *Mathematics of the USSR-Izvestiya*, 25(2):259, April 1985.
- [Gro07] Mikhael Gromov. *Metric Structures for Riemannian and Non-Riemannian Spaces*. Modern Birkhäuser Classics. Birkhäuser, Boston, 2007.
- [HMT20] David Hume, John M. Mackay, and Romain Tessera. Poincaré profiles of groups and spaces. *Revista Matemática Iberoamericana*, 36(6):1835–1886, March 2020.
- [Hum17] David Hume. A continuum of expanders. *Fundamenta Mathematicae*, 238:143–152, 2017.
- [IKT09] Adrian Ioana, Alexander Kechris, and Todor Tsankov. Subequivalence relations and positive-definite functions. *Groups, Geometry, and Dynamics*, pages 579–625, 2009.
- [Kid08] Yoshikata Kida. The mapping class group from the viewpoint of measure equivalence theory. *Memoirs of the American Mathematical Society*, 196(916):viii+190, 2008.
- [Kid10] Yoshikata Kida. Measure equivalence rigidity of the mapping class group. *Annals of Mathematics. Second Series*, 171(3):1851–1901, 2010.
- [KM04] A. S. Kechris and Benjamin D. Miller. *Topics in Orbit Equivalence*. Number 1852 in Lecture Notes in Mathematics. Springer, Berlin ; New York, 2004.
- [KV83] V. A. Kaimanovich and A. M. Vershik. Random Walks on Discrete Groups: Boundary and Entropy. *The Annals of Probability*, 11(3):457–490, August 1983.
- [LM18] François Le Maître. On a measurable analogue of small topological full groups. *Advances in Mathematics*, 332:235–286, July 2018.
- [LY15] R. Lyons and Peres Y. Poisson boundaries of lamplighter groups: Proof of the kaimanovich-vershik conjecture. 2015.
- [Now07] Piotr W. Nowak. On exactness and isoperimetric profiles of discrete groups. *Journal of Functional Analysis*, 243(1):323–344, February 2007.
- [OW80] Donald S. Ornstein and Benjamin Weiss. Ergodic theory of amenable group actions. I. The Rohlin lemma. *American Mathematical Society. Bulletin. New Series*, 2(1):161–164, 1980.
- [OW87] Donald S. Ornstein and Benjamin Weiss. Entropy and isomorphism theorems for actions of amenable groups. *Journal d’Analyse Mathématique*, 48(1):1–141, December 1987.
- [Pan89] Pierre Pansu. Métriques de Carnot-Carathéodory et Quasiisométries des Espaces Symétriques de rang un. *Annals of Mathematics*, 129(1):1–60, 1989.

- [Pit95] Christophe Pittet. Følner Sequences in Polycyclic Groups. *Revista Matemática Iberoamericana*, 11(3):675–685, December 1995.
- [Pit00] Ch Pittet. The isoperimetric profile of homogeneous Riemannian manifolds. *Journal of Differential Geometry*, 54(2):255–302, January 2000.
- [Sau02] Roman Sauer. *L²-Invariants of Groups and Discrete Measured Groupoids*. PhD thesis, Universität Münster, 2002.
- [Sha04] Yehuda Shalom. Harmonic analysis, cohomology, and the large-scale geometry of amenable groups. *Acta Mathematica*, 192(2):119–185, 2004.
- [Tes08] Romain Tessera. Large scale Sobolev inequalities on metric measure spaces and applications. *Revista Matemática Iberoamericana*, 24(3):825–864, April 2008.
- [V.A85] Kaimanovich V.A. Examples of noncommutative groups with nontrivial exit-boundary. *Journal of Soviet Mathematics volume*, 28:579–591, February 1985.
- [VS14] Kevin Verbeek and Subhash Suri. Metric Embedding, Hyperbolic Space, and Social Networks. In *Proceedings of the Thirtieth Annual Symposium on Computational Geometry, SOCG'14*, pages 501:501–501:510, New York, NY, USA, 2014. ACM.
- [Wei01] Benjamin Weiss. Monotileable amenable groups. In *Topology, Ergodic Theory, Real Algebraic Geometry: Rokhlin's Memorial*, pages 257–262. American Mathematical Soc., March 2001.
- [Woe05] Wolfgang Woess. Lamplighters, Diestel-Leader graphs, random walks, and harmonic functions. *Combin. Probab. Comput.*, 14(3):415–433, 2005.