

Multivariate geometric distributions, (logarithmically)
monotone sequences, and infinitely divisible laws
(with erratum by Natalia Shenkman)

Jan-Frederik Mai, Matthias Scherer, Natalia Shenkman*

*Department of Mathematics, Technische Universität München, Parkring 11, D-85748
Garching bei München, Germany*

Abstract

Two stochastic representations of multivariate geometric distributions are analyzed, both are obtained by lifting the lack-of-memory (LM) property of the univariate geometric law to the multivariate case. On the one hand, the narrow-sense multivariate geometric law can be considered a discrete equivalent of the well-studied Marshall-Olkin exponential law. On the other hand, the more general wide-sense geometric law is shown to be characterized by the LM property and can differ significantly from its continuous counterpart, e.g., by allowing for negative pairwise correlations.

For both families of distributions, their d -dimensional exchangeable subclass is characterized analytically via d -log-monotone, resp. d -monotone, sequences of parameters. Using this reparameterization, the subfamilies of distributions with conditionally i.i.d. components in the sense of de Finetti's theorem are determined. For these, a third stochastic construction based on a non-decreasing random walk is presented. The narrow-sense family is embedded in this construction when the increments of the involved random walk are infinitely divisible. The exchangeable narrow-sense law is furthermore shown to exhibit the multivariate right tail increasing (MRTI) dependence.

Keywords:

multivariate geometric law, lack-of-memory, exchangeability, completely monotone sequence, de Finetti's theorem, infinitely divisible law

2000 MSC: 62H05, 62H20

*Natalia Shenkman is solely responsible for the corrections and improvements marked in red. This includes a drastically simplified proof, the removal of two nonsensical proofs and the addition of some relevant references. The original manuscript was published in the *Journal of Multivariate Analysis* (2013).

Email addresses: jan-frederik.mai@xaia.com (Jan-Frederik Mai), scherer@tum.de (Matthias Scherer), shenkman@tum.de (Natalia Shenkman)

1. Introduction

The univariate geometric (resp. exponential) law is well known to be the only discrete (resp. continuous) distribution obeying the lack-of-memory (LM) property. Multivariate extensions have been studied in both cases. In the exponential case, an analytical derivation based on the multivariate continuous local LM property and probabilistic derivations based on exogenous exponential shock models are provided in [25]. All derivations are shown to lead to the same distribution. In the geometric case, multivariate extensions obtained as the waiting times for outcomes in a sequence of multinomial trials are introduced in [2]. In particular, [2] defines “a general multivariate geometric distribution” and presents a link to the Marshall-Olkin multivariate exponential distribution via multivariate geometric compounding of exponential random variables. Whereas [2] deals with probabilistic constructions of the multivariate geometric law, [26] and [29] are concerned with analytical derivations based on the LM property. [29] develop new classes of multivariate discrete survival functions adapted from monotonic behavior of a multivariate failure rate. In the process, they identify the families of multivariate geometric distributions with no-aging characteristic. Using the notation of this reference, survival functions belonging to both the IFR_3 and DFR_3 classes are precisely the survival functions satisfying the multivariate discrete local LM property. The results of [26] imply that, in contrast to the exponential case, the class of geometric distributions obtained using an exogenous fatal-shock model in the sense of [25], where the exponential shocks are replaced by geometric ones, does not comprise all solutions to the functional equation corresponding to the multivariate discrete local LM property. Therefore, they subdivide the set of multivariate geometric distributions into $\mathcal{G}^{\mathcal{N}}$ and $\mathcal{G}^{\mathcal{W}}$, abbreviating “narrow-sense” (\mathcal{N}) and “wide-sense” (\mathcal{W}), respectively. The fact that the geometric law requires a finer treatment compared with the exponential law is emphasized in [26]: “*the structure of families of multivariate geometric distributions is more complicated and more interesting than is the structure of families of multivariate exponential distributions.*”


In the recent literature, some attention has been drawn to the exchangeable and extendible subfamilies of multivariate distributions. In general, it is very difficult to verify analytically that a d -variate function F is a probability distribution (resp. survival probability) function. The major difficulty is to prove rectangular inequalities that guarantee positivity of the probability measure. The restriction to permutation invariant functions, corresponding to exchangeable probability distributions, can facilitate this task and lead to stunning mathematical links. For instance, if F is parameterized by a univariate function f , i.e. $F = F(f)$, the rectangular inequalities for F might translate into convenient analytic properties of f . These analytic properties might provide an illuminating loop back to probability theory and shed light on the probabilistic construction of F . In this spirit, [18, 24, 27, 28] characterize Archimedean copulas via monotonicity properties of their generator function. The latter is shown to be a characterizing transform of a univariate probability law which appears in the construction of Archimedean copulas. Similar results are known for spher-

ical and elliptical distributions, see [11], p. 26 ff., and [19, 35]. [22, 23, 30, 31] provide a link between Marshall-Olkin exponential distributions and completely monotone sequences. Relying on the results of [14], the references [22, 23] reveal astonishing interrelations with the theory of Lévy subordinators. A rigorous study of the exchangeable subsets of $\mathcal{G}^{\mathcal{N}}$ and $\mathcal{G}^{\mathcal{W}}$ discloses similar interesting interrelations with completely monotone sequences, random walks, and infinitely divisible laws. The present article contributes to the existing literature in the following way:

- **Stochastic models with limited memory:** It is shown that the family of general multivariate geometric distributions defined in [2] comprises precisely all solutions to the functional equation corresponding to the local discrete LM property and, thus, coincides with $\mathcal{G}^{\mathcal{W}}$ (see Theorem 2.6). Further, $\mathcal{G}^{\mathcal{N}}$ forms a proper subset of $\mathcal{G}^{\mathcal{W}}$. The survival functions for both families are computed in closed form (see Theorem 2.5).
- **Exchangeable subclasses:** The exchangeable subclasses are rewritten in terms of parameter sequences, truncated versions of which determine all lower-dimensional marginal distributions (see Remark 3.2). The set of d -dimensional exchangeable wide-sense distributions ($\mathcal{G}^{\mathcal{W},\mathcal{X}}$) is characterized by d -monotone sequences (\mathcal{M}_d), and the set of d -dimensional exchangeable narrow-sense distributions ($\mathcal{G}^{\mathcal{N},\mathcal{X}}$) is characterized by the (newly introduced) stronger notion of d -log-monotone sequences (\mathcal{LM}_d) (see Theorem 3.5).
- **Subclasses with conditionally i.i.d. components:** Starting from a d -variate exchangeable distribution, we investigate whether the distribution in question can be obtained by truncation of an infinite exchangeable sequence. This is solved analytically and corresponds to extendibility of the parameter sequence to a completely monotone sequence (\mathcal{M}_∞), respectively to a completely log-monotone sequence (\mathcal{LM}_∞) (see Corollary 4.1). For both the extendible wide-sense ($\mathcal{G}^{\mathcal{W},\mathcal{E}}$) and the extendible narrow-sense ($\mathcal{G}^{\mathcal{N},\mathcal{E}}$) families, a stochastic model with components that are i.i.d. conditioned on a non-decreasing random walk is constructed (see Theorem 4.6 (Esary and Marshall (1973))). In the universe of this model, the $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ -distribution arises as the proper subclass constructed by random walks with infinitely divisible increments. A natural link between \mathcal{LM}_∞ and infinitely divisible distributions is established (see Proposition 4.4 (Gnedin and Pitman (2008))). Interconnections with the results of [12, 13] on random integer partitions are highlighted.
- **Dependence properties:** Pairwise correlations are computed for both the wide- and narrow-sense geometric distribution (see Lemma 5.1). Interestingly, the wide-sense geometric distribution allows for negative correlations, whereas the narrow-sense geometric distribution can only model non-negative correlations. The exchangeable narrow-sense as well as the extendible wide-sense distributions are shown to exhibit the MRTI property (see Theorem 5.7 and Corollary 5.8).

- **Relations between subfamilies:** The following Venn diagram illustrates the relations that we identify between subfamilies of $\mathcal{G}^{\mathcal{W},\mathcal{X}}$. In

Figure 1: Characterization of multivariate exchangeable geometric distributions.



dimension 2, and only in dimension 2, the inner sets agree, i.e.

$$\mathcal{G}^{\mathcal{N},\mathcal{E}} = \mathcal{G}^{\mathcal{N},\mathcal{X}} = \mathcal{G}^{\mathcal{W},\mathcal{E}} = \mathcal{G}_{\text{MRTI}}^{\mathcal{W},\mathcal{X}} = \mathcal{G}^{\mathcal{W},\mathcal{X}}$$

with non-negative correlations, and form a proper subset of $\mathcal{G}^{\mathcal{W},\mathcal{X}}$.

2. Multivariate geometric distributions with limited memory

This section derives multivariate extensions of the univariate geometric distribution satisfying the local discrete LM property. To this end, we consider random variables τ taking values in $\mathbb{N} := \{1, 2, \dots\}$ and denote $\mathbb{N}_0 := \{0\} \cup \mathbb{N}$. The following definition is included for the sake of clarity.

Definition 2.1 (Discrete survival function). For $d \in \mathbb{N}$, the multi-indexed sequence $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$ is called a discrete survival function if there exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ supporting an \mathbb{N}^d -valued random vector (τ_1, \dots, τ_d) such that

$$\bar{F}_{n_1, \dots, n_d} = \mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d), \quad n_1, \dots, n_d \in \mathbb{N}_0.$$

A prominent example of a distribution on \mathbb{N} and the univariate starting point of the present investigation is the geometric law.

Example 2.2 (Geometric distribution). Let $p \in [0, 1)$. Considering a series of i.i.d. Bernoulli distributed random variables Z_1, Z_2, \dots with success probability $1 - p$, i.e. $\mathbb{P}(Z_1 = 1) = 1 - p = 1 - \mathbb{P}(Z_1 = 0)$, we denote by $\tau := \min\{n \in \mathbb{N} : Z_n = 1\}$ the first time of success. The distribution of τ is called geometric and denoted by $\text{Geo}(1 - p)$. Its corresponding survival function $\{\bar{F}_n\}_{n \in \mathbb{N}_0}$ is given by $\bar{F}_n := \mathbb{P}(\tau > n) = p^n$, $n \in \mathbb{N}_0$.

As already mentioned, the geometric distribution is the only distribution on \mathbb{N} satisfying the discrete LM property $\bar{F}_{n+m} = \bar{F}_n \bar{F}_m$, $n, m \in \mathbb{N}_0$. This renders the geometric distribution a discrete equivalent of the (continuous) exponential distribution. Moreover, it is easy to see that the geometric distribution is min-stable in the following sense: if $\tau_1 \sim \text{Geo}(1 - p_1)$ and $\tau_2 \sim \text{Geo}(1 - p_2)$ are independent, then $\min\{\tau_1, \tau_2\} \sim \text{Geo}(1 - p_1 p_2)$. This is another property shared with the exponential law. Considering limiting cases, $\tau \sim \text{Geo}(1)$ implies $\tau = 1$ almost surely. Further, a distribution with a point mass at infinity is conveniently interpreted as $\text{Geo}(0)$.

A first multivariate extension of the geometric law, preserving the local discrete LM property, is obtained using a discrete analogue of the Marshall–Olkin exponential shock model, see [25], where the exponentially distributed shocks are replaced by geometric ones. The resulting distribution is referred to as a multivariate geometric distribution in the narrow sense (\mathcal{G}^N). In the continuous case, the Marshall–Olkin exponential shock model generates precisely the class of all multivariate exponential distributions satisfying the local continuous LM property. However, the family of multivariate geometric distributions satisfying the local LM property has a more complicated and richer structure. In particular, it contains distributions which can not be obtained using a discrete analogue of the Marshall–Olkin model.

As a second multivariate extension of the geometric law, we study a multivariate geometric distribution introduced in Section 4 of [2]. We prove that this class of distributions comprises all solutions to the functional equation corresponding to the local discrete multivariate LM property. Consequently, it forms a proper superset of the parametric family of narrow-sense geometric distributions. Adopting the terminology of [26], geometric distributions obtained using the Arnold model will be called geometric in the wide sense (\mathcal{G}^W).

Definition 2.3 (Narrow- and wide-sense multivariate geometric law).
 (N) Let $p_I \in [0, 1]$, $\emptyset \neq I \subseteq \{1, \dots, d\}$, with $\prod_{I:k \in I} p_I < 1$ for $k = 1, \dots, d$, be given parameters. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space supporting a collection $\{E_I\}_{\emptyset \neq I \subseteq \{1, \dots, d\}}$ of independent geometric random variables, where $E_I \sim \text{Geo}(1 - p_I)$. Define the random vector (τ_1, \dots, τ_d) by

$$\tau_k := \min\{E_I : k \in I\}, \quad k = 1, \dots, d.$$

The distribution of (τ_1, \dots, τ_d) is called multivariate narrow-sense geometric with parameters $\mathbf{p} := \{p_I\}_{\emptyset \neq I \subseteq \{1, \dots, d\}}$ and is denoted by $\mathcal{G}^N(\mathbf{p})$.

(W) Let $\tilde{p}_I \in [0, 1]$, $I \subseteq \{1, \dots, d\}$, with $\sum_I \tilde{p}_I = 1$ and $\sum_{I:k \notin I} \tilde{p}_I < 1$ for $k = 1, \dots, d$, be given parameters. Consider an experiment with 2^d possible outcomes $I \subseteq \{1, \dots, d\}$, each of which occurs with probability \tilde{p}_I . The experiment is run repeatedly in independent and identically distributed trials. Further, let $\{\tilde{E}_I\}_{I \subseteq \{1, \dots, d\}}$ be a collection of random variables, where \tilde{E}_I represents the number of the experiment which first yielded the outcome I . Define the random vector (τ_1, \dots, τ_d) by

$$\tau_k := \min\{\tilde{E}_I : k \in I\}, \quad k = 1, \dots, d.$$

The distribution of (τ_1, \dots, τ_d) is called multivariate wide-sense geometric with parameters $\tilde{\mathbf{p}} := \{\tilde{p}_I\}_{I \subseteq \{1, \dots, d\}}$ and is denoted by $\mathcal{G}^{\mathcal{W}}(\tilde{\mathbf{p}})$.

Remark 2.4 (Comparing the stochastic models).

(\mathcal{N}) In the case of a narrow-sense geometric distribution, each component τ_k , $k = 1, \dots, d$, is defined as a minimum over 2^{d-1} independent geometric random variables indexed by the subsets $\emptyset \neq I \subseteq \{1, \dots, d\}$ with $k \in I$. By min-stability of the geometric law, τ_k is $\text{Geo}(1 - \prod_{I: k \in I} p_I)$ -distributed. The condition $\prod_{I: k \in I} p_I < 1$ ensures that τ_k is almost surely finite. The case where at least one τ_k is almost surely 1 can be excluded by requiring $p_I \neq 0$ for all I . The dependence between the components results from the overlapping subset indices in the definition of the τ_k 's.

(\mathcal{W}) In the case of a wide-sense geometric distribution, the shocks are geometric $\tilde{E}_I \sim \text{Geo}(\tilde{p}_I)$, $I \subseteq \{1, \dots, d\}$, and $\tau_k \sim \text{Geo}(1 - \sum_{I: k \notin I} \tilde{p}_I)$. Again, each τ_k is almost surely finite due to $\sum_{I: k \notin I} \tilde{p}_I < 1$ for $k = 1, \dots, d$, and $\tau_k = 1$ almost surely if $\sum_{I: k \notin I} \tilde{p}_I = 0$. The crucial difference between the two probabilistic models is that the geometric shocks E_I in the definition of the narrow-sense geometric distribution are independent, whereas the construction of a wide-sense multivariate geometric distribution is based on dependent geometric random variables \tilde{E}_I . Due to $\sum_I \tilde{p}_I = 1$ in the case of a wide-sense geometric distribution, both distributions $\mathcal{G}^{\mathcal{N}}$ and $\mathcal{G}^{\mathcal{W}}$ are parameterized by $2^d - 1$ parameters.

Theorems 2.5 and 2.6 characterize both distributions in terms of the survival function and the local LM property. In the following, $n_{(1)} \leq n_{(2)} \leq \dots \leq n_{(d)}$ denotes the ordered list of $n_1, \dots, n_d \in \mathbb{N}_0$.

Theorem 2.5 (Survival function and limited memory).

(\mathcal{N}) Let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{N}}(\mathbf{p})$. Then, (τ_1, \dots, τ_d) has support \mathbb{N}^d with discrete survival function $\{\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$, given by

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} := \mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) = \prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{i \in I} \{n_i\}}. \quad (1)$$

(\mathcal{W}) Let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{W}}(\tilde{\mathbf{p}})$. Then, (τ_1, \dots, τ_d) has support \mathbb{N}^d with discrete survival function $\{\bar{F}_{n_1, \dots, n_d}^{\mathcal{W}}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$, given by

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{W}} := \mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) = \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_n(i) \notin I \ \forall i=k, \dots, d}} \tilde{p}_I \right)^{n_{(k)} - n_{(k-1)}}, \quad (2)$$

with the convention $n_{(0)} := 0$, and $\pi_n: \{1, \dots, d\} \rightarrow \{1, \dots, d\}$ being a permutation depending on $\mathbf{n} = (n_1, \dots, n_d)$ such that $n_{\pi_n(1)} \leq n_{\pi_n(2)} \leq \dots \leq n_{\pi_n(d)}$.

Moreover, both distributions (1) and (2) satisfy the local discrete multivariate LM property

$$\mathbb{P}(\tau_{i_1} > n_{i_1} + m, \dots, \tau_{i_k} > n_{i_k} + m \mid \tau_{i_1} > m, \dots, \tau_{i_k} > m) = \mathbb{P}(\tau_{i_1} > n_{i_1}, \dots, \tau_{i_k} > n_{i_k}),$$

for $k = 1, \dots, d$, $1 \leq i_1, \dots, i_k \leq d$, $m, n_{i_1}, \dots, n_{i_k} \in \mathbb{N}_0$. (3)

PROOF. Refer to [Appendix A](#). \square

Theorem 2.6 (Uniqueness). *A distribution on \mathbb{N}^d satisfies the local discrete LM property (3) if and only if it is a d -variate wide-sense geometric distribution.*

PROOF. Refer to [Appendix A](#). \square

The inclusion $\mathcal{G}^{\mathcal{N}} \subsetneq \mathcal{G}^{\mathcal{W}}$ is immediate from Theorems 2.5 and 2.6, and proper in any dimension $d \geq 2$, see Example 3.8. Regarding a probabilistic meaning of the parameters $\tilde{\mathbf{p}} = \{\tilde{p}_I\}_{I \subseteq \{1, \dots, d\}}$ of $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{W}}(\tilde{\mathbf{p}})$, we have

$$\tilde{p}_I = \mathbb{P}(\tilde{E}_I = 1) = \mathbb{P}(\{\tau_i > 1 \forall i \notin I\} \cap \{\tau_i = 1 \forall i \in I\}), \quad I \subseteq \{1, \dots, d\}.$$

Example 2.7 (Three-variate geometric law).

(\mathcal{N}) Let $(\tau_1, \tau_2, \tau_3) \sim \mathcal{G}^{\mathcal{N}}(\{p_I\}_{\emptyset \neq I \subseteq \{1, 2, 3\}})$. Then, its corresponding survival function $\{\bar{F}_{n_1, n_2, n_3}^{\mathcal{N}}\}_{n_1, n_2, n_3 \in \mathbb{N}_0}$ is given by

$$\bar{F}_{n_1, n_2, n_3}^{\mathcal{N}} = p_{\{1\}}^{n_1} p_{\{2\}}^{n_2} p_{\{3\}}^{n_3} p_{\{1, 2\}}^{\max\{n_1, n_2\}} p_{\{1, 3\}}^{\max\{n_1, n_3\}} p_{\{2, 3\}}^{\max\{n_2, n_3\}} p_{\{1, 2, 3\}}^{n_{(3)}}. \quad (4)$$

(\mathcal{W}) Let $(\tau_1, \tau_2, \tau_3) \sim \mathcal{G}^{\mathcal{W}}(\{\tilde{p}_I\}_{I \subseteq \{1, 2, 3\}})$. Then, its corresponding survival function $\{\bar{F}_{n_1, n_2, n_3}^{\mathcal{W}}\}_{n_1, n_2, n_3 \in \mathbb{N}_0}$ is given by

$$\bar{F}_{n_1, n_2, n_3}^{\mathcal{W}} = \begin{cases} \tilde{p}_{\emptyset}^{n_1} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}})^{n_2 - n_1} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}} + \tilde{p}_{\{2\}} + \tilde{p}_{\{1, 2\}})^{n_3 - n_2}, & n_1 \leq n_2 \leq n_3, \\ \tilde{p}_{\emptyset}^{n_1} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}})^{n_3 - n_1} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}} + \tilde{p}_{\{3\}} + \tilde{p}_{\{1, 3\}})^{n_2 - n_3}, & n_1 \leq n_3 < n_2, \\ \tilde{p}_{\emptyset}^{n_2} (\tilde{p}_{\emptyset} + \tilde{p}_{\{2\}})^{n_1 - n_2} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}} + \tilde{p}_{\{2\}} + \tilde{p}_{\{1, 2\}})^{n_3 - n_1}, & n_2 < n_1 \leq n_3, \\ \tilde{p}_{\emptyset}^{n_2} (\tilde{p}_{\emptyset} + \tilde{p}_{\{2\}})^{n_3 - n_2} (\tilde{p}_{\emptyset} + \tilde{p}_{\{2\}} + \tilde{p}_{\{3\}} + \tilde{p}_{\{2, 3\}})^{n_1 - n_3}, & n_2 \leq n_3 < n_1, \\ \tilde{p}_{\emptyset}^{n_3} (\tilde{p}_{\emptyset} + \tilde{p}_{\{3\}})^{n_1 - n_3} (\tilde{p}_{\emptyset} + \tilde{p}_{\{1\}} + \tilde{p}_{\{3\}} + \tilde{p}_{\{1, 3\}})^{n_2 - n_1}, & n_3 < n_1 \leq n_2, \\ \tilde{p}_{\emptyset}^{n_3} (\tilde{p}_{\emptyset} + \tilde{p}_{\{3\}})^{n_2 - n_3} (\tilde{p}_{\emptyset} + \tilde{p}_{\{2\}} + \tilde{p}_{\{3\}} + \tilde{p}_{\{2, 3\}})^{n_1 - n_2}, & n_3 < n_2 < n_1. \end{cases} \quad (5)$$

An elementary example of a three-variate narrow-sense geometric distribution is (τ_1, τ_2, τ_3) with mutually independent $\tau_i \sim \text{Geo}(1 - p_i)$, $p_i \in [0, 1)$, $i = 1, 2, 3$, i.e.

$$\bar{F}_{n_1, n_2, n_3} = p_1^{n_1} p_2^{n_2} p_3^{n_3}. \quad (6)$$

The survival function in (6) can be represented as in (4) with parameters of a narrow-sense distribution $p_{\{i\}} = p_i$, $i = 1, 2, 3$, and $p_I = 0$ for $|I| > 1$. (6) can also be rewritten to resemble the survival function of a wide-sense distribution (5) by setting $\tilde{p}_I = \sum_{i=0}^{|I|} (-1)^{|I|+i} \sum_{\substack{J \subseteq I \\ |J|=i}} \prod_{\substack{j \in \{1, 2, 3\} \\ j \notin J}} p_j$, $I \subseteq \{1, 2, 3\}$.

For a discussion of the bivariate wide-sense geometric distribution, we refer the reader to [3].

Studying the exchangeable subfamilies of $\mathcal{G}^{\mathcal{N}}$ and $\mathcal{G}^{\mathcal{W}}$ reveals astonishing interrelations with completely monotone sequences and infinitely divisible laws. In particular, a convenient analytical characterization can be achieved for exchangeable geometric survival functions. Furthermore, exchangeable distributions might be of interest for practical reasons: in the non-exchangeable case, the number of parameters, namely $2^d - 1$ in dimension d , grows exponentially in d . The restriction to permutation invariant distributions allows to significantly reduce the number of parameters to d in dimension d .

3. Exchangeable subfamilies

This section provides an analytical characterization of exchangeable narrow- and wide-sense geometric distributions, i.e. distributions invariant under all permutations of their components. An exchangeability criterion for the distributions in (1) and (2) is given in Lemma 3.1.

Lemma 3.1 (Exchangeability criterion). *The d -dimensional survival functions (1) and (2) are exchangeable if and only if their parameters satisfy the following condition:*

$$|I_1| = |I_2| \Rightarrow p_{I_1} = p_{I_2}, \text{ resp. } |I_1| = |I_2| \Rightarrow \tilde{p}_{I_1} = \tilde{p}_{I_2}, \quad (7)$$

where $|I|$ denotes the cardinality of a set $I \subseteq \{1, \dots, d\}$.

PROOF. Refer to Appendix B. While the lemma is intuitive, the proofs provided in the appendix convey the impression of a certain complexity. They become trivial if the correct parametrization of the survival functions is invoked; see [36] and the proof of Lemma 2.1 in [37]. \square

Lemma 3.1 implies that in the exchangeable case, both distributions (1) and (2) are parameterized by only d parameters. Remark 3.2 carries out a reparametrization of exchangeable survival functions in terms of parameter sequences, truncated versions of which determine all lower-dimensional marginal distributions. Besides notational convenience, this reparametrization facilitates an analytical characterization of the corresponding survival function.

Remark 3.2 (Reparameterization).

(\mathcal{N}) For the exchangeable narrow-sense geometric distribution, the distinct parameters are p_1, \dots, p_d , where $p_k := p_{\{1, \dots, k\}}$ for $k = 1, \dots, d$. These parameters satisfy $p_1, \dots, p_d \in (0, 1]$ and $\prod_{i=1}^d p_i < 1$. A combinatorial exercise shows that in this case, the survival function (1) simplifies to

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d \left(\prod_{i=1}^{d-k+1} p_i^{\binom{d-k}{i-1}} \right)^{n_{(d-k+1)}}.$$

Introducing the notation

$$a_k := \prod_{i=1}^{d-k+1} p_i^{\binom{d-k}{i-1}}, \quad k = 1, \dots, d, \quad (8)$$

we obtain a convenient expression for the d -dimensional exchangeable narrow-sense geometric survival function:

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d a_k^{n_{(d-k+1)}}. \quad (9)$$

For a comparison of the narrow- with the wide-sense geometric distributions, it is more convenient to parameterize the survival function (9) in terms of the parameters $\{b_k\}_{k=1}^d$, given by

$$b_k := \prod_{i=1}^k a_i, \quad k = 1, \dots, d. \quad (10)$$

Denoting $b_0 := 1$ and $n_{(0)} := 0$, this gives $a_k = b_k/b_{k-1}$ for $k = 1, \dots, d$, and the survival function in (9) takes the form

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d (b_k/b_{k-1})^{n_{(d-k+1)}} = \prod_{k=1}^d b_k^{n_{(d-k+1)} - n_{(d-k)}}. \quad (11)$$

(\mathcal{W}) Analogously, the exchangeable wide-sense geometric distribution can be parameterized by d parameters $\tilde{p}_1, \dots, \tilde{p}_d$, where $\tilde{p}_1 := \tilde{p}_\emptyset$ and $\tilde{p}_k := \tilde{p}_{\{1, \dots, k-1\}}$ for $k = 2, \dots, d$. The parameter $\tilde{p}_{\{1, \dots, d\}}$ is determined by $\tilde{p}_1, \dots, \tilde{p}_d$ via the condition $\sum_I \tilde{p}_I = 1$. The parameters $\tilde{p}_1, \dots, \tilde{p}_d$ satisfy $\tilde{p}_1, \dots, \tilde{p}_d \in [0, 1]$, $\sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \leq 1$, and $\sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i < 1$. Further, the exchangeable wide-sense geometric survival function (2) can be conveniently written as

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{W}} = \prod_{k=1}^d \beta_k^{n_{(d-k+1)} - n_{(d-k)}}, \quad (12)$$

where the new parameters $\{\beta_k\}_{k=1}^d$ are given by

$$\beta_k := \sum_{i=1}^{d-k+1} \binom{d-k}{i-1} \tilde{p}_i, \quad k = 1, \dots, d. \quad (13)$$

There exists a fundamental connection between the parameter spaces of exchangeable multivariate narrow- and wide-sense geometric distributions and d -monotone sequences. On the one hand, the sequence of parameters $(1, \beta_1, \dots, \beta_d)$, given as in (13), of every exchangeable d -variate wide-sense geometric distribution is $(d+1)$ -monotone. On the other hand, each $(d+1)$ -monotone sequence

defines admissible parameters of an exchangeable d -variate geometric distribution satisfying the local LM property. The related concept of complete monotonicity has been studied in the literature, see e.g. [14, 21]. In the present article, we particularize d -monotonicity by introducing the stronger notion of d -log-monotonicity. The meaning of d -log-monotonicity is twofold. First, it characterizes the parameters $\{b_k\}_{k=1}^d$, given as in (10), of an exchangeable d -variate narrow-sense geometric distribution. Second, as we shall see in Section 4, the complete log-monotonicity, i.e. d -log-monotonicity for all $d \in \mathbb{N}_0$, uniquely characterizes the sequence of non-positive exponential moments of an *infinitely divisible* probability law on $[0, \infty]$. This result constitutes an analogue to Hausdorff's moment problem, see [15, 16], i.e. $\{\beta_k\}_{k=0}^\infty$ is completely monotone if and only if it is the sequence of moments of a probability measure on $[0, 1]$. Moreover, it links the infinitely extendible subclass of exchangeable narrow-sense geometric distributions to infinitely divisible distributions on $[0, \infty]$.

Definition 3.3 (d -monotone and d -log-monotone sequences). *A finite sequence $(x_0, \dots, x_{d-1}) \in \mathbb{R}^d$ is said to be d -monotone if it satisfies*

$$\nabla^j x_k := \sum_{i=0}^j (-1)^i \binom{j}{i} x_{k+i} \geq 0, \quad k = 0, 1, \dots, d-1, \quad j = 0, 1, \dots, d-k-1. \quad (14)$$

An infinite sequence $\{x_k\}_{k \in \mathbb{N}_0}$ is called completely monotone if $\nabla^j x_k \geq 0$ for all $k \in \mathbb{N}_0, j \in \mathbb{N}_0$. We define the set \mathcal{M}_d of d -monotone and \mathcal{M}_∞ of completely monotone sequences, starting with 1, by

$$\begin{aligned} \mathcal{M}_d &:= \{ \{x_k\}_{k=0}^{d-1} \in \mathbb{R}^d \mid x_0 = 1, x_1 < 1, \{x_k\}_{k=0}^{d-1} \text{ is } d\text{-monotone} \}, \\ \mathcal{M}_\infty &:= \{ \{x_k\}_{k \in \mathbb{N}_0} \in \mathbb{R}^{\mathbb{N}_0} \mid x_0 = 1, x_1 < 1, \{x_k\}_{k \in \mathbb{N}_0} \text{ is completely monotone} \}, \end{aligned}$$

respectively. Further, we define the set \mathcal{LM}_d of d -log-monotone and \mathcal{LM}_∞ of completely log-monotone sequences, starting with 1, by

$$\begin{aligned} \mathcal{LM}_d &:= \{ \{x_k\}_{k=0}^{d-1} \in \mathbb{R}_{>0}^d \mid x_0 = 1, x_1 < 1, \nabla^{d-k-1} \ln x_k \geq 0, k=0, 1, \dots, d-2 \}, \\ \mathcal{LM}_\infty &:= \{ \{x_k\}_{k \in \mathbb{N}_0} \in \mathbb{R}_{>0}^{\mathbb{N}_0} \mid x_0 = 1, x_1 < 1, \nabla^j \ln x_k \geq 0, k \in \mathbb{N}_0, j \in \mathbb{N} \}, \end{aligned}$$

respectively. Lastly, we define the set \mathcal{SM}_d of d -strong-monotone and \mathcal{SM}_∞ of completely strong-monotone sequences by

$$\begin{aligned} \mathcal{SM}_d &:= \{ \{x_k\}_{k=0}^{d-1} \in \mathbb{R}_{>0}^d \mid x_0 < 1, \{\ln x_k^{-1}\}_{k=0}^{d-1} \text{ is } d\text{-monotone} \}, \\ \mathcal{SM}_\infty &:= \{ \{x_k\}_{k \in \mathbb{N}_0} \in \mathbb{R}_{>0}^{\mathbb{N}_0} \mid x_0 < 1, \{\ln x_k^{-1}\}_{k \in \mathbb{N}_0} \text{ is completely monotone} \}, \end{aligned}$$

respectively.

For $1 \leq j \leq d-1-k$, we have $\nabla^j x_k = \nabla^{j-1} x_k - \nabla^{j-1} x_{k+1}$. Further, condition (14) is equivalent to $\nabla^{d-1-k} x_k \geq 0$ for $k = 0, 1, \dots, d-1$, see [34] or [22].

By definition, each sequence $\{x_k\}_{k=0}^{d-1} \in \mathcal{M}_d$ (resp. $\{x_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$) is $[0, 1]$ -valued and decreasing, whereas each sequence $\{x_k\}_{k=0}^{d-1} \in \mathcal{SM}_d$ (resp. $\{x_k\}_{k \in \mathbb{N}_0} \in \mathcal{SM}_\infty$) is $(0, 1]$ -valued and increasing. Further, in the definition of the set \mathcal{LM}_d , $\ln x_{d-1} \geq 0$ is the only missing condition for $\{\ln x_k\}_{k=0}^{d-1}$ to be d -monotone. Moreover, as we show later in this section, there exists a close relationship between the classes \mathcal{M}_d , \mathcal{LM}_d , and \mathcal{SM}_d , such as, e.g., $\mathcal{LM}_d \subsetneq \mathcal{M}_d$ and $\mathcal{LM}_\infty \subsetneq \mathcal{M}_\infty$, see Figure 1.

Trivially, each completely monotone sequence $\{x_k\}_{k \in \mathbb{N}_0}$ is d -monotone for all $d \in \mathbb{N}$, when truncated at x_{d-1} . Analogously, each completely log-monotone sequence $\{x_k\}_{k \in \mathbb{N}_0}$ is d -log-monotone for all $d \in \mathbb{N}$, when truncated at x_{d-1} . However, not every d -(log-) monotone sequence can be extended to a completely (log-)monotone sequence. Consider the following example.

Example 3.4 (Proper 3-monotone sequence). *Consider the sequence $(x_0, x_1, x_2) := (1, 1/2, 1/5)$. It is easy to check that $(x_0, x_1, x_2) \in \mathcal{M}_3$. Assume that (x_0, x_1, x_2) can be extended to an infinite completely monotone sequence $\{x_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$. Then, by Hausdorff's moment problem, $x_k = \mathbb{E}[\tau^k]$ for $k = 0, 1, 2$ and a $[0, 1]$ -valued random variable τ . Jensen's inequality gives a contradiction:*

$$\frac{1}{5} = x_2 = \mathbb{E}[\tau^2] \geq \mathbb{E}[\tau]^2 = x_1^2 = \frac{1}{4}.$$

Indeed, it follows from the Hankel determinants criterion, see [17] and [9], p. 20, that only $(1, 1/2, x_2) \in \mathcal{M}_3$ with $x_2 \geq 1/4$ can be extended to an infinite completely monotone sequence. In contrast, each $(1, x_1) \in \mathcal{M}_2$ is extendible to a completely monotone sequence.

The next theorem constitutes an analytical characterization of d -variate exchangeable narrow- and wide-sense geometric distributions in terms of d -monotone and d -log-monotone sequences, respectively. This facilitates a rigorous study of $\mathcal{GN}^{\mathcal{X}}$ and $\mathcal{GW}^{\mathcal{X}}$. In particular, it helps to determine the subclasses with conditionally i.i.d. components in the sense of de Finetti's theorem, see Section 4.

Theorem 3.5 (Characterization of the exchangeable geometric law).

(\mathcal{N}) *Consider the parametric family $\mathcal{GN}^{\mathcal{X}}$ of all exchangeable, non-degenerate d -dimensional narrow-sense geometric survival functions. Then,*

$$\begin{aligned} \mathcal{GN}^{\mathcal{X}} &= \left\{ \bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d a_k^{n(d-k+1)} \mid \{a_k\}_{k=1}^d \in \mathcal{SM}_d \right\} \\ &= \left\{ \bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d b_k^{n(d-k+1) - n(d-k)} \mid (1, b_1, \dots, b_d) \in \mathcal{LM}_{d+1} \right\}. \end{aligned}$$

The parameters $p_1, \dots, p_d \in (0, 1]$ with $\prod_{i=1}^d p_i < 1$ and $(a_1, \dots, a_d) \in \mathcal{SM}_d$ are related via

$$a_k = \prod_{i=1}^{d-k+1} p_i^{\binom{d-k}{i-1}}, \quad k = 1, \dots, d,$$

$$p_k = \prod_{i=1}^k a_{d-i+1}^{(-1)^{(k-i)} \binom{k-1}{i-1}}, \quad k = 1, \dots, d.$$

The parameters $\{b_k\}_{k=1}^d$ with $(1, b_1, \dots, b_d) \in \mathcal{LM}_{d+1}$ are given as in (10).

(W) Consider the parametric family $\mathcal{G}^{\mathcal{W}, \mathcal{X}}$ of all exchangeable d -dimensional wide-sense geometric survival functions. Then,

$$\mathcal{G}^{\mathcal{W}, \mathcal{X}} = \{\bar{F}_{n_1, \dots, n_d}^{\mathcal{W}} = \prod_{k=1}^d \beta_k^{n_{(d-k+1)} - n_{(d-k)}} \mid (1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}\}.$$

The parameters $\tilde{p}_1, \dots, \tilde{p}_d \in [0, 1]$ with $\sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \leq 1$, $\sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i < 1$, and $\{\beta_k\}_{k=1}^d$ with $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ are related via

$$\beta_k = \sum_{i=1}^{d-k+1} \binom{d-k}{i-1} \tilde{p}_i, \quad k = 1, \dots, d,$$

$$\tilde{p}_k = \nabla^{k-1} \beta_{d-k+1}, \quad k = 1, \dots, d.$$

PROOF. Refer to [Appendix B](#). □

Example 3.6 (Three-variate exchangeable geometric law).

Consider the survival function (6) of a three-variate distribution with mutually independent geometric marginals from [Example 2.7](#). In the exchangeable case, i.e. $p_1 = p_2 = p_3 =: p$, (6) simplifies to

$$\bar{F}_{n_1, n_2, n_3} = p^{n_1 + n_2 + n_3} = \prod_{k=1}^3 b_k^{n_{(4-k)} - n_{(3-k)}}, \quad (15)$$

with $b_i = p^i$, $i = 1, 2, 3$. It can be easily verified that $(1, p, p^2, p^3) \in \mathcal{LM}_4$.

Another example of an exchangeable three-variate narrow sense geometric distribution is (τ_1, τ_2, τ_3) with identically $\text{Geo}(1-p)$ -distributed marginals, $p \in [0, 1)$, and the dependence structure induced by the upper Fréchet-Hoeffding bound. The survival function of (τ_1, τ_2, τ_3) is then given by

$$\bar{F}_{n_1, n_2, n_3} = p^{n_{(3)}} = \prod_{k=1}^3 b_k^{n_{(4-k)} - n_{(3-k)}}, \quad (16)$$

with $b_i = p$, $i = 1, 2, 3$. Trivially, $(1, p, p, p) \in \mathcal{LM}_4$. Note that the lower Fréchet-Hoeffding bound with homogeneous geometric marginals does not lead to a wide-sense distribution in dimension $d = 2$. In the non-exchangeable case, neither the upper nor the lower Fréchet-Hoeffding bound with geometric marginals satisfy the local LM property.

Theorem 3.5 (\mathcal{N}) provides an analytical characterization of the exchangeable, non-degenerate narrow-sense geometric distribution. The only distribution not covered by Theorem 3.5 (\mathcal{N}) is the degenerate distribution with point mass at $\{1\}^d$. It can be obtained by setting $p_k = 0$ for at least one $k = 1, \dots, d$, or, equivalently, $a_1 = 0$ and $b_k = 0$ for all $k = 1, \dots, d$. The corresponding survival function is $\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \mathbf{1}_{\{n_1 = \dots = n_d = 0\}}$. In the context of a wide-sense geometric distribution, the degenerate case results from $p_k = 1$ for all $k = 1, \dots, d$, or, equivalently, $\beta_k = 0$ for all $k = 1, \dots, d$. Note also that each component of an exchangeable narrow-sense geometric distribution is $Geo(1 - a_1)$ -distributed with

$$a_1 = b_1 = \prod_{i=1}^d p_i^{\binom{d-1}{i-1}},$$

whereas each component of an exchangeable wide-sense geometric distribution is $Geo(1 - \beta_1)$ -distributed with $\beta_1 = \sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i$.

In Theorem 3.5 (\mathcal{W}), the original parameters \tilde{p}_k , $k = 1, \dots, d$, of an exchangeable wide-sense geometric distribution are explicitly given in terms of the $(k - 1)$ -fold difference operator of β_{d-k+1} . Analogous relationships can be established in the case of an exchangeable narrow-sense geometric distribution as well. Namely, for $k = 1, \dots, d$, it holds that

$$p_k = \exp(-\nabla^{k-1} \ln a_{d-k+1}^{-1}) = \exp(-\nabla^k \ln b_{d-k}).$$

Theorem 3.7 shows that the condition $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ in Theorem 3.5 (\mathcal{W}) is even *necessary and sufficient* for the multi-indexed sequence $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$ in (12) to define a discrete survival function.

Theorem 3.7 (Analytical characterization of the survival function).

The multi-indexed sequence $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$, given by

$$\bar{F}_{n_1, \dots, n_d} = \prod_{k=1}^d \beta_k^{n_{(d-k+1)} - n_{(d-k)}}, \quad n_1, \dots, n_d \in \mathbb{N}_0,$$

is a d -dimensional survival function if and only if $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$. In this case, $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$ defines an exchangeable wide-sense geometric distribution.

PROOF. Refer to Appendix B. The proof in the published manuscript is overly complicated since the key property of the distribution is not exploited. We have replaced it by a very simple argumentation based on Theorems 2.6 and 3.5. \square

The following example illustrates that $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distributions form a proper subclass of $\mathcal{G}^{\mathcal{W}, \mathcal{X}}$ -distributions.

Example 3.8 (Geometric in the wide-, but not in the narrow sense).

Consider again the sequence $(1, 1/2, 1/5) \in \mathcal{M}_3$ from Example 3.4. From Theorem 3.7, it follows that

$$\bar{F}_{n_1, n_2} = a_1^{\max(n_1, n_2)} a_2^{\min(n_1, n_2)}, \quad n_1, n_2 \in \mathbb{N}_0,$$

with $(a_1, a_2) = (1/2, 2/5)$, defines an exchangeable bivariate wide-sense geometric distribution. However, by Theorem 3.5, this distribution is not a $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distribution, since

$$\nabla^1 \ln a_1^{-1} = \ln \frac{4}{5} < 0,$$

and, consequently, $(a_1, a_2) \notin \mathcal{SM}_2$. Recall that the condition $(a_1, a_2) \in \mathcal{SM}_2$ ensures that the original parameters p_1, p_2 of the $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distribution are $(0, 1]$ -valued. The above choice of (a_1, a_2) leads to $p_1 = a_2 = 2/5$ and $p_2 = a_1/a_2 = 5/4 > 1$.

Next, we discuss conditions under which a d -variate exchangeable distribution can be obtained by truncation of an infinite exchangeable random sequence. By virtue of de Finetti's seminal theorem, this is equivalent to determining the subclasses with conditionally independent and identically distributed components.

4. Extendible subfamilies

In this section, we provide necessary and sufficient conditions on parameters of the survival function in (9) (resp. in (12)) to define a survival function of a narrow- (resp. wide-sense) geometric distribution in any dimension $d \in \mathbb{N}$. Consider the exchangeable random vector (τ_1, \dots, τ_d) , geometrically distributed (in the narrow or in the wide sense) with original parameters $p_{1,d}, \dots, p_{d,d}$. Clearly, e.g. by Theorem 3.5, every subvector has a geometric law as well. Denote the parameters of a subvector of (τ_1, \dots, τ_d) of length j , $j \in \{1, \dots, d\}$, by $p_{k,j}$, $k = 1, \dots, j$. For the sake of notational convenience, throughout Section 4, we omit the tilde symbol on the parameters $p_{k,j}$ of a wide-sense distribution. We depict the relation between the $p_{k,j}$, $j \in \{1, \dots, d\}$, $k = 1, \dots, j$, in the following triangular scheme:

$$\begin{array}{ccccccc}
 & & & & & & p_{1,1} \\
 & & & & & & p_{1,2} & p_{2,2} \\
 & & & & & & p_{1,3} & p_{2,3} & p_{3,3} \\
 & & & & & & \ddots & & \ddots \\
 & & & & & & p_{1,d-1} & \dots & p_{d-1,d-1} \\
 & & & & & & p_{1,d} & p_{2,d} & \dots & p_{d-1,d} & p_{d,d}
 \end{array}$$

By Remark 3.2, the parameters of two adjacent rows are related multiplicatively, via $p_{k,j} = p_{k,j+1} p_{k+1,j+1}$, for all $k = 1, \dots, j$, $j = 1, \dots, d-1$, in the case of a $\mathcal{G}^{\mathcal{N}, \mathcal{X}}(\mathbf{p})$ -distribution, and additively, via $p_{k,j} = p_{k,j+1} + p_{k+1,j+1}$, in

the case of a $\mathcal{G}^{\mathcal{W},\mathcal{X}}(\mathbf{p})$ -distribution. Recall that in the context of a $\mathcal{G}^{\mathcal{N},\mathcal{X}}(\mathbf{p})$ -distribution, the parameter sequence (a_1, \dots, a_j) , given by

$$a_k = p_{1,k} = \prod_{i=1}^{d-k+1} p_{i,d}^{\binom{d-k}{i-1}}, \quad k = 1, \dots, j,$$

determines all k -dimensional marginal distributions for $k = 1, \dots, j$, and each τ_k is distributed according to $Geo(1 - a_1)$. For a $\mathcal{G}^{\mathcal{W},\mathcal{X}}(\mathbf{p})$ -distribution, all k -dimensional marginals are determined by the parameter sequence $(\beta_1, \dots, \beta_j)$, where

$$\beta_k = p_{1,k} = \sum_{i=1}^{d-k+1} \binom{d-k}{i-1} p_{i,d}, \quad k = 1, \dots, j,$$

and each τ_k is distributed according to $Geo(1 - \beta_1)$.

Extendibility of (τ_1, \dots, τ_d) by one dimension to an exchangeable geometrically distributed $(\tau_1, \dots, \tau_d, \tau_{d+1})$ is equivalent to extendibility of a given triangle by an additional row $p_{1,d+1}, \dots, p_{d,d+1}, p_{d+1,d+1}$ without violating the multiplicative, resp. additive, structure. For a $\mathcal{G}^{\mathcal{N},\mathcal{X}}(\mathbf{p})$ -distribution, it is the case precisely when there exists a $q \in (0, 1]$ s.t.

$$q^{-1} \prod_{i=1}^{2k-1} p_{i,d}^{(-1)^{i+1}} \in (0, 1], \quad q \prod_{i=1}^{2k} p_{i,d}^{(-1)^i} \in (0, 1],$$

for all $k = 1, \dots, d/2$, if d is even, or $k = 1, \dots, ((d-1)/2 + 1)$, if d is odd. Similarly, for a $\mathcal{G}^{\mathcal{W},\mathcal{X}}(\mathbf{p})$ -distribution, extendibility by one dimension is equivalent to the existence of a $q \in [0, 1]$ s.t.

$$\sum_{i=1}^{2k-1} (-1)^{i+1} p_{i,d} - q \in [0, 1], \quad \sum_{i=1}^{2k} (-1)^i p_{i,d} + q \in [0, 1],$$

for all $k = 1, \dots, d/2$, if d is even, or $k = 1, \dots, ((d-1)/2 + 1)$, if d is odd. The extendibility of (τ_1, \dots, τ_d) to an *infinite* exchangeable sequence $\{\tau_k\}_{k \in \mathbb{N}}$, such that every finite subvector follows an exchangeable geometric distribution, corresponds to infinite extendibility of the associated triangle. This is equivalent to extendibility of the sequence $\{a_k\}_{k=1}^d$ to an infinite completely strong-monotone sequence $\{a_k\}_{k=1}^\infty \in \mathcal{SM}_\infty$, resp. extendibility of the sequence $(1, \beta_1, \dots, \beta_d)$ to an infinite completely monotone sequence. Corollary 4.1 summarizes these results.

Corollary 4.1 (Characterization of the infinitely extendible geometric law).

Denote $b_0 := 1$ and $\beta_0 := 1$.

(\mathcal{N}) Consider the parametric family $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ of all infinitely extendible, non-dege-

nerate d -dimensional narrow-sense geometric survival functions. Then,

$$\begin{aligned}\mathcal{G}^{\mathcal{N},\mathcal{E}} &= \{\bar{F}_{n_1,\dots,n_d}^{\mathcal{N}} = \prod_{k=1}^d a_k^{n(d-k+1)} \mid \{a_k\}_{k=1}^d \text{ is extendible to } \{a_k\}_{k \in \mathbb{N}} \in \mathcal{SM}_\infty\} \\ &= \{\bar{F}_{n_1,\dots,n_d}^{\mathcal{N}} = \prod_{k=1}^d b_k^{n(d-k+1)-n(d-k)} \mid \{b_k\}_{k=0}^d \text{ is extendible to } \{b_k\}_{k \in \mathbb{N}_0} \in \mathcal{LM}_\infty\}.\end{aligned}$$

(\mathcal{W}) Consider the parametric family $\mathcal{G}^{\mathcal{W},\mathcal{E}}$ of all infinitely extendible d -dimensional wide-sense geometric survival functions. Then,

$$\mathcal{G}^{\mathcal{W},\mathcal{E}} = \{\bar{F}_{n_1,\dots,n_d}^{\mathcal{W}} = \prod_{k=1}^d \beta_k^{n(d-k+1)-n(d-k)} \mid \{\beta_k\}_{k=0}^d \text{ is extendible to } \{\beta_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty\}.$$

PROOF. The assertion follows immediately from Theorem 3.5. \square

By Hausdorff's moment problem, the condition $\{\beta_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$ implies that $\{\beta_k\}_{k \in \mathbb{N}_0}$ is the sequence of moments of a probability measure on $[0, 1]$. In a similar manner, the completely log-monotone sequences $\{b_k\}_{k \in \mathbb{N}_0} \in \mathcal{LM}_\infty$ are related to infinitely divisible probability measures.

4.1. Relation to infinitely divisible distributions

The log-monotonicity of an infinite sequence is closely related to the existence of an infinitely divisible probability measure on $[0, \infty]$. Recall that a $[0, \infty]$ -valued random variable X is called infinitely divisible if for every $n \geq 2$ there exists a sequence of n independent and identically distributed $[0, \infty]$ -valued random variables, whose sum equals X in distribution.

Example 4.2 (Infinitely divisible distributions on $[0, \infty]$). Every infinitely divisible distribution, discrete or continuous, on $[0, \infty)$ can be modified to an infinitely divisible distribution on $[0, \infty]$, by weighting the probability mass function so that it integrates to some a , $a \in (0, 1)$, over $[0, \infty)$, and assigning the probability of $1 - a$ to the value ∞ . We illustrate this procedure taking the geometric distribution as an example. To this end, let $X \sim \text{Geo}(1-p)$. The distribution of X is infinitely divisible. More precisely, comparing the characteristic function of $X - 1$ with the one of the negative binomial distribution (see [32], p. 14), one immediately sees that for each $n \geq 2$, $X - 1$ equals in distribution the sum of n independent and identically negative binomially $\text{NB}(\frac{1}{n}, p)$ -distributed random variables. Now consider a modification \tilde{X} distributed on $\mathbb{N} \cup \{\infty\}$ according to $\tilde{X} \mid \tilde{X} < \infty \sim \text{Geo}(1-p)$, $\mathbb{P}(\tilde{X} < \infty) = a$, and $\mathbb{P}(\tilde{X} = \infty) = 1 - a$ for some $a \in (0, 1)$. Next, we show that \tilde{X} is infinitely divisible. Fix an arbitrary $n \geq 2$. Let Y_1, \dots, Y_n be independent and identically distributed random variables with $Y_1 \mid Y_1 < \infty \sim \text{NB}(\frac{1}{n}, p)$, $\mathbb{P}(Y_1 < \infty) = a^{\frac{1}{n}}$, and $\mathbb{P}(Y_1 = \infty) = 1 - a^{\frac{1}{n}}$. Then,

$$\mathbb{P}\left(1 + \sum_{i=1}^n Y_i = k\right) = \mathbb{P}\left(\sum_{i=1}^n Y_i = k - 1 \mid Y_i < \infty, i=1, \dots, n\right) \mathbb{P}(Y_1 < \infty)^n = \mathbb{P}(\tilde{X} = k),$$

and

$$\mathbb{P}\left(1 + \sum_{i=1}^n Y_i = \infty\right) = 1 - (1 - \mathbb{P}(Y_1 = \infty))^n = 1 - a = \mathbb{P}(\tilde{X} = \infty).$$

Thus, \tilde{X} equals in distribution $\sum_{i=1}^n (Y_i + 1/n)$, and the assertion follows.

The next lemma was originally used in the proof of Proposition 4.4 and is left here for the sake of our erratum.

Lemma 4.3 (Gnedin and Pitman [14]). *Let X be a $[0, \infty]$ -valued random variable with $\mathbb{P}(X < \infty) > 0$. Then, the following statements are equivalent.*

- (i) *The distribution of X is infinitely divisible on $[0, \infty]$.*
- (ii) *For all $k \in \mathbb{N}_0$, it holds that $\mathbb{E}[e^{-kX}] = e^{-\mu_k}$, where $\{-\nabla\mu_k\}_{k=0}^\infty$ is completely monotone and $\mu_0 = 0$.*

PROOF. This is simply a reformulation of Corollary 4.2 in [14] using their Definition 1.5. In the published manuscript, a nonsensical proof was given. \square

Proposition 4.4 is a simple corollary of Hausdorff's moment problem. It establishes a relationship between completely monotone sequences, completely log-monotone sequences, and the existence of an infinitely divisible probability measure.

Proposition 4.4 (Gnedin and Pitman [14]). *Let $\{b_k\}_{k \in \mathbb{N}_0}$ be sequence of strictly positive real numbers. The following statements are equivalent.*

- (i) $\{b_k\}_{k \in \mathbb{N}_0} \in \mathcal{LM}_\infty$.
- (ii) $\{b_k\}_{k \in \mathbb{N}_0}$ is given by

$$b_k = \int_{[0, \infty]} e^{-kx} \kappa(dx), \quad k \in \mathbb{N}_0,$$

for an infinitely divisible probability measure κ on $[0, \infty]$ with $\kappa(\{0\}) < 1$ and $\kappa(\{\infty\}) < 1$.

- (iii) $\{b_k^r\}_{k=0}^\infty \in \mathcal{M}_\infty$ for each fixed $r > 0$.
- (iv) $\{\sqrt[n]{b_k}\}_{k=0}^\infty \in \mathcal{M}_\infty$ for each fixed $n \in \mathbb{N}$.

PROOF. The equivalence of (i) and (ii) is simply a reformulation of Lemma 4.3 (in a context where we restrict ourselves to $b_1 < 1$, or, equivalently, to κ being not concentrated at 0). In the published manuscript, a nonsensical proof relying on the results of [21] was given. In Appendix C, we provide an alternative proof to the one given in [14]. \square

In part (ii) of Proposition 4.4, we use the convention $\infty \cdot 0 = 0$. The equivalence of (i) and (ii) constitutes an analogue of Hausdorff's moment problem. Results analogous to the equivalences of (ii), (iii), and (iv) for the Hausdorff moment sequence $\{b_k\}_{k=0}^\infty \in \mathcal{M}_\infty$ have been obtained for normalized Stieltjes as well as normalized Hamburger moment sequences. In our study, the complete log-monotonicity arises naturally as a characterizing property of parameter sequences of the $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ -law. Interestingly, it relates to the existing theory on logarithmically completely monotonic functions on $(0, \infty)$, for which, characterizations by completely monotonic functions, positive definite functions, or moment sequences resembling Proposition 4.4 are known. For an overview of the existing results on logarithmically completely monotonic functions and moment sequences, we refer to [4, 5].

As an immediate consequence to the finite-dimensional exchangeable case, the next corollary states that $(1, \beta_1, \beta_2, \dots) \in \mathcal{M}_\infty$ is *necessary and sufficient* for (12) to define a survival function in any dimension $d \in \mathbb{N}$.

Corollary 4.5 (Analytical characterization of the survival function).

The multi-indexed sequence $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$, given by

$$\bar{F}_{n_1, \dots, n_d} = \prod_{k=1}^d \beta_k^{n_{(d-k+1)} - n_{(d-k)}}, \quad n_1, \dots, n_d \in \mathbb{N}_0,$$

is a d -dimensional survival function for each $d \geq 2$ if and only if $(1, \beta_1, \beta_2, \dots) \in \mathcal{M}_\infty$. In this case, $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$ defines an infinitely extendible wide-sense geometric distribution.

PROOF. The corollary follows from Theorem 3.7. □

In view of de Finetti's theorem, extendible distributions are characterized by conditionally i.i.d. components. A stochastic representation in the sense of de Finetti's theorem can be achieved for both the $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ and $\mathcal{G}^{\mathcal{W},\mathcal{E}}$ families.

4.2. Stochastic representation in the sense of de Finetti's theorem

Theorem 4.6 provides an alternative stochastic model with components that are i.i.d. conditioned on a non-decreasing random walk for both the $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ and $\mathcal{G}^{\mathcal{W},\mathcal{E}}$ families. The $\mathcal{G}^{\mathcal{N},\mathcal{E}}$ -distribution corresponds to the proper subclass constructed by random walks with infinitely divisible increments.

Theorem 4.6 (Esary and Marshall [10]). Consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ supporting a sequence $\{X_k\}_{k \in \mathbb{N}}$ of i.i.d. $[0, \infty]$ -valued random variables with $\mathbb{P}(0 < X_1 \leq \infty) > 0$. Furthermore, let $\{E_k\}_{k \in \mathbb{N}}$ be an i.i.d. sequence of $\text{Exp}(1)$ -distributed random variables, independent of $\{X_k\}_{k \in \mathbb{N}}$. Define the random variables $\{\tau_k\}_{k \in \mathbb{N}}$ via

$$\tau_k := \min\{n \in \mathbb{N} : X_1 + \dots + X_n \geq E_k\}, \quad k \in \mathbb{N}.$$

(\mathcal{W}) For each $d \geq 2$, the random vector (τ_1, \dots, τ_d) has an exchangeable d -variate wide-sense geometric distribution with survival function (12), where

$$\beta_k = \mathbb{E}[e^{-k X_1}], \quad k \in \mathbb{N}.$$

(\mathcal{N}) If, additionally, the distribution of X_1 is infinitely divisible, then and only then, for each $d \geq 2$, the random vector (τ_1, \dots, τ_d) has an exchangeable narrow-sense geometric distribution with survival function (11), where

$$b_k = \mathbb{E}[e^{-k X_1}], \quad k \in \mathbb{N}.$$

Moreover, every extendible wide-sense (resp. narrow-sense) geometric distribution can be constructed using the probabilistic model in (\mathcal{W}) (resp. in (\mathcal{N})).

PROOF. Refer to [Appendix C](#). While we provide an alternative proof, the results found in [10] are more general. \square

In the situation of Theorem 4.6, one says that the geometrically distributed (τ_1, \dots, τ_d) is *extendible* to an infinite exchangeable sequence. By virtue of de Finetti's theorem, see [8] and [1], p. 20 ff., the random variables $\{\tau_k\}_{k \in \mathbb{N}}$ are i.i.d. conditioned on the σ -algebra generated by the sequence $\{X_k\}_{k \in \mathbb{N}}$. On the one hand, an extendible geometric law is parameterized by the sequence of moments of a single $[0, 1]$ -valued random variable. On the other hand, a single (infinitely divisible) random variable is sufficient for a simple and efficient construction of an extendible wide-sense (resp. narrow-sense) geometric law. This constitutes a practically convenient approach to provide low-parametric families of high-dimensional geometric distributions.

Example 4.7 (Three-variate extendible geometric law).

Consider the survival function (15) of a three-variate exchangeable distribution with mutually independent geometric marginals parameterized by the sequence $(1, p, p^2, p^3)$ from Example 3.6. (15) is infinitely extendible and can be constructed as in Theorem 4.6 (\mathcal{N}) using the degenerate distribution $X_1 := -\ln p$. A degenerate distribution is, trivially, infinitely divisible.

The survival function (16), parameterized by $(1, p, p, p)$ and corresponding to the upper Fréchet-Hoeffding bound from Example 3.6, is infinitely extendible as well. It can be constructed as in Theorem 4.6 (\mathcal{N}) using the infinitely divisible distribution $\mathbb{P}(X_1 = 0) = p = 1 - \mathbb{P}(X_1 = \infty)$.

Remark 4.8 (Link to the Bernoulli sieve). The construction of Theorem 4.6 involves a non-decreasing random walk with increments $\{X_k\}_{k \in \mathbb{N}}$. However, there is another possible construction which is motivated by coin tosses. It is called the Bernoulli sieve in [12] and defines $\{\tau_k\}_{k \in \mathbb{N}}$ as follows:

$$\tau_k := \min\{n \in \mathbb{N} : Z_k^{(n)} = 1\}, \quad k \in \mathbb{N},$$

where the sequences $\{Z_k^{(1)}\}_{k \in \mathbb{N}}, \{Z_k^{(2)}\}_{k \in \mathbb{N}}, \dots$ are independent and for each $n \in \mathbb{N}$, $\{Z_k^{(n)}\}_{k \in \mathbb{N}}$ is a sequence of Bernoulli random variables conditioned on a random success probability $1 - Y_n \in [0, 1]$. It is assumed that the sequence $\{Y_n\}_{n \in \mathbb{N}}$ is i.i.d. and $\mathbb{P}(Y_1 < 1) > 0$. The distribution of the $\{Y_k\}_{k \in \mathbb{N}}$ is precisely η from the proof of Theorem 4.6 above. In words, imagine a game with infinitely many players labeled by $k \in \mathbb{N}$. The game has (possibly) infinitely many rounds, and in each round a certain number of players drops out. In round $n \in \mathbb{N}$, a referee randomly picks a coin with probability of tails $1 - Y_n$. Then, for each player the coin is thrown once and the player drops out if she gets tails. The random variable τ_k denotes the drop out round of player k . The condition $\mathbb{P}(Y_1 < 1) > 0$ guarantees that each player drops out eventually.

Whereas the present article studies the distribution of (τ_1, \dots, τ_d) , [12] studies the distribution of (P_1, \dots, P_{K_d}) , where

$K_d := |\{\tau_1, \dots, \tau_d\}| = \#$ of rounds in which one of the first d players drops out,

and, with $\tau_{(1)} \leq \dots \leq \tau_{(d)}$ denoting the order statistics of (τ_1, \dots, τ_d) ,

$P_1 := \max\{j : \tau_{(j)} = \tau_{(1)}\} = \#$ of drop outs in the first of these rounds,

$P_2 := \max\{j : \tau_{(P_1+j)} = \tau_{(P_1+1)}\} = \#$ of drop outs in the second of these rounds,

\vdots

$P_{K_d} := \max\{j : \tau_{(P_{K_d-1}+j)} = \tau_{(P_{K_d-1}+1)}\} = \#$ of drop outs in the last of these rounds.

Notice that $P_1, \dots, P_{K_d} \in \mathbb{N}$ and $P_1 + \dots + P_{K_d} = d$, hence (P_1, \dots, P_{K_d}) defines a random partition of d , and vast literature can be found on the study of such objects. [13] extend the Bernoulli sieve to a “continuous” case using Lévy subordinators. Considering non-decreasing random walks as discrete equivalents of Lévy subordinators explains their appearance in Theorem 4.6. Furthermore, [23] show that the extendible subclass of Marshall–Olkin exponential distributions can be characterized by Lévy subordinators. In view of these references, the present article completes the picture, see Table 1.

Table 1: Stochastic objects with limited memory.

	<i>non-decreasing processes</i>	<i>random integer partitions</i>	<i>extendible multivariate laws</i>
<i>continuous</i>	<i>Lévy subordinators</i>	<i>regenerative composition structures, see [13]</i>	<i>Marshall–Olkin distributions, see [23]</i>
<i>discrete</i>	<i>random walks with non-negative increments</i>	<i>the Bernoulli sieve, see [12]</i>	<i>$\mathcal{G}^{\mathcal{W}, \varepsilon}$ and $\mathcal{G}^{\mathcal{N}, \varepsilon}$, the present article</i>

5. Dependence structure

In this section, we study the dependence structure of the wide- and the narrow-sense geometric distributions discussed in Sections 2, 3, and 4. Whereas the narrow-sense construction always leads to distributions with non-negatively correlated components, the more general wide-sense geometric distribution allows for negative correlations. Taking into account the specific form of the survival function, the explicit formulas for the correlation coefficients can be conveniently obtained by writing $\tau_i = \sum_{n=0}^{\infty} \mathbf{1}_{\{\tau_i > n\}}$ for \mathbb{N} -valued τ_i , $i = 1, 2$, and computing the mixed second moment via

$$\mathbb{E}[\tau_1 \tau_2] = \sum_{i \geq 0} \sum_{j \geq 0} \mathbb{P}(\tau_1 > i, \tau_2 > j).$$

The computation is tedious but straightforward, and therefore omitted. The formulas for the correlation coefficients for the wide- and the narrow-sense geometric distribution in dimension $d \geq 2$, in both exchangeable and general cases, are provided in Lemma 5.1. The conditions on the involved parameters are given in Theorem 3.5, in the case of an exchangeable distribution, and in Definition 2.3, in the general case.

Lemma 5.1 (Correlation coefficients).

(\mathcal{N}) Let $d \geq 2$ and let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{N}}(\mathbf{p})$. Then, for $n, m \in \{1, \dots, d\}$, $n \neq m$, the correlation coefficient $\text{Corr}[\tau_n, \tau_m]$ is given by

$$\text{Corr}[\tau_n, \tau_m] = \frac{\sqrt{\prod_{I: (n \in I, m \notin I) \cup (n \notin I, m \in I)} p_I (1 - \prod_{I: n, m \in I} p_I)}}{1 - \prod_{I: (n \in I) \cup (m \in I)} p_I}.$$

If, additionally, the distribution of (τ_1, \dots, τ_d) is exchangeable, then

$$\text{Corr}[\tau_n, \tau_m] = \frac{a_2 - a_1}{1 - a_1 a_2} = \frac{b_2 - b_1^2}{b_1(1 - b_2)}, \quad n, m \in \{1, \dots, d\}, n \neq m,$$

with a_1, a_2 and b_1, b_2 defined in (8) and (10), respectively.

(\mathcal{W}) Let $d \geq 2$ and let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{W}}(\tilde{\mathbf{p}})$. Then, for $n, m \in \{1, \dots, d\}$, $n \neq m$, such that $\sum_{I: n \notin I} \tilde{p}_I \neq 0$ and $\sum_{I: m \notin I} \tilde{p}_I \neq 0$, the correlation coefficient $\text{Corr}[\tau_n, \tau_m]$ is given by

$$\text{Corr}[\tau_n, \tau_m] = \frac{\sum_{I: n, m \notin I} \tilde{p}_I - \sum_{I: n \notin I} \sum_{J: m \notin J} \tilde{p}_I \tilde{p}_J}{(1 - \sum_{I: n, m \notin I} \tilde{p}_I) \sqrt{\sum_{I: n \notin I} \sum_{J: m \notin J} \tilde{p}_I \tilde{p}_J}}.$$

If, additionally, the distribution of (τ_1, \dots, τ_d) is exchangeable, then

$$\text{Corr}[\tau_n, \tau_m] = \frac{\beta_2 - \beta_1^2}{\beta_1(1 - \beta_2)}, \quad n, m \in \{1, \dots, d\}, n \neq m,$$

with β_1, β_2 defined in (13).

PROOF. A tedious but straightforward computation provides the results. \square

In part (W) of Lemma 5.1, we exclude the degenerate case $\sum_{I:n \notin I} \tilde{p}_I = 0$, $n \in \{1, \dots, d\}$, corresponding to $\tau_n = 1$ almost surely. For exchangeable (τ_1, \dots, τ_d) , this translates into $\beta_1 \neq 0$.

Example 5.2 (Geometric in the wide-, but not in the narrow sense). Consider again the sequence $(\beta_0, \beta_1, \beta_2) = (1, \frac{1}{2}, \frac{1}{5}) \in \mathcal{M}_3$ from Examples 3.4 and 3.8, defining parameters of a bivariate $\mathcal{G}^{\mathcal{W}, \mathcal{X}}$ -distribution. With Lemma 5.1 (W), we compute

$$\text{Corr}[\tau_1, \tau_2] = \frac{\beta_2 - \beta_1^2}{\beta_1(1 - \beta_2)} = -\frac{1}{8}.$$

In the case of a wide-sense geometric distribution, the correlation coefficient between every two components is bounded from below by -0.5 in any dimension $d \geq 2$. The correlation coefficient $\text{Corr}[\tau_n, \tau_m] = -0.5$ is obtained in the case $\tilde{p}_I = 2^{1-d}$ for all $I \subseteq \{1, \dots, d\}$ s.t. $n \in I$, $m \notin I$ or $m \in I$, $n \notin I$, and $\tilde{p}_I = 0$, otherwise. Note that due to $|\{I : n \in I, m \notin I\}| = 2^{d-2}$, these parameters satisfy the conditions of Definition 2.3 (W).

In the case of an exchangeable d -variate distribution, the correlation coefficient is bounded from below by $-1/d$ for any $d \geq 2$. This can be verified by observing that for $\beta_1 \neq 0$, the correlation coefficient

$$\text{Corr}[\tau_n, \tau_m] = \frac{\beta_2 - \beta_1^2}{\beta_1(1 - \beta_2)}, \quad n, m = 1, \dots, d,$$

is strictly increasing in β_2 and strictly decreasing in β_1 . Setting $\beta_2 = 0$, the condition $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ gives $\beta_k = 0$ for all $k = 2, \dots, d$ and $(1 - d\beta_1) \geq 0$. Hence, the minimum is attained at $(\beta_0, \dots, \beta_d) = (1, 1/d, 0, \dots, 0) \in \mathcal{M}_{d+1}$. Recall that the lower bound for the correlation coefficient between the components of an arbitrary exchangeable d -variate distribution is given by $-1/(d-1)$ for any $d \geq 2$, see [20] or [1], p. 8.

Remark 5.3 (Correlations induced by the Fréchet-Hoeffding bounds).

Let $\tau_1 \sim \text{Geo}(1 - p_1)$ and $\tau_2 \sim \text{Geo}(1 - p_2)$, $p_1, p_2 \in (0, 1)$. Assume that the dependence between τ_1 and τ_2 is now given by the lower Fréchet-Hoeffding bound, i.e.

$$\mathbb{P}(\tau_1 \leq n_1, \tau_2 \leq n_2) = \max\{0, 1 - p_1^{n_1} - p_2^{n_2}\}.$$

Then, the correlation coefficient $\text{Corr}[\tau_1, \tau_2]$ is given by

$$\text{Corr}[\tau_1, \tau_2] = \frac{(1-p_1)(1-p_2)}{\sqrt{p_1 p_2}} \left(\sum_{i \geq 0} \sum_{j \geq 0} \max\{0, p_1^i + p_2^j - 1\} - \frac{1}{(1-p_1)(1-p_2)} \right).$$

In the special case $p_1 + p_2 \leq 1$, the expression for the correlation coefficient simplifies to

$$\text{Corr}[\tau_n, \tau_m] = -\sqrt{p_1 p_2}.$$

Recall that correlations induced by the lower Fréchet-Hoeffding bound are always non-positive. In our case of geometric marginals, we observe $\text{Corr}[\tau_1, \tau_2] \rightarrow 0$ as $p_1 \rightarrow 0$, $p_2 \rightarrow 0$, and $\text{Corr}[\tau_1, \tau_2] \rightarrow -1$ as $p_1 \rightarrow 1$, $p_2 \rightarrow 1$. It is worth noticing that a distribution with non-degenerate geometric marginals and the lower Fréchet-Hoeffding copula is not geometric in the wide-sense by Theorem 2.6.

Now let the dependence between $\tau_k \sim \text{Geo}(1-p_k)$, $p_k \in (0, 1)$, $k \in \{1, \dots, d\}$, be given by the upper Fréchet-Hoeffding bound, i.e.

$$\mathbb{P}(\tau_1 \leq n_1, \dots, \tau_d \leq n_d) = \min \left\{ 1 - p_1^{n_1}, \dots, 1 - p_d^{n_d} \right\}.$$

Then, for $n, m \in \{1, \dots, d\}$, $n \neq m$, the correlation coefficient $\text{Corr}[\tau_n, \tau_m]$ is given by

$$\text{Corr}[\tau_n, \tau_m] = \frac{(1-p_n)(1-p_m)}{\sqrt{p_n p_m}} \left(\sum_{i \geq 0} \sum_{j \geq 0} \min \{ p_n^i, p_m^j \} - \frac{1}{(1-p_n)(1-p_m)} \right).$$

Recall that correlations induced by the upper Fréchet-Hoeffding bound are always non-negative. In the homogeneous case $p_1 = p_k$ for all $k = 2, \dots, d$, we obtain $\text{Corr}[\tau_n, \tau_m] = 1$ for all $p_1 \in (0, 1)$. More precisely,

$$\mathbb{E}[\tau_n \tau_m] = \sum_{i \geq 0} \sum_{j \geq 0} \min \{ p_1^i, p_1^j \} = \sum_{i \geq 0} \left(\sum_{j=0}^i p_1^i + \sum_{j=i+1}^{\infty} p_1^j \right) = \frac{1+p_1}{(1-p_1)^2},$$

and, hence,

$$\text{Corr}[\tau_n, \tau_m] = \left(\frac{1+p_1}{(1-p_1)^2} - \frac{1}{(1-p_1)^2} \right) \left(\frac{(1-p_1)^2}{p_1} \right) = 1.$$

Moreover, in this case, the corresponding distribution is an exchangeable narrow-sense geometric distribution with survival function

$$\bar{F}_{n_1, \dots, n_d} = 1 + \sum_{i=1}^d \sum_{\substack{\emptyset \neq I \subseteq \{1, \dots, d\} \\ |I|=i}} (-1)^i \mathbb{P}(\tau_j \leq n_j, j \in I) = p_1^{n(d)}.$$

The following example shows that the set of all bivariate $\mathcal{G}^{\mathcal{N}}$ -distributions is precisely the set of all bivariate $\mathcal{G}^{\mathcal{W}}$ -distributions with non-negative correlations.

Example 5.4 (Non-negative correlations $\Rightarrow \mathcal{G}^{\mathcal{N}}$ in $d = 2$). In view of Theorems 2.5 and 2.6, every $(\tau_1, \tau_2) \sim \mathcal{G}^{\mathcal{N}}$ follows a $\mathcal{G}^{\mathcal{W}}$ -distribution with non-negatively correlated components. We show that the converse also holds. To this end, let $(\tau_1, \tau_2) \sim \mathcal{G}^{\mathcal{W}}$ with $\text{Corr}[\tau_1, \tau_2] \geq 0$. By Theorem 2.5 (\mathcal{W}),

$$\mathbb{P}(\tau_1 > i, \tau_2 > j) = \begin{cases} (\tilde{p}_\emptyset + \tilde{p}_{\{2\}})^{i-j} \tilde{p}_\emptyset^j, & \text{if } i \geq j, \\ (\tilde{p}_\emptyset + \tilde{p}_{\{1\}})^{j-i} \tilde{p}_\emptyset^i, & \text{if } i < j, \end{cases}$$

with parameters satisfying $\sum_{I \subseteq \{1,2\}} \tilde{p}_I = 1$, $\sum_{I: k \notin I} \tilde{p}_I < 1$, for $k = 1, 2$, and $\tilde{p}_\emptyset - (\tilde{p}_\emptyset + \tilde{p}_{\{1\}})(\tilde{p}_\emptyset + \tilde{p}_{\{2\}}) \geq 0$, where the last condition is due to non-negativity of the correlation coefficient, see Lemma 5.1 (W). For $\tilde{p}_\emptyset > 0$, define

$$p_{\{1\}} := \frac{\tilde{p}_\emptyset}{\tilde{p}_\emptyset + \tilde{p}_{\{1\}}}, \quad p_{\{2\}} := \frac{\tilde{p}_\emptyset}{\tilde{p}_\emptyset + \tilde{p}_{\{2\}}}, \quad p_{\{1,2\}} := \frac{(\tilde{p}_\emptyset + \tilde{p}_{\{1\}})(\tilde{p}_\emptyset + \tilde{p}_{\{2\}})}{\tilde{p}_\emptyset}.$$

$p_{\{1\}}$, $p_{\{2\}}$, and $p_{\{1,2\}}$ are well-defined and satisfy the conditions of Definition 2.3 (N). Further,

$$\mathbb{P}(\tau_1 > i, \tau_2 > j) = \begin{cases} (p_{\{1\}}p_{\{1,2\}})^i p_{\{2\}}^j, & \text{if } i \geq j, \\ (p_{\{2\}}p_{\{1,2\}})^j p_{\{1\}}^i, & \text{if } i < j. \end{cases}$$

Hence, by Theorem 2.5 (N), $(\tau_1, \tau_2) \sim \mathcal{G}^{\mathcal{N}}(p_{\{1\}}, p_{\{2\}}, p_{\{1,2\}})$.

On the contrary, in dimension $d \geq 3$, the set of all d -variate $\mathcal{G}^{\mathcal{N}}$ -distributions is a proper subset of all d -variate $\mathcal{G}^{\mathcal{W}}$ -distributions with non-negative correlations.

Example 5.5 (Non-negative correlations $\not\Rightarrow \mathcal{G}^{\mathcal{N}}$ in $d \geq 3$). Let $d \geq 3$ and let X be Bernoulli distributed with success probability $1 - p = 0.9$. For $k = 0, \dots, d$, define $\beta_k := \mathbb{E}[e^{-kX}] = p + (1 - p)e^{-k}$. By Theorem 4.6 (W), $(\beta_0, \dots, \beta_d)$ defines parameters of a d -variate $\mathcal{G}^{\mathcal{W}, \mathcal{E}}$ -distribution. Using the formula for correlation coefficients from Lemma 5.1 (W),

$$\text{Corr}[\tau_n, \tau_m] = \frac{\beta_2 - \beta_1^2}{\beta_1(1 - \beta_2)} = \frac{0.1 + 0.9e^{-2} - (0.1 + 0.9e^{-1})^2}{(0.1 + 0.9e^{-1})(0.9 - 0.9e^{-2})} \approx 0.1072 > 0,$$

for $n, m \in \{1, \dots, d\}$, $n \neq m$. However, since

$$\nabla^3 \ln \beta_0 = \ln \frac{\beta_2^3}{\beta_1^3 \beta_3} \approx -0.06 < 0,$$

it follows that $(\beta_0, \dots, \beta_d) \notin \mathcal{LM}_{d+1}$, and, hence, in view of Theorem 3.5 (N), $(\beta_0, \dots, \beta_d)$ does not define parameters of a d -variate $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distribution. In this context, note that the Bernoulli distribution is not infinitely divisible, and, by Theorem 4.6 (N), $(\beta_0, \dots, \beta_d)$ defines parameters of a d -variate $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distribution for each $d \geq 2$ if and only if the distribution of X is infinitely divisible.

Remark 5.6 (Relationships between subfamilies of the geometric law). The relationships between exchangeable subfamilies of the multivariate geometric law with limited memory can be summarized as follows:

(i) $\mathcal{G}^{\mathcal{N}, \mathcal{E}} \subseteq \mathcal{G}^{\mathcal{N}, \mathcal{X}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{X}}$ and $\mathcal{G}^{\mathcal{N}, \mathcal{E}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{E}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{X}}$.

The inclusion $\mathcal{G}^{\mathcal{N}, \mathcal{X}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{X}}$ is proper in every dimension $d \geq 2$ by Example 3.8. The inclusion $\mathcal{G}^{\mathcal{W}, \mathcal{E}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{X}}$ is proper in every dimension $d \geq 2$ by Example 3.4.

(ii) In addition, for exchangeable bivariate geometric distributions:

$$\mathcal{G}^{\mathcal{N},\mathcal{X}} = \mathcal{G}^{\mathcal{N},\mathcal{E}} = \mathcal{G}^{\mathcal{W},\mathcal{E}} \subsetneq \mathcal{G}^{\mathcal{W},\mathcal{X}}.$$

More precisely, by Hausdorff's moment problem, $(1, \beta_1, \beta_2)$ is extendible to an infinite sequence $(1, \beta_1, \beta_2, \dots) \in \mathcal{M}_\infty$ if and only if $\beta_k = \mathbb{E}[\tau^k]$, $k = 1, 2$, for a $[0, 1]$ -valued random variable τ . Then, by Jensen's inequality $\beta_2 - \beta_1^2 \geq 0$. Hence, every $\mathcal{G}^{\mathcal{W},\mathcal{E}}$ -distribution has non-negatively correlated components, see Lemma 5.1 (\mathcal{W}). It follows by Example 5.4 that $\mathcal{G}^{\mathcal{W},\mathcal{E}} = \mathcal{G}^{\mathcal{N},\mathcal{E}} \subseteq \mathcal{G}^{\mathcal{N},\mathcal{X}}$ in dimension $d = 2$. It remains to justify that $\mathcal{G}^{\mathcal{N},\mathcal{E}} \supseteq \mathcal{G}^{\mathcal{N},\mathcal{X}}$, meaning that each sequence $(1, b_1, b_2) \in \mathcal{LM}_3$ is infinitely extendible to a sequence $(1, b_1, b_2, \dots) \in \mathcal{LM}_\infty$. Let $(1, b_1, b_2) \in \mathcal{LM}_3$. On the one hand, by 3-log-monotonicity of $(1, b_1, b_2)$, it holds that $b_1^2 \leq b_2 \leq b_1 < 1$. On the other hand, by Proposition 4.4, $(1, b_1, b_2) \in \mathcal{LM}_3$ is infinitely extendible to a sequence $(1, b_1, b_2, \dots) \in \mathcal{LM}_\infty$ if and only if $(1, b_1^r, b_2^r)$ is infinitely extendible to a sequence $(1, b_1^r, b_2^r, \dots) \in \mathcal{M}_\infty$ for each fixed $r > 0$. Finally, Hankel's criterion for infinite extendibility of d -monotone sequences, see [9], p. 20, implies that $(1, b_1^r, b_2^r)$ is infinitely extendible to $(1, b_1^r, b_2^r, \dots) \in \mathcal{M}_\infty$ if and only if $b_1^{2r} \leq b_2^r \leq b_1^r \leq 1$ for each fixed $r > 0$. This gives the assertion.

In dimension $d \geq 3$, none of the inclusions $\mathcal{G}^{\mathcal{W},\mathcal{E}} \subseteq \mathcal{G}^{\mathcal{N},\mathcal{X}}$ and $\mathcal{G}^{\mathcal{N},\mathcal{X}} \subseteq \mathcal{G}^{\mathcal{W},\mathcal{E}}$ holds. On the one hand, $\mathcal{G}^{\mathcal{W},\mathcal{E}} \not\subseteq \mathcal{G}^{\mathcal{N},\mathcal{X}}$ by Example 5.5. On the other hand, it can be shown that not every $(1, b_1, b_2, b_3) \in \mathcal{LM}_4$ is extendible to a sequence $(1, b_1, b_2, b_3, \dots) \in \mathcal{M}_\infty$.

The class of exchangeable narrow-sense geometric distributions possesses a strong positive dependence property, namely the multivariate right tail increasing (MRTI) dependence. MRTI implies, in particular, the positive upper orthant increasing ratio dependence, the positive upper orthant dependence, and, hence, non-negative correlations. For a detailed discussion of the notions of multivariate positive dependence, we refer to [6, 7, 38]. For the convenience of the reader, we recall that a random vector (τ_1, \dots, τ_d) with values in \mathbb{N}^d is said to be MRTI if for all permutations $\pi: \{1, \dots, d\} \rightarrow \{1, \dots, d\}$

$$\mathbb{P}(\tau_{\pi(i)} > n_i \mid \tau_{\pi(1)} > n_1, \dots, \tau_{\pi(i-1)} > n_{i-1})$$

is increasing in n_1, \dots, n_{i-1} for all $n_i, i \in \{2, \dots, d\}$.

We show in Theorem 5.7 that

$$\frac{a_k}{a_{k+1}} = \frac{b_k^2}{b_{k-1}b_{k+1}} \leq 1 \quad (\text{resp. } \frac{\beta_k^2}{\beta_{k-1}\beta_{k+1}} \leq 1), \quad k = 1, \dots, d-1,$$

is a necessary and sufficient condition on the parameters of a d -dimensional $\mathcal{G}^{\mathcal{N},\mathcal{X}}$ - (resp. $\mathcal{G}^{\mathcal{W},\mathcal{X}}$ -) survival function, given by (9) (resp. (12)), to be MRTI. In view of Theorem 3.5 (\mathcal{N}), it holds that $\{a_k\}_{k=1}^d \in \mathcal{SM}_d$ for any set of parameters $\{a_k\}_{k=1}^d$ of a d -variate $\mathcal{G}^{\mathcal{N},\mathcal{X}}$ -distribution. Hence, $\{a_k\}_{k=1}^d$ is non-decreasing, and it follows that every d -variate $\mathcal{G}^{\mathcal{N},\mathcal{X}}$ -distribution is MRTI positively dependent. The next theorem summarizes these results.

Theorem 5.7 (MRTI property for exchangeable distributions).

(\mathcal{N}) Every $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distributed (τ_1, \dots, τ_d) is MRTI positively dependent.

(\mathcal{W}) A $\mathcal{G}^{\mathcal{W}, \mathcal{X}}$ -distributed (τ_1, \dots, τ_d) is MRTI positively dependent if and only if the parameters $\{\beta_k\}_{k=0}^d$ of its corresponding survival function $\bar{F}^{\mathcal{W}}$, given by (12), satisfy $\frac{\beta_k^2}{\beta_{k-1}\beta_{k+1}} \leq 1$ for all $k = 1, \dots, d-1$.

PROOF. Refer to Appendix D. □

On the one hand, exchangeable wide-sense geometric distributions possess an MRTI positive dependence structure under the conditions given in Theorem 5.7 (\mathcal{W}). On the other hand, as illustrated in Example 5.2, exchangeable wide-sense geometric distributions in general allow for negative correlations. Extendible (narrow- and wide-sense) geometric distributions are always MRTI. This is the content of Corollary 5.8.

Corollary 5.8 (MRTI property for extendible distributions). *Every extendible geometric distribution satisfying the local LM property is MRTI positively dependent.*

PROOF. In view of Corollary 4.1 and Theorem 5.7 (\mathcal{W}), it remains to show that any $\{\beta_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$ satisfies $\beta_k^2 \leq \beta_{k-1}\beta_{k+1}$ for all $k \in \mathbb{N}$. By Hausdorff's moment problem, $\{\beta_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$ is equivalent to $\beta_k = \mathbb{E}[\tau^k]$ for $k \in \mathbb{N}_0$ and a $[0, 1]$ -valued random variable τ . The Cauchy-Schwarz inequality gives $\mathbb{E}[\tau^k]^2 \leq \mathbb{E}[\tau^{k-1}]\mathbb{E}[\tau^{k+1}]$ for all $k \in \mathbb{N}$, and the corollary follows. □

Appendix A. Proofs from Section 2

PROOF OF THEOREM 2.5. To prove (\mathcal{N}), let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{N}}(\mathbf{p})$. Check that

$$\mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) = \mathbb{P}(E_I > \max_{i \in I} \{n_i\}, \emptyset \neq I \subseteq \{1, \dots, d\}) = \prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{i \in I} \{n_i\}}.$$

Next, we show that the claimed survival function (1) satisfies the local LM property. Fix $k \in \{1, \dots, d\}$ and let $1 \leq i_1 < \dots < i_k \leq d$, $m, n_{i_1}, \dots, n_{i_k} \in \mathbb{N}_0$. Define $\tilde{n}_1, \dots, \tilde{n}_d$ by

$$\tilde{n}_j := \begin{cases} n_j, & j \in \{i_1, \dots, i_k\} \\ 0, & j \notin \{i_1, \dots, i_k\} \end{cases}.$$

Then,

$$\prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{j \in I} \{\tilde{n}_j + m\}} = \left(\prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{j \in I} \{\tilde{n}_j\}} \right) \left(\prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{j \in I} \{m\}} \right),$$

which implies the assertion.

To prove (\mathcal{W}), let $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{W}}(\tilde{\mathbf{p}})$. The event $\{\tau_1 > n_1, \dots, \tau_d > n_d\}$

means that the first $n_{(1)}$ outcomes are of type $I = \emptyset$, the next $n_{(2)} - n_{(1)}$ outcomes are $I \in \{\emptyset, \{\pi_n(1)\}\}$, the next $n_{(3)} - n_{(2)}$ outcomes are $I \in \{\emptyset, \{\pi_n(1)\}, \{\pi_n(2)\}, \{\pi_n(1), \pi_n(2)\}\}$, etc. Thus, with $n_{(0)} := 0$ we have

$$\mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) = \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_n(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I \right)^{n_{(k)} - n_{(k-1)}}.$$

Finally, we show that the survival function (2) satisfies the local LM property. Again, fix $k \in \{1, \dots, d\}$ and let $1 \leq i_1 < \dots < i_k \leq d$, $m, n_{i_1}, \dots, n_{i_k} \in \mathbb{N}_0$. With $\tilde{n}_1, \dots, \tilde{n}_d$ defined as above

$$\bar{F}_{\tilde{n}_1+m, \dots, \tilde{n}_d+m}^{\mathcal{W}} = \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_n(i) \notin I \forall i=1, \dots, d}} \tilde{p}_I \right)^m \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_n(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I \right)^{\tilde{n}_{(k)} - \tilde{n}_{(k-1)}} = \bar{F}_{m, \dots, m}^{\mathcal{W}} \bar{F}_{\tilde{n}_1, \dots, \tilde{n}_d}^{\mathcal{W}}.$$

Hence the theorem. \square

PROOF OF THEOREM 2.6. By Theorem 2.5, the d -variate wide-sense geometric distribution satisfies the local discrete LM property (3). Let (τ_1, \dots, τ_d) be a random vector on \mathbb{N}^d satisfying the local discrete LM property (3). For $I \subseteq \{1, \dots, d\}$, define

$$\tilde{p}_I = \mathbb{P}(\{\tau_i > 1 \forall i \notin I\} \cap \{\tau_i = 1 \forall i \in I\}). \quad (\text{A.1})$$

We show that (τ_1, \dots, τ_d) follows a $\mathcal{G}^{\mathcal{W}}(\{\tilde{p}_I\}_{I \subseteq \{1, \dots, d\}})$ -distribution with survival function (2). First, we prove that (i) for $k = 1, \dots, d$,

$$\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I = \mathbb{P}(\tau_{\pi(k)} > 1, \dots, \tau_{\pi(d)} > 1),$$

for any permutation $\pi: \{1, \dots, d\} \rightarrow \{1, \dots, d\}$, (ii) $\sum_I \tilde{p}_I = 1$, (iii) $\tilde{p}_I \in [0, 1]$, and (iv) $\sum_{I: k \notin I} \tilde{p}_I < 1$ for $k = 1, \dots, d$. To see that (i) holds, check that for $k = 1, \dots, d$

$$\begin{aligned} \sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I &= \sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi(i) \notin I \forall i=k, \dots, d}} \mathbb{P}(\{\tau_i > 1 \forall i \notin I\} \cap \{\tau_i = 1 \forall i \in I\}) \\ &= \mathbb{P}(\tau_1 > 1, \dots, \tau_d > 1) \\ &\quad + \sum_{n=1}^{k-1} \sum_{1 \leq i_1 < \dots < i_n \leq k-1} \mathbb{P}(\{\tau_j > 1 \forall j \notin \{\pi(i_1), \dots, \pi(i_n)\}\} \cap \{\tau_j = 1 \forall j \in \{\pi(i_1), \dots, \pi(i_n)\}\}) \\ &= \mathbb{P}(\tau_{\pi(k)} > 1, \dots, \tau_{\pi(d)} > 1), \end{aligned}$$

where the last equality is due to

$$\begin{aligned} &\bigcup_{\substack{1 \leq i_1 < \dots < i_n \leq k-1, \\ n=1, \dots, k-1}} \left\{ \{\tau_j > 1 \forall j \notin \{\pi(i_1), \dots, \pi(i_n)\}\} \cap \{\tau_j = 1 \forall j \in \{\pi(i_1), \dots, \pi(i_n)\}\} \right\} \cup \{\tau_1 > 1, \dots, \tau_d > 1\} \\ &= \{\tau_i > 1 \forall i \notin \{\pi(1), \dots, \pi(k-1)\}\} = \{\tau_{\pi(k)} > 1, \dots, \tau_{\pi(d)} > 1\}. \end{aligned}$$

(ii) follows immediately from (A.1) and the fact that (τ_1, \dots, τ_d) is \mathbb{N}^d -valued:

$$\sum_{I \subseteq \{1, \dots, d\}} \tilde{p}_I = \sum_{I \subseteq \{1, \dots, d\}} \mathbb{P}(\{\tau_i > 1 \ \forall i \notin I\} \cap \{\tau_i = 1 \ \forall i \in I\}) = \mathbb{P}(\tau_1 \geq 1, \dots, \tau_d \geq 1) = 1.$$

(iii) is trivial, and (iv) follows from the finiteness of the components τ_k , $k = 1, \dots, d$. More precisely, using (i), it follows for $k = 1, \dots, d$, that

$$\sum_{I: k \notin I} \tilde{p}_I = \mathbb{P}(\tau_k > 1) < 1,$$

since, otherwise, by the LM property, $\mathbb{P}(\tau_k > n) = \mathbb{P}(\tau_k > 1)^n = 1$ for all $n \in \mathbb{N}_0$, contradicting $\tau_k < \infty$ almost surely. Finally, we show that $(\tau_1, \dots, \tau_d) \sim \mathcal{G}^{\mathcal{W}}(\{\tilde{p}_I\}_{I \subseteq \{1, \dots, d\}})$ with survival function as in (2). By the local LM property (3),

$$\mathbb{P}(\tau_{i_1} > n, \dots, \tau_{i_k} > n) = \mathbb{P}(\tau_{i_1} > 1, \dots, \tau_{i_k} > 1)^n,$$

for $k = 1, \dots, d$, $1 \leq i_1 < \dots < i_k \leq d$, and $n \in \mathbb{N}_0$. From this, with $\pi_{\mathbf{n}}: \{1, \dots, d\} \rightarrow \{1, \dots, d\}$ being a permutation depending on $\mathbf{n} = (n_1, \dots, n_d)$ such that $n_{\pi_{\mathbf{n}}(1)} \leq n_{\pi_{\mathbf{n}}(2)} \leq \dots \leq n_{\pi_{\mathbf{n}}(d)}$, we derive

$$\begin{aligned} \mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) &= \mathbb{P}(\tau_{\pi_{\mathbf{n}}(1)} > n_{(1)}, \dots, \tau_{\pi_{\mathbf{n}}(d)} > n_{(d)}) \\ &= \mathbb{P}(\tau_{\pi_{\mathbf{n}}(1)} > n_{(1)}, \tau_{\pi_{\mathbf{n}}(2)} > n_{(1)} + (n_{(2)} - n_{(1)}), \dots, \tau_{\pi_{\mathbf{n}}(d)} > n_{(1)} + (n_{(d)} - n_{(1)})) \\ &= \mathbb{P}(\tau_{\pi_{\mathbf{n}}(1)} > n_{(1)}, \dots, \tau_{\pi_{\mathbf{n}}(d)} > n_{(1)}) \mathbb{P}(\tau_{\pi_{\mathbf{n}}(2)} > n_{(2)} - n_{(1)}, \dots, \tau_{\pi_{\mathbf{n}}(d)} > n_{(d)} - n_{(1)}) \\ &= \mathbb{P}(\tau_1 > 1, \dots, \tau_d > 1)^{n_{(1)}} \mathbb{P}(\tau_{\pi_{\mathbf{n}}(2)} > 1, \dots, \tau_{\pi_{\mathbf{n}}(d)} > 1)^{n_{(2)} - n_{(1)}} \\ &\quad \times \mathbb{P}(\tau_{\pi_{\mathbf{n}}(3)} > n_{(3)} - n_{(2)}, \dots, \tau_{\pi_{\mathbf{n}}(d)} > n_{(d)} - n_{(2)}) \\ &= \prod_{k=1}^d \mathbb{P}(\tau_{\pi_{\mathbf{n}}(k)} > 1, \dots, \tau_{\pi_{\mathbf{n}}(d)} > 1)^{n_{(k)} - n_{(k-1)}} = \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_{\mathbf{n}}(i) \notin I \ \forall i=k, \dots, d}} \tilde{p}_I \right)^{n_{(k)} - n_{(k-1)}}, \end{aligned}$$

where the last equality is due to (i). Definition 2.3 together with Theorem 2.5 give the result. \square

Appendix B. Proofs from Section 3

PROOF OF LEMMA 3.1. First, we prove the claim for the survival function in (1). Necessity is shown via induction over the cardinality of subsets of $\{1, \dots, d\}$. To begin with, it is shown that $p_{\{1\}} = \dots = p_{\{d\}}$. Exchangeability implies for each $k = 2, \dots, d$ that

$$\prod_{\substack{\emptyset \neq I \subseteq \{1, \dots, d\} \\ I \neq \{1\}}} p_I = \bar{F}_{0,1,\dots,1} = \bar{F}_{1,\dots,1,0,1,\dots,1} = \prod_{\substack{\emptyset \neq I \subseteq \{1, \dots, d\} \\ I \neq \{k\}}} p_I.$$

Division by $\prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I = \bar{F}_{1, \dots, 1}$ on both sides gives $p_{\{1\}} = p_{\{k\}}$. Concerning the induction step, assume the claim to be true for cardinalities less than or equal to $k < d$ and let $I_0 \subseteq \{1, \dots, d\}$ with $|I_0| = k + 1$. Define

$$n_j := \begin{cases} 1, & j \in I_0 \\ 0, & j \notin I_0 \end{cases}, \quad \tilde{n}_j := \begin{cases} 1, & j \in \{1, \dots, k+1\} \\ 0, & j \notin \{1, \dots, k+1\} \end{cases}.$$

From exchangeability it follows that

$$\prod_{\substack{\emptyset \neq I \subseteq \{1, \dots, d\} \\ I \not\subseteq I_0}} p_I = \bar{F}_{n_1, \dots, n_d} = \bar{F}_{\tilde{n}_1, \dots, \tilde{n}_d} = \prod_{\substack{\emptyset \neq I \subseteq \{1, \dots, d\} \\ I \not\subseteq \{1, \dots, k+1\}}} p_I.$$

Division by $\prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I = \bar{F}_{1, \dots, 1}$ on both sides implies

$$p_{I_0} \underbrace{\prod_{\substack{\emptyset \neq I \subseteq I_0 \\ |I| \leq k}} p_I}_{=:(*)} = p_{\{1, \dots, k+1\}} \underbrace{\prod_{\substack{\emptyset \neq I \subseteq \{1, \dots, k+1\} \\ |I| \leq k}} p_I}_{=:(**)}$$

where $(*) = (**)$ holds by induction hypothesis. Hence, $p_{\{1, \dots, k+1\}} = p_{I_0}$, and the necessity is shown.

To prove sufficiency, assume that the parameters satisfy (7) and let π denote any permutation on $\{1, \dots, d\}$. Then, using the notation $\pi(I) := \{\pi(i) : i \in I\}$,

$$\begin{aligned} \bar{F}_{n_{\pi(1)}, \dots, n_{\pi(d)}} &= \prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{i \in I} \{n_{\pi(i)}\}} = \prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_I^{\max_{i \in \pi(I)} \{n_i\}} \\ &= \prod_{\emptyset \neq I \subseteq \{1, \dots, d\}} p_{\pi(I)}^{\max_{i \in \pi(I)} \{n_i\}} = \bar{F}_{n_1, \dots, n_d}, \end{aligned}$$

where the third equality above follows from the assumption (7).

Now, we establish the claim for the survival function in (2). Necessity is shown, again, via induction over the cardinality of subsets of $\{1, \dots, d\}$. From $\bar{F}_{0, 1, \dots, 1} = \bar{F}_{1, 0, 1, \dots, 1} = \dots = \bar{F}_{1, \dots, 1, 0}$, it follows that $\tilde{p}_{\{1\}} = \dots = \tilde{p}_{\{d\}}$. Assume the claim is true for cardinalities less than or equal to $k < d$. From exchangeability of $\bar{F}_{n_1, \dots, n_d}$, it follows for all permutations π on $\{1, \dots, d\}$ that

$$\begin{aligned} \sum_{\substack{I \subseteq \{1, \dots, d\} \\ i \notin I \forall i = k+2, \dots, d}} \tilde{p}_I &= \mathbb{P}(\tau_1 > 0, \dots, \tau_{k+1} > 0, \tau_{k+2} > 1, \dots, \tau_d > 1) \\ &= \mathbb{P}(\tau_{\pi(1)} > 0, \dots, \tau_{\pi(k+1)} > 0, \tau_{\pi(k+2)} > 1, \dots, \tau_{\pi(d)} > 1) = \sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi(i) \notin I \forall i = k+2, \dots, d}} \tilde{p}_I. \end{aligned}$$

From this, using the induction hypothesis, we infer

$$\begin{aligned}
\tilde{p}_{\{1, \dots, k+1\}} &= \sum_{\substack{I \subseteq \{1, \dots, d\} \\ i \notin I \forall i=k+2, \dots, d}} \tilde{p}_I - \sum_{\substack{I \subseteq \{1, \dots, d\}, |I| \leq k \\ i \notin I \forall i=k+2, \dots, d}} \tilde{p}_I \\
&= \sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi(i) \notin I \forall i=k+2, \dots, d}} \tilde{p}_I - \sum_{\substack{I \subseteq \{1, \dots, d\}, |I| \leq k \\ \pi(i) \notin I \forall i=k+2, \dots, d}} \tilde{p}_I = \tilde{p}_{\{\pi(1), \dots, \pi(k+1)\}}.
\end{aligned}$$

Hence the necessity.

To prove sufficiency, assume that the parameters satisfy (7) and let $\tilde{\pi}$ be any permutation on $\{1, \dots, d\}$. Denoting $\tilde{\pi}(\mathbf{n}) = (n_{\tilde{\pi}(1)}, \dots, n_{\tilde{\pi}(d)})$, and with $\hat{\pi}: \{1, \dots, d\} \rightarrow \{1, \dots, d\}$ being a permutation depending on $\tilde{\pi}(\mathbf{n})$ such that $n_{\hat{\pi}(\tilde{\pi}(1))} \leq n_{\hat{\pi}(\tilde{\pi}(2))} \leq \dots \leq n_{\hat{\pi}(\tilde{\pi}(d))}$, we obtain

$$\begin{aligned}
\bar{F}_{n_{\tilde{\pi}(1)}, \dots, n_{\tilde{\pi}(d)}} &= \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \tilde{\pi}(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I \right)^{n_{\tilde{\pi}(k)} - n_{\tilde{\pi}(k-1)}} \\
&= \prod_{k=1}^d \left(\sum_{\substack{I \subseteq \{1, \dots, d\} \\ \pi_{\mathbf{n}}(i) \notin I \forall i=k, \dots, d}} \tilde{p}_I \right)^{n_{(k)} - n_{(k-1)}} = \bar{F}_{n_1, \dots, n_d},
\end{aligned}$$

where the second equality above follows from assumption (7) together with $n_{\tilde{\pi}(k)} = n_{(k)}$ for all $k = 1, \dots, d$. Hence the lemma. \square

PROOF OF THEOREM 3.5. We first show (\mathcal{N}) . From Theorem 2.5 (\mathcal{N}) together with (9), the d -dimensional non-degenerate exchangeable narrow-sense geometric survival function is given by

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{N}} = \prod_{k=1}^d \left(\prod_{i=1}^{d-k+1} p_i^{\binom{d-k}{i-1}} \right)^{n_{(d-k+1)}} = \prod_{k=1}^d a_k^{n_{(d-k+1)}},$$

with $p_k \in (0, 1]$ for all $k = 1, \dots, d$, $\prod_{i=1}^d p_i < 1$, and $\{a_k\}_{k=1}^d$ defined as in (8). The formula for p_k , $k = 1, \dots, d$, is justified by

$$\begin{aligned}
\prod_{i=1}^k a_{d-i+1}^{(-1)^{(k-i)} \binom{k-1}{i-1}} &= \prod_{i=1}^k \left(\prod_{j=1}^i p_j^{\binom{i-1}{j-1}} \right)^{(-1)^{(k-i)} \binom{k-1}{i-1}} = \prod_{i=1}^k \prod_{j=1}^i p_j^{(-1)^{(k-i)} \binom{k-j}{i-j} \binom{k-1}{j-1}} \\
&= \prod_{j=1}^k p_j^{(-1)^{(k)} \binom{k-1}{j-1} \sum_{i=j}^k (-1)^{(i)} \binom{k-j}{i-j}} = \prod_{j=1}^k p_j^{(-1)^{(k+j)} \binom{k-1}{j-1} \mathbf{1}_{\{k=j\}}} = p_k,
\end{aligned}$$

where the fourth equality follows from

$$\sum_{i=j}^k (-1)^{(i)} \binom{k-j}{i-j} = \sum_{l=0}^{k-j} (-1)^{(l+j)} \binom{k-j}{l} = (-1)^{(j)} \mathbf{1}_{\{k=j\}}.$$

Next, we show that $p_k \in (0, 1]$ for all $k = 1, \dots, d$ and $\prod_{i=1}^d p_i < 1$ if and only if $\{a_k\}_{k=1}^d \in \mathcal{SM}_d$. To this end, observe that for $k = 0, 1, \dots, d-1$,

$$\begin{aligned} \nabla^{k-1}(\ln a_{d-k+1}^{-1}) &= \sum_{i=0}^{k-1} (-1)^i \binom{k-1}{i} (-\ln a_{d-k+i+1}) \\ &= \sum_{i=0}^{k-1} (-1)^{k-i-1} \binom{k-1}{i} (-\ln a_{d-i}) = \sum_{i=1}^k (-1)^{k-i} \binom{k-1}{i-1} (-\ln a_{d-i+1}) \\ &= -\ln \left(\prod_{i=1}^k a_{d-i+1}^{(-1)^{(k-i)} \binom{k-1}{i-1}} \right) = -\ln p_k, \end{aligned}$$

and $\nabla^{k-1}(\ln a_{d-k+1}^{-1}) \geq 0$ for all $k=1, \dots, d$ is equivalent to $\nabla^{d-k-1}(-\ln a_{k+1}) \geq 0$ for all $k = 0, 1, \dots, d-1$. Taking into account that

$$a_1 = \prod_{i=1}^d p_i^{\binom{d-1}{i-1}},$$

it follows that $p_k \in (0, 1]$ for all $k = 1, \dots, d$ and $\prod_{i=1}^d p_i < 1$ if and only if $\{a_k\}_{k=1}^d \in \mathcal{SM}_d$. Furthermore, Theorem 2.5 (\mathcal{N}) together with (8) and (9) imply that from any finite sequence $\{a_k\}_{k=1}^d \in \mathcal{SM}_d$ one can construct an exchangeable d -variate narrow-sense geometric distribution by choosing the parameters p_1, \dots, p_d as

$$p_k := \prod_{i=1}^k a_{d-i+1}^{(-1)^{(k-i)} \binom{k-1}{i-1}}, \quad k = 1, \dots, d.$$

Finally, given $\{b_k\}_{k=1}^d$ as in (10), and denoting $b_0 := 1$, observe that

$$\nabla^{d-k} \ln a_k^{-1} = \nabla^{d-k} \ln b_{k-1} - \nabla^{d-k} \ln b_k = \nabla^{d-k+1} \ln b_{k-1}$$

for all $k = 1, \dots, d$. From this, we conclude that $\{b_k\}_{k=0}^d \in \mathcal{LM}_{d+1}$ if and only if $\{a_k\}_{k=1}^d \in \mathcal{SM}_d$. With (11), the assertion follows.

Next, we prove (\mathcal{W}). From Theorem 2.5 (\mathcal{W}) together with (12) and (13), the d -dimensional exchangeable wide-sense geometric survival function is given by

$$\bar{F}_{n_1, \dots, n_d}^{\mathcal{W}} = \prod_{k=1}^d \left(\sum_{i=1}^{d-k+1} \binom{d-k}{i-1} \tilde{p}_i \right)^{n_{(d-k+1)} - n_{(d-k)}} = \prod_{k=1}^d \beta_k^{n_{(d-k+1)} - n_{(d-k)}},$$

with $\tilde{p}_1, \dots, \tilde{p}_d \in [0, 1]$, $\sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \leq 1$, $\sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i < 1$, and $\{\beta_k\}_{k=1}^d$

defined as in (13). The formula for \tilde{p}_k , $k = 1, \dots, d$, is justified by

$$\begin{aligned}
\nabla^{k-1}\beta_{d-k+1} &= \sum_{i=0}^{k-1} (-1)^i \binom{k-1}{i} \beta_{d-k+i+1} = \sum_{i=0}^{k-1} (-1)^i \binom{k-1}{i} \sum_{j=1}^{k-i} \binom{k-i-1}{j-1} \tilde{p}_j \\
&= \sum_{j=1}^k \sum_{i=0}^{k-j} (-1)^i \binom{k-1}{i} \binom{k-i-1}{j-1} \tilde{p}_j = \sum_{j=1}^k \binom{k-1}{j-1} \tilde{p}_j \sum_{i=0}^{k-j} (-1)^i \binom{k-j}{i} \\
&= \sum_{j=1}^k \binom{k-1}{j-1} \tilde{p}_j \mathbf{1}_{\{k=j\}} = \tilde{p}_k.
\end{aligned}$$

Now, we infer $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ from $\tilde{p}_1, \dots, \tilde{p}_d \in [0, 1]$ with $\sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \leq 1$ and $\sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i < 1$, and vice versa. For $k = 1, \dots, d$, it holds that $\nabla^{d-k} \beta_k = \tilde{p}_{d-k+1}$. Further,

$$\begin{aligned}
\nabla^d \beta_0 &= \sum_{j=0}^d (-1)^j \binom{d}{j} \beta_j = \beta_0 + \sum_{j=1}^d (-1)^j \binom{d}{j} \beta_j = 1 + \sum_{j=1}^d (-1)^j \binom{d}{j} \sum_{i=1}^{d-j+1} \binom{d-j}{i-1} \tilde{p}_i \\
&= 1 + \sum_{i=1}^d \tilde{p}_i \sum_{j=1}^{d-i+1} (-1)^j \binom{d}{j} \binom{d-j}{i-1} = 1 + \sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \sum_{j=1}^{d-i+1} (-1)^j \binom{d-i+1}{j} \\
&= 1 + \sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i \left(\sum_{j=0}^{d-i+1} (-1)^j \binom{d-i+1}{j} - 1 \right) = 1 - \sum_{i=1}^d \binom{d}{i-1} \tilde{p}_i.
\end{aligned}$$

Lastly, $\beta_1 = \sum_{i=1}^d \binom{d-1}{i-1} \tilde{p}_i$, and, therefore, $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ gives precisely the conditions on $\tilde{p}_1, \dots, \tilde{p}_d$ as stated in the assertion. Furthermore, Theorem 2.5 (\mathcal{N}) together with (12) and (13) imply that from any finite sequence $\{\beta_k\}_{k=1}^d$ with $(1, \beta_1, \dots, \beta_d) \in \mathcal{M}_{d+1}$ one can construct an exchangeable d -dimensional wide-sense geometric distribution by choosing the parameters $\tilde{p}_1, \dots, \tilde{p}_d$ as

$$\tilde{p}_k = \nabla^{k-1} \beta_{d-k+1}, \quad k = 1, \dots, d.$$

This completes the proof. \square

PROOF OF THEOREM 3.7. Sufficiency follows directly from Theorem 3.5 (\mathcal{W}). To prove necessity, suppose that (9) defines a d -dimensional survival function. As is easily seen, it is exchangeable and satisfies the local discrete LM property (3). Hence, in view of Theorem 2.6, it is the survival function of a d -variate exchangeable wide-sense geometric distribution. The claim follows now from Theorem 3.5 (\mathcal{W}). \square

Appendix C. Proofs from Section 4

PROOF OF PROPOSITION 4.4. That (i) implies (iv) follows directly from Corollary 4.1, recalling that $\mathcal{G}^{\mathcal{N}, \mathcal{E}} \subseteq \mathcal{G}^{\mathcal{W}, \mathcal{E}}$, together with the fact that \mathcal{LM}_∞ is closed

under positive powers. The reverse implication is easy: since $\{b_k\}_{k \in \mathbb{N}_0}$ is strictly positive, it follows for all $k \in \mathbb{N}_0$ and all $j \in \mathbb{N}$ that

$$\nabla^j \ln b_k = \lim_{n \rightarrow \infty} n \nabla^j (\sqrt[n]{b_k} - 1) = \lim_{n \rightarrow \infty} n \nabla^j \sqrt[n]{b_k} \geq 0.$$

The equivalence of (iii) and (iv) is obtained from the closure of \mathcal{M}_∞ under limits and products. To see why (ii) implies (iv), note that from the infinite divisibility of κ , we have for all $n \in \mathbb{N}$ that

$$\sqrt[n]{b_k} = \int_{[0, \infty]} e^{-kx} \kappa_n(dx), \quad k \in \mathbb{N}_0, \quad (\text{C.1})$$

where κ_n is such that κ is the n -fold convolution on κ_n with itself. One verifies using a simple induction that for all $n, j \in \mathbb{N}$ and for all $k \in \mathbb{N}_0$,

$$\nabla^j \sqrt[n]{b_k} = \nabla^{j-1} \sqrt[n]{b_k} - \nabla^{j-1} \sqrt[n]{b_{k+1}} = \int_{[0, \infty]} e^{-kx} (1 - e^{-x})^j \kappa_n(dx) \geq 0,$$

implying that $\{\sqrt[n]{b_k}\}_{k \in \mathbb{N}_0}$ is completely monotone for all $n \in \mathbb{N}$. Moreover, $b_0 = 1$ since κ is a probability measure, $b_1 < 1$ because κ is not concentrated at 0, and $b_k > 0$ for all $k \in \mathbb{N}_0$ as κ is not concentrated at ∞ . Hence, $\{\sqrt[n]{b_k}\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$ for all $n \in \mathbb{N}$. Finally, to show that (iv) implies (ii), it is enough to notice that, in view of Hausdorff's moment problem [16], $\{\sqrt[n]{b_k}\}_{k \in \mathbb{N}_0}$ can be represented as the moment sequence (C.1) for some unique probability measure κ_n on $[0, \infty]$ for all $n \in \mathbb{N}$. Therefore, for all $n \in \mathbb{N}$, κ_1 is the n -fold convolution of κ_n with itself, meaning that κ_1 is infinitely divisible. And $b_1 \in (0, 1)$ yields $\kappa(\{0\}) < 1$ and $\kappa(\{\infty\}) < 1$. \square

PROOF OF THEOREM 4.6. We first show that the model in (\mathcal{W}) generates precisely the family of $\mathcal{G}^{\mathcal{W}, \mathcal{E}}$ -distributions. By Corollary 4.1 (\mathcal{W}), every $\mathcal{G}^{\mathcal{W}, \mathcal{E}}$ -distribution is characterized by an infinite sequence $(1, \beta_1, \beta_2, \dots) \in \mathcal{M}_\infty$. From the given sequence $(1, \beta_1, \beta_2, \dots) \in \mathcal{M}_\infty$, we construct an infinite exchangeable sequence $\{\tau_k\}_{k \in \mathbb{N}}$ of random variables such that (τ_1, \dots, τ_d) has the survival function (12) for each $d \geq 2$. To this end, denote $\beta_0 := 1$ and assume that $\{\beta_k\}_{k \in \mathbb{N}_0} \in \mathcal{M}_\infty$. By virtue of Hausdorff's moment problem, see [15, 16], there exists a unique probability law η on the unit interval $[0, 1]$ such that

$$\beta_k = \int_{[0, 1]} x^k \eta(dx), \quad k \in \mathbb{N}_0.$$

Now consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ supporting the following independent sequences of random variables:

- An i.i.d. sequence $\{Y_k\}_{k \in \mathbb{N}}$, where $Y_1 \sim \eta$.
- An i.i.d. sequence $\{E_k\}_{k \in \mathbb{N}}$, where $E_1 \sim \text{Exp}(1)$.

Define the i.i.d. sequence $\{X_k\}_{k \in \mathbb{N}}$ by $X_k := -\ln(Y_k)$, $k \in \mathbb{N}$. Notice that it may hold that $\mathbb{P}(Y_k = 0) > 0$, in which case we have $\mathbb{P}(X_k = \infty) > 0$ by conveniently defining $\ln 0 := -\infty$. Define the sequence $\{\tau_k\}_{k \in \mathbb{N}}$ by

$$\tau_k := \min\{n \in \mathbb{N} : X_1 + \dots + X_n \geq E_k\}, \quad k \in \mathbb{N}.$$

Notice that $\mathbb{P}(X_k = 0) < 1$ by the assumption $\beta_1 < 1$, implying that the τ_k are well-defined in \mathbb{N} . We compute for $n_1, \dots, n_d \in \mathbb{N}_0$ that

$$\begin{aligned} \mathbb{P}(\tau_1 > n_1, \dots, \tau_d > n_d) &= \mathbb{P}(X_1 + \dots + X_{n_1} < E_1, \dots, X_1 + \dots + X_{n_d} < E_d) \\ &= \mathbb{E}\left[\mathbb{P}(X_1 + \dots + X_{n_1} < E_1, \dots, X_1 + \dots + X_{n_d} < E_d \mid X_1, \dots, X_{n_d})\right] \\ &= \mathbb{E}\left[\prod_{k=1}^d \mathbb{P}(X_1 + \dots + X_{n_k} < E_k \mid X_1, \dots, X_{n_d})\right] = \mathbb{E}\left[\prod_{k=1}^d e^{-(X_1 + \dots + X_{n_k})}\right] \\ &= \mathbb{E}\left[e^{-d(X_1 + \dots + X_{n_1})} e^{-(d-1)(X_{n_1+1} + \dots + X_{n_2})} \dots e^{-(X_{n_{d-1}+1} + \dots + X_{n_d})}\right] \\ &= \prod_{k=1}^d \mathbb{E}\left[e^{-(d-k+1)(X_{n_{k-1}+1} + \dots + X_{n_k})}\right] = \prod_{k=1}^d \mathbb{E}\left[(e^{-X_1})^{(d-k+1)}\right]^{n_k - n_{k-1}} \\ &= \prod_{k=1}^d \mathbb{E}\left[(e^{-X_1})^k\right]^{n(d-k+1) - n(d-k)} = \prod_{k=1}^d \mathbb{E}[Y_1^k]^{n(d-k+1) - n(d-k)} = \prod_{k=1}^d \beta_k^{n(d-k+1) - n(d-k)}. \end{aligned}$$

Finally, by Corollary 4.1 (\mathcal{N}), the distribution of (τ_1, \dots, τ_d) is geometric in the narrow sense if and only if $(1, \beta_1, \beta_2, \dots) \in \mathcal{LM}_\infty$. In view of Proposition 4.4, it is the case precisely when the distribution of X_1 is infinitely divisible. \square

Appendix D. Proofs from Section 5

PROOF OF THEOREM 5.7. Let $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$, given by

$$\bar{F}_{n_1, \dots, n_d} = \prod_{k=1}^d a_k^{n(d-k+1)} = \prod_{k=1}^d b_k^{n(d-k+1) - n(d-k)},$$

be a d -dimensional survival function, where $b_0 := 1$ and $\{b_k\}_{k=1}^d$ is as in (10). We show that $\{\bar{F}_{n_1, \dots, n_d}\}_{n_1, \dots, n_d \in \mathbb{N}_0}$ is MRTI if and only if $\frac{a_k}{a_{k+1}} = \frac{b_k^2}{b_{k-1}b_{k+1}} \leq 1$ $\forall k = 1, \dots, d-1$.

$$\mathbb{P}(\tau_k > n_k \mid \tau_1 > n_1, \dots, \tau_{k-1} > n_{k-1}) \geq \mathbb{P}(\tau_k > n_k \mid \tau_1 > \tilde{n}_1, \dots, \tau_{k-1} > \tilde{n}_{k-1}) \quad (\text{D.1})$$

holds for all $n_i, \tilde{n}_i \in \mathbb{N}_0$ with $n_i \geq \tilde{n}_i$, $i = 1, \dots, k-1$, and all $k = 2, \dots, d$ if and only if $a_k/a_{k+1} \leq 1$. Fix an arbitrary $k \in \{2, \dots, d\}$. Define the sequence $\{T_i\}_{i=1}^{k-1}$ recursively by

$$\begin{aligned} T_i &:= T_{i-1} + \text{sort}(n_1, \dots, n_{k-1})[i] - \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[i] \\ &\quad + \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[i] - \text{sort}(n_1, \dots, n_k)[i], \end{aligned}$$

where $T_0 := 0$, and for $i \in \mathbb{N}$, $i \leq k$, $sort(n_1, \dots, n_k)[i]$ denotes the i -th reverse order statistic of the set $\{n_1, \dots, n_k\}$, so that $sort(n_1, \dots, n_k)[1] \geq sort(n_1, \dots, n_k)[2] \geq \dots \geq sort(n_1, \dots, n_k)[k]$. Observe that

$$\begin{aligned}
T_{k-1} &= \sum_{i=1}^{k-1} sort(n_1, \dots, n_{k-1})[i] - \sum_{i=1}^{k-1} sort(\tilde{n}_1, \dots, \tilde{n}_{k-1})[i] \\
&\quad + \sum_{i=1}^{k-1} sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[i] - \sum_{i=1}^{k-1} sort(n_1, \dots, n_k)[i] \\
&= \sum_{i=1}^{k-1} n_i + \sum_{i=1}^{k-1} \tilde{n}_i + n_k - \sum_{i=1}^k n_i - \sum_{i=1}^{k-1} \tilde{n}_i \\
&\quad + sort(n_1, \dots, n_k)[k] - sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[k] \\
&= sort(n_1, \dots, n_k)[k] - sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[k].
\end{aligned}$$

This allows us to write the ratio of conditional probabilities in (D.1) as

$$\begin{aligned}
\frac{\mathbb{P}(\tau_k > n_k \mid \tau_1 > n_1, \dots, \tau_{k-1} > n_{k-1})}{\mathbb{P}(\tau_k > n_k \mid \tau_1 > \tilde{n}_1, \dots, \tau_{k-1} > \tilde{n}_{k-1})} &= \frac{\prod_{i=1}^k a_i^{sort(n_1, \dots, n_k)[i]} \prod_{i=1}^{k-1} a_i^{sort(\tilde{n}_1, \dots, \tilde{n}_{k-1})[i]}}{\prod_{i=1}^{k-1} a_i^{sort(n_1, \dots, n_{k-1})[i]} \prod_{i=1}^k a_i^{sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[i]}} \\
&= a_k^{sort(n_1, \dots, n_k)[k] - sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[k]} \prod_{i=1}^{k-1} a_i^{T_{i-1} - T_i} = \left(\frac{a_2}{a_1}\right)^{T_1} \left(\frac{a_3}{a_2}\right)^{T_2} \dots \left(\frac{a_k}{a_{k-1}}\right)^{T_{k-1}}.
\end{aligned}$$

It can be easily seen that the terms T_1 and T_{k-1} are non-negative. We show that actually all elements of $\{T_i\}_{i=1}^{k-1}$ are non-negative, which results in

$$\mathbb{P}(\tau_k > n_k \mid \tau_1 > n_1, \dots, \tau_{k-1} > n_{k-1}) \geq \mathbb{P}(\tau_k > n_k \mid \tau_1 > \tilde{n}_1, \dots, \tau_{k-1} > \tilde{n}_{k-1})$$

if and only if $a_{i+1}/a_i \geq 1$, $i = 1, \dots, k-1$, as claimed. The equivalence holds, since for each $i \in \{2, \dots, k\}$,

$$\frac{\mathbb{P}(\tau_k > n_k \mid \tau_1 > n_1, \dots, \tau_{k-1} > n_{k-1})}{\mathbb{P}(\tau_k > n_k \mid \tau_1 > \tilde{n}_1, \dots, \tau_{k-1} > \tilde{n}_{k-1})} = \frac{a_i}{a_{i-1}}$$

by choosing $n_j = \tilde{n}_j = 2$ for all $j = 1, \dots, i-2$, $n_j = \tilde{n}_j = 0$ for all $j = i, \dots, k-1$, $n_{i-1} = n_k = 1$, and $\tilde{n}_{i-1} = 0$.

To complete the proof, we show that $\{T_i\}_{i=1}^{k-1}$ is non-negative. For this, order the sets $\{n_1, \dots, n_k\}$ and $\{\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k\}$. Then, there exist $j, l \in \{1, \dots, k\}$, $j \leq l$, such that

$$\begin{aligned}
n_k &= sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[j], \\
n_k &= sort(n_1, \dots, n_k)[l].
\end{aligned}$$

Notice that

$$\begin{aligned}
sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[m] &= sort(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m] && \forall m < j, \\
sort(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[m] &= sort(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m-1] && \forall m > j, \\
sort(n_1, \dots, n_k)[m] &= sort(n_1, \dots, n_{k-1})[m] && \forall m < l, \\
sort(n_1, \dots, n_k)[m] &= sort(n_1, \dots, n_{k-1})[m-1] && \forall m > l.
\end{aligned}$$

Hence, $T_i = 0$ for all $i < j$. For $i \geq l$, we compute

$$\begin{aligned}
T_i &= \sum_{m=1}^i \text{sort}(n_1, \dots, n_{k-1})[m] - \sum_{m=1}^i \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m] \\
&\quad + \sum_{m=1}^i \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1}, n_k)[m] - \sum_{m=1}^i \text{sort}(n_1, \dots, n_k)[m] \\
&= \sum_{m=1}^i \text{sort}(n_1, \dots, n_{k-1})[m] - \sum_{m=1}^i \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m] \\
&\quad + \sum_{m=1}^{j-1} \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m] + n_k + \sum_{m=j+1}^i \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[m-1] \\
&\quad - \sum_{m=1}^{l-1} \text{sort}(n_1, \dots, n_{k-1})[m] - n_k - \sum_{m=l+1}^i \text{sort}(n_1, \dots, n_{k-1})[m-1] \\
&= \text{sort}(n_1, \dots, n_{k-1})[i] - \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[i] \geq 0.
\end{aligned}$$

Finally, if $j < l$, then, arguing analogously, we obtain for $j \leq i < l$,

$$T_i = n_k - \text{sort}(\tilde{n}_1, \dots, \tilde{n}_{k-1})[i] \geq 0.$$

This verifies (W). (\mathcal{N}) follows by noticing that $\{a_k\}_{k=1}^d$ is non-decreasing for any set of parameters $\{a_k\}_{k=1}^d$ of a d -variate $\mathcal{G}^{\mathcal{N}, \mathcal{X}}$ -distribution. \square

References

- [1] ALDOUS D.J. (1985). Exchangeability and related topics, *Springer, École d'Été de Probabilités de Saint-Flour XIII-1983. Lecture Notes in Mathematics* **1117** pp. 1–198.
- [2] ARNOLD B.C. (1975). A characterization of the exponential distribution by multivariate geometric compounding, *Sankhyā: The Indian Journal of Statistics* **37:1** pp. 164–173.
- [3] AZLAROV T.A. and VOLODIN N.A. (1983). On the discrete analog of Marshall-Olkin's distribution, *Stability problems for stochastic models. Lecture Notes in Mathematics* **982** pp. 17–23.
- [4] BERG C. (2007). On powers of Stieltjes moment sequences, II, *Journal of Computational and Applied Mathematics* **199** pp. 23–38.
- [5] BERG C. (2008). Stieltjes-Pick-Bernstein-Schoenberg and their connection to complete monotonicity, *Positive Definite Functions. From Schoenberg to Space-Time Challenges*.
- [6] BLOCK H.W. and TING M.L. (1981). Some concepts of multivariate dependence, *Communications in Statistics A - Theory and Methods* **10** pp. 749–762.

- [7] COLANGELO A., SCARSINI M. and SHAKED M. (2005). Some notions of multivariate positive dependence, *Insurance: Mathematics and Economics* **37** pp. 13–26.
- [8] DE FINETTI B. (1937). La prévision: ses lois logiques, ses sources subjectives, *Annales de l'Institut Henri Poincaré* **7** pp. 1–68.
- [9] DETTE H. and STUDDEN W.J. (1997). The theory of canonical moments with applications in statistics, probability and analysis, *Wiley Series in Probability and Mathematical Statistics*.
- [10] ESARY, J.D. and MARSHALL, A.W. (1973). Multivariate geometric distributions generated by a cumulative damage process, *Naval Postgraduate School, Monterey, California*, <http://calhoun.nps.edu/handle/10945/30003>.
- [11] FANG K.T., KOTZ S. and NG K.W. (1990). *Symmetric Multivariate and Related Distributions*, Chapman and Hall, London.
- [12] GNEDIN A.V. (2004). The Bernoulli sieve, *Bernoulli* **10:1** pp. 79–96.
- [13] GNEDIN A.V. and PITMAN J. (2005). Regenerative composition structures, *Annals of Probability* **33:2** pp. 445–479.
- [14] GNEDIN A.V. and PITMAN J. (2008). Moments of convex distribution functions and completely alternating sequences, *Probability and Statistics: Essays in Honor of David A. Freedman* **2** pp. 30–41.
- [15] HAUSDORFF F. (1921). Summationsmethoden und Momentfolgen I, *Mathematische Zeitschrift* **9:3–4** pp. 74–109.
- [16] HAUSDORFF F. (1923). Momentenproblem für ein endliches Intervall, *Mathematische Zeitschrift* **16** pp. 220–248.
- [17] KARLIN S. and SHAPLEY L.S. (1953). Geometry of moment spaces, *Memoirs of the American Mathematical Society* **12**.
- [18] KIMBERLING C.H. (1974). A probabilistic interpretation of complete monotonicity, *Aequationes Mathematicae* **10** pp. 152–164.
- [19] KINGMAN J.F.C. (1972). On random sequences with spherical symmetry, *Biometrika* **59** pp. 492–494.
- [20] KINGMAN J.F.C. (1978). Uses of exchangeability, *Annals of Probability* **6:2** pp. 183–197.
- [21] LORCH L. and NEWMAN D.J. (1983). On the composition of completely monotonic functions and completely monotonic sequences and related questions, *Journal of the London Mathematical Society* **2:28–1** pp. 31–45.

- [22] MAI J.F. and SCHERER M. (2009). Lévy-frailty Copulas, *Journal of Multivariate Analysis* **100:7** pp. 1567–1585.
- [23] MAI J.F. and SCHERER M. (2011). Reparameterizing Marshall–Olkin copulas with applications to sampling, *Journal of Statistical Computation and Simulation* **81:1** pp. 59–78.
- [24] MALOV S.V. (2001). On finite-dimensional Archimedean copulas, *Asymptotic Methods in Probability and Statistics with Applications* pp. 19–35.
- [25] MARSHALL A.W. and OLKIN I. (1967). A multivariate exponential distribution, *Journal of the American Statistical Association* **62:317** pp. 30–44.
- [26] MARSHALL A.W. and OLKIN I. (1995). Multivariate exponential and geometric distributions with limited memory, *Journal of Multivariate Analysis* **53** pp. 110–125.
- [27] MCNEIL A.J. and NEŠLEHOVÁ J. (2009). Multivariate Archimedean copulas, d -monotone functions and l_1 -norm symmetric distributions, *Annals of Statistics* **37(5B)** pp. 3059–3097.
- [28] MORILLAS P.M. (2005). A characterization of absolutely monotonic (Δ) functions of a fixed order, *Publications de l'Institut Mathématique, Nouvelle série* **78** pp. 93–105.
- [29] NAIR N.U. and ASHA G. (1997). Some classes of multivariate life distributions in discrete time, *Journal of Multivariate Analysis* **62** pp. 181–189.
- [30] RESSEL P. (2011). Monotonicity properties of multivariate distribution and survival functions with an application to Lévy-frailty copulas, *Journal of Multivariate Analysis* **102(3)** pp. 393–404.
- [31] RESSEL P. (2012). Finite exchangeability, Lévy-frailty copulas and higher-order monotonic sequences, to appear in *Journal of Theoretical Probability*.
- [32] SATO K.I. (1999). Lévy processes and infinitely divisible distributions, *Cambridge University Press*.
- [33] SCHILLING R.L., SONG R. and VONDRACEK Z. (2010). *Bernstein Functions*, De Gruyter, Berlin.
- [34] SCHOENBERG I.J. (1932). On finite and infinite completely monotonic sequences, *Bulletin of the American Mathematical Society* **38:2** pp. 72–76.
- [35] SCHOENBERG I.J. (1938). Metric spaces and positive definite functions, *Transactions of the American Mathematical Society* **44** pp. 522–536.
- [36] SHENKMAN N. (2017). A natural parametrization of multivariate distributions with limited memory, *Journal of Multivariate Analysis* **155** pp. 234–251.

- [37] SHENKMAN N. (2020). A primer on the characterization of the exchangeable Marshall–Olkin copula via monotone sequences, *forthcoming in Brazilian Journal of Probability and Statistics*.
- [38] TONG Y.L. (1994). Some useful notions for studying stochastic inequalities in multivariate distributions, *Multivariate Analysis and Its Applications IMS Lecture Notes* **24** pp. 55–71.