

ON THE SUPERSINGULAR LOCUS OF THE SHIMURA VARIETY FOR $\mathrm{GU}(2, 2)$ OVER A RAMIFIED PRIME

YASUHIRO OKI

ABSTRACT. We study the structure of the supersingular locus of the Rapoport–Zink integral model of the Shimura variety for $\mathrm{GU}(2, 2)$ over a ramified odd prime with the special maximal parahoric level. We prove that the supersingular locus equals the disjoint union of two basic loci, one of which is contained in the flat locus, and the other is not. We also describe explicitly the structure of the basic loci. More precisely, the former one is purely 2-dimensional, and each irreducible component is birational to the Fermat surface. On the other hand, the latter one is purely 1-dimensional, and each irreducible component is birational to the projective line.

CONTENTS

1. Introduction	2
1.1. Main results	3
1.2. Results on the Rapoport–Zink spaces	4
2. Rapoport–Zink spaces for a ramified $\mathrm{GU}(2, 2)$	7
2.1. Local PEL datum	7
2.2. p -divisible groups and isocrystals with additional structures	8
2.3. Definition of the Rapoport–Zink spaces	9
2.4. The neutralness condition	10
3. Singularities of the Rapoport–Zink spaces	12
4. Deligne–Lusztig varieties for even special orthogonal groups	16
5. Bruhat–Tits stratification in the neutral case, I	20
5.1. Preliminaries	20
5.2. Hodge star operator	21
5.3. Exceptional isomorphism	23
5.4. Description of a subset of field valued points	24
5.5. Rapoport–Zink space for the quaternionic unitary group of degree 2	27
5.6. Vertex lattices, I	30
5.7. Connected components	32
5.8. Bruhat–Tits stratification, I	33
6. Bruhat–Tits stratification in the neutral case, II	35
6.1. Vertex lattices, II	35
6.2. Bruhat–Tits stratification, II	36
7. Bruhat–Tits stratification in the non-neutral case	37
8. Supersingular loci of unitary Shimura varieties	40
8.1. Rapoport–Zink integral models of unitary Shimura varieties	40
8.2. Non-emptiness of the basic Newton strata	41

2010 *Mathematics Subject Classification.* 11G18, 14G35.

Key words and phrases. Unitary Shimura varieties, Rapoport–Zink spaces, supersingular loci.

8.3. Structure of the supersingular loci

43

References

46

1. INTRODUCTION

This paper contributes to the study of bad reductions of integral models of unitary Shimura varieties. More precisely, we study the geometric structure of an important part, which is called the supersingular locus, of the geometric special fibers of the Rapoport–Zink integral models of Shimura varieties for some unitary groups over certain ramified primes.

Before describing our results, we recall notions of integral models and supersingular loci of Shimura varieties for unitary similitude groups. Let $L = \mathbb{Q}(\sqrt{\Delta})$ an imaginary quadratic field, where $\Delta \in \mathbb{Z}_{<0}$ is square-free. Fix an isomorphism $L \otimes_{\mathbb{Q}} \mathbb{R} \cong \mathbb{C}$, and let U be an L/\mathbb{Q} -hermitian space of signature (r, s) . Put $\mathbf{G} := \mathrm{GU}(U)$, the unitary similitude group of U , and denote by \mathbf{X} the Grassmanian of negative definite s -dimensional subspaces of $U \otimes_{\mathbb{Q}} \mathbb{R}$. Then the pair (\mathbf{G}, \mathbf{X}) is a Shimura datum. Hence, for a sufficiently small compact open subgroup K of $\mathbf{G}(\mathbb{A}_f)$ where \mathbb{A}_f^p is the finite adèle ring of \mathbb{Q} , we obtain the (canonical model of) Shimura variety $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X})$ over $E_{r,s}$, where

$$E_{r,s} := \begin{cases} \mathbb{Q} & \text{if } r = s, \\ L & \text{otherwise.} \end{cases}$$

It is known that $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X})$ is of dimension rs .

Next, we consider an integral model of $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X})$. Let ν be a place in $E_{r,s}$, and p a prime which lies below ν . We assume $p \neq 2$ if p does not split in L . Moreover, we assume that $K = K^p K_p$, where K^p is a sufficiently small compact open subgroup of $\mathbf{G}(\mathbb{A}_f^p)$, and K_p is a Bruhat–Tits subgroup, that is, a subgroup of $\mathbf{G}(\mathbb{Q}_p)$ which is a stabilizer of a self-dual lattice chain in $U \otimes_{\mathbb{Q}} \mathbb{Q}_p$ in the sense of [RZ96, Definition 3.13]. Here \mathbb{A}_f^p is the finite adèle ring of \mathbb{Q} without p -component. Then, Rapoport and Zink constructed in [RZ96] a scheme $\mathcal{S}_{K,U}$ over the ring of integers $O_{E_{r,s,\nu}}$ of $E_{r,s,\nu}$, which is defined as the moduli space of (families of) abelian varieties and additional structures. Here $E_{r,s,\nu}$ is the completion of E at ν . It is known that the generic fiber of $\mathcal{S}_{K,U}$ is isomorphic to $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X}) \otimes_{E_{r,s}} E_{r,s,\nu}$. We define the supersingular locus $\mathcal{S}_{K,U}^{\mathrm{ss}}$ of $\mathcal{S}_{K,U}$ as the locus of the geometric special fiber of $\mathcal{S}_{K,U}$ where the corresponding abelian varieties are supersingular.

It is known that studying the structure of supersingular loci is important for number theory. For example, it can be applied to study some relations between intersection multiplicities of special divisors of Shimura varieties and the first derivatives of Eisenstein series. If $\mathbf{G} = \mathrm{GU}(1, n-1)$, then it is conjectured by [KR14, Conjecture 14.5.1], and recently solved in [LZ19].

We recall some results on the descriptions of $\mathcal{S}_{K,U}^{\mathrm{ss}}$. First, suppose that p splits in L . It is known that $\mathcal{S}_{K,U}^{\mathrm{ss}}$ is non-empty only if $r = s$. If $(r, s) = (1, 1)$ and K_p is hyperspecial, then $\mathcal{S}_{K,U}^{\mathrm{ss}}$ is a finite set of $\overline{\mathbb{F}}_p$ -valued points. On the other hand, Fox [Fox18] (for $p \neq 2$) and Wang [Wan21] (for any p) gave explicit descriptions of $\mathcal{S}_{K,U}^{\mathrm{ss}}$ for $(r, s) = (2, 2)$ and K_p is hyperspecial. In this case, any irreducible component is birational to \mathbb{P}^1 .

Second, consider the case that p inerts in L . For $(r, s) = (1, n-1)$ and K_p is hyperspecial, the structure of $\mathcal{S}_{K,U}^{\mathrm{ss}}$ is given by Vollaard [Vol10] (for $n = 3$) and by Vollaard and Wedhorn [VW11] (for general n). Moreover, Cho [Cho18] analyzed in the case $(r, s) = (1, n-1)$ and K_p is a parahoric subgroup which is the stabilizer of a single lattice in $U \otimes_{\mathbb{Q}} \mathbb{Q}_p$. On the other hand, Howard and

Pappas [HP14] described concretely a structure of $\mathcal{S}_{K,U}^{\mathrm{ss}}$ when $(r, s) = (2, 2)$ and K is hyperspecial. In this case, every irreducible component is birational to the Fermat surface.

Third, we assume that p ramifies in L . In this case, $\mathcal{S}_{K,U}$ may not be flat over $O_{E_{r,s,\nu}}$. Let $\mathcal{S}_{K,V}^{\mathrm{loc}}$ be the scheme-theoretic closure of the generic fiber of $\mathcal{S}_{K,V}$. When $(r, s) = (1, n-1)$ and K_p is the stabilizer of a self-dual lattice, then Rapoport, Terstiege and Wilson [RTW14] considered a description of the structure of $\mathcal{S}_{K,U}^{\mathrm{ss}} \cap \mathcal{S}_{K,U}^{\mathrm{loc}}$. Note that K is not parahoric if n is even. On the other hand, Wu [Wu16] studied $\mathcal{S}_{K,U}^{\mathrm{ss}} \cap \mathcal{S}_{K,U}^{\mathrm{loc}}$ when $(r, s) = (1, n-1)$ and K_p is given by the stabilizer of a single lattice Λ in $V \otimes_{\mathbb{Q}} \mathbb{Q}_p$ satisfying $\Lambda^{\vee} = \sqrt{\Delta}^{-1} \Lambda$ if n is even, or $\Lambda \subset \Lambda^{\vee} \subset \sqrt{\Delta}^{-1} \Lambda$ and $\mathrm{length}(\Lambda^{\vee}/\Lambda) = n-1$ if n is odd. Here Λ^{\vee} is the dual lattice of Λ with respect to the hermitian form on $V \otimes_{\mathbb{Q}} \mathbb{Q}_p$. In this case, the special fiber of $\mathcal{S}_{K,V}^{\mathrm{loc}}$ is known to be smooth (which is called “exotic good reduction”).

In this paper, we consider the case where p ramifies in L and $(r, s) = (2, 2)$. For it, we also consider the similar problem on the corresponding Rapoport–Zink spaces. We explain our results in the sequel.

1.1. Main results. Keep the notations as above (in particular, p is an odd prime). We assume the following:

- $(r, s) = (2, 2)$,
- p is ramified in L and $\nu = p$,
- $\mathbf{G} \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is quasi-split over \mathbb{Q}_p ,
- K_p is a special maximal parahoric subgroup of $\mathbf{G}(\mathbb{Q}_p)$.

Then the reflex field of the Shimura datum (\mathbf{G}, \mathbf{X}) is $E_{2,2} = \mathbb{Q}$, and the integral model $\mathcal{S}_{K,U}$ of $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X}) \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is defined over \mathbb{Z}_p . Moreover, the \mathbb{Z}_p -scheme is defined as a moduli space of quadruples $(A, \iota, \lambda, \bar{\eta}^p)$, where

- A is an abelian 4-fold,
- $\iota: O \rightarrow \mathrm{End}(A)$ is a ring homomorphism,
- λ is a polarization of A such that

$$\mathrm{ord}_p(\#\ker(\lambda)) = \mathrm{ord}_p(\#\ker(\iota(\sqrt{\Delta}))),$$

- $\bar{\eta}^p$ is a K^p -level structure on A .

Here O is an order of L which is maximal at p . We will give a more precise definition of $\mathcal{S}_{K,U}$ in Definition 8.2.

Theorem 1.1. (Theorem 8.7 (i)–(iii))

- (i) *There is a decomposition into open and closed subschemes*

$$\mathcal{S}_{K,U} = \mathcal{S}_{K,U}^{\mathrm{loc}} \sqcup (\mathcal{S}_{K,U} \setminus \mathcal{S}_{K,U}^{\mathrm{loc}}).$$

- (ii) *The scheme $\mathcal{S}_{K,U}^{\mathrm{loc}}$ is flat over \mathbb{Z}_p and regular of dimension 5.*
 (iii) *The scheme $\mathcal{S}_{K,U} \setminus \mathcal{S}_{K,U}^{\mathrm{loc}}$ is smooth over \mathbb{F}_p and 3-dimensional.*

Theorem 1.2. (Theorem 8.8) *The supersingular locus $\mathcal{S}_{K,U}^{\mathrm{ss}}$ consists of two basic loci $\mathcal{S}_{K,U,[b_0]}$ and $\mathcal{S}_{K,U,[b_1]}$, where $\mathcal{S}_{K,U,[b_0]} \subset \mathcal{S}_{K,U}^{\mathrm{loc}}$, and $\mathcal{S}_{K,U,[b_1]} \subset \mathcal{S}_{K,U} \setminus \mathcal{S}_{K,U}^{\mathrm{loc}}$.*

We give concrete descriptions the structure of these two basic loci.

Theorem 1.3. (Theorem 8.10 (i), (ii))

- (i) The scheme $\mathcal{S}_{K,U,[b_0]}$ is purely 2-dimensional. Any irreducible component is birational to the Fermat surface F_p defined by

$$x_0^{p+1} + x_1^{p+1} + x_2^{p+1} + x_3^{p+1} = 0$$

in $\mathbb{P}_{\mathbb{F}_p}^3 = \text{Proj } \overline{\mathbb{F}_p}[x_0, x_1, x_2, x_3]$.

- (ii) Let C be an irreducible component of $\mathcal{S}_{K,U,[b_0]}$. Then, for any irreducible component $C' \neq C$ of $\mathcal{S}_{K,U,[b_0]}$, the intersection $C \cap C'$ is either the empty set, a single point or birational to $\mathbb{P}_{\mathbb{F}_p}^1$. Moreover, the following hold:

$$\begin{aligned} \#\{C' \mid C \cap C' \text{ is a single point}\} &\leq p(p+1)(p^2+1), \\ \#\{C' \mid C \cap C' \text{ is birational to } \mathbb{P}_{\mathbb{F}_p}^1\} &\leq (p+1)(p^2+1). \end{aligned}$$

Theorem 1.4. (Theorem 8.12)

- (i) The scheme $\mathcal{S}_{K,U,[b_1]}$ is purely 1-dimensional. Any irreducible component is birational to $\mathbb{P}_{\mathbb{F}_p}^1$.
- (ii) Let C be an irreducible component of $\mathcal{S}_{K,U,[b_1]}$. Then, for any irreducible component $C' \neq C$ of $\mathcal{S}_{K,U,[b_1]}$, the intersection $C \cap C'$ is either the empty set or a single point. Moreover, we have

$$\#\{C' \mid C \cap C' \text{ is a single point}\} \leq p(p^2+1).$$

Remark 1.5. We have more precise results on non-smooth points of $\mathcal{S}_{K,U}^{\text{loc}}$ and numbers of connected and irreducible components of $\mathcal{S}_{K,U,[b_j]}$. See Theorems 8.10, 8.11 and 8.13.

To prove Theorems 1.3 and 1.4, we deduce them from an explicit description of the underlying topological spaces of the corresponding Rapoport–Zink spaces by using the p -adic uniformization theorem [RZ96, Theorem 6.30]. Next, we explain the results on Rapoport–Zink spaces as mentioned above.

1.2. Results on the Rapoport–Zink spaces. Let F be a ramified quadratic extension of \mathbb{Q}_p , and O_F the ring of integers of F . We denote by $a \mapsto \bar{a}$ the non-trivial Galois automorphism of F over \mathbb{Q}_p . We fix a uniformizer ϖ of F . Put $W := W(\overline{\mathbb{F}_p})$, the ring of Witt vectors over $\overline{\mathbb{F}_p}$. Then, for a W -scheme S in which p is locally nilpotent, we consider a triple (X, ι, λ) , where

- X is a p -divisible group of dimension 4 and height 8 over S ,
- $\iota: O_F \rightarrow \text{End}(X)$ is a ring homomorphism,
- $\lambda: X \rightarrow X^\vee$ is a quasi-polarization,

satisfying the following conditions for any $a \in O_F$:

- $\det(T - \iota(a) \mid \text{Lie}(X)) = (T^2 - \text{Tr}_{F/\mathbb{Q}_p}(a)T + N_{F/\mathbb{Q}_p}(a))^2$,
- $\lambda \circ \iota(a) = \iota(\bar{a})^\vee \circ \lambda$.

Let us fix such a triple $b = (\mathbb{X}_0, \iota_0, \lambda_0)$ over $\text{Spec } \overline{\mathbb{F}_p}$ such that \mathbb{X}_0 is isoclinic of slope $1/2$ and λ_0 is an isogeny satisfying $\text{Ker}(\lambda_0) = \text{Ker } \iota_0(\varpi)$. We define $\mathcal{M}_{b,\mu}^{(0)}$ (the subscript μ will be introduced in Section 2.1) to be the deformation space of $(\mathbb{X}_0, \iota_0, \lambda_0)$ by O_F -linear quasi-isogenies. It is a formal scheme over $\text{Spf } W$, which is formally locally of finite type. We denote by $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ the underlying reduced subscheme of $\mathcal{M}_{b,\mu}^{(0)}$.

With a fixed b , we associate an F/\mathbb{Q}_p -hermitian space C_b of dimension 4. Then the \mathbb{Q}_p -valued points of the unitary group $U(C_b)(\mathbb{Q}_p)$ of C_b acts on $\mathcal{M}_{b,\mu}^{(0)}$. We say that a lattice T in C_b is a *vertex lattice* if $T \subset T^\vee \subset \varpi^{-1}T$. We call $t := \text{length}_{O_F}(T^\vee/T)$ the type of T . The type t is an even

integer satisfying $0 \leq t \leq 4$. We denote by $\text{VL}_b(t)$ the set of all vertex lattices of type t , and put $\text{VL}_b := \bigcup_{t=0}^2 \text{VL}_b(2t)$. We construct a locally closed stratification of $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ by means of vertex lattices in C_b .

We consider two cases whether C_b is split or not as an F/\mathbb{Q}_p -hermitian space. We say that b is μ -neutral if C_b is split.

Case 1. The μ -neutral case. In this case, $\text{VL}_b(2t)$ is non-empty for any $t \in \{0, 1, 2\}$. We introduce an order \leq on VL_b as follows. For distinct $T, T' \in \text{VL}_b$, we have $T < T'$ if one of the following holds:

- $T \in \text{VL}_b(4)$, $T' \in \text{VL}_b(0)$ and $T \subset T'$,
- $T \in \text{VL}_b(4)$, $T' \in \text{VL}_b(2)$ and $T \subset (T')^\vee$,
- $T \in \text{VL}_b(0)$, $T' \in \text{VL}_b(2)$ and $T \subset (T')^\vee$.

Moreover, put

$$d_b(T) := \begin{cases} 0 & \text{if } T \in \text{VL}_b(4), \\ 4 & \text{if } T \in \text{VL}_b(0), \\ 6 & \text{if } T \in \text{VL}_b(2). \end{cases}$$

With $T \in \text{VL}_b$, we associate a reduced locally closed subscheme $\text{BT}_{b,\mu,T}^{(0)}$ of $\mathcal{M}_{b,\mu}^{(0),\text{red}}$. See Section 6.2.

Theorem 1.6. *Suppose that b is μ -neutral.*

- (i) (Theorem 3.1 (i), Theorem 6.6 (iv)) *The formal scheme $\mathcal{M}_{b,\mu}^{(0)}$ is flat over $\text{Spf } W$, and regular of dimension 5. The non-formally smooth locus of $\mathcal{M}_{b,\mu}^{(0)}$ equals $\coprod_{T \in \text{VL}_b(4)} \text{BT}_{b,\mu,T}^{(0)}$.*
- (ii) (Theorem 5.29) *There is a decomposition into open and closed formal subschemes*

$$\mathcal{M}_{b,\mu}^{(0)} = \mathcal{M}_{b,\mu}^{(0,0)} \sqcup \mathcal{M}_{b,\mu}^{(0,1)}.$$

Moreover, $\mathcal{M}_{b,\mu}^{(0,i),\text{red}} := \mathcal{M}_{b,\mu}^{(0,i)} \cap \mathcal{M}_{b,\mu}^{\text{red}}$ are connected and isomorphic to each other.

- (iii) (Theorem 6.6 (i)) *There is a locally closed stratification*

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \coprod_{T \in \text{VL}_b} \text{BT}_{b,\mu,T}^{(0)}.$$

- (iv) (Theorem 6.6 (ii)) *For $x \in \text{VL}_b$, let $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ be the closure of $\text{BT}_{b,\mu,T}^{(0)}$ in $\mathcal{M}_{b,\mu}^{(0),\text{red}}$. Then we have*

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} = \coprod_{T' \leq T} \text{BT}_{b,\mu,T'}^{(0)}.$$

- (v) (Theorem 6.6 (iii), Theorem 6.7) *For $T \in \text{VL}_b$, every connected component of $\text{BT}_{b,\mu,T}^{(0)}$ is isomorphic to a generalized Deligne–Lusztig variety for the split $\text{SO}_{d_b(T)}$ (it is non-classical if $T \in \text{VL}_b(0) \sqcup \text{VL}_b(2)$). Moreover, the following hold:*

- *if $T \in \text{VL}_b(4)$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is a single point,*
- *if $T \in \text{VL}_b(0)$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is isomorphic to two copies of \mathbb{P}^1 ,*
- *if $T \in \text{VL}_b(2)$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is isomorphic to two copies of the Fermat surface F_p .*

In particular, $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ is purely 2-dimensional.

Case 2. The non- μ -neutral case. In this case, $\text{VL}_b(2t)$ is non-empty if and only if $t \in \{0, 1\}$. Moreover, put

$$d_b(T) := 4 - t(T).$$

With $T \in \text{VL}_b$, we associate a reduced locally closed subscheme $\text{BT}_{b,\mu,T}^{(0)}$ of $\mathcal{M}_{b,\mu}^{(0)}$. See Section 7.

Theorem 1.7. *Suppose that b is not μ -neutral.*

- (i) (Theorem 3.1 (ii)) *The formal scheme $\mathcal{M}_{b,\mu}^{(0)}$ equals $\mathcal{M}_{b,\mu}^{(0)} \times_{\text{Spf } W} \text{Spec } \overline{\mathbb{F}}_p$, which is formally smooth of dimension 3 over $\text{Spec } \overline{\mathbb{F}}_p$.*
- (ii) (Theorem 7.12) *There is a decomposition into open and closed formal subschemes*

$$\mathcal{M}_{b,\mu}^{(0)} = \mathcal{M}_{b,\mu}^{(0,0)} \sqcup \mathcal{M}_{b,\mu}^{(0,1)}.$$

Moreover, $\mathcal{M}_{b,\mu}^{(0,i),\text{red}} := \mathcal{M}_{b,\mu}^{(0,i)} \cap \mathcal{M}_{b,\mu}^{\text{red}}$ are connected and isomorphic to each other.

- (iii) *There is a locally closed stratification*

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \coprod_{T \in \text{VL}_b} \text{BT}_{b,\mu,T}^{(0)}$$

- (iv) (Theorem 7.8 (ii)) *For $T \in \text{VL}_b$, let $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ be the closure of $\text{BT}_{b,\mu,T}^{(0)}$ in $\mathcal{M}_{b,\mu}^{(0),\text{red}}$. Then we have*

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} = \coprod_{T' \subset T} \text{BT}_{b,\mu,T'}^{(0)}.$$

- (v) (Theorem 7.6, Theorem 7.13) *For $T \in \text{VL}_b$, the stratum $\text{BT}_{b,\mu,T}^{(0)}$ is isomorphic to the disjoint union of two (classical) Deligne–Lusztig varieties for the non-split $\text{SO}_{d_b(T)}$ associated with Coxeter elements. Moreover, the following hold:*

- *if $T \in \text{VL}_b(2)$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ consists of two points,*
- *if $T \in \text{VL}_b(0)$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is isomorphic to two copies of \mathbb{P}^1 .*

In particular, $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ is purely 1-dimensional.

First, we sketch the proof of Theorem 1.6. For the proof, we use the results of [Oki19] on a Rapoport–Zink space for $\text{GU}_2(D)$, the quaternionic unitary similitudes group of degree 2. Let $\mathcal{M}_D^{(0)}$ be the height 0 locus of the Rapoport–Zink space for $\text{GU}_2(D)$ appeared in [Oki19]. Then we construct a closed immersion of $\mathcal{M}_D^{(0)}$ into $\mathcal{M}_{b,\mu}^{(0)}$, and prove equality of topological spaces $U(C_b)(\mathbb{Q}_p)\mathcal{M}_D^{(0),\text{red}} = \mathcal{M}_{b,\mu}^{(0),\text{red}}$. On the other hand, in the same argument as [HP14], we construct a 6-dimensional quadratic space $\mathbf{L}_{b,0}$ over \mathbb{Q}_p as a subspace of $\text{End}^0(\mathbb{X}_0) := \text{End}(\mathbb{X}_0) \otimes_{\mathbb{Z}} \mathbb{Q}$ which realizes an exceptional isomorphism associated to the identity $A_3 = D_3$ of Dynkin diagrams. Moreover, we introduce the notion of the vertex lattices in $\mathbf{L}_{b,0}$. Then Theorem 1.6 (ii) follows by using some properties of vertex lattices in $\mathbf{L}_{b,0}$ and the closed immersion $\mathcal{M}_D^{(0)} \rightarrow \mathcal{M}_{b,\mu}^{(0)}$. Moreover, we attach a closed formal subscheme $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ of $\mathcal{M}_{b,\mu}^{(0,0)}$ for each vertex lattice Λ in $\mathbf{L}_{b,0}$, and construct a locally closed stratification of $\mathcal{M}_{b,\mu}^{(0,0),\text{red}}$ indexed by the vertex lattices in $\mathbf{L}_{b,0}$. Finally, we give a connection between the vertex lattices in $\mathbf{L}_{b,0}$ and those in C_b , and construct a locally closed stratification of $\mathcal{M}_{b,\mu}^{(0)}$ indexed by the vertex lattices in C_b by using the correspondence of vertex lattices as above and the stratification of $\mathcal{M}_{b,\mu}^{(0,0)}$.

Note that the use of the above exceptional isomorphism gives us a moduli description of any irreducible component of $\mathcal{M}_{b,\mu}^{(0)}$. We expect an application of such a result, for example an analogue of [Oki19, Section 8].

Second, we give an outline of the proof of Theorem 1.7. The proof is essentially the same as [Wu16]. In this case, we more directly construct a locally closed stratification indexed by the vertex lattices in C_b .

Remark 1.8. We can also describe the whole structure of the supersingular locus $\mathcal{S}_{K,U}^{\text{ss}}$ of the Rapoport–Zink integral model for $(r, s) = (1, 3)$ when p, L and K are as in Section 1.1. In this case, $\mathcal{S}_{K,U}^{\text{ss}} \cap \mathcal{S}_{K,U}^{\text{loc}}$ is studied by [Wu16]. It consists of two basic loci, one of which is $\mathcal{S}_{K,U}^{\text{ss}} \cap \mathcal{S}_{K,U}^{\text{loc}}$, and the other is $\mathcal{S}_{K,U}^{\text{ss}} \setminus \mathcal{S}_{K,U}^{\text{loc}}$. The former one is purely 1-dimensional, and each irreducible component is isomorphic to \mathbb{P}^1 . The latter one is purely 2-dimensional, and each irreducible component is isomorphic to the Fermat surface F_p .

On the other hand, it is natural to consider a description of the whole structure of the supersingular locus $\mathcal{S}_{K,U}^{\text{ss}}$ for $(r, s) = (1, 3)$ when p and L are as above, and K_p is the stabilizer of a *self-dual* lattice in $U \otimes_{\mathbb{Q}} \mathbb{Q}_p$. In this case, the structure of $\mathcal{S}_{K,U}^{\text{ss}} \cap \mathcal{S}_{K,U}^{\text{loc}}$ is determined by [RTW14]. To achieve this question, it suffices to study $\mathcal{S}_{K,U}^{\text{ss}}$ in the case where “ $(r, s) = (1, 3)$ ” is replaced with “ $(r, s) = (2, 2)$ ”. This is one of our future’s problem.

Organization of this paper. In Section 2, we define Rapoport–Zink spaces which will be considered in this paper. In Section 3, we study the singularities of the Rapoport–Zink spaces, and prove Theorem 3.1. In Section 4, we investigate the generalized Deligne–Lusztig varieties for even special orthogonal groups that are appeared in Theorems 1.6 and 1.7. In Sections 5 and 6, we consider the Rapoport–Zink space $\mathcal{M}_{b,\mu}$ in the μ -neutral case. More precisely, in Section 5, we construct a 6-dimensional quadratic space $\mathbf{L}_{b,0}$ over \mathbb{Q}_p , and give the definition of vertex lattices. Moreover, we construct the Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0,0)}$ indexed by vertex lattices in $\mathbf{L}_{b,0}$. In Section 6, we introduce the notion of the vertex lattices in C_b , and construct the Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0)}$. In particular, we prove Theorem 1.6 in this section. In Section 7, we construct the Bruhat–Tits stratification of the Rapoport–Zink space $\mathcal{M}_{b,\mu}$ in the non- μ -neutral case, and prove Theorem 1.7. In Section 8, we recall the precise definition of the Rapoport–Zink integral models of some Shimura varieties for (\mathbf{G}, \mathbf{X}) , and prove the results in Section 1.1.

Acknowledgment. I would like to thank my advisor Yoichi Mieda for his constant support and encouragement.

This work was carried out with the support from the Program for Leading Graduate Schools, MEXT, Japan. This work was also supported by the JSPS Research Fellowship for Young Scientists and KAKENHI Grant Number 19J21728.

Notations. We use the following notations in this paper.

- **p -adic fields and their extensions.** For a field k of characteristic p , let $W(k)$ be the Cohen ring over k . Note that it is the ring of Witt vectors if k is perfect. We denote by σ the p -th power Frobenius on k or its lift to $W(k)$. If $k = \overline{\mathbb{F}}_p$, write W and K_0 for $W(k)$ and $\text{Frac}(W(k))$ respectively. We normalize the p -adic valuations ord_p on \mathbb{Q}, \mathbb{Q}_p and $\text{Frac}(W(k))$ so that $\text{ord}_p(p) = 1$.
- For a scheme S , we denote by $\text{Irr}(S)$ the set of irreducible components of S .

2. RAPOPORT–ZINK SPACES FOR A RAMIFIED GU(2, 2)

2.1. Local PEL datum. We fix a datum which will be used until Section 7. Let p be an odd prime number, F a ramified quadratic extension field of \mathbb{Q}_p , and O_F the integer ring of F . Fix a uniformizer ϖ of O_F , and denote by $a \mapsto \bar{a}$ the non-trivial Galois automorphism of F over \mathbb{Q}_p . Moreover, let V be a 4-dimensional F -vector space, and $(,)$ a \mathbb{Q}_p -valued alternating form on V satisfying $(ax, y) = (x, \bar{a}y)$ for any $a \in F$ and $x, y \in V$. Note that there is a unique F/\mathbb{Q}_p -hermitian form \langle , \rangle on V such that

$$(,) = \frac{1}{2} \text{Tr}_{F/\mathbb{Q}_p}(\varpi^{-1} \langle , \rangle).$$

In this paper, we assume that the F/\mathbb{Q}_p -hermitian space $(V, \langle \cdot, \cdot \rangle)$ is split, that is, there is an F -basis $\mathbf{e}_1, \dots, \mathbf{e}_4$ of V such that $\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{i,5-j}$. We fix $\mathbf{e}_1, \dots, \mathbf{e}_4$ as above, and put

$$\mathbf{\Lambda} := O_F \mathbf{e}_1 \oplus O_F \mathbf{e}_2 \oplus O_F(\varpi \mathbf{e}_3) \oplus O_F(\varpi \mathbf{e}_4).$$

Then it is a ϖ -modular lattice, that is, $\mathbf{\Lambda}^\vee = \varpi^{-1} \mathbf{\Lambda}$.

We define an algebraic group G over \mathbb{Q}_p by

$$G(R) = \{(g, c) \in \mathrm{GL}_{F \otimes_{\mathbb{Q}_p} R}(V \otimes_{\mathbb{Q}_p} R) \times R^\times \mid (g(x), g(y)) = c(x, y) \text{ for all } x, y \in V \otimes_{\mathbb{Q}_p} R\}$$

for any \mathbb{Q}_p -algebra R . Note that G is reductive and quasi-split over \mathbb{Q}_p . We denote by $\mathrm{sim}_G: G \rightarrow \mathbb{G}_m$ the similitude character of G .

We regard G as a subgroup of $\mathrm{Res}_{F/\mathbb{Q}_p} \mathrm{GL}_F(V)$ by the fixed basis $\mathbf{e}_1, \dots, \mathbf{e}_4$, and put

$$\mu: \mathbb{G}_m \rightarrow G; z \mapsto \mathrm{diag}(1, 1, z, z).$$

2.2. p -divisible groups and isocrystals with additional structures. For a field extension $k/\overline{\mathbb{F}}_p$, let $F_{W(k)} := F \otimes_{\mathbb{Z}_p} W(k)$ and $O_{F, W(k)} := O_F \otimes_{\mathbb{Z}_p} W(k)$. Note that $F_{W(k)}$ is a field and $O_{\check{F}}$ is the integer ring of \check{F} . Especially, we write \check{F} and $O_{\check{F}}$ for $F_{W(\overline{\mathbb{F}}_p)}$ and $O_{F, W(\overline{\mathbb{F}}_p)}$ respectively.

We denote by σ the map $\mathrm{id}_F \otimes \sigma$ on $O_{F, W(k)}$. Moreover, we extend the map $a \mapsto \bar{a}$ to $F_{W(k)}$ as a $\mathrm{Frac} W(k)$ -linear map.

On the other hand, let Nilp be the category of W -schemes in which p is locally nilpotent. For $S \in \mathrm{Nilp}$, put $\overline{S} := S \times_{\mathrm{Spec} W} \mathrm{Spec} \overline{\mathbb{F}}_p$.

Definition 2.1. Let $S \in \mathrm{Nilp}$.

- (i) A p -divisible group with (G, μ) -structure over S is a triple (X, ι, λ) consisting of the following data:
- X is a p -divisible group over S of dimension 4 and height 8,
 - $\iota: O_F \rightarrow \mathrm{End}(X)$ is a ring homomorphism,
 - $\lambda: X \rightarrow X^\vee$ is a quasi-isogeny so that $\lambda^\vee = -\lambda$,
- satisfying the following conditions for any $a \in O_F$:
- (Kottwitz condition) $\det(T - \iota(a) \mid \mathrm{Lie}(X)) = (T^2 - \mathrm{Tr}_{F/\mathbb{Q}_p}(a)T + \mathrm{N}_{F/\mathbb{Q}_p}(a))^2$,
 - $\lambda \circ \iota(a) = \iota(\bar{a})^\vee \circ \lambda$.
- (ii) For two p -divisible groups with (G, μ) -structures (X, ι, λ) and (X', ι', λ') over S , a quasi-isogeny from (X, ι, λ) to (X', ι', λ') is an O_F -linear quasi-isogeny $\rho: X \rightarrow X'$ satisfying $\rho^\vee \circ \lambda' \circ \rho = c(\rho)\lambda$ for some locally constant function $c(\rho): \overline{S} \rightarrow \mathbb{Q}_p^\times$.

Remark 2.2. The cocharacter μ appears in the Kottwitz condition. In fact, the right-hand side can be written as $\det(a \mid V_0)$ in the sense of [RZ96, 3.23 a)], where

$$V_0 := \{x \in V \otimes_{\mathbb{Q}_p} K_0 \mid \mu(z)x = x \text{ for all } z \in \mathbb{G}_m\} = \check{F}\mathbf{e}_1 \oplus \check{F}\mathbf{e}_2.$$

It is convenient to introduce the notion of isocrystals with G -structures.

Definition 2.3. Let k be an algebraically closed field of characteristic p .

- (i) An isocrystal with G -structure over k is a triple $(N, \mathbf{F}_N, (\cdot, \cdot)_N)$, where
- N is a finite-dimensional $F \otimes_{\mathbb{Z}_p} W(k)$ -vector space,
 - $\mathbf{F}_N: N \rightarrow N$ is a σ -linear map,
 - $(\cdot, \cdot)_N: N \times N \rightarrow \mathrm{Frac} W(k)$ is a non-degenerate alternating form satisfying $(ax, y)_N = (x, \bar{a}y)_N$ for any $a \in F \otimes_{\mathbb{Z}_p} W(k)$.

- (ii) For two isocrystals with G -structures $(N, \mathbf{F}_N, (\cdot, \cdot)_N)$ and $(N', \mathbf{F}_{N'}, (\cdot, \cdot)_{N'})$, a morphism of isocrystals with G -structures is a homomorphism of $F \otimes_{\mathbb{Z}_p} W(k)$ -vector space $\varphi: N \xrightarrow{\cong} N'$ and $c \in (F \otimes_{\mathbb{Z}_p} W(k))^\times$ such that $\varphi \circ \mathbf{F}_N = \mathbf{F}_{N'} \circ \varphi$ and $(\varphi(x), \varphi(y))_{N'} = c(x, y)_N$ for any $x, y \in N$.

We can construct isocrystals with G -structure by elements of $G(K_0)$ or p -divisible groups with (G, μ) -structures as follow:

- (i) Set $k := \overline{\mathbb{F}}_p$. For $b \in G(K_0)$, we define an isocrystal over $\overline{\mathbb{F}}_p$ as follows:

$$(N_b, \mathbf{F}_b) := (V \otimes_{\mathbb{Q}_p} K_0, b(\mathrm{id} \otimes \sigma)).$$

Then $(N_b, \mathbf{F}_b, (\cdot, \cdot))$ is an isocrystal with G -structure over $\overline{\mathbb{F}}_p$ by the natural way. Then, for two elements b and b' of $G(K_0)$, there is an isomorphism between $(N_b, \mathbf{F}_b, (\cdot, \cdot))$ and $(N_{b'}, \mathbf{F}_{b'}, (\cdot, \cdot))$ if and only if b and b' are σ -conjugate, that is, there is $g \in G(K_0)$ such that $b' = gb\sigma(g)^{-1}$. Hence this correspondence induces a bijection between the set of all σ -conjugacy class of $G(K_0)$ and the set of isomorphism classes of isocrystals with G -structures over $\overline{\mathbb{F}}_p$.

- (ii) Let p -divisible group with G -structure (X, ι, λ) over k . Consider the covariant rational Dieudonné module $\mathbb{D}(X)_{\mathbb{Q}}$ of X . Then ι and λ induces an F -action and a non-degenerate alternating form on $\mathbb{D}(X)_{\mathbb{Q}}$ respectively, and $\mathbb{D}(X)_{\mathbb{Q}}$ becomes an isocrystal with G -structure over k by their structures.

2.3. Definition of the Rapoport–Zink spaces. Let $b \in G(K_0)$. Here we assume that (N_b, \mathbf{F}_b) is isoclinic of slope $1/2$. Take a p -divisible group with (G, μ) -structure $(\mathbb{X}_0, \iota_0, \lambda_0)$ over $\mathrm{Spec} \overline{\mathbb{F}}_p$ such that $\mathbb{D}(\mathbb{X}_0)_{\mathbb{Q}} \cong N_b$ as isocrystals with G -structures over $\overline{\mathbb{F}}_p$ and $\mathrm{Ker}(\lambda_0) = \mathrm{Ker}(\iota_0(\varpi))$. We define $\mathcal{M}_{b, \mu}$, the *Rapoport-Zink space for (G, b, μ)* , as the functor that parameterizes the equivalence classes of $(X, \iota, \lambda, \rho)$ for $S \in \mathrm{Nilp}_W$, where (X, ι, λ) is a p -divisible group with (G, μ) -structure over S satisfying $\mathrm{Ker}(\lambda) = \mathrm{Ker}(\iota(\varpi))$, and

$$\rho: X \times_S \overline{S} \rightarrow \mathbb{X}_0 \otimes_{\overline{\mathbb{F}}_p} \overline{S}$$

is a quasi-isogeny of p -divisible groups with (G, μ) -structures. Two p -divisible groups with G -structures with quasi-isogenies $(X, \iota, \lambda, \rho)$ and $(X', \iota', \lambda', \rho')$ are equivalent if $(\rho')^{-1} \circ \rho$ lifts to an isomorphism $X \rightarrow X'$ over S .

The functor $\mathcal{M}_{b, \mu}$ is representable by a formal scheme over $\mathrm{Spf} W$, which is formally locally of finite type. Moreover, there is a decomposition into open and closed formal subschemes

$$\mathcal{M}_{b, \mu} = \coprod_{i \in \mathbb{Z}} \mathcal{M}_{b, \mu}^{(i)},$$

where $\mathcal{M}_{b, \mu}^{(i)}$ is the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_{b, \mu}$ where $\mathrm{Im}(c(\rho)) \subset p^i \mathbb{Z}_p^\times$. These follow from [RZ96, Theorem 3.25].

We define another F/\mathbb{Q}_p -hermitian space C_b associated to b . Consider the isocrystal with G -structure $(N_b, \mathbf{F}_b, (\cdot, \cdot))$ over $\overline{\mathbb{F}}_p$. Fix a square root $\eta \in W^\times$ of $p^{-1}\varpi^2$, and put $\mathbf{F}_{b,0} := \eta\varpi^{-1}\mathbf{F}_b$. By [RTW14, p.1171], we have

$$(\mathbf{F}_{b,0}(x), \mathbf{F}_{b,0}(y)) = \sigma((x, y)), \quad \langle \mathbf{F}_{b,0}(x), \mathbf{F}_{b,0}(y) \rangle = \sigma(\langle x, y \rangle)$$

for any $x, y \in N_b$. Moreover, $(N_b, \mathbf{F}_{b,0})$ is isoclinic of slope 0. Let C_b be the $\mathbf{F}_{b,0}$ -fixed part of N_b . Then we can consider (\cdot, \cdot) and $\langle \cdot, \cdot \rangle$ on C_b . Moreover, $(C_b, \langle \cdot, \cdot \rangle)$ is an F/\mathbb{Q}_p -hermitian space of dimension 4.

We recall an algebraic group J_b over \mathbb{Q}_p defined by [RZ96, 1.12]. It is defined by

$$J_b(R) = \{g \in G(R \otimes_{\mathbb{Q}_p} K_0) \mid g \circ \mathbf{F}_b = \mathbf{F}_b \circ g\}$$

for any \mathbb{Q}_p -algebra R . Let $\text{sim}_{J_b}: J_b \rightarrow \mathbb{G}_m$ be the character induced by sim_G . As in [RZ96, 1.12], the group $J_b(\mathbb{Q}_p)$ acts on $\mathcal{M}_{b,\mu}$ by

$$J_b(\mathbb{Q}_p) \times \mathcal{M}_{b,\mu}(S) \rightarrow \mathcal{M}_{b,\mu}(S); (g, (X, \iota, \lambda, \rho)) \mapsto (X, \iota, \lambda, g \circ \rho).$$

This action induces an isomorphism $\mathcal{M}_{b,\mu}^{(0)} \cong \mathcal{M}_{b,\mu}^{(i)}$ for each $i \in \mathbb{Z}$. Moreover, $\mathcal{M}_{b,\mu}^{(0)}$ is stable under $J_b^1(\mathbb{Q}_p)$, where

$$J_b^1 := \{g \in J_b \mid \text{sim}_{J_b}(g) = 1\}.$$

The structure of J_b is as follows:

Proposition 2.4. ([RTW14, Proposition 1.5]) *There is an isomorphism of \mathbb{Q}_p -algebraic groups*

$$J_b \cong \text{GU}(C_b).$$

Moreover, sim_{J_b} equals the composition of the similitude character of $\text{GU}(C_b)$ and the above isomorphism.

Let $k/\overline{\mathbb{F}}_p$ be a field extension. We give a description of the set of k -rational points of $\mathcal{M}_{b,\mu}^{(0)}$ by means of linear algebra. Put $N_{b,W(k)} := N_b \otimes_W W(k)$, and let $\text{DiL}^\varpi(N_{b,W(k)})$ be the set all $O_{F,W(k)}$ -lattices M in $N_{b,W(k)}$ satisfying the following conditions:

- $M^\vee = \varpi^{-1}M$,
- $pM \subset \mathbf{F}_b^{-1}(pM) \subset M$,
- $\text{length}_{O_{F,W(k)}} M/\mathbf{F}_b^{-1}(pM) = 4$ and $\text{length}_{O_{F,W(k)}} \varpi M + \mathbf{F}_b^{-1}(pM)/\mathbf{F}_b^{-1}(pM) \leq 2$.

Proposition 2.5. *There is a $J_b^1(\mathbb{Q}_p)$ -equivariant bijection*

$$\mathcal{M}_{b,\mu}^{(0)}(k) \cong \text{DiL}^\varpi(N_{b,W(k)}).$$

Proof. These follow from the theory of windows. See [Zin01]. Note that the third condition on $\text{DiL}^\varpi(N_{b,W(k)})$ is equivalent to the Kottwitz condition. ■

2.4. The neutralness condition. Here we introduce the notion of the neutralness condition which is defined in [RV14, Definition 2.5]. For this, we recall the *Kottwitz map* defined by [Kot85]. Let $B(G)$ be the set of all σ -conjugacy classes of $G(K_0)$, and $\pi_1(G)$ the Borovoi's algebraic fundamental group of G . We denote by Γ the absolute Galois group of \mathbb{Q}_p . Then the Kottwitz map is the map

$$\overline{\kappa}_G: B(G) \rightarrow \pi_1(G)_\Gamma.$$

By [RR96, Theorem 1.15], it induces an isomorphism

$$B(G)_{\text{basic}} \xrightarrow{\cong} \pi_1(G)_\Gamma,$$

where $B(G)_{\text{basic}}$ is the set of σ -conjugacy classes of $G(K_0)$ which is basic in the sense of [Kot85].

We can find an explicit description of $\overline{\kappa}_G$ in [PR09, Section 1.2] and [RSZ17, Section 3.3]. More precisely, there is an isomorphism

$$\psi: \pi_1(G)_\Gamma \xrightarrow{\cong} \mathbb{Z} \times \mathbb{Z}/2.$$

Moreover, the composite $\psi \circ \overline{\kappa}_G$ is induced by

$$\kappa_G: G(K_0) \rightarrow \mathbb{Z} \times \mathbb{Z}/2; b \mapsto (\text{ord}_p(\text{sim}_G(b)), \det_{\check{F}}(b)/\text{sim}_G(b)^2 \bmod \varpi O_{\check{F}})$$

(note that $\det_{\check{F}}(g)/\text{sim}_G(g)^2$ has \check{F}/K_0 -norm 1).

Definition 2.6. Let μ be as in Section 2.1, and denote by μ^\natural the image of μ under the map

$$X_*(G) \rightarrow X_*(G^{\mathrm{ab}})_\Gamma \cong \pi_1(G)_\Gamma.$$

An element $b \in G(K_0)$ or $[b] \in B(G)$ is said to be μ -neutral if $\kappa_G([b]) = \mu^\natural$.

Remark 2.7. In [RV14, Definition 2.5], the neutral condition is defined among elements in $B(G)$ which is *acceptable* with respect to μ . However, the above definition makes sense for any $b \in B(G)$.

For $b \in G(K_0)$, put $[b] := \{gb\sigma(g)^{-1} \in G(K_0) \mid g \in G(K_0)\}$. We determine and consider the μ -neutral condition on the set

$$B(G)_{1,\mathrm{basic}} := \{[b] \in B(G)_{\mathrm{basic}} \mid \mathrm{ord}_p(\mathrm{sim}_G(b)) = 1\}.$$

Take $[b] \in B(G)_{1,\mathrm{basic}}$. Then it is represented by one of the following:

$$b_0 := \begin{pmatrix} & & & \varpi \\ & & \varpi & \\ & \varpi & & \\ \varpi & & & \end{pmatrix}, \quad b_1 := \begin{pmatrix} & & & \varpi \\ & \varpi & 0 & \\ & 0 & -\varpi & \\ \varpi & & & \end{pmatrix}$$

Hence we have an equality

$$B(G)_{1,\mathrm{basic}} = \{[b_0], [b_1]\}.$$

Note that $\kappa_G([b_j]) = (1, j)$ for $j \in \{0, 1\}$, and $(N_{b_j}, \mathbf{F}_{b_j})$ is isoclinic of slope $1/2$. On the other hand, we have $\psi(\mu^\natural) = (1, 0)$. Hence $[b]$ is μ -neutral if and only if $[b] = [b_0]$.

We also obtained a criterion for the basic condition by means of an isocrystal.

Proposition 2.8. *An element $b \in G(K_0)$ satisfies $[b] \in B(G)_{1,\mathrm{basic}}$ if and only if (N_b, \mathbf{F}_b) is isoclinic of slope $1/2$.*

Proof. By the argument as above, (N_b, \mathbf{F}_b) is isoclinic of slope $1/2$ if $[b] \in B(G)_{1,\mathrm{basic}}$. On the other hand, suppose that (N_b, \mathbf{F}_b) is isoclinic of slope $1/2$. Then it is clear that $\mathrm{ord}_p(\mathrm{sim}_G(b)) = 1$. On the other hand, Proposition 2.4 implies that J_b is an inner form, that is, b is basic. Hence we have $[b] \in B(G)_{1,\mathrm{basic}}$. \blacksquare

Remark 2.9. Both $[b_0]$ and $[b_1]$ are acceptable with respect to μ . This follows from the weak admissibility (this can be checked directly) and [RV14, Proposition 2.9].

Finally, we give a characterization of the μ -neutralness condition for $b \in G(K_0)$ satisfying $[b] \in B(G)_{1,\mathrm{basic}}$ by means of linear algebra.

Proposition 2.10. *Let $b \in G(K_0)$ satisfying $[b] \in B(G)_{1,\mathrm{basic}}$. Then the following are equivalent:*

- (i) $[b]$ is μ -neutral,
- (ii) $\det_{\mathbb{F}}(b)/\mathrm{sim}_G(b) \in 1 + \varpi O_{\mathbb{F}}$,
- (iii) C_b is split as an F/\mathbb{Q}_p -hermitian space,
- (iv) $\mathrm{length}_{O_{\mathbb{F}}} \varpi M + \mathbf{V}_b(M)/\mathbf{V}_b(M)$ is an even integer for any (some) $M \in \mathrm{DiL}^\varpi(N_b)$, where $\mathbf{V}_b := p\mathbf{F}_b^{-1}$.

Proof. The assertion (i) \iff (ii) follows from the description of the Kottwitz map as above. The assertion (iii) \iff (iv) follow from [RSZ17, Lemma 3.3] and the isomorphism

$$\mathbf{V}_b: M + \mathbf{F}_{b,0}(M)/M \xrightarrow{\cong} \varpi M + \mathbf{V}_b(M)/\mathbf{V}_b(M).$$

We prove (i) \iff (iv). We may assume that b equals either b_0 or b_1 . Then we have $\mathbf{\Lambda} \in \text{DiL}^\varpi(N_{b_j})$ and

$$\text{length}_{\mathcal{O}_{\overline{\mathbb{F}}}} \varpi \mathbf{\Lambda} + \mathbf{V}_{b_j}(\mathbf{\Lambda})/\mathbf{V}_{b_j}(\mathbf{\Lambda}) = j$$

for $j \in \{0, 1\}$. Hence the equivalence between (i) and (iv) follows. \blacksquare

3. SINGULARITIES OF THE RAPOPORT–ZINK SPACES

In this section, we prove the following:

Theorem 3.1. *Let $b \in G(K_0)$ be as in Section 2.3.*

- (i) *If b is μ -neutral, then the formal scheme $\mathcal{M}_{b,\mu}$ is regular. Moreover, it is formally smooth over $\text{Spf } W$ outside the discrete set of $\overline{\mathbb{F}}_p$ -rational points such that the corresponding p -divisible group with (G, μ) -structure $(X, \iota, \lambda, \rho)$ over $\overline{\mathbb{F}}_p$ satisfies $\iota(\varpi) = 0$ on $\text{Lie}(X)$.*
- (ii) *If b is not μ -neutral, then we have $\mathcal{M}_{b,\mu} = \mathcal{M}_{b,\mu} \times_{\text{Spf } W} \text{Spec } \overline{\mathbb{F}}_p$, and it is formally smooth of dimension 3 over $\text{Spec } \overline{\mathbb{F}}_p$.*

To prove Theorem 3.1, we determine the singularity of certain scheme, which is called the local model for (G, μ) . Let $\mathbf{\Lambda}$ be as in Section 2.1, and put $\mathbf{\Lambda}_0 := \varpi^{-1}\mathbf{\Lambda}$.

Definition 3.2. (i) We define an \mathbb{Z}_p -scheme M_μ^{naive} as the functor which parametrizes \mathcal{F} of $\mathbf{\Lambda}_{0, \mathcal{O}_S} := \mathbf{\Lambda}_0 \otimes_{\mathbb{Z}_p} \mathcal{O}_S$ satisfying the following conditions:

- $\det(T - a \mid \mathcal{F}) = (T^2 - \text{Tr}_{F/\mathbb{Q}_p}(a)T + \text{N}_{F/\mathbb{Q}_p}(a))^2$ for any $a \in \mathcal{O}_F$,
- \mathcal{F} is a maximally totally isotropic subspace with respect to the pairing

$$(1) \quad \mathbf{\Lambda}_{0, \mathcal{O}_S} \times \mathbf{\Lambda}_{0, \mathcal{O}_S} \xrightarrow{\varpi \times \text{id}} \mathbf{\Lambda}_{0, \mathcal{O}_S} \times \mathbf{\Lambda}_{0, \mathcal{O}_S} \xrightarrow{(\cdot, \cdot)} \mathcal{O}_S.$$

Note that M_μ^{naive} is representable by a projective scheme over \mathbb{Z}_p .

- (ii) We define M_μ^{loc} as the scheme theoretic closure of the generic fiber $M_{\mu, \mathbb{Q}_p}^{\text{naive}}$ of M_μ^{naive} .

We denote by $[\cdot, \cdot]_0$ the pairing (1), and define an affine group scheme \mathcal{G} over \mathbb{Z}_p by

$$\mathcal{G}(R) = \{(g, c) \in \text{GL}_{\mathcal{O}_F \otimes_{\mathbb{Z}_p} R}(\mathbf{\Lambda}_0 \otimes_{\mathbb{Z}_p} R) \times R^\times \mid [g(x), g(y)]_0 = c[x, y]_0 \text{ for all } x, y \in \mathbf{\Lambda}_0 \otimes_{\mathbb{Z}_p} R\}$$

for any \mathbb{Z}_p -algebra R . Note that we have $\mathcal{G}(\mathbb{Q}_p) = G(\mathbb{Q}_p)$, and $\mathcal{G}(\mathbb{Z}_p)$ is the stabilizer of $\mathbf{\Lambda}_0$ in $G(\mathbb{Q}_p)$. Moreover, there is an action of \mathcal{G} on M_μ^{naive} by

$$\mathcal{G}(S) \times M_\mu^{\text{naive}}(S) \rightarrow M_\mu^{\text{naive}}(S); (g, \mathcal{F}) \mapsto g(\mathcal{F})$$

for any \mathbb{Z}_p -scheme S .

Proposition 3.3.

- (i) *There is a decomposition into open and closed subschemes*

$$M_\mu^{\text{naive}} = M_\mu^{\text{loc}} \sqcup (M_\mu^{\text{naive}} \setminus M_\mu^{\text{loc}}).$$

- (ii) *The scheme M_μ^{loc} is regular and irreducible of dimension 5. The singular locus of M_μ^{loc} consists the unique \mathbb{F}_p -rational point y_0 such that the corresponding subspace \mathcal{F}_0 of $\mathbf{\Lambda}_{0, \mathbb{F}_p}$ satisfying $\varpi \mathcal{F}_0 = 0$. Moreover, there is an isomorphism*

$$\mathcal{O}_{M_\mu^{\text{loc}}, y_0} \cong (\mathbb{Z}_p[t_0, t_1, t_2, t_3, t_4]/(t_0^2 + t_1 t_2 + t_3 t_4 - \varpi^2))_{(p, t_0, t_1, t_2, t_3, t_4)}.$$

- (iii) *The scheme $M_\mu^{\text{naive}} \setminus M_\mu^{\text{loc}}$ is the locus of M_μ^{naive} where the rank of ϖ on the corresponding subspace equals 1. Moreover, it is a smooth \mathbb{F}_p -scheme of dimension 3.*

Proposition 3.3 \Rightarrow *Proposition 3.1*. By [RZ96, Theorem 3.33], there is a diagram

$$\mathcal{M}_{b,\mu} \xleftarrow{\varphi} \widetilde{\mathcal{M}}_{b,\mu} \xrightarrow{\psi} \widehat{M}_\mu^{\text{naive}},$$

where $\widehat{M}_\mu^{\text{naive}}$ is the p -adic completion of $M_\mu^{\text{naive}} \times_{\text{Spec } \mathbb{Z}_p} \text{Spec } W$. Note that the map φ is a \mathcal{G} -torsor, and ψ is formally smooth. Moreover, there is a decomposition of open closed formal subschemes corresponding to Proposition 3.3 (i):

$$\widehat{M}_\mu^{\text{naive}} = \widehat{M}_\mu^{\text{loc}} \sqcup (M_\mu^{\text{naive}} \setminus M_\mu^{\text{loc}}).$$

Now take $x \in \mathcal{M}_{b,\mu}(\mathbb{F}_p)$ and $y \in \widetilde{\mathcal{M}}_{b,\mu}(\overline{\mathbb{F}}_p)$ satisfying $\varphi(y) = x$. By Propositions 2.5 and 2.10, the image of y under ψ lies in $\widehat{M}_\mu^{\text{loc}}$ if b is μ -neutral, and $M_\mu^{\text{naive}} \setminus M_\mu^{\text{loc}}$ if b is not μ -neutral. Hence the assertions follow from Proposition 3.3 (ii), (iii). \blacksquare

We also obtain a moduli description of M_μ^{loc} as follows:

Corollary 3.4. The scheme M_μ^{loc} equals the spin local model M_μ^{spin} (see [PR09] or [Smi14] for the definition of the spin local model).

Proof. As explained in [Smi14, 2.4], M_μ^{spin} is a closed subscheme of M_μ^{naive} . Since $(M_\mu^{\text{spin}})^{\text{red}} = M_\mu^{\text{loc}}$ by [Smi14, Theorem 1.3], Proposition 3.3 (i) implies that the canonical closed immersion $M_\mu^{\text{spin}} \hookrightarrow M_\mu^{\text{naive}}$ factors through $M_\mu^{\text{loc}} \hookrightarrow M_\mu^{\text{naive}}$. Hence the assertion follows. \blacksquare

We turn to prove Proposition 3.3. For it, we recall the structure of the generic fiber of M_μ^{loc} .

Proposition 3.5. ([PR09, 1.5.3]) *The generic fiber $M_{\mu,\mathbb{Q}_p}^{\text{naive}}$ of M_μ^{naive} is connected and smooth of dimension 4 over \mathbb{Q}_p .*

We rewrite a \mathbb{Z}_p -basis of Λ_0 as follows:

$$\begin{aligned} \mathbf{f}_1 &:= \varpi^{-1} \mathbf{e}_1, & \mathbf{f}_2 &:= \varpi^{-1} \mathbf{e}_2, & \mathbf{f}_3 &:= \mathbf{e}_3, & \mathbf{f}_4 &:= \mathbf{e}_4, \\ \mathbf{f}_5 &:= \mathbf{e}_1, & \mathbf{f}_6 &:= \mathbf{e}_2, & \mathbf{f}_7 &:= \varpi \mathbf{e}_3, & \mathbf{f}_8 &:= \varpi \mathbf{e}_4. \end{aligned}$$

Then the Gram matrix of $[\ ,]_0$ with respect to the above basis is

$$\mathbf{J} = \begin{pmatrix} & & & J_2 \\ & & -J_2 & \\ & -J_2 & & \\ J_2 & & & \end{pmatrix},$$

where J_2 is the anti-diagonal matrix of size 2 with coefficient 1.

Let $\text{OGr}(\Lambda_0)$ be the Grassmanian of maximally totally isotropic subspaces of Λ_0 . Note that it has exactly two connected components. Then there is a natural closed immersion

$$M_\mu^{\text{naive}} \rightarrow \text{OGr}(\Lambda_0).$$

Since $M_{\mu,\mathbb{Q}_p}^{\text{naive}}$ is connected by Proposition 3.5, there is a unique connected component $\text{OGr}^+(\Lambda_0)$ of $\text{OGr}(\Lambda_0)$ containing the image of $M_{\mu,\mathbb{Q}_p}^{\text{naive}}$ under the above closed immersion. We denote by $\text{OGr}^-(\Lambda_0)$ the other connected component of $\text{OGr}(\Lambda_0)$.

We write $M_{\mu,\mathbb{F}_p}^{\text{naive}}$ for the special fiber of M_μ^{naive} . For $r \in \{0, 1, 2\}$, let C_r be the locus of $M_{\mu,\mathbb{F}_p}^{\text{naive}}$ where the rank of ϖ on the corresponding subspace equals r . Then we have a decomposition into locally closed subschemes

$$M_{\mu,\mathbb{F}_p}^{\text{naive}} = C_0 \sqcup C_1 \sqcup C_2.$$

Note that the above decomposition is also the decomposition into \mathcal{G} -orbits. Moreover, we have

$$M_{\mu, \mathbb{F}_p}^{\text{naive}} \cap \text{OGr}^+(\mathbf{\Lambda}_0) = C_0 \sqcup C_2, \quad M_{\mu, \mathbb{F}_p}^{\text{naive}} \cap \text{OGr}^-(\mathbf{\Lambda}_0) = C_1$$

by [PR09, 7.1.4]. In particular, C_1 is open and closed in $M_{\mu, \mathbb{F}_p}^{\text{naive}}$.

Remark 3.6. The spin local model M_{μ}^{spin} equals $M_{\mu, \mathbb{F}_p}^{\text{naive}} \cap \text{OGr}^+(\mathbf{\Lambda}_0) = C_0 \sqcup C_2$ by the definition of the spin condition. See [PR09, 7.1.4, 7.2.1].

Now put

$$\begin{aligned} \mathcal{F}_0 &:= \mathbb{F}_p \mathbf{f}_5 \oplus \mathbb{F}_p \mathbf{f}_6 \oplus \mathbb{F}_p \mathbf{f}_7 \oplus \mathbb{F}_p \mathbf{f}_8, \\ \mathcal{F}_1 &:= \mathbb{F}_p \mathbf{f}_1 \oplus \mathbb{F}_p \mathbf{f}_5 \oplus \mathbb{F}_p \mathbf{f}_6 \oplus \mathbb{F}_p \mathbf{f}_7, \\ \mathcal{F}_2 &:= \mathbb{F}_p \mathbf{f}_1 \oplus \mathbb{F}_p \mathbf{f}_2 \oplus \mathbb{F}_p \mathbf{f}_5 \oplus \mathbb{F}_p \mathbf{f}_6. \end{aligned}$$

Then we have $\mathcal{F}_r \in C_r(\mathbb{F}_p)$ for all r . Moreover, for $r \in \{0, 1, 2\}$, put

$$\mathcal{F}_r^c := \bigoplus_{1 \leq i \leq 8, e_i \notin \mathcal{F}_r} \mathbb{F}_p \mathbf{f}_i,$$

and define an open neighborhood U_r of \mathcal{F}_r in M_{μ}^{naive} by

$$U_r(S) = \{\mathcal{F}_{r,f} \in M_{\mu, \mathbb{F}_p}^{\text{naive}} \mid f \in \text{Hom}_{\text{O}_S}(\mathcal{F}_r, \mathcal{F}_r^c)\},$$

for any \mathbb{Z}_p -scheme S , where

$$\mathcal{F}_{r,f} := \{x + f(x) \in \mathbf{\Lambda}_{0, \text{O}_S} \mid x \in \mathcal{F}_r\}.$$

Then we can represent $\mathcal{F}_{r,f}$ as a column space with respect to $\mathbf{f}_1, \dots, \mathbf{f}_8$ whose submatrix with respect to $\{i \in \{1, \dots, 8\} \mid \mathbf{f}_i \notin \mathcal{F}_r\}$ is a unit matrix of size 4.

To study U_r , we express the isotropicity condition (with respect to $[\cdot, \cdot]_0$) and the ϖ -stability condition by means of column spaces. Let

$$\begin{pmatrix} X \\ Y \end{pmatrix} \in M_{\mu}.$$

Then the isotropicity condition is equivalent to

$$(2) \quad ({}^t X \quad {}^t Y) \mathbf{J} \begin{pmatrix} X \\ Y \end{pmatrix} = 0.$$

On the other hand, the ϖ -stability condition is equivalent to

$$(3) \quad \begin{pmatrix} \varpi^2 Y \\ X \end{pmatrix} = \begin{pmatrix} X \\ Y \end{pmatrix} R$$

for some 4×4 -matrix R . Moreover, the Kottwitz condition is equivalent to the equality

$$(4) \quad \det(T - R \mid \mathcal{F}) = (T - \varpi^2)^2.$$

Second, we consider the open set U_2 . Then U_2 can be written as

$$\begin{pmatrix} E_2 & 0 \\ X_1 & X_2 \\ 0 & E_2 \\ Y_1 & Y_2 \end{pmatrix},$$

where E_2 is the unit matrix of size 2. Then the matrix R in (3) equals

$$\begin{pmatrix} 0 & \varpi^2 E_2 \\ E_2 & 0 \end{pmatrix}.$$

Note that (4) is then satisfied. Moreover, (3) amounts to saying that

$$X_1 = Y_2, \quad X_2 = \varpi^2 Y_1.$$

On the other hand, if we write $Y_k = (y_{ij}^{(k)})_{1 \leq i, j \leq 2}$ for $k \in \{1, 2\}$, then (2) is equivalent to the following equalities:

$$y_{11}^{(1)} = -y_{22}^{(1)}, \quad y_{12}^{(1)} = y_{21}^{(1)} = 0, \quad y_{11}^{(2)} = y_{22}^{(2)}, \quad y_{12}^{(2)} = y_{21}^{(2)}.$$

Therefore we have $U_2 \cong \mathbb{Z}_p[y_{11}^{(1)}, y_{12}^{(1)}, y_{11}^{(2)}, y_{12}^{(2)}]$ as desired. \blacksquare

4. DELIGNE–LUSZTIG VARIETIES FOR EVEN SPECIAL ORTHOGONAL GROUPS

In this section, we study some generalized Deligne–Lusztig varieties for even special orthogonal groups.

First, we recall the definition of generalized Deligne–Lusztig varieties. Let G_0 be a connected reductive group of \mathbb{F}_p , and put $G := G_0 \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$. We denote by Φ the Frobenius of G . Moreover, fix a Φ -stable maximal torus T of G , and a Φ -stable Borel subgroup B containing T . Let $W = W(B, T)$ be the Weyl group associated to (B, T) , and write $\Delta^* = \Delta^*(B, T)$ for the set of simple roots in W .

For $I \subset \Delta^*$, let W_I be the subgroup of W generated by I . Moreover, put $P_I := BW_I B$, which is a parabolic subgroup of G . Then there is a canonical isomorphism $\pi: P_I \backslash G / P_{\Phi(I)} \xrightarrow{\cong} W_I \backslash W / W_{\Phi(I)}$. Hence we obtain a map

$$\text{inv}: G/P_I \times G/P_{\Phi(I)} \xrightarrow{(g, g') \mapsto g^{-1}g'} P_I \backslash G / P_{\Phi(I)} \xrightarrow{\pi} W_I \backslash W / W_{\Phi(I)}.$$

Definition 4.1. Let $I \subset \Delta^*$ and $w \in W_I \backslash W / W_{\Phi(I)}$. A *generalized Deligne–Lusztig variety* for (G_0, I, w) is a locally closed subscheme of G/P_I as follows:

$$X_{P_I}(w) := \{g \in G/P_I \mid \text{inv}(g, \Phi(g)) = w\}.$$

Here we consider them for even special orthogonal groups. Let $(\Omega_{2d,0}, [,])$ be a quadratic space over \mathbb{F}_p of dimension $2d \geq 4$, and put $\Omega_{2d} := \Omega_{2d,0} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$. Fix a basis $e_1, \dots, e_d, f_1, \dots, f_d$ of Ω_{2d} satisfying $[e_i, e_j] = [f_i, f_j] = 0$, $[e_i, f_j] = \delta_{ij}$, and

$$\Phi(e_i) = \begin{cases} e_i & \text{if } 1 \leq i \leq d-1, \\ e_d & \text{if } i = d \text{ and } \Omega_{2d,0} \text{ is split,} \\ f_d & \text{if } i = d \text{ and } \Omega_{2d,0} \text{ is non-split,} \end{cases} \quad \Phi(f_i) = \begin{cases} f_i & \text{if } 1 \leq i \leq d-1, \\ f_d & \text{if } i = d \text{ and } \Omega_{2d,0} \text{ is split,} \\ e_d & \text{if } i = d \text{ and } \Omega_{2d,0} \text{ is non-split.} \end{cases}$$

Moreover, put

$$\begin{aligned} \mathcal{L}_{2d,i} &:= \overline{\mathbb{F}}_p e_1 \oplus \cdots \oplus \overline{\mathbb{F}}_p e_i \quad (i \in \{1, \dots, d-1\}), \\ \mathcal{L}_{2d,d}^+ &:= \mathcal{L}_{2d,d-1} \oplus \overline{\mathbb{F}}_p e_d, \quad \mathcal{L}_{2d,d}^- := \mathcal{L}_{2d,d-1} \oplus \overline{\mathbb{F}}_p f_d. \end{aligned}$$

Put $G_{2d,0} := \mathrm{SO}(\Omega_{2d,0})$ and $G_{2d} := G_{2d,0} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}_p}$. We regard G_{2d} as a subgroup of $\mathrm{GL}(\Omega_{2d})$ by the above basis. Furthermore, let $B_{2d} \subset G_{2d}$ be the upper-half Borel subgroup, and $T_{2d} \subset G_{2d}$ the diagonal torus. Then there is an isomorphism

$$W_{2d} := W(B_{2d}, T_{2d}) \cong \{w \in \mathfrak{A}_{2d} \mid w(i) + w(2d+1-i) = 2d+1 \text{ for all } i \in \{1, \dots, 2d\}\}.$$

Moreover, we have $\Delta_{2d}^* := \Delta^*(B_{2d}, T_{2d}) = \{s_{2d,1}, \dots, s_{2d,d-2}, t_{2d}^+, t_{2d}^-\}$, where

- $s_{2d,i} = (i \ i+1)(2d-i \ 2d+1-i)$ for $i \in \{1, \dots, d-2\}$,
- $t_{2d}^+ = (d-1 \ d)(d+1 \ d+2)$,
- $t_{2d}^- = (d-1 \ d+1)(d \ d+2)$.

Note that we have $\Phi(s_{2d,i}) = s_{2d,i}$ for any i and

$$\Phi(t_{2d}^\pm) = \begin{cases} t_{2d}^\pm & \text{if } \Omega_{2d,0} \text{ is split,} \\ t_{2d}^\mp & \text{if } \Omega_{2d,0} \text{ is non-split.} \end{cases}$$

Definition 4.2. For a quadratic space Ω_0 over \mathbb{F}_p of dimension n , we denote by $\mathrm{OGr}(\Omega_0)_{\overline{\mathbb{F}_p}}$ be the moduli space of maximal totally isotropic subspaces of $\Omega_0 \otimes_{\mathbb{F}_p} \overline{\mathbb{F}_p}$. Note that all maximal totally isotropic subspaces of $\Omega_0 \otimes_{\mathbb{F}_p} \overline{\mathbb{F}_p}$ are $[n/2]$ -dimensional, where $[n/2] := \max\{n' \in \mathbb{Z} \mid n' \leq n/2\}$. Moreover, put

$$S_{\Omega_0, \overline{\mathbb{F}_p}} := \{\mathcal{L} \in \mathrm{OGr}(\Omega_0)_{\overline{\mathbb{F}_p}} \mid \mathrm{rank}(\mathcal{L} \cap (\mathrm{id} \otimes \sigma)_*(\mathcal{L})) \geq m\},$$

where

$$m := \begin{cases} \frac{n}{2} - 2 & \text{if } n \in 2\mathbb{Z} \text{ and } \Omega_0 \text{ is split,} \\ \frac{n}{2} - 1 & \text{if } n \in 2\mathbb{Z} \text{ and } \Omega_0 \text{ is non-split,} \\ \frac{n-1}{2} & \text{if } n \in \mathbb{Z} \setminus 2\mathbb{Z}, \end{cases}$$

Here $(\mathrm{id} \otimes \sigma)_*(\mathcal{L})$ is the submodule generated by $(\mathrm{id} \otimes \sigma)(\mathcal{L})$. It is a closed subscheme of $\mathrm{OGr}(\Omega_0)_{\overline{\mathbb{F}_p}}$ by definition.

We introduce subschemes of $\mathrm{OGr}(\Omega_0)_{\overline{\mathbb{F}_p}}$ and $S_{\Omega_0, \overline{\mathbb{F}_p}}$ in the case $n = 2d$, that is, $\Omega_0 = \Omega_{2d,0}$. Let $P_{2d,0}^+$ and $P_{2d,0}^-$ be the stabilizers of $\mathcal{L}_{2d,d}^+$ and $\mathcal{L}_{2d,d}^-$ in G_{2d} respectively, which are parabolic subgroups of G_{2d} . Then the maps $g \mapsto g(\mathcal{L}_{2d,d}^+)$ and $g \mapsto g(\mathcal{L}_{2d,d}^-)$ induce an isomorphism

$$(G_{2d}/P_{2d,0}^+) \sqcup (G_{2d}/P_{2d,0}^-) \cong \mathrm{OGr}(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}.$$

We denote by $\mathrm{OGr}^\pm(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}$ the image of $G_{2d}/P_{2d,0}^\pm$ under the isomorphism above. Note that $\mathrm{OGr}^\pm(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}$ are the connected components of $\mathrm{OGr}(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}$. Then, for two elements $\mathcal{L}, \mathcal{L}'$ in $\mathrm{OGr}(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}(k)$ where k is an algebraic closed field extension over $\overline{\mathbb{F}_p}$, \mathcal{L} and \mathcal{L}' are contained in the same connected component if and only if $\dim_k(\mathcal{L}/\mathcal{L} \cap \mathcal{L}') = \dim_k(\mathcal{L}'/\mathcal{L} \cap \mathcal{L}') \in 2\mathbb{Z}$. On the other hand, put $S_{\Omega_{2d,0}}^\pm := S_{\Omega_{2d,0}} \cap \mathrm{OGr}^\pm(\Omega_{2d,0})_{\overline{\mathbb{F}_p}}$.

Now we give a connection between $S_{\Omega_{2d,0}}$ and generalized Deligne–Lusztig varieties for $G_{2d,0}$.

Case 1. The case for $\Omega_{2d,0}$ split. For $i \in \mathbb{Z}_{\geq 0}$, put

$$P_{2d,i}^\pm := \begin{cases} P_{\{s_{2d,1}, \dots, s_{2d,d-2-i}, t_{2d}^\pm\}} & \text{if } i \leq d-3, \\ P_{\{t_{2d}^\pm\}} & \text{otherwise.} \end{cases}$$

Then the following hold:

- $P_{2d,0}^\pm$ is the stabilizer of $\mathcal{L}_{2d,d}^\pm$,
- $P_{2d,i}^\pm$ is the stabilizer of the flag $\mathcal{L}_{2d,d-1-i} \subset \dots \subset \mathcal{L}_{2d,d-2} \subset \mathcal{L}_{2d,d}^\pm$ if $1 \leq i \leq d-3$,
- $P_{2d,i}^\pm$ is the stabilizer of the flag $\mathcal{L}_{2d,1} \subset \dots \subset \mathcal{L}_{2d,d-2} \subset \mathcal{L}_{2d,d}^\pm$ if $i \geq d-2$.

Proposition 4.3.(i) *There is a locally closed stratification*

$$S_{\Omega_{2d,0}}^{\pm} = X_{P_{2d,0}^{\pm}}(\text{id}) \sqcup X_{P_{2d,0}^{\pm}}(t_{2d}^{\mp}).$$

Moreover, the closure of $X_{P_{2d,0}^{\pm}}(t_{2d}^{\mp})$ in $G_{2d}/P_{2d,0}^{\pm}$ equals $S_{\Omega_{2d,0}}^{\pm}$.

(ii) *The schemes $S_{\Omega_{2d,0}}^{\pm}$ are irreducible.*

Proof. (i): The proof is the same as [RTW14, Proposition 5.3] or [HP14, Lemma 3.7].

(ii): It suffices to prove the irreducibility of $X_{P_{2d,0}^{\pm}}(t_{2d}^{\mp})$ by (i). However, this follows from [BR06, Theorem 2]. ■

Proposition 4.4. *There is a locally closed stratification*

$$S_{\Omega_{6,0}}^{\pm} = X_{P_{6,0}^{\pm}}(\text{id}) \sqcup X_{P_{6,1}^{\pm}}(t_6^{\mp}) \sqcup X_{P_{6,1}^{\pm}}(t_6^{\mp} s_{6,1}).$$

More precisely, there are locally closed subschemes X_0^{\pm} and X_1^{\pm} of $S_{\Omega_{6,0}}^{\pm}$ satisfying $X_0^{\pm} \cong X_{P_{6,1}^{\pm}}(t_6^{\mp})$ and $X_1^{\pm} \cong X_{P_{6,1}^{\pm}}(t_6^{\mp} s_{6,1})$ under the canonical morphism $G_6/P_{6,0}^{\pm} \rightarrow G_6/P_{6,1}^{\pm}$. Furthermore, we have

$$\overline{X_{P_{6,1}^{\pm}}(t_6^{\mp})} = X_{P_{6,0}^{\pm}}(\text{id}) \sqcup X_{P_{6,1}^{\pm}}(t_6^{\mp}), \quad \overline{X_{P_{6,1}^{\pm}}(t_6^{\mp} s_1)} = S_{\Omega_{6,0}}^{\pm}.$$

Here we denote by \overline{X} the closure of a subset X of $G_6/P_{6,1}^{\pm}$.

Proof. This follows from the same argument as the proof of [RTW14, Proposition 5.5] by using Proposition 4.3. ■

Remark 4.5. If $d \geq 4$, then we have

$$S_{\Omega_{2d,0}}^{\pm} = X_{P_{2d,0}^{\pm}}(\text{id}) \sqcup X_{P_{2d,1}^{\pm}}(t_{2d}^{\mp}) \sqcup X_{P_{2d,1}^{\pm}}(t_{2d}^{\mp} s_{2d,d-2}) \sqcup X_{P_{2d,1}^{\pm}}(s_{2d,d-2} s_{2d,d-3} t_{2d}^{\mp} s_{2d,d-2}).$$

The variety $X_{P_{2d,1}^{\pm}}(s_{2d,d-2} s_{2d,d-3} t_{2d}^{\mp} s_{2d,d-2})$ parametrizes all flags $\mathcal{L}_{d-2} \subset \mathcal{L}_d$ in Ω_{2d} where $\text{rank } \mathcal{L}_i = i$ satisfying $\mathcal{L}_{d-2} = \mathcal{L}_d \cap \Phi(\mathcal{L}_d)$ and $\text{rank}(\mathcal{L}_{d-2}/\mathcal{L}_{d-2} \cap \Phi(\mathcal{L}_{d-2})) = 2$.

Next, we give a connection between generalized Deligne–Lusztig varieties for SO_{2d-1} . Let $\omega_d := e_d - f_d \in \Omega_{2d,0}$, which is an anisotropic vector. We denote by $\Omega_{2d-1,0}$ the perpendicular of ω_d in $\Omega_{2d,0}$. Then we have $\dim_{\mathbb{F}_p}(\Omega_{2d-1,0}) = 2d - 1$. Moreover, for $i \in \{1, \dots, d-1\}$, put

$$\mathcal{L}_{2d-1,i} := \mathcal{L}_{2d,i+1} \cap \Omega_{2d-1}.$$

On the other hand, put $G_{2d-1,0} := \text{SO}(\Omega_{2d-1,0})$ and $G_{2d-1} := G_{2d-1,0} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$. We regard G_{2d-1} as a subgroup of GL_{2d-1} by the basis $e_1, \dots, e_{d-1}, e_d + f_d, f_{d-1}, \dots, f_1$. Moreover, let $B_{2d-1} \subset G_{2d-1}$ be the upper-half Borel subgroup, and $T_{2d-1} \subset G_{2d-1}$ the diagonal torus. Then there is an isomorphism

$$W_{2d-1} := W(B_{2d-1}, T_{2d-1}) \cong \{w \in \mathfrak{S}_{2d-1} \mid w(i) + w(2d-i) = 2d \text{ for all } i \in \{1, \dots, 2d-1\}\}.$$

Moreover, we have $\Delta_{2d-1}^* := \Delta^*(B_{2d-1}, T_{2d-1}) = \{s_{2d-1,1}, \dots, s_{2d-1,d-2}, t_{2d-1}\}$, where

- $s_{2d-1,i} = (i \ i+1)(2d-i \ 2d+1-i)$ for $i \in \{1, \dots, d-2\}$,
- $t_{2d-1} = (d-1 \ d+1)$.

For $i \in \mathbb{Z}_{\geq 0}$, put

$$P_{2d-1,i} := \begin{cases} P_{\{s_{2d-1,1}, \dots, s_{2d-1,d-2-i}\}} & \text{if } i \leq d-3, \\ B_{2d-1} & \text{otherwise.} \end{cases}$$

Then $P_{2d-1, i}$ is the stabilizer of the flag $\mathcal{L}_{2d-1, d-1-i} \subset \cdots \subset \mathcal{L}_{2d-1, d-1}$. Moreover, there is an isomorphism

$$G_{2d-1}/P_{2d-1, 0} \xrightarrow{\cong} \mathrm{OGr}(\Omega_{2d-1, 0})_{\mathbb{F}_p}; g \mapsto g(\mathcal{L}_{2d-1, d-1}).$$

Lemma 4.6. *There is a commutative diagram*

$$\begin{array}{ccc} G_{2d}/P_{2d, 0}^{\pm} & \xrightarrow{g \mapsto g(\mathcal{L}_{2d, d}^{\pm})} & \mathrm{OGr}^{\pm}(\Omega_{2d, 0})_{\mathbb{F}_p} \\ \downarrow g \mapsto g|_{\Omega_{2d-1}} & & \downarrow \mathcal{L} \mapsto \mathcal{L} \cap \Omega_{2d-1} \\ G_{2d-1}/P_{2d-1, 0} & \xrightarrow{g \mapsto g(\mathcal{L}_{2d-1, d-1})} & \mathrm{OGr}(\Omega_{2d-1, 0})_{\mathbb{F}_p}, \end{array}$$

where the all maps are isomorphisms.

Proof. The commutativity of diagram is clear. The isomorphy of the horizontal maps are already mentioned. On the other hand, [HP14, Lemma 3.4] and the definition of $\mathrm{OGr}^{\pm}(\Omega_{2d, 0})_{\mathbb{F}_p}$ imply the isomorphy of the vertical map in the right-hand side. Hence the isomorphy of the remaining map follows. \blacksquare

Proposition 4.7.

(i) *For $d \leq 3$, the isomorphism in Lemma 4.6 induces isomorphisms*

$$S_{\Omega_{2d, 0}}^{\pm} \cong S_{\Omega_{2d-1, 0}}, \quad X_{P_{2d, 0}^{\pm}}(\mathrm{id}) \cong X_{P_{2d-1, 0}}(\mathrm{id}), \quad X_{P_{2d, 0}^{\pm}}(t_{2d}^{\mp}) \cong X_{P_{2d-1, 0}}(t_{2d-1}).$$

(ii) *The isomorphism $X_{P_{6, 0}^{\pm}}(t_6^{\mp}) \cong X_{P_{5, 0}}(t_5)$ in (i) induces isomorphisms*

$$X_{P_{6, 1}^{\pm}}(t_6^{\pm}) \cong X_{P_{5, 1}}(t_5), \quad X_{P_{6, 1}^{\pm}}(t_6^{\mp} s_{6, 1}) \cong X_{B_5}(t_5).$$

Proof. (i): First, we prove $S_{\Omega_{2d, 0}}^{\pm} \cong S_{\Omega_{2d-1, 0}}$. It is clear in the case $d = 2$. If $d = 3$, then we obtain a closed immersion $S_{\Omega_{2d-1, 0}} \rightarrow S_{\Omega_{2d, 0}}^{\pm}$. On the other hand, we have $\dim S_{\Omega_{2d-1, 0}} = 2$ by [Oki19, Propositions 6.5], and $S_{\Omega_{2d, 0}}^{\pm}$ is irreducible of dimension ≤ 2 by Proposition 4.3 (ii). Hence the above map $S_{\Omega_{2d-1, 0}} \rightarrow S_{\Omega_{2d, 0}}^{\pm}$ is an isomorphism.

Next, we prove the remaining isomorphisms. By [Oki19, Proposition 6.4 (ii)], there is a locally closed stratification

$$S_{\Omega_{2d, 0}}^{\pm} = X_{P_{2d, 0}^{\pm}}(\mathrm{id}) \sqcup X_{P_{2d, 0}^{\pm}}(t_d^{\mp}),$$

Hence it suffices to prove the isomorphism $X_{P_{2d, 0}^{\pm}}(\mathrm{id}) \cong X_{P_{2d-1, 0}}(\mathrm{id})$. However, this follows from the definitions of $X_{P_{2d-1, 0}}(\mathrm{id})$ and $\mathrm{OGr}^{\pm}(\Omega_{2d, 0})_{\mathbb{F}_p}$.

(ii): Set

$$\begin{aligned} S_{\Omega_{6, 0}}^{1, \pm} &:= \{(\mathcal{L}_1, \mathcal{L}_3) \in \mathbb{P}(\Omega_{6, 0})_{\mathbb{F}_p} \times \mathrm{OGr}^{\pm}(\Omega_{6, 0})_{\mathbb{F}_p} \mid \mathcal{L}_1 = \mathcal{L}_3 \cap \Phi(\mathcal{L}_3)\}, \\ S_{\Omega_{5, 0}}^1 &:= \{(\mathcal{L}'_1, \mathcal{L}'_2) \in \mathbb{P}(\Omega_{5, 0})_{\mathbb{F}_p} \times \mathrm{OGr}(\Omega_{5, 0})_{\mathbb{F}_p} \mid \mathcal{L}'_1 = \mathcal{L}'_2 \cap \Phi(\mathcal{L}'_2)\}. \end{aligned}$$

By the definitions of $X_{P_{6, 0}^{\pm}}(t_6^{\mp})$ and $X_{P_{5, 0}}(t_5)$, we obtain an isomorphism

$$S_{\Omega_{6, 0}}^{1, \pm} \cong X_{P_{6, 0}^{\pm}}(t_6^{\mp}) \cong X_{P_{5, 0}}(t_5) \cong S_{\Omega_{5, 0}}^1.$$

This isomorphism is equal to $(\mathcal{L}_1, \mathcal{L}_3) \mapsto (\mathcal{L}_1 \cap \Omega_5, \mathcal{L}_3 \cap \Omega_5)$, and hence we obtain $\mathcal{L}_1 \cap \Omega_5 = \mathcal{L}_1$. Therefore, for $(\mathcal{L}_1, \mathcal{L}_3) \in S_{\Omega_{6, 0}}^{1, \pm}$, we have $\Phi(\mathcal{L}_1) = \mathcal{L}_1$ if and only if $\Phi(\mathcal{L}_1 \cap \Omega_5) = \mathcal{L}_1 \cap \Omega_5$. This equivalence implies the desired isomorphisms. \blacksquare

Case 2. The case for $\Omega_{2d,0}$ non-split. For $i \in \mathbb{Z}_{\geq 0}$, put

$$P_{2d,i} := \begin{cases} P_{\{s_{2d,1}, \dots, s_{2d,d-2-i}\}} & \text{if } i \leq d-3, \\ B_{2d-1} & \text{otherwise.} \end{cases}$$

Then the following hold:

- $P_{2d,i}$ is the stabilizer of the flag $\mathcal{L}_{2d,d-1-i} \subset \dots \subset \mathcal{L}_{2d,d-1}$ if $0 \leq i \leq d-3$,
- $P_{2d,i}^\pm$ is the stabilizer of the flag $\mathcal{L}_{2d,1} \subset \dots \subset \mathcal{L}_{2d,d-1}$ if $i \geq d-2$.

Proposition 4.8. ([HP14, Proposition 3.8]) *There is a locally closed stratification*

$$S_{\Omega_{2d,0}} = \coprod_{i=0}^{d-1} (X_{P_{2d,i}}(t_{2d}^+ s_{2d,d-2} \cdots s_{2d,d-i}) \sqcup X_{P_{2d,i}}(t_{2d}^- s_{2d,d-2} \cdots s_{2d,d-i})).$$

Moreover, for any $r \in \{0, \dots, d-1\}$, the closure of $X_{P_{2d,r}}(t_{2d}^\pm s_{2d,d-2} \cdots s_{2d,d-r})$ in $G_{2d}/P_{2d,0}$ equals

$$\prod_{i=0}^r X_{P_{2d,i}}(t_{2d}^+ s_{2d,d-2} \cdots s_{2d,d-i}).$$

5. BRUHAT–TITS STRATIFICATION IN THE NEUTRAL CASE, I

In this section, we describe the structure of $\mathcal{M}_{b,\mu}^{(0)}$ for a μ -neutral b by means of certain 6-dimensional quadratic space over \mathbb{Q}_p .

5.1. Preliminaries. Here we recall the basic notions on Clifford algebras and spinor similitude groups.

Let L a quadratic space over a ring A . We denote by $C(L)$ the Clifford algebra of L . Then there is a $\mathbb{Z}/2$ -grading

$$C(L) = C^+(L) \oplus C^-(L),$$

where $C^+(L)$ is the 0-th part, which is an A -subalgebra of $C(L)$. The ring $C^+(L)$ is called the even Clifford algebra.

We regard L as an A -submodule of $C^-(L)$. Let $v \mapsto v'$ be the involution on $C(L)$ satisfying $(v_1 \cdots v_d)' = v_d \cdots v_1$ for $v_1, \dots, v_d \in L$. Then we define an affine group scheme $\mathrm{GSpin}(L)$ over A by

$$\mathrm{GSpin}(L)(R) = \{g \in C^+(L \otimes_A R)^\times \mid g(L \otimes_A R)g^{-1} = L \otimes_A R, g'g \in R^\times\}$$

for any A -algebra R . Moreover, denote by $\mathrm{Spin}(L)$ the derived group of $\mathrm{GSpin}(L)$.

If $R = k$ is a field, we define an action of $\mathrm{GSpin}(L)(k)$ on L by the exact sequence

$$1 \rightarrow \mathbb{G}_m \rightarrow \mathrm{GSpin}(L) \rightarrow \mathrm{SO}(L) \rightarrow 1,$$

where $\mathrm{GSpin}(L) \rightarrow \mathrm{SO}(L)$ is defined by $g \mapsto [v \mapsto gvg^{-1}]$. In particular, the homomorphism $\mathrm{GSpin}(L)(k) \rightarrow \mathrm{SO}(L)(k)$ is surjective by Hilbert Satz 90.

On the other hand, the above exact sequence induces another exact sequence

$$1 \rightarrow \mu_2 \rightarrow \mathrm{Spin}(L) \rightarrow \mathrm{SO}(L) \rightarrow 1.$$

Taking the Galois cohomology over k , we obtain an exact sequence of groups

$$1 \rightarrow \mu_2(k) \rightarrow \mathrm{Spin}(L)(k) \rightarrow \mathrm{SO}(L)(k) \xrightarrow{\theta_L} k^\times / (k^\times)^2 \rightarrow H^1(k, \mathrm{Spin}(L)).$$

We call the homomorphism θ_L as the *spinor norm* of L . Note that all exact sequences as appeared in this section are functorial with respect to homomorphisms of quadratic spaces.

Proposition 5.1. ([Kne65]) *If k is a p -adic field and $\dim_k(L) \geq 3$, then $H^1(k, \text{Spin}(L)) = 1$. In particular, the spinor norm θ_L of L is surjective.*

5.2. Hodge star operator. Here we will define a quadratic space $\mathbf{L}_{b,0}$ over \mathbb{Q}_p which will be used to describe $\mathcal{M}_{b,\mu}^{(0)}$.

Until Section 5.8, we fix $\varepsilon \in W^\times$ satisfying $\sigma(\varepsilon) = -\varepsilon$.

Lemma 5.2. *There is an \check{F} -basis e_1, e_2, e_3, e_4 such that*

$$\mathbf{F}(e_1) = \varpi e_3, \quad \mathbf{F}(e_2) = \varpi e_4, \quad \mathbf{F}(e_3) = p\varpi^{-1}e_1, \quad \mathbf{F}(e_4) = p\varpi^{-1}e_2.$$

and

$$\langle e_i, e_j \rangle = \begin{cases} \varpi^{-1} & \text{if } (i, j) = (1, 3), \\ \varpi & \text{if } (i, j) = (2, 4), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Put

$$b' := \begin{pmatrix} & & & -p \\ & & p & \\ & 1 & & \\ -1 & & & \end{pmatrix}.$$

Then we have $\det_{\check{F}}(b') / \text{sim}_G(b')^2 = 1$, which implies that $[b']$ is μ -neutral by Proposition 2.10. Hence we may assume $b = b'$. Then

$$e_1 := \mathbf{e}_1, \quad e_2 := \varpi \mathbf{e}_2, \quad e_3 := -\varpi^{-1} \mathbf{e}_4, \quad e_4 := \mathbf{e}_3$$

gives a desired \check{F} -basis. ■

From now on, replacing $(\mathbb{X}_0, \iota_0, \lambda_0)$ to (X, ι, λ) satisfying $(X, \iota, \lambda, \rho) \in \mathcal{M}_{b,\mu}^{(0)}$ if necessary, we assume

$$M_0 = O_{\check{F}}(\varpi e_1) \oplus O_{\check{F}}e_2 \oplus O_{\check{F}}(\varpi e_3) \oplus O_{\check{F}}e_4.$$

Definition 5.3. Put

$$\omega := e_1 \wedge e_2 \wedge e_3 \wedge e_4 \in \bigwedge_{\check{F}}^4 N_b,$$

where e_1, \dots, e_4 are as in Lemma 5.2. Then we define a *Hodge star operator* $v \mapsto v^*$ on $\bigwedge_{\check{F}}^2 N_b$ (with respect to ω) by $v' \wedge v^* = \langle v', v \rangle_2 \omega$ for any $v' \in \bigwedge_{\check{F}}^2 N_b$. Here $\langle \cdot, \cdot \rangle_2$ is the bilinear form on $\bigwedge_{\check{F}}^2 N_b$ satisfying

$$\langle x_1 \wedge x_2, y_1 \wedge y_2 \rangle_2 = \langle x_1, y_1 \rangle \langle x_2, y_2 \rangle - \langle x_1, y_2 \rangle \langle x_2, y_1 \rangle$$

for any $x_1, x_2, y_1, y_2 \in N_b$.

By definition, we have $(av)^* = \bar{a}v^*$ and $(v^*)^* = v$ for any $a \in \check{F}$ and $v \in \bigwedge_{\check{F}}^2 N_b$. Now put

$$\mathbf{L}_b := \left\{ v \in \bigwedge_{\check{F}}^2 N_b \mid v^* = v \right\}.$$

Then it is a K_0 -vector space of dimension 6 which is spanned by the following elements:

$$\begin{aligned} x_1 &:= e_1 \wedge e_2, & x_2 &:= e_3 \wedge e_4, \\ x_3 &:= e_1 \wedge e_3 - \varpi^{-2} e_2 \wedge e_4, & x_4 &:= \varpi e_1 \wedge e_3 + \varpi^{-1} e_2 \wedge e_4, \\ x_5 &:= e_1 \wedge e_4, & x_6 &:= e_3 \wedge e_2. \end{aligned}$$

Note that \mathbf{L}_b depends on the choice of e_1, \dots, e_4 , cf. [HP14, Section 2.2].

We regard $\bigwedge_{\check{F}}^2 N_b$ as a K_0 -subspace of $\text{End}_{K_0}(N_b)$ by the injection

$$\bigwedge_{\check{F}}^2 N_b \rightarrow \text{End}_{K_0}(N_b); x \wedge y \mapsto [z \mapsto \langle x, z \rangle y - \langle y, z \rangle x].$$

Note that the image of the above injection consists of K_0 -endomorphisms f of N_b such that $f(ax) = \bar{a}f(x)$ for $a \in \check{F}$ and $x \in N_b$.

We will frequently use the following relation between $v \mapsto v'$ on $C(\mathbf{L}_b)$ and bilinear forms on N_b :

Lemma 5.4. *Let $v \in \mathbf{L}_b$. Then we have $\langle vx, y \rangle = -\overline{\langle x, vy \rangle}$ and $(vx, y) = (x, vy)$ for any $x, y \in N_b$.*

Proof. The proof is the same as [Oki19, Lemma 3.6]. ■

Proposition 5.5.

(i) *For $v \in \mathbf{L}_b$, as an element in $\text{End}_{K_0}(N_b)$, we have*

$$v^2 = \langle v, v \rangle.$$

In particular, the map $Q: \mathbf{L}_b \rightarrow \text{End}_{K_0}(N_b); v \mapsto v^2$ is K_0 -valued quadratic form. We regard \mathbf{L}_b as a quadratic space over K_0 by Q , and put

$$[,]: \mathbf{L}_b \rightarrow \text{End}_{K_0}(N_b); (v, w) \mapsto \frac{1}{2}(v \circ w + w \circ v) = \frac{1}{2}(Q(v+w) - Q(v) - Q(w)).$$

(ii) *The W -submodule $L_0 := \{v \in \mathbf{L}_b \mid v(M_0) \subset M_0\}$ of \mathbf{L}_b is a W -lattice. Moreover, we have*

$$L_0^\vee = \{v \in \mathbf{L}_b \mid v(M_0) \subset \varpi^{-1}M_0\}, \quad \text{length}_W(L_0^\vee/L_0) = 1.$$

(iii) *The inclusion $L_0 \subset \text{End}_W(M_0)$ induces isomorphisms*

$$C(L_0) \cong \text{End}_W(M_0), \quad C^+(L_0) \cong \text{End}_{O_{\check{F}}}(M_0).$$

Proof. (i): This follows from the same argument as the proof of [HP14, Proposition 2.4 (2)] by considering $e_i \wedge e_j$ for all i, j .

(ii): By computation, the Gram matrix of $[,]$ with respect to the basis x_1, \dots, x_6 as above is as follows:

$$\begin{pmatrix} 0 & -\frac{1}{2} & & & & \\ -\frac{1}{2} & 0 & & & & \\ & & -\varpi^{-2} & 0 & & \\ & & 0 & 1 & & \\ & & & & -\frac{1}{2} & 0 \\ & & & & 0 & -\frac{1}{2} \end{pmatrix}.$$

Moreover, we have

$$L_0 = Wx_1 \oplus Wx_2 \oplus W(\varpi^2 x_3) \oplus Wx_4 \oplus Wx_5 \oplus Wx_6$$

by definition. Hence L_0 is a W -lattice in \mathbf{L}_b . Moreover, we have $L_0 \subset L_0^\vee$ and $\mathrm{length}_W(L_0^\vee/L_0) = 1$. On the other hand, if we put

$$L'_0 := \{v \in \mathbf{L}_b \mid v(M_0) \subset \varpi^{-1}M_0\},$$

then we have

$$L'_0 = Wx_1 \oplus Wx_2 \oplus Wx_3 \oplus Wx_4 \oplus Wx_5 \oplus Wx_6$$

and $L'_0 \subset L_0^\vee$ by definition. Since $\mathrm{length}_W(L_0^\vee/L_0) = 1$ and $L_0 \subsetneq L'_0$, we obtain the equality $L'_0 = L_0^\vee$ as desired.

(iii): This follows from the same argument as the proof of [HP14, Proposition 2.4 (4)]. \blacksquare

We define a σ -linear bijection

$$\Phi_b: \mathrm{End}_{K_0}(N_b) \rightarrow \mathrm{End}_{K_0}(N_b); f \mapsto \mathbf{F}_b \circ f \circ \mathbf{F}_b^{-1}.$$

Then the isocrystal $(\mathrm{End}_{K_0}(N_b), \Phi_b)$ over $\overline{\mathbb{F}}_p$ is isoclinic of slope 0. Moreover, there is an isomorphism $\mathrm{End}(N_b)^{\Phi_b} \cong \mathrm{End}^0(\mathbb{X}_0)$. On the other hand, we have

$$\Phi_b(x \wedge y) = p^{-1}\mathbf{F}_b(x) \wedge \mathbf{F}_b(y)$$

under the inclusion $\bigwedge_{\overline{\mathbb{F}}_p}^2 N_b \subset \mathrm{End}_{K_0}(N_b)$. We can prove that Φ_b commutes with \star by the same argument as [Oki19, Lemma 3.9]. Hence we obtain an isocrystal (\mathbf{L}_b, Φ_b) over $\overline{\mathbb{F}}_p$ which is isoclinic of slope 0. We denote by $\mathbf{L}_{b,0}$ the Φ_b -fixed part of \mathbf{L}_b . Then $(\mathbf{L}_{b,0}, Q)$ becomes a quadratic space of dimension 6 over \mathbb{Q}_p .

We compute a basis of $\mathbf{L}_{b,0}$ explicitly. The \mathbb{Q}_p -vector space $\mathbf{L}_{b,0}$ is spanned by the following elements:

$$\begin{aligned} y_1 &:= x_1 + p^{-1}\varpi^2 x_2, & y_2 &:= \varepsilon(x_1 - p^{-1}\varpi^2 x_2), \\ y_3 &:= \varepsilon x_3, & y_4 &:= \varepsilon x_4, \\ y_5 &:= x_5 + x_6, & y_6 &:= \varepsilon(x_5 - x_6). \end{aligned}$$

The Gram matrix of $[\cdot, \cdot]$ with respect to the basis y_1, \dots, y_6 is as follows:

$$\begin{pmatrix} -p^{-1}\varpi^2 & & & & & \\ & p^{-1}\varpi^2\varepsilon^2 & & & & \\ & & -\varpi^{-2}\varepsilon^2 & & & \\ & & & \varepsilon^2 & & \\ & & & & -1 & \\ & & & & & \varepsilon^2 \end{pmatrix}.$$

Hence the discriminant and the Hasse invariant of $(\mathbf{L}_{b,0}, [\cdot, \cdot])$ are $-\varpi^2$ and -1 respectively.

We will use the following in Section 5.6:

Lemma 5.6. *We have $\iota_0(\varpi) = -p\varepsilon^4 y_1 \cdots y_6$ in $C(\mathbf{L}_b) \cong \mathrm{End}_{K_0}(N_b)$.*

Proof. This follows from direct computation. \blacksquare

5.3. Exceptional isomorphism. We define a left action of $G(K_0)$ on $\mathrm{End}_{K_0}(N_b)$ by

$$G(K_0) \times \mathrm{End}_{K_0}(N_b) \rightarrow \mathrm{End}_{K_0}(N_b); (g, f) \mapsto g \cdot f := g \circ f \circ g^{-1}.$$

Then we have

$$(5) \quad g \cdot (x \wedge y) = \mathrm{sim}_G(g)^{-1}g(x) \wedge g(y)$$

for any $g \in G$ by the inclusion $\bigwedge_F^2 N_b$. On the other hand, let

$$G^0 := \{g \in G \mid \text{sim}_G(g)^2 = \det_F(g)\}.$$

Then, for $g \in G(K_0)$, the action $v \mapsto g \cdot v$ on $\bigwedge_F^2 N_b$ commutes with \star if and only if $g \in G^0(K_0)$. See [HP14, Section 2.3].

Proposition 5.7.

(i) *The isomorphism $C^+(\mathbf{L}_b) \cong \text{End}_{\mathbb{F}}(N_b)$ in Proposition 5.5 (iii) induces an isomorphism*

$$\text{GSpin}(\mathbf{L}_b) \cong G^0 \otimes_{\mathbb{Q}_p} K_0,$$

which is compatible with the actions on \mathbf{L}_b .

(ii) *For any algebraic closed field k of characteristic p , the isomorphism in (i) induces an isomorphism*

$$\text{Stab}_{\text{GSpin}(\mathbf{L}_b)(\text{Frac } W(k))}(L_{0,W(k)}) \cong \text{Stab}_{G^0(\text{Frac } W(k))}(M_{0,W(k)}).$$

Proof. (i): This follows from the same argument as the proof of [HP14, Proposition 2.7] by using Lemma 5.4.

(ii): This follows from (i) and the definition of L_0 . ■

We denote by J_b^{der} the derived group of J_b , and put

$$J_b^0 := J_b \cap \text{Res}_{K_0/\mathbb{Q}_p} G^0.$$

Corollary 5.8. *The isomorphism $\text{GSpin}(\mathbf{L}_b) \cong G^0$ in Proposition 5.7 induces an isomorphism*

$$\text{GSpin}(\mathbf{L}_{b,0}) \cong J_b^0$$

and a central isogeny

$$J_b^{\text{der}} \rightarrow \text{SO}(\mathbf{L}_{b,0}),$$

which are compatible with the actions on $\mathbf{L}_{b,0}$.

Proof. This follows from Proposition 5.7 (i). ■

5.4. Description of a subset of field valued points. Here we fix an algebraically closed field k of characteristic p . We set $M_{0,W(k)} := M_0 \otimes_W W(k)$.

Definition 5.9. For $j \in \mathbb{Z}/2$, put

$$\text{DiL}_{M_0,j}^{\varpi}(N_{b,W(k)}) := \{M \in \text{DiL}^{\varpi}(N_{b,W(k)}) \mid \text{length}_{O_{F,W(k)}}(M + M_{0,W(k)})/M_{0,W(k)} \equiv j \pmod{2}\}.$$

Moreover, let $\mathcal{M}_{b,\mu}^{(0,j)}(k)$ be the subset of $\mathcal{M}_{b,\mu}^{(0)}(k)$ corresponding to $\text{DiL}_{M_0,j}^{\varpi}(N_{b,W(k)})$ under the bijection $\mathcal{M}_{b,\mu}^{(0)} \cong \text{DiL}^{\varpi}(N_{b,W(k)})$ in Proposition 2.5.

By definition, we have

$$\mathcal{M}_{b,\mu}^{(0)}(k) = \mathcal{M}_{b,\mu}^{(0,0)}(k) \sqcup \mathcal{M}_{b,\mu}^{(0,1)}(k).$$

Moreover, the action of $s \in J_b(\mathbb{Q}_p)$ satisfying $\kappa_G(s) = (0, 1)$ induces a bijection

$$\mathcal{M}_{b,\mu}^{(0,0)}(k) \cong \mathcal{M}_{b,\mu}^{(0,1)}(k).$$

Note that $\mathcal{M}_{b,\mu}^{(0,j)}(k)$ is stable under field extension, that is, for an algebraically closed field k' containing k and $j \in \mathbb{Z}/2$, the image of the canonical map $\mathcal{M}_{b,\mu}^{(0,j)}(k) \rightarrow \mathcal{M}_{b,\mu}^{(0)}(k')$ lies in $\mathcal{M}_{b,\mu}^{(0,j)}(k')$.

Here we will describe the set $\mathcal{M}_{b,\mu}^{(0,0)}(k)$ by means of certain lattices in $\mathbf{L}_{b,W(k)} := \mathbf{L}_b \otimes_W W(k)$.

Definition 5.10. A special lattice in $\mathbf{L}_{b,W(k)}$ is a $W(k)$ -lattice L in $\mathbf{L}_{b,W(k)}$ satisfying

$$\mathrm{length}_{W(k)} L + \Phi_b(L)/L \leq 1.$$

We denote by $\mathrm{SpL}(\mathbf{L}_{b,W(k)})$ the set of all special lattices in $\mathbf{L}_{b,W(k)}$. We prove the following:

Proposition 5.11. For any $M \in \mathrm{DiL}_{M_0,0}^{\varpi}(N_{b,W(k)})$,

$$L(M) := \{v \in \mathbf{L}_{b,W(k)} \mid v(\mathbf{V}_b(M)) \subset \varpi^{-1}\mathbf{V}_b(M)\}$$

is an element of $\mathrm{SpL}(\mathbf{L}_{b,W(k)})$. Moreover, the map

$$\mathrm{DiL}_{M_0,0}^{\varpi}(N_{b,W(k)}) \rightarrow \mathrm{SpL}(\mathbf{L}_{b,W(k)}); M \mapsto L(M)$$

is a $J_b^{\mathrm{der}}(\mathbb{Q}_p)$ -equivariant bijection.

We give a proof of Proposition 5.11. Let $\mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)})$ be the set of ϖ -modular $O_{F,W(k)}$ -lattices in $N_{b,W(k)}$ satisfying $\mathrm{length}_{O_{F,W(k)}}(M + M_{0,W(k)})/M_{0,W(k)} \in 2\mathbb{Z}$.

Lemma 5.12. For $g \in G^{\mathrm{der}}(\mathrm{Frac} W(k))$, we have $g(M_{0,W(k)}) \in \mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)})$. Moreover, the map

$$G^{\mathrm{der}}(\mathrm{Frac} W(k))/K_{p,W(k)} \rightarrow \mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)})$$

is bijective, where $K_{p,W(k)} := \mathrm{Stab}_{G^{\mathrm{der}}(\mathrm{Frac} W(k))}(M_{0,W(k)})$.

Proof. The first assertion is a consequence of [RSZ17, Lemma 3.2]. The injectivity of the map is clear. To prove the surjectivity, take $M \in \mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)})$. By [Jac62, Proposition 8.1], there is $g \in G$ such that $\det_F(g) = 1$ and $M = g(M_{0,W(k)})$. Then we have $\det_F(g) \in 1 + \varpi O_{F,W(k)}$ by [RSZ17, Lemma 3.2]. On the other hand, Hilbert Satz 90 implies that there is $t \in G(\mathrm{Frac} W(k))$ such that $\mathrm{sim}_G(t) = 1$, $\det_F(t) = \det_F(g)$ and $t(M_{0,W(k)}) = M_{0,W(k)}$ (note that we use the condition $\det_F(g) \in 1 + \varpi O_{F,W(k)}$ to prove $t(M_{0,W(k)}) = M_{0,W(k)}$). Then we have $gt^{-1} \in G^{\mathrm{der}}(\mathrm{Frac} W(k))$ and $gt^{-1}(M_{0,W(k)}) = M$. \blacksquare

Let $\mathrm{Lat}^1(\mathbf{L}_{b,W(k)})$ be the set of $W(k)$ -lattices L in $\mathbf{L}_{b,W(k)}$ satisfying $L^\vee \subset L$ and

$$\mathrm{length}_{W(k)}(L/L^\vee) = 1.$$

Lemma 5.13.

- (i) For any $M \in \mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)})$, $L^\sharp(M) := \{v \in \mathbf{L}_{b,W(k)} \mid v(M) \subset \varpi^{-1}M\}$ is an element of $\mathrm{Lat}^1(\mathbf{L}_{b,W(k)})$. Moreover, we have $L^\sharp(M)^\vee = \{v \in \mathbf{L}_{b,W(k)} \mid v(M) \subset M\}$.
- (ii) The map

$$\mathrm{Lat}_{M_0,0}^{\varpi}(N_{b,W(k)}) \rightarrow \mathrm{Lat}^1(\mathbf{L}_{b,W(k)}); M \mapsto L^\sharp(M)$$

is a $G^{\mathrm{der}}(\mathrm{Frac} W(k))$ -equivariant bijection.

Proof. (i): This follows from Lemma 5.12 and Proposition 5.5 (ii).

(ii): The $G^{\mathrm{der}}(\mathrm{Frac} W(k))$ -equivalence is clear by definition. The injectivity follows from Lemma 5.12 and Proposition 5.7 (ii). The surjectivity follows from [Shi10, Lemma 29.9] (by considering dual lattices) and Proposition 5.7 (i). \blacksquare

We regard $G \otimes_{\mathbb{Q}_p} K_0$ as a subgroup of $\mathrm{Res}_{\check{F}/K_0} \mathrm{GL}(N_b)$ by the basis $\mathbf{e}_1, \dots, \mathbf{e}_4$, and put

$$w_0 := \varpi \mathrm{id}_{N_b}, \quad w_1 := \mathrm{diag}(1, 1, -\varpi^2, -\varpi^2).$$

Note that w_0 and w_1 are elements of $\varpi \cdot G^{\text{der}}(\text{Frac } W(k))$. We set

$$X(b_0)_{K_{p,W(k)}} := \left\{ g \in G^{\text{der}}(\text{Frac } W(k))/K_{p,W(k)} \left| g^{-1}b_0\sigma(g) \in \bigcup_{i \in \{0,1\}} K_{p,W(k)}w_iK_{p,W(k)} \right. \right\},$$

where $b_0 \in G(K_0)$ is an element defined in Section 2.4. Note that we have $\text{sim}_G(b_0)^2 = \det_F(b_0) = \varpi^2$. Then the map

$$X(b_0)_{K_{p,W(k)}} \rightarrow \text{DiL}_{M_0,0}^{\varpi}(N_{b,W(k)}); g \mapsto g(M_{0,W(k)})$$

is an isomorphism.

On the other hand, we regard $\text{SO}(\mathbf{L}_{b,0})$ as a subgroup of $\text{GL}(\mathbf{L}_{b,0})$ by the basis x_1, \dots, x_6 , and put

$$w'_0 := \text{id}_{\mathbf{L}_b}, \quad w'_1 := \text{diag}(\varpi^{-2}, \varpi^2, -1, -1, -1, -1).$$

We denote by $K'_{p,W(k)}$ the stabilizer of $L_{0,W(k)}$ in $\text{SO}(\mathbf{L}_b)(\text{Frac } W(k))$. Note that the surjection $\pi: G^0 \otimes_{\mathbb{Q}_p} K_0 \rightarrow \text{SO}(\mathbf{L}_b)$ maps $\varpi^{-1}w_i$ to w'_i for each $i \in \{0,1\}$. Now put $b'_0 := \pi(\varpi^{-1}b_0)$, and set

$$X(b'_0)_{K'_{p,W(k)}} := \left\{ h \in \text{SO}(\mathbf{L}_b)(\text{Frac } W(k))/K'_{p,W(k)} \left| h^{-1}b'_0\sigma(h) \in \bigcup_{i \in \{0,1\}} K_{p,W(k)}w'_iK_{p,W(k)} \right. \right\}.$$

Then the map

$$X(b'_0)_{K'_{p,W(k)}} \rightarrow \text{SpL}(\mathbf{L}_{b,W(k)}); h \mapsto h(L_{0,W(k)}^{\vee})$$

is bijective.

Lemma 5.14. *The surjection π induces an isomorphism*

$$X(b_0)_{K_{p,W(k)}} \xrightarrow{\cong} X(b'_0)_{K'_{p,W(k)}}.$$

such that the following diagram commutes:

$$\begin{array}{ccc} X(b_0)_{K_{p,W(k)}} & \xrightarrow{\cong} & X(b'_0)_{K'_{p,W(k)}} \\ \downarrow \cong & & \downarrow h \mapsto h(L_{0,W(k)}^{\vee}) \\ \text{DiL}_{M_0,0}^{\varpi}(N_{b,W(k)}) & \xrightarrow{M \mapsto L(M)} & \text{Lat}^1(\mathbf{L}_{b,W(k)}). \end{array}$$

Note that Lemma 5.14 immediately implies Proposition 5.11.

Proof of Lemma 5.14. Put

$$\text{SO}(\mathbf{L}_b)(\text{Frac } W(k))_1 := \pi(G^{\text{der}}(\text{Frac } W(k))), \quad K'_{p,W(k),1} := \pi(K_{p,W(k)}).$$

Define

$$X'(b'_0)_{K'_{p,W(k),1}} := \left\{ h \in \text{SO}(\mathbf{L}_b)(\text{Frac } W(k))_1/K'_{p,W(k),1} \left| h^{-1}b'_0\sigma(h) \in \bigcup_{i \in \{0,1\}} K'_{p,W(k),1}w'_iK'_{p,W(k),1} \right. \right\}.$$

Since $\text{Ker}(\pi|_{G^{\text{der}}(\text{Frac } W(k))}) = \{\pm \text{id}_{N_{b,W(k)}}\}$ and $\pi(\varpi^{-1}w_i) = w'_i$ for $i \in \{0,1\}$, the homomorphism π induces an isomorphism

$$X(b_0)_{K_{p,W(k)}} \cong X'(b'_0)_{K'_{p,W(k),1}}.$$

On the other hand, note that the sequence

$$G^{\text{der}}(\text{Frac } W(k)) \xrightarrow{\pi|_{G^{\text{der}}(K_0)}} \text{SO}(\mathbf{L}_b)(\text{Frac } W(k)) \xrightarrow{\theta_{\mathbf{L}_b,W(k)}} \text{Frac } W(k)^{\times}/(\text{Frac } W(k))^2 \cong \mathbb{Z}/2$$

is exact. Moreover, the direct computation implies the following:

$$\pi(\varpi \operatorname{id}_{N_{b,W(k)}}) = -\operatorname{id}_{\mathbf{L}_{b,W(k)}}, \quad \theta_{\mathbf{L}_{b,W(k)}}(-\operatorname{id}_{\mathbf{L}_{b,W(k)}}) \neq 0.$$

Hence we obtain equalities

$$\operatorname{SO}(\mathbf{L}_b)(\operatorname{Frac} W(k)) = \pi(G^{\operatorname{der}}(\operatorname{Frac} W(k)) \times \{\pm \operatorname{id}_{\mathbf{L}_{b,W(k)}}\}), \quad K'_{p,W(k)} = \pi(K_{p,W(k)} \times \{\pm \operatorname{id}_{\mathbf{L}_{b,W(k)}}\}).$$

Hence the canonical map $X(b_0)_{K_{p,W(k)},1} \rightarrow X(b_0)_{K_{p,W(k)}}$ is injective. It is also surjective since $\theta_{\mathbf{L}_b}(w'_i) = 0$ for $i \in \{0, 1\}$ and $\theta_{\mathbf{L}_{b,W(k)}}(h^{-1}b'_0\sigma(h)) = 0$ for any $h \in G(\operatorname{Frac} W(k))$.

Finally, the commutativity of the diagram is clear. \blacksquare

We give another application of the bijection $h \mapsto h(L_{0,W(k)}^\vee)$, which will be used in Section 5.6.

Proposition 5.15. *For $L \in \operatorname{SpL}(\mathbf{L}_{b,W(k)})$, we have $\Phi_b(L) \subset p^{-1}L^\vee$.*

Proof. Write $L = h(L_{0,W(k)}^\vee)$, where $h \in \operatorname{SO}(\mathbf{L}_b)(\operatorname{Frac} W(k))$ satisfies $h^{-1}b'_0\sigma(h) \in K_{p,W(k)}w'_iK_{p,W(k)}$ for some $i \in \{0, 1\}$. Note that $L^\vee = h(L_{0,W(k)})$. Write $h^{-1}b'_0\sigma(h) = h_1w'_1h_2$, where $h_1, h_2 \in K_{p,W(k)}$. By replacing h to hh_1 if necessary, we may assume $h_1 = \operatorname{id}_{\mathbf{L}_b}$.

If $i = 0$, then we have $\Phi_b(L) = L$, and hence the assertion is trivial. Otherwise, we have

$$\begin{aligned} w'_1(L_{0,W(k)}^\vee) &= W(p^{-1}x_1) \oplus Wx_2 \oplus Wx_3 \oplus Wx_4 \oplus Wx_5 \oplus Wx_6 \\ &\subset W(p^{-1}x_1) \oplus W(p^{-1}x_2) \oplus Wx_3 \oplus W(p^{-1}x_4) \oplus W(p^{-1}x_5) \oplus W(p^{-1}x_6) = p^{-1}L_0. \end{aligned}$$

Here x_1, \dots, x_6 is the basis of \mathbf{L}_b defined in Section 5.2. Using this inclusion, we have

$$\Phi_b(L) = h(h^{-1}b'_0\sigma(h))(L_{0,W(k)}^\vee) = hw'_1(L_{0,W(k)}^\vee) \subset h(p^{-1}L_{0,W(k)}) = p^{-1}L^\vee.$$

Therefore the assertion follows. \blacksquare

5.5. Rapoport–Zink space for the quaternionic unitary group of degree 2. To understand the structure of $\mathcal{M}_{b,\mu}^{(0,0)}$, we will use a closed immersion from the Rapoport–Zink space for the quaternionic unitary similitude group of degree 2. We recall the definition and the Bruhat–Tits stratification of it constructed by [Oki19].

From now on, put $D := F[y_6]$ and $O_D := O_F[y_6]$, which are subrings of $\operatorname{End}^0(\mathbb{X}_0)$. Note that D is the quaternion division algebra over \mathbb{Q}_p , and O_D is the maximal order of D . We define an involution $*$ on D by

$$*: D \rightarrow D; d \mapsto y_6(\operatorname{Trd}_{D/\mathbb{Q}_p}(d) - d)y_6^{-1}.$$

Let G_D be the quaternionic unitary similitude group of degree 2. We regard G_D as a subgroup of G by the natural way in the sequel. For $S \in \operatorname{Nilp}$, a p -divisible with G_D -structure is a triple (X, ι, λ) , where

- X is a p -divisible group over S of dimension 4 and height 8,
- $\iota: O_D \rightarrow \operatorname{End}(X)$ is a ring homomorphism,
- $\lambda: X \rightarrow X^\vee$ is a quasi-isogeny so that $\lambda^\vee = -\lambda$,

satisfying the following conditions for any $d \in O_D$;

- (Kottwitz condition) $\det(T - \iota(d) \mid \operatorname{Lie}(X)) = (T^2 - \operatorname{Trd}_{D/\mathbb{Q}_p}(d) + \operatorname{Nrd}_{D/\mathbb{Q}_p}(d))^2$,
- $\lambda \circ \iota(d) = \iota(d^*)^\vee \circ \lambda$.

We define a quasi-isogeny of p -divisible groups with G_D -structures by the same manner as Definition 2.1 (ii).

Now we define $\iota'_0: O_D \rightarrow \operatorname{End}(\mathbb{X}_0)$ by the composite of the actions of O_F and y_6 on \mathbb{X}_0 , and put $\lambda'_0 := (y_6 \circ \iota_0(\varpi))^\vee \circ \lambda_0$. Then $(\mathbb{X}_0, \iota'_0, \lambda'_0)$ is a p -divisible group with G_D -structure over $\operatorname{Spec} \overline{\mathbb{F}}_p$. Furthermore, λ'_0 is an isomorphism since $\operatorname{Ker}(\lambda_0) = \operatorname{Ker}(\iota_0(\varpi))$ and y_6 is an isomorphism. We

define \mathcal{M}_D as the functor which parametrizes $(X, \iota, \lambda, \rho)$ for $S \in \text{Nilp}$, where (X, ι, λ) is a p -divisible group over S of dimension 4 and height 8 such that λ is an isomorphism, and

$$\rho: X \times_S \overline{S} \rightarrow \mathbb{X}_0 \otimes_{\overline{\mathbb{F}}_p} \overline{S}$$

is a quasi-isogeny of p -divisible groups with G_D -structures. We define an equivalence of two p -divisible groups with G_D -structures by the same manner as that of two p -divisible groups with (G, μ) -structures. The representability of \mathcal{M}_D is a consequence of [RZ96, Theorem 3.25]. Moreover, there is a decomposition into open and closed formal subschemes

$$\mathcal{M}_D = \coprod_{i \in \mathbb{Z}} \mathcal{M}_D^{(i)},$$

where $\mathcal{M}_D^{(i)}$ is the locus $(X, \iota, \lambda, \rho)$ of \mathcal{M}_D for $\text{Im}(c(\rho)) \subset p^i \mathbb{Z}_p^\times$.

Let J_D be the algebraic group over \mathbb{Q}_p defined by

$$J_D(R) = \{(g, c) \in (\text{End}_{O_D}(\mathbb{X}_0) \otimes_{\mathbb{Z}_p} R)^\times \times R^\times \mid g^\vee \circ \lambda'_0 \circ g = c \lambda'_0\}$$

for any \mathbb{Q}_p -algebra R . The group $J_D(\mathbb{Q}_p)$ acts on \mathcal{M}_D by the same manner as the action of $J_b(\mathbb{Q}_p)$ on $\mathcal{M}_{b, \mu}$. It is known that there is an isomorphism $J_D \cong \text{GSp}_4$. See [RZ96, 1.42].

Next, we recall the Bruhat–Tits stratification of \mathcal{M}_D constructed in [Oki19, Section 5.3]. Put

$$\mathbf{L}_D := \iota'_0(y_6 \varpi)^{-1} \{f \in \text{End}_{O_D}^0(\mathbb{X}_0) \mid \text{Tr}_{D/\mathbb{Q}_p}(f) = 0, f^\vee \circ \lambda'_0 = \lambda'_0 \circ f\}.$$

We call a lattice Λ in \mathbf{L}_D is a *vertex lattice* if we have $p\Lambda \subset \Lambda^\vee \subset \Lambda$. If Λ is a vertex lattice in \mathbf{L}_D , we call $t(\Lambda) := \dim_{\mathbb{F}_p}(\Lambda/\Lambda^\vee)$ as the type of Λ . Note that we have $t(\Lambda) \in \{1, 3, 5\}$ by [Oki19, Lemma 5.9]. We denote by $\text{VL}(\mathbf{L}_D, t)$ the set of all vertex lattices in \mathbf{L}_D of type t for $t \in \{1, 3, 5\}$, and put $\text{VL}(\mathbf{L}_D) := \sqcup_{t \in \{1, 3, 5\}} \text{VL}(\mathbf{L}_D, t)$.

For $\Lambda \in \text{VL}(\mathbf{L}_D)$, let $\mathcal{M}_{D, \Lambda}^{(0)}$ be the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_D^{(0)}$ where $\rho^{-1} \circ \Lambda \circ \rho \subset \text{End}(X)$. It is a closed formal subscheme of $\mathcal{M}_D^{(0)}$.

Proposition 5.16. ([Oki19, Corollary 5.25, Theorems 5.26, 5.27])

(i) For $\Lambda, \Lambda' \in \text{VL}(\mathbf{L}_D)$, we have

$$\mathcal{M}_{D, \Lambda}^{(0)} \cap \mathcal{M}_{D, \Lambda'}^{(0)} = \begin{cases} \mathcal{M}_{D, \Lambda \cap \Lambda'}^{(0)} & \text{if } \Lambda \cap \Lambda' \in \text{VL}(\mathbf{L}_D), \\ \emptyset & \text{otherwise.} \end{cases}$$

(ii) For $\Lambda \in \text{VL}(\mathbf{L}_D)$, the formal scheme $\mathcal{M}_{D, \Lambda}^{(0)}$ is a reduced scheme over $\text{Spec } \overline{\mathbb{F}}_p$. Moreover, the following hold:

- if $t(\Lambda) = 1$, then $\mathcal{M}_{D, \Lambda}^{(0)}$ is a single point,
- if $t(\Lambda) = 3$, then $\mathcal{M}_{D, \Lambda}^{(0)}$ is isomorphic to $\mathbb{P}_{\overline{\mathbb{F}}_p}^1$,
- if $t(\Lambda) = 5$, then $\mathcal{M}_{D, \Lambda}^{(0)}$ is isomorphic to the Fermat surface F_p defined by

$$x_0^{p+1} + x_1^{p+1} + x_2^{p+1} + x_3^{p+1} = 0$$

$$\text{in } \mathbb{P}_{\overline{\mathbb{F}}_p}^3 = \text{Proj } \overline{\mathbb{F}}_p[x_0, x_1, x_2, x_3].$$

In particular, $\mathcal{M}_{D, \Lambda}^{(0)}$ is projective smooth and irreducible of dimension $(t(\Lambda) - 1)/2$.

(iii) We have equalities

$$\mathcal{M}_D^{(0)} = \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_D)} \mathcal{M}_{D, \Lambda}^{(0)} = \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_D, 5)} \mathcal{M}_{D, \Lambda}^{(0)}.$$

In particular, any irreducible component of $\mathcal{M}_D^{(0)}$ is of the form $\mathcal{M}_{D, \Lambda}^{(0)}$ where $\Lambda \in \mathrm{VL}(\mathbf{L}_D, 5)$, which has 2-dimensional.

We give a relation between \mathcal{M}_D and $\mathcal{M}_{b, \mu}$. Let $N_{b, W(k)}^0$ be the eigenspace of y_6 on $N_{b, W(k)}$ with respect to ε . Then the restriction to $N_{b, W(k)}^0$ of the alternating form $(,)$ on $N_{b, W(k)}$ is non-degenerate. Moreover, $N_{b, W(k)}^0$ is stable under the σ -linear map $\mathbf{F}_{b, 0}$.

Proposition 5.17. *The morphism*

$$i_{y_6} : \mathcal{M}_D \rightarrow \mathcal{M}_{b, \mu}; (X, \iota, \lambda, \rho) \mapsto (X, \iota|_{\mathcal{O}_F}, \lambda \circ (\rho^{-1} y_6 \circ \rho) \circ \iota(\varpi), \rho)$$

is a closed immersion. The image of the closed immersion i_{y_6} equals the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_{b, \mu}$ where $\rho^{-1} \circ y_6 \circ \rho \in \mathrm{End}(X)$.

To prove Proposition 5.17, we need to pay attention to the Kottwitz condition. Let $\mathcal{Z}(y_6)$ be the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_{b, \mu}$ where $\rho^{-1} \circ y_6 \circ \rho \in \mathrm{End}(X)$.

Lemma 5.18. *For any algebraically closed field k of characteristic p , the last assertion in Proposition 5.17 holds for the sets of k -valued points.*

Proof. Put $\mathcal{Z}^{(0)}(y_6) := \mathcal{Z}(y_6) \cap \mathcal{M}_{b, \mu}^{(0)}$. It suffices to show the equality

$$i_{y_6}(\mathcal{M}_D^{(0)})(k) = \mathcal{Z}^{(0)}(y_6)(k).$$

Let $\mathrm{DiL}_{y_6}^{\varpi}(N_{b, W(k)})$ be the subset of $\mathrm{DiL}^{\varpi}(N_{b, W(k)})$ consisting of y_6 -stable elements. Then the bijection in Proposition 2.5 induces a bijection

$$\mathcal{Z}^{(0)}(y_6)(k) \cong \mathrm{DiL}_{y_6}^{\varpi}(N_{b, W(k)}).$$

Put $N_{b, W(k)}^0 := N_b^0 \otimes_W W(k)$, and $\mathrm{Lat}^2(N_{b, k}^0)$ be the set of lattices M^0 in $N_{b, W(k)}^0$ satisfying the following:

- $M^0 \subset (M^0)^\vee \subset p^{-1}M^0$ and $\mathrm{length}_{W(k)}((M^0)^\vee/M^0) = 2$,
- $M^0 \subset \mathbf{F}_{b, 0}((M^0)^\vee) \subset p^{-1}M^0$,
- $p(M^0)^\vee \subset \mathbf{F}_{b, 0}(M^0) \subset (M^0)^\vee$.

Claim. For $M \in \mathrm{DiL}_{y_6}^{\varpi}(N_{b, W(k)})$, $M \cap N_{b, W(k)}^0$ is an element of $\mathrm{Lat}^2(N_{b, W(k)}^0)$.

We prove this claim. Since $(M^0)^\vee = (\varpi^{-1}M^0)^\vee$, all the inclusion relations hold. Moreover, we have

$$1 \leq \mathrm{length}_{W(k)}(\mathbf{F}_{b, 0}((M^0)^\vee)/M^0) \leq 3$$

by the equality $y_6^2 = \varepsilon^2$ and the Kottwitz condition for the p -divisible group with (G, μ) -structure over k corresponding to M . On the other hand, we have $\mathrm{length}_{W(k)}((M^0)^\vee/M^0) \in 2\mathbb{Z}$, and hence

$$\mathrm{length}_{W(k)}((M^0)^\vee/M^0) = \mathrm{length}_{W(k)}(\mathbf{F}_{b, 0}((M^0)^\vee)/M^0) = 2,$$

as desired.

By Claim, we obtain a map

$$\mathrm{DiL}_{y_6}^{\varpi}(N_{b, W(k)}) \rightarrow \mathrm{Lat}^2(N_{b, W(k)}^0); M \mapsto M \cap N_{b, W(k)}^0.$$

This is bijective since

$$\mathrm{Lat}^2(N_{b, W(k)}^0) \rightarrow \mathrm{DiL}_{y_6}^{\varpi}(N_{b, W(k)}); M^0 \mapsto M^0 \oplus \varpi((M^0)^\vee)$$

gives an inverse map. On the other hand, the bijection in Proposition 2.5 induces a bijection

$$i_{y_6}(\mathcal{M}_D^{(0)})(k) \xrightarrow{\cong} \mathrm{Lat}^2(N_{b, W(k)}^0),$$

cf. [Wan20, Proposition 2.4]. Hence we obtain $i_{y_6}(\mathcal{M}_D^{(0)})(k) = \mathcal{Z}^{(0)}(y_6)(k)$ as desired. \blacksquare

Proof of Proposition 5.17. The injectivity of i_{y_6} as a functor is a consequence of the rigidity of quasi-isogenies (the assertion after [RZ96, 2.9]). To prove the rest of the assertions, it suffices to show the equality $i_{y_6}(\mathcal{M}_D) = \mathcal{Z}(y_6)$. Indeed, this equality and the rigidity of quasi-isogenies implies that i_{y_6} is a closed immersion.

The inclusion $i_{y_6}(\mathcal{M}_D) \subset \mathcal{Z}(y_6)$ is clear. Let us prove the another inclusion. Let $x = (X, \iota, \lambda, \rho) \in \mathcal{Z}(y_6)(S)$ where $S \in \text{Nilp}$. By localization, we may assume that $S = \text{Spec } R$, where R is a local ring. For $i \in \{0, 1\}$, let $\text{Lie}^i(X)$ be the eigenspace of y_6 on $\text{Lie}(X)$ with respect to $(-1)^i \varepsilon$. Then the both are locally free of rank 2 by applying Lemma 5.18 for the base change of x to an algebraic closure of the residue field of R . Hence we have $x \in i_{y_6}(\mathcal{M}_D)(S)$. \blacksquare

Proposition 5.19. *For any algebraically closed field k of characteristic p , the set $i_{y_6}(\mathcal{M}_D^{(0)})(k)$ is contained in $\mathcal{M}_{b,\mu}^{(0,0)}(k)$.*

Proof. Take $x = (X, \iota, \lambda, \rho) \in \mathcal{M}_D^{(0)}(k)$, and let $M \in \text{DiL}_{M_0,0}^{\varpi}(N_{b,W(k)})$ be corresponding to $\iota_{y_6}(x) \in \mathcal{M}_{b,\mu}^{(0)}(k)$. Then there is $g \in G_D^{\text{der}}(\text{Frac } W(k))$ such that $M = g(M_{0,W(k)})$ (see also the proof of Lemma 5.18). Hence we obtain $M \in \text{DiL}^{\varpi}(N_{b,W(k)})$ by Lemma 5.12. \blacksquare

5.6. Vertex lattices, I.

Definition 5.20. A vertex lattice in $\mathbf{L}_{b,0}$ is a \mathbb{Z}_p -lattice Λ satisfying $p\Lambda \subset \Lambda^\vee \subset \Lambda$. We call $t(\Lambda) := \dim_{\mathbb{F}_p}(\Lambda/\Lambda^\vee)$ as the type of Λ .

We denote by $\text{VL}(\mathbf{L}_{b,0})$ the set of all vertex lattices in $\mathbf{L}_{b,0}$.

Lemma 5.21. *For $\Lambda \in \text{VL}(\mathbf{L}_{b,0})$, we have $t(\Lambda) \in \{1, 3, 5\}$.*

Proof. This follows by the same argument as the proof of [Oki19, Lemma 5.9]. \blacksquare

For $t \in \{1, 3, 5\}$, we denote by $\text{VL}(\mathbf{L}_{b,0}, t)$ the set of vertex lattices in $\mathbf{L}_{b,0}$ of type t .

We denote by $\mathcal{B}(\mathbf{L}_{b,0})$ the Bruhat–Tits building of $\text{SO}(\mathbf{L}_{b,0})(\mathbb{Q}_p)$.

Proposition 5.22. *There is an $\text{SO}(\mathbf{L}_{b,0})(\mathbb{Q}_p)$ -equivariant bijection between $\text{VL}(\mathbf{L}_{b,0})$ and the set of vertices in $\mathcal{B}(\mathbf{L}_{b,0})$ satisfying the following.*

- (i) *The set $\text{VL}(\mathbf{L}_{b,0}, 1) \sqcup \text{VL}(\mathbf{L}_{b,0}, 5)$ corresponds to the set of special vertices in \mathcal{B}_b .*
- (ii) *If $\Lambda_0, \Lambda_1 \in \text{VL}(\mathbf{L}_{b,0}, 3)$ are distinct, then the corresponding vertices in $\mathcal{B}(\mathbf{L}_{b,0})$ are adjacent if and only if $\Lambda_i \in \text{VL}(\mathbf{L}_{b,0}, t)$, $\Lambda_{1-i} \in \text{VL}(\mathbf{L}_{b,0}, t')$ and $\Lambda_i \subset \Lambda_{1-i}$ for some $i \in \{0, 1\}$ and $t, t' \in \{1, 3, 5\}$ satisfying $t < t'$.*

Proof. Let $\mathcal{B}'(\mathbf{L}_{b,0})$ be the Bruhat–Tits building of $\text{O}(\mathbf{L}_{b,0})(\mathbb{Q}_p)$. Then, by [AN02] (for $\mathcal{D} = \mathbb{Q}_p$, $(\bar{\cdot}, \epsilon) = (\text{id}, 1)$ and $(n, r) = (6, 2)$), all the assertions are true when we replace $\mathcal{B}(\mathbf{L}_{b,0})$ with $\mathcal{B}'(\mathbf{L}_{b,0})$. On the other hand, recall that $\text{SO}(\mathbf{L}_{b,0})$ is the identity component of $\text{O}(\mathbf{L}_{b,0})$, and the center of $\text{SO}(\mathbf{L}_{b,0})$ is $\{\pm \text{id}_{\mathbf{L}_{b,0}}\} \cong \mu_2$, which is finite over \mathbb{Q}_p . Hence [Tit79, §2.1, p.44] implies that there is a canonical isomorphism of simplicial complexes $\mathcal{B}(\mathbf{L}_{b,0}) \cong \mathcal{B}'(\mathbf{L}_{b,0})$, which is $\text{SO}(\mathbf{L}_{b,0})$ -equivariant. Therefore we obtain the desired assertions. \blacksquare

Lemma 5.23. *For $\Lambda, \Lambda' \in \text{VL}(\mathbf{L}_{b,0})$, there is a sequence of vertex lattices in $\mathbf{L}_{b,0}$*

$$\Lambda_0 = \Lambda, \Lambda_1, \dots, \Lambda_n = \Lambda'$$

such that $\Lambda_i \subset \Lambda_{i+1}$ or $\Lambda_i \supset \Lambda_{i+1}$ for each $i \in \{0, \dots, n-1\}$.

Proof. This follows from the same argument as [HP17, Proposition 5.1.5]. \blacksquare

Lemma 5.24. *The perpendicular of $\mathbb{Q}_p y_6$ in $\mathbf{L}_{b,0}$ equals \mathbf{L}_D .*

Proof. The proof is the same as that of [Oki19, Proposition 3.20] by using Lemma 5.6. \blacksquare

Lemma 5.25. *For $\Lambda \in \mathrm{VL}(\mathbf{L}_{b,0})$, there is $g \in J_b^{\mathrm{der}}(\mathbb{Q}_p)$ and $\Lambda' \in \mathrm{VL}(\mathbf{L}_D, t(\Lambda))$ satisfying*

$$g(\Lambda) = \Lambda' \oplus \mathbb{Z}_p y_6.$$

Proof. Take $\Lambda'' \in \mathrm{VL}(\mathbf{L}_D, t(\Lambda))$. By Lemma 5.24, we have $\Lambda'' \oplus \mathbb{Z}_p y_6 \in \mathrm{VL}(\mathbf{L}_{b,0}, t(\Lambda))$. Hence [Shi10, Lemma 29.9] implies that there is $g' \in \mathrm{SO}(\mathbf{L}_{b,0})$ such that $g'(\Lambda) = \Lambda'' \oplus \mathbb{Z}_p y_6$. On the other hand, since $\dim_{\mathbb{Q}_p}(\mathbf{L}_D) > 3$, Proposition 5.1 for \mathbf{L}_D implies that there is $h \in \mathrm{SO}(\mathbf{L}_D)$ such that $\theta_{\mathbf{L}_D}(h) = \theta_{\mathbf{L}_{b,0}}(g')$. Hence hg' is contained in the image of $J_b^{\mathrm{der}}(\mathbb{Q}_p) \rightarrow \mathrm{SO}(\mathbf{L}_{b,0})$ by Proposition 5.1 for $\mathbf{L}_{b,0}$. Therefore, the element $g \in J_b^{\mathrm{der}}(\mathbb{Q}_p)$ corresponding to hg' and the vertex lattice $\Lambda' := h(\Lambda'')$ give a desired assertion. \blacksquare

For a $W(k)$ -lattice L in $\mathbf{L}_{b,W(k)}$ and $r \in \mathbb{Z}_{\geq 0}$, we define another $W(k)$ -lattice in $\mathbf{L}_{b,W(k)}$ as follows:

$$L^{(r)} := L + \Phi_b(L) + \cdots + \Phi_b^r(L).$$

Proposition 5.26. *For $L \in \mathrm{SpL}(\mathbf{L}_{b,W(k)})$, there is the minimum integer $r \in \{0, 1, 2\}$ such that $L^{(r)}$ is Φ_b -stable. Then, we have $L^{(r)} \in \mathrm{VL}(\mathbf{L}_{b,0}, 2r + 1)$ and $(L^{(r)})^\vee = \{v \in L^\vee \mid \Phi_b(v) = v\}$.*

Proof. We follow the proof of [Cho18, Lemma 2.7]. Let

$$r_0 := \min\{r \in \mathbb{Z}_{\geq 0} \mid \Phi_b(L^{(r)}) = L^{(r)}\}, \quad r_0^\vee := \min\{r \in \mathbb{Z}_{\geq 0} \mid \Phi_b((L^\vee)^{(r)}) = (L^\vee)^{(r)}\}.$$

Note that the existences of r_0 are r_0^\vee are consequences of [RZ96, Proposition 2.17]. Moreover, the assertions $(L^{(r_0)})^\vee \subset L^{(r_0)}$ and $(L^{(r_0)})^\vee = \{v \in L^\vee \mid \Phi_b(v) = v\}$ are trivial by definition.

In the sequel, we prove $pL^{(r_0)} \subset (L^{(r_0)})^\vee$, which implies $r_0 \in \{0, 1, 2\}$ and $L^{(r_0)} \in \mathrm{VL}(\mathbf{L}_{b,0}, 2r_0 + 1)$.

Claim. We have $r_0 \leq r_0^\vee$.

By the same argument as the proof of [Cho18, Lemma 2.8], we have

$$L^{(i)} \cap \Phi_b(L^{(i)}) = \Phi_b(L^{(i-1)})$$

for $1 \leq i < r_0$. If $r_0^\vee < r_0$, then the above equality implies

$$L^\vee \subset \bigcap_{n \in \mathbb{Z}} \Phi_b^n(L^{(r_0^\vee)}) = \bigcap_{n \in \mathbb{Z}} \Phi_b^n(L) \subset L \cap \Phi_b(L) \subsetneq L.$$

Since $\mathrm{length}_{W(k)} L/L^\vee = 1$, we have $L^\vee = L \cap \Phi_b(L)$, and hence $(L^\vee)^{(1)} = L$ by taking duals. Therefore we obtain $r_0^\vee = r_0 + 1 > r_0$, which contradicts to the assumption.

By Proposition 5.15 and the definition of r_0 , we have

$$pL^{(r_0)} \subset \bigcap_{n \in \mathbb{Z}} \Phi_b^n((L^\vee)^{(r_0-1)}).$$

On the other hand, the same argument as the proof of [Cho18, Lemma 2.8] implies

$$(L^\vee)^{(i)} \cap \Phi_b((L^\vee)^{(i)}) = \Phi_b((L^\vee)^{(i-1)})$$

for $1 \leq i < r_0^\vee$. Since $r_0 - 1 < r_0^\vee$ by Claim, we obtain

$$\bigcap_{n \in \mathbb{Z}} \Phi_b^n((L^\vee)^{(r_0-1)}) = \bigcap_{n \in \mathbb{Z}} \Phi_b^n(L^\vee) = (L^{(r_0)})^\vee.$$

Therefore the assertion $pL^{(r_0)} \subset (L^{(r_0)})^\vee$ is obtained. \blacksquare

For $\Lambda \in \text{VL}(\mathbf{L}_{b,0})$, let $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}(k)$ be the subset of $\mathcal{M}_{b,\mu}^{(0,0)}(k)$ corresponding to $\{L \in \text{SpL}(\mathbf{L}_{b,W(k)}) \mid \Lambda^\vee \subset L\}$ under the bijection in Proposition 5.11.

Corollary 5.27. *We have equalities*

$$\mathcal{M}_{b,\mu}^{(0,0)}(k) = \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_{b,0})} \mathcal{M}_{b,\mu,\Lambda}^{(0,0)}(k) = \bigcup_{g \in J_b^{\text{der}}(\mathbb{Q}_p)} \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_D, 5)} g(i_{y_6}(\mathcal{M}_{D,\Lambda}^{(0)}))(k).$$

Proof. The first equality follows from Proposition 5.26. The second equality follows from Lemma 5.25, Corollary 5.8 and Proposition 5.16 (iii). \blacksquare

5.7. Connected components. Here we study the connected components of the underlying topological space $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ of $\mathcal{M}_{b,\mu}^{(0)}$. For $x \in \mathcal{M}_{b,\mu}^{(0)}$, we denote by $\overline{k(x)}$ an algebraic closure of the residue field of x .

Definition 5.28. For $j \in \mathbb{Z}/2$, we define a subset $\mathcal{M}_{b,\mu}^{(0,j),\text{red}}$ of the topological space $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ as follows:

$$\mathcal{M}_{b,\mu}^{(0,j),\text{red}} := \{x \in \mathcal{M}_{b,\mu}^{(0),\text{red}} \mid x \in \mathcal{M}_{b,\mu}^{(0,j),\text{red}}(\overline{k(x)})\}.$$

By definition, we have

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \mathcal{M}_{b,\mu}^{(0,0),\text{red}} \sqcup \mathcal{M}_{b,\mu}^{(0,1),\text{red}}.$$

Moreover, the action of $s \in J_b(\mathbb{Q}_p)$ satisfying $\kappa_{J_b}(s) = (0, 1)$ induces a bijection

$$\mathcal{M}_{b,\mu}^{(0,0),\text{red}} \cong \mathcal{M}_{b,\mu}^{(0,1),\text{red}}.$$

Here we prove the following:

Theorem 5.29. *The subsets $\mathcal{M}_{b,\mu}^{(0,0),\text{red}}$ and $\mathcal{M}_{b,\mu}^{(0,1),\text{red}}$ are the connected components in $\mathcal{M}_{b,\mu}^{(0),\text{red}}$.*

By Theorem 5.29, we obtain a decomposition into open and closed formal subschemes

$$\mathcal{M}_{b,\mu}^{(0)} = \mathcal{M}_{b,\mu}^{(0,0)} \sqcup \mathcal{M}_{b,\mu}^{(0,1)}$$

such that the underlying topological space of $\mathcal{M}_{b,\mu}^{(0,j)}$ is $\mathcal{M}_{b,\mu}^{(0,j),\text{red}}$ for each $j \in \mathbb{Z}/2$.

From now, we give a proof of Theorem 5.29.

Lemma 5.30. *We have an equality*

$$\mathcal{M}_{b,\mu}^{(0,0),\text{red}} = \bigcup_{g \in J_b^{\text{der}}(\mathbb{Q}_p)} \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_D, 5)} g(i_{y_6}(\mathcal{M}_{D,\Lambda}^{(0)})).$$

Proof. This follows from Corollary 5.27. \blacksquare

Corollary 5.31. *Any element of $\text{Irr}(\mathcal{M}_{b,\mu}^{(0),\text{red}})$ is of the form $g(i_{y_6}(\mathcal{M}_{D,\Lambda}^{(0)}))$ where $g \in J_b^1(\mathbb{Q}_p)$ and $\Lambda \in \text{VL}(\mathbf{L}_D, 5)$. In particular, any $C \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0)})$ satisfies either $C \subset \mathcal{M}_{b,\mu}^{(0,0),\text{red}}$ or $C \subset \mathcal{M}_{b,\mu}^{(0,1),\text{red}}$.*

Proof. By Lemma 5.30, we have

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \bigcup_{g \in J_b^1(\mathbb{Q}_p)} \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_D, 5)} g(i_{y_6}(\mathcal{M}_{D,\Lambda}^{(0)})).$$

Hence the assertion follows from the above equality. \blacksquare

Proof of Theorem 5.29. It suffices to show that $\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$ is connected and open in $\mathcal{M}_{b,\mu}^{(0),\mathrm{red}}$. Indeed, these imply the same assertion for $\mathcal{M}_{b,\mu}^{(0,1),\mathrm{red}}$, and hence the assertion follows.

First, we prove the connectedness of $\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$. By Proposition 5.16 (i) and Corollary 5.31, it suffices to show the following:

Claim. For any $g, g' \in J_b^{\mathrm{der}}(\mathbb{Q}_p)$ and $\Lambda, \Lambda' \in \mathrm{VL}(\mathbf{L}_D)$, there is a sequence

$$\Lambda_0 = g(\Lambda \oplus \mathbb{Z}_p y_6), \Lambda_1 = g_1(\Lambda_{0,1} \oplus \mathbb{Z}_p y_6), \dots, \Lambda_n = g'(\Lambda' \oplus \mathbb{Z}_p y_6)$$

where $g_i \in J_b^{\mathrm{der}}(\mathbb{Q}_p)$ and $\Lambda_{0,i} \in \mathrm{VL}(\mathbf{L}_D)$ such that $\Lambda_i \subset \Lambda_{i+1}$ or $\Lambda_i \supset \Lambda_{i+1}$ for each $i \in \{0, \dots, n-1\}$.

However, this claim follows from Lemma 5.23 since Lemma 5.25 implies the equality

$$\mathrm{VL}(\mathbf{L}_{b,0}) = \{g(\Lambda \oplus \mathbb{Z}_p y_6) \mid g \in J_b^{\mathrm{der}}(\mathbb{Q}_p), \Lambda \in \mathrm{VL}(\mathbf{L}_D)\}.$$

Second, we prove that $\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$ is open in $\mathcal{M}_{b,\mu}^{(0),\mathrm{red}}$. Let $x \in \mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$, and take a quasi-compact open subset V' of $\mathcal{M}_{b,\mu}^{(0),\mathrm{red}}$ containing x . Then there are only finitely many irreducible components which intersects with V' . Now let $\mathrm{Irr}(\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}})_{V',1}$ be the set of $C \in \mathrm{Irr}(\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}})$ satisfying $C \cap V' \neq \emptyset$ and $C \subset \mathcal{M}_{b,\mu}^{(0,1)}$. Then the set

$$U' := V' \setminus \bigcup_{C \in \mathrm{Irr}(\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}})_{V',1}} C.$$

is an open neighborhood of x contained in $\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$ by Corollary 5.31 and the finiteness of the set $\mathrm{Irr}(\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}})_{V',1}$. \blacksquare

5.8. Bruhat–Tits stratification, I. Here we will describe the structure of $\mathcal{M}_{b,\mu}^{(0,0),\mathrm{red}}$ by means of vertex lattices.

Definition 5.32. Let $\Lambda \in \mathrm{VL}(\mathbf{L}_{b,0})$. We define $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ as the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_{b,\mu}^{(0,0)}$ where $\rho^{-1} \circ \Lambda^\vee \circ \rho \subset \mathrm{End}(X)$. It is a closed formal subscheme of $\mathcal{M}_{b,\mu}^{(0,0)}$.

By definition, the set of k -valued points of $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ equals $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}(k)$ for any algebraically closed field of characteristic p .

Proposition 5.33.

(i) For $\Lambda \in \mathrm{VL}(\mathbf{L}_D)$, the closed immersion i_{y_6} in Proposition 5.17 induces an isomorphism

$$\mathcal{M}_{D,\Lambda}^{(0)} \cong \mathcal{M}_{b,\mu,\Lambda \oplus \mathbb{Z}_p y_6}^{(0,0)}.$$

(ii) For $\Lambda \in \mathrm{VL}(\mathbf{L}_{b,0})$, there is $g \in J_b^{\mathrm{der}}(\mathbb{Q}_p)$ and $\Lambda' \in \mathrm{VL}(\mathbf{L}_D, t(\Lambda))$ such that g induces an isomorphism

$$\mathcal{M}_{b,\mu,\Lambda}^{(0,0)} \cong \mathcal{M}_{b,\mu,\Lambda' \oplus \mathbb{Z}_p y_6}^{(0,0)}.$$

Proof. (i): This follows from the description of the image of i_{y_6} in Proposition 5.17.

(ii): This follows from Lemma 5.25 and Corollary 5.8. \blacksquare

Corollary 5.34. For $\Lambda \in \mathrm{VL}(\mathbf{L}_{b,0})$, the formal scheme $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ is a reduced scheme over $\mathrm{Spec} \overline{\mathbb{F}}_p$. Moreover, the following hold:

- if $t(\Lambda) = 1$, then $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ is a single point,

- if $t(\Lambda) = 3$, then $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ is isomorphic to $\mathbb{P}_{\mathbb{F}_p}^1$,
- if $t(\Lambda) = 5$, then $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$ is isomorphic to the Fermat surface F_p .

Proof. These follow from Propositions 5.16 (i) and 5.33. ■

Proposition 5.35.

(i) *We have an equality*

$$\mathcal{M}_{b,\mu}^{(0,0)} = \bigcup_{\Lambda \in \text{VL}(\mathbf{L}_{b,0})} \mathcal{M}_{b,\mu,\Lambda}^{(0,0)}.$$

(ii) *For $\Lambda, \Lambda' \in \text{VL}(\mathbf{L}_{b,0})$, we have*

$$\mathcal{M}_{b,\mu,\Lambda}^{(0,0)} \cap \mathcal{M}_{b,\mu,\Lambda'}^{(0,0)} = \begin{cases} \mathcal{M}_{b,\mu,\Lambda \cap \Lambda'}^{(0,0)} & \text{if } \Lambda \cap \Lambda' \in \text{VL}(\mathbf{L}_{b,0}), \\ \emptyset & \text{otherwise.} \end{cases}$$

Proof. The proof is the same as [Oki19, Theorem 5.26] by using Corollary 5.34. ■

For $\Lambda \in \text{VL}(\mathbf{L}_{b,0})$, we set

$$\text{BT}_{b,\mu,\Lambda}^{(0,0)} := \mathcal{M}_{b,\mu,\Lambda}^{(0,0)} \setminus \bigcup_{\Lambda' \subsetneq \Lambda} \mathcal{M}_{b,\mu,\Lambda'}^{(0,0)}.$$

Theorem 5.36.

(i) *We have*

$$\mathcal{M}_{b,\mu}^{(0,0),\text{red}} = \prod_{\Lambda \in \text{VL}(\mathbf{L}_{b,0})} \text{BT}_{b,\mu,\Lambda}^{(0,0)}.$$

(ii) *For $\Lambda \in \text{VL}(\mathbf{L}_{b,0})$, $\text{BT}_{b,\mu,\Lambda}^{(0,0)}$ is a reduced scheme over $\overline{\mathbb{F}}_p$. Moreover, we have*

$$\mathcal{M}_{b,\mu,\Lambda}^{(0,0),\text{red}} = \prod_{\Lambda' \subset \Lambda} \text{BT}_{b,\mu,\Lambda'}^{(0,0)},$$

and the closure of $\text{BT}_{b,\mu,\Lambda}^{(0,0)}$ equals $\mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$.

Proof. (i): This follows from Proposition 5.35 (i).

(ii): The reducedness of $\text{BT}_{b,\mu,\Lambda}^{(0,0)}$ is a consequence of Corollary 5.34. The rest of the assertions follow from Proposition 5.35 (ii) and the definition of $\text{BT}_{b,\mu,\Lambda}^{(0,0)}$. ■

We call the equality in Theorem 5.36 (i) the *Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0,0)}$* .

Theorem 5.37. *The non-formally smooth locus of $\mathcal{M}_{b,\mu}^{(0,0)}$ equals $\prod_{\Lambda \in \text{VL}(\mathbf{L}_{b,0,1})} \mathcal{M}_{b,\mu,\Lambda}^{(0,0)}$.*

Proof. This follows from the same argument as the proof of [Oki19, Theorem 5.28] by using Proposition 5.11 and Corollary 5.34. ■

Corollary 5.38.

- (i) *Each irreducible component of $\mathcal{M}_{b,\mu}^{(0,0)}$ contains exactly $(p+1)(p^2+1)$ -non-formally smooth points of $\mathcal{M}_{b,\mu}^{(0,0)}$.*
- (ii) *Each non-formally smooth point of $\mathcal{M}_{b,\mu}^{(0,0)}$ is contained exactly $(p+1)(p^2+1)$ -irreducible components.*

(iii) Let $C \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)})$. For $C' \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)}) \setminus \{C\}$, the intersection $C \cap C'$ is either the empty set, a single point or isomorphic to \mathbb{P}^1 . Moreover, the following hold:

$$\begin{aligned} \#\{C' \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)}) \mid C \cap C' \text{ is a single point}\} &= p(p+1)(p^2+1), \\ \#\{C' \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)}) \mid C \cap C' = \mathbb{P}_{\mathbb{F}_p}^1\} &= (p+1)(p^2+1) \end{aligned}$$

Proof. These follow from the Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0,0)}$ and Theorem 5.37. \blacksquare

6. BRUHAT–TITS STRATIFICATION IN THE NEUTRAL CASE, II

In this section, we give the definition of the vertex lattices in C_b , and give a connection between them and the vertex lattices in $\mathbf{L}_{b,0}$. Moreover, we construct a Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0)}$ indexed by the vertex lattices in C_b .

6.1. Vertex lattices, II.

Definition 6.1. An O_F -lattice T in C_b is called a *vertex lattice* if we have $T \subset T^\vee \subset \varpi^{-1}T$. We call $t(T) := \dim_{\mathbb{F}_p}(T^\vee/T)$ the *type* of T .

We denote by VL_b the set of vertex lattices. By [RTW14, Lemma 3.2, Lemma 3.3], we have

$$\text{VL}_b = \text{VL}_b(0) \sqcup \text{VL}_b(2) \sqcup \text{VL}_b(4),$$

where $\text{VL}_b(t) := \{T \in \text{VL}_b \mid t(T) = t\}$ for $t \in \{0, 2, 4\}$.

We introduce a partial order \leq on VL_b . For distinct $T, T' \in \text{VL}_b$, we have $T < T'$ if one of the following holds:

- $T \in \text{VL}_b(4)$, $T' \in \text{VL}_b(0)$ and $T \subset T'$,
- $T \in \text{VL}_b(4)$, $T' \in \text{VL}_b(2)$ and $T \subset (T')^\vee$,
- $T \in \text{VL}_b(0)$, $T' \in \text{VL}_b(2)$ and $T \subset (T')^\vee$.

We can interpret vertex lattices in C_b by means of the Bruhat–Tits building \mathcal{B}_b of $J_b^{\text{der}}(\mathbb{Q}_p) = \text{SU}(C_b)(\mathbb{Q}_p)$.

Proposition 6.2. ([RTW14, Proposition 3.4]) *There is a $J_b^{\text{der}}(\mathbb{Q}_p)$ -equivariant bijection between $\text{VL}_b(0) \sqcup \text{VL}_b(4)$ and the set of vertices in \mathcal{B}_b satisfying the following.*

- (i) *The set $\text{VL}_b(4)$ corresponds to the set of special vertices in \mathcal{B}_b .*
- (ii) *If $T_0, T_1 \in \text{VL}_b(0) \sqcup \text{VL}_b(4)$ are distinct, then the corresponding vertices in \mathcal{B}_b are adjacent if and only if either one of the following hold:*
 - $T_i \in \text{VL}_b(0)$, $T_{1-i} \in \text{VL}_b(4)$ and $T_{1-i} \subset T_i$ for some $i \in \{0, 1\}$,
 - $T_0, T_1 \in \text{VL}_b(4)$ and $T_0 + T_1 \in \text{VL}_b(2)$.

In particular, $\text{VL}_b(2)$ corresponds to the set of midpoints of edges connecting two adjacent special vertices in \mathcal{B}_b .

Let \mathbf{T}_0 be the $\mathbf{F}_{b,0}$ -fixed part of M_0 (note that M_0 is $\mathbf{F}_{b,0}$ -stable), and put

$$\text{VL}_{b,\mathbf{T}_0}(4) := \{T \in \text{VL}_b(4) \mid \text{length}_{O_F}(T + \mathbf{T}_0)/\mathbf{T}_0 \in 2\mathbb{Z}\}.$$

We also introduce an order \preceq on $\text{VL}_b(0) \sqcup \text{VL}_b(4)$. For distinct $T, T' \in \text{VL}_b(0) \sqcup \text{VL}_b(4)$, we have $T \prec T'$ if one of the following hold:

- $T \in \text{VL}_{b,\mathbf{T}_0}(4)$, $T' \in \text{VL}_b(0)$ and $T \subset T'$,
- $T \in \text{VL}_{b,\mathbf{T}_0}(4)$, $T' \in \text{VL}_b(4) \setminus \text{VL}_{b,\mathbf{T}_0}(4)$ and $T \subset \varpi^{-1}T'$,
- $T \in \text{VL}_b(0)$, $T' \in \text{VL}_b(4) \setminus \text{VL}_{b,\mathbf{T}_0}(4)$ and $T \subset \varpi^{-1}T'$.

Proposition 6.3. *There is a $J_b^{\text{der}}(\mathbb{Q}_p)$ -equivariant isomorphism of ordered sets*

$$(\text{VL}_b(0) \sqcup \text{VL}_b(4), \preceq) \cong (\text{VL}(\mathbf{L}_{b,0}), \subset)$$

which induces the following bijections:

$$\text{VL}_{b,\mathbf{T}_0}(4) \cong \text{VL}(\mathbf{L}_{b,0}, 1), \quad \text{VL}_b(0) \cong \text{VL}(\mathbf{L}_{b,0}, 3), \quad \text{VL}_b(4) \setminus \text{VL}_{b,\mathbf{T}_0}(4) \cong \text{VL}(\mathbf{L}_{b,0}, 5).$$

Proof. Since $J_b^{\text{der}} \rightarrow \text{SO}(\mathbf{L}_{b,0})$ is a central isogeny by Corollary 5.8, [BT84, §4.2.15] implies that there is a $J_b^{\text{der}}(\mathbb{Q}_p)$ -equivariant isomorphism of simplicial complexes

$$\mathcal{B}_b \cong \mathcal{B}(\mathbf{L}_{b,0}).$$

We may assume that \mathbf{T}_0 corresponds to a vertex lattice in $\mathbf{L}_{b,0}$ of type 1. Then, we obtain the desired isomorphism by Propositions 5.22 and 6.2. \blacksquare

6.2. Bruhat–Tits stratification, II.

Lemma 6.4. *Let $T \in \text{VL}_b$, and $a \in 1 + \varpi O_F$ a norm 1 element. Then there is $h \in J^1(\mathbb{Q}_p)$ such that $\det_F(h) = a$, $h(T) = T$.*

Proof. By Hilbert Satz 90, there is $t \in F^\times$ so that $a = t^{-1}\bar{t}$. By assumption on a , we may assume $t \in O_F^\times$. Take a basis of C_b whose Gram matrix with respect to \langle, \rangle is the anti-diagonal matrix of size 4 that has 1 at every non-zero entry. Then $h := \text{diag}(t, 1, 1, t^{-1}) \in J^1(\mathbb{Q}_p)$ satisfies $\det_F(h) = a$ and $h(T) = T$. \blacksquare

Definition 6.5. For $T \in \text{VL}_b$, we define a closed formal subscheme $\mathcal{M}_{b,\mu,T}^{(0)}$ of $\mathcal{M}_{b,\mu}^{(0)}$ as follows. Take $h \in J_b(\mathbb{Q}_p)$ satisfying $\kappa_{J_b}(h) = (0, 1)$.

- If $T \in \text{VL}_{b,\mathbf{T}_0}(4)$, then let $\mathcal{M}_{b,\mu,T}^{(0)} := \mathcal{M}_{b,\mu,\Lambda(T)}^{(0,0)}$.
- If $T \in \text{VL}_b(4) \setminus \text{VL}_{b,\mathbf{T}_0}(4)$, then let $\mathcal{M}_{b,\mu,T}^{(0)} := h \left(\mathcal{M}_{b,\mu,\Lambda(h^{-1}(T))}^{(0,0)} \right)$.
- If $T \in \text{VL}_b(0)$, then let $\mathcal{M}_{b,\mu,T}^{(0)} := \mathcal{M}_{b,\mu,\Lambda(T)}^{(0,0)} \sqcup h \left(\mathcal{M}_{b,\mu,\Lambda(h^{-1}(T))}^{(0,0)} \right)$.
- If $T \in \text{VL}_b(2)$, then let $\mathcal{M}_{b,\mu,T}^{(0)} := \mathcal{M}_{b,\mu,\Lambda(T_0)}^{(0,0)} \sqcup h \left(\mathcal{M}_{b,\mu,\Lambda(h^{-1}(T_1))}^{(0,0)} \right)$. Here $T_0 \in \text{VL}_b(4) \setminus \text{VL}_{b,\mathbf{T}_0}(4)$ and $T_1 \in \text{VL}_{b,\mathbf{T}_0}(4)$ satisfying $T_0 + T_1 = T$ (note that T_0 and T_1 are unique by Lemma 6.2 (ii)).

We also define a locally closed subscheme $\text{BT}_{b,\mu,T}^{(0)}$ of $\mathcal{M}_{b,\mu}^{(0)}$ by the same manner as above.

Note that $\mathcal{M}_{b,\mu,T}^{(0)}$ and $\text{BT}_{b,\mu,T}^{(0)}$ are independent of the choice of h by Lemma 6.4 and the equality $h(\mathcal{M}_{b,\mu,T'}^{(0)}) = \mathcal{M}_{b,\mu,h(T')}^{(0)}$ for $h \in J_b^{\text{der}}(\mathbb{Q}_p)$ and $T' \in \text{VL}(0) \sqcup \text{VL}(4)$.

Theorem 6.6.

(i) *We have*

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \coprod_{T \in \text{VL}_b} \text{BT}_{b,\mu,T}^{(0)}.$$

(ii) *For $T \in \text{VL}_b$, $\mathcal{M}_{b,\mu,T}^{(0)}$ and $\text{BT}_{b,\mu,T}^{(0)}$ are reduced schemes over $\overline{\mathbb{F}}_p$. Moreover, we have*

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} = \coprod_{T' \leq T} \text{BT}_{b,\mu,T'}^{(0)}.$$

(iii) *Let $T \in \text{VL}_b$.*

- *if $t(T) = 4$, then $\mathcal{M}_{b,\mu,T}^{(0)}$ is a single point,*

- if $t(T) = 0$, then $\mathcal{M}_{b,\mu,T}^{(0)}$ is isomorphic to two copies of $\mathbb{P}_{\mathbb{F}_p}^1$,
 - if $t(T) = 2$, then $\mathcal{M}_{b,\mu,T}^{(0)}$ is isomorphic to two copies of the Fermat surface F_p .
- (iv) There is a bijection between $\text{Irr}(\mathcal{M}_{b,\mu}^{(0),\text{red}})$ and $\text{VL}(4)$.
- (v) The non-formally smooth locus of $\mathcal{M}_{b,\mu}^{(0)}$ equals $\coprod_{T \in \text{VL}_b(4)} \mathcal{M}_{b,\mu,T}$.

We call the equality in Theorem 6.6 by the Bruhat–Tits stratification of $\mathcal{M}_{b,\mu}^{(0)}$.

Proof. The assertions (i)–(iii) follow from Theorem 5.36 and the definitions of $\mathcal{M}_{b,\mu,T}^{(0)}$ and $\text{BT}_{b,\mu,T}^{(0)}$. The assertion (iv) is a consequence of Corollary 5.31. The assertion (v) follows from Theorem 5.37. ■

Finally, we give a connection between the Bruhat–Tits strata of $\mathcal{M}_{b,\mu}^{(0)}$ and some generalized Deligne–Lusztig varieties for split even special orthogonal groups appeared in Proposition 4.7. For $T \in \text{VL}_b$, put

$$\mathbb{B}_b(T) := \begin{cases} T/\varpi T^\vee & \text{if } t(T) \in \{0, 4\}, \\ T^\vee/\varpi T & \text{if } t(T) = 2. \end{cases}$$

We regard it as a quadratic space over \mathbb{F}_p by $(x, y) \mapsto (\varpi x, y) \bmod p$. Then $\mathbb{B}_b(T)$ is a split quadratic space over \mathbb{F}_p of dimension

$$\begin{cases} 0 & \text{if } t(T) = 4, \\ 4 & \text{if } t(T) = 0, \\ 6 & \text{if } t(T) = 2. \end{cases}$$

Theorem 6.7. *Let $T \in \text{VL}_b$.*

- (i) *There is an isomorphism $\mathcal{M}_{b,\mu,T}^{(0)} \cong S_{\mathbb{B}_b(T)}$ (see Definition 4.2 for the definition of the right-hand side).*
- (ii) *If $t(T) = 0$, then there is an isomorphism $\text{BT}_{b,\mu,T}^{(0)} \cong X_{P_{4,0}^+}(t_4^-) \sqcup X_{P_{4,0}^-}(t_4^+)$.*
- (iii) *If $t(T) = 2$, then there is an isomorphism $\text{BT}_{b,\mu,T}^{(0)} \cong X_{P_{6,1}^+}(t_6^- s_{6,1}) \sqcup X_{P_{6,1}^-}(t_6^+ s_{6,1})$.*

In particular, every connected component of $\text{BT}_{b,\mu,T}^{(0)}$ is isomorphic to a generalized Deligne–Lusztig variety for $\text{SO}(\mathbb{B}_b(T))$, which is non-classical if $T \in \text{VL}_b(0) \sqcup \text{VL}_b(2)$.

Proof. The assertion (i) for $t(T) = 0$ is a consequence of Theorem 6.6 (iii). The remaining assertions follow from Propositions 4.7, 5.33 and [Oki19, Corollary 6.10]. ■

7. BRUHAT–TITS STRATIFICATION IN THE NON-NEUTRAL CASE

In this section, we define the notion of vertex lattices, and construct a stratification of $\mathcal{M}_{b,\mu}^{(0)}$ by means of the vertex lattices for a non- μ -neutral b . The method is almost the same as [Wu16] except for the handling of connected components.

Lemma 7.1. *For $M \in \text{DiL}^\varpi(N_{b,W(k)})$, we have $\text{length}_{O_{F,W(k)}}(M + \mathbf{F}_{b,0}(M))/M = 1$.*

Proof. This follows from Proposition 2.10 and the non- μ -neutralness of b . ■

Definition 7.2. A vertex lattice in C_b is an O_F -lattice T in C_b satisfying $T \subset T^\vee \subset \varpi^{-1}T$. We call $t(T) := \dim_{O_F}(T^\vee/T)$ as the type of T .

Proposition 7.3. *For $M \in \text{DiL}^\varpi(N_b)$, there is the minimum integer $r \in \{1, 2\}$ such that*

$$M^{(r)} := M + \mathbf{F}_{b,0}(M) + \cdots + \mathbf{F}_{b,0}^r(M)$$

is $\mathbf{F}_{b,0}$ -stable. Then we have $M^{(r)} \in \text{VL}_b(2r)$ and $(M^{(r)})^\vee = \{x \in M \mid \mathbf{F}_{b,0}(x) = x\}$.

Proof. This follows from the same argument as the proof of [RTW14, Proposition 4.3] (by considering dual lattices) by using Lemma 7.1. \blacksquare

We denote by M_0 the Dieudonné module of \mathbb{X}_0 in N_b .

Definition 7.4. Let $T \in \text{VL}_b$.

- (i) We denote by $(X_T, \iota_T, \lambda_T)$ the p -divisible group with (G, μ) -structure over $\overline{\mathbb{F}}_p$ corresponding to $T \otimes_{\mathbb{Z}_p} W$ by the covariant Dieudonné theory. Then λ_T is an isogeny whose kernel is contained in $\text{Ker}(\iota_T(\varpi))$. Moreover, there is a quasi-isogeny $\rho_T: \mathbb{X}_0 \rightarrow X_T$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathbb{X}_0 & \xrightarrow{\lambda_0} & \mathbb{X}_0^\vee \\ \rho_T \downarrow & & \uparrow \rho_T^\vee \\ X_T & \xrightarrow{\lambda_T} & X_T^\vee. \end{array}$$

- (ii) We define $\mathcal{M}_{b,\mu,T}^{(0)}$ as the locus $(X, \iota, \lambda, \rho)$ of $\mathcal{M}_{b,\mu}^{(0)}$ where the quasi-isogeny

$$X \times_S \overline{S} \xrightarrow{\rho} \mathbb{X}_0 \otimes_{\overline{\mathbb{F}}_p} \overline{S} \xrightarrow{\rho_T \otimes \text{id}_{\overline{S}}} X_T \otimes_{\overline{\mathbb{F}}_p} \overline{S}$$

lift to an isogeny. It is a closed formal subscheme of $\mathcal{M}_{b,\mu}^{(0)}$.

By definition, $\mathcal{M}_{b,\mu,T}^{(0)}(k)$ corresponds to the set $\{M \in \text{DiL}^\varpi(M_{0,W(k)}) \mid M \subset T \otimes_{\mathbb{Z}_p} W(k)\}$ under the bijection $\mathcal{M}_{b,\mu}^{(0)}(k) \cong \text{DiL}^\varpi(M_{0,W(k)})$ for any field extension $k/\overline{\mathbb{F}}_p$.

We study the structure of $\mathcal{M}_{b,\mu,T}^{(0)}$ for $T \in \text{VL}_b$. For $T \in \text{VL}_b$, put $\mathbb{B}_b(T) := T/\varpi T^\vee$. We regard it as a quadratic space over \mathbb{F}_p by $(x, y) \mapsto (\varpi x, y) \bmod p$. Then it is a non-split quadratic space of dimension $4 - t(T)$ over \mathbb{F}_p .

Proposition 7.5. *For $T \in \text{VL}_b$, there is an isomorphism $\mathcal{M}_{b,\mu,T}^{(0),\text{red}} \cong S_{\mathbb{B}_b(T)}$ (see Definition 4.2 for the definition of the right-hand side).*

Proof. Let $(X, \iota, \lambda, \rho) \in \mathcal{M}_{b,\mu,T}^{(0)}(S)$ where $S \in \text{Nilp}$ is reduced (in particular, we have $S = \overline{S}$). Then $(\rho_T \circ \rho)^\vee$ is an isogeny whose kernel is contained in $\text{Ker}(\iota_T(\varpi) \circ \lambda_T^{-1})$ (note that $\iota_T(\varpi) \circ \lambda_T^{-1}$ is an isogeny). By [VW11, Corollary 4.7], $\text{Ker}(D((\rho_T \circ \rho)^\vee))$ is an \mathcal{O}_S -direct summand of

$$\text{Ker}(D(\iota_T(\varpi) \circ \lambda_T^{-1}))_S = \mathbb{B}_b(T) \otimes_{\mathbb{F}_p} \mathcal{O}_S$$

(here we use $S = \overline{S}$). Moreover, it is of rank $2 - t(T)/2$ since $\text{height}(\iota_T(\varpi) \circ \lambda_T^{-1}) = 2 \text{height}((\rho_T \circ \rho)^\vee)$. Hence we obtain a morphism

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} \rightarrow \text{OGr}(\mathbb{B}_b(T))$$

By Lemma 7.1, this morphism induces a morphism

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} \rightarrow S_{\mathbb{B}_b(T), \overline{\mathbb{F}}_p},$$

which is bijective on the sets of k -valued points for any field extension $k/\overline{\mathbb{F}}_p$. Since $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is projective and $S_{\mathbb{B}_b(T),\overline{\mathbb{F}}_p}$ is smooth, the induced morphism is an isomorphism by the same argument as the proof of [HP14, Theorem 3.9]. \blacksquare

Theorem 7.6. *Let $T \in \text{VL}_b$.*

- (i) *If $t(T) = 2$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ consists of two points.*
- (ii) *If $t(T) = 0$, then $\mathcal{M}_{b,\mu,T}^{(0),\text{red}}$ is isomorphic to two copies of $\mathbb{P}_{\mathbb{F}_p}^1$.*

Proof. These follow from Proposition 7.5 and [HP14, Section 3.3]. \blacksquare

Proposition 7.7.

- (i) *We have an equality*

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \bigcup_{T \in \text{VL}_b} \mathcal{M}_{b,\mu,T}^{(0),\text{red}}.$$

- (ii) *For $T, T' \in \text{VL}_b$, we have $\mathcal{M}_{b,\mu,T'}^{(0)} \subset \mathcal{M}_{b,\mu,T}^{(0)}$ if and only if $T' \subset T$.*

Proof. These follow from the same argument as the proof of [Vol10, Proposition 2.5] by using Proposition 7.3 and Theorem 7.6. \blacksquare

For $T \in \text{VL}_b$, set

$$\text{BT}_{b,\mu,T}^{(0)} := \mathcal{M}_{b,\mu,T}^{(0),\text{red}} \setminus \bigcup_{T' \subsetneq T} \mathcal{M}_{b,\mu,T'}^{(0),\text{red}}.$$

Theorem 7.8.

- (i) *We have an equality*

$$\mathcal{M}_{b,\mu}^{(0),\text{red}} = \prod_{T \in \text{VL}_b} \text{BT}_{b,\mu,T}^{(0)}.$$

- (ii) *For $T, T' \in \text{VL}_b$, we have*

$$\mathcal{M}_{b,\mu,T}^{(0),\text{red}} = \prod_{T' \subset T} \text{BT}_{b,\mu,T'}^{(0)}.$$

Proof. The proof is the same as Theorem 5.36 by using Proposition 7.7. \blacksquare

The equality in Theorem 7.8 is called the *Bruhat–Tits stratification* of $\mathcal{M}_{b,\mu}^{(0)}$.

For $j \in \mathbb{Z}/2$, we define a subset $\text{DiL}_{M_0,j}^{\varpi}(k)$ of $\text{DiL}^{\varpi}(M_0, W(k))$ and a subset $\mathcal{M}_{b,\mu}^{(0,j)}(k)$ of $\mathcal{M}_{b,\mu}^{(0)}$ by the same way as Definition 5.9 for any field extension $k/\overline{\mathbb{F}}_p$. Moreover, we define a subspace $\mathcal{M}_{b,\mu}^{(0,j)}$ of $M_{b,\mu}^{(0)}$ by the same manner as Definition 5.28.

Lemma 7.9. *For $T \in \text{VL}_b$, we have a decomposition into open and closed subsets*

$$\mathcal{M}_{b,\mu,T}^{(0)} = \mathcal{M}_{b,\mu,T}^{(0,0)} \sqcup \mathcal{M}_{b,\mu,T}^{(0,1)},$$

where $\mathcal{M}_{b,\mu,T}^{(0,j)} := \mathcal{M}_{b,\mu,T}^{(0)} \cap \mathcal{M}_{b,\mu}^{(0,j)}$. In particular, the decomposition as above becomes as schemes.

Proof. This follows from the definition of $\mathcal{M}_{b,\mu,T}^{(0,j)}$ and Proposition 7.5. \blacksquare

Corollary 7.10. *Let $C \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)})$. For $C' \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)}) \setminus \{C\}$, the intersection $C \cap C'$ is either the empty set or a single point. Moreover, we have*

$$\#\{C' \in \text{Irr}(\mathcal{M}_{b,\mu}^{(0,0)}) \mid C \cap C' \text{ is a single point}\} = p(p^2 + 1)$$

Proof. This follows from Proposition 7.7 and Lemma 7.9. ■

Next, we study the connected components of $\mathcal{M}_{b,\mu}^{(0),\text{red}}$.

Lemma 7.11. *Any irreducible component of $\mathcal{M}_{b,\mu}^{(0),\text{red}}$ is of the form $\mathcal{M}_{b,\mu,T}^{(0,0),\text{red}}$ or $\mathcal{M}_{b,\mu,T}^{(0,1)}$, where $T \in \text{VL}_b(0)$.*

Proof. This follows from Proposition 7.7 and Lemma 7.9. ■

Theorem 7.12. *The subsets $\mathcal{M}_{b,\mu}^{(0,0),\text{red}}$ and $\mathcal{M}_{b,\mu}^{(0,1),\text{red}}$ are the connected components of $\mathcal{M}_{b,\mu}^{(0),\text{red}}$.*

Proof. The proof is the same as Theorem 5.29 by using Lemma 7.11. ■

Finally, we give a relation between the Bruhat–Tits strata of $\mathcal{M}_{b,\mu}^{(0)}$ and some Deligne–Lusztig varieties for even special orthogonal groups appeared in Proposition 4.8.

Theorem 7.13. *For $T \in \text{VL}_b$, there is an isomorphism $\text{BT}_{b,\mu,T}^{(0)} \cong X_{B_{d_b(T)}}(w_{d_b(T)}^+) \sqcup X_{B_{d_b(T)}}(w_{d_b(T)}^-)$. In particular, $\text{BT}_{b,\mu,T}^{(0)}$ is isomorphic to the disjoint union of two Deligne–Lusztig varieties for $\text{SO}(\mathbb{B}_b(T))$ associated with Coxeter elements.*

Proof. This follows from Propositions 4.8 and 7.5. ■

Remark 7.14. We use the notations in Section 2.1. We have $G \otimes_{\mathbb{Q}_p} F \cong \text{GL}_{4,F} \otimes_F \mathbb{G}_{m,F}$. Set

$$\mu_1: \mathbb{G}_{m,F} \rightarrow G \otimes_{\mathbb{Q}} F; z \mapsto (\text{diag}(1, 1, 1, z), z).$$

Then a non- μ -neutral b is μ_1 -neutral. Moreover, we have $\mathcal{M}_{b,\mu} = \mathcal{M}_{b,\mu_1}$, where \mathcal{M}_{b,μ_1} is the Rapoport–Zink space for (G, b, μ_1) defined by the same manner as $\mathcal{M}_{b,\mu}$. In [Wu16], he first determined the connected components $\mathcal{M}_{b,\mu_1}^{(0,0)}$ and $\mathcal{M}_{b,\mu_1}^{(0,1)}$ of $\mathcal{M}_{b,\mu_1}^{(0)}$, and constructed the Bruhat–Tits stratification of $\mathcal{M}_{b,\mu_1}^{(0,0)}$. However, it seems to need a modification for determining $\mathcal{M}_{b,\mu_1}^{(0,j)}$.

Remark 7.15. Our method in this section can be generalized for the Rapoport–Zink space for $\text{GU}(1, n-1)$ with even n and a neutral acceptable b appeared in [Wu16]. The results are exactly the same as [Wu16, Theorem 1].

8. SUPERSINGULAR LOCI OF UNITARY SHIMURA VARIETIES

In this section, we study the supersingular loci of Shimura varieties for unitary similitude group of signature $(2, 2)$ associated to an imaginary quadratic field L over an odd prime p which ramifies in L .

8.1. Rapoport–Zink integral models of unitary Shimura varieties. Let $L = \mathbb{Q}(\sqrt{\Delta})$ be an imaginary quadratic field, where $\Delta \in \mathbb{Z}_{<0}$ is square-free. We denote by $a \mapsto \bar{a}$ the non-trivial Galois automorphism of L/\mathbb{Q} . Throughout of this section, we fix an injection $\varphi: L \hookrightarrow \overline{\mathbb{Q}}$ and an isomorphism $\overline{\mathbb{Q}_p} \cong \mathbb{C}$. Let $(U, \langle \cdot, \cdot \rangle_U)$ be an L/\mathbb{Q} -hermitian space of signature $(2, 2)$. For each prime ℓ , put

$$\text{inv}_\ell(U) := \det(\langle x_i, x_j \rangle_U)_{1 \leq i, j \leq 4} \in \mathbb{Q}_\ell^\times / N_{L_\ell/\mathbb{Q}_\ell}(L_\ell^\times),$$

where $L_\ell := L \otimes_{\mathbb{Q}} \mathbb{Q}_\ell$ and x_1, \dots, x_4 is an L_ℓ -basis of $U_\ell := U \otimes_{\mathbb{Q}} \mathbb{Q}_\ell$. Note that U_ℓ is split as an L_ℓ/\mathbb{Q}_ℓ -hermitian space if and only if $\text{inv}_\ell(U) = 1$.

Let $p > 2$ be an odd prime number. Here we assume that p ramifies in L and $\text{inv}_p(U) = 1$.

Put

$$(\cdot, \cdot)_U := \frac{1}{2} \text{Tr}_{L/\mathbb{Q}}(\langle \cdot, \cdot \rangle_U).$$

Then we have $(ax, y)_U = (x, \bar{a}y)_U$ for $x, y \in U$ and $a \in L$. We define an algebraic group \mathbf{G} over \mathbb{Q} by

$$\mathbf{G}(R) = \{(g, c) \in \mathrm{GL}_{L \otimes_{\mathbb{Q}} R}(U \otimes_{\mathbb{Q}} R) \times R^\times \mid (g(x), g(y))_U = c(x, y)_U \text{ for any } x, y \in U\}$$

for any \mathbb{Q} -algebra R . It is a reductive connected group over \mathbb{Q} . Note that $\mathbf{G} \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is quasi-split over \mathbb{Q} since $\mathrm{inv}_p(U) = 1$.

Definition 8.1. Let S be an \mathbb{Z}_p -scheme. A *polarized L -abelian variety of signature $(2, 2)$ over S* is a triple (A, ι, λ) , where

- A is an abelian variety of dimension 4 over S ,
- $\iota: O \rightarrow \mathrm{End}(A)$ is a ring homomorphism,
- $\lambda: A \rightarrow A^\vee$ is a polarization,

satisfying the following conditions for any $a \in O_L$:

- $\det(T - \iota(a) \mid \mathrm{Lie}(A)) = (T - \varphi(a))^2(T - \varphi(\bar{a}))^2$,
- $\lambda \circ \iota(a) = \iota(\bar{a})^\vee \circ \lambda$.

From now on, we fix an order O of L which is maximal at p , and put $O_p := O \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Let K^p be a sufficiently small compact open subgroup of $\mathbf{G}(\mathbb{A}_f^p)$, where \mathbb{A}_f^p is the finite adèle ring without p -component. On the other hand, let K_p be a special maximal parahoric subgroup of $\mathbf{G}(\mathbb{Q}_p)$. More precisely, it is a stabilizer of a $\sqrt{\Delta}$ -modular O_p -lattice in U_p .

Definition 8.2. We define an \mathbb{Z}_p -scheme $\mathcal{S}_{K,U}$ as the moduli space of the functor which parametrizes equivalence classes of $(A, \iota, \lambda, \bar{\eta}^p)$ for any noetherian scheme S , where

- (A, ι, λ) is a polarized L -abelian variety of signature $(2, 2)$ over S satisfying

$$\mathrm{ord}_p(\#\ker(\lambda)) = \mathrm{ord}_p(\#\ker(\iota(\sqrt{\Delta}))),$$

- $\bar{\eta}^p$ is a K^p -level structure, that is, a $\pi_1(S, \bar{s})$ -invariant K^p -orbit of an isomorphism

$$H_1(A_{\bar{s}}, \mathbb{A}_f^p) \xrightarrow{\cong} U \otimes_{\mathbb{Q}} \mathbb{A}_f^p$$

which respects $(L \otimes_{\mathbb{Q}} \mathbb{A}_f^p)/\mathbb{A}_f^p$ -hermitian forms up to an $(\mathbb{A}_f^p)^\times$ -multiple. Here \bar{s} is a geometric point of S .

Two quadruples $(A, \iota, \lambda, \bar{\eta}^p)$ and $(A', \iota', \lambda', (\bar{\eta}')^p)$ are equivalent if there is a prime-to- p isogeny $A \rightarrow A'$ which respects O_L -actions, polarizations and K^p -level structures.

Set $\mathbb{S} := \mathrm{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$. Fix a \mathbb{C} -basis e_1, \dots, e_4 of $U \otimes_{\mathbb{Q}} \mathbb{R}$ whose Gram matrix of $\langle \cdot, \cdot \rangle_U$ is $\mathrm{diag}(1, 1 - 1, -1)$, and regard $\mathbf{G} \otimes_{\mathbb{Q}} \mathbb{R}$ as a subgroup of $\mathrm{Res}_{\mathbb{C}/\mathbb{R}} \mathrm{GL}_4$ by the above basis. Let \mathbf{X} the $\mathbf{G}(\mathbb{R})$ -conjugacy class in $\mathrm{Hom}(\mathbb{S}, \mathbf{G} \otimes_{\mathbb{Q}} \mathbb{R})$ containing

$$h_U: \mathbb{S} \rightarrow \mathbf{G} \otimes_{\mathbb{Q}} \mathbb{R}; z \mapsto \mathrm{diag}(z^{(2)}, \bar{z}^{(2)}).$$

Then (\mathbf{G}, \mathbf{X}) is a Shimura datum whose reflex field is \mathbb{Q} . Moreover, the generic fiber of $\mathcal{S}_{K,U}$ equals $\mathrm{Sh}_K(\mathbf{G}, \mathbf{X}) \otimes_{\mathbb{Q}} \mathbb{Q}_p$. This follows from that the Hasse principal holds for \mathbf{G} , which is pointed out in [Vol10, Section 6.3].

8.2. Non-emptiness of the basic Newton strata. Put $F := L_p$, $O_F := O \otimes_{\mathbb{Z}} \mathbb{Z}_p$ and $V := U_p$. Set $G := \mathbf{G} \otimes_{\mathbb{Q}} \mathbb{Q}_p$, and let μ be the cocharacter of G over \mathbb{Q}_p associated to the cocharacter

$$\mathbb{G}_m \xrightarrow{z \mapsto (z, 1)} \mathbb{G}_m \otimes_{\mathbb{C}} \mathbb{G}_m \cong \mathbb{S} \otimes_{\mathbb{R}} \mathbb{C} \xrightarrow{h_{U, \mathbb{C}}} \mathbf{G} \otimes_{\mathbb{Q}} \mathbb{C}$$

and the fixed isomorphism $\overline{\mathbb{Q}}_p \cong \mathbb{C}$. Then the datum (F, O_F, V, G, μ) coincides with the one defined in Section 2.1. Hence we can apply the results in Sections 2–7.

We study the Newton strata of the geometric special fiber $\mathcal{S}_{K,U,\overline{\mathbb{F}}_p} := \mathcal{S}_{K,U} \otimes_{\mathbb{Z}_p} \overline{\mathbb{F}}_p$ of $\mathcal{S}_{K,U}$.

Definition 8.3. For $[b] \in B(G)$, we define $\mathcal{S}_{K,U,[b]}$ as the a locally closed subscheme of $\mathcal{S}_{K,U,\overline{\mathbb{F}}_p}$ by

$$\mathcal{S}_{K,U,[b]}(k) = \{(A, \iota, \lambda, \overline{\eta}^p) \in \mathcal{S}_{K,U}(k) \mid \mathbb{D}(A)_{\mathbb{Q}} \cong N_b \text{ for some (all) } b \in [b]\}$$

for any algebraically closed field k of characteristic p . We call $\mathcal{S}_{K,U,[b]}$ the *Newton stratum* of $\mathcal{S}_{K,U,\overline{\mathbb{F}}_p}$ with respect to $[b]$.

The Newton stratum $\mathcal{S}_{K,U,[b]}$ is non-empty only if $\text{ord}_p(\text{sim}_G(b)) = 1$ for some (all) $b \in [b]$. Moreover, it is closed in $\mathcal{S}_{K,U,\overline{\mathbb{F}}_p}$ if $[b] \in B(G)_{1,\text{basic}}$.

Let b_0 and b_1 be elements of $G(K_0)$ defined in Section 2.4. We have $[b_j] \in B(G)_{1,\text{basic}}$ for $j \in \{0, 1\}$. We prove the following:

Proposition 8.4. *The Newton strata $\mathcal{S}_{K,U,[b_0]}$ and $\mathcal{S}_{K,U,[b_1]}$ are non-empty.*

To prove Proposition 8.4, we introduce the notion of invariants for abelian varieties with \mathbf{G} -structures, which are essentially defined in [KR15, Section 3].

Definition 8.5. Let k be an algebraically closed field of characteristic p , and (A, ι, λ) a polarized L -abelian variety with signature $(2, 2)$ over k .

- For a prime $\ell \neq p$, consider the rational Tate module $V_{\ell}A$ of A . Then ι and λ induces an L_{ℓ} -action and a non-degenerate alternating form $(,)_{\ell}$ on $V_{\ell}A$. Moreover, there is an $L_{\ell}/\mathbb{Q}_{\ell}$ -hermitian form \langle , \rangle_{ℓ} on $V_{\ell}A$ satisfying the same formula as $(,)_U$. Put

$$\text{inv}_{\ell}(A, \iota, \lambda) := \det_{L_{\ell}}(\langle x_i, x_j \rangle_{\ell})_{1 \leq i, j \leq 4} \in \mathbb{Q}_{\ell}^{\times} / N_{L_{\ell}/\mathbb{Q}_{\ell}}(L_{\ell}^{\times}),$$

where x_1, \dots, x_4 is an L_{ℓ} -basis of $V_{\ell}A$. It is independent of the choice of x_1, \dots, x_4 .

- For a prime p , we consider the rational Dieudonné module $\mathbb{D}(A)_{\mathbb{Q}}$ with L -action and non-degenerate alternating form $(,)_p$, cf. Section 2.2 (i). Then there is an $(L \otimes_{\mathbb{Z}} W(k))/\text{Frac } W(k)$ -hermitian form \langle , \rangle_p on $\mathbb{D}(A)_{\mathbb{Q}}$ satisfying the same relation as $(,)_U$. Consider the isocrystal

$$(S(A), \mathbf{F}_{S(A)}) := \left(\bigwedge_{L \otimes_{\mathbb{Z}} W(k)}^4 \mathbb{D}(A)_{\mathbb{Q}}, \bigwedge_{L \otimes_{\mathbb{Z}} W(k)}^4 \mathbf{F} \right)$$

with the $(L \otimes_{\mathbb{Z}} W(k))/\text{Frac } W(k)$ -hermitian form $\langle , \rangle_{S(A)}$ associated to \langle , \rangle_p . Then there is $z \in S(A) \setminus \{0\}$ satisfying $\mathbf{F}_{S(A)}(z) = p^2 z$. Now put

$$\text{inv}_p(A, \iota, \lambda) := \langle z, z \rangle_{S(A)} \in \mathbb{Q}_p^{\times} / N_{L_p/\mathbb{Q}_p}(L_p^{\times}).$$

It is independent of the choice of z .

It suffices to show the following:

Proposition 8.6.

- There is a polarized O_L -abelian variety $(A_0, \iota_0, \lambda_0)$ of signature $(2, 2)$ over $\overline{\mathbb{F}}_p$ such that $\text{inv}_{\ell}(A_0, \iota_0, \lambda_0) = \text{inv}_{\ell}(U)$ for any prime ℓ .
- There is a polarized O_L -abelian variety $(A_1, \iota_1, \lambda_1)$ of signature $(2, 2)$ over $\overline{\mathbb{F}}_p$ such that $\text{inv}_{\ell}(A_1, \iota_1, \lambda_1) = \text{inv}_{\ell}(U)$ for any prime $\ell \neq p$, and $\text{inv}_p(A_1, \iota_1, \lambda_1) \neq \text{inv}_p(U)$.

Proposition 8.6 \Rightarrow *Proposition 8.4*. For $j \in \{0, 1\}$, let $(A_j, \iota_j, \lambda_j)$ be as in Proposition 8.6. By replacing A_j to an isogeneous one, we may assume that $\mathbb{D}(A_j)$ is a ϖ -modular lattice in $\mathbb{D}(A_j)_{\mathbb{Q}}$.

Note that there is an isomorphism of isocrystals with G -structures $\mathbb{D}(A_j)_{\mathbb{Q}} \cong N_{b_j}$ over $\overline{\mathbb{F}}_p$. This follows from the equality of invariants at p .

We construct an isomorphism $\eta^p: H_1(A_j, \mathbb{A}_f^p) \xrightarrow{\cong} U \otimes_{\mathbb{Q}} \mathbb{A}_f^p$ for $j \in \{0, 1\}$. Let S be the set of primes ℓ satisfying at least one of the following:

- ℓ ramifies in L ,
- $O \otimes_{\mathbb{Z}} \mathbb{Z}_{\ell}$ is not a maximal order in L_{ℓ} ,
- $\text{inv}_{\ell}(V) \neq 1$,
- $\ell \mid \deg(\lambda)$,
- $\ell = 2$.

Put $\widehat{\mathbb{Z}}^S := \prod_{\ell \notin S} \mathbb{Z}_{\ell}$. Then there is a lattice Λ_0 in U and an isomorphism of $(O_L \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}^S) / \widehat{\mathbb{Z}}^S$ -hermitian spaces

$$\eta_j^S: \prod_{\ell \notin S} T_{\ell} A \xrightarrow{\cong} \Lambda_0 \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}^S.$$

On the other hand, there is an isometry of $L_{\ell}/\mathbb{Q}_{\ell}$ -hermitian spaces

$$\eta_{j,\ell}: V_{\ell} A_j \xrightarrow{\cong} U_{\ell}$$

by the equalities of invariants at all $\ell \in S \setminus \{p\}$. Combining with η_j^S and $\eta_{j,\ell}$ for $\ell \in S \setminus \{p\}$, we obtain an isomorphism of $(L \otimes_{\mathbb{Q}} \mathbb{A}_f^p) / \mathbb{A}_f^p$ -hermitian spaces

$$\eta_j^p: H_1(A_j, \mathbb{A}_f^p) \xrightarrow{\cong} U \otimes_{\mathbb{Q}} \mathbb{A}_f^p.$$

Therefore $(A_j, \iota_j, \lambda_j, \eta_j^p \bmod K^p)$ gives an $\overline{\mathbb{F}}_p$ -valued point of $\mathcal{S}_{K,U,[b_j]}$. ■

Proof of Proposition 8.6. (i): Let E be a supersingular elliptic curve over $\overline{\mathbb{F}}_p$. Fix an injection $\iota_E: O_L \rightarrow \text{End}(E)$ (note that $\text{End}(E)$ is a maximal order of the quaternion algebra over \mathbb{Q} which is ramified exactly at p and ∞) and a principal polarization λ_E of E . Then we have $\lambda_E \circ \iota_E(a) = \iota_E(\bar{a})^{\vee} \circ \lambda_E$ for any $a \in O_L$.

There is $c \in \mathbb{Q}^{\times}$ so that $c = \text{inv}_{\ell}(U)$ in $\mathbb{Q}_{\ell}^{\times} / N_{L_{\ell}/\mathbb{Q}_{\ell}}(L_{\ell}^{\times})$ for any prime ℓ of \mathbb{Q} . In particular, we have $c > 0$ and $c \in N_{L_p/\mathbb{Q}_p}(L_p^{\times})$. By replacing c to ce for some $e \in N_{L/\mathbb{Q}}(L^{\times})$ if necessary, we may assume that $\text{ord}_p(c) = 0$. Write $c = c_1/c_2$, where $c_i \in \mathbb{Z}_{>0}$ and $p \nmid c_i$. Then, $(E^4, \iota_E^4, (c_1 \lambda_E) \times (c_2 \lambda_E)^3)$ is a desired polarized L -abelian 4-fold over $\overline{\mathbb{F}}_p$.

(ii): Write $\text{Hom}(L, \overline{\mathbb{Q}}_p) = \{\varphi_1, \varphi_2\}$. By the theory of complex multiplication [ST61], for $i \in \{1, 2\}$, there is an elliptic curve \mathcal{E}_i over \mathbb{Z}_p with an action ι_i on \mathcal{E}_i such that $\mathcal{E}_{i,\overline{\mathbb{F}}_p}$ is supersingular, and L acts on $\text{Lie}(\mathcal{E}_{i,\overline{\mathbb{Q}}_p})$ by φ_i . Put $\mathcal{A} := \mathcal{E}_1 \times \mathcal{E}_2$, and define an action ι of O_L on \mathcal{A} by $\iota_1 \times \iota_2$. Then we have

$$\det(T - \iota(a) \mid \text{Lie}(\mathcal{A}_{\mathbb{Q}_p})) = T^2 - \text{Tr}_{L/\mathbb{Q}}(a)T + N_{L/\mathbb{Q}}(a)$$

for any $a \in O_L$. By the same argument as the proof of [KR15, Lemma 3.5], there is a polarization λ of \mathcal{A} such that $\ker(\lambda) = \ker(\iota(\sqrt{\Delta}))$. Note that we have $\text{inv}_{\ell}(\mathcal{A}_{\overline{\mathbb{F}}_p}, \iota_{\overline{\mathbb{F}}_p}, \lambda_{\overline{\mathbb{F}}_p}) = 1$ for any prime ℓ . On the other hand, by the same argument in the proof of (i), there is a polarized L -abelian surface (A', ι', λ') over $\overline{\mathbb{F}}_p$ such that $\text{inv}_{\ell}(A', \iota', \lambda') = \text{inv}_{\ell}(U)$ for any prime $\ell \neq p$ and $\text{inv}_p(A', \iota', \lambda') \neq \text{inv}_p(U)$. Hence $(\mathcal{A}_{\overline{\mathbb{F}}_p} \times A', \iota_{\overline{\mathbb{F}}_p} \times \iota', \lambda_{\overline{\mathbb{F}}_p} \times \lambda')$ is a desired polarized O_L -abelian 4-fold over $\overline{\mathbb{F}}_p$. ■

8.3. Structure of the supersingular loci. We define the supersingular locus $\mathcal{S}_{K,U}^{\text{ss}}$ as the reduced closed subscheme of $\mathcal{S}_{K,U,\overline{\mathbb{F}}_p}$ satisfying

$$\mathcal{S}_{K,U}^{\text{ss}}(k) = \{(A, \iota, \lambda, \overline{\eta}^p) \in \mathcal{S}_{K,U}(k) \mid A \text{ is supersingular}\}$$

for any algebraically closed field k of characteristic p .

Theorem 8.7.

(i) *There is a decomposition into open and closed subschemes*

$$\mathcal{S}_{K,U} = \mathcal{S}_{K,U}^{\text{loc}} \sqcup (\mathcal{S}_{K,U} \setminus \mathcal{S}_{K,U}^{\text{loc}}).$$

(ii) *The scheme $\mathcal{S}_{K,U}^{\text{loc}}$ is flat over \mathbb{Z}_p and regular of dimension 5.*

(iii) *The scheme $\mathcal{S}_{K,U} \setminus \mathcal{S}_{K,U}^{\text{loc}}$ is smooth over \mathbb{F}_p and 3-dimensional.*

(iv) *The non-smooth locus of $\mathcal{S}_{K,U}^{\text{loc}} \otimes_{\mathbb{Z}_p} W$ is contained in $\mathcal{S}_{K,U}^{\text{ss}}$.*

Proof. These follow from the same argument as [Oki19, Theorem 7.5] using Proposition 3.3 and [Hai05, Theorem 6.4]. ■

Theorem 8.8. *The supersingular locus $\mathcal{S}_{K,U}^{\text{ss}}$ equals $\mathcal{S}_{K,U,[b_0]} \sqcup \mathcal{S}_{K,U,[b_1]}$.*

Proof. This follows from the equality $B(G)_{1,\text{basic}} = \{[b_0], [b_1]\}$ proved in Section 2.4 and Proposition 2.8. ■

We give a connection between $\mathcal{S}_{K,U}^{\text{ss}}$ the Rapoport–Zink spaces $\mathcal{M}_{b_0,\mu}$ and $\mathcal{M}_{b_1,\mu}$. For $j \in \{0, 1\}$, let $\widehat{\mathcal{S}}_{K,U,[b_j]}$ be the completion of $\mathcal{S}_{K,U} \otimes_{\mathbb{Z}_p} W$ along $\mathcal{S}_{K,U,[b_j]}$.

Proposition 8.9. *Let $j \in \{0, 1\}$. Then there is an isomorphism*

$$I_j(\mathbb{Q}) \backslash (\mathcal{M}_{b_j,\mu} \times \mathbf{G}(\mathbb{A}_f^p) / K^p) \xrightarrow{\cong} \widehat{\mathcal{S}}_{K,U,[b_j]}$$

of formal schemes over $\text{Spf } W$. Here I_j is an algebraic group over \mathbb{Q} defined by

$$I_j(R) = \{(g, c) \in (\text{End}_{O_L}(A_j) \otimes_{\mathbb{Z}} R) \times R^\times \mid g^\vee \circ \lambda_j \circ g = c\lambda_j\}$$

for any \mathbb{Q} -algebra R .

Proof. If we can apply the p -adic uniformization theorem [RZ96, Theorem 6.30], then the assertion follows. For this, it suffices to show the non-emptiness of $\mathcal{S}_{K,U,[b_j]}$ and that the Hasse principle for \mathbf{G} holds. However, the first assertion follows from Proposition 8.4, and the second assertion is already pointed out in Section 8.1. ■

Now we can prove the theorems on explicit descriptions of basic loci in Section 1.1 by using Proposition 8.9 and the result on the Rapoport–Zink spaces.

Theorem 8.10.

(i) *The scheme $\mathcal{S}_{K,U,[b_0]}$ is purely 2-dimensional. Any irreducible component is birational to the Fermat surface*

$$x_0^{p+1} + x_1^{p+1} + x_2^{p+1} + x_3^{p+1} = 0$$

in $\mathbb{P}_{\mathbb{F}_p}^3 = \text{Proj } \overline{\mathbb{F}_p}[x_0, x_1, x_2, x_3]$.

(ii) *Let C be an irreducible component of $\mathcal{S}_{K,U,[b_0]}$. Then, for any irreducible component $C' \neq C$ of $\mathcal{S}_{K,U,[b_0]}$, the intersection $C \cap C'$ is either the empty set, a single point or birational to $\mathbb{P}_{\mathbb{F}_p}^1$. Moreover, the following hold:*

$$\#\{C' \in \text{Irr}(\mathcal{S}_{K,U,[b_0]}) \setminus \{C\} \mid C \cap C' \text{ is a single point}\} \leq p(p+1)(p^2+1),$$

$$\#\{C' \in \text{Irr}(\mathcal{S}_{K,U,[b_0]}) \setminus \{C\} \mid C \cap C' \text{ is birational to } \mathbb{P}_{\mathbb{F}_p}^1\} \leq (p+1)(p^2+1).$$

(iii) *Each $C \in \text{Irr}(\mathcal{S}_{K,U,[b_0]})$ contains at most $(p+1)(p^2+1)$ -non-smooth points of $\mathcal{S}_{K,U}$.*

- (iv) *Each non-smooth point of $\mathcal{S}_{K,U}$ is contained in at most $(p+1)$ -irreducible components of $\mathcal{S}_{K,U,[b_0]}$.*

Proof. This follows from the same argument as the proof of [Oki19, Theorem 7.15] by using Theorems 8.9, 6.6 and Corollary 5.38. \blacksquare

Theorem 8.11.

- (i) *The number of connected components of $\mathcal{S}_{K,U,[b_0]}$ equals the cardinality of*

$$I_0(\mathbb{Q}) \backslash (J_{b_0}^{0,0} \backslash J_{b_0}(\mathbb{Q}_p) \times \mathbf{G}(\mathbb{A}_f^p) / K^p),$$

where $J_{b_0}^{0,0} := \{g \in J_{b_0}^0(\mathbb{Q}_p) \mid \text{ord}_p(\text{sim}(g)) = 0\}$.

- (ii) *The number of irreducible components of $\mathcal{S}_{K,U,[b_0]}$ equals the cardinality of*

$$I_0(\mathbb{Q}) \backslash (J_{b_0}(\mathbb{Q}_p) / K_{b_0,4} \times \mathbf{G}(\mathbb{A}_f^p) / K^p),$$

where $K_{b_0,4}$ is the stabilizer of a vertex lattice in $\text{VL}_{b_0}(4)$.

- (iii) *The number of non-smooth points of $\mathcal{S}_{K,U}$ equals the cardinality of*

$$I_0(\mathbb{Q}) \backslash (J_{b_0} / K_{b_0,4} \times \mathbf{G}(\mathbb{A}_f^p) / K^p),$$

where $K_{b_0,4}$ is the same as one in (ii).

Proof. (i): The proof is the same as [Oki19, Theorem 7.13 (i)] by using Theorems 8.9 and 5.29.

(ii): The proof is the same as [Oki19, Theorem 7.13 (ii)] by using Theorems 8.9 and 6.6 (iv).

(iii): This follows from the same argument as the proof of [Oki19, Theorem 7.13 (iii)] by using Theorems 8.9 and 6.6 (v). \blacksquare

Theorem 8.12.

- (i) *The scheme $\mathcal{S}_{K,U,[b_1]}$ is purely 1-dimensional. Any irreducible component is birational to $\mathbb{P}_{\mathbb{F}_p}^1$.*

- (ii) *Let C be an irreducible component of $\mathcal{S}_{K,U,[b_1]}$. Then, for any irreducible component $C' \neq C$ of $\mathcal{S}_{K,U,[b_1]}$, the intersection $C \cap C'$ is either the empty set or a single point. Moreover, we have*

$$\#\{C' \in \text{Irr}(\mathcal{S}_{K,U,[b_1]}) \setminus \{C\} \mid C \cap C' \text{ is a single point}\} \leq p(p^2 + 1).$$

Proof. This follows from the same argument as the proof of [Oki19, Theorem 7.15] by using Theorems 8.9, 7.6 and Corollary 7.10. \blacksquare

Theorem 8.13.

- (i) *The number of connected components of $\mathcal{S}_{K,U,[b_1]}$ equals the cardinality of*

$$I_1(\mathbb{Q}) \backslash (J_{b_1}^{0,0} \backslash J_{b_1}(\mathbb{Q}_p) \times \mathbf{G}(\mathbb{A}_f^p) / K^p),$$

where $J_{b_1}^{0,0} := \{g \in J_{b_1}^0(\mathbb{Q}_p) \mid \text{ord}_p(\text{sim}(g)) = 0\}$.

- (ii) *The number of irreducible components of $\mathcal{S}_{K,U,[b_1]}$ equals the cardinality of*

$$I_1(\mathbb{Q}) \backslash (J_{b_1}(\mathbb{Q}_p) / K_{b_1,0} \times \mathbf{G}(\mathbb{A}_f^p) / K^p),$$

where $K_{b_1,0}$ is the stabilizer of a vertex lattice in $\text{VL}_{b_1}(0)$.

Proof. (i): The proof is the same as [Oki19, Theorem 7.13 (i)] by using Theorems 8.9 and 7.12.

(ii): This follows from the same argument as the proof of [Oki19, Theorem 7.13 (ii)] by using Theorem 8.9 and Corollary 7.11. \blacksquare

REFERENCES

- [AN02] P. Abramenko, G. Nebe, *Lattice chain models for affine buildings of classical groups*, Math. Ann. **322** (2002), 537–562.
- [BR06] C. Bonnafé, R. Rouquier, *On the irreducibility of Deligne-Lusztig varieties*, C. R. Math. Acad. Sci. Paris **343**, vol. 1 (2006), 37–39.
- [BT84] F. Bruhat, J. Tits, *Groupes réductifs sur un corps local, II: Schémas en groupes. Existence d’une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. **60** (1984), 197–376.
- [Cho18] S. Cho, *The basic locus of the unitary Shimura variety with parahoric level structure, and special cycles*, preprint, arXiv:1807.09997, 2018.
- [Fox18] M. Fox, *The GL_4 -Rapoport–Zink space*, to appear in Int. Math. Res. Not., arXiv:1811.02040, 2018.
- [Hai05] T. J. Haines, *Introduction to Shimura varieties with bad reduction of parahoric type*, in *Harmonic analysis, the trace formula, and Shimura varieties, Proc. Clay Mathematics Institute 2003 Summer School, The Fields Institute Toronto, 2–27, June 2003*, Clay Mathematical Proceedings, vol. 4, eds J. Arthur, D. Ellwood and R. E. Kottwitz (American Mathematical Society/Clay Mathematics Institute, Providence, RI/Cambridge, MA, 2005), 583–658.
- [HP14] B. Howard, G. Pappas, *On the supersingular locus of $GU(2, 2)$ Shimura variety*, Algebra Number Theory **8** (2014), 1659–1699.
- [HP17] B. Howard, G. Pappas, *Rapoport–Zink spaces for spinor groups*, Compos. Math. **153** (2017), 1050–1118.
- [Jac62] R. Jacobowitz, *Hermitian forms over local fields*, Amer. J. Math. **84** (1962), 441–465.
- [Kne65] M. Kneser, *Galois-Kohomologie halbeinfacher algebraischer Gruppen über p -adischen Körpern, I*, Math. Z. **88** (1965), 40–47, II, *ibid* **89** (1965), 250–272.
- [Kot85] R. E. Kottwitz, *Isocrystals with additional structure*, Compos. Math. **56** (1985), 201–220.
- [Kot92] R. E. Kottwitz, *Points on some Shimura varieties over finite fields*, J. Amer. Math. Soc. **5** (1992), no. 2, 373–444.
- [KR14] S. S. Kudla, M. Rapoport, *Special cycles on unitary Shimura varieties II: global theory*, J. Reine Angew. Math. **697** (2014), 91–157.
- [KR15] S. S. Kudla, M. Rapoport, *New cases of p -adic uniformization*, Astérisque **370** (2015), 207–241.
- [LZ19] C. Li, W. Zhang, *Kudla–Rapoport cycles and derivatives of local densities*, to appear in J. Amer. Math. Soc., preprint, arXiv:1908.01701, 2019.
- [Oki19] Y. Oki, *On supersingular loci of Shimura varieties for quaternionic unitary groups of degree 2*, to appear in manuscripta math, arXiv:1907.07026, 2019.
- [PR09] G. Pappas, M. Rapoport, *Local models in the ramified case III. unitary groups*, J. Inst. Math. Jessieu **8** (2009), no. 3, 507–564.
- [RR96] M. Rapoport, M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, Compos. Math. **103** (1996), no. 2, 153–181.
- [RSZ17] M. Rapoport, B. Smithling, W. Zhang, *On the arithmetic transfer conjecture for exotic smooth formal moduli spaces*, Duke Math. J. **166** (2017), no. 12, 2183–2336.
- [RTW14] M. Rapoport, U. Terstiege, S. Wilson, *The supersingular locus of the Shimura variety for $GU(1, n - 1)$ over a ramified prime*, Math. Z. **276** (2014), no. 3–4, 1165–1188.
- [RV14] M. Rapoport, E. Viehmann, *Towards a theory of local Shimura varieties*, Münster J. Math. **7** (2014), 273–326.
- [RZ96] M. Rapoport, Th. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996.
- [Shi10] G. Shimura, *Arithmetic of quadratic forms*, Springer Monographs in Mathematics, Springer, New York, 2010.
- [ST61] G. Shimura, Y. Taniyama, *Complex multiplication of abelian varieties and its applications to number theory*, Publ. Math. Soc. Japan, no. 6, 1961.
- [Smi14] B. Smithling, *Topological flatness of local models for ramified unitary groups II. the even dimensional case*, J. Inst. Math. Jessieu **13** (2014), no. 2, 303–393.
- [Tit79] J. Tits, *Reductive groups over local fields*, in *Automorphic forms, representations and L -functions (Proc. Symp. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, 29–69, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979.

- [Vol10] I. Vollaard, *The supersingular locus of the Shimura variety for $GU(1, s)$* , *Canad. J. Math.* **62** (2010), no. 3, 668–720.
- [VW11] I. Vollaard, T. Wedhorn, *The supersingular locus of the Shimura variety for $GU(1, n - 1)$ II*, *Invent. Math.* **184** (2011), no. 3, 591–627.
- [Wu16] H. Wu, *The supersingular locus of the Shimura varieties with exotic good reduction*, preprint, arXiv:1609.08775, 2016.
- [Wan20] H. Wang, *On the Bruhat–Tits stratification of a quaternionic unitary Rapoport–Zink space*, *Math. Ann.* **376** (2020), no. 3–4, 1107–1144.
- [Wan21] H. Wang, *On the Bruhat–Tits stratification for $GU(2, 2)$ type Rapoport–Zink space: unramified case*, *Ann. Math. Qué.* **45** (2021), no. 1, 19–44.
- [Zin01] Th. Zink, *Windows for displays of p -divisible groups*, pp. 491–518 in *Moduli of abelian varieties* (Texel, 1999), edited by C. Faber et al., *Progr. Math.* **195**, Birkhäuser, Basel, 2001.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN.

Email address: oki@ms.u-tokyo.ac.jp