

# $n$ -exact Character Graphs

Mahdi Ebrahimi\*

*School of Mathematics, Institute for Research in Fundamental Sciences (IPM),  
P.O. Box: 19395-5746, Tehran, Iran*

## Abstract

Let  $\Gamma$  be a finite simple graph. If for some integer  $n \geq 4$ ,  $\Gamma$  is a  $K_n$ -free graph whose complement has an odd cycle of length at least  $2n - 5$ , then we say that  $\Gamma$  is an  $n$ -exact graph. For a finite group  $G$ , let  $\Delta(G)$  denote the character graph built on the set of degrees of the irreducible complex characters of  $G$ . In this paper, we prove that the order of an  $n$ -exact character graph is at most  $2n - 1$ . Also we determine the structure of all finite groups  $G$  with extremal  $n$ -exact character graph  $\Delta(G)$ .

**Keywords:** Character graph, Character degree,  $n$ -exact graph, Odd cycle.  
**AMS Subject Classification Number:** 20C15, 05C38, 05C25.

## 1 Introduction

Let  $G$  be a finite group and  $R(G)$  be the solvable radical of  $G$ . Also let  $\text{cd}(G)$  be the set of all character degrees of  $G$ , that is,  $\text{cd}(G) = \{\chi(1) \mid \chi \in \text{Irr}(G)\}$ , where  $\text{Irr}(G)$  is the set of all complex irreducible characters of  $G$ . The set of prime divisors of character degrees of  $G$  is denoted by  $\rho(G)$ . It is well known that the character degree set  $\text{cd}(G)$  may be used to provide information on the structure of the group  $G$ . For example, a result due to Berkovich [5] says that if a prime  $p$  divides every non-linear character degree of a group  $G$ , then  $G$  is solvable.

A useful way to study the character degree set of a finite group  $G$  is to associate a graph to  $\text{cd}(G)$ . One of these graphs is the character graph  $\Delta(G)$  of  $G$  [13]. Its vertex set is  $\rho(G)$  and two vertices  $p$  and  $q$  are joined by an edge

---

\*m.ebrahimi.math@ipm.ir

if the product  $pq$  divides some character degree of  $G$ . We refer the readers to a survey by Lewis [10] for results concerning this graph and related topics.

There is an interesting conjecture on the structure of  $\Delta(G)$  which is posed by Akhlaghi and Tong-Viet in [4]:

**Conjecture.** Let  $G$  be a finite group such that for some positive integer  $n$ ,  $\Delta(G)$  is  $K_n$ -free. If  $G$  is solvable, then  $|\rho(G)| \leq 2n - 2$ , and if  $G$  is non-solvable, then  $|\rho(G)| \leq 2n - 1$ .

In [1], it was shown that for a solvable group  $G$ , the complement of  $\Delta(G)$  does not have an odd cycle, in other word, it is bipartite. This result guarantees this conjecture for solvable groups. The conjecture still is not proved for non-solvable groups, however the results in [15], [4] and [3] say that it is true when  $n \in \{3, 4, 5\}$ . Now let  $\Gamma$  be a finite simple graph. If for some integer  $n \geq 4$ ,  $\Gamma$  is a  $K_n$ -free graph whose complement has an odd cycle of length at least  $2n - 5$ , then we say that  $\Gamma$  is an  $n$ -exact graph. For example,  $K_4$ -free graphs with non-bipartite complement are 4-exact. In this paper, we wish to show that the conjecture is true for a finite group whose character graph is  $n$ -exact, for some integer  $n \geq 4$ .

**Theorem A.** *Let  $G$  be a finite group, and  $n \geq 4$  be an integer. If  $\Delta(G)$  is an  $n$ -exact graph, then there exists a normal subgroup  $R(G) < M \leq G$  so that  $G/R(G)$  is an almost simple group with socle  $S := M/R(G) \cong \text{PSL}_2(q)$ , where  $q$  is a prime power, and  $|\rho(G)| \leq 2n - 1$ .*

The solvable group  $G$  is said to be disconnected if  $\Delta(G)$  is disconnected. Also  $G$  is called of disconnected Type  $n$  if  $G$  satisfies the hypotheses of Example 2.n in [9]. Let  $n \geq 4$  be an integer and  $\Gamma$  be a finite simple graph with vertex set  $V$ . The cardinality of  $V$  is called the order of  $\Gamma$ . An  $n$ -exact graph  $\Gamma$  with minimum or maximum possible order is called an extremal  $n$ -exact graph. Note that for an integer  $n \geq 1$ , the set of prime divisors of  $n$  is denoted by  $\pi(n)$ . Now we are ready to state the next result of this paper.

**Corollary B.** *Let  $G$  be a finite group with extremal  $n$ -exact character graph  $\Delta(G)$ , for some integer  $n \geq 4$ . Then one of the following cases occurs:*

- a)**  $|\rho(G)| = 2n - 5$  and for some integer  $\alpha \geq 2$ ,  $G \cong \text{PSL}_2(2^\alpha) \times R(G)$ , where  $|\pi(2^\alpha \pm 1)| = n - 3$  and  $R(G)$  is abelian.
- b)**  $|\rho(G)| = 2n - 1$  and for some integer  $\alpha \geq 2$ ,  $G \cong \text{PSL}_2(2^\alpha) \times R(G)$ , where  $|\pi(2^\alpha \pm 1)| = n - 3, n - 2$  or  $n - 1$ . Also
- i)** If  $|\pi(2^\alpha \pm 1)| = n - 3$ , then for some disconnected groups  $A$  and  $B$  of disconnected Types 1 or 4,  $R(G) \cong A \times B$ ,  $|\rho(A)| = |\rho(B)| = 2$  and  $\rho(G) = \pi(S) \uplus \rho(A) \uplus \rho(B)$ .

- ii) If  $|\pi(2^\alpha \pm 1)| = n - 2$ , then  $R(G)$  is a disconnected group of disconnected Type 1 or 4,  $|\rho(R(G))| = 2$  and  $\rho(G) = \pi(S) \uplus \rho(R(G))$ .*
- iii) If  $|\pi(2^\alpha \pm 1)| = n - 1$ , then  $R(G)$  is abelian and  $\rho(G) = \pi(S)$ .*

## 2 Preliminaries

In this paper, all groups are assumed to be finite and all graphs are simple and finite. For a finite group  $G$ , the set of prime divisors of  $|G|$  is denoted by  $\pi(G)$ . If  $H \leq G$  and  $\theta \in \text{Irr}(H)$ , we denote by  $\text{Irr}(G|\theta)$  the set of irreducible characters of  $G$  lying over  $\theta$  and define  $\text{cd}(G|\theta) := \{\chi(1) \mid \chi \in \text{Irr}(G|\theta)\}$ . We frequently use, Gallagher's Theorem which is corollary 6.17 of [8].

**Lemma 2.1.** *Let  $N \triangleleft G$  and  $\varphi \in \text{Irr}(N)$ . Then for every  $\chi \in \text{Irr}(G|\varphi)$ ,  $\chi(1)/\varphi(1)$  divides  $[G : N]$ .*

Let  $\Gamma$  be a graph with vertex set  $V(\Gamma)$  and edge set  $E(\Gamma)$ . The complement of  $\Gamma$  and the induced subgraph of  $\Gamma$  on  $X \subseteq V(\Gamma)$  are denoted by  $\Gamma^c$  and  $\Gamma[X]$ , respectively. If  $E(\Gamma) = \emptyset$ ,  $\Gamma$  is called an empty graph. When  $n := |V(\Gamma)|$ , any cycle of  $\Gamma$  of length  $n$  is called a Hamilton cycle. We say that  $\Gamma$  is Hamiltonian if it contains a Hamilton cycle. We use the notations  $K_n$  for a complete graph with  $n$  vertices and  $C_n$  for a cycle of length  $n$ . If for some integer  $n \geq 2$ ,  $\Gamma$  does not contain a copy of  $K_n$  as an induced subgraph, then  $\Gamma$  is called a  $K_n$ -free graph. We now state some relevant results on character graphs needed in the next sections.

**Lemma 2.2.** [1] *Let  $G$  be a solvable group. Then  $\Delta(G)^c$  is bipartite.*

**Lemma 2.3.** [11] *Let  $G$  be a solvable group with  $\Delta(G) \cong C_4$ . Then  $G \cong A \times B$ , where  $A$  and  $B$  are disconnected groups. Also for some distinct primes  $p, q, r$  and  $s$ ,  $\rho(A) = \{p, q\}$  and  $\rho(B) = \{r, s\}$ .*

**Lemma 2.4.** [11] *Let  $G$  be a solvable group. If  $\Delta(G)$  has at least 4 vertices, then either  $\Delta(G)$  contains a triangle or  $\Delta(G) \cong C_4$ .*

The structure of the character graphs of  $\text{PSL}_2(q)$  and  ${}^2B_2(q^2)$  are determined as follows:

**Lemma 2.5.** [17] *Let  $G \cong \text{PSL}_2(q)$ , where  $q \geq 4$  is a power of a prime  $p$ .*

**a)** *If  $q$  is even, then  $\Delta(G)$  has three connected components,  $\{2\}$ ,  $\pi(q - 1)$  and  $\pi(q + 1)$ , and each component is a complete graph.*

**b)** *If  $q > 5$  is odd, then  $\Delta(G)$  has two connected components,  $\{p\}$  and  $\pi((q - 1)(q + 1))$ .*

*i) The connected component  $\pi((q - 1)(q + 1))$  is a complete graph if and only*

if  $q - 1$  or  $q + 1$  is a power of 2.

ii) If neither of  $q - 1$  or  $q + 1$  is a power of 2, then  $\pi((q - 1)(q + 1))$  can be partitioned as  $\{2\} \cup M \cup P$ , where  $M = \pi(q - 1) - \{2\}$  and  $P = \pi(q + 1) - \{2\}$  are both non-empty sets. The subgraph of  $\Delta(G)$  corresponding to each of the subsets  $M, P$  is complete, all primes are adjacent to 2, and no prime in  $M$  is adjacent to any prime in  $P$ .

**Lemma 2.6.** [16] Let  $G \cong^2 B_2(q^2)$ , where  $q^2 = 2^{2m+1}$  and  $m \geq 1$ . The set of primes  $\rho(G)$  can be partitioned as  $\rho(G) = \{2\} \cup \pi(q^2 - 1) \cup \pi(q^4 + 1)$ . The subgraph of  $\Delta(G)$  on  $\rho(G) - \{2\}$  is complete and 2 is adjacent in  $\Delta(G)$  to precisely the primes in  $\pi(q^2 - 1)$ .

When  $\Delta(G)^c$  is not a bipartite graph, then there exists a useful restriction on the structure of  $G$  as follows:

**Lemma 2.7.** [2] Let  $G$  be a finite group and  $\pi$  be a subset of the vertex set of  $\Delta(G)$  such that  $|\pi| > 1$  is an odd number. Then  $\pi$  is the set of vertices of a cycle in  $\Delta(G)^c$  if and only if  $O^{\pi'}(G) = S \times A$ , where  $A$  is abelian,  $S \cong \text{SL}_2(u^\alpha)$  or  $S \cong \text{PSL}_2(u^\alpha)$  for a prime  $u \in \pi$  and a positive integer  $\alpha$ , and the primes in  $\pi - \{u\}$  are alternately odd divisors of  $u^\alpha + 1$  and  $u^\alpha - 1$ .

**Lemma 2.8.** [12] Let  $p$  be a prime,  $f \geq 2$  be an integer,  $q = p^f \geq 5$  and  $S \cong \text{PSL}_2(q)$ . If  $q \neq 9$  and  $S \leq G \leq \text{Aut}(S)$ , then  $G$  has irreducible characters of degrees  $(q + 1)[G : G \cap \text{PGL}_2(q)]$  and  $(q - 1)[G : G \cap \text{PGL}_2(q)]$ .

Now we present some facts on the structure of character graphs with non-bipartite complement.

**Lemma 2.9.** [7] Let  $G$  be a finite group,  $R(G) < M \leq G$ ,  $S := M/R(G)$  be isomorphic to  $\text{PSL}_2(q)$ , where for some prime  $p$  and positive integer  $f \geq 1$ ,  $q = p^f$ ,  $|\pi(S)| \geq 4$  and  $S \leq G/R(G) \leq \text{Aut}(S)$ . Also let  $\theta \in \text{Irr}(R(G))$ . If  $\Delta(G)^c$  is not bipartite, then  $\theta$  is  $M$ -invariant.

**Lemma 2.10.** [7] Assume that  $f \geq 2$  is an integer,  $q = 2^f$ ,  $S \cong \text{PSL}_2(q)$  and  $G$  is a finite group such that  $G/R(G) = S$ . If  $\Delta(G)[\pi(S)] = \Delta(S)$ , then  $G \cong S \times R(G)$ .

**Lemma 2.11.** [6] Let  $G$  be a finite group. The complement of the character graph  $\Delta(G)$  is a non-bipartite Hamiltonian graph if and only if  $G \cong \text{SL}_2(2^f) \times A$ , where  $f \geq 2$  is an integer,  $||\pi(2^f + 1)| - |\pi(2^f - 1)|| \leq 1$  and  $A$  is an abelian group.

We end this section with the following result.

**Lemma 2.12.** [7] Let  $G$  be a disconnected group. If  $|\rho(G)| = 2$  and  $2 \notin \rho(G)$ , then  $G$  is of disconnected Type 1 or 4.

### 3 The proof of Main results

In this section, we wish to prove our main results. When  $G$  is a finite group with 4-exact character graph, then using [4], [7] and Lemma 2.11 we have nothing to prove. Hence in the sequel, we assume that  $G$  is a finite group such that for some integer  $n \geq 5$ ,  $\Delta(G)$  is  $n$ -exact.

**Lemma 3.1.** *Suppose  $G$  is a finite group and  $\Delta(G)$  is an  $n$ -exact graph, for some integer  $n \geq 5$ . Then*

**a)** *There exists a normal subgroup  $R(G) < M \leq G$  so that  $G/R(G)$  is an almost simple group with socle  $S := M/R(G) \cong \text{PSL}_2(u^\alpha)$ , where  $u$  is a prime,  $\alpha$  is a positive integer and  $u^\alpha > 5$ .*

**b)** *For some  $\pi_0 \subseteq \pi(S)$  and integer  $m \geq n - 3$ ,  $\Delta(G)[\pi_0]$  precisely contains three connected components  $\Delta(G)[\{u\}]$  and  $\Delta(G)[\pi(u^\alpha \pm 1) \cap \pi_0] \cong K_m$ .*

**c)** *If  $\pi$  is the set of vertices of an odd cycle in  $\Delta(G)^c$ , then  $u \in \pi \subseteq \pi(S)$ .*

*Now let  $\pi_R := \rho(R(G)) - \pi(G/R(G))$ ,  $\pi_M := \pi([G : M]) - \rho(M)$  and  $\pi_F = \pi_R \cup \pi_M$ .*

**d)**  *$\Delta(G)[\pi_M]$  is a complete graph with at most two vertices and every  $p \in \pi_F$  is adjacent to all vertices in  $\pi(u^{2\alpha} - 1)$ .*

**e)** *If  $u = 2$  and  $\pi([G : M]) \subseteq \pi_F$ , then for every  $\theta \in \text{Irr}(R(G))$ ,  $\Delta(G)[\pi(\theta(1)[G : M](u^\alpha \pm 1))]$  are complete.*

*Proof.* **a,b)** Since  $\Delta(G)$  is an  $n$ -exact graph, for some  $\pi_0 \subseteq \rho(G)$ ,  $\pi_0$  is the set of vertices of an odd cycle in  $\Delta(G)^c$  and  $|\pi_0| \geq 2n - 5$ . By Lemma 2.7,  $N := O^{\pi_0}(G) = R \times A$ , where  $A$  is abelian,  $R \cong \text{SL}_2(u^\alpha)$  or  $R \cong \text{PSL}_2(u^\alpha)$  for a prime  $u \in \pi_0$  and a positive integer  $\alpha$ , and the primes in  $\pi_0 - \{u\}$  are alternately odd divisors of  $u^\alpha + 1$  and  $u^\alpha - 1$ . Hence  $m := |\pi(u^\alpha \pm 1) \cap \pi_0| \geq n - 3$  and  $\Delta(G)[\pi(u^\alpha \pm 1) \cap \pi_0] \cong K_m$ . Assume that for some  $\epsilon \in \{\pm 1\}$ , there exist  $x \in (\pi(u^\alpha + \epsilon) \cap \pi_0) \cup \{u\}$  and  $y \in \pi(u^\alpha - \epsilon) \cap \pi_0$  so that  $x$  and  $y$  are adjacent vertices in  $\Delta(G)$ . Then for some  $\chi \in \text{Irr}(G)$ ,  $xy$  divides  $\chi(1)$ . No let  $\theta \in \text{Irr}(N)$  be a constituent of  $\chi_N$ . By Lemma 2.1,  $xy$  divides  $\theta(1)$ . It is a contradiction. Thus  $\Delta(G)[\{u\}]$  and  $\Delta(G)[\pi(u^\alpha \pm 1) \cap \pi_0]$  are precisely connected components of  $\Delta(G)[\pi_0]$ . Let  $M := NR(G)$ . Then  $S := M/R(G) \cong N/R(N) \cong \text{PSL}_2(u^\alpha)$  is a non-abelian minimal normal subgroup of  $G/R(G)$ .

Let  $C/R(G) = C_{G/R(G)}(M/R(G))$ . We claim that  $C = R(G)$  and  $G/R(G)$  is an almost simple group with socle  $S = M/R(G)$ . Suppose on the contrary that  $C \neq R(G)$  and let  $L/R(G)$  be a chief factor of  $G$  with  $L \leq C$ . Then  $L/R(G) \cong T^k$ , for some non-abelian simple group  $T$  and some integer  $k \geq 1$ . As  $L \leq C$ ,  $LM/R(G) \cong L/R(G) \times M/R(G) \cong S \times T^k$ . Let  $F := \pi(S) \cap \pi(T)$ . Note that  $\pi_0 \cap \rho(T) = \emptyset$ ,  $|\rho(T)| \geq 3$  and  $2 \in F$ . Now one of the following cases occurs:

Case 1.  $|F| = 1$ : Then  $F = \{2\}$ . Hence 3 does not divide  $|T|$ . Therefore by [14],  $T$  is a Suzuki simple group. By Lemma 2.6,  $\Delta(T) \subseteq \Delta(S \times T^k)$  contains a triangle. Thus  $\Delta(S \times T^k)[\pi(u^\alpha + 1) \cup \rho(T)]$  contains a copy of  $K_n$  and it is a contradiction.

Case 2.  $|F| \geq 2$ : Then it is easy to see that  $\Delta(S \times T^k)[\pi(u^\alpha + 1) \cup \rho(T)]$  contains a copy of  $K_n$  and it is again a contradiction.

c) Let  $\pi \subseteq \rho(G)$  be the set of vertices of an odd cycle in  $\Delta(G)^c$ . Then using Lemma 2.7,  $N_1 := O^{\pi'}(G) = R_1 \times A_1$ , where  $A_1$  is abelian,  $R_1 \cong \text{SL}_2(u_1^{\alpha_1})$  or  $R_1 \cong \text{PSL}_2(u_1^{\alpha_1})$  for a prime  $u_1 \in \pi$  and a positive integer  $\alpha_1$ , and the primes in  $\pi - \{u_1\}$  are alternately odd divisors of  $u_1^{\alpha_1} + 1$  and  $u_1^{\alpha_1} - 1$ . Thus  $S_1 := N_1 R(G)/R(G) \cong \text{PSL}_2(u_1^{\alpha_1})$  is a minimal normal subgroup of  $G/R(G)$  and  $u_1 \in \pi \subseteq \pi(S_1)$ . Hence by part (a),  $S = S_1$  and  $u = u_1$ .

d) Using Lemma 2.8, Lemma 2.9 and this fact that  $\text{SL}_2(u^\alpha)$  is the schur representation of  $S$ ,  $\Delta(G)[\pi_M]$  is complete and every  $p \in \pi_F$  is adjacent to all vertices in  $\pi(u^{2\alpha} - 1)$ . If  $|\pi_M| \geq 3$ , then  $\Delta(G)[\pi_M \cup \pi(u^\alpha + 1)]$  contains a copy of  $K_n$  and it is a contradiction. Thus  $|\pi_M| \leq 2$ .

e) Let  $d = [G : M]$ . Then  $G/R(G) \cong S \rtimes \mathbb{Z}_d$ . Note that  $\pi(d) \cap \pi(S) = \emptyset$ . Therefore as the schur multiplier of  $S$  is trivial, the schur multiplier of  $G/R(G)$  is too. Hence by Lemma 2.8 and Gallagher's Theorem, for every  $\theta \in \text{Irr}(R(G))$ ,  $\theta(1)d(u^\alpha \pm 1) \in \text{cd}(G)$  and it completes the proof.  $\square$

**Proof of Theorem A.** By Lemma 3.1 (a,b), there exists a normal subgroup  $R(G) < M \leq G$  so that  $G/R(G)$  is an almost simple group with socle  $S := M/R(G) \cong \text{PSL}_2(u^\alpha)$ , where  $u$  is a prime,  $\alpha$  is a positive integer and  $u^\alpha > 5$ . Also for some  $\pi_0 \subseteq \pi(S)$  and integer  $m \geq n - 3$ ,  $\Delta(G)[\pi_0]$  precisely contains three connected components  $\Delta(G)[\{u\}]$  and  $\Delta(G)[\pi(u^\alpha \pm 1) \cap \pi_0] \cong K_m$ . It is easy to see that  $\rho(G) = \pi_F \uplus \pi(S)$ . Using Lemma 3.1 (d),  $\Delta(G)[\pi_M]$  is a complete graph with at most two vertices and every  $p \in \pi_F$  is adjacent to all vertices in  $\pi(u^{2\alpha} - 1)$ . Now one of the following cases occurs:

Case 1:  $|\pi(u^\alpha \pm 1)| = n - 3$ . Then by Lemma 2.5,  $u = 2$  and  $\Delta(G)[\pi_0] = \Delta(S)$ . We claim that  $|\pi_F| \leq 4$ . On the contrary, we assume that  $|\pi_F| \geq 5$ . Let  $d := [G : M]$ . Then using Lemma 2.8,  $\pi(d) \cap \pi(S) = \emptyset$  and  $\pi(d) \subseteq \pi_F$ . If  $\pi_M \neq \emptyset$ , then as  $|\pi_M| \leq 2$ , using Lemmas 2.2 and 3.1 (e),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + 1)]$  contains a copy of  $K_n$  which is impossible. Thus  $\pi_M = \emptyset$  and  $\pi_F = \pi_R$ . Therefore using Lemma 2.2,  $\Delta(G)[\pi_F]$  contains a triangle and by Lemma 3.1 (e), we again obtain a contradiction. Thus  $|\pi_F| \leq 4$  and  $|\rho(G)| \leq 2n - 1$ .

Case 2:  $|\pi(u^\alpha \pm 1)| \leq n - 2$  and for some  $\epsilon \in \{\pm 1\}$ ,  $|\pi(u^\alpha + \epsilon)| = n - 2$ . If  $|\pi_F| \geq 3$ , then using Lemma 3.1 (c),  $\Delta(G)[\pi_F]$  is a non-empty graph and  $\Delta(G)[\pi_F \cup \pi(u^\alpha + \epsilon)]$  contains a copy of  $K_n$  which is a contradiction. Thus  $|\pi_F| \leq 2$  and  $|\rho(G)| \leq 2n - 1$ .

Case 3: For some  $\epsilon \in \{\pm 1\}$ ,  $|\pi(u^\alpha + \epsilon)| = n - 1$ . Then by Lemma 3.1 (d),

$\pi_F = \emptyset$  and so  $|\rho(G)| \leq 2n - 1$ . □

### 3.1 Proof of Corollary B.

Let  $G$  be a finite group with extremal  $n$ -exact character graph  $\Delta(G)$ , for some integer  $n \geq 5$ . Then using Theorem A,  $|\rho(G)| = 2n - 5$  or  $2n - 1$ . If  $|\rho(G)| = 2n - 5$ , then by Lemma 2.11, we are done. Thus we can assume that  $|\rho(G)| = 2n - 1$ . By Lemma 3.1, there exists a normal subgroup  $R(G) < M \leq G$  so that  $G/R(G)$  is an almost simple group with socle  $S := M/R(G) \cong \text{PSL}_2(u^\alpha)$ , where  $u$  is a prime,  $\alpha$  is a positive integer and  $u^\alpha > 5$ . Also for some  $\pi_0 \subseteq \pi(S)$  and integer  $m \geq n - 3$ ,  $\Delta(G)[\pi_0]$  precisely contains three connected components  $\Delta(G)[\{u\}]$  and  $\Delta(G)[\pi(u^\alpha \pm 1) \cap \pi_0] \cong K_m$ . Since  $\Delta(G)$  is  $K_n$ -free,  $\Delta(S)$  is too. The structure of  $\Delta(S)$  is determined by Lemma 2.5.

**Lemma 3.2.** *Let  $\Delta(G)[\pi_F]$  be a non-empty graph and  $|\pi(S)| \geq 2n - 4$ . Then  $G$  does not exist.*

*Proof.* Since  $|\pi(S)| \geq 2n - 4$ , there exists  $\epsilon \in \{\pm 1\}$  such that  $|\pi(u^\alpha + \epsilon)| \geq n - 2$ . Thus using Lemma 3.1 (d),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + \epsilon)]$  contains a copy of  $K_n$  and it is a contradiction. □

**Lemma 3.3.** *If  $|\pi(u^\alpha \pm 1)| = n - 3$ , then  $u = 2$  and  $G \cong S \times A \times B$ , where  $A$  and  $B$  are disconnected groups of disconnected Types 1 or 4,  $|\rho(A)| = |\rho(B)| = 2$  and  $\rho(G) = \pi(S) \uplus \rho(A) \uplus \rho(B)$ .*

*Proof.* Since  $|\pi(u^\alpha \pm 1)| = n - 3$ ,  $\pi(S) = \pi_0$ . Thus  $|\pi_F| = 4$ ,  $u = 2$  and  $\Delta(G)[\pi(S)] = \Delta(S)$ . If  $d := [G : M]$ , then using Lemma 2.8,  $\pi(d) \cap \pi(S) = \emptyset$  and so  $\pi(d) \subseteq \pi_F$ . Now we claim that  $d = 1$  and  $G = M$ . On the contrary, we assume that  $d \neq 1$ . If  $|\pi(d)| \geq 2$ , then by Lemma 3.1 (e),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + 1)]$  contains a copy of  $K_n$  which is a contradiction. Also if  $|\pi(d)| = 1$ , then using Lemma 2.2,  $\Delta(G)[\pi_R]$  is non-empty and by Lemma 3.1 (e),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + 1)]$  again contains a copy of  $K_n$  and it is a contradiction. Therefore  $d = 1$  and  $G = M$ . Hence by Lemma 2.10,  $G \cong S \times R(G)$ . Thus as  $\Delta(G)[\pi(S)] = \Delta(S)$ ,  $\pi(S) \cap \rho(R(G)) = \emptyset$  and  $\rho(R(G)) = \pi_R = \pi_F$ . Also as  $\Delta(G)$  is  $K_n$ -free,  $\Delta(R(G))$  is  $K_3$ -free. Hence using Lemma 2.4,  $\Delta(R(G)) \cong C_4$ . Thus by Lemma 2.3,  $R(G) \cong A \times B$ , where  $A$  and  $B$  are disconnected groups with  $|\rho(A)| = |\rho(B)| = 2$ . Therefor by Lemma 2.12,  $A$  and  $B$  are of disconnected Types 1 or 4 and the proof is completed. □

**Lemma 3.4.** *If  $|\pi(u^\alpha \pm 1)| = n - 2$  and  $u = 2$ , then  $G \cong S \times R(G)$ , where  $R(G)$  is a disconnected group of disconnected Type 1 or 4. Also  $|\rho(R(G))| = 2$  and  $\rho(G) = \pi(S) \uplus \rho(R(G))$ .*

*Proof.* Since  $|\rho(G)| = 2n - 1$ ,  $|\pi_F| = 2$ . If  $\pi_F = \pi_M$ , then using Lemma 3.1 (d),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + 1)]$  is a copy of  $K_n$  which is a contradiction. Hence  $\pi_R$  is non-empty. Note that there exist  $x \in \pi(u^\alpha + 1)$  and  $y \in \pi(u^\alpha - 1)$  such that  $\pi(u^\alpha + 1) = \pi_0 \cup \{x\}$  and  $\pi(u^\alpha - 1) = \pi_0 \cup \{y\}$ . Let  $d := [G : M]$ . We claim that  $x, y \notin \pi(d)$ . On the contrary, we assume that  $z \in \{x, y\} \cap \pi(d)$ . Then using Lemmas 2.8 and 3.1 (d),  $\Delta(G)[\pi_R \cup \{z\} \cup \pi(u^{2\alpha} - 1)]$  contains a copy of  $K_n$  which is impossible. Thus  $x, y \notin \pi(d)$ . Suppose  $p \in \pi_0 \cap \pi(d)$ . Then by Lemma 2.8,  $p$  is adjacent to all vertices in  $\pi_0 - \{u, p\}$ . It is a contradiction with the structure of  $\Delta(G[\pi_0])$ . Therefore  $\pi(d) \subseteq \pi_F$ . If  $d \neq 1$ , then by Lemma 3.1 (e),  $\Delta(G)[\pi_F \cup \pi(u^\alpha + 1)]$  contains a copy of  $K_n$  which is a contradiction. Thus  $d = 1$  and  $G = M$ . Hence there exist distinct primes  $q$  and  $q'$  such that  $\pi_F = \pi_R = \{q, q'\}$ . Note that as the schur multiplier of  $S$  is trivial, by Gallagher's Theorem,  $q$  and  $q'$  are adjacent to all vertices in  $\pi(S)$ . Now we claim that  $\Delta(G)[\pi(S)] = \Delta(S)$ . On the contrary, we assume that there exist adjacent vertices  $a, b \in \pi(S)$  in  $\Delta(G)$  such that  $a$  and  $b$  are non-adjacent vertices in  $\Delta(S)$ . Then for some  $\chi \in \text{Irr}(G)$ ,  $ab$  divides  $\chi(1)$ . Let  $\theta \in \text{Irr}(R(G))$  be a constituent of  $\chi_{R(G)}$ . Then using Lemma 2.9,  $\theta$  is  $G$ -invariant. Thus as the Schur multiplier of  $S$  is trivial, by Gallagher's Theorem,  $\text{cd}(G|\theta) = \{m\theta(1) \mid m \in \text{cd}(S)\}$ . Hence as  $\chi(1) \in \text{cd}(G|\theta)$ ,  $\theta(1)$  is divisible by  $a$  or  $b$ . Without loss of generality, we assume that  $a$  divides  $\theta(1)$ . There exists  $\epsilon \in \{\pm 1\}$  so that  $a \notin \pi(u^\alpha + \epsilon)$ . Thus the induced subgraph of  $\Delta(G)$  on  $X := \pi(u^\alpha + \epsilon) \cup \{a, q\}$  is a copy of  $K_n$  and it is a contradiction. Therefore  $\Delta(G)[\pi(S)] = \Delta(S)$ . Hence by Lemma 2.10,  $G \cong S \times R(G)$ . If  $\rho(R(G)) \cap \pi(S) \neq \emptyset$ , then the induced subgraph of  $\Delta(G)$  on  $\pi(S) \cup \{q\}$  contains a copy of  $K_n$  and it is a contradiction. Thus as  $\Delta(G)$  is  $K_n$ -free, it is easy to see that  $\rho(R(G)) = \pi_R = \{q, q'\}$  and  $\Delta(R(G)) \cong K_2^c$ . Therefore by Lemma 2.12,  $R(G)$  is a disconnected group of disconnected Type 1 or 4. Finally,  $\rho(G) = \pi(S) \uplus \rho(R(G))$  and the proof is completed.  $\square$

**Lemma 3.5.** *If  $|\pi(u^\alpha \pm 1)| = n - 1$  and  $u = 2$ , then  $G \cong S \times R(G)$ , where  $R(G)$  is abelian and  $\rho(G) = \pi(S)$ .*

*Proof.* Since  $|\rho(G)| = 2n - 1$ ,  $\rho(G) = \pi(S)$ . If  $G \neq M$ , then using Lemma 2.8,  $\Delta(G)$  contains a copy of  $K_n$  which is impossible. Thus  $G = M$ . We claim that  $\Delta(G)[\pi(S)] = \Delta(S)$ . On the contrary, we assume that there exist non-adjacent vertices  $x$  and  $y$  in  $\Delta(S)$  such that  $x$  and  $y$  are adjacent vertices in  $\Delta(G)$ . Thus for some  $\chi \in \text{Irr}(G)$ ,  $\chi(1)$  is divisible by  $xy$ . Let  $\theta \in \text{Irr}(R(G))$  be a constituent of  $\chi_{R(G)}$ . Using Lemma 2.9,  $\theta$  is  $G$ -invariant. Thus as the Schur multiplier of  $S$  is trivial,  $\theta$  extends to  $G$ . By Gallagher's Theorem,  $\text{cd}(G|\theta) = \{m\theta(1) \mid m \in \text{cd}(S)\}$ . Thus as  $\chi(1) \in \text{cd}(G|\theta)$  is divisible by  $xy$ ,  $\theta(1)$  is divisible by  $x$  or  $y$ . Hence one of the character degrees  $\theta(1)(u^\alpha - 1)$  or  $\theta(1)(u^\alpha + 1)$  of  $G$  is divisible by  $n$  distinct primes which is a contradiction.



Therefore  $\Delta(G)[\pi(S)] = \Delta(S)$ . Thus using Lemma 2.10,  $G \cong S \times R(G)$ . Hence as  $\Delta(G)$  is  $K_n$ -free,  $R(G)$  is abelian and we are done.  $\square$

**Lemma 3.6.** *If either  $u = 2$  and  $|\pi(u^\alpha + 1)| \neq |\pi(u^\alpha - 1)|$ , or  $u$  is odd, then  $G$  does not exist.*

*Proof.* By Lemma 2.5, one of the following cases occurs:

Case 1: Either  $|\pi(u^\alpha \pm 1)| = n - 2$  and  $u$  is odd, or for some  $\epsilon \in \{\pm 1\}$ ,  $|\pi(u^\alpha + \epsilon)| = n - 3$  and  $|\pi(u^\alpha - \epsilon)| = n - 2$ : Since  $|\rho(G)| = 2n - 1$ ,  $|\pi_F| = 3$ . Thus by Lemma 3.1 (c),  $\Delta(G)[\pi_F]$  is a non-empty graph. Hence as  $|\pi(S)| = 2n - 4$ , using Lemma 3.2, we are done.

Case 2: for some  $\epsilon \in \{\pm 1\}$ ,  $|\pi(u^\alpha - \epsilon)| = n - 1$  and either  $|\pi(u^\alpha + \epsilon)| \leq n - 2$  and  $u = 2$ , or  $|\pi(u^\alpha + \epsilon)| = n - 2$  or  $n - 1$  and  $u$  is odd: Since  $|\rho(G)| = 2n - 1$ ,  $\pi_F \neq \emptyset$ . Thus using Lemma 3.1 (d), the induced subgraph of  $\Delta(G)$  on  $\pi_F \cup \pi(u^\alpha - \epsilon)$  contains a copy of  $K_n$  and it is a contradiction.  $\square$

## Acknowledgements

This research was supported in part by a grant from School of Mathematics, Institute for Research in Fundamental Sciences (IPM).

## References

- [1] Z. Akhlaghi, C. Casolo, S. Dolfi, K. Khedri and E. Pacifici. On the character degree graph of solvable groups, Proc. Amer. Math. Soc., 146 (2018), 1505-1513.
- [2] Z. Akhlaghi, C. Casolo, S. Dolfi, E. Pacifici, L. Sanus, On the character degree graph of finite groups, Annali di Mat. Pura Appl., (2019), DOI: 10.1007/s10231-019-00833-0.
- [3] Z. Akhlaghi, K. Khedri and B. Taeri, Finite groups with  $K_5$ -free prime graphs, Comm. Algebra, (2018), DOI: 10.1080/00927872.2018.1501576.
- [4] Z. Akhlaghi, H.P. Tong-Viet, Finite groups with  $K_4$ -free prime graphs, Algebr. Represent. Theory, 18(1) (2015), 235-256.
- [5] Y. Berkovich, Finite groups with small sums of degrees of some non-linear irreducible characters, J. Algebra, 171 (1995), 426-443.
- [6] M. Ebrahimi, Character graphs with non-bipartite Hamiltonian complement, Bull. Aust. Math. Soc., (2019), DOI: 10.1017/S0004972719001163.

- [7] M. Ebrahimi,  $k_4$ -free character graphs with seven vertices, *comm. Algebra*, (2019) DOI: 10.1080/00927872.2019.1670197.
- [8] I.M. Isaacs, *Character Theory of Finite Groups*, AMS Chelsea Publishing, Providence, RI, 2006. Corrected reprint of the 1976 original [Academic Press. New York; MR0460423].
- [9] M.L. Lewis, Solvable groups whose degree graphs have two connected components, *J. Group Theory*, 4(3) (2001), 255-275.
- [10] M.L. Lewis, An overview of graphs associated with character degrees and conjugacy class sizes in finite groups, *Rocky Mountain, J. Math.*, 38 (1) (2008), 175-211.
- [11] M.L. Lewis, Q. Meng, Square character degree graphs yield direct products, *J. Algebra*, 349 (2012), 185-200.
- [12] M.L. Lewis, D.L. White, Non-solvable groups with no prime dividing three character degrees, *J. Algebra*, 336 (2011), 158-183.
- [13] O. Manz, R. Staszewski, W. Willems, On the number of components of a graph related to character degrees, *Proc. Amer. Math. Soc.*, 103 (1) (1988), 31-37.
- [14] I. Toborg, R. Waldecker, Finite simple  $3'$ -groups are cyclic or Suzuki groups, *Arch. Math. (Basel)*, 102 (2014), no.4, 301-312.
- [15] H.P. Tong-Viet, Groups whose prime graphs have no triangles, *J. Algebra*, 378 (2013), 196-206.
- [16] D.L. White, Degree graphs of simple groups of exceptional Lie type, *Comm. Algebra*, 32 (9) (2004), 3641-3649.
- [17] D.L. White, Degree graphs of simple linear and unitary groups, *Comm. Algebra*, 34 (8) (2006), 2907-2921.