

SPECTRAL RECIPROCITY VIA INTEGRAL REPRESENTATIONS

RAMON M. NUNES

ABSTRACT. We prove a spectral reciprocity formula for automorphic forms on $GL(2)$ over a number field that is reminiscent of the one found by Blomer and Khan. Our approach uses period representations of L -functions and the language of automorphic representations.

1. INTRODUCTION

In the past few years, some attention has been given to spectral reciprocity formulae. There are two main reasons that help understand this recent interest. The first one is that they give a somewhat conceptual way of summarizing a technique often used in dealing with problems on families of GL_2 L -functions in which one uses the Kuznetsov formula on both directions in order to estimate a moment of these L -functions. The second one comes from their satisfying intrinsic nature relating objects that have no *a priori* reason to be linked.

The first versions of these formulae ([BK17], [AK18] and [BK18]) used Classical techniques such as the Voronoi summation formula and the Kuznetsov formula). Starting from [Zac18], it became clear that an adelic approach could be of interest. Not only this renders generalization to number fields almost immediate, it can also avoid some of the combinatorial difficulties appearing when applying the Voronoi formula.

In this article we use the theory of adelic automorphic representations and integral representations of Rankin-Selberg L -functions to deduce a result on number fields of similar flavor to that of [BK17, Theorem 1]. With respect to Blomer and Khan's result, our result has the advantage of being valid for number fields. On the other hand we need to make some technical restrictions that prevent us from having a full generalization of their result. For the moment our results only work when the fixed GL_3 form is cuspidal (which unfortunately is the less interesting case) and our formula only contemplates forms that are spherical at every infinity place. When $F = \mathbb{Q}$, this corresponds to even Maass forms.

1.1. Statement of results. Let F be a number field, with ring of integers \mathfrak{o}_F . Let Π be a cuspidal automorphic representation of $GL(3)$ over F . For each automorphic representation π of $GL(2)$, we consider the completed L -functions

$$\Lambda(s, \pi), \Lambda(s, \text{Ad}, \pi) \text{ and } \Lambda(s, \Pi \times \pi).$$

These are, respectively, the Hecke L -function and the Adjoint L -function of π , and the Rankin-Selberg L -function of $\Pi \times \pi$.

Date: November 8, 2021.

This work was partially supported by the DFG-SNF lead agency program grant 200021L-153647.

Remark 1.1. *All the Rankin-Selberg L -functions in this article are product of local L -functions that are naively defined as in (14). These coincide with the actual local L -functions à la Langlands at all the unramified places but might differ at the ramified ones.*

Let ξ_F denote the completed Dedekind zeta function of F and let $\xi_F^*(1)$ denote its residue at the pole in $s = 1$. Let \mathcal{A} denote the collection of automorphic representations for $\mathrm{GL}(2)$ over F , both cuspidal and Eisenstein and let $H : \mathcal{A} \rightarrow \mathbb{C}$ denote an admissible weight function as defined in §6.1. We consider the following sums:

$$\mathcal{C}(H) = \sum_{\pi \in \mathcal{C}(S)} \frac{\Lambda(s, \Pi \times \pi) \Lambda(w, \pi)}{\Lambda(1, \mathrm{Ad}, \pi)} H(\pi)$$

and

$$\mathcal{E}(H) = \xi_F^*(1) \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \frac{\Lambda(s, \Pi \times \pi(\omega, it)) \Lambda(w, \pi(\omega, it))}{\Lambda^*(1, \mathrm{Ad}, \pi(\omega, it))} H(\pi(\omega, it)) \frac{dt}{2\pi}, \quad (1)$$

where S is any finite set of places containing all the archimedean ones and those for which H_v is ramified. Moreover, $\mathcal{C}(S)$ (resp. $\Xi(S)$) denotes the collection of cuspidal automorphic representations of $\mathrm{GL}(2)$ (resp. unitary idele characters) over F that are unramified everywhere outside S . Finally, $\pi(\omega, it)$ is as defined in §3.2. The main object of study in this work is the following “moment”:

$$\mathcal{M}(H) := \mathcal{C}(H) + \mathcal{E}(H),$$

We remark that the values of $\Lambda(s, \Pi \times \pi(\omega, it))$, $\Lambda(w, \pi(\omega, it))$ and $\Lambda^*(1, \mathrm{Ad}, \pi(\omega, it))$ can be given in terms of simpler L -functions as follows:

$$\begin{aligned} \Lambda(s, \Pi \times \pi(\omega, it)) &= \Lambda(s + it, \Pi \times \omega) \Lambda(s - it, \Pi \times \bar{\omega}) \\ \Lambda(w, \pi(\omega, it)) &= \Lambda(w + it, \omega) \Lambda(w - it, \bar{\omega}), \\ \Lambda^*(1, \mathrm{Ad}, \pi(\omega, it)) &= \mathrm{Res}_{s=1} [\Lambda(s + 2it, \omega^2) \Lambda(s - 2it, \bar{\omega}^2) \xi_K(s)] \\ &= \Lambda(1 + 2it, \omega^2) \Lambda(1 - 2it, \bar{\omega}^2) \xi_K^*(1), \end{aligned}$$

where $\Lambda(s, \Pi \times \omega)$ and $\Lambda(s, \omega)$ are the completed Rankin-Selberg L -function of $\Pi \times \omega$ and the Dirichlet L -function of ω , respectively.

We start with the following result which can be seen as a pre-spectral reciprocity formula.

Theorem 1.1. *Let $s, w \in \mathbb{C}$ and define*

$$(s', w') := \left(\frac{1 + w - s}{2}, \frac{3s + w - 1}{2} \right). \quad (2)$$

Let H be a (s, w) – admissible weight function and \check{H} be the (s', w') – admissible weight function given by (26). Let S be a finite set of places containing all the archimedean ones and those for which H_v is ramified. Suppose the real parts of s , w , s' and w' are sufficiently large. Then we have the relation

$$\mathcal{M}(H) = \mathcal{M}(\check{H}).$$

Theorem 1.1 is a completely symmetrical formula but only holds when the real part of the parameters s, w, s' and w' are sufficiently large. In order to obtain a formula that also holds at the central point $s = w = s' = w' = \frac{1}{2}$, we need to analytically continue the term $\mathcal{E}(H)$. This is done in section 8 under a technical condition enclosed in Hypothesis 1. In order to ensure that this hypothesis is satisfied we make a simplifying assumptions which, we believe, can be verified by appealing to heavier machinery on the theory of automorphic forms.

We suppose that Φ_v is spherical for every archimedean place v . This restricts the spectral sum to the automorphic representations that are unramified at every *archimedean* place (in the case $F = \mathbb{Q}$, this means that we are turning our attention to even Maass forms). In fact it trivializes, for every archimedean place v , the weight H_v into the characteristic function of unramified representations of $\mathrm{GL}_2(F_v)$.

Spectral Reciprocity. Let Π be an everywhere unramified cuspidal automorphic representation for $\mathrm{GL}(3)$ over F . Let $s, w \in \mathbb{C}$ let \mathfrak{q} and \mathfrak{l} be coprime ideals and finally, we write $U_\infty := \prod_{v|\infty} \{y \in F_v^*; |y_v| = 1\}$. Let H be the (s, w) – admissible weight function with kernel $\Phi = \Phi^{\mathfrak{q}, \mathfrak{l}}$ (see Section 7). We also define the following modified Hecke eigenvalues for every ideal \mathfrak{l} of \mathfrak{o}_F :

$$\widehat{\lambda}_\pi(\mathfrak{l}, w) := \sum_{\mathfrak{a}\mathfrak{b}=\mathfrak{l}} \frac{\mu(\mathfrak{a})}{(N\mathfrak{a})^w} \lambda_\pi(\mathfrak{b}) = \prod_{\substack{\mathfrak{p}^{n_{\mathfrak{p}}} || \mathfrak{l} \\ n_{\mathfrak{p}} \geq 1}} \left(\lambda_\pi(\mathfrak{p}^{n_{\mathfrak{p}}}) - \frac{\lambda_\pi(\mathfrak{p}^{n_{\mathfrak{p}}-1})}{(N\mathfrak{p})^w} \right). \quad (3)$$

It follows from (29), Proposition 7.1 and Proposition 7.3, that we have

$$H(\pi) = \begin{cases} \frac{\widehat{\lambda}_\pi(\mathfrak{l}, w)}{(N\mathfrak{l})^w} \frac{h_{\mathfrak{q}}(s, w; \Pi, \pi)}{N\mathfrak{q}}, & \pi \in \mathcal{A}_{\mathfrak{q}}(\infty), \\ 0, & \text{otherwise.} \end{cases}$$

where $h_{\mathfrak{q}}(s, w; \Pi, \pi) = 1$ if $\mathrm{cond}(\pi) = \mathfrak{q}$ and in general, $h_{\mathfrak{q}}(s, w; \Pi, \pi) \ll (N\mathfrak{q})^{\theta+\epsilon}$. Here and thereafter, θ (not to be confused with ϑ) denotes an approximation towards the Ramanujan conjecture for Π , i.e. we take θ to be any number such that for any place v , the satake parameters α_v^i for Π_v satisfy $|\alpha_{v,i}|_v \leq p_v^\theta$.

Hence, with this choice of Φ , we have

$$\mathcal{M}(H) = \mathcal{M}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}),$$

where

$$\mathcal{M}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) := \mathcal{C}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) + \mathcal{E}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}),$$

with

$$\mathcal{C}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) = \frac{1}{N\mathfrak{q}} \sum_{\substack{\pi \text{ cusp}^0 \\ \mathrm{cond}(\pi) | \mathfrak{q}}} \frac{\Lambda(s, \Pi \times \pi) \Lambda(w, \pi)}{\Lambda(1, \mathrm{Ad}, \pi)} \frac{\widehat{\lambda}_\pi(\mathfrak{l}, w)}{(N\mathfrak{l})^w} h_{\mathfrak{q}}(s, w; \Pi, \pi)$$

and

$$\mathcal{E}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) = \frac{\xi_F^*(1)}{N\mathfrak{q}} \sum_{\substack{\omega \in F^\times U_\infty \backslash \mathbb{A}_{(1)}^\times \\ \text{cond}(\omega)^2 | \mathfrak{q}}} \int_{-\infty}^{\infty} \frac{\Lambda(s, \Pi \times \pi(\omega, it)) \Lambda(w, \pi(\omega, it))}{\Lambda^*(1, \text{Ad}, \pi(\omega, it))} \\ \times \frac{\widehat{\lambda}_{\pi(\omega, it)}(\mathfrak{l}, w)}{(N\mathfrak{l})^w} h_{\mathfrak{q}}(s, w; \Pi, \pi(\omega, it)) \frac{dt}{2\pi},$$

The notation cusp^0 denotes that we are restricting to forms that are unramified at every archimedean place and the analogous role in the Eisenstein part is played by quotienting by U_∞ . Finally, we let

$$\mathcal{N}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) := \mathcal{R}(\check{H}) - \mathcal{R}(H),$$

where \mathcal{R} is given by (42) and H is as above.

In our next theorem and thereafter, θ (not to be confused with ϑ) denotes an approximation towards the Ramanujan conjecture for Π , i.e. we take θ to be any number such that for any place v , the satake parameters α_v^i for Π_v satisfy $|\alpha_{v,i}|_v \leq p_v^\theta$.

Theorem 1.2. *Let Π be an everywhere unramified cuspidal automorphic representation of $\text{GL}(3)$ over F . Suppose \mathfrak{q} and \mathfrak{l} are unramified, coprime ideals and that $\frac{1}{2} \leq \Re(s) \leq \Re(w) < \frac{3}{4}$. Then we have*

$$\mathcal{M}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) = \mathcal{N}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) + \mathcal{M}_0(\Pi, s', w', \mathfrak{l}, \mathfrak{q}),$$

where s', w' are as in (2). Moreover $h_{\mathfrak{q}}(s, w; \Pi, \pi) = 1$ when $\text{Cond}(\pi) = \mathfrak{q}$, $h_{\mathfrak{q}}(s, w; \Pi, \pi) \ll (N\mathfrak{q})^{\theta+\epsilon}$ for any $\epsilon > 0$ in general, and also,

$$\mathcal{N}_0(\Pi, s, w, \mathfrak{q}, \mathfrak{l}) \ll (N\mathfrak{q})^{\theta-1+\epsilon} + (N\mathfrak{l})^{\theta-1+\epsilon}. \quad (4)$$

As an application, We may deduce a non-vanishing result which is similar in spirit to [Kha12, Theorem 1.2], where we prove an asymptotic formula for a family of forms of prime level \mathfrak{p} and let $N\mathfrak{p}$ tend to infinity. It is maybe worth mentioning that although the results are similar, Khan's result concerns modular forms of sufficiently large weight k for $\text{GL}(2)$ over the field of rationals, while our result holds for everywhere unramified forms for $\text{GL}(2)$ over an arbitrary number field.

Corollary 1.3. *Let Π be an unramified cuspidal automorphic representation of $\text{GL}(3)$ over F and let \mathfrak{p} be an unramified prime ideal of \mathfrak{o}_F . Then for every $\epsilon > 0$, we have*

$$\frac{1}{N\mathfrak{p}} \sum_{\substack{\pi \text{ cusp}^0 \\ \text{cond}(\pi) = \mathfrak{p}}} \frac{\Lambda(\frac{1}{2}, \Pi \times \pi) \Lambda(\frac{1}{2}, \pi)}{\Lambda(1, \text{Ad}, \pi)} = \frac{4\xi_F^*(1)}{\xi_F(2)} \Lambda(1, \Pi) \Lambda(1, \bar{\Pi}) + O_\epsilon \left((N\mathfrak{p})^{\vartheta - \frac{1}{2} + \epsilon} \right).$$

In particular, for $N\mathfrak{p}$ sufficiently large, there is at least one automorphic representation π of conductor \mathfrak{p} , unramified for every archimedean place and such that $\Lambda(\frac{1}{2}, \Pi \times \pi)$ and $\Lambda(\frac{1}{2}, \pi)$ are both non-zero.

ACKNOWLEDGEMENTS

This work benefited from discussions I had with many people including Philippe Michel, Raphaël Zacharias and Han Wu. I take this opportunity to thank them.

2. NOTATIONS

Number fields and completions. In this section and throughout the paper, F will denote a fixed number field with ring of intergers \mathfrak{o}_F and discriminant d_F . For v a place of F , we let F_v be the completion of F at the place v . If v is non-Archimedean, we write \mathfrak{o}_v for the ring of integers in F_v , \mathfrak{m}_v for its maximal ideal and ϖ_v for its uniformizer. The adèle ring of F is denoted by \mathbb{A} , its unit group is denoted by \mathbb{A}^\times and finally, $\mathbb{A}_{(1)}^\times$ denotes the ideles of norm 1. We also fix once and for all, an isomorphism $\mathbb{A}^\times \simeq \mathbb{A}_{(1)}^\times \times \mathbb{R}_{>0}$.

Additive characters. We let $\psi = \otimes_v \psi_v$ be the additive character $\psi = \psi_{\mathbb{Q}} \circ \text{Tr}_{F/\mathbb{Q}}$ where $\text{Tr}_{F/\mathbb{Q}}$ is the trace map and $\psi_{\mathbb{Q}}$ is the classical additive character on $\mathbb{A}_{\mathbb{Q}}$ which is trivial on \mathbb{Q} . Let d_v be the conductor of ψ_v , *i.e.*: the smallest non-negative integer such that ψ_v is trivial on $\mathfrak{m}_v^{d_v}$. Notice that $d_v = 0$ for every finite place not dividing the discriminant and we have the relation $d_F = \prod_v p_v^{d_v}$, where p_v is the size of the residual field of F_v given by $p_v := |\mathfrak{o}_v/\mathfrak{m}_v|$.

Measures. In the group \mathbb{A} we use a product measure $dx = \prod_v dx_v$, where for each v this denotes a self dual measure with respect to ψ_v . This means that for real v , dx_v is the Lebesgue measure on \mathbb{R} , for complex v , dx_v is twice the Lebesgue measure on \mathbb{C} and for each finite v , this is a Haar measure on F_v giving measure $p_v^{-d_v/2}$ to the compact subgroup \mathfrak{o}_v . As for the multiplicative group F_v^\times we use Michel-Venkatesh's normalization and define $d^\times x_v := \zeta_{F_v}(1) \frac{dx_v}{|x_v|}$.

3. PRELIMINARIES ON AUTOMORPHIC REPRESENTATIONS

In the course of studying automorphic forms in $\text{GL}(n)$, it will be important to distinguish a few of its subgroups. For any unitary ring R with R^\times being its invertible elements, we let $Z_n(R)$ denote the group of central matrices (*i.e.* non-trivial multiples of the identity). $N_n(R)$ denote the maximal unipotent group formed by matrices with entries 1 on the diagonal and 0 below and we let A_n denote the diagonal matrices with lower-right entry 1. In other words,

$$Z_n(R) := \left\{ \begin{pmatrix} u & & & \\ & u & & \\ & & \ddots & \\ & & & u \end{pmatrix}; u \in R^* \right\},$$

$$N_n(R) := \left\{ \begin{pmatrix} 1 & x_{1,2} & & & x_{1,n} \\ & 1 & x_{2,3} & & \\ & & \ddots & \ddots & \\ & & & 1 & x_{n-1,n} \\ & & & & 1 \end{pmatrix}; x_{i,j} \in R, 1 \leq i < j \leq n \right\},$$

and

$$A_n(R) := \left\{ \begin{pmatrix} y_{n-1} \cdots y_1 & & & & \\ & y_{n-2} \cdots y_1 & & & \\ & & \ddots & & \\ & & & y_1 & \\ & & & & 1 \end{pmatrix}; y_i \in R^\times, 1 \leq i \leq n-1 \right\}.$$

We extend our additive character to N_n in the following way: If $n \in \mathbb{N}_{n-1}(\mathbb{A})$ is such that the entries right above the diagonal are $x_{1,2}, \dots, x_{n-1,n}$, then $\psi(n) := \psi(x_{1,2} + \dots + x_{n-1,n})$ and similarly for ψ_v . We can extend the measures on the local fields F_v and their unit groups F_v^\times to measures on the groups $Z_n(F_v)$, $N_n(F_v)$ and $A_n(F_v)$ using the coordinates used in their definitions. In our work, we only need to integrate on subgroups of GL_2 , we shall define these measures only in this case. Let

$$z(u) = \begin{pmatrix} u & \\ & u \end{pmatrix}, n(x) = \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix}, a(y) = \begin{pmatrix} y & \\ & 1 \end{pmatrix}.$$

As u, y run through F_v^\times and x through F_v , these functions give parametrizations of the groups $Z_2(F_v)$, $N_2(F_v)$ and $A_2(F_v)$. Thus we can give them the same measures as those of F_v^\times and F_v .

Moreover, let K_v denote a maximal compact subgroup of $\mathrm{GL}_2(F_v)$ given by

$$K_v := \begin{cases} O(2), & \text{if } F_v = \mathbb{R}, \\ U(2), & \text{if } F_v = \mathbb{C}, \\ \mathrm{GL}_2(\mathfrak{o}_v), & v < \infty. \end{cases}$$

We can now define a Haar measure on $\mathrm{GL}_2(F_v)$ by appealing to the Iwasawa decomposition. Let dk be a Haar probability measure on K_v and consider the surjective map

$$\begin{aligned} Z_2(F_v) \times N_2(F_v) \times A_2(F_v) \times K_v &\rightarrow \mathrm{GL}_2(F_v) \\ (z, n, a, k) &\mapsto z n a k. \end{aligned}$$

and let dg_v be the pull-back by this map of the measure

$$d^\times u \times dx \times \frac{d^\times y}{|y|_v} \times dk.$$

In particular, for GL_2 we have that for every measurable function on $\mathrm{GL}_2(F_v)$, we have

$$\int_{\mathrm{GL}_2(F_v)} f(g_v) dg_v = \int_{K_v} \int_{F_v^\times} \int_{F_v} \int_{F_v^\times} f \left(\begin{pmatrix} u & \\ & u \end{pmatrix} \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \begin{pmatrix} y & \\ & 1 \end{pmatrix} k \right) d^\times u dx \frac{d^\times y}{|y|_v} dk.$$

Similarly, we shall consider measures on the quotients $N_2(F_v)\backslash\mathrm{GL}_2(F_v)$ and $Z_2(F_v)\backslash\mathrm{GL}_2(F_v)$ by omitting the terms dx and $d^\times u$ respectively. Now, given a group G for which we have attached a haar measure dg_v to $G(F_v)$, we attach to $G(\mathbb{A})$ the product measure $dg = \prod_v dg_v$.

Let for a ring R , $\mathrm{PGL}_n(R) := Z_n(R)\backslash\mathrm{GL}_n(R)$. Since $\mathrm{PGL}_2(F) \hookrightarrow \mathrm{PGL}_2(\mathbb{A})$ discretely, we may transpose the measure of $\mathrm{PGL}_2(\mathbb{A})$ to a measure on

$$\mathbf{X} := \mathrm{PGL}_2(F)\backslash\mathrm{PGL}_2(\mathbb{A}) = Z_2(\mathbb{A})\mathrm{GL}_2(F)\backslash\mathrm{GL}_2(\mathbb{A}),$$

which turns out to have finite total measure $\mathrm{vol}(\mathbf{X}) < +\infty$.

3.1. Automorphic representations for $\mathrm{GL}(2)$. Consider the Hilbert space $L^2(\mathbf{X})$ with an action of $\mathrm{GL}_2(\mathbb{A})$ given by multiplication on the right and associated $\mathrm{GL}_2(\mathbb{A})$ -invariant inner product given by

$$\langle \phi_1, \phi_2 \rangle_{L^2(\mathbf{X})} = \int_{\mathbf{X}} \phi_1(g) \overline{\phi_2(g)} dg. \quad (5)$$

It is well-known that this space decomposes as

$$L^2(\mathbf{X}) = L^2_{\mathrm{cusp}}(\mathbf{X}) \oplus L^2_{\mathrm{res}}(\mathbf{X}) \oplus L^2_{\mathrm{cont}}(\mathbf{X}), \quad (6)$$

where $L^2_{\mathrm{cusp}}(\mathbf{X})$ denotes the closed subspace of cuspidal functions given by the functions satisfying the relation

$$\int_{N_2(F)\backslash N_2(\mathbb{A})} \phi(n g) dn = 0,$$

$L^2_{\mathrm{res}}(\mathbf{X})$ is the residual spectrum consisting of all the one-dimensional subrepresentations of $L^2(\mathbf{X})$ and $L^2_{\mathrm{cont}}(\mathbf{X})$ is expressed in terms of Eisenstein series which we discuss further below. Moreover, $L^2_{\mathrm{cusp}}(\mathbf{X})$ decomposes as a direct sum of irreducible $\mathrm{GL}_2(\mathbb{A})$ -representations, which are called the cuspidal automorphic representations.

3.2. Induced representations and Eisenstein series. Given a character $\omega : F^\times \backslash \mathbb{A}^\times \rightarrow \mathbb{C}^\times$, we denote by $\pi(\omega)$ the representation of $\mathrm{GL}_2(\mathbb{A})$ which is unitarily induced from the character $\omega \boxplus \omega^{-1}$ of $B_2(\mathbb{A})$, *i.e.* the space of functions f on $\mathrm{GL}_2(\mathbb{A})$ such that

$$f \left(\begin{pmatrix} a & b \\ & d \end{pmatrix} g \right) = |a/d|^{1/2} \omega(a) \omega^{-1}(d) f(g), \quad \langle f, f \rangle_{\mathrm{Ind}} < +\infty$$

where $K := \prod_v K_v$ and

$$\langle f_1, f_2 \rangle_{\mathrm{Ind}} := \int_K f_1(k) \overline{f_2(k)} dk < +\infty. \quad (7)$$

Given such f and $s \in \mathbb{C}$ we define an Eisenstein series by a process of analytic continuation. It is given by the following series, as long as it converges:

$$\mathrm{Eis}(f)(g) := \sum_{\gamma \in B_2(F)\backslash\mathrm{GL}_2(F)} f(\gamma g),$$

where for a ring R ,

$$B_2(R) := \left\{ \begin{pmatrix} a & b \\ & d \end{pmatrix}; a, d \in R^\times, b \in R \right\}.$$

It will be useful to rewrite the inner product (7) as an inner product between two Eisenstein series. In other words, for $f_1, f_2 \in \pi(\omega)$ and $\phi_i = \text{Eis}(f_i)$, $i = 1, 2$, we write

$$\langle \phi_1, \phi_2 \rangle_{\text{Eis}} := \langle f_1, f_2 \rangle_{\text{Ind}}. \quad (8)$$

Finally, for a complex parameter s , we use the notation $\pi(\omega, s) := \pi(\omega \times |\cdot|^s)$.

3.3. Spectral decomposition for a fixed level. We already encountered a decomposition of $L^2(\mathbf{X})$ in (6), but in practice we will start from a function in $L^2(\mathbf{X})$ which is right invariant by a large compact set $K \subset \text{GL}_2(\mathbb{A})$ and moreover we will need more uniformity than simply L^2 -convergence. In the following, we write down a more precise form of this decomposition for functions in $\mathcal{C}^\infty(\mathbf{X}/K)$, where K is a compact group of the form

$$K^S := \prod_{v \notin S} K_v,$$

In the next proposition, we write down a spectral decomposition for functions on \mathbf{X} that are right invariant by K^S . The only intervening representations are those that are unramified outside S . That means $\pi \in C(s)$ or $\pi = \pi(\omega, it)$ for $\omega \in \Xi(s)$. For each cuspidal (resp. Eisenstein) automorphic representation π , we let $\mathcal{B}_c(\pi)$ (resp. $\mathcal{B}_e(\pi)$) denote an orthonormal basis of the realization of π with respect to the inner product $\langle \cdot, \cdot \rangle_{L^2(\mathbf{X})}$ (resp. $\langle \cdot, \cdot \rangle_{\text{Eis}}$). We may now state the following rephrasing of [Wu14, Theorem 2.16]:

Proposition 3.1. *Let $F \in \mathcal{C}^\infty(\mathbf{X}/K_0(\mathfrak{q}))$ and of rapid decay, then*

$$\begin{aligned} F(g) = & \sum_{\pi \in C(s)} \sum_{\phi \in \mathcal{B}_c(\pi)} \langle F, \phi \rangle \phi(g) + \text{vol}(\mathbf{X})^{-1} \sum_{\chi^2=1} \langle F, \chi \circ \det \rangle \chi(\det g) \\ & + \frac{1}{4\pi} \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \sum_{\phi \in \mathcal{B}_e(\pi(\omega, it))} \langle F, \phi \rangle \phi(g) dt, \end{aligned}$$

where the convergence is absolute and uniform for g on any compact subset of \mathbf{X} .

4. PERIODS AND L FUNCTIONS

In this section, we consider irreducible automorphic representations π of $\text{GL}_n(\mathbb{A})$ and the period integrals related to some Rankin-Selberg L -functions. We shall be mainly concerned with the generic representations, which are those admitting a Whittaker model. We start with general values of n and later we specify for $n \leq 3$, which are the ones concerned by our main results. We shall not distinguish between a representation π and its space of smooth vectors V_π^∞ . An automorphic form ϕ will always denote a smooth vector in an irreducible automorphic representation.

4.1. Whittaker functions. Let π be a generic automorphic representation of $\mathrm{GL}_n(\mathbb{A})$ and let $\phi \in \pi$ be an automorphic form. Let $W_\phi : \mathrm{GL}_n(\mathbb{A}) \rightarrow \mathbb{C}$ be the Whittaker function of ϕ given by

$$W_\phi(g) = \int_{N_n(F) \backslash N_n(\mathbb{A})} \phi(n g) \psi^{-1}(n) dn.$$

It satisfies $W_\phi(n g) = \psi(n) W_\phi(g)$ for all $n \in N_n(\mathbb{A})$.

We can also define Whittaker functions for smooth generic admissible representations of GL_n over local fields. Moreover, given a generic irreducible admissible representation of $\mathrm{GL}_n(\mathbb{A})$ we might write down an isomorphism $\pi \simeq \otimes'_v \pi_v$ where for each v , π_v is a local generic admissible representation of $\mathrm{GL}_n(F_v)$ and we might define Whittaker functions for each local representation such that for every $\phi \in \pi$ such that if $\phi = \otimes'_v \phi_v$ through the above isomorphism, we have

$$W_\phi(g) = \prod_v W_{\phi_v}(g_v), \quad g = (g_v)_v \in \mathrm{GL}_n(\mathbb{A}). \quad (9)$$

In fact, the map $\phi \mapsto W_\phi$ is an intertwiner between π and

$$\mathcal{W}(\pi, \psi) := \{W_\phi; \phi \in \pi\},$$

the so-called Whittaker model of π . We similarly define the local Whittaker models $\mathcal{W}(\pi_v, \psi_v)$. Later on, it will be convenient to exchange freely between a representation and its associated Whittaker model. The importance of the latter comes from its close relation to local L -functions, as we will see in §4.2.

It is also important to consider Whittaker functions with respect to the inverse character $\psi' = \psi^{-1}$, so we analogously define W'_ϕ and W'_{ϕ_v} by replacing ψ_v by $\psi'_v = \psi_v^{-1}$ and ψ by $\psi' = \prod_v \psi'_v$ in all the previous definitions. It follows from uniqueness of local Whittaker functions that we must have the relation

$$W'_{\phi_v} = \overline{W_{\phi_v}} \text{ for all places } v \text{ of } F. \quad (10)$$

If a local generic admissible representation π_v is unramified, this means that in π_v there exists a vector which is right invariant by the action of $\mathrm{GL}_n(\mathfrak{o}_v)$. Such a vector is called *spherical* and spherical vectors are unique up to multiplication by scalars. We say that a vector $\phi_v \in \pi_v$ is *normalized spherical* if it is spherical and moreover its associated Whittaker function W_{ϕ_v} satisfies $W_{\phi_v}(e) = 1$, where $e \in \mathrm{GL}_n(F_v)$ denotes the identity.

4.2. Integral representations of $\mathrm{GL}_{n+1} \times \mathrm{GL}_n$ L -functions. This theory is very similar to (the adelic version of) Hecke's theory of L -functions for GL_2 . We start with Π and π irreducible automorphic representations of $\mathrm{GL}_{n+1}(\mathbb{A})$ and $\mathrm{GL}_n(\mathbb{A})$ respectively and let $\Phi \in \Pi$, $\phi \in \pi$ be automorphic forms. Suppose momentarily that Φ is a cusp form and hence rapidly decreasing. We can thus consider for every $s \in \mathbb{C}$ the following integral

$$I(s, \Phi, \phi) := \int_{\mathrm{GL}_n(F) \backslash \mathrm{GL}_n(\mathbb{A})} \Phi \left(\begin{matrix} h & \\ & 1 \end{matrix} \right) \phi(h) |\det h|^{s-\frac{1}{2}} dh.$$

It follows from the Whittaker decomposition of cusp forms (see [Cog07, Theorem 1.1]) that if Φ is a cuspidal function

$$I(s, \Phi, \phi) = \Psi(s, W_\Phi, W'_\phi), \quad (\operatorname{Re}(s) \gg 1), \quad (11)$$

where

$$\Psi(s, W, W') := \int_{N_n(\mathbb{A}) \backslash \operatorname{GL}_n(\mathbb{A})} W \begin{pmatrix} h & \\ & 1 \end{pmatrix} W'(h) |\det h|^{s-\frac{1}{2}} dh \quad (12)$$

Our next result gives some of the good properties of $\Psi(s, W, W')$, *i.e.* convergence, boundedness in vertical strips and the fact that it factors into local integrals whenever Φ and ϕ also factor. The first part follows from gauge estimates for Whittaker functions and the last part is [Cog07, Theorem 3.3].

Proposition 4.1. *Let $\Phi = \otimes'_v \Phi_v \in \Pi$ and $\phi = \otimes'_v \phi_v \in \pi$ be cusp forms. Let W_{Φ_v} and W'_{ϕ_v} be as in §4.1. Then, for s sufficiently large, $\Psi(s, W_\Phi, W'_\phi)$ converges and we have the factorization*

$$\Psi(s, W_\Phi, W'_\phi) = \prod_v \Psi_v(s, W_{\Phi_v}, W'_{\phi_v}),$$

where

$$\Psi_v(s, W, W') := \int_{N_n(F_v) \backslash \operatorname{GL}_n(F_v)} W \begin{pmatrix} h_v & \\ & 1 \end{pmatrix} W'(h_v) |\det h_v|^{s-\frac{1}{2}} dh_v. \quad (13)$$

Moreover, if both Π_v and π_v are unramified and Φ_v and ϕ_v are normalized spherical,

$$\Psi_v(s, W_{\Phi_v}, W'_{\phi_v}) = L(s, \Pi_v \times \pi_v).$$

Remark 4.1. *We shall use Proposition 4.1 for $n = 2$ and $n = 1$. In the case $n = 1$ we shall write $I(s, \phi)$, $\Psi(s, W_\phi)$ and $\Psi_v(s, W_{\phi_v})$ instead of $I(s, \phi, \mathbf{1})$, $\Psi(s, W_\phi, W_{\mathbf{1}_v})$ and $\Psi_v(s, W_{\phi_v}, W_{\mathbf{1}})$, where $\mathbf{1}$ and $\mathbf{1}_v$ denote the constant functions on $\operatorname{GL}_1(\mathbb{A})$ and $\operatorname{GL}_1(F_v)$ respectively.*

4.3. Newforms. For an irreducible generic representation π_v of $\operatorname{GL}_n(F_v)$ we shall need to consider its associated newform (essential vector). They were first introduced by Jacquet, Piatetski-Shapiro and Shalika in [JPSS81] and generalize the normalized spherical vectors for ramified representations. In the course of our local computations in section 7, we shall need an explicit description of the values on diagonal matrices of newforms for GL_2 and GL_3 . The next proposition follows from [Miy14, Theorem 4.1] and the relation between Satake parameters and Hecke eigenvalues:

Proposition 4.2. *Let Π_v (resp. π_v) be an irreducible generic representation for $\operatorname{GL}_3(F_v)$ (resp. $\operatorname{GL}_3(F_v)$). Let W_{Π_v} and W_{π_v} be the associated newforms. Let*

$$\lambda_{\pi_v}(\nu) := p_v^{\frac{\nu}{2}} W_{\pi_v} \begin{pmatrix} \varpi_v^\nu & \\ & 1 \end{pmatrix}, \quad A_{\Pi_v}(\nu_2, \nu_1) := p_v^{\nu_1 + \nu_2} W_{\Pi_v} \begin{pmatrix} \varpi_v^{\nu_1 + \nu_2} & & \\ & \varpi_v^{\nu_1} & \\ & & 1 \end{pmatrix}.$$

Then $\lambda_{\pi_v}(\nu)$ and $A_{\Pi_v}(\nu_1, \nu_2)$ vanish unless $\nu \geq 0$ and $\nu_1, \nu_2 \geq 0$, respectively. Furthermore if π_v is the local component of an automorphic form, $\lambda_{\pi_v}(\nu) = \lambda_\pi(\mathfrak{p}_v^\nu)$ is the \mathfrak{p}_v^ν -th Hecke eigenvalue of π .

Proposition 4.1 above tells us how the integral $\Psi(s, W, W')$ factors as a product of local integrals and ensures that for almost every place v (the unramified ones), we obtain the desired local L function. As for the remaining places, we need to calculate them for some specific choices of Φ_v and ϕ_v in section 7. For some of these calculation we may appeal to the calculations of [BKL19]:

Theorem A. ([BKL19, Theorem 1.1]) *Let v be a finite place of F and let Π_v and π_v be generic irreducible admissible representations of $\mathrm{GL}_3(F_v)$ and $\mathrm{GL}_2(F_v)$ respectively. Let $\{\alpha_i\}_{i=1}^3$ and $\{\gamma_j\}_{j=1}^2$ denote the Langlands parameters of Π_v and π_v , respectively, and let $L(s, \Pi_v \times \pi_v)$ be the naive Rankin–Selberg L -factor defined by*

$$L(s, \Pi_v \times \pi_v) = \prod_{i=3}^n \prod_{j=1}^2 (1 - \alpha_i \gamma_j p^{-s})^{-1}. \quad (14)$$

Suppose that $\mathrm{cond}(\pi_v) = n > 0$. Then, if W_{Π_v} , W'_{π_v} are the newvectors of Π_v and π_v respectively then

$$\Psi_v(s, W_{\Pi_v}^{(n)}, W'_{\pi_v}) = \frac{p}{p+1} L(s, \Pi_v \times \pi_v),$$

where n is such that $\mathrm{cond} \pi_v = \mathfrak{m}_v^n$, and

$$W_{\Pi_v}^{(n)} := \sum_{\beta \in \mathfrak{o}_v / \mathfrak{m}_v^n} \Pi \begin{pmatrix} 1 & \beta \varpi_v^{-n} & \\ & 1 & \\ & & 1 \end{pmatrix} W_{\Pi_v}. \quad (15)$$

4.4. Relation between inner products on $\mathrm{GL}(2)$. Let π be a generic automorphic representation of $\mathrm{GL}(2)$ over F . We define a $\mathrm{GL}_2(\mathbb{A})$ invariant inner product on the representation space of π as follows: If π is cuspidal, then we may see V_π as a subset of $L^2(\mathbf{X})$. So that π inherits the inner product from $L^2(\mathbf{X})$ given by (5). If π is Eisenstein we cannot see the representation of π as a subset of $L^2(\mathbf{X})$ and hence we equip it with the inner product given by (8).

There is however another way of defining an inner product in these representations which is independent of whether π is cuspidal or Eisenstein. This is done by using the Whittaker model.

For each place v , we may construct a $\mathrm{GL}_2(F_v)$ –invariant inner form on $\mathcal{W}(\pi_v, \psi_v)$ by letting

$$\vartheta_v(W_1, W_2) = \frac{\int_{F_v^\times} W_1(a(y_v)) \overline{W_2(a(y_v))} d^\times y_v}{\zeta_v(1) L_v(1, \mathrm{Ad} \pi) / \zeta_v(2)}, \quad (16)$$

The fact that the numerator of (16) is indeed right $\mathrm{GL}_2(F_v)$ –invariant follows from the theory of the Kirillov model and the inclusion of the denominator is to ensure the following property: Whenever π_v is unramified and W is normalized spherical for π_v , we have $\vartheta_v(W, W) = 1$. Finally, for $\phi_1 = \otimes \phi_{1,v}$ and $\phi_2 = \otimes \phi_{2,v}$ we define the *canonical* inner product by the formula

$$\langle \phi_1, \phi_2 \rangle_{can} := 2d_F^{1/2} \Lambda^*(1, \mathrm{Ad} \pi) \times \prod_v \vartheta_v(W_{\phi_{1,v}}, W_{\phi_{2,v}}). \quad (17)$$

Since every two $\mathrm{GL}_2(\mathbb{A})$ -invariant inner products in π must be equal up to multiplication by scalars, it follows that we can compare the canonical inner product with the ones introduced earlier for cuspidal and Eisenstein representations. Indeed, the theory of Rankin-Selberg convolutions in the cuspidal case and a direct computation in the Eisenstein case gives us the following relation:

$$\langle \phi_1, \phi_2 \rangle_{can} = \begin{cases} \langle \phi_1, \phi_2 \rangle_{L^2(\mathbf{X})}, & \text{if } \pi \text{ is cuspidal,} \\ 2\xi_F^*(1) \langle \phi_1, \phi_2 \rangle_{\mathrm{Eis}}, & \text{if } \pi \text{ is Eisenstein.} \end{cases} \quad (18)$$

5. ABSTRACT RECIPROCITY

In this section we show an identity between two periods. At this point no attempt is made in order to relate it to moments of L -functions. The proof is a rather simple matrix computation.

Suppose $\Phi \in \mathcal{C}^\infty(Z_3(\mathbb{A})\mathrm{GL}_3(F)\backslash\mathrm{GL}_3(\mathbb{A}))$ satisfies, for every $g \in \mathrm{GL}_3(\mathbb{A})$ and $A > 0$, the inequalities

$$\Phi \left(\begin{pmatrix} yu & & \\ & u & \\ & & 1 \end{pmatrix} g \right) \leq C(g) \left(|y| + \frac{1}{|y|} \right)^{-A} \left(|u| + \frac{1}{|u|} \right)^{-A}. \quad (19)$$

Then, we let

$$\mathcal{A}_s \Phi(h) := |\det h|^{s-\frac{1}{2}} \int_{F^\times \backslash \mathbb{A}^\times} \Phi \left(\begin{pmatrix} z(u)h & & \\ & & \\ & & 1 \end{pmatrix} \right) |u|^{2s-1} d^\times u. \quad (20)$$

We observe that the hypothesis (19) ensures that $y \mapsto \mathcal{A}_s \Phi \left(\begin{pmatrix} y & \\ & 1 \end{pmatrix} h \right)$ is of rapid decay as $|y| \rightarrow 0$ or $+\infty$. Before we proceed, we recall the following definition from §4.2 (see Remark 4.1):

$$I(s, \phi) := \int_{F^\times \backslash \mathbb{A}^\times} \phi(a(y)) |y|^{s-\frac{1}{2}} d^\times y,$$

whenever it converges. We are now ready to state the main result of this section:

Proposition 5.1. *Let $\Phi \in \mathcal{C}^\infty(Z_3(\mathbb{A})\mathrm{GL}_3(F)\backslash\mathrm{GL}_3(\mathbb{A}))$ satisfying (19). Then, for every $s, w \in \mathbb{C}$, we have the reciprocity relation*

$$I(w, \mathcal{A}_s \Phi) = I(w', \mathcal{A}_{s'} \check{\Phi}),$$

where $s' = \frac{1}{2}(w - s + 1)$, $w' = \frac{1}{2}(3s + w - 1)$, and $\check{\Phi}$ is given by

$$\check{\Phi} = \Pi(w_{23}) \cdot \Phi, \quad w_{23} = \begin{pmatrix} 1 & & \\ & & \\ & & 1 \end{pmatrix}. \quad (21)$$

By definition,

$$I(w, \mathcal{A}_s \Phi) = \int_{F^\times \backslash \mathbb{A}^\times} \int_{F^\times \backslash \mathbb{A}^\times} \Phi \left(\begin{pmatrix} z(u)a(y) & & \\ & & \\ & & 1 \end{pmatrix} \right) |u|^{2s-1} |y|^{s+w-1} d^\times u d^\times y. \quad (22)$$

Now, since Φ is left invariant by $Z_3(\mathbb{A})G_3(F)$, we see that for every $u, y \in \mathbb{A}^\times$, one has

$$\begin{aligned} \Phi \left(\begin{pmatrix} uy & & \\ & u & \\ & & 1 \end{pmatrix} \right) &= \Phi \left(\begin{pmatrix} u & & \\ & u & \\ & & u \end{pmatrix} w_{23} \begin{pmatrix} y & & \\ & u^{-1} & \\ & & 1 \end{pmatrix} w_{23} \right) \\ &= \Phi \left(\begin{pmatrix} y & & \\ & u^{-1} & \\ & & 1 \end{pmatrix} w_{23} \right) = \check{\Phi} \left(\begin{pmatrix} y & & \\ & u^{-1} & \\ & & 1 \end{pmatrix} \right). \end{aligned}$$

Applying this to (22) and making the change of variables $(u, y) = (u'^{-1}, u'y')$ gives the result.

6. SPECTRAL EXPANSION OF THE PERIOD

In this section we will give a spectral decomposition of the period $I(w, \mathcal{A}_s \Phi)$. We start by highlighting the kind of weight functions that we will encounter.

Let Π be an automorphic cuspidal representation for GL_3 over F and let $\Phi = \otimes_v \Phi_v \in \Pi$ be a cusp form. Let S be a finite set of places containing all archimedean places and all the places for which Φ is not normalized spherical. Notice that since Φ is a cusp form, it satisfies (19) and hence, $\mathcal{A}_s \Phi$ is of rapid decay and cuspidal. We can thus spectrally decompose it as in Proposition 3.1):

$$\begin{aligned} \mathcal{A}_s \Phi(h) &= \sum_{\pi \in C(S)} \sum_{\phi \in \mathcal{B}_c(\pi)} \langle \mathcal{A}_s \Phi, \phi \rangle \phi(h) + \mathrm{vol}(\mathbf{X})^{-1} \sum_{\omega^2=1} \langle \mathcal{A}_s \Phi, \omega \circ \det \rangle \omega(\det g) \\ &\quad + \frac{1}{4\pi} \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \sum_{\phi \in \mathcal{B}_c(\pi(\omega, it))} \langle \mathcal{A}_s \Phi, \phi \rangle \phi(h) dt. \end{aligned}$$

By integrating both sides of the above expression against an additive character and over the compact set $N_2(F) \backslash N_2(\mathbb{A})$, we get the following relation for Whittaker functions:

$$W_{\mathcal{A}_s \Phi}(h) = \sum_{\pi \in C(S)} \sum_{\phi \in \mathcal{B}_c(\pi)} \langle \mathcal{A}_s \Phi, \phi \rangle W_\phi(h) + \frac{1}{4\pi} \sum_{\pi \in \Xi(S)} \int_{-\infty}^{\infty} \sum_{\phi \in \mathcal{B}_c(\pi(\omega, it))} \langle \mathcal{A}_s \Phi, \phi \rangle W_\phi(h) dt,$$

Notice that since the one-dimensional representations are not generic, they do not contribute to the above expression. Now, because of rapid decay of the Whittaker functions W_ϕ as $|y| \rightarrow +\infty$, then if we take $\mathrm{Re}(w)$ sufficiently large, we obtain

$$\begin{aligned} \Psi(w, \mathcal{A}_s \Phi) &= \sum_{\pi \in C(S)} \sum_{\phi \in \mathcal{B}_c(\pi)} \langle \mathcal{A}_s \Phi, \phi \rangle \Psi(w, W_\phi) \\ &\quad + \frac{1}{4\pi} \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \sum_{\phi \in \mathcal{B}_e(\pi(\omega, it))} \langle \mathcal{A}_s \Phi, \phi \rangle \Psi(w, W_\phi) dt. \end{aligned} \quad (23)$$

Since $\mathcal{A}_s \Phi$ is a cuspidal function for GL_2 , It follows by (11) that

$$\int_{\mathbb{A}^\times} W_{\mathcal{A}_s \Phi}(a(y)) |y|^{w-\frac{1}{2}} = I(w, \mathcal{A}_s \Phi),$$

The next step is to realize the terms $\langle \mathcal{A}_s \Phi, \phi \rangle$ and $\Psi(w, W_\phi)$ as a product of local integrals. First, it follows from Proposition 4.1 that

$$\Psi(w, W_\phi) = \prod_v \Psi_v(w, W_{\phi_v}).$$

Moreover, from the definition of $\mathcal{A}_s \Phi$, we deduce, after changing variables that

$$\begin{aligned} \langle \mathcal{A}_s \Phi, \phi \rangle &= \int_{\mathbf{X}} |\det h|^{s-\frac{1}{2}} \left(\int_{F^\times \backslash \mathbb{A}^\times} \Phi \begin{pmatrix} hz(u) & \\ & 1 \end{pmatrix} |u|^{2s-1} du \right) \overline{\phi(h)} dh \\ &= \int_{\mathrm{GL}_2(F) \backslash \mathrm{GL}_2(\mathbb{A})} \Phi \begin{pmatrix} h & \\ & 1 \end{pmatrix} \overline{\phi(h)} |\det h|^{s-\frac{1}{2}} dh = I(s, \Phi, \overline{\phi}). \end{aligned}$$

Since Φ is a cusp form on $\mathrm{GL}(3)$, it follows from, (11), and again by Proposition 4.1, that for $\mathrm{Re}(s)$ sufficiently large,

$$I(s, \Phi, \overline{\phi}) = \Psi(s, W_\Phi, W'_\phi) = \prod_v \Psi_v(s, W_{\Phi_v}, W'_{\phi_v}),$$

where $\Psi_v(s, W, W')$ is given by (13). As a consequence, we have

We summarize the results of this section in the following proposition:

$$\begin{aligned} I(w, \mathcal{A}_s \Phi) &= \sum_{\pi \in C(S)} \sum_{\phi \in \mathcal{B}_c(\pi)} \prod_v \Psi_v \left(s, W_{\Phi_v}, W'_{\phi_v} \right) \prod_v \Psi_v(s, W_{\phi_v}) \\ &\quad + \frac{1}{4\pi} \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \sum_{\phi \in \mathcal{B}_e(\pi(\omega, it))} \prod_v \Psi_v \left(s, W_{\Phi_v}, W'_{\phi_v} \right) \prod_v \Psi_v(s, W_{\phi_v}) dt. \end{aligned} \quad (24)$$

For each generic automorphic representation π we will now construct an orthonormal basis for V_π which is formed exclusively by pure tensors: We start by choosing for each place v , an orthogonal basis $\mathcal{B}^W(\pi_v)$ of the space $\mathcal{W}(\pi_v, \psi_v)$. Consider now the elements $\phi = \otimes_v \phi_v$ such that for every finite v , W_{ϕ_v} lies in $\mathcal{B}^W(\pi_v)$ and for all but finitely many v , W_{ϕ_v} is normalized spherical. This provides us with an *orthogonal* basis for V_π . In order to get an *orthonormal* basis is to multiply these vectors by the correcting factors coming from (18). Applying this principle to (24) leads the following Proposition:

Proposition 6.1. *Let Π be a cuspidal automorphic representation and let $\Phi = \otimes_v \Phi_v \in \Pi$ be a cusp form. Then, we have*

$$2d_F^{1/2}I(w, \mathcal{A}_s\Phi) = \sum_{\pi \in C(S)} \frac{\Lambda(s, \Pi \times \pi, s)\Lambda(w, \pi)}{\Lambda(1, \text{Ad } \pi)} H(\pi) \\ + \xi_F^*(1) \sum_{\omega \in \Xi(S)} \int_{-\infty}^{\infty} \Lambda^*(1, \text{Ad } \pi(\omega, it)) H(\pi(\omega, it)) \frac{dt}{2\pi},$$

where $H(\pi) = \prod_v H_v(\pi_v)$ and H_v depends on Φ_v , s and w and is given by

$$H_v(\pi_v) := \sum_{W \in \mathcal{B}^W(\pi_v)} \frac{\Psi_v(s, W_{\Phi_v}, \overline{W})\Psi_v(w, W)}{L_v(s, \Pi_v \times \pi_v)L_v(w, \pi_v)}. \quad (25)$$

Theorem 1.1 now follows directly from Proposition 6.1 and (5.1). In the next section we will make choices for the cusp form Φ and calculate the associated weight function $H(\pi)$. These computations along with Proposition 5.1 and Proposition 6.1 would be enough to deduce a version of Theorem 1.2 for large values of $\Re(s)$ and $\Re(w)$. The final touch will be added in Section 8, where we analytically continue the moment decomposition of $I(w, \mathcal{A}_s\Phi)$ to $\frac{1}{2} \leq \Re(s), \Re(w) < 1$. Before we move on to the next section, it is important to establish some definitions about the weight functions introduced above.

6.1. Admissible weight functions. In view of Proposition 6.1 we are led to consider the class of functions H such that

$$H(\Pi) = \prod_v H_v(\pi_v)$$

and H_v is given by (25) where $\Phi = \otimes_v \Phi_v \in \Pi$ is an automorphic form and $s, w \in \mathbb{C}$. We will say that such an H is the (s, w) – *admissible weight function* with *kernel* Φ .

In a context where the pair (s, w) is clear, we shall speak simply of an *admissible weight function*. In order to avoid mentioning the kernel Φ , we will say that H_v is unramified if and only if Φ_v is spherical. We will see in section 7, if H_v is unramified, then it is a scalar multiple of the characteristic function of the unramified representations of $\text{GL}_2(F_v)$.

Finally, given $s, w \in \mathbb{C}$, if H is the (s, w) – admissible weight function with kernel Φ , we let \check{H} is the (s', w') – admissible weight function associated to $\check{\Phi}$, where s' and w' are as in (2) and $\check{\Phi}$ is as in (21). In other words,

$$\check{H}(\pi) = \check{H}_v(\pi_v),$$

where

$$\check{H}_v(\pi) := \sum_{W \in \mathcal{B}^W(\pi_v)} \frac{\Psi_v(s', W_{\check{\Phi}_v}, \overline{W})\Psi_v(w', W)}{L_v(s', \Pi_v \times \pi_v)L_v(w', \pi_v)}. \quad (26)$$

Remark 6.1. Notice that we are slightly abusing notation in the definition of \check{H} since we actually went through the triple (s, w, Φ) used to define H . So that whenever we talk about the transformation $H \mapsto \check{H}$ we should actually think of the map $(s, w, \Phi) \mapsto (s', w', \check{\Phi})$. This is not very serious and in particular, while proving Theorem 1.2, we will explicitly describe the kernel function Φ instead of the weight function H .

7. LOCAL COMPUTATIONS

Let Π be an unramified cuspidal automorphic representation of GL_3 over F . For all v , we let Φ_v^0 correspond to the normalized spherical vector in the Whittaker model. Let \mathfrak{q} and \mathfrak{l} be two coprime integral ideals of F . Let $\Phi^{\mathfrak{q}, \mathfrak{l}} = \otimes_v \Phi_v^{\mathfrak{q}, \mathfrak{l}}$ be such that for all $v \nmid \mathfrak{q}\mathfrak{l}$, we have $\Phi_v^{\mathfrak{q}, \mathfrak{l}} = \Phi_v^0$, for $v \mid \mathfrak{q}$,

$$\Phi_v^{\mathfrak{q}, \mathfrak{l}}(g) := \frac{1}{p_v^n} \sum_{\beta \in \mathfrak{m}_v^{-n}/\mathfrak{o}_v} \Phi_v^0 \left(g \begin{pmatrix} 1 & & \beta \\ & 1 & \\ & & 1 \end{pmatrix} \right), \quad (27)$$

where $n = v(\mathfrak{q})$, and, finally, for $v \mid \mathfrak{l}$,

$$\Phi_v^{\mathfrak{q}, \mathfrak{l}}(g) := \frac{1}{p_v^m} \sum_{\beta \in \mathfrak{m}_v^{-m}/\mathfrak{o}_v} \Phi_v^0 \left(g \begin{pmatrix} 1 & & \beta \\ & 1 & \\ & & 1 \end{pmatrix} \right),$$

with $m = v(\mathfrak{l})$.

In this section we compute H_v for all values of v . By construction of $\Phi^{\mathfrak{q}, \mathfrak{l}}$, we make take $S = \{v \mid \mathfrak{q}\} \cup \{v \mid \mathfrak{l}\} \cup \{v \mid \infty\}$. If $v \notin S$, we know that π_v is unramified and $\Phi_v^{\mathfrak{q}, \mathfrak{l}} = \Phi_v^0$ is spherical. We may thus restrict the sum defining $H_v(\pi_v)$ to a basis of space of vectors that are right invariant by K_v . This space is one dimensional and the last part of Proposition 4.1 gives $H_v(\pi_v) = 1$ in that case. We consider the cases where $v \mid \mathfrak{l}$, $v \mid \mathfrak{q}$ and $v \mid \infty$, in that order.

7.1. Non-archimedean case I: $v \mid \mathfrak{l}$. Recall that

$$H_v(\pi_v) = \sum_{W \in \mathcal{B}^W(\pi_v)} \frac{\Psi_v(s, W_{\Phi_v}, \overline{W}) \Psi_v(w, W)}{L_v(s, \Pi_v \times \pi_v) L_v(w, \pi_v)},$$

where

$$\Phi_v := \frac{1}{p_v^n} \sum_{\beta \in \mathfrak{m}_v^{-m}/\mathfrak{o}_v} \Phi_v^0 \left(g \begin{pmatrix} 1 & & \beta \\ & 1 & \\ & & 1 \end{pmatrix} \right), \quad (m = v(\mathfrak{l})).$$

Even though this is not at all obvious, we will show that H_v vanishes unless π_v is unramified. The first step is noticing that by right $\mathrm{GL}_2(F_v)$ invariance of the Haar measure,

$$\Psi \left(s, \Pi_v \begin{pmatrix} h & \\ & 1 \end{pmatrix} W_{\Phi_v}, \overline{W} \right) = \Psi \left(s, W_{\Phi_v}, \overline{\pi_v(h)W} \right).$$

Now we notice that given a basis $\mathcal{B}^W(\pi_v)$ of $\mathcal{W}(\pi_v, \psi_v)$, we may create a different one by considering the set

$$\{\pi_v(h) \cdot W, W \in \mathcal{B}^W(\pi_v)\},$$

for some fixed element $h \in \mathrm{GL}_2(F_v)$. Applying this trick for $h = \begin{pmatrix} 1 & \beta \\ & 1 \end{pmatrix}$ for $\beta \in \mathfrak{m}_v^{-n}/\mathfrak{o}_v$, we deduce that

$$H_v(\pi_v) = \sum_{W \in \mathcal{B}^W(\pi_v)} \frac{\Psi_v(s, W_{\Pi_v}, \overline{W}) \Psi_v(w, W^{(n)})}{L_v(s, \Pi_v \times \pi_v) L_v(w, \pi_v)},$$

where

$$W^{(n)}(h) := \sum_{\beta \in \mathfrak{m}_v^{-n}/\mathfrak{o}_v} W \left(h \begin{pmatrix} 1 & -\beta \\ & 1 \end{pmatrix} \right).$$

Since $W \mapsto \overline{\Psi_v(s, W_{\Pi_v}, \overline{W})}$ is a right K_v invariant linear functional, it vanishes unless π_v is unramified. Moreover, this functional is invariant by orthogonal projection into the space $\mathcal{W}(\pi_v, \psi_v)^{K_v}$ of right K_v -invariant vectors of $\mathcal{W}(\pi_v, \psi_v)$. As a consequence, we may restrict the sum defining H_v to a sum over a basis of $\mathcal{W}(\pi_v, \psi_v)^{K_v}$ which is a 1-dimensional space that can be spanned by the normalized spherical vector. Now, by the end of Proposition 4.1, $\Psi(s, W_{\Pi_v}, \overline{W_{\pi_v}}) = L_v(s, \Pi_v \times \pi_v)$ and by Proposition 4.2,

$$\begin{aligned} \Psi_v(w, W_{\pi_v}^{(n)}) &= \int_{F_v^\times} \delta_{v(y) \geq m} W_{\pi_v} \begin{pmatrix} y & \\ & 1 \end{pmatrix} d^\times y, \\ &= \sum_{\mu \geq m} \frac{\lambda_{\pi_v}(\mu)}{p_v^{\mu w}} = p_v^{-mw} \left(\lambda_{\pi_v}(m) - \frac{\lambda_{\pi_v}(m)}{p_v^w} \right) L_v(w, \pi_v). \end{aligned} \quad (28)$$

In particular, if $\pi \simeq \otimes'_v \pi_v$ is an automorphic representation, we obtain

$$\prod_{v|\mathfrak{q}} H_v(\pi_v) = \frac{\widehat{\lambda}_\pi(\mathfrak{l}, w)}{(N\mathfrak{l})^w}. \quad (29)$$

7.2. Non-archimedean case II: $v \mid \mathfrak{q}$. We now suppose that $v \mid \mathfrak{q}$. By our choice of Φ_v , we have

$$W_{\Phi_v}(g) = \frac{1}{p_v^n} \sum_{\beta \in \mathfrak{m}_v^{-n}/\mathfrak{o}_v} W_{\Pi_v} \left(g \begin{pmatrix} 1 & & \beta \\ & 1 & \\ & & 1 \end{pmatrix} \right), \quad (n = v(\mathfrak{q})),$$

where W_{Π_v} is the normalized spherical vector. We will show that if we put $H_v(\pi_v)$ vanishes unless $\mathrm{cond}(\pi_v) \leq n$, that $H_v(\pi_v) \ll_\epsilon p^{n(\theta-1+\epsilon)}$ and if $\mathrm{cond}(\pi_v) = n$, $H_v(\pi_v) = p_v^{-n}$.

We start by fixing an orthonormal basis for $\mathcal{W}(\pi_v, \psi_v)$. We first notice that by a result of Casselman [Cas73], if we let $W_0 = W_{\pi_v}$ be the newvector and for each $j \geq 0$, we let

$$W_j := \pi_v \begin{pmatrix} 1 & \\ & \varpi_v \end{pmatrix} W_0, \quad (30)$$

then $\{W_0, W_1, W_2, \dots\}$ is a basis for $\mathcal{W}(\pi_v, \psi_v)$. Moreover, for each $j \geq 0$, $\{W_0, W_1, \dots, W_j\}$ is a basis for the $K_v[n_0 + j]$ -invariant vectors in $\mathcal{W}(\pi_v, \psi_v)$, where $n_0 = \text{cond}(\pi_v)$. Here and thereafter, $K_v[n]$ denotes the subgroup of K_v given by the matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $c \in \mathfrak{m}_v^n$.

We now construct an *orthonormal* basis by employing the process of Gram-Schmidt. This method is the local counterpart of the one found in [BM15].

Let $\delta_{\pi_v} = \delta_{n_0=0}$. Let $\alpha_{\pi_v} := \frac{\lambda_{\pi_v}}{\sqrt{p}(1+\frac{\delta_{\pi_v}}{p})}$, and let

$$\xi_{\pi_v}(0,0) = 1, \quad \xi_{\pi_v}(1,1) = \frac{1}{\sqrt{1-\alpha_{\pi_v}^2}}, \quad \xi_{\pi_v}(1,0) = -\alpha_{\pi_v} p^{1/2} \xi_{\pi_v}(1,1),$$

and

$$\xi_{\pi_v}(j,j) = \frac{1}{\sqrt{1-\alpha_{\pi_v}^2} \sqrt{1-\frac{\delta_{\pi_v}}{p^2}}}, \quad \xi_{\pi_v}(j,j-1) = -\lambda_{\pi_v} \xi_{\pi_v}(j,j-1), \quad \xi_{\pi_v}(j,j-2) = \delta_{\pi_v} \xi_{\pi_v}(j,j).$$

Finally, we put $\xi_{\pi_v}(j,k) = 0$ for $k \leq j-2$. It from a direct computation that

$$\xi_{\pi_v}(j,k) \ll p^{j\epsilon} p^{(j-k)\vartheta}. \quad (31)$$

More importantly, let

$$\widetilde{W}_j := \frac{1}{\langle W_0, W_0 \rangle} \sum_{k=1}^j \xi_{\pi_v}(j,k) p^{\frac{k-j}{2}} W_k. \quad (32)$$

Then one may see that $\{\widetilde{W}_0, \widetilde{W}_1, \widetilde{W}_2, \dots\}$ is an orthonormal basis of $\mathcal{W}(\pi_v, \psi_v)$. We choose $\mathcal{B}^W(\pi_v) = \{\widetilde{W}_0, \widetilde{W}_1, \widetilde{W}_2, \dots\}$ and proceed to explicitly compute $H_v(\pi_v)$.

By definition we have

$$\begin{aligned} W_{\Phi_v} \begin{pmatrix} a & b \\ c & d \\ & & 1 \end{pmatrix} &= \frac{1}{p_v^n} \sum_{\beta \in \mathfrak{m}_v^{-m}/\mathfrak{o}_v} \psi(\beta c) W_{\Pi_v} \begin{pmatrix} a & b \\ c & d \\ & & 1 \end{pmatrix} \\ &= \delta_{v(c) \geq n} W_{\Pi_v} \begin{pmatrix} a & b \\ c & d \\ & & 1 \end{pmatrix}. \end{aligned}$$

Hence, if we write $h = z(u)n(x)a(y)k$, with $x \in F_v$, $u, y \in F_v^\times$ and $k = (k_{ij}) \in K_v$, then W_{Φ_v} vanishes unless $v(uk_{21}) \geq n$. Letting $d_1 := \min(n, v(u))$ and $d_2 := n - d_1$, we see that this is equivalent to k belonging to $K_v[d_2]$. This allows us to write

$$\Psi_v(s, W_{\Phi_v}, \overline{W}) = \sum_{d_1+d_2=n} \sum_{\min(\nu_1, n)=d_1} p^{-2\nu_1(s-\frac{1}{2})} \Psi_{\nu_1, d_2}(W), \quad (33)$$

where

$$\Psi_{\nu_1}(W) = \int_{F_v^\times} \int_{K_v[d_2]} W_{\Pi_v}(z(\varpi_v^{\nu_1} a(y)) \overline{W}(a(y)k)) |y|^{s-\frac{3}{2}} d^\times y dk.$$

Here we employed our choice of Haar measure on $N_2(F_v) \backslash \mathrm{GL}_2(F_v)$ and used that W_{Π_v} is spherical and W is invariant by the center. Now, if $W = \widetilde{W}_j$ is an element of our basis, given by (32), then it follows that

$$\int_{K_v[f]} \widetilde{W}_j(hk) dk = \begin{cases} \mathrm{vol}(K_v[f]) \widetilde{W}_j(h), & \text{if } j + n_0 \leq f, \\ 0, & \text{otherwise.} \end{cases} \quad (34)$$

We reason as follows: On the one hand, for every j , \widetilde{W}_j is $K_v[n_0+j]$ -invariant and is orthogonal to $\mathcal{W}(\pi_v, \psi_v)^{K_v[n_0+j-1]}$. Additionally, the operator

$$W \mapsto \frac{1}{K_v[f]} \int_{K_v[f]} \pi_v(k) W dk$$

is the orthogonal projection into the space of $K_v[f]$ -invariant vectors.

Applying (33) and (34) to the definition of $H_v(\pi_v)$ and changing order of summation, we are led to

$$\begin{aligned} H_v(\pi_v) &= \frac{1}{L_v(s, \Pi_v \times \pi_v) L_v(w, \pi_v)} \sum_{d_1+d_2=n} \sum_{j \leq d_2-d_0} \mathrm{vol}(K_v[j]) \sum_{\min(\nu_1, n)=d_1} p^{-2\nu_1(s-1/2)} \\ &\quad \times \int_{F_v^\times} W_{\Pi_v} \left(\begin{pmatrix} z(\varpi_v^{\nu_1}) a(y) & \\ & 1 \end{pmatrix} \overline{W}_j(a(y)) |y|^{s-3/2} d^\times y \Psi_v(w, \widetilde{W}_j) \right). \end{aligned} \quad (35)$$

By letting $\lambda_{\pi_v, j}(\mu) = \widetilde{W}_j(a(\varpi_v^\mu)) p^{\mu/2}$ and using Proposition 4.2, we see that

$$\int_{F_v^\times} W_{\Pi_v} \left(\begin{pmatrix} z(\varpi_v^{\nu_1}) a(y) & \\ & 1 \end{pmatrix} \overline{W}_j(a(y)) |y|^{s-3/2} d^\times y = \sum_{\nu_2 \geq 0} A_{\pi_v}(\nu_2, \nu_1) \overline{\lambda_{\pi_v, j}(\nu_2)} p^{-\nu_1} p^{-\nu_2 s}$$

and also,

$$\Psi_v(w, \overline{W}_j) = \sum_{\mu \geq 0} \lambda_{\pi_v, j}(\mu) p^{-\mu w}.$$

Inserting our last findings in (35), we deduce that

$$\begin{aligned} H_v(\pi_v) &= \frac{1}{L_v(s, \Pi_v \times \pi_v) L_v(w, \pi_v)} \sum_{d_1+d_2=n} \frac{1}{\mathfrak{u}(p^{d_2})} \sum_{j \leq d_2-d_0} \\ &\quad \sum_{\min(\nu_1, n)=d_1} \sum_{\nu_2 \geq 0} \sum_{\mu \geq 0} \frac{A_{\Pi_v}(\nu_2, \nu_1) \overline{\lambda_{\pi_v, j}(\nu_2)} \lambda_{\pi_v, j}(\mu)}{p^{(2\nu_1+\nu_2)s} p^{\mu w}}. \end{aligned}$$

We now notice that Proposition 4.2 together with (30) and (32) imply that

$$\lambda_{\pi_v, j}(\nu) = (\mathbf{u}(p^{n_0})p^{-n_0})^{1/2} \sum_{k=1}^{\min(j, k)} \xi_{\pi_v}(j, k) p^{k-\frac{j}{2}} \lambda_{\pi_v}(\nu - k).$$

As a consequence, we deduce, after changing variables that

$$\begin{aligned} H_v(\pi_v) &= \frac{\mathbf{u}(p^{n_0})p^{-n_0}}{L_v(s, \Pi_v \times \pi_v) L_v(w, \pi_v)} \sum_{d_1+d_2=n} \frac{1}{\mathbf{u}(p^{d_2})} \sum_{j \leq d_2 - n_0} p^{-j} \sum_{k_1, k_2 \leq j} \xi_{\pi_v}(j, k_1) \xi_{\pi_v}(j, k_2) \\ &\quad \times p^{k_1(1-w)} p^{k_2(1-s)} p^{-2d_1 s} \sum_{\min(\nu_1, d_2)=0} \sum_{\nu_2 \geq 0} \frac{A_{\Pi_v}(\nu_2 + k_2, \nu_1 + d_1) \overline{\lambda_{\pi_v}(\nu_2)}}{p^{(2\nu_1 + \nu_2)s}} \sum_{\mu \geq 0} \frac{\lambda_{\pi_v}(\mu)}{p^{\mu w}}. \end{aligned}$$

We recognize the last sum as $L_v(w, \pi_v)$, so that

$$\begin{aligned} H_v(\pi_v) &= \frac{\mathbf{u}(p^{n_0})p^{-n_0}}{L_v(s, \Pi_v \times \pi_v)} \sum_{d_1+d_2=n} \frac{1}{\mathbf{u}(p^{d_2})} \sum_{j \leq d_2 - n_0} p^{-j} \sum_{k_1, k_2 \leq j} \xi_{\pi_v}(j, k_1) \xi_{\pi_v}(j, k_2) \\ &\quad \times p^{k_1(1-w)} p^{k_2(1-s)} p^{-2d_1 s} \sum_{\min(\nu_1, d_2)=0} \sum_{\nu_2 \geq 0} \frac{A_{\Pi_v}(\nu_2 + k_2, \nu_1 + d_1) \overline{\lambda_{\pi_v}(\nu_2)}}{p^{(2\nu_1 + \nu_2)s}}. \quad (36) \end{aligned}$$

We are now ready to prove the following proposition

Proposition 7.1. *Let $\Phi_v = \Phi_v^{\mathfrak{q}, l}$ be as in (27) and let $n_0 = v(\mathfrak{q})$ then for every $\epsilon > 0$ there exists $\delta > 0$ such that*

- (i) $H_v(\pi_v)$ vanishes if $\text{cond } \pi_v > n$.
- (ii) $H_v(\pi_v) = p^{-n}$ if $\text{cond } \pi_v = n$.
- (iii) $H_v(\pi_v) \ll_{\epsilon} p^{n(\theta-1+\epsilon)}$ in general for $\text{Re}(s), \text{Re}(w) > \frac{1}{2} - \delta$.

The first assertion follows by observing that if $n_0 > n$ then the sum over j in (36) will vanish independently of the value of d_2 .

The second one holds because if $n_0 = n$, we automatically have $d_2 = n$ and $d_1 = j = k_1 = k_2 = 0$ and hence

$$H_v(\pi_v) = \frac{1}{L_v(s, \Pi_v \times \pi_v)} \sum_{\nu_2 \geq 0} A_{\Pi_v}(\nu_2, 0) \lambda_{\pi_v}(\nu_2),$$

by our "normalization". Finally, (iii) follows by applying the estimate in (31) to (36), which gives

$$H_v(\pi_v) \ll_{\epsilon} p^{n(-1+\epsilon)} \sum_{d_1+d_2=n} \sum_{j=0}^{d_2-n_0} \sum_{k_1, k_2=0}^j p^{(k_1+k_2-2j)(\frac{1}{2}-\vartheta)} p^{(d_1+k_2)\theta} \ll_{\epsilon} p^{n(\theta-1+\epsilon)}.$$

7.3. Meromorphic continuation with respect to the spectral parameter. let v be a non-archimedean place of F . We let for any idele character ω ,

$$G_{\omega,v}(s, w, t) := H_v(\pi(\omega, it)_v), \quad (37)$$

where $\pi(\mathbf{1}, it)_v$ is as in §3.2. It follows from the proof of Proposition 7.1 that given $\eta < \frac{1}{2}$, there exists $\delta > 0$ such that for

$$|\operatorname{Im}(t)| < \eta, \quad \operatorname{Re}(s), \Re(w) < \frac{1}{2} - \delta,$$

the sum defining $G_{\omega,v}(s, w, t)$ converges and by classical results of complex analysis, it defines a holomorphic function in the s and w variables. We will now show that we can extend it to a larger region and that it is also holomorphic in the t variable. Our starting point is the expression (36) for $H_v(\pi_v)$, from which we can see that $G_{\omega,v}(s, w, t)$ is a linear combination of terms of the shape

$$L_{\omega,k_2,d_1,d_2}(s, t) := \frac{1}{L_v(s, \Pi_v \times \pi_v)} \sum_{\min(\nu_1, d_2)=0} \sum_{\nu_2 \geq 0} \frac{A_{\Pi_v}(\nu_2 + k_2, \nu_1 + d_1) \overline{\lambda_{\pi(\omega, it)_v}(\nu_2)}}{p_v^{(2\nu_1 + \nu_2)s}},$$

with coefficients given by meromorphic functions in the variables s, w and t . The only possible polar divisors occur for t satisfying $\omega_v(\varpi_v)^2 p_v^{2it} = p_v^{\pm 1}$, due to the term $(1 - \alpha_{\pi_v}^2)^{-1}$ appearing in $\xi_{\pi_v}(j, k_1) \xi_{\pi_v}(j, k_2)$. Moreover, it follows from [BK17, Lemma 14] that for any $\epsilon > 0$, there exists $\delta > 0$ such that $L_{\omega,k_2,d_1,d_2}(s, t)$ admits a holomorphic continuation to the region

$$\operatorname{Re}(s) > \frac{1}{4} - \delta \quad \operatorname{Re}(s) \pm \operatorname{Im}(t) > -\delta, \quad (38)$$

and satisfies

$$L_{\omega,k_2,d_1,d_2}(s, w, t) \ll p_v^{(d_1+k_2)(\theta+\epsilon)}. \quad (39)$$

As a consequence, we may deduce that $G_{\omega,v}(s, w, t)$ admits meromorphic continuation to the region (38).

Now, suppose $\omega = \mathbf{1}$ is the trivial character. We shall now consider the two-variable function obtained from $F_{\mathbf{1}}$ and given by

$$D_v(s, w) := G_{\mathbf{1}}(s, w, i(1-w)). \quad (40)$$

From what we have shown for $G_{\mathbf{1},v}$ we know that for sufficiently small $\delta > 0$, in the region

$$\frac{1}{2} - \delta < \operatorname{Re}(s), \operatorname{Re}(w) \leq 1, \quad (41)$$

the only possible polar divisors are at the values of w such that $p_v^{2-2w} = p_v$. We will show that these poles cannot occur. To see this, we let

$$E_{j,k_2}(w) := \xi_{\pi_v}(j, k_2) \sum_{k_1=0}^j \xi_{\pi_v}(j, k_1) p_v^{k_1(1-w)},$$

where $\pi_v = \pi(\mathbf{1}, (1-w)/i)$. An easy computation shows that

$$E_{j,k_2}(w) = \begin{cases} 1, & \text{if } j = 0, \\ \frac{t_{j,k_2}(w)}{(1-p_v^{2w-3})}, & \text{if } j = 1, \\ 0, & \text{otherwise,} \end{cases}$$

where t_{j,k_2} is an entire function. Moreover, it follows from (31) (Replacing ϑ by $1 - \Re(w)$), (36) and (39) that

$$D_v(s, w) \ll_{s,w,\epsilon} p^{n\epsilon} \sum_{d_1+d_2=n} \sum_{j=0}^{d_2-n_0} p_v^{j(1-2\Re(w))} + p_v^{j(1-\Re(s)-\Re(w))} \ll_{\epsilon} p_v^{n(\theta+\epsilon)},$$

for s and w satisfying (41). We now summarize our findings about $G_{\omega,v}(s, w, t)$ and $D_v(s, w)$:

Proposition 7.2. *Let ω be an idele character of F and let G_{ω} be the function given by (37), where H_v is the (s, w) – local admissible weight function with kernel $\Phi_v^{\mathfrak{q},1}$, and $\Phi_v^{\mathfrak{q},1}$ is as in §7. Then $G_{\omega,v}$ admits meromorphic continuation to the region (38) with polar divisor of the form $t = t_0$ where t_0 is a solution to $\omega_v(\varpi_v)^2 p_v^{2it_0} = p_v^{\pm 1}$. Moreover, if D_v is the function given by (40), then D_v admits a holomorphic continuation to the region (41) where it satisfies*

$$D_v(s, w) \ll_{s,w,\epsilon} p_v^{n(\theta+\epsilon)}.$$

7.4. Local computations, the archimedean case. The analysis of the archimedean weight functions $\mathcal{Z}_{\Phi_v, s, w}^v(\pi)$ is somewhat of a different nature from the non-archimedean case. In here we make the simplest choice imaginable. Namely we impose that for every archimedean place v , Π_v is unramified and Φ_v corresponds to the normalized spherical vector. As a consequence it easily follows that $H_v(\pi_v)$ vanishes unless π_v is itself unramified. If we suppose π_v is unramified and if we choose a basis of $B^W(\pi_v)$ such that each term corresponds to a different K -type, then there will be at most one element W for which the period $\Psi_v(s, W_{\Phi_v}, W)$ is non-vanishing and it must be a spherical vector for π_v .

Moreover, it follows from Stade's formula [Sta01, Theorem 3.4] that under suitable normalizations,

$$\Psi_v(s, W_{\Pi_v}, \overline{W}) = L_v(s, \Pi_v \times \pi_v), \quad \Psi_v(w, W) = L_v(w, \pi_v).$$

where $W = W_{\pi_v} \in \mathcal{W}(\pi_v, \psi_v)$ is spherical and such that $\vartheta_v(W, W) = 1$. In particular, the following holds:

Proposition 7.3. *Let v be an archimedean place of F . Let Π_v be an irreducible admissible generic representation for $\mathrm{GL}_3(F_v)$, then there exists a Whittaker function $W_{\Pi_v} \in \mathcal{W}(\Pi_v, \psi_v)$ such that for every irreducible admissible generic representation for $\mathrm{GL}_2(F_v)$, we have*

$$H(\pi_v) = \begin{cases} 1, & \text{if } \pi_v \text{ is unramified,} \\ 0, & \text{otherwise.} \end{cases}$$

8. ANALYTIC CONTINUATION OF THE EISENSTEIN PART

The result of our next Proposition will be subject to Hypothesis 1, whose verification in the case where $\Phi = \Phi^{\mathfrak{q}}$ follows from the main results of Section 7.

Before we state this hypothesis, we shall add one extra piece of notation. Let $G_{\omega,v}$ be as in (37) and put

$$G_{\omega}(s, w, t) = \prod_v G_{\omega,v}(s, w, t).$$

We are now ready to state the following:

Hypothesis 1. *Suppose that there exists $\delta > 0$ such that for every place v and every idele character ω and every $s, w \in \mathbb{C}$, the function*

$$t \mapsto G_{\omega,v}(s, w, t)$$

is holomorphic for $|\operatorname{Im}(t)| < \delta$.

The next proposition treats the meromorphic continuation of term $\mathcal{E}(H)$ for values of s and w with real parts smaller than 1. The proof follows the same lines as those of [BK17, Lemma 16] and [BK18, Lemma 3].

Proposition 8.1. *Suppose that Π is a cuspidal automorphic representation and let $\Phi \in \Pi$ be an automorphic form satisfying Hypothesis 1 for some $\delta > 0$. Let $\mathcal{E}(H)$ be given by (1), defined initially for $\operatorname{Re}(s), \operatorname{Re}(w) \gg 1$. It admits a meromorphic continuation to $\operatorname{Re}(s), \operatorname{Re}(w) \geq \frac{1}{2} - \epsilon$ for some $\epsilon > 0$ with at most finitely many polar divisors. If $\frac{1}{2} \leq \operatorname{Re}(s), \operatorname{Re}(w) < 1$, its analytic continuation is given by $\mathcal{E}(H) + \mathcal{R}(H)$, where*

$$\mathcal{R}(H) = \frac{1}{N_{\mathfrak{q}}} \sum_{\pm} \operatorname{res}_{t=\pm(1-w)/i} (\pm i) \frac{\Lambda(s+it, \Pi) \Lambda(s-it, \Pi) \xi_F(w+it) \xi_F(w-it)}{\xi_F(1+2it) \xi_F(1-2it)} G_{\mathbf{1}}(s, w, t). \quad (42)$$

Proof. Let $\delta > 0$ to be chosen later. We use the non-vanishing of Dirichlet L -functions $\Lambda(s, \omega)$ at $\operatorname{Re}(s) = 1$ and continuity to define a continuous function $\sigma : \mathbb{R} \mapsto (0, \delta)$ so that neither $\Lambda(1 - 2\sigma + 2it, \omega^2)$ nor $G_{\omega}(s, w, t - i\sigma)$ have a poles for $0 \leq \sigma < \sigma(t)$.

We start by noticing that we can *analytically* continue $\mathcal{E}(H)$ to $\operatorname{Re}(s), \operatorname{Re}(w) > 1$, since in that region, one does not encounter any poles of $\Lambda(w, \pi(\mathbf{1}, it))$ for any t . Now, suppose that

$$1 < \operatorname{Re}(s) < 1 + \sigma(\operatorname{Im}(s)) \text{ and } 1 < \operatorname{Re}(w) < 1 + \sigma(\operatorname{Im}(w)).$$

We shift the contour of the integral defining $\mathcal{E}(H)$ down to $\operatorname{Im} t = -\sigma(\operatorname{Re}(t))$. We pick up a pole of $\Lambda(w - it, \omega)$ when ω is the trivial character and $w - it = 1$.

We observe that in view of our choice for σ , the resulting integral defines a holomorphic function in the region

$$\begin{cases} 1 - \sigma(\operatorname{Im}(s)) < \operatorname{Re}(s) < 1 + \sigma(\operatorname{Im}(s)), \\ 1 - \sigma(\operatorname{Im}(t)) < \operatorname{Re}(w) < 1 + \sigma(\operatorname{Im}(w)). \end{cases}$$

Take now s and w satisfying $1 - \sigma(\operatorname{Im}(s)) < \operatorname{Re}(s) < 1$ and $1 - \sigma(\operatorname{Im}(t)) < \operatorname{Re}(w) < 1$. We may shift the contour back to the real line and pick a new pole when ω is trivial and at $w + it =$

1. This proves the desired formula for $1 - \sigma(\text{Im}(s)) < \text{Re}(s) < 1$ and $1 - \sigma(\text{Im}(t)) < \text{Re}(w)$ and it follows in general by analytic continuation to all s, w such that $\frac{1}{2} - \delta < \text{Re}(s), \text{Re}(w) < 1$ by Proposition 7.2. \square

9. CONCLUSION

In this section we put together the results of the last three sections in order to deduce Theorem 1.2. We have seen in Proposition 6.1 that for sufficiently large values of $\Re(s)$ and $\Re(w)$, we have the relation

$$2d_F^{1/2} I(w, \mathcal{A}_s \Phi) = \mathcal{C}(H) + \mathcal{E}(H).$$

If we assume that H satisfies Hypothesis 1, then we may apply Proposition 8 and deduce that for $\frac{1}{2} - \delta < \text{Re}(s), \text{Re}(w) < 1$, we have

$$2d_F^{1/2} I(w, \mathcal{A}_s \Phi) = \mathcal{C}(H) + \mathcal{E}(H) + \mathcal{R}(H). \quad (43)$$

Now suppose that \check{H} also satisfies Hypothesis (1) and furthermore, suppose $\frac{1}{2} < \text{Re}(s), \text{Re}(w) \leq \frac{3}{4}$. The last assertion implies that

$$\frac{1}{2} \leq \text{Re}(s'), \text{Re}(w') < 1,$$

from what we deduce that (43) also holds with H, s and w replaced by \check{H}, s' and w' , respectively.

The main assertion of Theorem 1.2 is now a direct consequence of Proposition 5.1 and the description of the local weights $H_v(\pi_v)$ from §7.1 and §7.2.

9.1. Proof of Corollary 1.3. We use Theorem 1.2 with $s = w = 1/2$, $\mathfrak{l} = \mathfrak{o}_F$ and $\mathfrak{q} = \mathfrak{p}$, a prime ideal. We obtain that

$$\mathcal{M}(H) = \mathcal{M}(\check{H}) + \mathcal{R}(\check{H}) - \mathcal{R}(H).$$

The last part of Proposition 8 tell us that we have:

$$\mathcal{R}(\check{H}) = \frac{4\xi_F^*(1)}{\xi_F(2)} \Lambda(1, \Pi) \Lambda(1, \bar{\Pi}).$$

Moreover, in view of Proposition 7.1, we may obtain that

$$\mathcal{M}(H) = \frac{1}{N\mathfrak{p}} \sum_{\substack{\pi \text{ cusp}^0 \\ \text{cond}(\pi) = \mathfrak{p}}} \frac{\Lambda(\frac{1}{2}, \Pi \times \pi) \Lambda(\frac{1}{2}, \pi)}{\Lambda(1, \text{Ad}, \pi)} + O\left((N\mathfrak{p})^{\theta-1} \mathcal{M}^*\right),$$

and

$$\mathcal{R}(H) \ll (N\mathfrak{p})^{\theta-1}.$$

where

$$\begin{aligned} \mathcal{M}^* := & \sum_{\substack{\pi \text{ cusp}^0 \\ \text{cond}(\pi)=\mathfrak{o}_F}} \frac{|\Lambda(\frac{1}{2}, \Pi \times \pi)\Lambda(\frac{1}{2}, \pi)|}{|\Lambda(1, \text{Ad}, \pi)|} \\ & + \sum_{\substack{\omega \in F^\times \widehat{U_\infty} \backslash \mathbb{A}_{(1)}^\times \\ \text{cond}(\omega)=1}} \int_{-\infty}^{\infty} \frac{|\Lambda(\frac{1}{2}, \Pi \times \pi(\omega, it))\Lambda(\frac{1}{2}, \pi(\omega, it))|}{|\Lambda^*(1, \text{Ad}, \pi(\omega, it))|} \frac{dt}{2\pi}. \end{aligned}$$

Finally, it is easy to see that we also have the bound

$$\mathcal{M}(H) \ll (N\mathfrak{p})^{\vartheta-1/2} \mathcal{M}^*.$$

Corollary 1.3 is now a consequence of $\mathcal{M}^* \ll 1$. In other words, we just need to ensure that it converges since it is clearly independent of \mathfrak{p} .

To see that, we notice that the finite part of $\frac{\Lambda(\frac{1}{2}, \Pi \times \pi)\Lambda(\frac{1}{2}, \pi)}{\Lambda(1, \text{Ad}, \pi)}$ is bounded polynomially in terms of the eigenvalues of π_v for archimedean v and that by Stirling's formula, we have for archimedean v :

$$\frac{L(\frac{1}{2}, \Pi_v \times \pi_v)L(\frac{1}{2}, \pi_v)}{L(1, \text{Ad}, \pi_v)} \ll |t_{\pi_v}|^C e^{-2c_{F_v} \pi |t_{\pi_v}|},$$

for $\pi_v = \pi(| \cdot |^{t_{\pi_v}})$, where

$$c_{F_v} = \begin{cases} 1, & \text{if } F_v = \mathbb{R}, \\ 2, & \text{if } F_v = \mathbb{C}. \end{cases}$$

This implies that the factor $\frac{\Lambda(\frac{1}{2}, \Pi \times \pi)\Lambda(\frac{1}{2}, \pi)}{\Lambda(1, \text{Ad}, \pi)}$ decay exponentially as the t_{π_v} grow. We now deduce the convergence of the left-hand side of \mathcal{M}^* by appealing to the Weyl law for $GL(2)$ over number fields (cf. [Pal12, Theorem 3.2.1]).

REFERENCES

- [AK18] Nickolas Andersen and Eren Mehmet Kiral. Level reciprocity in the twisted second moment of rankin-selberg L-functions. *preprint: (arXiv:1801.06089)*, 2018.
- [BK17] Valentin Blomer and Rizwanur Khan. Twisted moments of L-functions and spectral reciprocity. *preprint: (arXiv:1706.01245)*, 2017.
- [BK18] Valentin Blomer and Rizwanur Khan. Uniform subconvexity and symmetry breaking reciprocity. *preprint: (arXiv:1804.01602)*, 2018.
- [BKL19] Andrew Booker, Muthu Krishnamurthy, and Min Lee. Test vectors for rankin-selberg L-functions. *preprint: (arXiv:1903.03458)*, 2019.
- [BM15] Valentin Blomer and Djordje Milićević. The second moment of twisted modular l-functions. *Geometric and Functional Analysis*, 25(2):453–516, 2015.
- [Cas73] William Casselman. On some results of atkin and lehrer. *Math. Ann.*, 201:301–314, 1973.

- [Cog07] James W. Cogdell. L -functions and converse theorems for GL_n . In *Automorphic forms and applications*, volume 12 of *IAS/Park City Math. Ser.*, pages 97–177. Amer. Math. Soc., Providence, RI, 2007.
- [JPSS81] H. Jacquet, I. I. Piatetski-Shapiro, and J. Shalika. Conducteur des représentations du groupe linéaire. *Math. Ann.*, 256(2):199–214, 1981.
- [Kha12] Rizwanur Khan. Simultaneous non-vanishing of $GL(3) \times GL(2)$ and $GL(2)$ L -functions. *Math. Proc. Cambridge Philos. Soc.*, 152(3):535–553, 2012.
- [Miy14] Michitaka Miyauchi. Whittaker functions associated to newforms for $GL(n)$ over p -adic fields. *J. Math. Soc. Japan*, 66(1):17–24, 2014.
- [Pal12] Marc Palm. Explicit $GL(2)$ trace formulas and uniform, mixed weyl laws. *Göttingen: Univ. Göttingen (Diss.)*, 2012.
- [Sta01] Eric Stade. Mellin transforms of $GL(n, \mathbb{R})$ whittaker functions. *American journal of mathematics*, 123(1):121–161, 2001.
- [Wu14] Han Wu. Burgess-like subconvex bounds for $GL_2 \times GL_1$. *Geom. Funct. Anal.*, 24(3):968–1036, 2014.
- [Zac18] Raphael Zacharias. Periods and $GL_2 \times GL_1$ reciprocity. *preprint: (arXiv:1809.10593)*, 2018.

RAMON M. NUNES // DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ (UFC).
BLOCO 914, CAMPUS DO PICI // BLOCO 914 // CEP 60455-760 FORTALEZA - CE, BRASIL
E-mail address: `ramon@mat.ufc.br`