

# PURELY INSEPARABLE RICHELLOT ISOGENIES

BRADLEY W. BROCK AND EVERETT W. HOWE

ABSTRACT. We show that if  $C$  is a supersingular genus-2 curve over an algebraically-closed field of characteristic 2, then there are infinitely many Richelot isogenies starting from  $C$ . This is in contrast to what happens with non-supersingular curves in characteristic 2, or to arbitrary curves in characteristic not 2: In these situations, there are at most fifteen Richelot isogenies starting from a given genus-2 curve.

More specifically, we show that if  $C_1$  and  $C_2$  are two arbitrary supersingular genus-2 curves over an algebraically-closed field of characteristic 2, then there are exactly sixty Richelot isogenies from  $C_1$  to  $C_2$ , unless either  $C_1$  or  $C_2$  is isomorphic to the curve  $y^2 + y = x^5$ . In that case, there are either twelve or four Richelot isogenies from  $C_1$  to  $C_2$ , depending on whether  $C_1$  is isomorphic to  $C_2$ . (Here we count Richelot isogenies up to isomorphism.) We give explicit constructions that produce all of the Richelot isogenies between two supersingular curves.

## 1. INTRODUCTION

Let  $k$  be an algebraically closed field of characteristic 2. In this paper we prove that there are Richelot isogenies connecting any two supersingular genus-2 curves over  $k$ . More specifically:

**Theorem 1.1.** *Let  $C_1$  and  $C_2$  be two supersingular genus-2 curves over  $k$ . If neither  $C_1$  nor  $C_2$  is isomorphic to the curve  $y^2 + y = x^5$ , then there are exactly sixty Richelot isogenies from  $C_1$  to  $C_2$ , up to isomorphism. If exactly one of  $C_1$  or  $C_2$  is isomorphic to the special curve, then there are twelve Richelot isogenies from  $C_1$  to  $C_2$ , up to isomorphism. If both  $C_1$  and  $C_2$  are isomorphic to the special curve, then there are four Richelot isogenies from  $C_1$  to  $C_2$ , up to isomorphism.*

In Section 5 we provide explicit constructions that give all of the Richelot isogenies connecting two supersingular curves.

To explain the significance of our result, let us give some background information about Richelot isogenies. Let  $k$  be an algebraically closed field of arbitrary characteristic and let  $J_1$  and  $J_2$  be abelian surfaces over  $k$  with principal polarizations  $\lambda_1$  and  $\lambda_2$ . (We view a polarization of an abelian variety  $A$  as an isogeny from  $A$  to its dual variety, rather than as a line bundle.) A *Richelot isogeny* from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$  is an isogeny  $\varphi : J_1 \rightarrow J_2$  that fits in a diagram

$$(1) \quad \begin{array}{ccc} J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\ \varphi \downarrow & & \uparrow \widehat{\varphi} \\ J_2 & \xrightarrow{\lambda_2} & \widehat{J}_2 \end{array}$$

where  $\widehat{\varphi}$  is the dual isogeny of  $\varphi$ . If  $C_1$  and  $C_2$  are genus-2 curves over  $k$ , a *Richelot isogeny* from  $C_1$  to  $C_2$  is defined to be a Richelot isogeny from the Jacobian of  $C_1$  to the Jacobian of  $C_2$ .

Up to automorphisms of  $(J_2, \lambda_2)$ , a Richelot isogeny is determined by its kernel in  $J_1$ . In characteristic not 2, this kernel is an order-4 subgroup of the 2-torsion of  $J_1(k)$  that is isotropic with respect to the Weil pairing on  $J_1[2]$ . Conversely, every maximal isotropic subgroup of  $J_1[2](k)$

Date: 5 February 2020.

2010 Mathematics Subject Classification. Primary 14K02; Secondary 14H40, 14H45.

Key words and phrases. Richelot isogeny, characteristic 2, supersingular, inseparable.

gives rise to an isogeny from  $(J_1, \lambda_1)$  to some principally-polarized variety  $(J_2, \lambda_2)$  (see [8, Proposition 16.8, p. 135]).

We say that two Richelot isogenies  $\varphi$  and  $\psi$  from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$  are *isomorphic* to one another if there are automorphisms  $\beta_1$  of  $(J_1, \lambda_1)$  and  $\beta_2$  of  $(J_2, \lambda_2)$  such that  $\psi = \beta_2 \circ \varphi \circ \beta_1$ . Thus we see that in characteristic not 2 there is a bijection between the set of isomorphism classes of Richelot isogenies starting from  $(J_1, \lambda_1)$  and the set of orbits of maximal isotropic subgroups of  $J_1[2](k)$  under the action of the automorphism group of  $(J_1, \lambda_1)$ . Typically this automorphism group acts trivially on the set of maximal isotropic subgroups, and in this case the number of Richelot isogenies starting from  $(J_1, \lambda_1)$  is just the number of maximal isotropic subgroups of  $J_1[2](k)$ .

An easy calculation shows that there are 15 maximal isotropic subgroups of a 4-dimensional  $\mathbb{F}_2$ -vector space with a nondegenerate alternating pairing. Thus we see that for a typical principally-polarized surface  $(J_1, \lambda_1)$  in characteristic not 2, there are 15 non-isomorphic Richelot isogenies starting from  $(J_1, \lambda_1)$ .

In characteristic 2 life is a little different, because the 2-torsion of  $J_1$  can no longer be understood simply in terms of the points of order 2 in  $J_1(k)$ . Instead, we must consider  $J_1[2]$  as a group scheme. Indeed, the bijection we mentioned two paragraphs ago is really between the set of isomorphism classes of Richelot isogenies starting from  $(J_1, \lambda_1)$  and the set of orbits of maximal isotropic subgroup schemes of  $J_1[2]$  under the action of the automorphism group  $\text{Aut}(J_1, \lambda_1)$ . Let us see what this means in characteristic 2.

Suppose that  $(J_1, \lambda_1)$  is a principally-polarized abelian surface over an algebraically-closed field  $k$  of characteristic 2. The rank of the group scheme  $J_1[2]$  is 16, and  $J_1[2]$  (like all finite commutative group schemes over perfect fields) can be decomposed into a product

$$J_1[2] = A_{\text{rr}} \times A_{\text{rl}} \times A_{\text{lr}} \times A_{\text{ll}},$$

where  $A_{\text{rr}}$  is a reduced group scheme with reduced dual,  $A_{\text{rl}}$  is a reduced group scheme with local dual,  $A_{\text{lr}}$  is a local group scheme with reduced dual, and  $A_{\text{ll}}$  is a local group scheme with local dual [12, Corollary, p. 52]. Since  $J_1[2]$  is a 2-torsion group scheme in characteristic 2, its reduced-reduced factor is trivial. And since  $J_1[2]$  is self-dual (via the Weil pairing, obtained from the principal polarization  $\lambda_1$ ), its reduced-local and local-reduced parts are dual to one another. Thus, there are three possibilities:

- $A_{\text{ll}}$  has rank 1 and  $A_{\text{rl}}$  and  $A_{\text{lr}}$  have rank 4 (the *ordinary* case);
- $A_{\text{ll}}$  has rank 4 and  $A_{\text{rl}}$  and  $A_{\text{lr}}$  have rank 2 (the *almost ordinary* case);
- $A_{\text{ll}}$  has rank 16 and  $A_{\text{rl}}$  and  $A_{\text{lr}}$  have rank 1 (the *supersingular* case).

Consider the ordinary case, where  $J_1[2]$  is isomorphic to the product of a rank-4 reduced group scheme  $A_{\text{rl}}$  with a rank-4 local group scheme  $A_{\text{lr}}$ . The kernel of a Richelot isogeny starting at  $J_1$  can be also be written as a product  $R \times L$  of a reduced group scheme  $R$  with a local group scheme  $L$ , and there are three possible shapes for these groups:

- $R = 0$  and  $L = A_{\text{lr}}$ ;
- $R = A_{\text{rl}}$  and  $L = 0$ ;
- $R(k)$  is  $\{0, T\}$  for one of the three points  $T$  of order 2 in  $J_1(k)$ , and  $L$  is the unique rank-2 sub-group scheme of  $A_{\text{lr}}$  that pairs trivially with  $R$  under the Weil pairing.

The first possibility is the Frobenius isogeny starting at  $(J_1, \lambda_1)$ , and the second is the Verschiebung. The third possibility gives rise to three Richelot isogenies, corresponding to the three points of order 2 in  $J_1(k)$ . These isogenies can be understood by lifting to characteristic 0 via the Serre–Tate canonical lift.

For the almost-ordinary case, we note that a result of Manin [7, Theorem 4.1, pp. 72–73] shows that the rank-4 group scheme  $A_{\text{ll}}$  that appears in the product decomposition of  $J_1[2]$  is isomorphic to the 2-torsion subgroup  $E[2]$  of the unique supersingular elliptic curve  $E$  over  $k$ , and so

there is a unique isotropic subgroup  $S$  of  $A_{\text{II}}$  of rank 2. Thus, the only maximal isotropic subgroups of  $A_{\text{rI}} \times A_{\text{lr}} \times A_{\text{II}}$  are  $A_{\text{rI}} \times S$  and  $A_{\text{lr}} \times S$ , so there are only two Richelot isogenies starting from  $(J_1, \lambda_1)$ . The one with kernel  $A_{\text{lr}} \times S$  is the Frobenius, and the one with kernel  $A_{\text{rI}} \times S$  is Verschiebung.

Note that in every case we have discussed so far, a given polarized abelian surface is the source for finitely many Richelot isogenies: fifteen in characteristic not 2, five for ordinary surfaces in characteristic 2, and two for almost-ordinary surfaces in characteristic 2. That is what makes Theorem 1.1 somewhat surprising: Since there are Richelot isogenies between every pair of supersingular curves, there are *infinitely many* Richelot isogenies starting from a given supersingular curve.

Our definition of a Richelot isogeny is of course a modern one. Richelot's original papers [9, 10], published in 1836 and 1837, were concerned with the evaluation of "ultra-elliptic integrals". In 1865, Königsberger [6] interpreted Richelot's work in terms of duplication formulæ for two-variable theta functions. In characteristic 0, this formulation is essentially the same as our formulation in terms of isogenies of abelian varieties. See Bost and Mestre [1] for more information.

This paper is structured as follows. In Section 2 we review a few facts about supersingular genus-2 curves in characteristic 2. In Section 3 we restate our main theorem in slightly different terms, together with three lemmas used in its proof. In Section 4 we prove the theorem, and in Section 5 we give explicit constructions that, in every case, provide all of the Richelot isogenies between two curves.

## 2. SUPERSINGULAR GENUS-2 CURVES IN CHARACTERISTIC 2

In this section we present some background material and easily-proven results about supersingular genus-2 curves in characteristic 2.

Recall that the Igusa invariants  $[J_2 : J_4 : J_6 : J_8 : J_{10}]$  of a genus-2 curve over  $k$  form an element of a weighted projective space over  $k$ , where each  $J_i$  has weight  $i$ , and where we have  $J_2 J_6 = J_4^2$  and  $J_{10} \neq 0$ .

**Lemma 2.1.** *Let  $C$  be an arbitrary genus-2 curve over an algebraically-closed field  $k$  of characteristic 2, with Igusa invariants  $[J_2 : J_4 : J_6 : J_8 : J_{10}]$ .*

1.  $C$  is supersingular if and only if  $J_2 = J_4 = J_6 = 0$ .
2. If  $C$  is supersingular and  $J_8 \neq 0$ , then
  - (a)  $C$  is isomorphic to the curve  $y^2 + y = Ax^5 + Ax^3$ , where  $A^{16} = J_8^5 / J_{10}^4$ .
  - (b)  $\text{Aut } C$  is a group of order 32.
  - (c) More specifically: For every root  $t$  of

$$T^{16} + T^8 + A^{-2}T^2 + A^{-3}T = T(T^5 + T + A^{-1})(T^{10} + T^6 + A^{-1}T^5 + A^{-2}),$$

there are two polynomials  $f$  of degree at most 2 such that  $(x, y) \mapsto (x + t, y + f(x))$  is an automorphism of  $C$ . Every automorphism of  $C$  arises in this way. The automorphisms with  $t = 0$  are the identity and the hyperelliptic involution  $\iota$ ; the automorphisms with  $t$  a root of  $T^5 + T + A^{-1}$  have order 2; the automorphisms with  $t$  a root of  $T^{10} + T^6 + A^{-1}T^5 + A^{-2}$  have order 4, and their squares are the hyperelliptic involution. Two automorphisms of order 4 are conjugate if and only if they come from the same value of  $t$ .

3. If  $C$  is supersingular and  $J_8 = 0$ , then
  - (a)  $C$  is isomorphic to the curve  $y^2 + y = x^5$ .
  - (b)  $\text{Aut } C$  is a group of order 160.
  - (c) More specifically: for every fifth root of unity  $\zeta$  and every root  $t$  of

$$T^{16} + T = T(T^5 + 1)(T^{10} + T^5 + 1),$$

there are two polynomials  $f$  of degree at most 2 such that  $(x, y) \mapsto (\zeta x + t, y + f(x))$  is an automorphism of  $C$ . Every automorphism of  $C$  arises in this way. The automorphisms with  $\zeta \neq 1$  have order divisible by 5; the automorphisms with  $\zeta = 1$  and  $t = 0$  are the identity and the hyperelliptic involution  $\iota$ ; the automorphisms with  $\zeta = 1$  and  $t^5 = 1$  have order 2; the automorphisms with  $\zeta = 1$  and with  $t$  a root of  $T^{10} + T^5 + 1$  have order 4, and their squares are the hyperelliptic involution. Two automorphisms of order 4 are conjugate if and only if they come from values of  $t$  having the same fifth power.

*Proof.* First we show that every supersingular genus-2 curve over  $k$  can be written in the form  $y^2 + y = f$  for a quintic polynomial  $f$ .

Every genus-2 curve over  $k$  has a model, nonsingular in the affine plane, of the form  $y^2 + hy = f$  for polynomials  $h, f \in k[x]$  with  $f$  of degree 5 and  $h$  nonzero of degree at most 2. (The nonsingularity is equivalent to  $h$  and  $(h')^2 f + (f')^2$  being coprime.) The desingularization of this model has a single point at infinity, which we denote  $\infty$ . If  $h$  is nonconstant, say with a root  $a$ , then the degree-0 divisor  $D = (a, \sqrt{f(a)}) - \infty$  represents a 2-torsion point on the Jacobian, because  $2D$  is the divisor of the function  $x - a$ . Therefore, for a supersingular curve  $C$ , the polynomial  $h$  must be a nonzero constant. By scaling  $y$  appropriately, we may assume that  $h = 1$ .

By explicit computation (which we leave to the reader and their computer algebra system) we see that the Igusa invariants of a curve of the form  $y^2 + y = (\text{quintic})$  satisfy  $J_2 = J_4 = J_6 = 0$ . These conditions define an irreducible 1-dimensional subvariety of the moduli space of curves, and we have just shown that it contains the subvariety of supersingular curves. But the moduli space of supersingular principally-polarized abelian surfaces is one-dimensional and closed in the moduli space of all principally-polarized abelian surfaces [5, Theorem 7(i), p. 163], so every curve with  $J_2 = J_4 = J_6 = 0$  must be supersingular.

This completes the proof of the first statement. However, the appeal to a deep theorem about the dimension of components of moduli spaces of abelian varieties with a given  $p$ -rank might be unsatisfying, especially since there are elementary arguments that will serve instead. So let us show more explicitly that every genus-2 curve with  $J_2 = J_4 = J_6 = 0$  must be supersingular. (The argument also foreshadows reasoning about genus-4 curves that appears in Section 5.4.)

Let  $P = (r, s)$  be any affine point on the hyperelliptic curve  $y^2 + y = x^5$  with  $r \neq 0$  and let  $P' = (r, s + 1)$  be the image of  $P$  under the hyperelliptic involution. Set  $b = (r^5 + 1)/r$  and consider the curve

$$C_b: y^2 + y = x^5 + bx^3.$$

We check that there is an involution  $\alpha: C_b \rightarrow C_b$  given by

$$(x, y) \mapsto (x + r^2, y + rx^2 + r^3x + s),$$

that the functions  $u = x^2 + r^2x$  and  $v = y + (x/r^2)(rx^2 + r^3x + s)$  are stable under this involution, and that these functions satisfy  $v^2 + v = u^3/r^2 + (s/r^4)u$ , so that the quotient of  $C_b$  by the involution  $\alpha$  is the supersingular elliptic curve  $E_P$  given by  $y^2 + y = x^3/r^2 + (s/r^4)x$ . Repeating this computation with  $P'$  (that is, replacing  $s$  with  $s + 1$ ), we construct another involution  $\alpha'$  of  $C_b$ , and we find the quotient of  $C_b$  by this involution is the supersingular elliptic curve  $E_{P'}$  given by  $y^2 + y = x^3/r^2 + ((s + 1)/r^4)x$ . We compute that  $\alpha$  and  $\alpha'$  commute with one another, and we

note that their product is the hyperelliptic involution  $\iota$  of  $C_b$ . It follows that we have a diagram

$$\begin{array}{ccccc}
 & & C_b & & \\
 & \langle \alpha \rangle & \downarrow & \langle \alpha' \rangle & \\
 E_P & & \mathbb{P}^1 & & E_{P'} \\
 & \searrow & \downarrow & \swarrow & \\
 & & \mathbb{P}^1 & & 
 \end{array}$$

that exhibits  $C_b$  as a Galois  $V_4$ -extension of  $\mathbb{P}^1$ . This diagram shows that the Jacobian of  $C_b$  is isogenous to the product  $E_P \times E_{P'}$ , so  $C_b$  is supersingular. The vector of Igusa invariants of  $C_b$  is  $[0:0:0:b^8:1]$ . Thus, every genus-2 curve over  $k$  with  $J_2 = J_4 = J_6 = 0$  is supersingular.

Next we note that two vectors of Igusa invariants  $[0:0:0:J_8:J_{10}]$  and  $[0:0:0:J'_8:J'_{10}]$  (with  $J_{10}$  and  $J'_{10}$  nonzero) are equal if and only if  $J_8^5/J_{10}^4 = (J'_8)^5/(J'_{10})^4$ , so supersingular curves over  $k$  are classified up to isomorphism by the invariant  $I := J_8^5/J_{10}^4$ .

Let us turn to statement (2). Given any nonzero  $A$ , we compute that the Igusa invariants of the curve  $C_A: y^2 + y = Ax^5 + Ax^3$  are  $[0:0:0:A^8:A^6]$ , so the invariant  $I(C_A)$  is equal to  $A^{16}$ , which is equal to the invariant of  $I(C)$ . Thus,  $C$  is isomorphic to  $C_A$ .

Suppose  $\alpha$  is an automorphism of the curve  $C_A$ . The hyperelliptic involution  $\iota$  of  $C_A$  is central in the automorphism group, so  $\alpha$  commutes with  $\iota$ , and it is easy to check that therefore  $\alpha$  must be of the form  $(x, y) \mapsto (g(x), y + f(x))$  for rational functions  $f(x)$  and  $g(x)$  in  $k(x)$ . The rational function  $g(x)$  must define an automorphism of  $\mathbb{P}^1$  that fixes the unique ramification point of the double cover  $C_A \rightarrow \mathbb{P}^1$  (which is  $\infty$ ) so  $g(x) = ax + t$  for some  $a, t \in k$  with  $a \neq 0$ . From the condition that

$$(y + f(x))^2 + (y + f(x)) = A(ax + t)^5 + A(ax + t)^3$$

we find that

$$f(x)^2 + f(x) = A(a^5 - 1)x^5 + Aa^4tx^4 + A(a^3 - 1)x^3 + Aa^2tx^2 + Aa(t^4 + t^2)x + (At^5 + At^3).$$

This shows that  $a^5 = a^3 = 1$  so that  $a = 1$ , and also that the rational function  $f(x)$  is a polynomial of degree at most 2. Writing  $f(x) = f_2x^2 + f_1x + f_0$ , we find that

$$f_2^2x^4 + (f_1^2 + f_2)x^2 + f_1x + (f_0^2 + f_0) = Atx^4 + Atx^2 + A(t^4 + t^2)x + (At^5 + At^3),$$

so that

$$\begin{aligned}
 f_0^2 + f_0 &= At^5 + At^3 \\
 f_1 &= At^4 + At^2 \\
 f_2 &= f_1^2 + At = A^2t^8 + A^2t^4 + At \\
 f_2^2 &= At.
 \end{aligned}$$

The first of these equations always has two solutions. The remaining three can be solved if and only if  $(A^2t^8 + A^2t^4 + At)^2 = At$ , which can be rewritten as

$$A^4t^{16} + A^4t^8 + A^2t^2 + At = 0,$$

or

$$t^{16} + t^8 + A^{-2}t^2 + A^{-3}t = 0.$$

Thus, every automorphism gives a root  $t$  of the polynomial in statement 2(c), and every such root  $t$  gives two polynomials  $f(x) = f_2x^2 + f_1x + f_0$  such that  $(x, y) \mapsto (x + t, y + f(x))$  is an automorphism. Note that this already proves statement 2(b).

When  $t = 0$  the automorphisms one obtains are the identity and the hyperelliptic involution  $\iota$ . Suppose  $\alpha: (x, y) \mapsto (x + t, y + f_2x^2 + f_1x + f_0)$  is an automorphism with  $t \neq 0$ . We compute that  $\alpha^2$  is given by  $(x, y) \mapsto (x, y + f_2t^2 + f_1t)$ , and using the formulæ for  $f_1$  and  $f_2$  we see that

$$f_2t^2 + f_1t = A^2t^{10} + A^2t^6 + At^5 = At^5(At^5 + At + 1).$$

If  $t^5 + t + A^{-1} = 0$  then  $f_2t^2 + f_1t = 0$  and  $\alpha^2$  is the identity. If  $A^2t^{10} + A^2t^6 + At^5 + 1 = 0$  then  $f_2t^2 + f_1t = 1$  and  $\alpha^2 = \iota$ , so  $\alpha$  has order 4.

It is clear that conjugate automorphisms have the same value of  $t$ . Let  $\alpha$  be any an automorphism of order 4. Consider the automorphisms  $\beta$  of order 4 that do *not* conjugate  $\alpha$  to  $\alpha^3$ ; for such a  $\beta$  we must have  $\alpha\beta = \beta\alpha$ , and we see that

$$(\alpha\beta)^2 = \alpha^2\beta^2 = \iota^2 = 1.$$

This shows that  $\beta \mapsto \alpha\beta$  is an injective map from the set of  $\beta$  of order 4 that commute with  $\alpha$  to the set of automorphisms of order at most 2. Since there are 20 automorphisms of order 4 and only 12 of order at most 2, we see that there must be a  $\beta$  of order 4 that does *not* commute with  $\alpha$ , and that therefore conjugates  $\alpha$  to  $\alpha^3$ . Thus, there is a bijection between the conjugacy classes of automorphisms of order 4 and the roots of  $T^{10} + T^6 + A^{-1}T^5 + A^{-2}$ .

The proof of statement (3) is analogous, and we leave it to the reader. The main difference is that instead of finding that  $a^5 = a^3 = 1$ , we only have that  $a^5 = 1$ , so that  $g(x)$  can be of the form  $\zeta x + t$  for a 5th root of unity  $\zeta$ . We note that all of the automorphisms of  $y^2 + y = x^5$  can be defined over  $\mathbb{F}_{16}$ .  $\square$

The invariant we used in the proof of Lemma 2.1 is useful enough to deserve a name.

**Definition 2.2.** *If  $C$  is a supersingular genus-2 curve in characteristic 2, with Igusa invariants  $[0 : 0 : 0 : J_8 : J_{10}]$ , we define the supersingular invariant of  $C$  to be the quantity  $I(C) = J_8^5 / J_{10}^4$ .*

We close with some useful statements about the finite simple sub-group schemes of the square of the unique supersingular elliptic curve  $E$  over  $k$ . These statements can be found in [3, §2].

Let  $\alpha_2$  be the unique simple local-local group scheme over  $k$ ; the group scheme  $\alpha_2$  is the kernel of Frobenius on the additive group over  $k$ . Note that  $\text{Hom}(\alpha_2, \alpha_2)(k) = k$ , and that there is a unique copy of  $\alpha_2$  in  $E$ . Fix an embedding  $e: \alpha_2 \rightarrow E$ .

Given two elements  $i, j \in k$ , not both 0, we get an embedding

$$\alpha_2 \xrightarrow{(i,j)} \alpha_2 \times \alpha_2 \xrightarrow{e \times e} E \times E$$

of  $\alpha_2$  in  $E \times E$ . The images we get from a pair  $(i, j)$  and a pair  $(i', j')$  are the same if and only if  $[i : j] = [i' : j']$  as elements of  $\mathbb{P}^1(k)$ . (And the image does not depend on the choice of  $e$ .)

The quotient of  $E \times E$  by  $[i : j](\alpha_2)$  is isomorphic to  $E \times E$  if and only if  $[i : j]$  lies in  $\mathbb{P}^1(\mathbb{F}_4)$ . Otherwise, the quotient contains a unique copy of  $\alpha_2$ . The quotient, divided by this unique copy of  $\alpha_2$ , is  $E \times E$ .

The Jacobian of a supersingular genus-2 curve over  $k$  is isomorphic to  $E \times E / [i : j](\alpha_2)$  for some  $[i : j] \in \mathbb{P}^1(k)$  that does not lie in  $\mathbb{P}^1(\mathbb{F}_4)$ .

### 3. THE THEOREM, AND A FEW LEMMAS

Let  $k$  be an algebraically closed field of characteristic 2. Using the terminology from the preceding section, we can restate our main theorem as follows:

**Theorem 3.1.** *Let  $C_1$  and  $C_2$  be two supersingular genus-2 curves over  $k$ , with supersingular invariants  $I_1$  and  $I_2$  and with Jacobians  $(J_1, \lambda_1)$  and  $(J_2, \lambda_2)$ . If  $I_1$  and  $I_2$  are both nonzero, there are exactly sixty Richelot isogenies from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$ , up to isomorphism. If exactly one of  $I_1$  and  $I_2$  is zero, there are*

exactly twelve Richelot isogenies from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$ , up to isomorphism. If both  $I_1$  and  $I_2$  are zero, there are exactly four Richelot isogenies from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$ , up to isomorphism.

Supersingular curves have Jacobians isogenous to products of supersingular elliptic curves. The following lemmas tell us a little more about the smallest isogenies  $E \times E \rightarrow J$  and  $J \rightarrow E \times E$  and about polarizations on  $E \times E$  obtained from  $J$ . To state the lemmas, we must first set some notation.

Let  $E$  be the unique supersingular elliptic curve over  $k$ . The endomorphism ring of  $E$  is a maximal order in the quaternion algebra  $H$  over  $\mathbb{Q}$  that is ramified at 2 and infinity. We write  $H = \mathbb{Q}\langle i, j \rangle$  where  $i^2 = j^2 = -1$  and  $ij = -ji$ , and we may take  $\text{End } E$  to be the order  $\mathcal{O}$  containing  $i, j$ , and  $(1 + i + j + ij)/2$ . There is a unique two-sided prime  $\mathfrak{p}$  of  $\mathcal{O}$  lying above 2; the quotient  $\mathcal{O}/\mathfrak{p}$  is isomorphic to  $\mathbb{F}_4$ .

The elliptic curve  $E$  has a unique principal polarization  $p: E \rightarrow \widehat{E}$ . We let  $P: E \times E \rightarrow \widehat{E} \times \widehat{E}$  be the product polarization  $p \times p$ .

Let  $M$  be the endomorphism of  $E \times E$  given by the matrix

$$\begin{bmatrix} 2 & 1+i \\ 1-i & 2 \end{bmatrix}$$

in  $M_2(\mathcal{O})$ , and let  $M'$  be the endomorphism

$$\begin{bmatrix} 2 & -1-i \\ -1+i & 2 \end{bmatrix}.$$

**Lemma 3.2.** *Let  $J$  be a 2-dimensional supersingular Jacobian over  $k$  and let  $\psi_0$  be a 2-isogeny from  $J$  to  $E \times E$ . Then every 2-isogeny  $\psi: J \rightarrow E \times E$  is of the form  $\alpha \circ \psi_0$  for an automorphism  $\alpha$  of  $E \times E$ .*

*Proof.* As we noted above,  $J$  has only one subgroupscheme of rank 2, so  $\psi$  and  $\psi_0$  have the same kernel. Therefore  $\psi$  factors through  $\psi_0$ .  $\square$

**Lemma 3.3.** *Let  $J$  be a 2-dimensional supersingular Jacobian over  $k$  and let  $\varphi_0$  be a 2-isogeny from  $E \times E$  to  $J$ . Then every 2-isogeny  $\varphi: E \times E \rightarrow J$  is of the form  $\varphi_0 \circ \alpha$  for an automorphism  $\alpha$  of  $E \times E$ .*

*Proof.* The 2-isogenies from  $E \times E$  to  $J$  correspond by duality to the 2-isogenies from  $\widehat{J}$  to  $\widehat{E} \times \widehat{E}$ . Since  $\widehat{J} \cong J$  and  $\widehat{E} \times \widehat{E} \cong E \times E$ , Lemma 3.3 is just a restatement of Lemma 3.2.  $\square$

**Lemma 3.4.** *Let  $J$  be a 2-dimensional supersingular Jacobian over  $k$ . There is an isogeny  $\varphi: E \times E \rightarrow J$  that pulls the canonical polarization of  $J$  back to  $M \circ P$ , and there is an isogeny  $\varphi': E \times E \rightarrow J$  that pulls the canonical polarization back to  $M' \circ P$ .*

*Proof.* Let  $\varphi$  be any 2-isogeny from  $E \times E$  to  $J$ . Then  $\varphi$  pulls back the canonical polarization of  $J$  to some degree-4 polarization of  $E \times E$  with kernel  $\alpha_2 \times \alpha_2$ . But from [3] we know that the number of isomorphism classes of such polarizations is equal to the class number  $H_2(1, 2)$  of the non-principal genus in  $H \times H$ , and this class number is 1. So we can modify  $\varphi$  by automorphisms of  $E \times E$  in order to get the pullback of the canonical polarization to be  $M \circ P$  or  $M' \circ P$ .  $\square$

#### 4. PROOF OF THE THEOREM

Let  $(J_1, \lambda_1)$  and  $(J_2, \lambda_2)$  be the canonically polarized Jacobians of  $C_1$  and  $C_2$ , respectively. From Lemma 3.4 we know that there is a degree-2 isogeny  $\varphi_2: E \times E \rightarrow J_2$  that pulls the principal polarization  $\lambda_2$  back to the degree-4 polarization  $P \circ M$ , and there is a degree-2 isogeny  $\varphi_1: E \times E \rightarrow J_1$  that pulls the principal polarization  $\lambda_1$  back to the degree-4 polarization  $P \circ M'$ .

Let  $\psi_1$  be the composite isogeny

$$J_1 \xrightarrow{\lambda_1} \widehat{J}_1 \xrightarrow{\widehat{\varphi}_1} \widehat{E} \times \widehat{E} \xrightarrow{P^{-1}} E \times E.$$

**Claim 4.1.** Every Richelot isogeny from  $C_1$  to  $C_2$  can be written as a composition  $\varphi_2 \circ \alpha \circ \psi_1$  for some automorphism  $\alpha$  of the polarized variety  $(E \times E, P \circ M)$ , and every such composition is a Richelot isogeny.

*Proof.* Let us begin by showing that the diagram

$$(2) \quad \begin{array}{ccc} J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\ \psi_1 \downarrow & & \uparrow \widehat{\psi}_1 \\ E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \end{array}$$

is commutative. Using the fact that  $\psi_1 = P^{-1} \circ \widehat{\varphi}_1 \circ \lambda_1$ , we check that

$$\begin{aligned} \widehat{\psi}_1 \circ P \circ M \circ \psi_1 &= \widehat{\lambda}_1 \circ \varphi_1 \circ \widehat{P}^{-1} \circ P \circ M \circ P^{-1} \circ \widehat{\varphi}_1 \circ \lambda_1 \\ &= \widehat{\lambda}_1 \circ \varphi_1 \circ M \circ P^{-1} \circ \widehat{\varphi}_1 \circ \lambda_1 \\ &\quad \text{[because } \widehat{P} = P, \text{ since } P \text{ is a polarization]} \\ &= \widehat{\lambda}_1 \circ \varphi_1 \circ M \circ P^{-1} \circ \widehat{\varphi}_1 \circ \lambda_1 \circ \varphi_1 \circ \varphi_1^{-1} \\ &= \widehat{\lambda}_1 \circ \varphi_1 \circ M \circ P^{-1} \circ P \circ M' \circ \varphi_1^{-1} \\ &\quad \text{[because } \varphi_1 \text{ pulls } \lambda_1 \text{ back to } P \circ M'] \\ &= \widehat{\lambda}_1 \circ \varphi_1 \circ M \circ M' \circ \varphi_1^{-1} \\ &= \widehat{\lambda}_1 \circ \varphi_1 \circ 2 \circ \varphi_1^{-1} \\ &\quad \text{[because } M \circ M' \text{ is multiplication by 2]} \\ &= 2 \circ \widehat{\lambda}_1 \\ &= 2 \circ \lambda_1 \quad \text{[because } \lambda_1 = \widehat{\lambda}_1]. \end{aligned}$$

Also note that the diagram

$$(3) \quad \begin{array}{ccc} E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\ \varphi_2 \downarrow & & \uparrow \widehat{\varphi}_2 \\ J_2 & \xrightarrow{\lambda_2} & \widehat{J}_2 \end{array}$$

is commutative, by the definition of  $\varphi_2$ .

Now suppose that  $\alpha$  is an automorphism of  $(E \times E, P \circ M)$ . This means that the diagram

$$(4) \quad \begin{array}{ccc} E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\ \alpha \downarrow & & \uparrow \widehat{\alpha} \\ E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \end{array}$$

is commutative. Stacking Diagrams (2), (4), and (3) on top of one another, we get Diagram (1). Thus, given  $\alpha$ , we get a Richelot isogeny from  $C_1$  to  $C_2$ .

Now suppose we are given a Richelot isogeny  $\varphi$  from  $C_1$  to  $C_2$ . The kernel of  $\varphi$  is a local-local subgroupscheme of  $J_1[2]$ . This kernel must contain a copy of  $\alpha_2$ , and there is a unique  $\alpha_2$  in  $J_1[2]$ . So the isogeny  $\varphi : J_1 \rightarrow J_2$  must factor through  $\psi_1$ , say  $\varphi = \gamma \circ \psi_1$  for an isogeny  $\gamma : E \times E \rightarrow J_2$ ,



and it follows that the middle arrow of the following diagram is  $P \circ M$ :

$$\begin{array}{ccc}
 J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\
 \psi_1 \downarrow & & \uparrow \widehat{\psi}_1 \\
 E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\
 \gamma \downarrow & & \uparrow \widehat{\gamma} \\
 J_2 & \xrightarrow{\lambda_2} & \widehat{J}_2
 \end{array}$$

Lemma 3.3 shows that  $\gamma$  is  $\varphi_2 \circ \alpha$  for some automorphism  $\alpha$  of  $E \times E$ , so we can further expand the diagram to get

$$\begin{array}{ccc}
 J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\
 \psi_1 \downarrow & & \uparrow \widehat{\psi}_1 \\
 E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\
 \alpha \downarrow & & \uparrow \widehat{\alpha} \\
 E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\
 \varphi_2 \downarrow & & \uparrow \widehat{\varphi}_2 \\
 J_2 & \xrightarrow{\lambda_2} & \widehat{J}_2
 \end{array}$$

The middle part of the diagram shows that  $\alpha$  is an automorphism of the principally-polarized variety  $(E \times E, P \circ M)$ . This proves the claim.  $\square$

Every automorphism  $\alpha$  of  $E \times E$  can be written as an invertible matrix  $A$  in  $M_2(\mathcal{O})$ . The automorphism  $\alpha$  respects the polarization  $P \circ M$  if and only if we have  $M = A^*MA$ , where  $A^*$  is the conjugate transpose of  $A$ . (This is because the Rosati involution on  $\text{End } E \times E$  determined by the product polarization  $P$  is equal to the conjugate transpose.) It is not hard to compute (with Magma, for example) that there are 1920 invertible matrices  $A$  in  $M_2(\mathcal{O})$  that satisfy this condition. Thus,  $\#\text{Aut}(E \times E, P \circ M) = 1920$ .

Suppose  $\beta$  is an automorphism of  $(J_1, \lambda_1)$ . Then  $\beta$  must take the unique copy of  $\alpha_2$  in  $J_1$  to itself, so  $\beta$  gives an automorphism of the polarized variety  $(E \times E, P \circ M)$ . This gives us an (injective) homomorphism

$$\text{Aut}(J_1, \lambda_1) \rightarrow \text{Aut}(E \times E, P \circ M)$$

that we denote by  $F_1$ .

Now suppose  $A$  is an automorphism of  $(E \times E, P \circ M)$ . If  $A$  takes  $\ker \varphi_2$  to itself, then  $A$  descends to give an automorphism of  $(J_2, \lambda_2)$ . Lemmas 1.4 and 1.5 of [4] show that every automorphism of  $(J_2, \lambda_2)$  comes from a unique such  $A$ . Thus, we get an (injective) homomorphism

$$\text{Aut}(J_2, \lambda_2) \rightarrow \text{Aut}(E \times E, P \circ M)$$

that we denote by  $F_2$ .

Claim 4.1 leads us to this key observation: The set of Richelot isogenies from  $C_1$  to  $C_2$  (up to isomorphism) is in bijection with the orbits of  $\text{Aut}(E \times E, P \circ M)$  under the combined actions of Image  $F_1$  on the right and Image  $F_2$  on the left. To make use of this observation we have to say a little more about  $F_1$  and  $F_2$ . First we look at  $F_2$ .

Suppose  $A$  is an automorphism of  $E \times E$ . How does  $A$  permute the various copies of  $\alpha_2$  sitting inside  $E \times E$ ? Recall that the copies of  $\alpha_2$  correspond to elements  $[i : j]$  of  $\mathbb{P}^1(k)$ . Let  $B$  in  $\text{GL}_2 \mathbb{F}_4$  be the reduction of  $A$  modulo the two-sided prime  $\mathfrak{p}$  over 2. Using Dieudonné modules, it is not hard

to show that  $A$  takes  $[i : j](\alpha_2)$  to  $[i' : j'](\alpha_2)$ , where  $[i' : j']$  is  $B[i : j]$ , under the natural action of  $\mathrm{GL}_2$  on  $\mathbb{P}^1$ . (A description of the Dieudonné module of  $E$  and of the action of  $\mathrm{End} E$  on this module is given in Section 4.2 of [11].)

By explicit calculation, we find that the image of  $\mathrm{Aut}(E \times E, P \circ M)$  in  $\mathrm{GL}_2 \mathbb{F}_4$  is  $\mathrm{SL}_2 \mathbb{F}_4$ . Let  $G$  be the kernel of the reduction map from  $\mathrm{Aut}(E \times E, P \circ M)$  to  $\mathrm{SL}_2 \mathbb{F}_4$ , so that  $\#G = 32$ . We see that the image of  $F_2$  contains  $G$ , and is equal to  $G$  if  $C_2$  is not the special curve  $y^2 + y = x^5$ .

We can also say a little more about  $F_1$ . The Frobenius is a Richelot isogeny from  $C_1$  to  $C'_1$  (where  $C'_1$  denotes the curve defined by the same equations as  $C_1$ , but with all coefficients squared), so we can always draw a diagram

$$\begin{array}{ccc}
 J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\
 \psi_1 \downarrow & & \uparrow \widehat{\psi}_1 \\
 E \times E & \xrightarrow{P \circ M} & \widehat{E} \times \widehat{E} \\
 \downarrow & & \uparrow \\
 J'_1 & \xrightarrow{\lambda'_1} & \widehat{J}'_1
 \end{array}$$

where the map  $J_1 \rightarrow J'_1$  on the left is Frobenius. It follows that the image of  $F_1$  contains  $G$ , and is equal to  $G$  unless  $C_1$  is the special curve  $y^2 + y = x^5$ .

Now we use the key observation above to complete the proof of the theorem. If neither  $C_1$  nor  $C_2$  is the special curve, then the images of  $F_1$  and  $F_2$  are both equal to the normal subgroup  $G$  of  $\mathrm{Aut}(E \times E, P \circ M)$ , so the double cosets

$$G \backslash \mathrm{Aut}(E \times E, P \circ M) / G$$

are in bijection with  $\mathrm{SL}_2 \mathbb{F}_4$ , a group with 60 elements.

If exactly one of  $C_1$  or  $C_2$  is isomorphic to the special curve, then the Richelot isogenies are in bijection with the cosets of  $\mathrm{SL}_2 \mathbb{F}_4$  by a non-normal subgroup of order 5. There are 12 orbits.

If  $C_1$  and  $C_2$  are both isomorphic to the special curve, then the Richelot isogenies are in bijection with the double cosets of  $\mathrm{SL}_2 \mathbb{F}_4$  by a non-normal subgroup of order 5 on the left, and a (possibly different) such subgroup on the right. By direct calculation, we find that no matter what the subgroups, there are 4 orbits.

This proves the theorem.  $\square$

## 5. EXPLICIT CONSTRUCTIONS FOR PURELY INSEPARABLE RICHELOT ISOGENIES


Theorem 1.1 tells us how many Richelot isogenies there are between two supersingular genus-2 curves over an algebraically-closed field  $k$  of characteristic 2. In this section we produce explicit constructions that exhibit all of these isogenies.

In Sections 5.1 and 5.2 we present two constructions of Richelot isogenies between supersingular genus-2 curves. In Section 5.3 we discuss the Frobenius and Verschiebung isogenies, and we prove that for supersingular Jacobians they are not isomorphic to one another. In Section 5.4 we show how to parametrize the Richelot isogenies discussed in Section 5.1, and in Section 5.5 we prove that the constructions we have presented account for all of the Richelot isogenies between two supersingular Jacobians.

**5.1. The dihedral construction.** Let  $D$  be a supersingular hyperelliptic curve of genus 4 over  $k$ , and suppose  $\alpha$  and  $\beta$  are involutions of  $D$  that anti-commute — that is, suppose  $\alpha\beta = \iota\beta\alpha$ , where  $\iota$  is the hyperelliptic involution on  $D$ .

Let  $G$  be the subgroup of the automorphism group  $\mathrm{Aut} D$  of  $D$  generated by the involutions  $\alpha$  and  $\beta$ . The anti-commutation of  $\alpha$  and  $\beta$  shows that  $G$  is a dihedral group of order 8, so we can draw a diagram of the intermediate curves in the Galois cover  $D \rightarrow D/G \cong \mathbb{P}^1$  as follows:


(5)



Here the curves  $C_1$  and  $C'_1$  are isomorphic to one another because  $\alpha$  and  $i\alpha$  are conjugate to one another in  $G$ ; likewise,  $C_2$  and  $C'_2$  are isomorphic to one another.

From this large  $D_8$  diagram we can extract the  $V_4$  diagram from the upper left:

(6)




This abelian diagram shows that the Jacobian of  $D$  decomposes (up to isogeny) as the sum of the Jacobians of  $C_1$  and  $C'_1$ , so that, in particular, the genus of  $C_1$  must be 2. Likewise, the genus of  $C_2$  must be 2. Also, since  $D$  is supersingular, so must be  $C_1$  and  $C_2$ .

Let  $\varphi_1$  be the natural degree-2 map from  $D$  to  $C_1 = D/\langle \alpha \rangle$ , and let  $\varphi_2$  be the natural map from  $D$  to  $C_2 = D/\langle \beta \rangle$ . Each of the curves  $C_1$ ,  $C_2$ , and  $D$  has a unique Weierstrass point, and each map  $\varphi_i$  takes the Weierstrass point of  $D$  to that of  $C_i$ . Let  $J$  be the Jacobian of  $D$ , and for each  $i$  let  $J_i$  be the Jacobian of  $C_i$ .

**Proposition 5.1.** *The map  $\varphi_{2*}\varphi_1^* : J_1 \rightarrow J_2$  is a Richelot isogeny with dual isogeny  $\varphi_{1*}\varphi_2^*$ , and the curve  $D$  and the involutions  $\alpha$  and  $\beta$  can be recovered (up to isomorphism) from this isogeny.*

*Proof.* Note that the homomorphism  $\varphi_1^*\varphi_{1*}$  of  $J$  is simply  $1 + \alpha^*$ , and likewise  $\varphi_2^*\varphi_{2*} = 1 + \beta^*$ . If we let  $J_\alpha$  be the subvariety of  $J$  where  $\alpha^*$  acts trivially, then we have a diagram



For every point  $P \in J_\alpha(k)$  we have

$$\begin{aligned} (1 + \alpha^*)(1 + \beta^*)(P) &= (1 + \alpha^* + \beta^* + \alpha^*\beta^*)(P) \\ &= (1 + \alpha^* + \beta^* - \beta^*\alpha^*)(P) \\ &= 2P + \beta^*(P) - \beta^*(P) \\ &= 2P, \end{aligned}$$

so the composite map  $J_\alpha \rightarrow J \rightarrow J_\alpha$  on the bottom of the diagram is multiplication by 2. It follows that the composite map  $J_1 \rightarrow J_2 \rightarrow J_1$  at the top of the diagram is also multiplication by 2.

Let  $\lambda$ ,  $\lambda_1$ , and  $\lambda_2$  be the canonical principal polarizations of  $J$ ,  $J_1$ , and  $J_2$ , respectively. Lemma 4.4 (p. 186) of [2] shows that the homomorphisms<sup>1</sup>  $\varphi_{1*}$  and  $\varphi_1^*$  are dual to one another, in the sense that

$$\widehat{\varphi_{1*}} = \lambda \varphi_1^* \lambda_1^{-1} \quad \text{and} \quad \widehat{\varphi_1^*} = \lambda_1 \varphi_{1*} \lambda^{-1},$$

and the same holds for  $\varphi_{2*}$  and  $\varphi_2^*$ . In the preceding paragraph we showed that  $(\varphi_{1*} \varphi_2^*)(\varphi_{2*} \varphi_1^*) = 2$  on  $J_1$ , and using the duality we just mentioned we find

$$(\lambda_1^{-1} \widehat{\varphi_1^*} \lambda \lambda^{-1} \widehat{\varphi_{2*}} \lambda_2)(\varphi_{2*} \varphi_1^*) = 2,$$

so that

$$(\lambda_1^{-1} \widehat{\varphi_{2*} \varphi_1^*} \lambda_2)(\varphi_{2*} \varphi_1^*) = 2,$$

and therefore

$$\widehat{\varphi_{2*} \varphi_1^*} \lambda_2(\varphi_{2*} \varphi_1^*) = 2\lambda_1.$$

This is exactly what is shown in Diagram (1), so  $\varphi_{2*} \varphi_1^*$  is a Richelot isogeny from  $(J_1, \lambda_1)$  to  $(J_2, \lambda_2)$ .

Similarly,  $\varphi_{1*} \varphi_2^*$  is a Richelot isogeny from  $(J_2, \lambda_2)$  to  $(J_1, \lambda_1)$ , and the two isogenies are dual to one another because their compositions in both orders are equal to multiplication by 2.

Now we turn to the final statement of the proposition.

Suppose we are given an isogeny  $\psi: J_1 \rightarrow J_2$ . Every such isogeny comes from a correspondence on  $C_1 \times C_2$ , that is, a divisor on  $C_1 \times C_2$  that does not consist solely of horizontal and vertical components. We will construct a *particular* correspondence that represents  $\psi$ .

Let  $W_1$  and  $W_2$  be the Weierstrass points on  $C_1$  and  $C_2$ , respectively, and let  $P \in C_1(k)$  be an arbitrary point with  $P \neq W_1$ . The isogeny  $\psi$  takes the class of the degree-0 divisor  $P - W_1$  on  $C_1$  to a point on the Jacobian of  $C_2$ , and every such point other than the identity can be represented as the class of a divisor  $Q + R - 2W_2$  in a unique way.<sup>2</sup> For every  $P$  such that  $[P - W_1]$  is not in the kernel of  $\psi$  we let  $\overline{\psi}(P)$  be the set  $\{Q, R\}$ , where  $Q$  and  $R$  are the points of  $C_2$  such that  $\psi(P - W_1) = [Q + R - 2W_2]$ ; for  $P$  such that  $[P - W_1]$  is in the kernel of  $\psi$  we take  $\overline{\psi}(P)$  to be the empty set. Let  $\mathcal{W}_0$  be the divisor on  $C_1 \times C_2$  consisting of the Zariski closure of the union over all  $P$  the sets  $\{(P, Q) : Q \in \overline{\psi}(P)\}$ . If there is a  $P$  such that  $\#\overline{\psi}(P) = 2$ , we take  $\mathcal{W} = \mathcal{W}_0$ ; if there is no such  $P$ , then we take  $\mathcal{W} = 2\mathcal{W}_0$ . Clearly the isogeny defined by  $\mathcal{W}$  is equal to  $\psi$ .

Note that the divisor  $\mathcal{W}$  is either an irreducible curve, or the union of two distinct irreducible curves, or twice a single irreducible curve.

What does this construction produce in our case, where  $\psi = \varphi_{2*} \varphi_1^*$ ? Let  $W$  be the Weierstrass point on  $D$ , and for a given  $P \in C_1(k)$  let  $Q'$  and  $R'$  be the two points of  $D(k)$  that map to  $P$ . The pullback (via  $\varphi_1$ ) of the divisor  $P - W_1$  is equal to  $Q' + R' - 2W$ , and the push-forward (via  $\varphi_2$ ) of this divisor is  $\varphi_2(Q') + \varphi_2(R') - 2W_2$ . Thus, the divisor  $\mathcal{W}$  on  $C_1 \times C_2$  is nothing other than the image of  $D$  under the map  $\varphi_1 \times \varphi_2$ , so  $\mathcal{W}$  is also a curve. But since the degree-2 maps  $\varphi_1: D \rightarrow C_1$  and  $\varphi_2: D \rightarrow C_2$  factor through  $\mathcal{W}$ , we see that either  $D$  is birationally equivalent to  $\mathcal{W}$  and the projection maps from  $\mathcal{W}$  to  $C_1$  and  $C_2$  have degree 2, or  $\mathcal{W}$  is birationally equivalent to both  $C_1$  and  $C_2$  and the projection maps have degree 1. The latter possibility is inconsistent with  $\varphi_{2*} \varphi_1^*$  having degree 4, so the map  $\varphi_1 \times \varphi_2: D \rightarrow C_1 \times C_2$  gives a birational equivalence between  $D$  and  $\mathcal{W}$ . Under this equivalence, the projection maps from  $\mathcal{W}$  to  $C_1$  and  $C_2$  correspond to  $\varphi_1$  and  $\varphi_2$ .

Thus,  $D$  and the double covers  $\varphi_1$  and  $\varphi_2$  can be recovered, up to isomorphism, from the Richelot isogeny  $\varphi_{2*} \varphi_1^*$ , and the involutions  $\alpha$  and  $\beta$  are determined by  $\varphi_1$  and  $\varphi_2$ . This proves the proposition.  $\square$

<sup>1</sup> The cited lemma mistakenly refers to ‘‘isogenies’’ when it should refer to ‘‘homomorphisms.’’

<sup>2</sup> The identity has an infinite number of such representations; it is equal to the class of  $Q + \iota(Q) - 2W_2$  for any point  $Q$ .

**5.2. The degenerate construction.** Suppose  $C_1$  is a supersingular genus-2 curve over  $k$ , and let  $\gamma$  be an automorphism of  $C_1$  such that  $\gamma^2$  is the hyperelliptic involution  $\iota$ . Associated to  $\gamma$  we have the pull-back automorphism  $\gamma^*$  of the Jacobian  $J_1$  of  $C_1$ .

**Proposition 5.2.** *The map  $1 + \gamma^*: J_1 \rightarrow J_1$  is a Richelot isogeny with dual isogeny  $1 - \gamma^*$ , and the automorphism  $\gamma$  can be recovered (up to conjugation in the automorphism group of  $J_1$ ) from this isogeny. This isogeny cannot be obtained from the construction of Proposition 5.1.*

*Proof.* Let  $\gamma_* \in \text{Aut } J_1$  be the push-forward of  $\gamma$ . The composition  $\gamma_*\gamma^*$  is simply multiplication by the degree of  $\gamma$ , which is 1; furthermore, since  $\iota^* = -1$  we have  $(\gamma^*)^2 = -1$ , so that  $\gamma_* = -\gamma^*$ . Thus we have a commutative diagram

$$\begin{array}{ccc} J_1 & \xrightarrow{2} & J_1 \\ 1+\gamma^* \downarrow & & \uparrow 1+\gamma_* \\ J_1 & \xrightarrow[\sim]{1} & J_1. \end{array}$$

From [2, Lemma 4.4, p. 186] we know that the  $\gamma_* = \lambda_1^{-1} \widehat{\gamma^*} \lambda_1$ , where  $\lambda_1$  is the canonical principal polarization of  $J_1$  and where  $\widehat{\gamma^*}$  is the dual isogeny of  $\gamma^*$ . This means that we can extend the preceding diagram to get

$$\begin{array}{ccccc} J_1 & \xrightarrow{2} & J_1 & \xrightarrow{\lambda_1} & \widehat{J}_1 \\ 1+\gamma^* \downarrow & & \uparrow 1+\gamma_* & & \uparrow 1+\widehat{\gamma^*} \\ J_1 & \xrightarrow[\sim]{1} & J_1 & \xrightarrow{\lambda_1} & \widehat{J}_1. \end{array}$$

The outer portion of this diagram is precisely Diagram (1), so we find that  $1 + \gamma^*$  is a Richelot isogeny from  $(J_1, \lambda_1)$  to itself. The dual isogeny is clearly  $1 - \gamma^*$ . Also, from the isogeny  $1 + \gamma^*$  we can recover the automorphism  $\gamma^*$  (up to conjugation), and Torelli's theorem tells us that this specifies  $\gamma$  (up to conjugation).

To show that this isogeny cannot be produced from the construction of the preceding section, we repeat the construction of the correspondence  $\mathcal{W} \subset C_1 \times C_1$  given in the proof of Proposition 5.1. If  $W_1$  is the Weierstrass point of  $C_1$  and  $P$  is an arbitrary point of  $C_1$  such that  $[P - W_1]$  is not in the kernel of  $1 + \gamma^*$ , we find that

$$(1 + \gamma^*)([P - W_1]) = [P + \gamma^{-1}(P) - 2W_1],$$


so the divisor  $\mathcal{W}$  is the union of the graph of the identity of  $C_1$  and the graph of the automorphism  $\gamma^{-1}$  of  $C_1$ . In particular,  $\mathcal{W}$  is the union of two irreducible curves (each isomorphic to  $C_1$ ). For the Richelot isogenies in Proposition 5.1, the divisor  $\mathcal{W}$  was an irreducible curve. Thus, no isogeny can be produced by both of these constructions.  $\square$

As we have just proven, these degenerate isogenies do not come from the dihedral construction described in the preceding section. However, if we generalize the dihedral construction to allow genus-4 "curves"  $D$  that are *not* irreducible, then the degenerate isogenies fit into the same framework. Let us sketch here how this works.

Let  $D$  be the disjoint union of two copies of  $C_1$ , which we denote by writing  $D = C_1 \amalg C_1$ . An automorphism of  $D$  can either swap the copies of  $C_1$  or not. Given two automorphisms  $f$  and  $g$  of  $C_1$ , we will denote by  $\begin{bmatrix} f & 0 \\ 0 & g \end{bmatrix}$  the automorphism that sends the first component to itself by  $f$  and the second component to itself by  $g$ ; we will denote by  $\begin{bmatrix} 0 & g \\ f & 0 \end{bmatrix}$  the automorphism that sends the first component to the second via  $f$  and the second component to the first via  $g$ .

Suppose  $\gamma$  is an automorphism of  $C_1$  with  $\gamma^2 = \iota$ . Let  $\alpha = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ , let  $\beta = \begin{bmatrix} 0 & \gamma \\ \iota\gamma & 0 \end{bmatrix}$ , and let  $I = \begin{bmatrix} \iota & 0 \\ 0 & \iota \end{bmatrix}$ . Note that  $I^*$  acts as  $-1$  on the Jacobian  $J = J_1 \oplus J_1$  of  $D$ , so  $I$  plays the role of the hyperelliptic involution; it is central in the automorphism group of  $D$ .

We see that  $\alpha$  and  $\beta$  are involutions that anti-commute with one another, in the sense that  $\alpha\beta = I\beta\alpha$ , and the subgroup  $G$  of  $\text{Aut } D$  generated by them is dihedral of order 8. As in Section 5.1, we get a Galois cover  $D \rightarrow D/G \cong \mathbb{P}^1$ :



If we let  $\text{Id}$  denote the identity map on  $C_1$ , then the four maps from  $C_1 \amalg C_1$  to  $C_1$  in this diagram are, from left to right, isomorphic to  $(\iota, \iota)$ ,  $(\text{Id}, \text{Id})$ ,  $(\iota\gamma, \text{Id})$ , and  $(\gamma, \iota)$ . In the notation of Section 5.1, we have  $\varphi_1 = (\text{Id}, \text{Id})$  and  $\varphi_2 = (\iota\gamma, \text{Id})$ .

Now we can compute where the homomorphism  $\varphi_{2*}\varphi_1^*: J_1 \rightarrow J_1$  sends the class of a divisor  $P - W_1$ . We see that  $\varphi_1^*(P)$  consists of two points:  $P$  on one copy of  $C_1$  in  $C_1 \amalg C_1$ , and  $P$  on the other copy of  $C_1$ . Applying  $\varphi_2$  to this divisor gives  $P + \iota\gamma(P)$ . Similarly, we find that  $W_1$  gets sent to  $2W_1$ , so

$$\varphi_{2*}\varphi_1^*([P - W_1]) = [P + \iota\gamma(P) - 2W_1].$$

But  $\gamma^*(P) = \iota\gamma(P)$ , so we find that  $1 + \gamma^* = \varphi_{2*}\varphi_1^*$ , and the Richelot isogeny we produced with the degenerate construction can be viewed as coming from a generalized version of the dihedral construction.

**5.3. Frobenius and Verschiebung.** To be complete, we will make a few remarks on the Richelot isogenies given by Frobenius and Verschiebung. Let  $C_1$  be an arbitrary supersingular genus-2 curve over  $k$ , say given by an equation  $y^2 = x^5 + bx^3$ , and let  $C_2$  be the curve  $y^2 = x^5 + b^2x^3$ . Let  $(J_1, \lambda_1)$  and  $(J_2, \lambda_2)$  be the polarized Jacobians of these curves. The Frobenius morphism  $\varphi$  from  $C_1$  to  $C_2$  takes a point  $(x, y)$  to the point  $(x^2, y^2)$ , and the push-forward of this map is an isogeny  $F: J_1 \rightarrow J_2$  also called the Frobenius. The isogeny  $V: J_2 \rightarrow J_1$  given by  $V = \lambda_1^{-1}\widehat{F}\lambda_2$  is called the Verschiebung. We have a diagram

$$\begin{array}{ccc} J_1 & \xrightarrow{2\lambda_1} & \widehat{J}_1 \\ F \downarrow & & \uparrow \widehat{F} \\ J_2 & \xrightarrow{\lambda_2} & \widehat{J}_2 \end{array}$$

that shows that the Frobenius is a Richelot isogeny and that its dual isogeny is the Verschiebung.

If we apply the construction of the divisor  $\mathcal{W}$  from the proof of Proposition 5.1 to the isogeny  $\psi = F$ , we find that  $\overline{\psi}(P) = \{\varphi(P), W_2\}$  for every  $P \neq W_1$  in  $C_1(k)$ , so that  $\mathcal{W} \subset C_1 \times C_2$  is the union of the graph of  $\varphi$  and a “horizontal” copy of  $C_1$ .

Similarly, if we construct  $\mathcal{W} \subset C_2 \times C_1$  from the Verschiebung  $V: J_2 \rightarrow J_1$ , we obtain the transpose of the graph of  $\varphi$  together with a “horizontal” copy of  $C_2$ .

We see that neither Frobenius nor Verschiebung can be constructed from the dihedral construction or from the degenerate construction from the preceding sections.

There is one remaining point to consider: Are the Frobenius and Verschiebung isogenies starting from a given Jacobian ever isomorphic to one another? The following theorem says that the answer is no.

**Theorem 5.3.** *Let  $C$  be a supersingular genus-2 curve over  $k$  with supersingular invariant  $I$ , let  $C^{(2)}$  and  $C^{(1/2)}$  be the curves with invariants  $I^2$  and  $I^{1/2}$ , respectively, and let  $J$ ,  $J^{(2)}$ , and  $J^{(1/2)}$  be the Jacobians of these curves. Then Frobenius isogeny  $F: J \rightarrow J^{(2)}$  is never isomorphic to the Verschiebung isogeny  $v: J \rightarrow J^{(1/2)}$ .*

*Proof.* Suppose, to get a contradiction, that  $F$  and  $V$  are isomorphic. This means that there are isomorphisms  $a: J \rightarrow J$  and  $b: J^{(2)} \rightarrow J^{(1/2)}$  of polarized Jacobians such that  $V = b \circ F \circ a$ . Let  $F'$  be the Frobenius morphism from  $J^{(1/2)}$  to  $J$ . By definition,  $F'V$  is multiplication by 2 on  $J$ .

Torelli's theorem tells us that there are isomorphisms  $\alpha: C \rightarrow C$  and  $\beta: C^{(2)} \rightarrow C^{(1/2)}$  such that  $a = \alpha_*$  and  $b = \beta_*$ . Also, if we let  $\varphi: C \rightarrow C^{(2)}$  and  $\varphi': C^{(1/2)} \rightarrow C$  be the Frobenius morphisms of curves, then  $F = \varphi_*$  and  $F' = \varphi'_*$ .

Let  $\infty$  be the infinite point on  $C$  and let  $P$  be any point on  $C$  other than  $\infty$ . Then in the Jacobian of  $C$  we have

$$\begin{aligned} [2P - 2\infty] &= 2[P - \infty] \\ &= (F' \circ V)([P - \infty]) \\ &= (F' \circ b \circ F \circ a)([P - \infty]) \\ &= (\varphi'_* \circ \beta_* \circ \varphi_* \circ \alpha_*)([P - \infty]) \\ &= [(\varphi' \circ \beta \circ \varphi \circ \alpha)(P) - \infty] \\ &= [Q - \infty] \\ &= [Q + \infty - 2\infty] \end{aligned}$$

where  $Q = (\varphi' \circ \beta \circ \varphi \circ \alpha)(P)$ . But since  $P \neq \infty$ , this equality contradicts the fact (noted above) that every nonzero point on  $J$  has a unique representation as the class of a divisor of the form  $[R + S - 2\infty]$ .  $\square$

**5.4. Parametrizing dihedral diagrams.** In this section we investigate how to create a dihedral diagram like Diagram (5) starting from two given double covers of  $\mathbb{P}^1$  by supersingular genus-2 curves; that is, given  $C_1 \rightarrow \mathbb{P}^1$  and  $C_2 \rightarrow \mathbb{P}^1$ , we want to determine what other information is necessary and sufficient to specify a diagram like Diagram (5) up to isomorphism.

Since we are working up to isomorphism, we may demand that the double covers  $C_1 \rightarrow \mathbb{P}^1$  and  $C_2 \rightarrow \mathbb{P}^1$  be in the standard forms specified in Lemma 2.1. For now, let us assume that the supersingular invariants of the two curves are nonzero, so that we can write the double cover  $C_1 \rightarrow \mathbb{P}^1$  as  $y^2 + y = A^2(x^5 + x^3)$  and the double cover  $C_2 \rightarrow \mathbb{P}^1$  as  $v^2 + v = B^2(u^5 + u^3)$  for some nonzero  $A$  and  $B \in k$ . (We use  $A^2$  and  $B^2$  instead of simply  $A$  and  $B$  in order to avoid a proliferation of square roots in various formulæ to come.)

**Theorem 5.4.** *Up to isomorphism, diagrams in the form of Diagram (5) with  $C_1$  and  $C_2$  given by  $y^2 + y = A^2(x^5 + x^3)$  and  $v^2 + v = B^2(u^5 + u^3)$ , respectively, are in bijection with pairs  $(a, c)$  of nonzero elements of  $k$  satisfying*


$$(7) \quad 0 = B^4 a^8 + B^2 a^6 c + a^4 + A^2 a c^6 + A^4 c^8 + c^4$$

$$(8) \quad 0 = A^2 B^6 a^{10} + B^2 a^5 + A^2 B^2 a^4 c^2 + A^2 B^2 a^2 c^4 + A^6 B^2 c^{10} + A^2 c^5$$

$$(9) \quad 0 \neq B^2 a^5 + A^2 c^5.$$

*Proof.* Suppose we have a diagram like Diagram (5) that contains our given maps  $C_1 \rightarrow \mathbb{P}^1$  and  $C_2 \rightarrow \mathbb{P}^1$ . We begin by focusing on a subdiagram, looking especially at the relationships among the copies of  $\mathbb{P}^1$ :

(10)



Since (as we saw in Section 5.1) the hyperelliptic curve  $D$  is supersingular, the double cover  $D \rightarrow \mathbb{P}^1$  of the top  $\mathbb{P}^1$  is ramified at only one point. The point  $\infty$  in the left  $\mathbb{P}^1$  ramifies in  $C_1 \rightarrow \mathbb{P}^1$ , so if two points of the top  $\mathbb{P}^1$  mapped to  $\infty$  in the left  $\mathbb{P}^1$ , both of those points would ramify in the double cover  $D \rightarrow \mathbb{P}^1$ , a contradiction. Therefore  $\infty$  ramifies in the top left cover  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ , and similarly in the top right cover  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ . It follows that  $\infty$  in the left  $\mathbb{P}^1$  and  $\infty$  in the right  $\mathbb{P}^1$  map to the same point in the bottom  $\mathbb{P}^1$ , and this point in the bottom  $\mathbb{P}^1$  ramifies going up to the left and going up to the right.

We can choose a coordinate  $z$  on the bottom  $\mathbb{P}^1$  so that this common image point is  $\infty$ . Then the bottom left map  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$  is of the form  $z = a^2x^2 + b^2x + e_1^2$  and the bottom right map  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$  is  $z = c^2u^2 + d^2u + e_2^2$ . (We again take the coefficients to be squares to avoid square root signs later.) Setting  $e = e_1 + e_2$ , we find that the  $V_4$ -diagram of copies of  $\mathbb{P}^1$  is specified by the relation

$$(11) \quad a^2x^2 + b^2x + c^2u^2 + d^2u + e^2 = 0,$$

for some  $a, b, c, d, e \in k$ . We note that  $a, b, c$ , and  $d$  must all be nonzero in order for the bottom left and bottom right covers  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$  to be separable maps of degree 2. The conic defined by this equation is the  $\mathbb{P}^1$  that appears in the top of the  $V_4$  diagram, so the conic must be nonsingular, which means simply that  $ad^2 + b^2c \neq 0$ .

(At this point, the coefficients in Equation (11) are only specified up to a scaling factor; in a moment we will see that there is a unique way of scaling the coefficients so that we have both  $b = Ac^2$  and  $d = Ba^2$ . Scaled in this way, the coefficients  $a$  and  $c$  will give the  $(a, c)$  pair specified in the statement of the theorem.)

If we take  $t = ax + cu$ , then  $t$  generates the function field of the top  $\mathbb{P}^1$ , and we have

$$(12) \quad x = (ct^2 + d^2t + ce^2)/(b^2c + d^2a)$$

$$(13) \quad u = (at^2 + b^2t + ae^2)/(b^2c + d^2a).$$

Equation (11), together with the equations for the double covers  $C_1 \rightarrow \mathbb{P}^1$  and  $C_2 \rightarrow \mathbb{P}^1$ , defines a 1-dimensional variety. We still need to check that this variety is isomorphic to the curve  $D$  and that we get the  $D_8$  extension given in Diagram (5). As a first step, we take a look at the maps that are supposed to give the anti-commuting involutions  $\alpha$  and  $\beta$  of  $D$ .

Recall that Diagram (6) shows a  $V_4$  extension extracted from Diagram (5). The middle extension  $C_1 \rightarrow \mathbb{P}^1$  in Diagram (6) is the Artin–Schreier extension

$$y^2 + y = A^2x^5 + A^2x^3.$$



The extension  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$  on the right is given by Equation (11), which we can scale and rewrite as an Artin–Schreier equation:

$$\left(\frac{c^2}{d^2}u\right)^2 + \left(\frac{c^2}{d^2}u\right) = \frac{a^2c^2}{d^4}x^2 + \frac{b^2c^2}{d^4}x + \frac{e^2c^2}{d^4}.$$

Therefore the double cover  $C'_1 \rightarrow \mathbb{P}^1$  is given by

$$\left(y + \frac{c^2}{d^2}u\right)^2 + \left(y + \frac{c^2}{d^2}u\right) = A^2x^5 + A^2x^3 + \frac{a^2c^2}{d^4}x^2 + \frac{b^2c^2}{d^4}x + \frac{e^2c^2}{d^4}.$$

Consider the involution  $\beta$  of  $D$ , which is supposed to fix  $C_2$  and take  $C_1$  to  $C'_1$ . On the  $V_4$  diagram of copies of  $\mathbb{P}^1$ , the involution  $\beta$  must then act trivially on the rightmost  $\mathbb{P}^1$  and nontrivially on the leftmost  $\mathbb{P}^1$ . Since  $\beta$  fixes  $C_2$  we must have  $\beta^*u = u$  and  $\beta^*v = v$ . Since  $\beta$  acts nontrivially on the leftmost  $\mathbb{P}^1$  but fixes the bottom  $\mathbb{P}^1$ , we must have  $\beta^*x = x + b^2/a^2$ . And since  $\beta$  takes  $C_1$  to  $C'_1$ , we must have

$$\beta^*y = y + \frac{c^2}{d^2}u + F$$

for some  $F \in k(x)$ . Applying  $\beta^*$  to  $y^2 + y = A^2x^5 + A^2x^3$  we find that

$$\left(y + \frac{c^2}{d^2}u + F\right)^2 + \left(y + \frac{c^2}{d^2}u + F\right) = A^2\left(x + \frac{b^2}{a^2}\right)^5 + A^2\left(x + \frac{b^2}{a^2}\right)^3,$$

which simplifies to

$$F^2 + F = \frac{A^2b^2}{a^2}x^4 + \left(\frac{A^2b^2}{a^2} + \frac{a^2c^2}{d^4}\right)x^2 + \left(\frac{A^2b^8}{a^8} + \frac{A^2b^4}{a^4} + \frac{b^2c^2}{d^4}\right)x + \left(\frac{A^2b^{10}}{a^{10}} + \frac{A^2b^6}{a^6} + \frac{e^2c^2}{d^4}\right).$$

Therefore  $F = f_2x^2 + f_1x + f_0$  for some  $f_2, f_1, f_0 \in k$ . Looking at the coefficient of  $x^4$  in the preceding equality, we find that  $f_2 = Ab/a$ . Looking at the coefficient of  $x$ , we find

$$f_1 = \frac{A^2b^8}{a^8} + \frac{A^2b^4}{a^4} + \frac{b^2c^2}{d^4}.$$

We also have the condition that  $\beta$  is an involution, which implies that

$$f_2x^2 + f_1x + f_0 + f_2\left(x + \frac{b^2}{a^2}\right)^2 + f_1\left(x + \frac{b^2}{a^2}\right) + f_0 = 0;$$

this simplifies to  $a^2f_1 + b^2f_2 = 0$ . Combining these three relations, we find that

$$(14) \quad 0 = A^2a^4b^2d^4 + A^2b^6d^4 + Aa^5bd^4 + a^8c^2.$$

It is not hard to check that that if this relation holds (and if we choose an appropriate value for  $f_0$ ), then indeed  $F^2 + F$  is equal to the appropriate polynomial expression in  $x$  given above.

The same argument, applied to the involution  $\alpha$ , shows that we must have

$$(15) \quad 0 = B^2b^4c^4d^2 + B^2b^4d^6 + Bb^4c^5d + a^2c^8.$$

Next we check that the extension of the top  $\mathbb{P}^1$  in Diagram (10) we get from  $C_1$  is isomorphic to the one we get from  $C_2$ . This condition simply means that the Artin–Schreier extension of  $k(t)$  we get from Equation (12) and  $y^2 + y = A^2x^5 + A^2x^3$  should be isomorphic to the Artin–Schreier extension we get from Equation (13) and  $v^2 + v = B^2u^5 + B^2u^3$ . This means that the expression

$$S = A^2x^5 + A^2x^3 + B^2u^5 + B^2u^3,$$

viewed as a polynomial in  $t$ , should be expressible as  $T^2 + T$  for some  $T \in k[t]$ . If we write

$$S = s_{10}t^{10} + s_9t^9 + s_8t^8 + \cdots + s_1t + s_0,$$

then  $S$  can be written  $T^2 + T$  if and only if we have

$$(16) \quad 0 = s_9$$

$$(17) \quad 0 = s_7$$

$$(18) \quad 0 = s_{10} + s_5^2$$

$$(19) \quad 0 = s_6 + s_3^2$$

$$(20) \quad 0 = s_8 + s_4^2 + s_2^4 + s_1^8.$$

Consider Equation (16). With a computer algebra system it is easy to check that

$$s_9 = \frac{(Ba^2b + Ac^2d)^2}{(ad^2 + b^2c)^5},$$

so Equation (16) is equivalent to  $Ba^2b = Ac^2d$ . Now, all of our conditions on the variables  $a$ ,  $b$ ,  $c$ ,  $d$ , and  $e$  have been homogeneous, because these variables appear as coefficients in the conic defined by Equation (11). The condition that  $Ba^2b = Ac^2d$  gives us an opportunity to remove this homogeneity in a natural way: We scale all five variables by the (unique) factor that makes the condition  $b = Ac^2$  hold. Then the relation  $Ba^2b = Ac^2d$  from Equation (16) shows that we also have  $d = Ba^2$ . This allows us to eliminate  $b$  and  $d$  from our equations. In particular, we find that Equations (14) and (15) become

$$(21) \quad 0 = A^4B^8a^{10} + A^4B^4a^2c^4 + A^4B^2c^5 + 1$$

$$(22) \quad 0 = A^2B^4a^5 + A^4B^4a^4c^2 + A^8B^4c^{10} + 1.$$

Also note that the condition that the conic given by Equation (11) be nonsingular — namely, that  $ad^2 + b^2c \neq 0$  — becomes Inequality (9).

With these new scaled variables, we continue to check the conditions required for  $S$  to be of the form  $T^2 + T$ . Again, it is simplest to use a computer algebra system. We find that Equation (17) is automatically satisfied. We find that

$$s_{10} + s_5^2 = \frac{(A^2B^6a^{10} + B^2a^5 + A^2B^2a^4c^2 + A^2B^2a^2c^4 + A^6B^2c^{10} + A^2c^5)^2}{(B^2a^5 + A^2c^5)^6},$$

so Equation (18) is equivalent to Equation (8) from the statement of the theorem.

To analyze Equation (19), we rely on our computer algebra system once again. First of all, we find that Equation (19) holds if and only if

$$(23) \quad 0 = A^4B^{12}a^{24} + A^2B^{10}a^{21}c + B^8a^{18} + A^4B^8a^{16}c^6 + A^2B^6a^{15}c^3 \\ + A^2B^6a^{13}c^5 + A^4B^4a^{10}c^8 + A^4B^4a^8c^{10} + A^8B^4a^6c^{16} \\ + A^6B^2a^5c^{13} + A^6B^2a^3c^{15} + A^{10}B^2ac^{21} + A^{12}B^4c^{24} + A^8c^{18}.$$

For a moment, let us view  $a$ ,  $c$ ,  $A$ , and  $B$  as indeterminates. Let  $t_1$ ,  $t_2$ , and  $t_3$  be the polynomials in  $\mathbb{F}_2[a, c, A, B]$  that appear on the right-hand sides of Equations (21), (22), and (23), respectively, and let  $r_1$  be the polynomial that appears on the right-hand side of Equation (7). We check via computer algebra that the ideal of  $\mathbb{F}_2[a, c, A, B]$  generated by  $t_1$ ,  $t_2$ , and  $t_3$  is equal to the ideal generated by  $t_1$ ,  $t_2$ , and  $r_1$ . Thus, demanding that Equations (21), (22), and (23) hold is equivalent to demanding that Equations (21), (22), and (7) hold.

We turn to our final condition on creating a diagram: Equation (20). We find that

$$s_8 + s_4^2 + s_2^4 + s_1^8 = \frac{(p_{16}e^{16} + p_8e^8 + p_4e^4 + p_2e^2 + p_1e + p_0)^2}{(B^2a^5 + A^2c^5)^{24}},$$

where

$$\begin{aligned}
p_{16} &= (B^2a^5 + A^2B^2a^4c^2 + A^2B^2a^2c^4 + A^2c^5)^4 \\
p_8 &= (B^2a^3 + A^2c^3)^2(B^2a^5 + A^2c^5)^6 \\
p_4 &= A^4B^4a^2c^2(B^2a^5 + A^2c^5)^6(B^2a^7 + A^2c^7)^2 \\
p_2 &= (B^2a^5 + A^2c^5)^8 \\
&\quad \cdot (A^2B^6a^{11}c + B^4a^8 + A^4B^4a^6c^6 + A^2B^2a^5c^3 + A^2B^2a^3c^5 + A^6B^2ac^{11} + A^4c^8) \\
p_1 &= (B^2a^5 + A^2c^5)^{10} \\
p_0 &= A^2B^2ac(B^2a^5 + A^2c^5)^9(B^2a^7 + A^2c^7).
\end{aligned}$$

However, we note that the term  $B^2a^5 + A^2B^2a^4c^2 + A^2B^2a^2c^4 + A^2c^5$  appearing in  $p_{16}$  is equal to

$$A^2B^2(B^2a^5 + A^2c^5)^2 + r_2,$$

where  $r_2$  is the expression on the right-hand side of Equation (8), and the term

$$A^2B^6a^{11}c + B^4a^8 + A^4B^4a^6c^6 + A^2B^2a^5c^3 + A^2B^2a^3c^5 + A^6B^2ac^{11} + A^4c^8$$

appearing in  $p_2$  is equal to

$$(A^2B^2a^3c^3 + a^2 + c^2)^2 + r_1 + acr_2,$$

so

$$p_{16}e^{16} + p_8e^8 + p_4e^4 + p_2e^2 + p_1e + p_0 = (B^2a^5 + A^2c^5)^6(q_{16}e^{16} + q_8e^8 + q_4e^4 + q_2e^2 + q_1e + q_0)$$

where

$$\begin{aligned}
q_{16} &= A^8B^8(B^2a^5 + A^2c^5)^2 \\
q_8 &= (B^2a^3 + A^2c^3)^2 \\
q_4 &= A^4B^4a^2c^2(B^2a^7 + A^2c^7)^2 \\
q_2 &= (B^2a^5 + A^2c^5)^2(A^2B^2a^3c^3 + a^2 + c^2)^2 \\
q_1 &= (B^2a^5 + A^2c^5)^4 \\
q_0 &= A^2B^2ac(B^2a^5 + A^2c^5)^3(B^2a^7 + A^2c^7).
\end{aligned}$$

For a given choice of  $a$  and  $c$ , a value of  $e$  satisfies Equation 20 if and only if it is a root of

$$(24) \quad q_{16}T^{16} + q_8T^8 + q_4T^4 + q_2T^2 + q_1T + q_0 = 0.$$

Note that  $q_{16}$  and  $q_1$  are both nonzero, so for a given choice of  $a$  and  $c$  there are exactly 16 values of  $e \in k$  that satisfy Equation 20. We will show that different choices for  $e$  lead to isomorphic diagrams.

Lemma 2.1 says that for every root  $x_0$  of  $f := T^{16} + T^8 + A^{-4}T^2 + A^{-6}T$  there are two automorphisms of  $C_1$  that lie over the automorphism  $x \mapsto x + x_0$  of  $\mathbb{P}^1$ . (We use  $A^{-4}$  and  $A^{-6}$  as coefficients in the polynomial  $f$  instead of the  $A^{-2}$  and  $A^{-3}$  that appear in the lemma because our curve uses an  $A^2$  where the curve in the lemma uses an  $A$ .) Shifting  $x$  by  $x_0$  changes the equation for the conic in Equation (11) by changing  $e$  to  $e + e_0$ , where

$$e_0 = ax_0 + b\sqrt{x_0} = ax_0 + c^2A\sqrt{x_0}.$$

Since  $e + e_0$  must also be a root of Equation (24), we see that  $e_0$  must be a root of the additive polynomial  $g := q_{16}T^{16} + q_8T^8 + q_4T^4 + q_2T^2 + q_1T$ . Thus, the map  $x_0 \mapsto ax_0 + Ac^2\sqrt{x_0}$  is a homomorphism from the (additive group of the) roots of the polynomial  $f$  to those of  $g$ . If this map is surjective, then the isomorphism class of the diagram obtained from a given choice of  $a, c$ ,

and  $e$  does not depend on the choice of  $e$ , so it will suffice for us to show that no nonzero root  $x_0$  of  $f$  satisfies  $ax_0 + Ac^2\sqrt{x_0} = 0$ .

Suppose, to obtain a contradiction, that  $x_0$  is a nonzero root of  $f$  such that  $ax_0 + Ac^2\sqrt{x_0} = 0$ . Then  $x_0 = A^2c^4/a^2$ , and plugging this expression into  $f$  we find that

$$(25) \quad \begin{aligned} 0 &= a^{30}c^4 + A^4a^{28}c^8 + A^{20}a^{16}c^{32} + A^{36}c^{64} \\ &= c^4(a^5 + A^2a^4c^2 + A^6c^{10})^2(a^{10} + A^6a^5c^{10} + A^8a^4c^{12} + A^{12}c^{20})^2. \end{aligned}$$

The first factor on the right-hand side of Equation (25) is nonzero because  $c \neq 0$ . The second factor is also nonzero, because from Equation (22) we see that

$$a^5 + A^2a^4c^2 + A^6c^{10} = \frac{A^2B^4a^5 + A^4B^4a^4c^2 + A^8B^4c^{10}}{A^2B^4} = \frac{1}{A^2B^4} \neq 0.$$

And the third factor is nonzero as well, because we have

$$B^4(a^{10} + A^6a^5c^{10} + A^8a^4c^{12} + A^{12}c^{20}) = (B^2a^5 + A^2c^5)^2 + A^4c^{10}(A^2B^4a^5 + A^4B^4a^4c^2 + A^8B^4c^{10} + 1),$$

and the first term on the right-hand side is nonzero by Inequality (9) while the second term is zero by Equation (22). Thus, the right-hand side of Equation (25) is nonzero, while the left-hand side is zero. This contradiction shows that for given values of  $a$  and  $c$ , the isomorphism class of Diagram (5) does not depend on the choice of  $e$  satisfying Equation (20).

In short, when  $C_1$  and  $C_2$  are supersingular genus-2 curves over  $k$  specified by nonzero  $A, B \in k$  as in the statement of the theorem, then every diagram of the shape of Diagram (5) gives rise to a unique pair  $(a, c)$  of nonzero elements of  $k$  that satisfies Equations (7) and (8), Inequality (9), and Equations (21) and (22). And our analysis works in reverse, as well: Given nonzero  $a$  and  $c$  in  $k$  that satisfy these four equalities and one inequality, we set  $b = c^2A$  and  $d = a^2B$  and choose a value of  $e$  such that Equation (24) holds — and then the  $V_4$  diagram of copies of  $\mathbb{P}^1$  specified by the conic in Equation (11) extends to give a  $D_8$  diagram like Diagram (5).

To complete the proof, we need only show that any pair  $(a, c)$  of nonzero elements of  $k$  that satisfies Equations (7) and (8) and Inequality (9) also satisfies Equations (21) and (22). If we let  $r_1, r_2,$  and  $r_3$  be the quantities on the right-hand sides of (7), (8), and (9), and if we let  $t_1$  and  $t_2$  be the quantities on the right-hand sides of (21) and (22), we see that

$$\begin{aligned} r_3^2t_1 &= r_2^2 + A^6B^2c^{10}r_2 + A^4B^4a^4c^4r_1 \\ r_3^2t_2 &= r_2^2 + A^2B^6a^{10}r_2 + A^4B^4a^4c^4r_1. \end{aligned}$$

Since  $r_1 = r_2 = 0$  and  $r_3 \neq 0$ , we find that  $t_1 = t_2 = 0$ .  $\square$

The same strategy leads to the following two results, which deal with the cases where one or both of  $C_1$  and  $C_2$  have supersingular invariant equal to 0; we leave the details of the proofs to the reader.

**Theorem 5.5.** *Up to isomorphism, diagrams in the form of Diagram (5) with  $C_1$  and  $C_2$  given by  $y^2 + y = A^2(x^5 + x^3)$  and  $v^2 + v = u^5$ , respectively, are in bijection with orbits of pairs  $(a, c)$  of nonzero elements of  $k$  satisfying*

$$(26) \quad 0 = a^6c + a^4 + A^2ac^6 + A^4c^8$$

$$(27) \quad 0 = A^2a^{10} + a^5 + A^2a^4c^2 + A^6c^{10} + A^2c^5$$

$$(28) \quad 0 \neq a^5 + c^5A^2,$$

under the action of the fifth roots of unity given by  $\zeta \cdot (a, c) = (\zeta a, \zeta^3 c)$ .  $\square$

**Theorem 5.6.** *There are no diagrams in the form of Diagram (5) when  $C_1$  and  $C_2$  are given by  $y^2 + y = x^5$  and  $v^2 + v = u^5$ , respectively.  $\square$*

**Remark 5.7.** We will make one comment about a possibly non-obvious point in the proof of Theorem 5.5. As in the proof of Theorem 5.4, we derive equations from the requirement that there be anti-commuting involutions of  $D$ ; for Theorem 5.5, the analogs of Equations (21) and (22) are

$$(29) \quad 0 = A^4 a^{10} + A^4 c^5 + 1$$

$$(30) \quad 0 = A^2 a^5 + A^4 a^4 c^2 + A^8 c^{10} + 1.$$

If we let  $t_1$  and  $t_2$  be the right-hand sides of these equations, and if we let  $r_1$ ,  $r_2$ , and  $r_3$  be the right-hand sides of (26), (27), and (28), then we have

$$r_3 t_1 = A^2 c^2 (A^2 a^5 + A^4 c^5 + 1) r_1 + (A^2 a^5 + A^2 a c^3 + A^4 c^5 + 1) r_2$$

$$r_3 t_2 = A^2 c^2 (A^2 a^5 + A^4 c^5 + 1) r_1 + (A^2 a c^3 + 1) r_2.$$

Thus, Equations (26) and (27) and Inequality (28) imply Equations (29) and (30).

**5.5. Counting isogenies.** Now we are in a position to show that the dihedral construction, the degenerate construction, the Frobenius, and the Verschiebung account for all of the Richelot isogenies between two supersingular curves. First we look at the case of two curves whose supersingular invariants are nonzero, so that according to Theorem 1.1 there are 60 isogenies to account for.

**Theorem 5.8.** *Let  $A$  and  $B$  be nonzero elements of  $k$ , and let  $C_1$  and  $C_2$  be the curves given by  $y^2 + y = A^2(x^5 + x^3)$  and  $v^2 + v = B^2(u^5 + u^3)$ , respectively. The number of isomorphism classes of Richelot isogenies from the Jacobian of  $C_1$  to the Jacobian of  $C_2$  coming from the dihedral construction, from the degenerate construction, from Frobenius, and from Verschiebung are as given in Table 1, and these account for all of the Richelot isogenies from the Jacobian of  $C_1$  to the Jacobian of  $C_2$ .*

TABLE 1. The number of isomorphism classes of Richelot isogenies from the Jacobian of  $y^2 + y = A^2(x^5 + x^3)$  to the Jacobian of  $v^2 + v = B^2(u^5 + u^3)$  coming from the dihedral construction, from the degenerate construction, from Frobenius, and from Verschiebung, depending on whether  $A = B$ ,  $A = B^2$ , and  $B = A^2$ . The fourth row gives the case where  $A$  and  $B$  are distinct primitive cube roots of unity, and the sixth row gives the case when  $A = B = 1$ .

Conditions on $A$ and $B$ :			Number of isogenies coming from...			
$A = B?$	$A = B^2?$	$B = A^2?$	Dihedral	Degen.	Frob.	Ver.
no	no	no	60	0	0	0
no	no	yes	59	0	1	0
no	yes	no	59	0	0	1
no	yes	yes	58	0	1	1
yes	no	no	50	10	0	0
yes	yes	yes	48	10	1	1

*Proof.* First let us consider the number of isogenies coming from the dihedral construction. From Proposition 5.1 and Theorem 5.4 we know that the number of isomorphism classes of Richelot isogenies from  $C_1$  to  $C_2$  is given by the number of pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (7) and (8) and Inequality (9).

As in the proof of Theorem 5.4, we view  $a$ ,  $c$ ,  $A$ , and  $B$  as indeterminates, we let  $r_1$  and  $r_2$  be the polynomials in  $\mathbb{F}_2[a, c, A, B]$  that appear on the right-hand sides of Equations (7) and (8), respectively, and we let  $r_3$  be the polynomial on the right-hand side of Inequality (9).

Given two elements  $g_1, g_2$  in  $\mathbb{F}_2[a, c, A, B]$  and a variable  $v \in \{a, c, A, B\}$ , we let  $\text{Res}_v(g_1, g_2)$  denote the resultant of  $g_1$  and  $g_2$  with respect to the variable  $v$  and we let  $\text{Disc}_v(g_1)$  denote the discriminant of  $g_1$  with respect to the variable  $v$ . We compute that

$$(31) \quad \text{Res}_c(r_1, r_2) = A^{24}(A+B)^8 a^{20} f,$$

where

$$\begin{aligned} f = & A^{24}B^{48}a^{60} + A^{18}B^{36}a^{45} + A^{16}B^{32}a^{40} + A^{14}B^{28}a^{33} + A^{12}B^{24}a^{28} \\ & + A^{10}B^{20}a^{25} + A^{10}B^{20}a^{21} + A^8B^{16}a^{20} + A^8B^{16}a^{18} + A^8B^{16}a^{16} \\ & + A^8B^{16}a^{12} + A^6B^{12}a^9 + A^4B^8a^8 + A^4B^8a^6 + A^4B^8a^4 \\ & + A^2B^4a^5 + A^2B^4a^3 + A^2B^4a + (A+B^2)^4. \end{aligned}$$

We further compute that

$$(32) \quad \text{Disc}_a f = A^{1416}B^{2832}(A+B^2)^{176}.$$

Suppose now we are given specific nonzero values of  $A$  and  $B$  in  $k$  with  $A \neq B^2$  and  $B \neq A^2$  and  $A \neq B$ . From Equations (31) and (32) we find that there are exactly 60 nonzero values of  $a \in k$  for which there exists a  $c \in k$  with  $r_1(a, c) = r_2(a, c) = 0$ . We claim that each of these values of  $c$  is nonzero. To see this, note that if  $c$  were 0 we would have  $0 = r_1(a, 0) = a^4(Ba+1)^4$  and  $0 = r_2(a, 0) = B^2a^5(A^2B^4a^5+1)$ ; the first relation gives  $a = 1/B$  and then the second gives  $B = A^2$ , a contradiction. Thus there are exactly 60 pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (7) and (8). We must still consider whether any of these pairs fails to satisfy Inequality (9); that is, we must check to see whether  $r_3 = 0$  for any of these pairs.

We check that  $\text{Res}_c(r_2, r_3) = A^{22}B^{14}(A+B)^4a^{20}$ , and since we have assumed that  $A \neq B$ , every one of the 60 pairs  $(a, c)$  satisfies Inequality (9). This gives us the count of dihedral Richelot isogenies on the first line of Table 1.

To see what happens when  $A \neq B^2$  and  $B \neq A^2$  but  $A = B$ , we repeat the computations we have above, but this time we work in the polynomial ring  $\mathbb{F}_2[a, c, A]$  and set  $B = A$ . (Note that the assumptions that  $A \neq B^2$  and  $B \neq A^2$  imply that  $A \neq 1$ .) When  $A = B$ , the polynomial  $r_3$  has a factor  $a+c$ , so we must avoid solutions with  $a=c$ . The polynomials  $r_1$  and  $r_2$  also have factors of  $a+c$ , and we write  $r_1 = (a+c)u_1$  and  $r_2 = (a+c)u_2$  for polynomials  $u_1, u_2 \in \mathbb{F}_2[a, c, A]$ .

We compute that

$$\text{Res}_c(u_1, u_2) = A^{20}a^{12}f_1f_2,$$

where

$$\begin{aligned} f_1 = & A^{24}a^{20} + A^{16}a^{12} + A^{14}a^{11} + A^{12}a^{10} \\ & + A^{12}a^8 + A^8a^6 + A^6a^3 + A^2a + (A+1)^4 \\ f_2 = & A^{32}a^{30} + A^{28}a^{26} + A^{26}a^{25} + A^{22}a^{21} + A^{20}a^{18} + A^{18}a^{17} \\ & + A^{14}a^{15} + A^{14}a^{13} + A^{10}a^{11} + (A^{12} + A^8)a^{10} + A^{10}a^9 \\ & + A^6a^7 + (A^8 + A^4)a^6 + (A^6 + A^2)a^5 + A^4a^4 + A^2a^3 + 1, \end{aligned}$$

and we find that

$$\text{Disc}_a f_1f_2 = A^{2660}(A+1)^{120}.$$

For a given value of  $A = B$  in  $k$ , with  $A \neq 0, 1$ , we find that there are exactly 50 nonzero values of  $a \in k$  for which there exists a  $c \in k$  with  $u_1(a, c) = u_2(a, c) = 0$ . The same argument we used before shows that none of these values of  $c$  can be equal to 0. We check that

$$\text{Res}_c(u_2, r_3) = A^{36}a^{24},$$

so every one of these 50 pairs  $(a, c)$  satisfies Inequality (9). This gives us the count of dihedral Richelot isogenies on the fifth line of Table 1.

Now let us consider the case where  $B = A^2$  but  $A \neq B^2$  (so that  $A \neq B$  as well). In this case, Equation (32) still shows that the discriminant of  $f$  is nonzero, so we again have 60 distinct nonzero values of  $a$  for which there exists a  $c \in k$  with  $r_1(a, c) = r_2(a, c) = 0$ . However, we check that now there is a (unique) nonzero value of  $a$  satisfying  $r_1(a, 0) = r_2(a, 0) = 0$ , namely  $a = 1/A^2$ . This shows that when  $B = A^2$  and  $A \neq B^2$  we have exactly 59 pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (7) and (8) and Inequality (9). This gives us the count of dihedral Richelot isogenies on the second line of Table 1. By symmetry, we get the count for the third line as well.

Now suppose  $A = B^2$  and  $B = A^2$  but  $A \neq B$ ; this happens precisely when  $A$  and  $B$  are distinct cube roots of unity. In this case we can explicitly compute the pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (7) and (8) and Inequality (9), and we find there are 58 such pairs. This gives us the count of dihedral Richelot isogenies on the fourth line of Table 1.

Likewise, when  $A = B^2$  and  $B = A^2$  and  $A = B$  (that is, when  $A = B = 1$ ), we find that there are 48 pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (7) and (8) and Inequality (9). This gives us the count of dihedral Richelot isogenies on the last line of Table 1.

We turn now to the question of counting degenerate Richelot isogenies. Degenerate isogenies require that  $C_1 \cong C_2$ ; that is, that  $A = B$ . This explains the four zero entries in the ‘‘Degenerate’’ column of Table 1.

Suppose  $A = B$ , and let  $C$  be the curve  $C_1 \cong C_2$ . We see from Proposition 5.2 that the number of degenerate Richelot isogenies is equal to the number of conjugacy classes of elements  $\gamma \in \text{Aut } C$  with  $\gamma^2 = \iota$ , the hyperelliptic involution. Statement 2(c) of Lemma 2.1 says that there are exactly 10 such conjugacy classes, and this gives us the count of degenerate Richelot isogenies on the last two lines of the table.

The entries for Frobenius and Verschiebung are self-explanatory.

We see that for each pair  $(A, B)$ , we can account for 60 distinct Richelot isogenies, the number given by Theorem 1.1.  $\square$

**Theorem 5.9.** *Let  $A$  be a nonzero elements of  $k$ , and let  $C_1$  and  $C_2$  be the curves given by  $y^2 + y = A^2(x^5 + x^3)$  and  $v^2 + v = u^5$ , respectively. Then up to isomorphism, there are 12 Richelot isogenies from the Jacobian of  $C_1$  to the Jacobian of  $C_2$  coming from the dihedral construction, and these account for all of the Richelot isogenies from the Jacobian of  $C_1$  to the Jacobian of  $C_2$ .*

*Proof.* From Proposition 5.1 and Theorem 5.5 we know that the number of isomorphism classes of Richelot isogenies from  $C_1$  to  $C_2$  is given by the number of orbits of pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (26) and (27) and Inequality (28), under the action of the fifth roots of unity given by  $\zeta \cdot (a, c) = (\zeta a, \zeta^3 c)$ .

As before, we view  $a$ ,  $c$ , and  $A$  as indeterminates, we let  $r_1$  and  $r_2$  be the polynomials in  $\mathbb{F}_2[a, c, A]$  that appear on the right-hand sides of Equations (26) and (27), respectively, and we let  $r_3$  be the polynomial on the right-hand side of Inequality (28).

We compute that  $\text{Res}_c(r_1, r_2) = A^{34}a^{20}f$ , where

$$f = A^{22}a^{60} + A^{16}a^{45} + A^{14}a^{40} + A^8a^{25} + A^6a^{20} + a^5 + A^2,$$

and we note that  $\text{Disc}_a f = A^{1356}$ . From this we see that given any specific nonzero value of  $A$ , there are exactly 60 nonzero values of  $a \in k$  for which there exists a  $c \in k$  with  $r_1(a, c) = r_2(a, c) = 0$ . Since  $r_1(a, 0) = a^4$ , we see immediately that these  $c$  must be nonzero, so we get 60 pairs  $(a, c)$  of nonzero elements of  $k$  that satisfy Equations (26) and (27). Since  $\text{Res}_c(r_2, r_3) = A^{26}a^{30}$ , each of these pairs also satisfies Inequality (28).

Under the action given in Theorem 5.5, the group of fifth roots of unity acts without fixed points on the set of these pairs, so there are 12 orbits. Therefore there are 12 isomorphism classes

of Richelot isogeny from  $C_1$  to  $C_2$  given by the dihedral construction, and, by Theorem 1.1, this accounts for all of the Richelot isogenies between these curves.  $\square$

**Theorem 5.10.** *Let  $C$  be the curve given by  $y^2 + y = x^5$ . There are exactly 2 isomorphism classes of Richelot isogeny from the Jacobian of  $C$  to itself coming from the degenerate construction, and these two degenerate isogenies, together with the Frobenius and the Verschiebung, represent all of the isomorphism classes of Richelot isogeny from the Jacobian of  $C$  to itself.*

*Proof.* Theorem 5.6 says that there are no Richelot isogenies from the Jacobian of  $C$  to itself coming from the dihedral construction. Proposition 5.2 and part 3(c) of Lemma 2.1 show that there are exactly 2 isomorphism classes of Richelot isogeny from the Jacobian of  $C$  to itself coming from the degenerate construction. The Frobenius and the Verschiebung are also Richelot isogenies from the Jacobian of  $C$  to itself, and Theorem 5.3 says that they are not isomorphic to one another. This gives us a total of four isomorphism classes of Richelot isogeny from the Jacobian of  $C$  to itself, and Theorem 1.1 says that there are no others.  $\square$

#### REFERENCES

- [1] Jean-Benoît Bost and Jean-François Mestre, *Moyenne arithmético-géométrique et périodes des courbes de genre 1 et 2*, Gaz. Math. (1988), no. 38, 36–64.
- [2] Everett W. Howe and Kristin E. Lauter, *New methods for bounding the number of points on curves over finite fields*, Geometry and arithmetic (C. Faber, G. Farkas, and R. de Jong, eds.), EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2012, pp. 173–212.
- [3] Tomoyoshi Ibukiyama, Toshiyuki Katsura, and Frans Oort, *Supersingular curves of genus two and class numbers*, Compositio Math. **57** (1986), no. 2, 127–152.
- [4] Toshiyuki Katsura and Frans Oort, *Families of supersingular abelian surfaces*, Compositio Math. **62** (1987), no. 2, 107–167.
- [5] Neal Koblitz,  *$p$ -adic variation of the zeta-function over families of varieties defined over finite fields*, Compositio Math. **31** (1975), no. 2, 119–218.
- [6] L. Königsberger, *Über die Transformation der Abelschen Functionen erster Ordnung*, J. Reine Angew. Math. **64** (1865), 17–42.
- [7] Yu. I. Manin, *Theory of commutative formal groups over fields of finite characteristic*, Russian Math. Surveys **18** (1963), no. 6, 1–83, translated by P. M. Cohn.
- [8] J. S. Milne, *Abelian varieties*, Arithmetic geometry (Storrs, Conn., 1984) (G. Cornell and J.H. Silverman, eds.), Springer, New York, 1986, pp. 103–150.
- [9] Frédéric Richelot, *Essai sur une méthode générale pour déterminer la valeur des intégrales ultra-elliptiques, fondée sur des transformations remarquables de ces transcendentes*, C. R. Acad. Sci. Paris **2** (1836), 622–627.
- [10] Fried. Jul. Richelot, *De transformatione integralium Abelianorum primi ordinis commentatio*, J. Reine Angew. Math. **16** (1837), 221–341.
- [11] William C. Waterhouse, *Abelian varieties over finite fields*, Ann. Sci. École Norm. Sup. (4) **2** (1969), 521–560.
- [12] ———, *Introduction to affine group schemes*, Graduate Texts in Mathematics, vol. 66, Springer-Verlag, New York, 1979.

CENTER FOR COMMUNICATIONS RESEARCH, 805 BUNN DRIVE, PRINCETON, NJ 08540-1966 USA

E-mail address: [brock@idaccr.org](mailto:brock@idaccr.org)

UNAFFILIATED MATHEMATICIAN, SAN DIEGO, CA, USA

URL: <http://ewhowe.com>

E-mail address: [however@alumni.caltech.edu](mailto:however@alumni.caltech.edu)