

TERNARY QUADRATIC FORMS REPRESENTING SAME INTEGERS

JANGWON JU

ABSTRACT. In 1997, Kaplansky conjectured that if two positive definite ternary quadratic forms with integer coefficients have perfectly identical integral representations, then they are isometric, both regular, or included either of two families of ternary quadratic forms. In this article, we prove the existence of pairs of ternary quadratic forms representing same integers which are not in the Kaplansky's list.

1. INTRODUCTION

For a positive definite quadratic form f of rank r with integer coefficients and a nonnegative integer n , we define a set

$$R(n, f) = \{(x_1, x_2, \dots, x_r) \in \mathbb{Z}^r \mid f(x_1, x_2, \dots, x_r) = n\}.$$

We also define $r(n, f) = |R(n, f)|$. It is well-known that if f is positive definite, then $r(n, f)$ is always finite. Determining $r(n, f)$ for an arbitrary positive definite quadratic form f is quite an old problem which is still widely open.

For a quadratic form f of rank r , the theta series of f is given by

$$\theta_f(z) = \sum_{\mathbf{x} \in \mathbb{Z}^r} q^{f(\mathbf{x})} = \sum_{n=0}^{\infty} r(n, f) q^n \quad (q = e^{2\pi iz}).$$

It is well known that $\theta_f(z)$ is a modular form of weight $\frac{r}{2}$. If two quadratic forms f and g are isometric over \mathbb{Z} , then clearly $\theta_f(z) = \theta_g(z)$. However, the converse is not true in general. In [17], Witt proved that there are two positive definite even unimodular quadratic forms of dimension 16 which have the same theta series. After that, such examples were found even in dimension 12, 8, and 4 (for details, see [10], [9] and [15]). In [2], Conway and Sloane found a 4parameter family of pairs of quaternary quadratic forms with same theta series. Furthermore, it was proved that the Conway-Sloane tetralattice pairs are not isometric over \mathbb{Z} (for details, see [1]). However, it is well known that unary and binary positive definite quadratic forms are determined up to integral equivalence by their theta series. Furthermore, Schiemann proved that if two positive definite ternary quadratic forms have same theta series, then they are isometric over \mathbb{Z} (for details, see [16]).

2010 *Mathematics Subject Classification.* 11E12, 11E20.

Key words and phrases. Representations of ternary quadratic forms, Kaplansky conjecture.

This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (NRF-2019R1F1A1064037).

For a positive definite quadratic form f with integer coefficients, we define the set $Q(f)$ of all nonnegative integers n such that $f(x, y, z) = n$ has an integer solution. For quadratic forms f_1, f_2, g_1 and g_2 , we write $\{f_1, g_1\} \simeq \{f_2, g_2\}$ if $f_1 \simeq f_2$ and $g_1 \simeq g_2$ or $f_1 \simeq g_2$ and $g_1 \simeq f_2$.

In 1938, Delone proved in [3] that for two binary quadratic forms f and g , if $Q(f) = Q(g)$, then they are isometric over \mathbb{Z} except the case when

$$\{f, g\} \simeq \{x^2 + xy + y^2, x^2 + 3y^2\}.$$

In 1997, Kaplansky conjectured in his letter to Schiemann that for two ternary quadratic forms f and g , if $Q(f) = Q(g)$, then one of the following holds:

- (i) $f \simeq g$;
- (ii) f and g are both regular;
- (iii) $\{f, g\} \simeq \{ax^2 + by^2 + bz^2 + byz, ax^2 + by^2 + 3bz^2\}$ for some $a, b \in \mathbb{Z}$;
- (iv) $\{f, g\} \simeq \{ax^2 + ay^2 + az^2 + byz + bxz + bxy, ax^2 + (2a-b)y^2 + (2a+b)z^2 + 2bxz\}$ for some $a, b \in \mathbb{Z}$.

Here, a quadratic form f is called regular if it represents all integers that are locally represented by f . It was proved that the conjecture holds if only diagonal ternary quadratic forms are considered (for details, see [4]).

Recently, Oishi-Tomiyasu gave a list of candidates of sets of ternary quadratic forms violating Kaplansky conjecture. Actually, it was proved that ternary quadratic forms contained in each set S_i for $i = 1, \dots, 15$ in Table 1.1 which are not isometric each other and are not regular represent same integers up to 3,000,000. Furthermore, she provided a modified version of Kaplansky conjecture (for details, see [13]). All sets in Table 1.1 except S_{13} was firstly provided by Jagy and he checked that each quadratic form in the sets represents same integers up to 1,000,000.

In this article, we prove the existence of pairs of ternary quadratic forms representing same integers which are not in the Kaplansky's list. Actually, we prove that each of two quadratic forms contained in S_4, S_6, S_7 and S_8 in Table 1.1 represents same integers.

Let

$$f(x_1, x_2, \dots, x_r) = \sum_{1 \leq i, j \leq r} a_{ij} x_i x_j \quad (a_{ij} = a_{ji} \in \mathbb{Z})$$

be a positive definite integral quadratic form. The corresponding integral symmetric matrix of f is defined by $M_f = (a_{ij})$. For an integer k , we say k is *represented by* f if the equation $f(x_1, x_2, \dots, x_m) = k$ has an integer solution.

Any unexplained notations and terminologies can be found in [8] or [14].

2. GENERAL TOOLS

In [7], [11], and [12], we developed a method that determines whether or not integers in an arithmetic progression are represented by some particular ternary quadratic form. We briefly introduce this method for those who are unfamiliar with it.

Let d be a positive integer and let a be a nonnegative integer ($a \leq d$). We define

$$S_{d,a} = \{dn + a \mid n \in \mathbb{N} \cup \{0\}\}.$$

Table 1.1. Ternary quadratic forms with same representations

S_1	$5x^2 + 8y^2 + 8z^2 - 5yz - xz - 4xy,$	$5x^2 + 5y^2 + 8z^2 - yz - 4xz - 2xy$
S_2	$3x^2 + 4y^2 + 7z^2 - yz,$	$3x^2 + 4y^2 + 7z^2 + 4yz + 3xz + 3xy$
S_3	$x^2 + 4y^2 + 7z^2 - yz,$	$x^2 + 4y^2 + 5z^2 - yz - xz$
S_4	$4x^2 + 7y^2 + 25z^2 - 4yz - 2xz - 2xy,$	$4x^2 + 7y^2 + 7z^2 + 5yz + 2xz + 2xy$
S_5	$2x^2 + 6y^2 + 41z^2 - 3yz - xz,$	$2x^2 + 2y^2 + 41z^2 + yz + 2xz + 2xy$
S_6	$2x^2 + 6y^2 + 14z^2 - 3yz - xz,$	$2x^2 + 2y^2 + 14z^2 + yz + 2xz + 2xy$
S_7	$2x^2 + 4y^2 + 8z^2 + 4yz + xz + xy,$	$2x^2 + 2y^2 + 4z^2 - yz - 2xz$
S_8	$5x^2 + 5y^2 + 8z^2 - 4xz - 3xy,$	$5x^2 + 7y^2 + 7z^2 + 6yz + xz + 5xy$
S_9	$3x^2 + 3y^2 + 7z^2 + yz + 2xz + xy,$	$3x^2 + 5y^2 + 5z^2 + 3yz + xz + 3xy$
S_{10}	$5x^2 + 5y^2 + 8z^2 - yz - 2xz - 4xy,$	$5x^2 + 5y^2 + 6z^2 - 3xz - 2xy$
S_{11}	$2x^2 + 4y^2 + 7z^2 - xz - xy,$	$2x^2 + 4y^2 + 7z^2 + 4yz + 2xz + xy$
S_{12}	$4x^2 + 6y^2 + 7z^2 + 3yz + 2xz + 3xy,$	$4x^2 + 4y^2 + 6z^2 - 3xz - 2xy$
S_{13}	$5x^2 + 12y^2 + 28z^2 - 4xz - 4xy,$	$5x^2 + 12y^2 + 24z^2 - 8yz - 4xy,$
	$5x^2 + 12y^2 + 21z^2 - 4yz - 2xz - 4xy,$	$5x^2 + 12y^2 + 12z^2 - 4xz - 4xy$
S_{14}	$3x^2 + 5y^2 + 7z^2 - 2yz - 2xy,$	$3x^2 + 5y^2 + 6z^2 - 2xz - 2xy,$
	$3x^2 + 5y^2 + 6z^2 + 4yz + 2xz + 2xy,$	$3x^2 + 3y^2 + 5z^2 - 2yz - 2xz$
S_{15}	$3x^2 + 5y^2 + 5z^2 + 5yz + 2xz + 3xy,$	$3x^2 + 3y^2 + 5z^2 - yz - 2xz - xy,$
	$3x^2 + 3y^2 + 5z^2 + 2yz + 3xz + xy,$	$3x^2 + 3y^2 + 3z^2 + yz + xz + 3xy$

For two integral ternary quadratic forms f, g , we define

$$R(g, d, a) = \{v \in (\mathbb{Z}/d\mathbb{Z})^3 \mid vM_gv^t \equiv a \pmod{d}\}$$

and

$$R(f, g, d) = \{T \in M_3(\mathbb{Z}) \mid T^t M_f T = d^2 M_g\}.$$

A coset (or, a vector in the coset) $v \in R(g, d, a)$ is said to be *good* with respect to f, g, d , and a if there is a $T \in R(f, g, d)$ such that $\frac{1}{d} \cdot vT^t \in \mathbb{Z}^3$. The set of all good vectors in $R(g, d, a)$ is denoted by $R_f(g, d, a)$. If $R(g, d, a) \setminus R_f(g, d, a) = \emptyset$, then we write

$$g <_{d,a} f.$$

If $g <_{d,a} f$, then by Lemma 2.2 of [11], we have

$$(2.1) \quad S_{d,a} \cap Q(g) \subset Q(f).$$

In general, if d is large, then it is hard to compute the set $R(g, d, a) \setminus R_f(g, d, a)$ exactly by hand. A MAGMA based computer program for computing the set could be available upon request to the author.

3. REPRESENTATIONS OF TERNARY QUADRATIC FORMS

In this section, we prove that each of two quadratic forms contained in S_4, S_6, S_7 and S_8 in Table 1.1 represents same integers.

For some technical reason, we consider integral quadratic forms obtained from quadratic forms in Table 1.1 by scaling 2. Let m be a positive integer. Note that for two quadratic forms f and g , $Q(f) = Q(g)$ if and only if $Q(mf) = Q(mg)$.

(i) Quadratic forms in S_4 . Let $f(x, y, z) = 8x^2 + 14y^2 + 50z^2 - 8yz - 4xz - 4xy$ and $g(x, y, z) = 8x^2 + 14y^2 + 14z^2 + 10yz + 4xz + 4xy$. Then each symmetric matrix corresponding to f and g is as follows:

$$M_f = \begin{pmatrix} 8 & -2 & -2 \\ -2 & 14 & -4 \\ -2 & -4 & 50 \end{pmatrix} \quad \text{and} \quad M_g = \begin{pmatrix} 8 & 2 & 2 \\ 2 & 14 & 5 \\ 2 & 5 & 14 \end{pmatrix}.$$

One may easily show that for $T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -2 \\ 0 & -1 & 1 \end{pmatrix}$, $T^t M_g T = M_f$, which implies that f is a subform of g . Therefore, $Q(f) \subset Q(g)$.

It is enough to show that every positive even integer represented by g is also represented by f . One may easily compute that

$$R(g, 4, 0) = \{(v_1, v_2, v_3) \in (\mathbb{Z}/4\mathbb{Z})^3 \mid v_2 \equiv 0 \pmod{2}, v_3 \equiv 0 \pmod{2}\}.$$

Furthermore, note that $R(f, g, 4)$ consists of 8 matrices, and in particular, it contains the following matrix:

$$T_1 = \begin{pmatrix} 4 & 2 & 2 \\ 0 & 4 & 2 \\ 0 & 0 & 2 \end{pmatrix}.$$

Then, we know that $\frac{1}{4} \cdot (v_1, v_2, v_3) \cdot T_1^t \in \mathbb{Z}^3$, for any $(v_1, v_2, v_3) \in R(g, 4, 0)$. Therefore, $g <_{4,0} f$ holds. Then by (2.1), every positive integer congruent to 0 modulo 4 which is represented by g is also represented by f . As a sample, for $(1, 4, 2) \in \mathbb{Z}^3$, note that $g(1, 4, 2) = 392$ and $(1, 4, 2)$ is congruent to $(1, 0, 2)$ modulo 4 which is contained in $R(g, 4, 0)$. Then

$$f(4, 5, 1) = f\left(\frac{1}{4} \cdot (1, 4, 2) \cdot T_1^t\right) = g(1, 4, 2) = 392.$$

Similarly, one may easily show that

$$g <_{12,6} f, \quad g <_{12,10} f.$$

In fact, $Q(g) \cap \{12n + 10 \mid n \in \mathbb{N} \cup \{0\}\} = \emptyset$. However, we consider this case for completeness of the proof. It also happens for some other cases that we will consider later. Therefore, by (2.1) every positive integer congruent to 6 or 10 modulo 12 which is represented by g is also represented by f . Now, for some nonnegative integer n_1 and $v \in \mathbb{Z}^3$, assume that $g(v) = 12n_1 + 2$. One may easily check that there are exactly 864 vectors in $R(g, 12, 2)$ and 144 matrices in $R(f, g, 12)$. Furthermore, all vectors in $R(g, 12, 2)$ are good vectors with respect to $f, g, 12$ and 2 except 32 vectors. Note that

$$\begin{aligned} & R(g, 12, 2) \setminus R_f(g, 12, 2) \\ &= \{(v_1, v_2, v_3) \in (\mathbb{Z}/12\mathbb{Z})^3 \mid v_1 \not\equiv 0 \pmod{3}, v_2 \equiv \pm 3 \pmod{12}, v_3 \equiv \pm 3 \pmod{12}\}. \end{aligned}$$

Now, define

$$\tilde{T} = \begin{pmatrix} 12 & 6 & 2 \\ 0 & 0 & 12 \\ 0 & -12 & -8 \end{pmatrix}.$$

Note that $\tilde{T}^t M_g \tilde{T} = 12^2 M_g$. Then one may easily check that for any vector $u \in \mathbb{Z}^3$ such that $u \pmod{12} \in R(g, 12, 2) \setminus R_f(g, 12, 2)$,

$$(3.1) \quad \frac{1}{12} \cdot u \tilde{T}^t \in \mathbb{Z}^3.$$

All computations were done by a computer program based on MAGMA. If $v \pmod{12}$ is a good vector with respect to $f, g, 12$ and 2 , then there is a $T_2 \in R(f, g, 12)$ such that $\frac{1}{12} \cdot v T_2^t \in \mathbb{Z}^3$ and $f(\frac{1}{12} \cdot v T_2^t) = 12n_1 + 2$. Assume that $v \pmod{12}$ is not a good vector. Then by (3.1), we know that $\frac{1}{12} \cdot v \tilde{T}^t \in \mathbb{Z}^3$. If $\frac{1}{12} \cdot v \tilde{T}^t \pmod{12} \in R_f(g, 12, 2)$, then we are done. If $\frac{1}{12} \cdot v \tilde{T}^t \pmod{12}$ is not good, then $(\frac{1}{12})^2 \cdot v (\tilde{T}^t)^2 \in \mathbb{Z}^3$ by (3.1). Now inductively, we may assume that for any positive integer m ,

$$\left(\frac{1}{12}\right)^m \cdot v (\tilde{T}^t)^m \in \mathbb{Z}^3 \text{ and } \left(\frac{1}{12}\right)^m \cdot v (\tilde{T}^t)^m \pmod{12} \in R(g, 12, 2) \setminus R_f(g, 12, 2).$$

Since there are only finitely many integer solution of $g(x, y, z) = 12n_1 + 2$ and $\frac{1}{12} \tilde{T}$ has an infinite order, it is impossible, unless v is an eigenvector of \tilde{T} . Note that $(\pm 1, 0, 0)$ are the only integral primitive eigenvectors of \tilde{T} . Hence, if $12n_1 + 2$ is not of the form $g(\pm t, 0, 0) = 8t^2$ for some positive integer t , then it is also represented by f . Furthermore, $8t^2$ is represented by f since 8 is represented by f . Therefore $12n_1 + 2$ is represented by f . Hence, every positive even integer which is represented by g is also represented by f . Therefore, f and g have perfectly identical integral representations.

(ii) Quadratic forms in S_6 . Let $f(x, y, z) = 4x^2 + 12y^2 + 28z^2 - 6yz - 2xz$ and $g(x, y, z) = 4x^2 + 4y^2 + 28z^2 + 2yz + 4xz + 4xy$. One may easily show that f is a subform of g . Therefore $Q(f) \subset Q(g)$.

It is enough to show that every positive even integer represented by g is also represented by f . One may easily show that

$$g <_{4,2} f, \quad g <_{8,0} f, \quad g <_{24,12} f, \quad g <_{24,20} f, \quad g <_{48,4} f, \quad \text{and} \quad g <_{48,28} f.$$

Therefore, we know that $Q(g) \subset Q(f)$ by (2.1). Hence, f and g have perfectly identical integral representations.

(iii) Quadratic forms in S_7 . Let $f(x, y, z) = 4x^2 + 8y^2 + 16z^2 + 8yz + 2xz + 2xy$ and $g(x, y, z) = 4x^2 + 4y^2 + 8z^2 - 2yz - 4xz$. One may easily show that f is a subform of g . Therefore, $Q(f) \subset Q(g)$.

It is enough to show that every positive even integer represented by g is also represented by f . One may easily show that

$$g <_{4,2} f, \quad g <_{24,0} f, \quad g <_{24,4} f, \\ g <_{24,8} f, \quad g <_{24,12} f, \quad g <_{24,16} f, \quad \text{and} \quad g <_{24,20} f.$$

Therefore, we know that $Q(g) \subset Q(f)$ by (2.1). Hence, f and g have perfectly identical integral representations.

(iv) Quadratic forms in S_8 . Let $f(x, y, z) = 10x^2 + 10y^2 + 16z^2 - 8xz - 6xy$ and $g(x, y, z) = 10x^2 + 14y^2 + 14z^2 + 12yz + 2xz + 10xy$. One may easily show that

$$g <_{4,0} f, \quad g <_{12,2} f, \quad g <_{12,6} f, \quad g <_{36,10} f, \quad g <_{36,22} f, \quad \text{and} \quad g <_{36,34} f.$$

Therefore, we know that $Q(g) \subset Q(f)$ by (2.1).

Conversely, one may easily show that

$$f \prec_{4,0} g, \quad f \prec_{12,2} g, \quad f \prec_{12,6} g, \quad f \prec_{36,10} g, \quad f \prec_{36,22} g, \quad \text{and} \quad f \prec_{36,34} g.$$

Therefore, we know that $Q(f) \subset Q(g)$ by (2.1). Hence, f and g have perfectly identical integral representations.

Remark 3.1. (1) For the other pairs in Table 1.1 not considered in this article, it seems to be inadequate to apply our method.

(2) In 1981, Hsia conjectured in [5] that two positive ternary quadratic forms in the same genus are equivalent if they primitively represent the same integers. Jagy found two candidates of pairs of ternary quadratic forms violating Hsia's conjecture. Actually, He proved that each of two quadratic forms contained in S_8 and S_9 in Table 1.1 which are contained in same genus primitively represents same integers up to 1,000,000 (see [6]).

REFERENCES

- [1] J. Cervino and G. Hein, *The Conway-Sloane tetralattice pairs are non-isometric*, Adv. Math. **228**(2011), 153-166.
- [2] J. H. Conway and N. J. A. Sloane, *Fourdimensional lattices with the Same theta series*, Int. Math. Res. Not. **4**(1992), 93-96.
- [3] B. N. Delone, *Geometry of positive quadratic forms*, Uspehi Math. Nauk. **4**(1938), 102-164.
- [4] T. Do, *Ternary quadratic forms and modular forms of half integral weight*, Master thesis, University of Auckland.
- [5] J. S. Hsia, *Regular positive ternary quadratic forms*, Mathematika **28**(1981), 231-238.
- [6] W. C. Jagy, *personal communication*, March 26, 2020.
- [7] J. Ju, *Universal mixed sums of generalized 4- and 8-gonal numbers*, to appear in Int. J. Number Theory.
- [8] Y. Kitaoka, *Arithmetic of quadratic forms*, Cambridge University Press, 1993.
- [9] Y. Kitaoka, *Positive definite quadratic forms with the same representation numbers*, Arch. Math. **28**(1977), 495-497.
- [10] M. Kneser, *Lineare Relationen zwischen Darstellungszahlen quadratischer Formen*, Math. Ann. **168**(1967), 3139.
- [11] B.-K. Oh, *Regular positive ternary quadratic forms*, Acta Arith. **147**(2011), 233-243.
- [12] B.-K. Oh, *Ternary universal sums of generalized pentagonal numbers*, J. Korean Math. Soc. **48**(2011), 837-847.
- [13] R. Oishi-Tomiyasu, *On positive-definite ternary quadratic forms with the same representations over \mathbb{Z}* , arXiv:1703.08854v11.
- [14] O. T. O'Meara, *Introduction to quadratic forms*, Springer Verlag, New York, 1963.
- [15] A. Schiemann, *Ein Beispiel positiv definiter quadratischer Formen der Dimension 4 mit gleichen Darstellungszahlen*, Arch. Math. **54**(1990), 372-375.
- [16] A. Schiemann, *Ternary positive definite quadratic forms are determined by their theta series*, Math. Ann. **308**(1997), 507-517.
- [17] E. Witt, *Eine Identität zwischen Modulformen zweiten Grades*, Abh. Math. Sem. Hansischen Univ. **14**(1941), 323-337.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ULSAN, ULSAN 44610, REPUBLIC OF KOREA
E-mail address: jangwonju@ulsan.ac.kr