

**ON SOME IMAGINARY TRIQUADRATIC NUMBER FIELDS  $k$   
WITH  $\text{Cl}_2(k) \simeq (2, 4)$  OR  $(2, 2, 2)$**

ABDELMALEK AZIZI, MOHAMED MAHMOUD CHEMS-EDDIN,  
AND ABDELKADER ZEKHNINI

ABSTRACT. Let  $d$  be a square free integer and  $L_d := \mathbb{Q}(\zeta_8, \sqrt{d})$ . In the present work we determine all the fields  $L_d$  such that the 2-class group,  $\text{Cl}_2(L_d)$ , of  $L_d$  is of type  $(2, 4)$  or  $(2, 2, 2)$ .

## 1. INTRODUCTION

Let  $k$  be a number field and  $\text{Cl}_2(k)$  its 2-class group, that is the 2-Sylow subgroup of its ideal class group  $\text{Cl}(k)$ . The problem of determining the structure of  $\text{Cl}_2(k)$  is one of the most interesting problems of algebraic number theory, accordingly many mathematicians treated this problem for some number fields of degree 2, 4. For example, in [17] using binary quadratic forms theory, Kaplan, determined the 2-class group of some quadratic number fields. The authors of [10, 11] used genus theory and class field theory to characterize those imaginary quadratic number fields,  $k$ , with 2-class group of type  $(2, 2^m)$  or  $(2, 2, 2)$  and the 2-rank of the class group of its Hilbert 2-class field equal to 2. In [5], using units and the 2-part of the class number of subextensions of  $k$ , the authors determined the 2-class group of some real biquadratic number fields  $k = \mathbb{Q}(\sqrt{m}, \sqrt{d})$  with  $d$  is an odd square free integer. Using similar techniques, the paper [7] characterizes all the fields  $k = \mathbb{Q}(i, \sqrt{d})$  such that  $\text{Cl}_2(k)$  is of type  $(2, 4)$  or  $(2, 2, 2)$  (here  $(a_1, \dots, a_r)$  denotes the direct sum of cyclic groups of order  $a_i$ , for  $i = 1, \dots, r$ ). Whenever  $k$  is an imaginary multiquadratic number field, this problem is strongly related to the units of  $k$  and the class number of the 2-part of the class numbers of its sub-extensions as we will see later. This paper is, actually, a continuation and extension of our earlier work [4], in which we determined the rank of the 2-class group of all fields of the form  $L_d := \mathbb{Q}(\zeta_8, \sqrt{d})$  with  $d$  a positive square free integer and moreover we determined all fields  $L_d$  for which the 2-class group,  $\text{Cl}_2(L_d)$ , is of type  $(2, 2)$ . In this work, we are interested in determining all positive square free integers  $d$  satisfying  $\text{Cl}_2(L_d)$  is of type  $(2, 4)$  or  $(2, 2, 2)$ .

### NOTATIONS

The next notations will be used for the rest of this article:

---

2000 *Mathematics Subject Classification.* 11R11; 11R16; 11R18; 11R27; 11R29.

*Key words and phrases.* 2-group rank; 2-class group; imaginary triquadratic number fields.

- $d$ : A positive odd square free integer,
- $L_d$ :  $\mathbb{Q}(\zeta_8, \sqrt{d})$ ,
- $\text{Cl}_2(L)$ : The 2-class group of some number field  $L$ ,
- $h(L)$ : The class number of the number field  $L$ ,
- $h_2(L)$ : The 2-class number of the number field  $L$ ,
- $h_2(m)$ : The 2-class number of the quadratic field  $\mathbb{Q}(\sqrt{m})$ ,
- $E_L$ : The unit group of any number field  $L$ ,
- $\varepsilon_m$ : The fundamental unit of  $\mathbb{Q}(\sqrt{m})$ ,
- $W_L$ : The set of roots of unity contained in a field  $L$ ,
- $\omega_L$ : The cardinality of  $W_L$ ,
- $L^+$ : The maximal real subfield of an imaginary number field  $L$ ,
- $Q_L$ : The Hasse's index, that is  $[E_L : W_L E_{L^+}]$ , if  $L/L^+$  is CM,
- $Q(L/k)$ : The unit index of a  $V_4$ -extension  $L/k$ ,
- $q(L) := [E_L : \prod_i E_{k_i}]$ , with  $k_i$  are the quadratic subfields of  $L$ .

## 2. PRELIMINARIES

Let us first collect some results and definitions that we will need in the sequel. Recall that a field  $K$  is said to be a CM-field if it is a totally complex quadratic extension of a totally real number field. Note also that a  $V_4$ -extension  $K/k$  (i.e., a normal extension of number fields with  $\text{Gal}(K/k) = V_4$ , where  $V_4$  is the Klein's four-group) is called  $V_4$ -extension of CM-fields if exactly two of its three quadratic sub-extensions are CM-fields. Let us next recall the class number formula for a  $V_4$ -extension of CM-fields:

**Proposition 1** ([20]). *Let  $L/K$  be a  $V_4$ -extension of CM-fields, then*

$$h(L) = \frac{Q_L}{Q_{K_1} Q_{K_2}} \cdot \frac{\omega_L}{\omega_{K_1} \omega_{K_2}} \cdot \frac{h(K_1)h(K_2)h(L^+)}{h(K)^2}.$$

Where  $K_1, K_2, L^+$  are the three sub-extensions of  $L/K$ , with  $K_1$  and  $K_2$  are CM-fields.

The following class number formula for a multiquadratic number field is usually attributed to Kuroda [19], but it goes back to Herglotz [15] (cf. [12, p. 27]).

**Proposition 2.** *Let  $K$  be a multiquadratic number field of degree  $2^n$ ,  $n \in \mathbb{N}$ , and  $k_i$  the  $s = 2^n - 1$  quadratic subfields of  $K$ . Then*

$$h(K) = \frac{1}{2^v} [E_K : \prod_{i=1}^s E_{k_i}] \prod_{i=1}^s h(k_i),$$

with

$$v = \begin{cases} n(2^{n-1} - 1); & \text{if } K \text{ is real,} \\ (n-1)(2^{n-2} - 1) + 2^{n-1} - 1 & \text{if } K \text{ is imaginary.} \end{cases}$$

Continue with the next formula called Kuroda's class number formula for a  $V_4$ -extension  $K/k$ .

**Proposition 3** ([21], p. 247). *Let  $K/k$  be a  $V_4$ -extension. Then we have :*

$$h(K) = \begin{cases} \frac{1}{4} \cdot Q(K/k) \cdot \prod_{i=1}^3 h(k_i) & \text{if } k = \mathbb{Q} \text{ and } K \text{ is real,} \\ \frac{1}{2} \cdot Q(K/k) \cdot \prod_{i=1}^3 h(k_i) & \text{if } k = \mathbb{Q} \text{ and } K \text{ is imaginary,} \\ \frac{1}{4} \cdot Q(K/k) \cdot \prod_{i=1}^3 h(k_i)/h(k)^2 & \text{if } k \text{ is an imaginary quadratic extension of } \mathbb{Q}. \end{cases}$$

Where  $k_i$  are the 3 subextensions of  $K/k$ .

### 3. FIELDS $L_d$ FOR WHICH $\text{Cl}_2(L_d)$ IS OF TYPE $(2, 4)$

Our goal in this section is to determine all fields  $L_d$  for which  $\text{Cl}_2(L_d) \simeq (2, 4)$ . Recall first the definition of the rational biquadratic residue symbol:

For a prime  $p \equiv 1 \pmod{4}$  and a quadratic residue  $a \pmod{p}$ ,  $\left(\frac{a}{p}\right)_4$  will denote the rational biquadratic residue symbol defined by

$$\left(\frac{a}{p}\right)_4 = \pm 1 \equiv a^{\frac{p-1}{4}} \pmod{p}.$$

Moreover, for an integer  $a \equiv 1 \pmod{8}$ , the symbol  $\left(\frac{a}{2}\right)_4$  is defined by

$$\left(\frac{a}{2}\right)_4 = 1 \text{ if } a \equiv 1 \pmod{16}, \quad \left(\frac{a}{2}\right)_4 = -1 \text{ if } a \equiv 9 \pmod{16}.$$

It turns out that  $\left(\frac{a}{2}\right)_4 = (-1)^{\frac{a-1}{8}}$ .

In all this section, let  $p, p_i, q$  and  $q_i$  be prime integers such that  $p \equiv p_i \equiv 1 \pmod{4}$  and  $q \equiv q_i \equiv 3 \pmod{4}$  with  $i \in \mathbb{N}^*$ . The following lemma is proved in our earlier paper [4, Theorem 5.6].

**Lemma 1.** *The rank of the 2-class group  $\text{Cl}_2(L_d)$  of  $L_d$  equals 2 if and only if  $d$  takes one of the following forms:*

1.  $d = q_1 q_2$  with  $q_1 \equiv q_2 \equiv 3 \pmod{8}$ .
2.  $d = p_1 p_2$  with  $p_1 \equiv p_2 \equiv 5 \pmod{8}$ .
3.  $d = q_1 q_2$  with  $q_1 \equiv 3 \pmod{8}$  and  $q_2 \equiv 7 \pmod{8}$ .
4.  $d = pq$  with  $p \equiv 5 \pmod{8}$  and  $q \equiv 7 \pmod{8}$ .
5.  $d = p$  with  $p \equiv 1 \pmod{8}$  and  $\left[ p \equiv 9 \pmod{16} \text{ or } \left(\frac{2}{p}\right)_4 \neq \left(\frac{p}{2}\right)_4 \right]$ .

We need also the following result.

**Lemma 2.** *Let  $d = p_1 p_2$ , where  $p_1, p_2$  are two rational primes such that  $p_i \equiv 5 \pmod{8}$ . Then  $h_2(L_d)$  is divisible by 16.*

*Proof.* We have  $L_d = \mathbb{Q}(\sqrt{p_1 p_2}, \sqrt{2}, i)$  is an imaginary multiquadratic number field of degree  $2^3$ . So by Proposition 2 we have

$$h(L_d) = \frac{1}{2^5} q(L_d) h(p_1 p_2) h(-p_1 p_2) h(2p_1 p_2) h(-2p_1 p_2) h(2) h(-2) h(-1).$$

It is known that  $h_2(2)$ ,  $h_2(-2)$  and  $h_2(-1)$  are equal to 1, and by [17, p. 350]  $h_2(-2p_1 p_2) = 4$ . On the other hand, [9, Theorem 2.2] implies that  $q(L_d)$  is a power of 2. So by passing to the 2-part in the above equation, we get

$$h_2(L_d) = \frac{1}{2^3} q(L_d) h_2(p_1 p_2) h_2(-p_1 p_2) h_2(2p_1 p_2). \quad (1)$$

Note that:

- By [14, Corollary 18.4],  $h_2(p_1 p_2)$  is divisible by 2.
- By [17, pp. 348-349] (Propositions  $B'_1$  and  $B'_4$ ),  $h_2(-p_1 p_2)$  is divisible by 8.
- By [14, Corollaries 18.4, 19.7 and 19.8 ],  $h_2(2p_1 p_2)$  is divisible by 4.

On one hand,  $\zeta_8 = \frac{1+i}{\sqrt{2}} \in E_{L_d}$ . On the other hand, letting  $k_i$  be the quadratic subfields of  $L_d$ , one gets easily

$$\prod_i E_{k_i} = \langle i, \varepsilon_2, \varepsilon_{-2}, \varepsilon_{p_1 p_2}, \varepsilon_{2p_1 p_2}, \varepsilon_{-p_1 p_2}, \varepsilon_{-2p_1 p_2} \rangle.$$

So the 8-th root of unity  $\zeta_8 \notin \prod_i E_{k_i}$ . Thus  $\bar{1}$  and  $\bar{\zeta}_8$  are two distinct cosets in the quotient  $E_{L_d} / \prod_i E_{k_i}$ . Thus,  $q(L_d)$  is divisible by 2. It follows by the equality (1) above that  $h_2(L_d)$  is divisible by  $\frac{1}{2^3} \cdot 2 \cdot 2 \cdot 8 \cdot 4 = 16$  as we wished to prove.  $\square$

**Example 1.** Let  $d = p_1 p_2$  be as in the above Lemma. Using PARI/GP calculator version 2.9.4 (64bit), Dec 20, 2017, for  $5 \leq p_i \leq 200$ ,  $i = 1, 2$ , we could not find a field  $L_d$  such that  $h_2(L_d) = 16$ . We have the following examples.

1. For  $p_1 = 13$  and  $p_2 = 5$ , we have  $h_2(L_{13.5}) = 32$ .
2. For  $p_1 = 37$  and  $p_2 = 53$ , we have  $h_2(L_{37.53}) = 64$ .

**Remark 1.** With hypothesis and notations of Lemma 2, we find in [9] a unit group of  $L_d$ .

**Lemma 3** ([25], Lemma 2). *Let  $d \equiv 1 \pmod{4}$  be a positive square free integer and  $\varepsilon_d = x + y\sqrt{d}$  be the fundamental unit of  $\mathbb{Q}(\sqrt{d})$ . Assume  $N(\varepsilon_d) = 1$ , then*

1.  $x + 1$  and  $x - 1$  are not squares in  $\mathbb{N}$ , i.e.,  $2\varepsilon_d$  is not a square in  $\mathbb{Q}(\sqrt{d})$ .
2. For all prime  $p$  dividing  $d$ ,  $p(x + 1)$  and  $p(x - 1)$  are not squares in  $\mathbb{N}$ .

**Lemma 4.** *Let  $d = q_1 q_2$ , with  $q_1 \equiv q_2 \equiv 3 \pmod{8}$  are two primes such that  $\begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = 1$ . Then we have  $E_{L_d} = \langle \zeta_8, \varepsilon_2, \varepsilon_{q_1 q_2}, \sqrt{\varepsilon_{2q_1 q_2}} \rangle$ , and thus  $q(L_d) = 4$ .*

*Proof.* As  $q_1 q_2 \equiv 1 \pmod{8}$ , we claim that the unit  $\varepsilon_{q_1 q_2}$  can be written as  $\varepsilon_{q_1 q_2} = a + b\sqrt{q_1 q_2}$ , where  $a$  and  $b$  are two integers. Indeed, suppose that  $\varepsilon_{q_1 q_2} = (\alpha + \beta\sqrt{d})/2$  where  $\alpha, \beta$  are two integers. Since  $N(\varepsilon_{q_1 q_2}) = 1$ , one deduces that  $\alpha^2 - 4 = \beta^2 d$ , hence  $\alpha^2 - 4 \equiv \beta^2 \pmod{8}$ . On the other hand, if

we suppose that  $\alpha$  and  $\beta$  are odd, then  $\alpha^2 \equiv \beta^2 \equiv 1 \pmod{8}$ , but this implies the contradiction  $-3 \equiv 1 \pmod{8}$ . Thus  $\alpha$  and  $\beta$  are even and our claim is established.

It is known that  $N(\varepsilon_{q_1q_2}) = 1$ , so by the unique factorization in  $\mathbb{Z}$  and Lemma 3 one gets

$$(1) : \begin{cases} a + 1 = 2q_1b_1^2 \\ a - 1 = 2q_2b_2^2 \end{cases} \quad \text{or} \quad (2) : \begin{cases} a - 1 = 2q_1b_1^2 \\ a + 1 = 2q_2b_2^2 \end{cases}$$

for some integers  $b_1$  and  $b_2$  such that  $b = 2b_1b_2$ .

If the system (2) holds, then

$$-1 = \left(\frac{2q_1}{q_2}\right) = \left(\frac{a-1}{q_2}\right) = \left(\frac{a+1-2}{q_2}\right) = \left(\frac{-2}{q_2}\right) = 1.$$

Which is absurd. Therefore  $\begin{cases} a + 1 = 2q_1b_1^2 \\ a - 1 = 2q_2b_2^2 \end{cases}$ . Thus,  $\sqrt{\varepsilon_{q_1q_2}} = b_1\sqrt{q_1} + b_2\sqrt{q_2}$ . So  $\varepsilon_d$  is not a square in  $L_d^+$ .

We have  $\varepsilon_{2d}$  has a positive norm. Put  $\varepsilon_{2d} = x + y\sqrt{2q_1q_2}$ . We similarly show that  $\sqrt{2\varepsilon_{2d}} = y_1 + y_2\sqrt{2q_1q_2} \in L_d^+$ , for some integers  $y_1$  and  $y_2$ . Therefore,  $\varepsilon_{2d}$  is a square in  $L_d^+$  since  $\sqrt{2} \in L_d^+$ . As  $\varepsilon_2$  has a negative norm, then using the algorithm described in [24] (or in [2, p. 113]) we have  $\{\varepsilon_2, \varepsilon_{q_1q_2}, \sqrt{\varepsilon_{2q_1q_2}}\}$  is a fundamental system of unit of  $L_d^+$ . Hence the result follows easily by [3, Proposition 2].  $\square$

To continue, consider the following parameters:

- For a rational prime  $p$  such that  $p \equiv 1 \pmod{8}$ , set  $p = u^2 - 2v^2$  where  $u$  and  $v$  are two positive integers such that  $u \equiv 1 \pmod{8}$  (for the existence of  $u$  and  $v$  see [22]).
- For two primes  $q_1$  and  $q_2$  such that  $q_1 \equiv q_2 \equiv 3 \pmod{8}$ ,  $\left(\frac{q_1}{q_2}\right) = 1$ , there exist five integers  $X, Y, k, l$  and  $m$  such that  $2q_2 = k^2X^2 + 2lXY + 2mY^2$  and  $q_1 = l^2 - 2k^2m$  (cf. [17, p. 356]).

**Theorem 1.** *Let  $d$  be an odd positive square free integer. Then  $\text{Cl}_2(L_d) \simeq (2, 4)$  if and only if  $d$  takes one of the two following forms:*

1.  $d = p \equiv 9 \pmod{16}$  is a prime such that  $\left(\frac{2}{p}\right)_4 \neq \left(\frac{p}{2}\right)_4$  and  $\left(\frac{u}{p}\right)_4 = -1$ .
2.  $d = q_1q_2$ , with  $q_1 \equiv q_2 \equiv 3 \pmod{8}$  are primes such that  $\left(\frac{q_1}{q_2}\right) = 1$  and

$$\left(\frac{-2}{|k^2X + lY|}\right) = -1.$$

*Proof.* It suffices to determine for which forms of  $d$  appearing in Lemma 1, we have  $h_2(L_d) = 8$ . Let us firstly eliminate some cases.

- By [4, Propositions 5.13 and 5.14],  $h_2(L_d) \neq 8$  whenever  $d$  takes forms in the third and the fourth item of Lemma 1.

- The form of  $d$  in the second item of Lemma 1 is also eliminated by Lemma 2.
- Note that if  $p \equiv 1 \pmod{16}$  and  $\left(\frac{2}{p}\right)_4 \neq \left(\frac{p}{2}\right)_4$ , then by [4, Theorem 5.7],  $h_2(L_p) \neq 8$ .

Note that by results in the beginning of Section 3,  $p \equiv 9 \pmod{16}$  implies that  $\left(\frac{p}{2}\right)_4 = -1$ . Hence, by Lemma 1, we have to check the following cases.

- (I)  $d = p$ , is a prime such that  $\left(\frac{2}{p}\right)_4 = \left(\frac{p}{2}\right)_4 = -1$ .
- (II)  $d = p$ , is a prime such that  $p \equiv 9 \pmod{16}$  and  $\left(\frac{2}{p}\right)_4 \neq \left(\frac{p}{2}\right)_4$ .
- (III)  $d = q_1q_2$ , for two primes  $q_1$  and  $q_2$  such that  $q_1 \equiv q_2 \equiv 3 \pmod{8}$ .

Let  $p \equiv 1 \pmod{8}$  be a prime. Set  $L_p^+ = \mathbb{Q}(\sqrt{2}, \sqrt{p})$ ,  $K = \mathbb{Q}(\sqrt{2}, i)$  and  $K' = \mathbb{Q}(\sqrt{2}, \sqrt{-p})$ . By applying Proposition 1 to the extension  $L_p/\mathbb{Q}(\sqrt{2})$ , we have

$$h(L_p) = \frac{Q_{L_p} \omega_{L_p} h(L_p^+)h(K)h(K')}{Q_K Q_{K'} \omega_K \omega_{K'} h(\mathbb{Q}(\sqrt{2}))^2}.$$

We have  $h(\mathbb{Q}(\sqrt{2})) = h(K) = 1$ . By [6, Théorème 3],  $Q_{L_p} = 1$  and by [4, Lemma 2.5]  $Q_K = 1$ . Since  $\omega_{L_p} = \omega_K = 8$  and  $\omega_{K'} = 2$ , then by passing to the 2-part in the above equality we get

$$h_2(L_p) = \frac{1}{2Q_{K'}} h_2(L_p^+) h_2(K'). \quad (2)$$

As  $\varepsilon_2$  has a negative norm, so by the item (2) of Section 3 of [3] we obtain that  $E_{K'} = \langle -1, \varepsilon_2 \rangle$ . This in turn implies that  $q(K') = Q_{K'} = 1$ . From which we infer, by Proposition 2, that  $h_2(K') = \frac{1}{2} \cdot 1 \cdot h_2(2)h_2(-p)h_2(-2p) = \frac{1}{2}h_2(-p)h_2(-2p)$ . It follows, by the equality (2), that

$$h_2(L_p) = \frac{1}{4} h_2(L_p^+) h_2(-p) h_2(-2p). \quad (3)$$

Note that from [13, Theorem 2] and the proof of [13, Theorem 1], one deduces easily that  $h_2(-p) = 4$  if and only if  $\left(\frac{2}{p}\right)_4 \neq \left(\frac{p}{2}\right)_4$ .

- Suppose that  $d$  takes the form (I). So 8 divides  $h_2(-p)$ . Note also that, by [16, p. 596],  $h_2(-2p)$  is divisible by 4, and by [18, Theorem 2]  $h_2(L_p^+)$  is even. It follows by the equality (3) that 16 divides  $h_2(L_p)$ . So this case is eliminated

- Suppose that  $d$  takes the form (II). Thus  $\left(\frac{2}{p}\right)_4 = 1$  and  $\left(\frac{p}{2}\right)_4 = -1$ . So, by [18, Theorem 2],  $h_2(L_p^+) = 1$ . Therefore, by the equality (3) and the note below it, we have  $h_2(L_p) = h_2(-2p)$ .  
Keep the notations of [16, p. 601] and let  $p = a^2 + b^2 = 2e^2 - d^2$ , with  $e \geq 1$ . By using notations in the proof of [13, Theorem 1] and [16, p. 601] we easily deduce that  $1 = \left(\frac{2}{p}\right)_4 = (-1)^{b/4}$ , so  $b \equiv 0 \pmod{8}$ . Therefore, by [16, Théorème 3],  $h_2(-2p) \equiv 0 \pmod{8}$ . It follows, by [22, Theorem 2],  $h_2(-2p) = 8$  if and only if  $\left(\frac{u}{p}\right)_4 = -1$ . So the first item of our theorem.
- Suppose that  $d$  takes the third form (III). Without loss of generality we may assume that  $\left(\frac{q_1}{q_2}\right) = 1$ . By Proposition 2 we have

$$h(L_d) = \frac{1}{2^5} q(L_d) h(q_1 q_2) h(-q_1 q_2) h(2q_1 q_2) h(-2q_1 q_2) h(2) h(-2) h(-1).$$

By [14, Corollary 18.4]  $h_2(q_1 q_2) = 1$  and by [17, pp. 345, 354] we have  $h_2(2q_1 q_2) = 2$  and  $h_2(-q_1 q_2) = 4$  respectively. It is known that  $h(2) = h(-2) = h(-1) = 1$ . Since by Lemma 4  $q(L_d) = 4$ , then by passing to the 2-part in the above equality we get

$$h_2(L_d) = \frac{1}{2^5} \cdot 4 \cdot 1 \cdot 4 \cdot 2 \cdot h_2(-2q_1 q_2) = h_2(-2q_1 q_2).$$

Therefore, by [17, p. 357] and Lemma 1,  $\text{Cl}_2(L_d) = (2, 4)$  if and only if  $h_2(-2q_1 q_2) = 8$  which is equivalent to  $\left(\frac{-2}{|k^2 X + lY|}\right) = -1$ . This achieves the proof.  $\square$

Let  $\ell$  be a positive integer. For a finite abelian group  $G$ , its  $2^\ell$ -rank is defined as  $r_{2^\ell}(G) = \dim_{\mathbb{F}_2}(2^{\ell-1}G/2^\ell G)$ . Or equivalently looking at the decomposition of the group  $G$  into cyclic groups as  $G = \prod_i C_{n_i}$ , the  $2^\ell$ -rank of  $G$  equals the number of  $n_i$ 's divisible by  $2^\ell$ .

**Corollary 1.** *Let  $p$  be a rational prime such that  $p \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p}\right)_4 = \left(\frac{p}{2}\right)_4 = -1$ , then the 8-rank of  $\text{Cl}_2(L_p)$  equals 1.*

*Proof.* By [6, Théorème 10], we deduce that the 4-rank of  $\text{Cl}_2(L_p)$  equals 1, and by the proof of Theorem 1,  $h_2(L_p)$  is divisible by 16, so the result.  $\square$

From the proof of Theorem 1 we get

**Corollary 2.** *Let  $q_1$  and  $q_2$  be two rational primes such that  $q_1 \equiv q_2 \equiv 3 \pmod{8}$ , then  $h_2(L_{q_1q_2}) = h_2(-2q_1q_2)$ .*

**Example 2.** For all the examples below, we used PARI/GP calculator version 2.9.4 (64bit), Dec 20, 2017.

1. For  $p = 89$ ,  $u = 17$  and  $v = 10$ , we have  $p = u^2 - 2v^2$ ,  $\left(\frac{2}{p}\right)_4 = -\left(\frac{u}{p}\right)_4 = 1$  and  $\text{Cl}_2(L_{89}) \simeq (2, 4)$ .
2. For  $q_1 = 11$ ,  $q_2 = 19$ ,  $k = 1$ ,  $l = 3$ ,  $m = -1$ ,  $X = 4$  and  $Y = 1$ , we have  $q_1 = l^2 - 2k^2m$ ,  $2q_2 = k^2X^2 + 2lXY + 2mY^2$ ,  $\left(\frac{-2}{|k^2X + lY|}\right) = \left(\frac{-2}{7}\right) = -1$  and  $\text{Cl}_2(L_{11.19}) \simeq (2, 4)$ .

#### 4. FIELDS $L_d$ FOR WHICH $\text{Cl}_2(L_d)$ IS OF TYPE $(2, 2, 2)$

In this section we determine all fields  $L_d$  such that  $\text{Cl}_2(L_d) \simeq (2, 2, 2)$ . Keep the above notations:  $p, p_i, q$  and  $q_i$  are prime integers satisfying  $p \equiv p_i \equiv 1 \pmod{4}$  and  $q \equiv q_i \equiv 3 \pmod{4}$  with  $i \in \mathbb{N}^*$ . From section 4 of [4], it is easy to deduce the following result:

**Lemma 5.** *The rank of the 2-class group  $\text{Cl}_2(L_d)$  of  $L_d$  equals 3 if and only if  $d$  takes one of the following forms.*

1.  $d = p$  with  $p \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p}\right)_4 = \left(\frac{p}{2}\right)_4 = 1$ .
2.  $d = q_1q_2$  with  $q_1 \equiv q_2 \equiv 7 \pmod{8}$ .
3.  $d = qp$  with  $q \equiv 3 \pmod{8}$  and  $p \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p}\right)_4 = -1$ .
4.  $d = p_1p_2$  with  $p_1 \equiv 5 \pmod{8}$ ,  $p_2 \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p_2}\right)_4 \neq \left(\frac{p_2}{2}\right)_4$ .
5.  $d = q_1q_2p$  with  $q_1 \equiv q_2 \equiv 3 \pmod{8}$  and  $p \equiv 5 \pmod{8}$ .
6.  $d = qp_1p_2$  with  $q \equiv 3 \pmod{8}$  and  $p_1 \equiv p_2 \equiv 5 \pmod{8}$ .

We will need the following lemmas.

**Lemma 6.** *Let  $d = q_1q_2$ , with  $q_1 \equiv q_2 \equiv 7 \pmod{8}$  are two primes such that  $\left(\frac{q_1}{q_2}\right) = 1$ . Then we have  $E_{L_d} = \langle \zeta_8, \varepsilon_2, \varepsilon_{q_1q_2}, \sqrt{\varepsilon_{2q_1q_2}}$  or  $\sqrt{\varepsilon_{q_1q_2}\varepsilon_{2q_1q_2}} \rangle$ , and thus  $q(L_d) = 4$ .*

*Proof.* Similar to the proof of Lemma 4. □

**Lemma 7.** 1. *Let  $d = p$ , for a prime  $p \equiv 1 \pmod{8}$  such that  $\left(\frac{2}{p}\right)_4 = \left(\frac{p}{2}\right)_4 = 1$ .*

*Then,  $h_2(L_p) \equiv 0 \pmod{16}$ , and thus  $\text{Cl}_2(L_p)$  is not elementary.*

2. *Let  $d = q_1q_2$  with  $q_1$  and  $q_2$  are two rational primes such that  $q_1 \equiv q_2 \equiv 7 \pmod{8}$ . Then,  $h_2(L_{q_1q_2}) \equiv 0 \pmod{32}$ , and thus  $\text{Cl}_2(L_{q_1q_2})$  is not elementary.*



3. Let  $d = p_1 p_2$  with  $p_1 \equiv 5 \pmod{8}$ ,  $p_2 \equiv 1 \pmod{8}$  are two primes such that  $\left(\frac{p_2}{p_1}\right) = 1$ . Then,  $h_2(L_{p_1 p_2}) \equiv 0 \pmod{32}$ .

*Proof.* 1. By equality (3) in the proof of Theorem 1 we have:

$$h_2(L_p) = \frac{1}{4} h_2(L_p^+) h_2(-p) h_2(-2p).$$

Note that by [16, p. 596],  $h_2(-2p)$  is divisible by 4. Since for a prime  $p' \equiv 1 \pmod{8}$ ,  $h_2(-p') = 4$  if and only if  $\left(\frac{2}{p'}\right) \neq \left(\frac{p'}{2}\right)$  (see the proof of Theorem 1), then  $h_2(-p)$  is divisible by 8 (in fact  $\left(\frac{2}{p}\right) = \left(\frac{p}{2}\right)$  and by [14, Corollaries 18.4 and 19.6]  $h_2(-p)$  is divisible by 4). Thus,  $h_2(L_p)$  is divisible by  $8 \cdot h_2(L_p^+)$ . By [18, Theorem 2],  $h_2(L_p^+)$  is even. From which we infer that  $h_2(L_p)$  is divisible by 16. So we have the first item.

2. By Proposition 2 we have

$$h(L_{q_1 q_2}) = \frac{1}{2^5} q(L_{q_1 q_2}) h(q_1 q_2) h(-q_1 q_2) h(2q_1 q_2) h(-2q_1 q_2) h(2) h(-2) h(-1).$$

On one hand, by [17, p. 345],  $h_2(2q_1 q_2)$  is divisible by 4, by [17, pp. 354, 356],  $h_2(-q_1 q_2)$  and  $h_2(-2q_1 q_2)$  are both divisible by 8, and, by [14, Corollary 18.4],  $h_2(q_1 q_2) = 1$ . On the other hand, by Lemma 6,  $q(L_{q_1 q_2}) = 4$ . From all these results, it follows by passing to the 2-part in the above equality that  $h_2(L_d)$  is divisible by  $\frac{1}{2^5} \cdot 4 \cdot 8 \cdot 4 \cdot 8 = 32$ . And the second item follows.


3. We proceed as in the proof the second item. □

**Example 3.** Let  $d = p$  be as in the first item of the above Lemma. Using PARI/GP calculator version 2.9.4 (64bit), Dec 20, 2017, for  $3 \leq p \leq 10^4$ , we could not find a field  $L_d$  such that  $h_2(L_d) = 16$ . We have the following examples.

1. For  $p = 113$ , we have  $\left(\frac{2}{p}\right) = \left(\frac{p}{2}\right) = 1$  and  $h_2(L_{113}) = 64$ .
2. For  $p = 337$ , we have  $\left(\frac{2}{p}\right) = \left(\frac{p}{2}\right) = 1$  and  $h_2(L_{337}) = 32$ .
3. For  $q_1 = 7$  and  $q_2 = 31$ , we have  $h_2(L_{7.31}) = 32$ .
4. For  $q_1 = 7$  and  $q_2 = 23$ , we have  $h_2(L_{7.23}) = 64$ .

**Lemma 8.** 1. If  $d = q_1 q_2 p$  with  $q_1 \equiv q_2 \equiv 3 \pmod{8}$  and  $p \equiv 5 \pmod{8}$ , then  $h_2(L_{q_1 q_2 p}) \equiv 0 \pmod{32}$ . So  $\text{Cl}_2(L_{q_1 q_2 p})$  is not elementary.  
 2. If  $d = q p_1 p_2$  with  $q \equiv 3 \pmod{8}$  and  $p_1 \equiv p_2 \equiv 5 \pmod{8}$ , then  $h_2(L_{q p_1 p_2}) \equiv 0 \pmod{32}$ . So  $\text{Cl}_2(L_{q p_1 p_2})$  is not elementary.

*Proof.* 1. Consider the following diagram (Figure 1 below):

FIGURE 1.  $L_{q_1 q_2 p}/\mathbb{Q}(i)$ .

By applying Proposition 3 to the  $V_4$ -extension  $L_{q_1 q_2 p}/\mathbb{Q}(i)$  we get:

$$h(L_d) = \frac{1}{4} Q(L_d/\mathbb{Q}(i)) h(K_1) h(K_2) h(K_3) / h(\mathbb{Q}(i))^2.$$

It is known that  $h(K_1) = h(\mathbb{Q}(i)) = 1$  (in fact,  $K_1 = \mathbb{Q}(\zeta_8)$ ), so

$$h(L_d) = \frac{1}{4} Q(L_d/\mathbb{Q}(i)) h(K_2) h(K_3). \quad (4)$$

Note that by [23, Proposition 2], the ranks of the 2-class groups of  $K_2$  and  $K_3$  equal 2 and 3 respectively. The author of [1] determined all fields  $\mathbb{Q}(\sqrt{d}, i)$  for which the 2-class group is of type  $(2, 2)$ . By checking all the results of this last reference, we deduce that 2-class group of  $K_2$  is not of type  $(2, 2)$ . So 8 divides the class number of  $K_2$ .

On the other hand, by [7, Théorème 5.3] the 2-class group of  $K_3$  is not of type  $(2, 2, 2)$ . So the class number of  $K_2$  is divisible by 16. Hence by equality (4)  $h(L_d)$  is divisible by  $\frac{1}{4} \cdot 8 \cdot 16 = 32$ . So the first item.

2. We similarly prove the second item by using [7, Théorème 5.3] and [23, Proposition 2]. □

**Example 4.** Let  $d = qp_1 p_2$  be as in the second item of the above Lemma. Using PARI/GP calculator version 2.9.4 (64bit), Dec 20, 2017, for  $3 \leq q, p_i \leq 100$ , we could not find a field  $L_d$  such that  $h_2(L_d) = 32$ . We have the following examples.

1. For  $q_1 = 3, q_2 = 11$  and  $p = 5$ , we have  $h_2(L_{3.11.5}) = 32$ .
2. For  $q_1 = 3, q_2 = 11$  and  $p = 13$ , we have  $h_2(L_{3.11.13}) = 128$ .
3. For  $q = 3, p_1 = 5$  and  $p_2 = 13$ , we have  $h_2(L_{3.5.13}) = 64$ .
4. For  $q = 3, p_1 = 5$  and  $p_2 = 29$ , we have  $h_2(L_{3.5.29}) = 128$ .

**Lemma 9.** Let  $d = qp$  with  $q \equiv 3 \pmod{8}$ ,  $p \equiv 1 \pmod{8}$  and  $\left(\frac{p}{q}\right) = -1$ . Then

$E_{L_d} = \langle \zeta_8, \varepsilon_2, \sqrt{\varepsilon_{pq}}, \sqrt{\varepsilon_{2pq}} \rangle$ , and thus  $q(L_d) = 8$ .

*Proof.* We proceed as in the proof of Lemma 4. □

Now we are able to state the main theorem of this subsection.

**Theorem 2.** *Let  $d$  be a square free integer, then  $\text{Cl}_2(L_d) \simeq (2, 2, 2)$  if and only if  $d$  takes one of the two following forms:*

1.  $d = p_1 p_2$  with  $p_1 \equiv 5 \pmod{8}$ ,  $p_2 \equiv 1 \pmod{8}$ ,  $\left(\frac{2}{p_2}\right)_4 \neq \left(\frac{p_2}{2}\right)_4$  and  $\left(\frac{p_2}{p_1}\right) = -1$ .
2.  $d = qp$  with  $q \equiv 3 \pmod{8}$ ,  $p \equiv 1 \pmod{8}$ ,  $\left(\frac{2}{p}\right)_4 = -1$  and  $\left(\frac{p}{q}\right) = -1$ .

*Proof.* It suffices to determine which forms of  $d$  appearing in Lemma 5, we have  $h_2(L_d) = 8$ . Let us start, as above, by eliminating certain inconvenient cases.

- The forms of  $d$  in the first and the second items of Lemma 5 are eliminated by Lemma 7.
- The forms of  $d$  in the two last items of Lemma 5 are eliminated by Lemma 8.

It follows that it suffices to check the two following cases:

- (I)  $d = p_1 p_2$  with  $p_1 \equiv 5 \pmod{8}$ ,  $p_2 \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p_2}\right)_4 \neq \left(\frac{p_2}{2}\right)_4$ .
- (II)  $d = qp$  with  $q \equiv 3 \pmod{8}$  and  $p \equiv 1 \pmod{8}$  and  $\left(\frac{2}{p}\right)_4 = -1$ .

- Suppose that  $d$  takes the form (I), then [8, Theorem 4.2] and the last assertion of Lemma 7 give the first item.
- Suppose now that  $d$  takes the form (II). Consider the following diagram (Figure 2 below):




FIGURE 2.  $L_d/\mathbb{Q}(\sqrt{2})$ .

By Proposition 1 we have:

$$h(L_d) = \frac{Q_{L_d} \omega_{L_d} h(L_d^+) h(K_1) h(K_2)}{Q_{K_1} Q_{K_2} \omega_{K_1} \omega_{K_2} h(\mathbb{Q}(\sqrt{2}))^2}. \quad (5)$$

Note that  $h(K_2) = h(\mathbb{Q}(\sqrt{2})) = 1$ . We have,  $d\varepsilon_2$  is not a square in  $\mathbb{Q}(\sqrt{2})$  since otherwise we will get, for some  $\alpha$  in  $\mathbb{Q}(\sqrt{2})$ ,  $d\varepsilon_2 = \alpha^2$ , then  $N_{\mathbb{Q}(\sqrt{2})/\mathbb{Q}}(d\varepsilon_2) = -d^2 = N_{\mathbb{Q}(\sqrt{2})/\mathbb{Q}}(\alpha)^2$ . Which is false. It follows by [3, Proposition 3], that  $\{\varepsilon_2\}$

is a fundamental system of units of  $K_1$ , i.e.,  $Q_{K_1} = 1$ . Since  $Q_{K_2} = 1$  (cf. [4, Lemma 2.5]),  $\omega_{L_d} = \omega_{K_2} = 8$  and  $\omega_{K_1} = 2$ , then by passing to the 2-part in the equality (5), we get  $h_2(L_d) = \frac{Q_{L_d}}{2} h_2(L_d^+) h_2(K_1)$ . By Proposition 3, we have  $h_2(K_1) = \frac{1}{2} h_2(-2d) h_2(-d) h_2(2)$ . So

$$h_2(L_d) = \frac{Q_{L_d}}{4} h_2(L_d^+) h_2(-2d) h_2(-d). \quad (6)$$

Note that, by Lemma 9, if  $\left(\frac{p}{q}\right) = -1$ , then  $Q_{L_d} = 1$ . Note also that,

- (1) from [5, Théorème 2] and its proof, we deduce that  $h_2(L_d^+)$  is divisible by 4 and  $h_2(L_d^+) = 4$  if and only if  $\left(\frac{p}{q}\right) = -1$ ,
- (2) from [14, Corollaries 18.5 and 19.6], we deduce that  $h_2(-d)$  is even and  $h_2(-d) = 2$  if and only if  $\left(\frac{p}{q}\right) = -1$ ,
- (3) from [14, Corollaries 18.5 and 19.6] and [17, p. 353], we deduce that  $h_2(-2d)$  is divisible by 4 if and only if  $\left(\frac{p}{q}\right) = -1$ .

Hence plugging all of these results into equality (6), one gets the second item. Which completes the proof.  $\square$

**Example 5.** For all the examples below, we used PARI/GP calculator version 2.9.4 (64bit), Dec 20, 2017.

1. Let  $p_1 = 29$  and  $p_2 = 17$ . We have  $p_2 \equiv 1 \pmod{8}$ ,  $\left(\frac{2}{p_2}\right)_4 \neq \left(\frac{p_2}{2}\right)_4$ ,  $\left(\frac{p_2}{p_1}\right) = -1$  and  $\text{Cl}_2(L_{29.17}) \simeq (2, 2, 2)$ .
2. Let  $q = 11$  and  $p = 17$ . We have  $\left(\frac{2}{p}\right)_4 = -1$ ,  $\left(\frac{p}{q}\right) = -1$  and  $\text{Cl}_2(L_{11.17}) \simeq (2, 2, 2)$ .

#### ACKNOWLEDGMENT

We would like to thank the unknown referee for his/her several helpful suggestions that helped us to improve our paper, and for calling our attention to the missing details. Many thanks are also due to Professor Elliot Benjamin for his important remarks on this paper.

#### REFERENCES

- [1] Azizi A., *Sur le 2-groupe de classes d'idéaux de  $\mathbb{Q}(\sqrt{d}, i)$* , Rend. Circ. Mat. Palermo, **48** (1999), 71–92.
- [2] Azizi A., *Sur les unités de certains corps de nombres de degré 8 sur  $\mathbb{Q}$* , Ann. Sci. Math. Québec, **29** (2005), 111–129.
- [3] Azizi A., *Unités de certains corps de nombres imaginaires et abéliens sur  $\mathbb{Q}$* , Ann. Sci. Math. Québec, **23** (1999), 15–21.

- [4] Azizi A., Chems-Eddin M.M. and Zekhnini A., *On the rank of the 2-class group of some imaginary triquadratic number fields*. [arXiv:1905.01225v3](https://arxiv.org/abs/1905.01225v3).
- [5] Azizi A. et Mouhib A., *Capitulation des 2-classes d'idéaux de  $\mathbb{Q}(\sqrt{2}, \sqrt{d})$  où  $d$  est un entier naturel sans facteurs carrés*, Acta. Arith., **109** (2003), 27-63.
- [6] Azizi A. et Taous M., *Capitulation des 2-classes d'idéaux de  $k = \mathbb{Q}(\sqrt{2p}, i)$* , Acta. Arith., **131** (2008), 103-123.
- [7] Azizi A. et Taous M., *Déterminations des corps  $k = \mathbb{Q}(\sqrt{d}, \sqrt{-1})$  dont les 2-groupes de classes sont de type  $(2, 4)$  ou  $(2, 2, 2)$* , Rend. Istit. Mat. Univ. Trieste, **40** (2008), 93-116.
- [8] Azizi A., Zekhnini A. and Taous M., *On the unit index of some real biquadratic number fields*, Turk. J. Math., **42** (2018), 703-715.
- [9] Azizi A., Zekhnini A. and Taous M., *On the strongly ambiguous classes of  $\mathbb{k}/\mathbb{Q}$  where  $\mathbb{k} = \mathbb{Q}(\sqrt{2p_1p_2}, i)$* , Asian-Eur. J. Math., **7** (2014), 1-26.
- [10] Benjamin E., Lemmermeyer F. and Snyder C., *Imaginary quadratic fields  $k$  with  $\text{Cl}_2(k) \simeq (2, 2^m)$  and rank  $\text{Cl}_2(k^1) = 2$* , Paci. J. Math., **198** (2001), 15-31.
- [11] Benjamin E., Lemmermeyer F. and Snyder C., *Imaginary quadratic fields  $k$  with  $\text{Cl}_2(k) \simeq (2, 2, 2)$* , J. Number Theory, **103** (2003), 38-70.
- [12] Benjamin E., Lemmermeyer F. and Snyder C., *On the unit group of some multiquadratic number fields*, Paci. J. Math., **230** (2007), 27-40.
- [13] Brown E., *The class number of  $\mathbb{Q}(\sqrt{-p})$  for  $p \equiv 1 \pmod{8}$  a prime*, Proc. Amer. Math. Soc., **31** (1972), 381-383.
- [14] Conner P. E. and Hurrelbrink J., *Class number parity*, Ser. Pure. Math., Vol 8, World Scientific, 1988.
- [15] Herglotz G., *Über einen Dirichletschen Satz*, Math. Z., **12** (1922), 255-261.
- [16] Kaplan P., *Divisibilité par 8 du nombre des classes des corps quadratiques dont le 2-groupe des classes est cyclique, et réciprocité biquadratique*, J. Math. Soc. Japan, **25** (1973), 596-608.
- [17] Kaplan P., *Sur le 2-groupe des classes d'idéaux des corps quadratiques*, J. Reine Angew. Math., **1** (1976), 313-363.
- [18] Kučera R., *On the parity of the class number of biquadratic field*, J. Number theory, **52** (1995), 43-52.
- [19] Kuroda S., *Über die Klassenzahlen algebraischer Zahlkörper*, Nagoya Math. J., **1** (1950), 1-10.
- [20] Lemmermeyer F., *Ideal class groups of cyclotomic number fields I*, Acta. Arith., **72** (1995), 347-359.
- [21] Lemmermeyer F., *Kuroda's class number formula*, Acta. Arith., **66** (1994), 245-260.
- [22] Leonard P. A. and Williams K.S., *On the divisibility of the class numbers of  $\mathbb{Q}(\sqrt{-d})$  and  $\mathbb{Q}(\sqrt{-2d})$  by 16*, Can. Math. Bull., **25** (1982), 200-206.
- [23] McCall T.M., Parry C.J. and Ranalli R.R., *Imaginary bicyclic biquadratic fields with cyclic 2-class group*, J. Number Theory, **53** (1995), 88-99.
- [24] Wada H., *On the class number and the unit group of certain algebraic number fields*, J. Fac. Sci. Univ. Tokyo, **13** (1966), 201-209.
- [25] Zekhnini A., Azizi A. and Taous M., *On the generators of the 2-class group of the field  $\mathbb{Q}(\sqrt{q_1q_2p}, i)$  Correction to Theorem 3 of [5]*, IJPAM., **103** (2015), 99-107.

ABDELMALEK AZIZI: MOHAMMED FIRST UNIVERSITY, MATHEMATICS DEPARTMENT, SCIENCES FACULTY, OUJDA, MOROCCO

*Email address:* `abdelmalekazizi@yahoo.fr`

MOHAMED MAHMOUD CHEMS-EDDIN: MOHAMMED FIRST UNIVERSITY, MATHEMATICS DEPARTMENT, SCIENCES FACULTY, OUJDA, MOROCCO

*Email address:* `2m.chemseddin@gmail.com`

ABDELKADER ZEKHNINI: MOHAMMED FIRST UNIVERSITY, MATHEMATICS DEPARTMENT, PLURIDISCIPLINARY FACULTY, NADOR, MOROCCO

*Email address:* `zekha1@yahoo.fr`