

Compatible rewriting of noncommutative polynomials for proving operator identities

Cyrille Chenavier, Clemens Hofstadler,
Clemens G. Raab, and Georg Regensburger*

Abstract

The goal of this paper is to prove operator identities using equalities between noncommutative polynomials. In general, a polynomial expression is not valid in terms of operators, since it may not be compatible with domains and codomains of the corresponding operators. Recently, some of the authors introduced a framework based on labelled quivers to rigorously translate polynomial identities to operator identities. In the present paper, we extend and adapt the framework to the context of rewriting and polynomial reduction. We give a sufficient condition on the polynomials used for rewriting to ensure that standard polynomial reduction automatically respects domains and codomains of operators. Finally, we adapt the noncommutative Buchberger procedure to compute additional compatible polynomials for rewriting. In the package `OperatorGB`, we also provide an implementation of the concepts developed.

Keywords Rewriting, noncommutative polynomials, quiver representations, automated proofs, completion

1 Introduction

Properties of linear operators can often be expressed in terms of identities they satisfy. Algebraically, these identities can be represented in terms of noncommutative polynomials in some set X . The elements of X correspond to basic operators and polynomial multiplication models composition of operators. Based on this, proving that a claimed operator identity follows from assumed identities corresponds to the polynomial f , associated to the claim, lying in the ideal generated by the set F of polynomials associated to the assumptions. However, ideal membership $f \in (F)$ is not enough for proving an operator identity in general, since computations with noncommutative polynomials ignore compatibility conditions between domains and codomains of the operators.

In order to represent domains and codomains of the operators, we use the framework introduced recently in [17]. So, we consider a quiver (i.e., a directed

*This work was supported by the Austrian Science Fund (FWF): P 27229, P 32301, and P 32952

{cyrille.chenavier,clemens.hofstadler,clemens.raab,georg.regensburger}@jku.at
Institute for Algebra, Johannes Kepler University, Altenberger Straße 69, Linz, Austria

multigraph) Q , where vertices correspond to functional spaces, edges correspond to basic operators between those spaces and are labelled with symbols from X . Then, paths in Q correspond to composition of basic operators and induce monomials over X that are compatible with Q . Note that we can allow the same label for different edges if the corresponding operators satisfy the same identities in F . For instance, differential and integral operators can act on different functional spaces, as illustrated in our running example below. For formal details and relevant notions, see Sections 2 and 4. Informally, a polynomial is compatible with the quiver if it makes sense in terms of operators and f is called a Q -consequence of F if it can be obtained from F by doing computations using compatible polynomials only. This means that these computations also make sense in terms of operators.

Obviously, the claim f and the assumptions F have to be compatible with Q . In [17], it was shown that f is a Q -consequence of F if $f \in (F)$ and each element of F is uniformly compatible, which means that all its monomials can be assigned the same combinations of domains and codomains. This is in particular the case when each edge has a unique label and polynomials do not have a constant term. Note that ideal membership can be checked independently of Q and is undecidable in general. In practice, it can often be checked by computing a (partial) noncommutative Gröbner basis G of F and reducing f to zero by G , see [15]. The package `OperatorGB` [11] can check compatibility of polynomials with quivers and, based on partial Gröbner bases, can compute explicit representations of polynomials in terms of generators of the ideal. Versions for `MATHEMATICA` and `SAGEMATH` can be obtained at:

<http://gregensburger.com/softw/OperatorGB>

In this paper, we generalize the formal definition of Q -consequences to the case when elements of F are compatible but not necessarily uniformly compatible, see Section 3. Then, we show in Section 4 that being a Q -consequence implies that the corresponding operator identity can indeed be proven by computations with operators. Since elements of F do not have to be uniformly compatible, we impose in Section 5 restrictions on the polynomial rewriting, so that it respects the quiver. For the same reason, we also impose restrictions on the computation of partial Gröbner bases in Section 6. Based on such a partial Gröbner basis, one often can prove algorithmically that f is a Q -consequence of F just by standard polynomial reduction. To this end, we also extend the package `OperatorGB`.

Gröbner bases for noncommutative polynomials have been applied to operator identities in the pioneering work [10, 9], where Gröbner bases are used to simplify matrix identities in linear systems theory. In [8, 13], the main strategy for solving matrix equations, coming from factorization of engineering systems and matrix completion problems, is to apply Gröbner bases with respect to an ordering appropriate for elimination. The same approach was used in [18] to compute Green's operators for linear two-point boundary problems with constant coefficients.

If edges of the quiver have unique labels, it has been observed in the literature that the operations used in the noncommutative analog of Buchberger's algorithm respect compatibility of polynomials with domains and codomains of operators, cf. [9, Thm. 25]. See also Remark 6 and Theorem 5 for a formal

statement using the framework of the present paper. For an analogous observation in the context of path algebras, see [16, Sec. 47.10]. We were informed in personal communication that questions related to proving operator identities via computations of Gröbner bases are also addressed in [14].

Alternatively, computations with operators can also be modelled by partial algebras arising from diagrams, for which an analogous notion of Gröbner bases was sketched in [1, Sec. 9] and developed in [3]. Moreover, generalizations of Gröbner bases and syzygies are considered in [7], where higher-dimensional linear rewriting systems are introduced for rewriting of operators with domains and codomains.

We conclude this section with a small running example that we use throughout the paper to illustrate the notions that we introduce from practical point of view. A MATHEMATICA notebook that illustrates the use of the new functionality of the package using this running example can be obtained at the webpage mentioned above.

Example 1. Consider the inhomogeneous linear differential equation

$$y''(x) + A_1(x)y'(x) + A_0(x)y(x) = r(x)$$

and assume that it can be factored into the two first-order equations

$$y'(x) - B_2(x)y(x) = z(x) \quad \text{and} \quad z'(x) - B_1(x)z(x) = r(x).$$

It is well-known that a particular solution is given by the nested integral

$$y(x) = H_2(x) \int_{x_2}^x H_2(t)^{-1} H_1(t) \int_{x_1}^t H_1(u)^{-1} r(u) du dt, \quad (1)$$

where $H_i(x)$ is a solution of $y'(x) - B_i(x)y(x) = 0$ such that $H_i(x)^{-1}$ exists. In order to translate this claim into an operator identity, let us consider the differentiation $\partial : y(x) \mapsto y'(x)$ and the two integrations

$$\int_1 : y(x) \mapsto \int_{x_1}^x y(t) dt \quad \text{and} \quad \int_2 : y(x) \mapsto \int_{x_2}^x y(t) dt.$$

Moreover, any function $F(x)$ induces a multiplication operator $F : y(x) \mapsto y(x)F(x)$ and \cdot denotes the composition of operators. Thus, the factored differential equation and the solution correspond to the following operators

$$L := (\partial - B_1) \cdot (\partial - B_2), \quad S := H_2 \cdot \int_2 \cdot H_2^{-1} \cdot H_1 \cdot \int_1 \cdot H_1^{-1}$$

and the claim corresponds to the identity $L \cdot S = \text{id}$. In terms of functions, this means that $y(x) = (Sr)(x)$ is a solution of

$$(Ly)(x) = r(x). \quad (2)$$

Using the Leibniz rule, H_i being a solution of the factor differential equation corresponds to

$$\partial \cdot H_i = H_i \cdot \partial + B_i \cdot H_i$$

and the invertibility corresponds to $H_i \cdot H_i^{-1} = \text{id}$. The last fact we use for proving the claim is the fundamental theorem of calculus, which corresponds to

$$\partial \cdot \int_1 = \text{id}, \quad \partial \cdot \int_2 = \text{id}.$$

In Example 2, we will show how these operator identities can be translated into noncommutative polynomials that are compatible with a quiver.

2 Preliminaries

In this section, we recall the main definitions and basic facts from [17] that formalize compatibility of polynomials with a labelled quiver.

We fix a commutative ring R with unit as well as a set X . We consider the free noncommutative algebra $R\langle X \rangle$ generated by the alphabet X : it can be regarded as the ring of noncommutative polynomials in the set of indeterminates X with coefficients in R , where indeterminates commute with coefficients but not with each other. The monomials are words $x_1 \dots x_n \in \langle X \rangle$, $x_i \in X$, including the empty word 1. Every polynomial $f \in R\langle X \rangle$ has a unique representation as a sum

$$f = \sum_{m \in \langle X \rangle} c_m m$$

with coefficients $c_m \in R$, such that only finitely many coefficients are nonzero, and its support is defined as

$$\text{supp}(f) := \{m \in \langle X \rangle \mid c_m \neq 0\},$$

where c_m are as above.

Recall that a quiver is a tuple (V, E, s, t) , where V is a set of vertices, E is a set of edges, and $s, t : E \rightarrow V$ are source and target maps, that are extended to all paths $p = e_n \dots e_1$ by letting $s(p) = s(e_1)$ and $t(p) = t(e_n)$. For every vertex $v \in V$, there is a distinct path ϵ_v that starts and ends in v without passing through any edge, and which acts as a local identity on paths p , that is $\epsilon_{t(p)} p = p = p \epsilon_{s(p)}$. A labelled quiver, $Q = (V, E, X, s, t, l)$ is a quiver equipped with a label function $l : E \rightarrow X$ of edges into the alphabet X . We extend l into a function from paths to monomials by letting $l(p) = l(e_n) \dots l(e_1) \in \langle X \rangle$, and $l(\epsilon_v) = 1$ is the empty word for every vertex v . From now on, we fix a labelled quiver $Q = (V, E, X, s, t, l)$.

Definition 1. *Given a labelled quiver and a monomial m , we define the set of signatures of m as*

$$\sigma(m) := \{(s(p), t(p)) \mid p \text{ a path in } Q \text{ with } l(p) = m\} \subseteq V \times V.$$

A polynomial $f \in R\langle X \rangle$ is said to be compatible with Q if its set of signatures $\sigma(f)$ is non empty, where:

$$\sigma(f) := \bigcap_{m \in \text{supp}(f)} \sigma(m) \subseteq V \times V.$$

Finally, we denote by $s(f)$ and $t(f)$ the images of $\sigma(f)$ through the natural projections of $V \times V$ on V .

Note that we have $\sigma(0) = V \times V$ and $\sigma(1) = \{(v, v) \mid v \in V\}$.

Computing with compatible polynomials does not always result in compatible polynomials. However, under some conditions, the sum and product of compatible polynomials are compatible as well. The following properties of signatures are straightforward to prove; see also Lemmas 10 and 11 in [17].

Lemma 1. *Let $f, g \in R\langle X \rangle$ be compatible with Q . Then,*

1. If $\sigma(f) \cap \sigma(g) \neq \emptyset$, then $f + g$ is compatible with Q and $\sigma(f + g) \supseteq \sigma(f) \cap \sigma(g)$.
2. If $s(f) \cap t(g) \neq \emptyset$, then fg is compatible with Q and

$$\sigma(fg) \supseteq \{(u, w) \in s(g) \times t(f) \mid \exists v \in s(f) \cap t(g) : (u, v) \in \sigma(g) \wedge (v, w) \in \sigma(f)\}.$$

We use the following conventions when we draw labelled quivers: we do not give names to vertices and edges, but denote them by a bullet and an arrow oriented from its source to its target, respectively, and the label of an edge is simply written above the arrow representing this edge.


Example 2. Let us continue the running example. The Leibniz rule and invertibility for H_1 and H_2 and the fundamental theorem of calculus correspond to the following noncommutative polynomials in $\mathbb{Z}\langle X \rangle$, where $X = \{h_1, h_2, b_1, b_2, \tilde{h}_1, \tilde{h}_2, i, d\}$.

$$\begin{aligned} f_1 &= dh_1 - h_1d - b_1h_1, & f_2 &= dh_2 - h_2d - b_2h_2, \\ f_3 &= h_1\tilde{h}_1 - 1, & f_4 &= h_2\tilde{h}_2 - 1, \\ f_5 &= di - 1 \end{aligned}$$

We collect these polynomials in the set $F := \{f_1, \dots, f_5\}$. Notice that we represent the two integrals by a single indeterminate, so we only need one polynomial for the fundamental theorem of calculus. The claim corresponds to

$$f := (d - b_1)(d - b_2)h_2i\tilde{h}_2h_1i\tilde{h}_1 - 1.$$

Since integration and differentiation decrease and increase the regularity of functions, it is natural to consider the following labelled quiver with 3 vertices (more details are given Section 4) with labels in the alphabet X .



Either directly or by the package, we check that f and each element of F is compatible with the quiver. Denoting the vertices from left to right by v_1, v_2, v_3 , we obtain the following signatures.

$$\begin{aligned} \sigma(f_1) &= \{(v_2, v_3)\}, & \sigma(f_2) &= \{(v_1, v_2)\}, \\ \sigma(f_3) &= \{(v_3, v_3)\}, & \sigma(f_4) &= \{(v_2, v_2)\}, \\ \sigma(f_5) &= \{(v_2, v_2), (v_3, v_3)\}, \\ \sigma(f) &= \{(v_3, v_3)\} \end{aligned}$$

To determine $\sigma(f)$, for example, notice that $\sigma(h_2i\tilde{h}_2h_1i\tilde{h}_1) = \{(v_3, v_1)\}$ and that $\sigma(dd) = \sigma(b_1d) = \sigma(db_2) = \sigma(b_1b_2) = \{(v_1, v_3)\}$ and recall that $\sigma(1)$ contains all pairs of the form (v_i, v_i) .

3 Q-consequences

The following definition characterizes the situations when a representation of the claim in terms of the assumptions is also valid in terms of operators. This generalizes the notion of Q -consequence given in [17]. Throughout the section, we fix a labelled quiver Q with labels in a set X .

Definition 2. A Q -consequence of $F \subseteq R\langle X \rangle$ is a polynomial $f \in R\langle X \rangle$, compatible with Q , such that there exist $g_i \in F$, $a_i, b_i \in R\langle X \rangle$, $1 \leq i \leq n$, such that

$$f = \sum_{i=1}^n a_i g_i b_i, \quad (3)$$

and for every $(u, v) \in \sigma(f)$ and every i , there exist vertices u_i, v_i such that $(u, u_i) \in \sigma(b_i)$, $(u_i, v_i) \in \sigma(g_i)$ and $(v_i, v) \in \sigma(a_i)$.

The conditions on the signatures mean that there exist three paths in the quiver as illustrated in the following diagram.

$$\begin{array}{ccc} u & \xrightarrow{f} & v \\ b_i \downarrow & & \uparrow a_i \\ u_i & \xrightarrow{g_i} & v_i \end{array}$$

Proving that a given representation (3) satisfies the required conditions of the above definition is straightforward. In Proposition 1, we give an alternative criterion for Q -consequences. This criterion will play an important role later in Section 5 on rewriting. Before, we need the following lemma.

Lemma 2. Let $m \in \langle X \rangle$ be a monomial and $g \in R\langle X \rangle$ be a polynomial such that $\sigma(m) \subseteq \sigma(g)$. Then, for all monomials $a, b \in \langle X \rangle$, we have $\sigma(amb) \subseteq \sigma(agb)$. Moreover, for every $(u, v) \in \sigma(amb)$, there exist two vertices \tilde{u}, \tilde{v} such that $(u, \tilde{u}) \in \sigma(b)$, $(\tilde{u}, \tilde{v}) \in \sigma(g)$, and $(\tilde{v}, v) \in \sigma(a)$.

Proof. For every $(u, v) \in \sigma(amb)$, there exists a path from u to v with label amb . We split this path in 3 parts: the first part β has label b , the third part α has label a , and the second part has label m . Since $\sigma(m) \subseteq \sigma(g)$, for every $\tilde{m} \in \text{supp}(g)$, there also exists a path γ from $\tilde{u} := t(\beta)$ to $\tilde{v} := s(\alpha)$ with label \tilde{m} , as pictured on the following diagram

$$u \xrightarrow{b} \tilde{u} \begin{array}{c} \xrightarrow{m} \\ \xrightarrow{\tilde{m}} \end{array} \tilde{v} \xrightarrow{a} v$$

Hence, $a\tilde{m}b$ is the label of $\alpha\gamma\beta$. Consequently, $\sigma(amb) \subseteq \sigma(a\tilde{m}b)$ for every $\tilde{m} \in \text{supp}(g)$, and $(\tilde{u}, \tilde{v}) \in \sigma(g)$. \square

Proposition 1. Let $F \subseteq R\langle X \rangle$ be a set of polynomials such that for every $g \in F$, there exists $m_g \in \text{supp}(g)$ such that $\sigma(m_g) \subseteq \sigma(g)$. Let $f \in R\langle X \rangle$ be a compatible polynomial such that there exist $\lambda_i \in R$, $g_i \in F$, $a_i, b_i \in \langle X \rangle$, $1 \leq i \leq n$, such that

$$f = \sum_{i=1}^n \lambda_i a_i g_i b_i, \quad (4)$$

and for each i , we have $\sigma(f) \subseteq \sigma(a_i m_{g_i} b_i)$. Then, f is a Q -consequence of F .

Proof. By hypotheses, f is compatible and for every $(u, v) \in \sigma(f)$ and for every $1 \leq i \leq n$, we have $(u, v) \in \sigma(a_i m_{g_i} b_i)$. Hence, using the hypothesis $\sigma(m_{g_i}) \subseteq \sigma(g_i)$, from Lemma 2, there exist vertices u_i and v_i such that $(u, u_i) \in \sigma(b_i)$, $(u_i, v_i) \in \sigma(g_i)$ and $(v_i, v) \in \sigma(a_i)$. As a consequence, f is a Q -consequence of F . \square

Note that if for $m_g \in \text{supp}(g)$, we have $\sigma(m_g) \subseteq \sigma(g)$, then $\sigma(m_g) = \sigma(g)$ holds by definition.

Example 3. Let us continue Example 2. We show that f is a Q -consequence of F by considering the following representation:

$$f = f_1 i \tilde{h}_1 + (d - b_1) f_2 i \tilde{h}_2 h_1 i \tilde{h}_1 + f_3 + (d - b_1) f_4 h_1 i \tilde{h}_1 + (d - b_1) h_2 f_5 \tilde{h}_2 h_1 i \tilde{h}_1 + h_1 f_5 \tilde{h}_1. \quad (5)$$

Such a representation can be obtained with the package by tracking cofactors in polynomial reduction w.r.t. a monomial order. Here, we consider a degree-lexicographic order such that d is greater than h_i 's and b_i 's. Then, f can be reduced to zero using F , which gives (5). Now, we have to check assumptions on signatures, either by checking Definition 2 or the assumptions of Proposition 1, both options are implemented in the package. For applying Proposition 1 by hand, we can choose $m_{f_1} = d h_1$, $m_{f_2} = h_2 d$, $m_{f_3} = h_1 \tilde{h}_1$, $m_{f_4} = h_2 \tilde{h}_2$, and $m_{f_5} = d i$, which satisfy $m_{f_i} \in \text{supp}(f_i)$ and $\sigma(m_{f_i}) = \sigma(f_i)$. Expanding (5) in the form (4), we may check that $\sigma(a_i m_{g_i} b_i) = \{(v_3, v_3)\} = \sigma(f)$ for every summand in the representation (4), which proves that f is a Q -consequence of F .

To conclude this section, we prove that the property of being a Q -consequence is transitive, which we will exploit in Section 6.

Theorem 1. Let $F, G \subseteq R\langle X \rangle$ be sets of polynomials such that each element of G is a Q -consequence of F . Then, any Q -consequence of G is also a Q -consequence of F .

Proof. Let h be a Q -consequence of G , so that it is compatible with Q . Moreover, $h = \sum_i a_i g_i b_i$, with $g_i \in G$ and $a_i, b_i \in R\langle X \rangle$ such that for every $(u, v) \in \sigma(h)$ and every i , there exist vertices u_i, v_i such that $(u, u_i) \in \sigma(b_i)$, $(u_i, v_i) \in \sigma(g_i)$ and $(v_i, v) \in \sigma(a_i)$. Since every element of G is a Q -consequence of F , for each g_i , there exist $a_{i,j}, b_{i,j} \in R\langle X \rangle$ and $f_{i,j} \in F$ such that $g_i = \sum_j a_{i,j} f_{i,j} b_{i,j}$ and for every $(u_i, v_i) \in \sigma(g_i)$ and every j , there exist $(u_{i,j}, v_{i,j}) \in \sigma(f_{i,j})$ such that $(u_i, u_{i,j}) \in \sigma(b_{i,j})$ and $(v_{i,j}, v_i) \in \sigma(a_{i,j})$. All together, we have

$$h = \sum_i \sum_j a_i a_{i,j} f_{i,j} b_{i,j} b_i.$$

For every j , u_i and v_i belong to $s(b_{i,j}) \cap t(b_i)$ and $s(a_i) \cap t(a_{i,j})$, respectively, so that from Point 2 of Lemma 1, $(u, u_{i,j}) \in \sigma(b_{i,j} b_i)$ and $(v_{i,j}, v) \in \sigma(a_i a_{i,j})$, respectively. Hence, h is a Q -consequence of F . \square

4 Realizations

In this section, we formalize the translation of polynomials to operators by substituting variables by basic operators. In particular, we show in Theorem 2

that being a Q -consequence is enough to ensure that the corresponding operator identity can be inferred from the assumed operator identities. To this end, we summarize the relevant notions and basic facts from [17, Section 5].

For a quiver (V, E, s, t) and a ring R , (\mathcal{M}, φ) is called a *representation* of the quiver (V, E, s, t) , if $\mathcal{M} = (\mathcal{M}_v)_{v \in V}$ is a family of R -modules and φ is a map that assigns to each $e \in E$ a R -linear map $\varphi(e) : \mathcal{M}_{s(e)} \rightarrow \mathcal{M}_{t(e)}$, see e.g. [5, 6]. Note that any nonempty path $e_n \dots e_1$ in the quiver induces a R -linear map $\varphi(e_n) \dots \varphi(e_1)$, since the maps $\varphi(e_{i+1})$ and $\varphi(e_i)$ can be composed for every $i \in \{1, \dots, n-1\}$ by definition of φ . Similarly, for every $v \in V$, the empty path ϵ_v induces the identity map on \mathcal{M}_v .

Remark 1. *All notions and results of this section naturally generalize to R -linear categories by considering objects and morphisms in such a category instead of R -modules and R -linear maps, respectively. For more details, see Section 5.2 in [17].*

Definition 3. *Let R be a ring and let Q be a labelled quiver with labelling l . We call a representation (\mathcal{M}, φ) of Q consistent with the labelling l if for any two nonempty paths $p = e_n \dots e_1$ and $q = d_n \dots d_1$ in Q with the same source and target, equality of labels $l(p) = l(q)$ implies $\varphi(e_n) \dots \varphi(e_1) = \varphi(d_n) \dots \varphi(d_1)$ as R -linear maps.*

Remark 2. *If all paths with the same source and target have distinct labels, then every representation of that labelled quiver is consistent with its labelling. In particular, this holds if for every vertex all outgoing edges have distinct labels or analogously for incoming edges. These sufficient conditions can be verified without the need for considering all possible paths.*

For Definition 4 and Lemma 3, we fix a ring R , a labelled quiver $Q = (V, E, X, s, t, l)$ and a consistent representation $\mathcal{R} = (\mathcal{M}, \varphi)$ of Q . In order to define realizations of a polynomial, we first need to introduce some notations. Given two vertices v, w , we write $R\langle X \rangle_{v,w}$ for the set of polynomials $f \in R\langle X \rangle$ such that $(v, w) \in \sigma(f)$. From Point 1 of Lemma 1, $R\langle X \rangle_{v,w}$ is a module, and it is clear that this module is free with basis the set of monomials m such that $(v, w) \in \sigma(m)$. We also denote by $\text{Hom}_R(\mathcal{M}_v, \mathcal{M}_w)$ the set of R -linear maps from \mathcal{M}_v to \mathcal{M}_w .

Definition 4. *For $v, w \in V$, we define the R -linear map $\varphi_{v,w} : R\langle X \rangle_{v,w} \rightarrow \text{Hom}_R(\mathcal{M}_v, \mathcal{M}_w)$ by*

$$\varphi_{v,w}(l(e_n \dots e_1)) := \varphi(e_n) \dots \varphi(e_1)$$

for all nonempty paths $e_n \dots e_1$ in Q from v to w and, if $v = w$, also by $\varphi_{v,v}(1) := \text{id}_{\mathcal{M}_v}$. For all $f \in R\langle X \rangle_{v,w}$, we call the R -linear map $\varphi_{v,w}(f)$ a realization of f w.r.t. the representation \mathcal{R} of Q .

Notice that the map $\varphi_{v,w}$ is well-defined since, by consistency of \mathcal{R} , for every monomial $m \in R\langle X \rangle_{v,w}$, its realization $\varphi_{v,w}(m)$ does not depend on the path from v to w with label m .

In the proof of Theorem 2, we use an intermediate result given in [17, Lemma 31], whose statement is the following.

Lemma 3. *Let $u, v, w \in V$. Then, for all $f \in R\langle X \rangle_{v,w}$ and $g \in R\langle X \rangle_{u,v}$, we have that $fg \in R\langle X \rangle_{u,w}$ and*

$$\varphi_{u,w}(fg) = \varphi_{v,w}(f) \cdot \varphi_{u,v}(g).$$

Theorem 2. *Let $F \subseteq R\langle X \rangle$ be a set of polynomials and let Q be a labelled quiver with labels in X . If a polynomial $f \in R\langle X \rangle$ is a Q -consequence of F , then for all consistent representations of the quiver Q such that all realizations of all elements of F are zero, all realizations of f are zero.*

Proof. Assume that f is a Q -consequence, so that it is compatible with Q and it can be written in the form $\sum a_i g_i b_i$, such that for each $(u, v) \in \sigma(f)$ and each i , there exist vertices u_i, v_i such that $(u, u_i) \in \sigma(b_i)$, $(u_i, v_i) \in \sigma(g_i)$ and $(v_i, v) \in \sigma(a_i)$. Let us fix a consistent representation $\mathcal{R} = (\mathcal{M}, \varphi)$ of Q . By linearity of $\varphi_{u,v}$ and from Lemma 3, we have

$$\varphi_{u,v}(f) = \sum \varphi_{u,v}(a_i g_i b_i) = \sum \varphi_{v_i,v}(a_i) \cdot \varphi_{u_i,v_i}(g_i) \cdot \varphi_{u,u_i}(b_i).$$

Hence, if all realizations of all elements of F are zero, then $\varphi_{u,v}(f) = 0$, which means that all realizations of f w.r.t \mathcal{R} are zero. \square

Example 4. *We finish our proof of (2) by considering certain representations of the quiver of Example 2. For a nonnegative integer k and an open interval $I \subseteq \mathbb{R}$, we assign the spaces $C^k(I)$, $C^{k+1}(I)$, and $C^{k+2}(I)$ to the vertices from right to left. Hence, differentiation and integration induce operators $\partial : C^{k+1}(I) \rightarrow C^k(I)$, $\partial : C^{k+2}(I) \rightarrow C^{k+1}(I)$, $\int_1 : C^k(I) \rightarrow C^{k+1}(I)$, and $\int_2 : C^{k+1}(I) \rightarrow C^{k+2}(I)$. We also assume the following regularity of functions: B_1 is C^k , H_1 and B_2 are C^{k+1} and H_2 is C^{k+2} on I . Then, the natural representation associated with these operators is consistent. Moreover, we have seen in Example 3 that f is a Q -consequence of F . Since all realizations of f_i 's are zero, by Theorem 2, all realizations of f are zero. In particular, for every nonnegative integer k and every $r(x) \in C^k(I)$, the function $y(x)$ defined by (1) is a solution of the inhomogeneous differential equation (2).*

Instead of considering scalar differential equations we could consider differential systems of the form (2) for vector-valued functions $y(x)$ of arbitrary dimension n . More explicitly, we can also consider coefficients $B_1(x), B_2(x)$ as $n \times n$ matrices, $r(x)$ as a vector of dimension n , and $H_1(x)$ and $H_2(x)$ as fundamental matrix solutions of the homogeneous systems $y'(x) - B_i(x)y(x) = 0$. We still obtain consistent representations of the quiver where the vertices are mapped to $C^k(I)^n, C^{k+1}(I)^n$ and $C^{k+2}(I)^n$, respectively. Then, Theorem 2 immediately proves that the function $y(x)$ defined by (1) is a solution of the inhomogeneous differential equation (2). Similarly, analogous statements for other suitable functional spaces can be proven just by choosing different representations of the quiver.

5 Compatible rewriting

In this section, we give conditions on polynomials such that rewriting to zero of a compatible polynomial by them proves that it is a Q -consequence. First, we recall from [17, Definition 2] a general notion of rewriting one polynomial by

another in terms of an arbitrary monomial division. Notice that the standard polynomial reduction is a particular case, where m is the leading monomial of g w.r.t. a monomial order and λ is such that amb is cancelled in (6).

Definition 5. Let $g \in R\langle X \rangle$ be a polynomial and let $m \in \text{supp}(g)$. Let $f \in R\langle X \rangle$ be a polynomial such that m divides some monomial $m_f \in \text{supp}(f)$, i.e., $m_f = amb$ for monomials $a, b \in \langle X \rangle$. For every $\lambda \in R$, we say that f can be rewritten to

$$h := f + \lambda agb, \quad (6)$$

using (g, m) .

We fix a labelled quiver Q with labels in X . It turns out that to obtain Q -consequences using rewriting (Theorem 3), we need to choose suitable divisor monomials such that signatures only increase. In particular, this is the case when divisor monomials have minimal signature, as stated in the following lemma.

Lemma 4. Let $g \in R\langle X \rangle$ be a polynomial and let $m \in \text{supp}(g)$ be such that $\sigma(m) = \sigma(g)$. If f can be rewritten to $h = f + \lambda agb$ using (g, m) , then

$$\sigma(f) \subseteq \sigma(h) \quad \text{and} \quad \sigma(f) \subseteq \sigma(amb).$$

Proof. By definition of signatures, $\sigma(f) \subseteq \sigma(amb)$. By Lemma 2 and from $\sigma(m) = \sigma(g)$, we have $\sigma(amb) \subseteq \sigma(agb)$. Altogether, $\sigma(f)$ is included in $\sigma(agb)$, which itself is contained in $\sigma(\lambda agb)$. From 1. of Lemma 1, we deduce $\sigma(f) \subseteq \sigma(h)$. \square

Now, we define the rewriting relation induced by a fixed choice of divisor monomials and its compatibility with a quiver. For any rewriting relation we denote single rewriting steps by \rightarrow and the reflexive transitive closure by $\overset{*}{\rightarrow}$.

Definition 6. Let $G \subseteq R\langle X \rangle$ be a set of polynomials and let $\text{DM} : G \rightarrow \mathcal{P}(\langle X \rangle)$ be a function from G to the power set of $\langle X \rangle$, such that $\text{DM}(g) \subseteq \text{supp}(g)$, for every $g \in G$.

1. For $g \in G$, we say that $m \in \text{DM}(g)$ is a divisor monomial of g w.r.t. DM .
2. We say that f rewrites to h by (G, DM) , denoted as $f \rightarrow_{G, \text{DM}} h$, if there exists $g \in G$ and a divisor monomial $m \in \text{DM}(g)$ such that f can be rewritten to h using (g, m) .
3. We say that DM is compatible with a labelled quiver Q if for every $g \in G$ and every $m \in \text{DM}(g)$, we have $\sigma(m) = \sigma(g)$.

From now on, we fix a set of polynomials $G \subseteq R\langle X \rangle$ as well as a map DM selecting divisor monomials.

Remark 3. Notice that there exist two extreme cases for the definition of DM :

1. DM selects exactly one monomial for each $g \in G$, for instance, the leading monomial $\text{LM}(g)$ w.r.t. a monomial order, see the example in Section 6.

2. All monomials in $\text{supp}(g)$ are divisor monomials. Then, $\rightarrow_{G, \text{DM}}$ coincides with the rewriting relation introduced in [17, Definition 2], for which ideal membership is equivalent to reduction to zero [17, Lemma 4]. Moreover, if such a DM is compatible with Q , then all polynomials in G are uniformly compatible, i.e., every monomial of a polynomial has the same signature. The following theorem gives a generalization of Corollary 17 in [17].

Theorem 3. *Let $G \subseteq R\langle X \rangle$ be a set of polynomials and let DM be a function selecting divisor monomials as in Definition 6. Let $f \in R\langle X \rangle$ be a polynomial such that $f \xrightarrow{*}_{G, \text{DM}} 0$. Then, for every labelled quiver Q with labels X such that DM is compatible with Q , we have that*

$$f \text{ is compatible with } Q \iff f \text{ is a } Q\text{-consequence of } G.$$

Proof. Since f rewrites to zero, there exists a sequence $f = h_0 \rightarrow h_1 \rightarrow \dots \rightarrow h_n = 0$. Hence, there exist $\lambda_i \in R$, $a_i, b_i \in \langle X \rangle$, $g_i \in G$, and $m_i \in \text{DM}(g_i)$ such that $h_i = h_{i-1} + \lambda_i a_i g_i b_i$ and $a_i m_i b_i \in \text{supp}(h_{i-1})$. Hence, f can be written as $f = \sum_{i=1}^n -\lambda_i a_i g_i b_i$. From Lemma 4, we conclude inductively that $\sigma(f) \subseteq \sigma(h_{i-1}) \subseteq \sigma(a_i m_i b_i)$. Hence, if f is compatible with Q , then f is a Q -consequence of G by Proposition 1. Conversely, if f is a Q -consequence of G , then it is compatible by definition. \square

Example 5. *Let us translate Example 3 in the language introduced in this section. The leading monomials w.r.t. the degree-lexicographic order used in that example can be understood as the divisor monomials selected by the function DM defined on F such that $\text{DM}(f_i) = \{\text{LM}(f_i)\}$ holds for all i . In particular,*

$$\begin{aligned} \text{DM}(f_1) &= \{dh_1\}, & \text{DM}(f_2) &= \{dh_2\}, \\ \text{DM}(f_3) &= \{h_1\tilde{h}_1\}, & \text{DM}(f_4) &= \{h_2\tilde{h}_2\}, & \text{DM}(f_5) &= \{di\}. \end{aligned}$$

Then, DM is not compatible with Q , since $\sigma(f_2) = \{(v_1, v_2)\}$ is not equal to $\sigma(dh_2) = \{(v_1, v_2), (v_2, v_3)\}$. Hence, we cannot apply Theorem 3 to show that f is a Q -consequence of F even though $f \xrightarrow{}_{F, \text{DM}} 0$. So, we need to look at the explicit representation of f induced by this reduction, which was already done in Example 3. In order to apply Theorem 3, we need to redefine DM so that it is compatible with Q . In particular, we need to impose $\text{DM}(f_2) \subseteq \{h_2d, b_2h_2\}$. If $b_2h_2 \in \text{DM}(f_2)$, then $f \xrightarrow{*}_{F, \text{DM}} 0$, which gives another proof that f is a Q -consequence of F based on Theorem 3. Otherwise, if $\text{DM}(f_2) = \{h_2d\}$, then f is irreducible w.r.t. $\xrightarrow{*}_{F, \text{DM}}$. Therefore, we need to complete F with Q -consequences of it such that DM remains compatible with Q and f reduces to zero, which is the topic of the next section.*

6 Compatible reductions and partial Gröbner bases

In this section, we discuss standard noncommutative polynomial reduction as a special case of the rewriting approach from the previous section. Since in the noncommutative case, Gröbner bases are not necessarily finite, see [15], we also have to work with partial Gröbner bases which are obtained by finitely

many iterations of the Buchberger procedure. We adapt the noncommutative Buchberger procedure for computing (partial) Gröbner bases that can be used for compatible rewriting.

In what follows, R is assumed to be a field \mathbb{K} and we fix a monomial order \leq on $\langle X \rangle$, that is, a well-founded total order compatible with multiplication on $\langle X \rangle$. We also fix a labelled quiver Q with labels in X and a set of polynomials $F \subseteq \mathbb{K}\langle X \rangle$. Given a set of polynomials $G \subseteq \mathbb{K}\langle X \rangle$, one step of the standard polynomial reduction w.r.t. G is denoted by $f \rightarrow_G h$.

As explained in Remark 3, the monomial order induces the DM function that selects leading monomials of a set $G \subseteq \mathbb{K}\langle X \rangle$. This DM function is compatible with Q if and only if all elements of G are Q -order compatible in the following sense.

Definition 7. *A compatible polynomial f is said to be Q -order compatible if $\sigma(\text{LM}(f)) = \sigma(f)$.*

By transitivity of Q -consequences, see Theorem 1, and Theorem 3, we obtain the following statement.

Corollary 1. *Let $F \subseteq \mathbb{K}\langle X \rangle$, $G \subseteq (F)$, and $f \in \mathbb{K}\langle X \rangle$ such that $f \xrightarrow{*}_G 0$. Then, for all labelled quivers Q such that all elements of G are both Q -consequences of F and Q -order compatible, we have*

$$f \text{ is compatible with } Q \iff f \text{ is a } Q\text{-consequence of } F.$$

Remark 4. *For polynomials, being Q -order compatible can also be interpreted in terms of a partial monomial order. Given $m, m' \in \langle X \rangle$, we define $m \leq_Q m'$ if $m \leq m'$ and $\sigma(m') \subseteq \sigma(m)$. The partial order \leq_Q respects multiplication of monomials since, by Lemma 2, $\sigma(m') \subseteq \sigma(m)$ implies $\sigma(am'b) \subseteq \sigma(amb)$ for all $a, b \in \langle X \rangle$. Then, f is Q -order compatible if and only if $\text{supp}(f)$ admits a greatest element for \leq_Q .*

Candidates for G as in Corollary 1 are partial Gröbner bases that are computed by the noncommutative Buchberger procedure [4, 15]. However, in view of the assumptions, we only add reduced S -polynomials that are both Q -consequences of F and Q -order compatible in each iteration. Checking Q -order compatibility is easy. Selecting Q -consequences is harder since we do not want to use explicit representations as in Definition 2. Instead, we propose a simpler criterion based on the following lemma and discussion.

First, we recall some terminology and fix notations for S -polynomials. Let $G \subseteq \mathbb{K}\langle X \rangle$. *Ambiguities* of G defined in [1], also called compositions in [2], are given by minimal overlaps or inclusions of the two leading monomials $\text{LM}(g)$ and $\text{LM}(g')$, where g and g' belong to G . Formally, each ambiguity can be described by a 6-tuple $\mathbf{a} = (g, g', a, b, a', b')$, where a, b, a', b' are monomials such that, among other conditions, we have

$$a \text{LM}(g)b = a' \text{LM}(g')b'.$$

This monomial is called the *source* of \mathbf{a} and the S -polynomial of \mathbf{a} is $\text{SP}(\mathbf{a}) := agb - a'g'b'$, cf. [15].

Lemma 5. *Let $G \subseteq \mathbb{K}\langle X \rangle$ be a set of Q -order compatible polynomials and let s be a S -polynomial of G with source a compatible monomial $m \in \langle X \rangle$. Then*

$\sigma(m) \subseteq \sigma(s)$. If moreover, $s \xrightarrow{*}_G \hat{s}$ with $\sigma(\hat{s}) \subseteq \sigma(m)$, then $\sigma(s) = \sigma(\hat{s}) = \sigma(m)$ and \hat{s} is a Q -consequence of G .

Proof. Since s is a S -polynomial of G of source m , there exist $g, g' \in G$ and monomials $a, a', b, b' \in \langle X \rangle$ such that $s = (m - agb) - (m - a'g'b')$ with $a \text{LM}(g)b = a' \text{LM}(g')b' = m$.

Let us prove the first assertion. The polynomials g and g' being Q -order compatible, we have $\sigma(\text{LM}(g)) = \sigma(g)$ and $\sigma(\text{LM}(g')) = \sigma(g')$. Hence, from Lemma 2, we have

$$\begin{aligned} \sigma(m) &= \sigma(a \text{LM}(g)b) \subseteq \sigma(agb), \\ \sigma(m) &= \sigma(a' \text{LM}(g')b') \subseteq \sigma(a'g'b'). \end{aligned} \quad (7)$$

From this and $s = a'g'b' - agb$, we get that $\sigma(m) \subseteq \sigma(s)$.

Now, we assume that $s \xrightarrow{*}_G \hat{s}$, that is there is a rewriting sequence

$$s_0 = s \rightarrow_G s_1 \rightarrow_G \cdots \rightarrow_G s_n = \hat{s},$$

so that there exist $g_i \in G$, $a_i, b_i \in \langle X \rangle$ and $\lambda_i \in \mathbb{K}$, $1 \leq i \leq n$, such that $a_i \text{LM}(g_i)b_i \in \text{supp}(s_i)$ and $s_{i+1} = s_i + \lambda_i a_i g_i b_i$, so that we have $\hat{s} - s = \sum_i \lambda_i a_i g_i b_i$ and

$$\hat{s} = \sum_{i=1}^n \lambda_i a_i g_i b_i + a'g'b' - agb. \quad (8)$$

Using inductively $s_i \rightarrow_G s_{i+1}$ and Lemma 4, we get

$$\sigma(s) \subseteq \sigma(\hat{s}) \quad \text{and} \quad \sigma(s) \subseteq \sigma(a_i \text{LM}(g_i)b_i), \quad (9)$$

so that we have

$$\sigma(m) \subseteq \sigma(s) \subseteq \sigma(\hat{s}).$$

If, moreover $\sigma(\hat{s}) \subseteq \sigma(m)$, then we get the following sequence of inclusions:

$$\sigma(\hat{s}) \subseteq \sigma(m) \subseteq \sigma(s) \subseteq \sigma(\hat{s}).$$

Hence, the equality $\sigma(\hat{s}) = \sigma(s) = \sigma(m)$ holds. Now, we show that \hat{s} is a Q -consequence using Proposition 1. Since the elements of G are Q -order compatible, we have $\sigma(\text{LM}(\tilde{g})) = \sigma(\tilde{g})$, for all $\tilde{g} \in G$. Moreover, since m is compatible and $\sigma(\hat{s}) = \sigma(m)$, \hat{s} is compatible. Finally, from (7) and (9), we have the following inclusions: $\sigma(\hat{s}) \subseteq \sigma(a \text{LM}(g)b)$, $\sigma(\hat{s}) \subseteq \sigma(a' \text{LM}(g')b')$ and $\sigma(\hat{s}) \subseteq \sigma(a_i \text{LM}(g_i)b_i)$. As a conclusion, \hat{s} is a Q -consequence of G . \square

Starting with a set of Q -order compatible polynomials F , we apply this lemma in the case where G is the partial Gröbner basis computed in the current iteration of the completion procedure. In particular, if a reduced S -polynomial \hat{s} satisfies $\sigma(\hat{s}) \subseteq \sigma(m)$ as in the lemma, then it is a Q -consequence of G . By transitivity, it is then also a Q -consequence of F , which follows from the following observation.

Remark 5. Consider a set $F \subseteq \mathbb{K}\langle X \rangle$ of compatible polynomials and a family of sets G_i inductively defined by $G_0 = \emptyset$ and $G_{i+1} = G_i \cup \{g_{i+1}\}$, where g_{i+1} is a Q -consequence of $F \cup G_i$. Using inductively transitivity of Q -consequences proven in Theorem 1, we obtain that, for each i , all elements of $F \cup G_i$ are Q -consequences of F .

In summary, we obtain the following adaptation of the noncommutative version of Buchberger's procedure for computing a partial Gröbner basis composed of elements that are both Q -consequences of F and Q -order compatible. At each step, we select an S -polynomial s whose source m is a compatible monomial, and we keep a reduced form \hat{s} only if it is Q -order compatible and $\sigma(\hat{s}) \subseteq \sigma(m)$. This procedure is implemented in the MATHEMATICA package `OperatorGB`. Note that since the Buchberger procedure does not terminate in general for noncommutative polynomials, also our adaptation of it is not guaranteed to terminate.

Notice that the completion procedure described above can be slightly generalized by not necessarily computing reduced forms of S -polynomials. Instead, we only reduce an S -polynomial as long as it remains a Q -consequence, see the discussion above, and it remains Q -order compatible. This is stated formally in Procedure 1.

Procedure 1 Q -order compatible completion

Input: $F \subseteq \mathbb{K}\langle X \rangle$, a labelled quiver Q with labels in X , and a monomial order \leq such that every $f \in F$ is Q -order compatible

Output: $G \supseteq F$ a set of Q -consequences of F that are Q -order compatible

```

1:  $P :=$  ambiguities of  $F$ ;  $G := F$ 
2: while  $P \neq \emptyset$  do
3:   choose  $\mathbf{a} \in P$ 
4:    $P := P \setminus \{\mathbf{a}\}$ ;  $s := \text{SP}(\mathbf{a})$ ;  $m :=$  the source of  $\mathbf{a}$ 
5:   if  $m$  is compatible and  $\sigma(s) \subseteq \sigma(m)$  and  $s$  is  $Q$ -order compatible then
6:     while  $\exists s' : s \rightarrow_G s'$  do
7:       if  $s' = 0$  then
8:         go to 2 (i.e., break the outer if statement)
9:       else if  $\sigma(s') \subseteq \sigma(m)$  and  $s'$  is  $Q$ -order compatible then
10:         $s := s'$ 
11:       else
12:         go to 15 (i.e., break the inner while loop)
13:       end if
14:     end while
15:      $G := G \cup \{s\}$ 
16:      $P := P \cup \{\text{ambiguities created by } s\}$ 
17:   end if
18: end while
19: return  $G$ 

```

Due to the checks in line 9, each element g of the output G of the procedure is both a Q -consequence of F and Q -order compatible. In summary, we have shown that our procedure is correct.

Theorem 4. *Let $F \subseteq \mathbb{K}\langle X \rangle$ be a set of polynomials, let Q be a labelled quiver with labels in X and let \leq be a monomial order such that each element of F is Q -order compatible. Then, each element of the output G of Procedure 1 is both a Q -consequence of F and Q -order compatible.*

Example 6. *Let us continue Example 5 in the case $\text{DM}(f_2) = \{h_2d\}$. For that, we consider the field $\mathbb{K} = \mathbb{Q}$ and a degree-lexicographic order such that $d < b_2 < h_2$ and d is greater than b_1 and h_1 . Then, choosing the ambiguity*

$(f_2, f_5, 1, i, h_2, 1)$, the first iteration of the outer loop in Procedure 1 yields $G := F \cup \{b_2h_2i - dh_2i + h_2\}$. With this G , we have $f \xrightarrow{*}_G 0$. From this reduction to 0, and since f is compatible with Q , f is a Q -consequence of F by Corollary 1. These computations can also be done by the package.

Remark 6. We consider the special case when all edges of Q have unique labels. Then, all non-constant monomials have at most one element in their signature. Therefore, every compatible polynomial is Q -order compatible for any monomial order, since the monomial 1 is the smallest. Moreover, one can show easily that the source of an ambiguity of two polynomials is compatible whenever these two polynomials are compatible with Q . In addition, from Lemma 4, it follows that polynomial reduction of compatible polynomials by compatible ones does not change the signature unless the result of the reduction lies in \mathbb{K} . Altogether, Procedure 1 reduces to the standard Buchberger procedure (i.e., without checking signatures and compatibility during computation) as long as no S -polynomial is (or is reduced to) a nonzero constant. In other words, we have the following theorem, which, together with Theorem 2, gives a generalization of Theorem 1 in [17].

Theorem 5. Assume that edges of Q have unique labels. Let $F \subseteq \mathbb{K}\langle X \rangle$ be a set of compatible polynomials and let G be a (partial) Gröbner basis computed by the standard Buchberger procedure (i.e., disregarding Q during computation). If G does not contain a constant polynomial, then for every polynomial $f \in \mathbb{K}\langle X \rangle$ such that $f \xrightarrow{*}_G 0$, we have

$$f \text{ is compatible with } Q \iff f \text{ is a } Q\text{-consequence of } F.$$

Moreover, if $1 \notin (F)$, then this equivalence holds for every $f \in (F)$.

7 Summary and discussion

By Theorem 2, for proving new operator identities from known ones, it suffices to show that the corresponding polynomials are Q -consequences. In practice, there are several options to prove that a compatible polynomial f is a Q -consequence of some set F of compatible polynomials. Each of these options can be turned into a certificate that f is a Q -consequence of F . Given an explicit representation of f in terms of F of the form (3), one can either check Definition 2 directly, or expand cofactors into monomials and apply Proposition 1. Alternatively, using compatible rewriting, if $f \xrightarrow{*}_{F, \text{DM}} 0$ and the selection of divisor monomials by DM is compatible with Q , then f is a Q -consequence by Theorem 3. Altogether, from Theorems 2 and 3, we immediately obtain the following.

Corollary 2. Let F be a set of polynomials, DM a function selecting divisor monomials, and f a polynomial such that $f \xrightarrow{*}_{F, \text{DM}} 0$. Then, for all labelled quivers Q such that f , F , and DM are compatible with Q and for all consistent representations of Q such that all realizations of all elements of F are zero, all realizations of f are zero.

Note that rewriting to zero w.r.t. F and DM is independent of the quiver Q . In particular, if the above corollary is interpreted in terms of R -linear categories, the main result of [17], Theorem 32, is obtained as a special case by Remark 3.

More generally, if one cannot verify that f can be rewritten to zero by F , there still might exist a set G of Q -consequences of F with divisor monomials selected by some DM such that $f \xrightarrow{*}_{G, \text{DM}} 0$ and Theorem 1 can be applied. Algorithmically, based on suitable monomial orderings, Procedure 1 produces candidates G such that Corollary 1 can be used to prove that f is a Q -consequence of F by standard polynomial reduction.

Notice that Procedure 1 can be extended in various directions. For example, in order to systematically generate more Q -consequences, reduced S -polynomials that are not Q -order compatible could be collected in a separate set, which should not be used for constructing and reducing new S -polynomials. Instead of fixing a monomial ordering from the beginning, one might start with a partial ordering that is then extended during the completion procedure in order to make obtained S -polynomials Q -order compatible. More generally, without any partial ordering on monomials, one might even consider compatible functions DM which not necessarily select only one divisor monomial per polynomial and aim at completing the induced rewriting relation. However, termination of such rewriting relations is an issue. Finally, another topic for future research is to generalize the results of this paper to tensor reduction systems used for modelling linear operators as described in [12].

References

- [1] George M. Bergman. The diamond lemma for ring theory. *Adv. in Math.*, 29(2):178–218, 1978.
- [2] Leonid A. Bokut'. Imbeddings into simple associative algebras. *Algebra i Logika*, 15(2):117–142, 245, 1976.
- [3] Leonid A. Bokut, Yuqun Chen, and Yu Li. Gröbner-Shirshov bases for categories. In *Operads and universal algebra*, volume 9 of *Nankai Ser. Pure Appl. Math. Theoret. Phys.*, pages 1–23. World Sci. Publ., Hackensack, NJ, 2012.
- [4] Bruno Buchberger. Ein Algorithmus zum Auffinden der Basiselemente des Restklassenrings nach einem nulldimensionalen Polynomideal. *Universität Innsbruck, Austria, Ph. D. Thesis*, 1965. English translation in *J. Symbolic Comput.* 41 (2006), 475–511.
- [5] Harm Derksen and Jerzy Weyman. Quiver representations. *Notices Amer. Math. Soc.*, 52(2):200–206, 2005.
- [6] Peter B. Gothen and Alastair D. King. Homological algebra of twisted quiver bundles. *J. London Math. Soc. (2)*, 71(1):85–99, 2005.
- [7] Yves Guiraud, Eric Hoffbeck, and Philippe Malbos. Convergent presentations and polygraphic resolutions of associative algebras. *Math. Z.*, 293(1-2):113–179, 2019.
- [8] J. William Helton and Mark Stankus. Computer assistance for “discovering” formulas in system engineering and operator theory. *J. Funct. Anal.*, 161:289–363, 1999.

- [9] J. William Helton, Mark Stankus, and John J. Wavrik. Computer simplification of formulas in linear systems theory. *IEEE Trans. Automat. Control*, 43(3):302–314, 1998.
- [10] J. William Helton and John J. Wavrik. Rules for computer simplification of the formulas in operator model theory and linear systems. In *Nonselfadjoint operators and related topics (Beer Sheva, 1992)*, pages 325–354. Birkhäuser, Basel, 1994.
- [11] Clemens Hofstadler, Clemens G Raab, and Georg Regensburger. Certifying operator identities via noncommutative gröbner bases. *ACM Communications in Computer Algebra*, 53(2):49–52, 2019.
- [12] Jamal Hossein Poor, Clemens G. Raab, and Georg Regensburger. Algorithmic operator algebras via normal forms in tensor rings. *J. Symbolic Comput.*, 85:247–274, 2018.
- [13] F. Dell Kronewitter. Using noncommutative Gröbner bases in solving partially prescribed matrix inverse completion problems. *Linear Algebra Appl.*, 338:171–199, 2001.
- [14] Viktor Levandovskyy and Leonard Schmitz. Algorithmic algebraic proofs of identities between not only matrices. 2019. In preparation.
- [15] Teo Mora. An introduction to commutative and noncommutative Gröbner bases. *Theoret. Comput. Sci.*, 134(1):131–173, 1994.
- [16] Teo Mora. *Solving polynomial equation systems. Vol. IV. Buchberger theory and beyond*, volume 158 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2016.
- [17] Clemens G Raab, Georg Regensburger, and Jamal Hossein Poor. Formal proofs of operator identities by a single formal computation. *arXiv:1910.06165 [math.RA]*, 2019.
- [18] Markus Rosenkranz, Bruno Buchberger, and Heinz W. Engl. Solving linear boundary value problems via non-commutative Gröbner bases. *Appl. Anal.*, 82:655–675, 2003.