

On two congruences involving Franel numbers

Ji-Cai Liu

Department of Mathematics, Wenzhou University, Wenzhou 325035, PR China
jcliu2016@gmail.com

Abstract. Via symbolic summation method, we establish the following series for π^2 :

$$\sum_{k=1}^{\infty} \frac{H_k - 2H_{2k}}{(-3)^k k} = \frac{\pi^2}{18},$$

where $H_k = \sum_{j=1}^k 1/j$. We also derive a p -adic congruence related to this series. As an application, we prove two congruences involving Franel numbers, one of which was originally conjectured by Sun.

Keywords: Congruences; Harmonic numbers; Franel numbers; Bernoulli polynomials

MR Subject Classifications: 11A07, 05A19, 33F10

1 Introduction

It is well-known that (see [16, (3)])

$$\sum_{k=1}^{\infty} \frac{1}{k^2 \binom{2k}{k}} = \frac{\pi^2}{18}, \quad (1.1)$$

which can be derived from the following familiar power series expansion:

$$2(\arcsin(x/2))^2 = \sum_{k=1}^{\infty} \frac{x^{2k}}{k^2 \binom{2k}{k}}.$$

Recall that the Euler numbers and Bernoulli polynomials are defined as

$$\frac{2}{e^x + e^{-x}} = \sum_{n=0}^{\infty} E_n \frac{x^n}{n!}$$

and

$$\frac{xe^{tx}}{e^x - 1} = \sum_{k=0}^{\infty} B_k(t) \frac{x^k}{k!}.$$

In 2011, Sun [19, (1.3)] showed that (1.1) possesses the following interesting p -adic analogue:

$$\sum_{k=1}^{(p-1)/2} \frac{1}{k^2 \binom{2k}{k}} \equiv (-1)^{(p-1)/2} \frac{4}{3} E_{p-3} \pmod{p},$$

for any prime $p \geq 5$. Mattarei and Tauraso [15, Theorem 6.1] investigated congruence properties for the polynomial $\sum_{k=1}^{p-1} k^{-2} \binom{2k}{k}^{-1} t^k$.

In recent decades, infinite sums on harmonic numbers related to powers of π have been widely studied. For instance, D. Borwein and J.M. Borwein [1] proved that

$$\sum_{k=1}^{\infty} \frac{H_k^2}{k^2} = \frac{17}{360} \pi^4,$$

where the k th generalized harmonic number is given by

$$H_k^{(r)} = \sum_{j=1}^k \frac{1}{j^r},$$

with the convention that $H_k = H_k^{(1)}$. It is worth mentioning that Sun [20, 25] conjectured and proved many infinite identities involving harmonic numbers related to powers of π . For more series for π and related congruences as well as q -congruences, one can refer to [2, 3, 6–9, 13, 19, 21, 27, 28].

The first aim of the paper is to establish the following series for π^2 .

Theorem 1.1 *We have*

$$\sum_{k=1}^{\infty} \frac{H_k - 2H_{2k}}{(-3)^k k} = \frac{\pi^2}{18}. \quad (1.2)$$

Our proof of (1.2) makes use of a finite identity, which is derived from symbolic summation method. We also show that (1.2) has the following p -adic analogue.

Theorem 1.2 *For any prime $p \geq 5$, we have*

$$\sum_{k=1}^{(p-1)/2} \frac{H_k - 2H_{2k}}{(-3)^k k} \equiv \frac{1}{6} \left(\frac{p}{3}\right) B_{p-2} \left(\frac{1}{3}\right) \pmod{p}, \quad (1.3)$$

where $\left(\frac{\cdot}{3}\right)$ denotes the Legendre symbol.

In 1894, Franel [4] found that the sums of cubes of binomial coefficients:

$$f_n = \sum_{k=0}^n \binom{n}{k}^3$$

satisfy the recurrence:

$$(n+1)^2 f_{n+1} = (7n^2 + 7n + 2)f_n + 8n^2 f_{n-1}.$$

The numbers f_n are known as Franel numbers, which also appear in Strehl's identity (see [18]):

$$f_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n} = \sum_{k=0}^n \binom{n}{k} \binom{k}{n-k} \binom{2k}{k}. \quad (1.4)$$

Since the appearance of the Franel numbers, some interesting congruence properties have been gradually discovered (see [5, 11, 22, 23]). For instance, Sun [22, Theorem 1.1] proved that for any prime $p \geq 5$,

$$\sum_{k=0}^{p-1} (-1)^k f_k \equiv \left(\frac{p}{3}\right) \pmod{p^2}.$$

The second aim of the paper is prove the following two congruences involving Franel numbers and harmonic numbers.

Theorem 1.3 *For any prime $p \geq 5$, we have*

$$\sum_{k=0}^{p-1} (-1)^k f_k H_k^{(2)} \equiv \frac{1}{2} B_{p-2} \left(\frac{1}{3}\right) \pmod{p}, \quad (1.5)$$

$$\sum_{k=0}^{p-1} (-1)^k f_k H_k \equiv -2 \left(\frac{p}{3}\right) q_p(3) \pmod{p}, \quad (1.6)$$

where $q_p(3)$ is the Fermat quotient $(3^{p-1} - 1)/p$.

Note that (1.5) was originally conjectured by Sun [24, Conjecture 57]. Our proof of (1.5) is based on Theorem 1.2. We shall prove Theorems 1.1–1.3 in Sections 2–4, respectively.

2 Proof of Theorem 1.1

Before proving Theorem 1.1, we require the following finite identity.

Lemma 2.1 *For any non-negative integer n , we have*

$$\sum_{k=0}^n (-4)^k \binom{n}{k} \sum_{j=1}^k \frac{1}{j^2 \binom{2j}{j}} = (-3)^n \sum_{k=1}^n \frac{H_k - 2H_{2k}}{(-3)^k k}. \quad (2.1)$$

Proof. By using symbolic summation package **Sigma** due to Schneider [17], we find that both sides of (2.1) satisfy the same recurrence:

$$3(n+1)(2n+3)S(n) - (2n+5)(5n+9)S(n+1) + (2n^2+17n+29)S(n+2) + (n+3)(2n+5)S(n+3) = 0.$$

It is trivial to check that both sides of (2.1) are equal for $n = 0, 1, 2$. □

Proof of (1.2). By (2.1), we have

$$\begin{aligned} \sum_{k=1}^n \frac{H_k - 2H_{2k}}{(-3)^k k} &= \left(-\frac{1}{3}\right)^n \sum_{j=1}^n \frac{1}{j^2 \binom{2j}{j}} \sum_{k=j}^n (-4)^k \binom{n}{k} \\ &= \left(-\frac{1}{3}\right)^n \sum_{j=1}^n \frac{1}{j^2 \binom{2j}{j}} \left((-3)^n - \sum_{k=0}^{j-1} (-4)^k \binom{n}{k} \right) \\ &= \sum_{j=1}^n \frac{1}{j^2 \binom{2j}{j}} \left(1 - \sum_{k=0}^{j-1} \frac{(-4)^k}{(-3)^n} \binom{n}{k} \right). \end{aligned} \quad (2.2)$$

It is obvious that for $0 \leq k \leq j-1$,

$$\lim_{n \rightarrow \infty} \frac{(-4)^k}{(-3)^n} \binom{n}{k} = 0,$$

and so

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{j-1} \frac{(-4)^k}{(-3)^n} \binom{n}{k} = 0. \quad (2.3)$$

Letting $n \rightarrow \infty$ on both sides of (2.2) and noting (2.3), we arrive at

$$\sum_{k=1}^{\infty} \frac{H_k - 2H_{2k}}{(-3)^k k} = \sum_{j=1}^{\infty} \frac{1}{j^2 \binom{2j}{j}}. \quad (2.4)$$

Then the proof of (1.2) follows from (1.1) and (2.4). □

3 Proof of Theorem 1.2

In order to prove Theorem 1.2, we need three preliminary results.

Lemma 3.1 *For any non-negative integer n , we have*

$$\binom{2n}{n} \sum_{k=1}^n \frac{1}{k^2 \binom{2k}{k}} = \binom{2n}{n} H_n^{(2)} + \sum_{k=1}^n \frac{-3 \left(\frac{k}{3}\right)^2 + 2}{k^2} \binom{2n}{n+k}. \quad (3.1)$$

Proof. An identity due to Mattarei and Tauraso [15, (22)] says

$$\binom{2n}{n} \sum_{k=1}^n \frac{t^k}{k^2 \binom{2k}{k}} = \binom{2n}{n} H_n^{(2)} + \sum_{k=1}^n \frac{v_k(t-2)}{k^2} \binom{2n}{n+k}, \quad (3.2)$$

where the Lucas sequences $\{v_k(t)\}_{k \geq 0}$ are defined by

$$v_0(t) = 2, \quad v_1(t) = t, \quad \text{and} \quad v_k(t) = tv_{k-1}(t) - v_{k-2}(t) \quad \text{for } k \geq 2.$$

It is not hard to check that

$$v_k(-1) = -3 \left(\frac{k}{3} \right)^2 + 2. \quad (3.3)$$

Letting $t = 1$ in (3.2) and noting (3.3), we reach (3.1). \square

Lemma 3.2 *For any prime $p \geq 5$, we have*

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k} H_k^{(2)} \equiv -\frac{1}{6} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \quad (3.4)$$

Proof. We begin with the following polynomial congruence [15, (37)]:

$$\sum_{k=1}^{p-1} t^{p-k} \binom{2k}{k} H_k^{(2)} \equiv -2t \sum_{k=1}^{p-1} \frac{u_k(2-t)}{k^2} \pmod{p}, \quad (3.5)$$

where the Lucas sequences $\{u_k(t)\}_{k \geq 0}$ are defined by

$$u_0(t) = 0, \quad u_1(t) = 1, \quad \text{and} \quad u_k(t) = tu_{k-1}(t) - u_{k-2}(t) \quad \text{for } k \geq 2.$$

It is trivial to check that

$$u_k(1) = \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{2}, \quad (3.6)$$

where $\lfloor x \rfloor$ denotes the integral part of real x . Letting $t = 1$ in (3.5) and using (3.6) gives

$$\sum_{k=1}^{p-1} \binom{2k}{k} H_k^{(2)} \equiv - \sum_{k=1}^{p-1} \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{k^2} \pmod{p}. \quad (3.7)$$

Note that

$$\begin{aligned}
& \sum_{k=1}^{p-1} \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{k^2} \\
&= 2 \left(\sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 1 \pmod{6}}} \frac{1}{k^2} + \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2 \pmod{6}}} \frac{1}{k^2} - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 4 \pmod{6}}} \frac{1}{k^2} - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 5 \pmod{6}}} \frac{1}{k^2} \right) \\
&= 2H_{p-1}^{(2)} - 2 \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 0 \pmod{3}}} \frac{1}{k^2} - 4 \left(\sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 4 \pmod{6}}} \frac{1}{k^2} + \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 5 \pmod{6}}} \frac{1}{k^2} \right) \\
&= 2H_{p-1}^{(2)} - \frac{2}{9} H_{\lfloor p/3 \rfloor}^{(2)} - \frac{1}{9} \left(\sum_{k=1}^{\lfloor \frac{p+1}{6} \rfloor} \frac{1}{(k-1/3)^2} + \sum_{k=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(k-1/6)^2} \right). \tag{3.8}
\end{aligned}$$

If $p \equiv 1 \pmod{6}$, then $\lfloor p/6 \rfloor \equiv -1/6 \pmod{p}$, and so

$$\begin{aligned}
& \sum_{k=1}^{\lfloor \frac{p+1}{6} \rfloor} \frac{1}{(k-1/3)^2} + \sum_{k=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(k-1/6)^2} \\
&\equiv \sum_{k=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(2\lfloor p/6 \rfloor + k)^2} + \sum_{k=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(\lfloor p/6 \rfloor + k)^2} \pmod{p} \\
&= H_{(p-1)/2}^{(2)} - H_{\lfloor p/6 \rfloor}^{(2)}. \tag{3.9}
\end{aligned}$$

It follows from (3.8) and (3.9) that

$$\sum_{k=1}^{p-1} \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{k^2} \equiv 2H_{p-1}^{(2)} - \frac{2}{9} H_{\lfloor p/3 \rfloor}^{(2)} - \frac{1}{9} H_{(p-1)/2}^{(2)} + \frac{1}{9} H_{\lfloor p/6 \rfloor}^{(2)} \pmod{p}. \tag{3.10}$$

By Wolstenholme's theorem [10, page 114], [19, Lemma 2.4], [12, (9)] and the congruence

below equation [12, (47)], we have

$$H_{p-1}^{(2)} \equiv 0 \pmod{p}, \quad (3.11)$$

$$H_{(p-1)/2}^{(2)} \equiv 0 \pmod{p}, \quad (3.12)$$

$$H_{\lfloor p/3 \rfloor}^{(2)} \equiv \frac{1}{2} \left(\frac{p}{3} \right) B_{p-2} \left(\frac{1}{3} \right) \pmod{p}, \quad (3.13)$$

$$H_{\lfloor p/6 \rfloor}^{(2)} \equiv 5H_{\lfloor p/3 \rfloor}^{(2)} \stackrel{(3.13)}{\equiv} \frac{5}{2} \left(\frac{p}{3} \right) B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \quad (3.14)$$

Substituting (3.11)–(3.14) into (3.10) gives

$$\sum_{k=1}^{p-1} \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{k^2} \equiv \frac{1}{6} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \quad (3.15)$$

If $p \equiv 5 \pmod{6}$, in a similar way, we can also prove that

$$\sum_{k=1}^{p-1} \frac{(-1)^{\lfloor \frac{k}{3} \rfloor} + (-1)^{\lfloor \frac{k-1}{3} \rfloor}}{k^2} \equiv \frac{1}{6} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \quad (3.16)$$

Finally, combining (3.7), (3.15) and (3.16), we arrive at

$$\sum_{k=1}^{p-1} \binom{2k}{k} H_k^{(2)} \equiv -\frac{1}{6} B_{p-2} \left(\frac{1}{3} \right) \pmod{p},$$

as desired. □

Lemma 3.3 *Let $p \geq 5$ be a prime. Then for $0 \leq d \leq \frac{p-1}{2}$, we have*

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k+d} \equiv \binom{p-d}{3} - \binom{d}{3} \pmod{p}. \quad (3.17)$$

Proof. From [26, Theorem 4.2], we deduce that for $0 \leq d \leq n$,

$$\sum_{k=0}^{n-1} \binom{2k}{k+d} = \sum_{k=0}^{n-d} \binom{n-d-k}{3} \binom{2n}{k}.$$

Letting $n = (p+1)/2$ in the above gives

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k+d} = \sum_{k=0}^{(p+1)/2-d} \binom{(p+1)/2-d-k}{3} \binom{p+1}{k}.$$

Since for $2 \leq k \leq (p+1)/2 - d$, we have

$$\binom{p+1}{k} \equiv 0 \pmod{p},$$

and so

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k+d} \equiv \binom{(p+1)/2-d}{3} + \binom{(p+1)/2-d-1}{3} \pmod{p}. \quad (3.18)$$

Next, we distinguish two cases $p \equiv 1 \pmod{6}$ and $p \equiv 5 \pmod{6}$ to prove (3.17). In both cases, we can show that

$$\binom{(p+1)/2-d}{3} + \binom{(p+1)/2-d-1}{3} \equiv \binom{p-d}{3} - \binom{d}{3} \pmod{p}. \quad (3.19)$$

Finally, combining (3.18) and (3.19), we complete the proof of (3.17). \square

Now we are ready to prove Theorem 1.2.

Proof of (1.3). Letting $n = \frac{p-1}{2}$ in (2.1) and noting that

$$\binom{(p-1)/2}{k} \equiv \frac{\binom{2k}{k}}{(-4)^k} \pmod{p} \quad \text{and} \quad (-3)^{(p-1)/2} \equiv \left(\frac{p}{3}\right) \pmod{p},$$

we obtain

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k} \sum_{j=1}^k \frac{1}{j^2 \binom{2j}{j}} \equiv \left(\frac{p}{3}\right) \sum_{k=1}^{(p-1)/2} \frac{H_k - 2H_{2k}}{(-3)^k k} \pmod{p}.$$

In order to prove (1.3), it suffices to show that

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k} \sum_{j=1}^k \frac{1}{j^2 \binom{2j}{j}} \equiv \frac{1}{6} B_{p-2} \left(\frac{1}{3}\right) \pmod{p}. \quad (3.20)$$

By (3.1), (3.4) and (3.17), we have

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \binom{2k}{k} \sum_{j=1}^k \frac{1}{j^2 \binom{2j}{j}} \\ & \equiv -\frac{1}{6} B_{p-2} \left(\frac{1}{3}\right) + \sum_{k=0}^{(p-1)/2} \sum_{j=1}^k \frac{-3 \left(\frac{j}{3}\right)^2 + 2}{j^2} \binom{2k}{k+j} \pmod{p} \\ & = -\frac{1}{6} B_{p-2} \left(\frac{1}{3}\right) + \sum_{j=1}^{(p-1)/2} \frac{-3 \left(\frac{j}{3}\right)^2 + 2}{j^2} \sum_{k=0}^{(p-1)/2} \binom{2k}{k+j} \\ & \equiv -\frac{1}{6} B_{p-2} \left(\frac{1}{3}\right) + \sum_{j=1}^{(p-1)/2} \frac{(-3 \left(\frac{j}{3}\right)^2 + 2) \left(\binom{p-j}{3} - \binom{j}{3}\right)}{j^2} \pmod{p}. \end{aligned} \quad (3.21)$$

If $p \equiv 1 \pmod{6}$, then $\lfloor p/6 \rfloor \equiv -1/6 \pmod{p}$, and so

$$\begin{aligned}
& \sum_{j=1}^{(p-1)/2} \frac{\left(-3 \left(\frac{j}{3}\right)^2 + 2\right) \left(\left(\frac{p-j}{3}\right) - \left(\frac{j}{3}\right)\right)}{j^2} \\
&= \sum_{\substack{1 \leq j \leq (p-1)/2 \\ j \equiv 1 \pmod{3}}} \frac{1}{j^2} + 2 \sum_{\substack{1 \leq j \leq (p-1)/2 \\ j \equiv 0 \pmod{3}}} \frac{1}{j^2} \\
&= H_{(p-1)/2}^{(2)} + \frac{1}{9} H_{\lfloor p/6 \rfloor}^{(2)} - \sum_{\substack{1 \leq j \leq (p-1)/2 \\ j \equiv 2 \pmod{3}}} \frac{1}{j^2}. \tag{3.22}
\end{aligned}$$

Note that

$$\begin{aligned}
\sum_{\substack{1 \leq j \leq (p-1)/2 \\ j \equiv 2 \pmod{3}}} \frac{1}{j^2} &= \frac{1}{9} \sum_{j=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(j - 1/3)^2} \\
&\equiv \frac{1}{9} \sum_{j=1}^{\lfloor \frac{p}{6} \rfloor} \frac{1}{(j + 2\lfloor p/6 \rfloor)^2} \pmod{p} \\
&= \frac{1}{9} \left(H_{(p-1)/2}^{(2)} - H_{\lfloor p/3 \rfloor}^{(2)} \right). \tag{3.23}
\end{aligned}$$

It follows from (3.12), (3.13), (3.14), (3.22) and (3.23) that

$$\begin{aligned}
\sum_{j=1}^{(p-1)/2} \frac{\left(-3 \left(\frac{j}{3}\right)^2 + 2\right) \left(\left(\frac{p-j}{3}\right) - \left(\frac{j}{3}\right)\right)}{j^2} &\equiv \frac{8}{9} H_{(p-1)/2}^{(2)} + \frac{1}{9} H_{\lfloor p/3 \rfloor}^{(2)} + \frac{1}{9} H_{\lfloor p/6 \rfloor}^{(2)} \\
&\equiv \frac{1}{3} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \tag{3.24}
\end{aligned}$$

If $p \equiv 5 \pmod{6}$, in a similar way, we can also prove that

$$\sum_{j=1}^{(p-1)/2} \frac{\left(-3 \left(\frac{j}{3}\right)^2 + 2\right) \left(\left(\frac{p-j}{3}\right) - \left(\frac{j}{3}\right)\right)}{j^2} \equiv \frac{1}{3} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}. \tag{3.25}$$

Finally, combining (3.21), (3.24) and (3.25), we reach (3.20). \square

4 Proof of Theorem 1.3

To prove Theorem 1.3, we require the following two identities.

Lemma 4.1 *For any non-negative integer n , we have*

$$\sum_{k=n}^{2n} (-1)^k \binom{k}{n} \binom{n}{k-n} H_k^{(2)} = 3 \sum_{k=1}^n \frac{1}{k^2 \binom{2k}{k}}, \quad (4.1)$$

$$\sum_{k=n}^{2n} (-1)^k \binom{k}{n} \binom{n}{k-n} H_k = 2H_n. \quad (4.2)$$

Proof. The identities (4.1) and (4.2) possess the same proofs as (2.1), and we omit the details. \square

Proof of (1.5). By (1.4), we have

$$\begin{aligned} \sum_{k=0}^{p-1} (-1)^k f_k H_k^{(2)} &= \sum_{k=0}^{p-1} (-1)^k H_k^{(2)} \sum_{j=0}^k \binom{k}{j} \binom{j}{k-j} \binom{2j}{j} \\ &= \sum_{j=0}^{p-1} \binom{2j}{j} \sum_{k=0}^{p-1} (-1)^k \binom{k}{j} \binom{j}{k-j} H_k^{(2)} \\ &\equiv \sum_{j=0}^{(p-1)/2} \binom{2j}{j} \sum_{k=0}^{p-1} (-1)^k \binom{k}{j} \binom{j}{k-j} H_k^{(2)} \pmod{p}, \end{aligned} \quad (4.3)$$

because $\binom{2j}{j} \equiv 0 \pmod{p}$ for $(p+1)/2 \leq j \leq p-1$.

Note that for $0 \leq j \leq (p-1)/2$,

$$\begin{aligned} \sum_{k=0}^{p-1} (-1)^k \binom{k}{j} \binom{j}{k-j} H_k^{(2)} &= \sum_{k=0}^{2j} (-1)^k \binom{k}{j} \binom{j}{k-j} H_k^{(2)} \\ &\stackrel{(4.1)}{=} 3 \sum_{k=1}^j \frac{1}{k^2 \binom{2k}{k}}. \end{aligned} \quad (4.4)$$

It follows from (4.3), (4.4) and (3.20) that

$$\begin{aligned} \sum_{k=0}^{p-1} (-1)^k f_k H_k^{(2)} &\equiv 3 \sum_{j=0}^{(p-1)/2} \binom{2j}{j} \sum_{k=1}^j \frac{1}{k^2 \binom{2k}{k}} \\ &\equiv \frac{1}{2} B_{p-2} \left(\frac{1}{3} \right) \pmod{p}, \end{aligned}$$

as desired. \square

Proof of (1.6). Similarly to the proof of (1.5), by using (4.2), we can show that

$$\sum_{k=0}^{p-1} (-1)^k f_k H_k \equiv 2 \sum_{j=0}^{(p-1)/2} \binom{2j}{j} H_j \pmod{p}.$$

Combining the above and the following congruence (see [14, page 528]):

$$\sum_{j=0}^{(p-1)/2} \binom{2j}{j} H_j \equiv -\left(\frac{p}{3}\right) q_p(3) \pmod{p},$$

we complete the proof of (1.6). □

Acknowledgments. This work was supported by the National Natural Science Foundation of China (grant 11801417).

References

- [1] D. Borwein and J.M. Borwein, On an intriguing integral and some series related to $\zeta(4)$, Proc. Amer. Math. Soc. 123 (1995), 1191–1198.
- [2] H.H. Chan, S.H. Chan and Z. Liu, Domb’s numbers and Ramanujan-Sato type series for $1/\pi$, Adv. Math. 186 (2004), 396–410.
- [3] H.H. Chan, J. Wan and W. Zudilin, Legendre polynomials and Ramanujan-type series for $1/\pi$, Israel J. Math. 194 (2013), 183–207.
- [4] J. Franel, On a question of Laisant, L’Intermédiaire des Mathématiciens 1 (1894), 45–47.
- [5] V.J.W. Guo, Proof of two conjectures of Sun on congruences for Franel numbers, Integral Transforms Spec. Funct. 24 (2013), 532–539.
- [6] V.J.W. Guo, Proof of a generalization of the (B.2) supercongruence of Van Hamme through a q -microscope, Adv. in Appl. Math., to appear.
- [7] V.J.W. Guo and M. J. Schlosser, Some new q -congruences for truncated basic hypergeometric series: even powers, Results Math. 75 (2020), Art. 1.
- [8] V.J.W. Guo and M.J. Schlosser, Some q -supercongruences from transformation formulas for basic hypergeometric series, Constr. Approx., to appear.
- [9] V.J.W. Guo and W. Zudilin, A q -microscope for supercongruences, Adv. Math. 346 (2019), 329–358.
- [10] G.H. Hardy and E.M. Wright, An Introduction to the Theory of Numbers, sixth ed., Oxford University Press, Oxford, 2008.
- [11] F. Jarvis and H.A. Verrill, Supercongruences for the Catalan-Larcombe-French numbers, Ramanujan J. 22 (2010), 171–186.
- [12] E. Lehmer, On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson, Ann. Math. 39 (1938), 350–360.
- [13] J.-C. Liu and F. Petrov, Congruences on sums of q -binomial coefficients, Adv. in Appl. Math. 116 (2020), 102003.

- [14] G.-S. Mao and Z.-W. Sun, Two congruences involving harmonic numbers with applications, *Int. J. Number Theory* 12 (2016), 527–539.
- [15] S. Mattarei and R. Tauraso, Congruences for central binomial sums and finite polylogarithms, *J. Number Theory* 133 (2013), 131–157.
- [16] A. van der Poorten, A proof that Euler missed...Apéry's proof of the irrationality of $\zeta(3)$, *Math. Intelligencer* 1 (1978/79), 195–203.
- [17] C. Schneider, Symbolic summation assists combinatorics, *Sém. Lothar. Combin.* 56 (2007), B56b, 36 pp.
- [18] V. Strehl, Binomial sums and identities, *Maple Technical Newsletter* 10 (1993), 37–49.
- [19] Z.-W. Sun, Super congruences and Euler numbers, *Sci. China Math.* 54 (2011), 2509–2535.
- [20] Z.-W. Sun, List of conjectural series for powers of π and other constants, preprint (2011), arXiv:1102.5649.
- [21] Z.-W. Sun, Arithmetic theory of harmonic numbers, *Proc. Amer. Math. Soc.* 140 (2012), 415–428.
- [22] Z.-W. Sun, Congruences for Franel numbers, *Adv. in Appl. Math.* 51 (2013), 524–535.
- [23] Z.-W. Sun, Connections between $p = x^2 + 3y^2$ and Franel numbers, *J. Number Theory* 133 (2013), 2914–2928.
- [24] Z.-W. Sun, Open conjectures on congruences, *J. Nanjing Univ. Math. Biquart.* 36 (2019), 1–99.
- [25] Z.-W. Sun, New series for powers of π and related congruences, preprint (2019), arXiv:1911.05456.
- [26] R. Tauraso, q -Analogues of some congruences involving Catalan numbers, *Adv. in Appl. Math.* 48 (2012), 603–614.
- [27] X. Wang and X. Yuan, Infinite summation formulas involving Riemann-Zeta function, preprint (2019), arXiv:1908.09468.
- [28] W. Zudilin, Ramanujan-type supercongruences, *J. Number Theory* 129 (2009), 1848–1857.