

Hurwitz trees and deformations of Artin-Schreier covers

Huy Dang

ABSTRACT

Let R be a complete discrete valuation ring of equal characteristic $p > 0$. In this paper, we first generalize the notion of Hurwitz trees in mixed characteristic to Artin-Schreier covers (\mathbb{Z}/p -covers in characteristic p) of a formal disc over R . It is a combinatorial-differential object with the shape of the dual graph of a cover's semi-stable model, and is endowed with essential degeneration data, measured by Kato's refined Swan conductors, of that cover. We then show how the existence of a deformation between two given Artin-Schreier covers is completely determined by the existence of certain Hurwitz trees. Finally, using the Hurwitz trees criteria, we improve a known result about the connectedness of the moduli space of Artin-Schreier curves of fixed genus.

Contents

1	Introduction	2
1.1	Structure	3
2	Artin-Schreier Covers	3
2.1	Artin-Schreier Theory	3
2.2	Deformations of Artin-Schreier covers	5
2.3	The moduli space of Artin-Schreier covers and their deformations . . .	6
2.4	Reduction to the local deformation problem	7
2.5	Birational deformation and the different criterion	8
3	Degeneration of \mathbb{Z}/p-covers	9
3.1	Discs and annuli	9
3.2	Semistable model and a partition of a disc	10
3.3	Reduction of covers	11
3.4	Refined Swan conductors	12
4	Hurwitz trees of Artin-Schreier covers	20
4.1	Hurwitz tree	20
4.2	The Hurwitz tree associated to a \mathbb{Z}/p -cover of a disc.	22
5	Construction of \mathbb{Z}/p-covers from a Hurwitz tree	24
5.1	Realization of a vertex	24
5.2	Realization of an edge	25
5.3	Realization of a leaf	25

2010 *Mathematics Subject Classification* 14H30

Keywords: Artin-Schreier theory, moduli space, refined Swan conductor, good reduction

The author is supported by NSF-DMS grants 1602054 and 1900396.

5.4	Gluing the boundaries together	26
5.5	Proof of Theorem 5.1	27
5.6	Hurwitz trees and deformations	28
6	Applications	30
6.1	Some general exact differential form results	30
6.2	Disconnectedness of \mathcal{AS}_g	33
	References	34

1. Introduction

Throughout this paper, we assume that k is an algebraically closed field of characteristic $p > 0$. We use the notation $\{ \dots \}$ to denote a multi-set. An *Artin-Schreier curve* is a smooth, projective, connected k -curve Y , which is a \mathbb{Z}/p -cover of the projective line \mathbb{P}_k^1 . The moduli space of Artin-Schreier k -curves of fixed genus g , which we denote by \mathcal{AS}_g , has the property that there are curves with different *branching data* (e.g., different number of branch points) which lie in the same connected component [PZ12]. That motivates the study of equal characteristic deformations of these covers. In [Dan20], by explicitly constructing some local deformations, it was shown that \mathcal{AS}_g is connected when g is sufficiently large.

THEOREM 1.1 cf. [Dan20, Theorem 1.1]. – *When $p = 3$, \mathcal{AS}_g is always connected.*
 – *When $p = 5$, \mathcal{AS}_g is connected for any $g \geq 14$ and $g = 0, 2$. It is disconnected if $g = 4, 6, 8$.*
 – *When $p > 5$, \mathcal{AS}_g is connected if $g \geq \frac{(p^3 - 2p^2 + p - 8)(p - 1)}{8}$ and $g \leq \frac{p - 1}{2}$. It is disconnected if $\frac{p - 1}{2} < g \leq \frac{(p - 1)^2}{2}$.*

Observe that, when $p = 5$, the only cases that the theorem does not cover are $g = 10$ and $g = 12$ (\mathcal{AS}_g is empty otherwise). The techniques from [Dan20] are inadequate to study these moduli spaces, as well as other cases. In this paper, with the aim to improve that result, we study local deformations in more detail using the notion of *Hurwitz tree*. A quick overview is given below.

Let R be a complete discrete valuation ring with characteristic $p > 0$ residue field. When R is of mixed characteristic (for example $R = W(k)$ where k is a perfect field of characteristic $p > 0$), a G -cover of the formal disc $\text{Spec } R[[X]]$ gives rise to a combinatorial object called a *Hurwitz tree*, which has the shape of the dual graph of the semistable model associated to the cover (see e.g., [BW09], [Hen00]). The existence of such a tree, along with some other conditions, is necessary for a cover in characteristic p to "lift" to characteristic 0. In particular, when $G \cong \mathbb{Z}/p$, Henrio proves that the lifts of a G -cover can be classified by Hurwitz trees of certain forms associated to the cover [Hen00]. Moreover, by generalizing Henrio's technique, Bouw and Wewers prove an analog of Henrio's lifting result for $G \cong \mathbb{Z}/p \rtimes \mathbb{Z}/m$ where m is prime to p [BW06], and show that all D_p -covers (where $p \neq 2$) lift.

In this paper, we generalize the notion of Hurwitz tree to the case where R is a complete discrete valuation ring of equal characteristic, specifically, $R = k[[t]]$, and $G \cong \mathbb{Z}/p$. The equal characteristic degeneration of \mathbb{Z}/p -covers was well-studied by Maugeais and Saïdi in [Mau03] and [Sai07]. Using their results together with Henrio's idea in [Hen00], we show that the existence of a flat deformation between Artin-Schreier covers of given branching data equates to the existence of a Hurwitz tree that satisfies certain criteria that are imposed by these data. Below is the main result of this paper.

THEOREM 1.2. *Suppose $d, d_1, \dots, d_r \not\equiv 0 \pmod{p}$ are integers such that $\sum_{i=1}^r (d_i + 1) = d + 1$.*

Fix a projective line $\mathbb{P}_{k[[t]]}^1$ and a k -point b of $\mathbb{P}_{k[[t]]}^1$. Then there exists a \mathbb{Z}/p -Galois cover of $\mathbb{P}_{k[[t]]}^1$ whose special fiber is an Artin-Schreier cover over k , branched only at b with ramification jump d and whose generic fiber is an Artin-Schreier cover of $\mathbb{P}_{k((t))}^1$ branched at r points that specialize to b and which have ramification jumps d_1, d_2, \dots, d_r if and only if there exists a Hurwitz tree of type $\{d_1 + 1, d_2 + 1, \dots, d_n + 1\}$.

Type, which will be defined in Definition 4.2, is a combinatorial invariant of a Hurwitz tree. Applying the Hurwitz tree criteria, we improve Theorem 1.1 as below.

THEOREM 1.3. – When $p = 5$, \mathcal{AS}_g is connected if and only if $g \geq 14$ or $g = 0, 2$.
 – When $p > 5$, \mathcal{AS}_g is disconnected if $(p - 1)/2 < g \leq (p - 1)(p - 2)$.

Remark 1.4. The cases that remain unknown are when $p > 5$ and $(p - 1)(p - 2) < g \leq (p^3 - 2p^2 + p - 8)(p - 1)/8$. Note that when p and g are fixed, one can apply Theorem 1.2 and the computation strategy from Section 6.2 to obtain a full pictures of how the irreducible components of \mathcal{AS}_g fit together, hence also determine the connectedness of \mathcal{AS}_g . Our current program and computers cannot handle the calculations when p and g are large, though.

1.1 Structure

Section 2 gives a quick overview of Artin-Schreier theory, the moduli space of Artin-Schreier covers of fixed genus, and how to partition the space by the branching data of the points. In the same section, we link the geometry of \mathcal{AS}_g with the deformation of \mathbb{Z}/p -covers (of genus g), and reduce Theorem 1.2 to a local one (Theorem 2.20). In §3, we examine the degeneration of étale \mathbb{Z}/p -torsors on a disc using the language of Kato’s refined Swan conductors. They are crucial to our approach, and distinguish this paper from [Dan20]. Section 4 introduces the notion of Hurwitz tree and describes how to derive such a tree from a given \mathbb{Z}/p -deformation (§4.2), thus proving the forward direction of Theorem 1.2. Section 5 considers the inverse process of §4.2, constructing a \mathbb{Z}/p -deformation from a given Hurwitz tree, hence completing the proof of Theorem 1.2. We then recover some known results about deformations using the new technique in §6.1. Finally, the proof of Theorem 1.3 is given in §6.2.1.

ACKNOWLEDGEMENTS

The author would like to express his great gratitude to his advisor, Andrew Obus, for his guidance and very careful reading of the draft of this paper. He thanks Florian Pop for useful suggestions and proofreading an earlier version of this paper. He thanks Gjergji Zaimi, Fedor Petrov, Felipe Voloch, and many people who joined the discussion on this thread <https://mathoverflow.net/questions>. Finally, he thanks the anonymous referees for their careful reading and their useful comments. This work is funded by the Simons Foundation Grant Targeted for Institute of Mathematics, Vietnam Academy of Science and Technology, and NSF-DMS grants 1602054 and 1900396.

2. Artin-Schreier Covers

2.1 Artin-Schreier Theory

Let K be a field of characteristic $p > 0$. If L is a separable extension of K of degree p , then the classical Artin-Schreier theory says that $L = K(\alpha)$ where α is a root of a polynomial equation $y^p - y = a$ for some $a \in K$ (see [Lan02, Section VI.6]).

Let Y be an Artin-Schreier k -curve. Then, by definition, there is a \mathbb{Z}/p -cover $\phi: Y \rightarrow \mathbb{P}_k^1$ with an affine equation of the form

$$y^p - y = f(x) \quad (1)$$

for some non-constant rational function $f(x) \in k(x)$. Equation (1) is called an *Artin-Schreier equation*. Furthermore, a cover $\phi' : Y' \rightarrow \mathbb{P}_k^1$ defined by $y^p - y = g(x)$ is isomorphic to ϕ if and only if $g(x) = f(x) + l(x)^p - l(x)$ for some $h(x) \in K(x)$. We call the map \wp sending $l(x) \mapsto l(x)^p - l(x)$ the *Artin-Schreier operator*. At each ramification point, there is a filtration of *ramification groups in upper numbering* [Ser79, IV]. In our case, as the inertia group is \mathbb{Z}/p , the filtration has only one jump, which we call the *ramification jump* at the corresponding branch point.

Example 2.1. Suppose $p = 5$. The cover of Y of \mathbb{P}_k^1 defined by

$$y^5 - y = \frac{1}{x^5} + \frac{1}{(x-1)^2}, \quad (2)$$

is an Artin-Schreier curve. Note that the term $1/x^5$ is a 5th-power. Hence, by the above discussion, one may add $\wp((-1/x)) = (-1/x)^5 - (-1/x)$ to the right-hand-side of (2). The result is an Artin-Schreier equation of the form

$$y^5 - y = -\frac{1}{x} + \frac{1}{(x-1)^2}. \quad (3)$$

Remark 2.2. We say the Artin-Schreier equation (3) has *reduced form*. That means the partial fraction decomposition of the right-hand-side of the equation only consists of terms of prime-to- p degree. When k is algebraically closed, any Artin-Schreier cover can be represented by a rational function $f(x)$ of reduced form.

The rational function $f(x)$ in (1) tells us everything about the ramification data of the cover ϕ . Suppose $f(x)$ has r poles $\mathcal{B} := \{B_1, \dots, B_r\}$ on \mathbb{P}_k^1 . Let d_j be the order of the pole of $f(x)$ at B_j . One may assume that $f(x)$ is in reduced form. Hence, the number d_j is prime to p . It is an easy exercise to show that \mathcal{B} is the collection of branch points of ϕ , and d_j is the ramification jump at B_j . We call $h_j := d_j + 1$ the *conductor* at B_j . Then $h_j \geq 2$ and $h_j \not\equiv 1 \pmod{p}$. Moreover, the ramification divisor of ϕ is

$$D := \sum_{j=1}^r (p-1)h_j Q_j \quad (4)$$

where Q_j is the ramification point above B_j ([Ser79], IV, Proposition 4). Hence, the degree of the different is $\sum_{j=1}^r (p-1)h_j$. Applying the Riemann-Hurwitz formula ([Har77], IV, Corollary 2.4), we obtain the following lemma.

LEMMA 2.3 [PZ12, Lemma 2.6]. *The genus of Y is $g_Y = ((\sum_{j=1}^r h_j) - 2)(p-1)/2$.*

DEFINITION 2.4. We call the $r \times 1$ matrix $[h_1, \dots, h_r]^T$ the *branching datum* of ϕ . For instance, the cover in Example 2.1 has branching datum $[2, 3]^T$. Throughout the paper, we define d by $g = d(p-1)/2$. So, we have the identities:

$$d = \left(\sum_{j=1}^r h_j \right) - 2 \text{ and } \sum_{j=1}^r h_j = d + 2 = 2g/(p-1) + 2.$$

Remark 2.5. The above lemma shows that all the Artin-Schreier k -curves with the same genus g have the same sum of conductors $d + 2$. That is the essential difference between an Artin-Schreier cover and a \mathbb{Z}/p -cover in characteristic different from p . For each \mathbb{Z}/p -tamely-ramified-cover, a branch point contributes $p - 1$ to the degree of its ramification divisor. We thus say a branch point of a wildly ramified cover contributes more to the degree of the different. Therefore, every \mathbb{Z}/p -tamely-ramified-cover of genus g must have the same number of branch points. It hence makes sense to group Artin-Schreier covers of the same genus by their branching data. This idea is utilized by Pries and Zhu in [PZ12], and will be discussed in the next section.

Remark 2.6. When the Galois group is \mathbb{Z}/p^n , the Artin-Schreier theory is generalized by the Artin-Schreier-Witt theory [Lor08, §26] [Lan02, §4]. It says that a \mathbb{Z}/p^n -cover of \mathbb{P}_k^1 is determined by some certain length- n -Witt-vector over K . We will discuss the deformations of this family of covers in a forthcoming paper.

2.2 Deformations of Artin-Schreier covers

Suppose $C \xrightarrow{\phi} \mathbb{P}_k^1$ is a G -Galois cover over k , where C is a smooth, projective, connected k -curve. Suppose, moreover, that A is a Noetherian, complete k -algebra with residue field k . Let

$$\text{Def}_\phi : \text{Alg}/k \rightarrow \text{Set}$$

be the functor which to any $A, f : A \rightarrow k \in \text{Alg}/k$ associates classes of G -Galois covers $\mathcal{C} \xrightarrow{\psi} \mathbb{P}_A^1$ that make the following cartesian diagram commute

$$\begin{array}{ccc} C & \longrightarrow & \mathcal{C} \\ \downarrow \phi & & \downarrow \psi \\ \mathbb{P}_k^1 & \longrightarrow & \mathbb{P}_A^1 \\ \downarrow & & \downarrow \\ \text{Spec } k & \xrightarrow{f} & \text{Spec } A, \end{array} \quad (5)$$

and so that the G -action on \mathcal{C} induces the original action on C . We say ψ is a *deformation* of ϕ over A , or ϕ is deformed (over A) to ψ . For more details, see [BM00, §2]. In this paper, we focus on the case where $G = \mathbb{Z}/p$ and $A = k[[t]]$.

Remark 2.7. One unique aspect of characteristic p is that there exist flat deformations of a wildly ramified cover over rings of equal characteristic so that the number of branch points changes, but the genus does not. That gives us a way to investigate a cover in characteristic p : finding a connection of it with a slightly different one via equal characteristic deformation. For example, using some (equal characteristic) deformations of \mathbb{Z}/p^n -covers [Pop14, Lemma 3.2], Pop reduced the lifting problem for cyclic groups (a.k.a., the *Oort conjecture*) to the case that had been solved by Obus and Wewers in [OW14]. Another perk of studying these deformations is understanding the geometry of the moduli space that parameterizes Galois covers. That will be discussed further in §2.3.

Remark 2.8. Suppose $\phi : Y \rightarrow \mathbb{P}_k^1$ is a \mathbb{Z}/p -cover that branched at r points $\{P_1, \dots, P_r\}$ with conductor h_i at P_i . Suppose, moreover, that ψ is a smooth deformation of ϕ over a complete discrete valuation ring of *equal characteristic* whose generic fiber ψ_η has branch locus $\{P_{1,1}, \dots, P_{1,m_1}, \dots, P_{r,1}, \dots, P_{r,m_r}\}$, where each $P_{i,j}$ reduces to P_i and has conductor $h_{i,j}$. Then it follows from the

discussion in Remark 2.5 that $\sum_{i=1}^r \sum_{j=1}^{m_i} h_{i,j} = \sum_{i=1}^r h_i$. We say the deformation ψ has *type*

$$[h_1, \dots, h_r]^\top \rightarrow [h_{1,1}, \dots, h_{1,m_1}, h_{2,1}, \dots, h_{r,1}, \dots, h_{r,m_r}]^\top.$$

2.3 The moduli space of Artin-Schreier covers and their deformations

In [PZ12], the authors introduce the moduli space of Artin-Schreier k -curves of genus g , which they denote by \mathcal{AS}_g . They enumerate a family of locally closed strata of \mathcal{AS}_g by partitions of the integer $d + 2$ such that no entries are congruent to 1 modulo p , i.e., all the possible branching data of Artin-Schreier curves of genus g . We call the collection of those partitions Ω_{d+2} . For instance, the partition $\vec{E} = \{h_1, \dots, h_r\}$ of $d + 2$ is associated with the stratum $\Gamma_{\vec{E}}$, which is the collection of all the points of \mathcal{AS}_g that represent Artin-Schreier curves with branching datum $[h_1, \dots, h_r]^\top$. We write $\vec{E}_1 \prec \vec{E}_2$ if the latter one is a refinement of the former one.

Example 2.9. Suppose $p = 5$ and $g = 14$. Then $d = 7$, and the strata of \mathcal{AS}_g correspond to the following partitions of $d + 2$: $\{9\}, \{7, 2\}, \{5, 4\}, \{5, 2, 2\}, \{4, 3, 2\}, \{3, 3, 3\}$, and $\{3, 2, 2, 2\}$.

The following result shows that one can relate the geometry of \mathcal{AS}_g with the existence of equal characteristic deformations between curves in different strata (hence have distinct branching data). We say there *always* exist deformations of type $[\vec{E}_1]^\top \rightarrow [\vec{E}_2]^\top$ if, given an arbitrary Artin-Schreier curve of branching datum $[\vec{E}_1]^\top$, we can deform it to one with branching datum $[\vec{E}_2]^\top$.

PROPOSITION 2.10 [Dan20, Proposition 3.4]. *Suppose \vec{E}_1 and \vec{E}_2 are two partitions of $d + 2$. Then the stratum $\Gamma_{\vec{E}_1}$ is contained in the closure of $\Gamma_{\vec{E}_2}$ if and only if there exists a deformation over $k[[t]]$ from a point in $\Gamma_{\vec{E}_1}$ to one in $\Gamma_{\vec{E}_2}$. In particular, there always exist deformations of type $[\vec{E}_1]^\top \rightarrow [\vec{E}_2]^\top$ if one can find a deformation of such type.*

Furthermore, the next result shows that the closure of a stratum is simply a union of strata.

PROPOSITION 2.11 [Dan20, Corollary 3.6]. *Let \vec{E}_1 and \vec{E}_2 be as in the previous proposition. Suppose $\Gamma_{\vec{E}_1}$ is not in the closure of $\Gamma_{\vec{E}_2}$. Then $\Gamma_{\vec{E}_1}$ is disjoint from the closure of $\Gamma_{\vec{E}_2}$.*


Therefore, understanding these deformations can give us a full picture of the moduli space \mathcal{AS}_g . We thus want to answer the following question.

QUESTION 2.12 *Deformation of Artin-Schreier covers problem.* Suppose we are given \vec{E}_1 and \vec{E}_2 in Ω_{d+2} . Does there exist a deformation over $k[[t]]$ of type $[\vec{E}_1]^\top \rightarrow [\vec{E}_2]^\top$?

2.3.1 The graph C_d Let us fix a prime p and construct a directed graph C_d (where d is as in Definition 2.4). The vertices of the graph correspond to the partitions \vec{E} in Ω_d . There is an arrow from \vec{E} to \vec{E}' if and only if $\vec{E} \prec \vec{E}'$, and $\Gamma_{\vec{E}}$ lies in the closure $\Gamma_{\vec{E}'}$.

In general topology, if one irreducible subset of a space lies in the closure of another, then they are contained in the same connected component of that space. Thus, if C_d is connected, then so is \mathcal{AS}_g . It is straightforward to check that the converse also holds. From now on, we will study the geometry of C_d .

Example 2.13. Suppose $p = 5$ and $g = 14$. Then $d = 7$. Using the information from [Dan20, Theorem 3.10], we draw a subgraph of C_7 in Figure 1. As the subgraph is connected, the graph C_7 is connected, and so is \mathcal{AS}_{14} .


 FIGURE 1. A subgraph of C_7

Remark 2.14. One can apply the results later in this paper (Proposition 5.14 and Proposition 6.2) to show that the above diagram is the complete C_7 . Thus, we can read from the graph that the irreducible components of \mathcal{AS}_{14} are the closures of the following strata: $\Gamma_{\{5,4\}}$, $\Gamma_{\{3,3,3\}}$, $\Gamma_{\{4,3,2\}}$, $\Gamma_{\{5,2,2\}}$, and $\Gamma_{\{3,2,2,2\}}$. Furthermore, the intersection of $\bar{\Gamma}_{\{3,3,3\}}$ and $\bar{\Gamma}_{\{3,2,2,2\}}$ is $\Gamma_{\{9\}}$, and the intersection of $\bar{\Gamma}_{\{4,3,2\}}$ and $\bar{\Gamma}_{\{5,2,2\}}$ is $\Gamma_{\{7,2\}}$ ($\bar{\Gamma}$ denotes the closure of the strata Γ).

2.4 Reduction to the local deformation problem

We first state a fact about germs of Artin-Schreier curves, which implies that, locally, they are easy to control.

PROPOSITION 2.15. *Suppose $\phi : Y \rightarrow \mathbb{P}_k^1$ is an Artin-Schreier cover and $P \in \mathbb{P}_k^1$ is a branch point of ϕ . Then the localization of ϕ at P is determined by the ramification jump at P .*

Proof. See Lemma 2.1.2 of [Pri03]. □

That means a \mathbb{Z}/p -cover of $\text{Spec } k[[x]]$ of conductor h , up to a change of the variable x , is isomorphic to one defined by

$$y^p - y = \frac{1}{x^{h-1}}.$$

The following local-global principle type result will help us to reduce our study of Artin-Schreier deformations to the local case.

PROPOSITION 2.16 c.f. [Dan20, Proposition 3.4]. *Suppose $\vec{E}_1 = \{h_1, h_2, \dots, h_n\}$ and \vec{E}_2 are in Ω_h . Then there exists a deformation of type $[\vec{E}_1]^\top \rightarrow [\vec{E}_2]^\top$ if and only if there exist n partitions $\{h_i\} \prec \vec{E}_{i,2} := \{h_{i,1}, \dots, h_{i,m_i}\} \subseteq \vec{E}_2$ ($1 \leq i \leq n$), where the $\vec{E}_{i,2}$'s partition \vec{E}_2 , and n deformations $\psi_1, \psi_2, \dots, \psi_n$ over $k[[t]]$, where ψ_i has type $[h_i] \rightarrow [h_{i,1}, \dots, h_{i,m_i}]^\top$. In particular, a deformation of this type exists only if $\vec{E}_1 \prec \vec{E}_2$.*

Example 2.17. Suppose $p = 5$. Then, by Proposition 2.16, there exists a deformation (over $k[[t]]$) of type $[7, 4]^\top \rightarrow [4, 3, 2, 2]^\top$ if and only if there exist either two deformations of types $[7] \rightarrow [3, 2, 2]^\top$ and $[4] \rightarrow [4]$ (trivial deformation), or two deformations of types $[7] \rightarrow [4, 3]^\top$ and $[4] \rightarrow [2, 2]^\top$.

Remark 2.18. It follows immediately from the above proposition that, if we have a deformation as in Remark 2.8, then $\sum_{j=1}^{m_i} h_{i,j} = h_i$ for all i .

Proposition 2.15 and Proposition 2.16 suggest that we may assume ϕ is a \mathbb{Z}/p -extension $k[[z]]/k[[x]]$ branched only at $x = 0$ with conductor h . Therefore, we may also think of a deformation ϕ over $k[[t]]$ as a \mathbb{Z}/p -cover $k[[t]][[Z]]/k[[t]][[X]]$. We thus reduce Question 2.12 to the following.

QUESTION 2.19 Local deformation of Artin-Schreier covers. Suppose $\{h\} \prec \{h_1, \dots, h_r\} \in \Omega_h$ and ϕ is a \mathbb{Z}/p -cover $k[[z]]/k[[x]]$ given by $y^p - y = \frac{1}{x^{h-1}}$. Define $R := k[[t]]$. Does there exist a deformation $R[[Z]]/R[[X]]$ of ϕ with generic branching datum $[h_1, \dots, h_r]^T$?

Hence, the question can be fully answered by the following local version of Theorem 1.2.

THEOREM 2.20. *Let $\phi : k[[z]] \rightarrow k[[x]]$ be a local G -cover with conductor d . Then there exists a deformation of ϕ over $k[[t]]$ of type $[d + 1] \rightarrow [d_1 + 1, \dots, d_r + 1]^T$ if and only if there exists a Hurwitz tree of type $\{d_1 + 1, \dots, d_r + 1\}$.*

2.5 Birational deformation and the different criterion

Usually, when dealing with Galois extensions of $k[[x]]$, it will be more convenient to deal with extensions of fraction fields than extensions of rings. So we will often want to think of a Galois ring extension in terms of the associated extension of fraction fields.

DEFINITION 2.21. Suppose $A/k[[x]]$ is a local G -extension. Suppose, moreover, that $M/\text{Frac}(R[[X]])$ where $R/k[[t]]$ finite, is a G -extension, and A_R is the integral closure of $R[[X]]$ in M . We say $M/\text{Frac}(R[[X]])$ is a *birational deformation* of $A/k[[x]]$ if

- (i) The integral closure of $A_R \otimes_R k$ is isomorphic to A , and
- (ii) The G -action on $\text{Frac}(A) = \text{Frac}(A_R \otimes_R k)$ induced from that on A_R restricts to the given G -action on A .

The following criterion is extremely useful for seeing when a birational deformation is actually a deformation (i.e., when $A_R \otimes_R k$ is already integrally closed, this isomorphic to A).

PROPOSITION 2.22 (The different criterion) [GM98, I,3.4]. *Suppose $A_R/R[[X]]$ is a birational deformation of the G -Galois extension $A/k[[x]]$. Let $K = \text{Frac } R$, let δ_η be the degree of the different of $(A_R \otimes_R K)(R[[X]] \otimes_R K)$, and let δ_s be the degree of the different of $A/k[[x]]$. Then $\delta_s \leq \delta_\eta$, and equality holds if and only if $A_R/R[[X]]$ is a deformation of $A/k[[x]]$.*

Example 2.23. Let $p = 5$, $g = 10$, hence, $d = 5$. Consider the \mathbb{Z}/p -cover ψ given by the normalization of $\mathbb{P}_{k[[t]]}^1$ over the extension of its fraction field defined by the following equation:

$$Y^5 - Y = \frac{-2X + t^5}{(-2)X^5(X - t^5)^2} = H(X, t). \quad (6)$$

The special fiber is birational to the $\mathbb{Z}/5$ -cover

$$y^5 - y = \frac{1}{x^6},$$

which is branched at 0 with conductor 7.

On the generic fiber, when $t \neq 0$, the partial fraction decomposition of $H(X, t)$ is of the form:

$$\frac{-1}{2t^5X^5} + \frac{1}{2t^{15}X^3} + \frac{1}{t^{20}X^2} + \frac{3}{2t^{25}X} + \frac{1}{2t^{20}(X - t^5)^2} - \frac{3}{2t^{25}(X - t^5)}.$$

Adding $\frac{1}{2t^5X^5} - \frac{1}{2tX}$ to $H(X, t)$, we get

$$\frac{-1}{2tX} + \frac{1}{2t^{15}X^3} + \frac{1}{t^{20}X^2} + \frac{3}{2t^{25}X} + \frac{1}{2t^{20}(X-t^5)^2} - \frac{3}{2t^{25}(X-t^5)},$$

which is a reduced form. Hence, the generic fiber is branched at two points $X = 0$ and $X = t$, which have conductors 4 and 3, respectively. It then follows from Proposition 2.22 that ψ is a deformation of type $[7] \rightarrow [4, 3]^\top$.

Example 2.24. One can check, in the same fashion as above, that the $\mathbb{Z}/5$ -cover given by

$$Y^5 - Y = \frac{-2X + t^{10}}{(-2)X^5(X-t^{10})^2(X-t^5)^5}$$

is a deformation of $y^5 - y = \frac{1}{x^4}$ over $k[[t]]$, where the generic branch points $0, t^{10}$, and t^5 have conductors 4, 3, and 5, respectively. Hence, it is a deformation of type $[12] \rightarrow [4, 3, 5]^\top$.

Remark 2.25. Suppose ϕ is a Galois cover over k . We can also apply the criterion to determine whether a deformation ψ of ϕ over a ring R of mixed characteristic (e.g., R is a finite extension of $W(k)$, where $W(k)$ is a Witt vector over k) is smooth. The lifting problem for Galois covers concerns the existence of a Galois cover in characteristic 0 that reduces to a given one in characteristic p . See [OW14, §4] for some explicit lifts of local Artin-Schreier covers and \mathbb{Z}/p^2 -covers. Some good expositions of the problem are [Obu12], [BW06], [OW14], and [Wea18]. Recall that the notion of Hurwitz tree, which we will discuss in §4, is first introduced in [Hen00] to tackle the lifting problem for Artin-Schreier covers. Theorem 1.2 is the exact equal characteristic analog of [Hen00, Théorème de réalisation].

3. Degeneration of \mathbb{Z}/p -covers

In this section, we study the degeneration of \mathbb{Z}/p -covers of $\text{Spec } R[[X]]$, where R is a complete discrete valuation of equal characteristic and with uniformizer π . Normalize the canonical valuation on R so that $v(\pi) = 1$. Set $K := \text{Frac } R$ and $\mathbb{K} := \text{Frac } R[[X]]$. We first introduce a geometric interpretation of $\text{Spec } R[[X]]$ using the language of non-archimedean geometry.

3.1 Discs and annuli

Firstly, we identify the K -analytic points of $\text{Spec } R[[X]]$ with

$$D = \{\mathbf{u} \in (\mathbb{A}_K^1)^{\text{an}} \mid v(\mathbf{u}) > 0\}$$

by plugging in $X = \mathbf{u}$. We call X a *parameter* for the *open unit disc* D with center 0.

Let $R\{X\} \subsetneq R[[X]]$ consist of power series for which the coefficients tend to 0, i.e.,

$$R\{X\} = \left\{ \sum_{i \geq m} a_i X^i \mid \lim_{i \rightarrow \infty} v(a_i) = \infty \right\}$$

As before, K -points of a *closed unit disc* $\text{Spec } R\{X\}$ can be identified with

$$\mathcal{D} = \{\mathbf{u} \in (\mathbb{A}_K^1)^{\text{an}} \mid v(\mathbf{u}) \geq 0\}.$$

The *boundary* of the open (or closed) unit disc is represented by the scheme $\text{Spec } R[[X^{-1}]]\{X\}$. Note that the ring $S := R[[X^{-1}]]\{X\}$ is a complete discrete valuation ring with residue field $\bar{S} = k((x))$, uniformizing element π and fraction field $S \otimes_R K$.

For $r \in \mathbb{Q}_{\geq 0}$, and $a \in K$ such that $|a| = r$, the open (resp. closed) disc of radius r is characterized by the scheme $\text{Spec } R[[a^{-1}X]]$ (resp. $\text{Spec } R\{a^{-1}X\}$). Its set of K -points is isomorphic to the open (resp. closed) unit disc under the map $X \mapsto a^{-1}X$. Denote by

$$\mathcal{D}[s, z] := \{u \in (\mathbb{A}_K^1)^{\text{an}} \mid v(u - z) \geq s\},$$

where $z \in (\mathbb{A}_K^1)^{\text{an}}$, the closed disc of radius p^{-s} centered at $X - z$, and $\mathcal{D}[s] := \mathcal{D}[s, 0]$. One can associate with $\mathcal{D}[s, z]$ the "Gauss valuation" $v_{s,z}$ that is defined by

$$v_{s,z}(f) = \inf_{a \in \mathcal{D}[s,z]} (v(f(a))),$$

for each $f \in \mathbb{K}^\times$. This is a discrete valuation on \mathbb{K} which extends the valuation v on K , and has the property $v_{s,z}(X - z) = s$. We denote by κ_s the residue field of \mathbb{K} with respect to the valuation $v_{s,z}$. That is the function field of the canonical reduction $\overline{\mathcal{D}}[s, z]$ of $\mathcal{D}[s, z]$. In fact, $\overline{\mathcal{D}}[s, z]$ is isomorphic to the affine line over k with function field $\kappa_{s,z} = k(x_{s,z})$, where $x_{s,z}$ is the image of $\pi^{-s}(X - z)$ in $\kappa_{s,z}$. For a closed point $\bar{x} \in \overline{\mathcal{D}}[s, z]$, we let $\text{ord}_{\bar{x}} : \kappa_{s,z}^\times \rightarrow \mathbb{Z}$ denote the normalized discrete valuation corresponding to the specialization of \bar{x} on $\overline{\mathcal{D}}[s, z]$. We let ord_∞ denote the unique normalized discrete valuation on $\kappa_{s,z}$ corresponding to the "point at infinity".

The *open annulus of thickness ϵ* is described by $\text{Spec } R[[X, U]] / (XU - a)$, where $a \in K$ is such that $v(a) = \epsilon$. The open annulus has two boundaries, one given by $\text{Spec } R[[X]][X^{-1}]$ and one given by $\text{Spec } R[[U]][U^{-1}]$. Note that two annuli over R are isomorphic if and only if they have the same thickness.

For $F \in \mathbb{K}$, $z \in (\mathbb{A}_K^1)^{\text{an}}$, and $s \in \mathbb{Q}_{\geq 0}$, we let $[F]_{s,z}$ denotes the image of $\pi^{-v_{s,z}(F)}F$ in the residue field $\kappa_{s,z}$.

3.2 Semistable model and a partition of a disc

Consider the open unit disc $D := \text{Spec } R[[X]]$, and suppose we are given $x_{1,K}, \dots, x_{r,K}$ in $D(K)$, with $r \geq 2$. We can think of $x_{1,K}, \dots, x_{r,K}$ as elements of the maximal ideal of R . Let D^{st} be a modification of D such that

- the exceptional divisor \overline{D} of the blow-up $D^{\text{st}} \rightarrow \text{Spec } R[[X]]$ is a semistable curve over k ,
- the fixed points $x_{b,K}$ specialize to pairwise distinct smooth points x_b on \overline{D} , and
- if \bar{x}_0 denotes the unique point on \overline{D} which lies in the closure of $D^{\text{st}} \otimes k \setminus \overline{D}$,

then $(\overline{D}; \bar{x}_0, (x_b))$ is stably marked. We call D^{st} the *semi-stable model of the marked disc* $(D; x_1, \dots, x_r)$, and $(\overline{D}; \bar{x}_0, (x_b))$ its *special fiber*. The dual graph of the special fiber is a tree whose *leaves* correspond to the marked points and whose *root* corresponds to \bar{x}_0 .

Example 3.1. Suppose we are given a $\mathbb{Z}/5$ -cover (in characteristic 5) of \mathbb{P}_k^1 that has four branch points $X = 0$, $X = t^5$, $X = t^5(1 + t^5)$, and $X = t^{10}$. The left graph of Figure 2 represents a semi-stable model of the open unit disc $D = \text{Spec } R[[X]]$ marked by the branch points of the cover. The tree on the right is its dual graph.

We associate with each edge an annulus. In this example, e_0 corresponds to the spectrum of $R[[X, X_1]] / (XX_1 - t^5)$ and e_1 resembles $\text{Spec } R[[X_1^{-1} - 1, V]] / (V(X_1^{-1} - 1) - t^5)$. We say e_0 has *thickness 1*, which is the thickness of the associated annulus divided by $p = 5$. Moreover, each vertex, which is not the root of the tree, is associated with a *punctured disc*. For instance, the vertex v_1 corresponds to $\text{Spec } R\{X_1^{-1}, X_1, (X_1^{-1} - 1)^{-1}\}$. That can also be thought of as the complement of the open discs $\text{Spec } R[[X_1]]$ and $\text{Spec } R[[X_1^{-1} - 1]]$ inside the closed disc $\text{Spec } R\{X_1^{-1}\}$. The root is linked with $\text{Spec } R\{X\}$. Finally, each leaf is associated with an open disc. To illustrate, one cor-

related with \bar{t}^5 is $\text{Spec } \mathbb{R}[[t^{-5}(X - t^5)]]$. Table 1 shows where the K -points of the disc specialize.

Note that, on the punctured disc corresponding to v_1 , the marked points t^5 and $t^5 + t^{10}$ (resp. 0 and t^{10}) specialize to $\bar{1}$ (resp. to $\bar{0}$). We call the directions on the dual graph from v_1 toward e_2, e_1 , and e_0 the *directions* corresponding to (or with respect to) $\bar{0}, \bar{1}$, and $\bar{\infty}$, respectively.




FIGURE 2. The special fiber \bar{D} and its dual graph

Subscheme \bar{V} of \bar{D}	Points of $D(K)$ that specialize to \bar{V}	Associated algebraic object
$\bar{\infty}$	$\{Y \mid 0 < v(Y) < 5\}$	$\mathbb{R}[[X, X_1]] / (XX_1 - t^5)$
$\bar{D}_3 \cap \bar{D}_1$	$\{Y \mid 5 < v(Y) < 10\}$	$\mathbb{R}[[X_1^{-1}, X_2]] / (X_1^{-1}X_2 - t^5)$
$\bar{D}_3 \setminus \bar{D}_1$	$\{Y \mid v(Y) \geq 10\}$	$\mathbb{R}\{X_2^{-1}\}$
$\bar{D}_2 \cap \bar{D}_1$	$\{Y \mid 5 < v(Y - t^5) < 10\}$	$\mathbb{R}[[X_1^{-1} - 1, V]] / (V(X_1^{-1} - 1) - t^5)$
$\bar{D}_2 \setminus \bar{D}_1$	$\{Y \mid v(Y - t^5) \geq 10 \wedge v(Y) = 5\}$	$\mathbb{R}\{V^{-1}\}$
$\bar{D}_1 \setminus (\bar{D}_3 \cup \bar{D}_2 \cup \{\bar{\infty}\})$	$\{Y \mid v(Y) = 5 \wedge v(Y - t^5) = 5\}$	$\mathbb{R}\{X_1^{-1}, X_1, (X_1^{-1} - 1)^{-1}\}$

TABLE 1. Partitions of $D(K)$

3.3 Reduction of covers

Let $\psi : \mathbb{R}[[Z]] \rightarrow \mathbb{R}[[X]]$ be a Galois cover. After enlarging our ground field K , we may assume that ψ is *weakly unramified* with respect to the Gauss valuation v_0 (that maps X to 0), see [Epp73]. By definition, this means that for all extensions w of v_0 to the function field of $\mathbb{R}[[Z]]$, the ramification index $e(w/v_0)$ is equal to 1. It then follows that the special fiber $\text{Spec } \mathbb{R}[[Z]] \otimes_{\mathbb{R}} k$ is reduced.

DEFINITION 3.2. We say that the cover ψ has *étale reduction* if the induced map $\text{Spec } \mathbb{R}[[Z]] \otimes_{\mathbb{R}} k \xrightarrow{\phi} \text{Spec } \mathbb{R}[[X]] \otimes_{\mathbb{R}} k$ is generically étale.

DEFINITION 3.3. If ψ has étale reduction, we call ϕ the *reduction* of ψ , and ψ a *deformation* of ϕ over \mathbb{R} . We say ψ has *good reduction* if it has étale reduction, and ϕ is smooth.

Recall Proposition 2.15 shows that a local Artin-Schreier cover is determined by the conductor of its unique branch point. We therefore reformulate Question 2.19 as follows.

QUESTION 3.4. Suppose $\{h_1, \dots, h_r\} \in \Omega_h$. Does there exist a \mathbb{Z}/p -cover of $\text{Spec } \mathbb{R}[[X]]$ with good reduction that has generic branching datum $[h_1, \dots, h_r]^T$?

3.4 Refined Swan conductors

Suppose ψ is a cyclic cover of a closed disc $\text{Spec } R\{X\}$ with valuation ν_0 . After enlarging R , we may assume that ψ is weakly unramified with respect to ν_0 . The set of all branch points is called the *branch locus* of ψ and is denoted by $\mathbb{B}(\psi)$.

We define two invariants that measure the ramification of ψ with respect to the valuation ν_0 . The *depth* is

$$\delta(\psi) := \text{sw}(\psi)/p \in \mathbb{Q}_{\geq 0},$$

where $\text{sw}(\psi)$ is the Swan conductor defined identically to one for mixed characteristic [Wew14, Definition 3.3] of the character associated to ψ . The rational number $\delta(\psi)$ is equal to 0 if and only if ψ is unramified. If this is the case, then its reduction ϕ is well-defined. In particular, if ψ is of order p and $\delta(\psi) = 0$, then there exists $u \in \kappa \setminus \wp\kappa$ such that ϕ is defined by $y^p - y = u$. We call u the *reduction* of ψ . Recall that, by Artin-Schreier theory, u is unique up to adding an element of the form $\wp(a) = a^p - a$, where $a \in \kappa$. We say ψ is *radical* if $\delta(\psi) > 0$.

Suppose $\delta(\psi) > 0$ and $\pi \in R$ with $\nu(\pi) = 1$. Then we can define the *differential Swan conductor* or *differential conductor* with respect to π

$$\omega(\psi) := \text{dsw}(\psi) \in \Omega_{\kappa}^1$$

in the same way as [Wew14, Definition 3.9] (for mixed characteristic). It is derived from the *refined Swan conductor* $\text{rsw}(\psi)$ and depends on the choice of π . In particular, we have the relation

$$\text{rsw}(\psi) = \pi^{-\text{sw}(\psi)} \otimes \text{dsw}(\psi) \in \mathfrak{m}^{-\text{sw}(\psi)} \otimes_{\mathcal{O}_{\kappa}} \Omega_{\kappa}^1, \quad (7)$$

where $\mathfrak{m} = (\pi)$. Note that $\text{rsw}(\psi)$ does not depend on the choice of π . In this paper, we always pick $\pi = t$.

We call $\delta(\psi|_{\mathcal{D}[s,z]})$ (resp. $\omega(\psi|_{\mathcal{D}[s,z]})$) the depth Swan conductor (resp. the differential Swan conductor) at the *place* $[s, z]$, or just at the place s when $z = 0$. We call the ordered pair $(\delta(\psi), \omega(\psi))$ when $\delta(\psi) > 0$, or $(\delta(\psi), u)$ when $\delta(\psi) = 0$ and u the reduction of ψ , the *degeneration type* of ψ . We usually refer to $\delta(\psi)$ and $\omega(\psi)$ as the *refined Swan conductors* of ψ .

Suppose $\bar{x} \in \bar{\mathcal{D}}$ or $\bar{x} = \infty$, and let $\text{ord}_{\bar{x}}: \kappa^{\times} \rightarrow \mathbb{Z}$ be a normalized discrete valuation whose restriction to κ is trivial. Then the composite of ν with $\text{ord}_{\bar{x}}$ is a valuation on K of rank 2, which we denote by $\mathbb{K}^{\times} \rightarrow \mathbb{Q} \times \mathbb{Z}$. In [Kat87], Kato defines a Swan conductor $\text{sw}_{\psi}^{\mathbb{K}}(\bar{x}) \in \mathbb{Q}_{\geq 0} \times \mathbb{Z}$. Its first component is equal to $\text{sw}(\psi)$. We define the *boundary Swan conductor* with respect to \bar{x}

$$\text{sw}_{\psi}(\bar{x}) \in \mathbb{Z}$$

as the second component of $\text{sw}_{\psi}^{\mathbb{K}}(\bar{x})$. Geometrically, it gives the instantaneous rate of change of the depth in the direction (mentioned in Example 3.1) corresponding to \bar{x} . See Remark 3.5 and §3.4.4 for further discussion.

Remark 3.5. The invariant $\text{sw}_{\psi}(\bar{x})$ is determined by $\delta(\psi)$ and $\omega(\psi)$ as follows:

- (i) If $\delta(\psi) = 0$, then

$$\text{sw}_{\psi}(\bar{x}) = \text{sw}_{\phi}(\bar{x})$$

where ϕ is the reduction of ψ and $\text{sw}_{\phi}(\bar{x})$ is the usual Swan conductor of ϕ with respect to the valuation $\text{ord}_{\bar{x}}$ [Ser79, IV, §2]. That follows immediately from the definitions. We thus have $\text{sw}_{\psi}(\bar{x}) \geq 0$ and $\text{sw}_{\psi}(\bar{x}) = 0$ if and only if ϕ is unramified with respect to $\text{ord}_{\bar{x}}$. Note that, if ϕ is a one point \mathbb{Z}/p -cover of $\text{Spec } k[[x]]$ with ramification jump d at $\bar{0}$, then $\text{sw}_{\phi}(\bar{0})$ is equal to d [Ser79, Cor. 2 to Th. 1]. Hence, a \mathbb{Z}/p -cover ψ of $R[[X]]$ is a deformation of ϕ

over R only if $\text{sw}_\psi(\bar{0}) = d$.

(ii) If $\delta(\psi) > 0$, then we have

$$\text{sw}_\psi(\bar{x}) = -\text{ord}_{\bar{x}}(\omega(\psi)) - 1$$

This follows from [Kat87, Corollary 4.6].

3.4.1 Vanishing cycle formula This section adapts [OW14, §5.3.3] and generalizes it to the equal characteristic case.

DEFINITION 3.6. A G -cover $\psi : Y \rightarrow C \cong \mathbb{P}_k^1 = \text{Proj } K[X]$ is called *admissible* if its branch locus $\mathbb{B}(\psi)$ is contained in the open disc D corresponding to $\text{Spec } R[[X]]$.

By §2.4, we can restrict our study to the case where ψ is admissible.

An affinoid subdomain $\mathcal{D} \subseteq C^{\text{an}}$ gives rise to a blowup $C'_R \rightarrow C_R$ with the following properties ([BL93]): C'_R is a semistable curve whose special fiber $\bar{C}' := C'_R \otimes_R k$ consists of two smooth irreducible components that meet in exactly one point. The first component is the strict transform of \bar{C} , which we may identify with \bar{C} . The second component is the exceptional divisor \bar{Z} of the blow-up $C'_R \rightarrow C_R$. It is isomorphic to the projective line over k and intersects \bar{C} in the distinguished point \bar{x}_0 . By construction, the complement $\bar{Z}^0 := \bar{Z} \setminus \{\bar{x}_0\}$ is identified with the canonical reduction $\bar{\mathcal{D}}$ of the affinoid \mathcal{D} . This means that the discrete valuation on K corresponding to the prime divisor $\bar{Z} \subset C'_R$ is equivalent to the Gauss valuation ν associated with \mathcal{D} (see. §3.2) and that its residue field κ may be identified with the function field of \bar{Z} .

Let Y'_R denote the normalization of C'_R in Y . We obtain the following commutative diagram

$$\begin{array}{ccc} Y'_R & \longrightarrow & Y_R \\ \downarrow & & \downarrow \\ C'_R & \longrightarrow & C_R \end{array}$$

in which the vertical maps are finite G -covers and each horizontal map is the composition of an a blowup with a normalization. Let $\bar{W} \subset Y'_R$ be the exceptional divisor of $Y'_R \rightarrow Y_R$. After enlarging the ground field K , we may assume that ψ is weakly unramified with respect to ν (the valuation corresponding to \mathcal{D}). Note that this holds if and only if \bar{W} is reduced. That condition allows us to define the refined Swan conductors for the restriction of ψ to \mathcal{D} as in §3.4. We now choose a closed point $\bar{x} \in \bar{Z}^0 = \bar{\mathcal{D}}$ and a point $\bar{y} \in \bar{W}$ lying over \bar{x} . We let

$$\mathcal{U}(\mathcal{D}, \bar{x}) :=]\bar{x}[_{\mathcal{D}}$$

denote the residue class of \bar{x} on the affinoid \mathcal{D} . Clearly, $\mathcal{U}(\mathcal{D}, \bar{x})$ is isomorphic to the open unit disc. Finally, we let $q : \widetilde{\bar{W}} \rightarrow \bar{W}$ denote the normalization of \bar{W} and set

$$\delta_{\bar{y}} := \dim_{\kappa}(q_* \mathcal{O}_{\widetilde{\bar{W}}} / \mathcal{O}_{\bar{W}})_{\bar{y}}.$$

Then $\delta_{\bar{y}} \geq 0$ and $\delta_{\bar{y}} = 0$ if and only if $\bar{y} \in \bar{W}$ is a smooth point.

The above notation extends to the case $\mathcal{D} = C$ as follows. If $\mathcal{D} = C$, then we let $\bar{Z} := \bar{C}$ denote the special fiber of the smooth model C_R of C and $\bar{W} := \bar{Y}$ the special fiber of Y . We set $\bar{x} := \bar{x}_0$ and choose an arbitrary point $\bar{y} \in \bar{W}$ above \bar{x}_0 . The residue class $\mathcal{U}(\mathcal{D}, \bar{x})$ is now equal to the open disc D , and the invariant $\delta_{\bar{y}}$ is defined in the same way as for $\mathcal{D} \subsetneq C$.

DEFINITION 3.7. Suppose ψ is a \mathbb{Z}/p of C , $\mathcal{D} \subseteq C$ is an affinoid, and $\bar{x} \in \overline{\mathcal{D}}$. We define

$$\mathfrak{C}(\mathcal{D}, \psi, \bar{x}) := \left\{ \sum_z h_z \mid z \in \mathbb{B}(\psi) \cap \mathcal{U}(\mathcal{D}, \bar{x}) \right\}.$$

where $\mathbb{B}(\psi)$ is the set of branch points of ψ , and h_z is the conductor (which is the ramification jump plus one) of the branch point z .

PROPOSITION 3.8. *With the notation as above, we have*

$$\text{sw}_{\psi|_{\mathcal{D}}}(\bar{x}) = \mathfrak{C}(\mathcal{D}, \psi, \bar{x}) - 1 - \frac{2\delta_{\bar{y}}}{p-1}.$$

Proof. The proposition is the equal characteristic analog of [OW16, Proposition 2.14], where everything but the characteristic of R is identical. One thus can imitate the proof of [OW16, Proposition 2.7] (which gives a more general version of [OW16, Proposition 2.14]) when $n = 1$. The only difference is $|\mathbb{B} \cap \mathcal{U}(\bar{x})|$ in Obus and Wewers' formula is the integer $\mathfrak{C}(\mathcal{D}, \psi, \bar{x})$ in our situation, as they are both equal to $\sum_{y \in \text{Ram}(f) \cap V} R_y$ in [CTT16, Theorem 6.2.3], which translates to $\sum_{w \in Y \mid \psi(w) \in \mathbb{B} \cap \mathcal{U}(\bar{x})} R_w$ in ours. That number R_w is, in turn, equal to the degree of the different of the corresponding local extension [CTT16, Theorem 4.6.4], hence is $(p-1)h_{\psi(w)}$ (recall that $h_{\psi(w)}$ is the conductor of the branch point $\psi(w)$). \square

We then obtain the following results.

COROLLARY 3.9. *Let ψ be an admissible cyclic cover of C of order p . Then*

$$\mathfrak{C}(\mathcal{D}, \psi, \bar{0}) = \sum_{z \in \mathbb{B}(\psi)} h_z \geq \text{sw}_{\psi|_{\mathcal{D}}}(\bar{0}) + 1.$$

Also, the cover ψ has good reduction if and only if $\delta(\psi) = 0$ and equality holds above.

Proof. The proof adapts the one for [OW14, Corollary 5.12]. The inequality in the first assertion follows immediately from Proposition 3.8 since $\mathbb{B}(\psi) \subsetneq \mathcal{D} = \mathcal{U}(\mathcal{D}[0], \bar{0})$ (because we assume ψ is admissible). In addition, the cover ψ has good reduction if and only if $\delta(\psi) = 0$ and $\bar{Y} = \overline{W}$ is smooth in any point \bar{y} above the distinguished point $\bar{0}$. The latter condition is equivalent to $\delta_{\bar{y}} = 0$. Thus, the rest also follows from Proposition 3.8. \square

REMARK 3.10. In the situation of Corollary 3.9, when ψ has good reduction, the reduction ϕ of its restriction to \mathcal{D} is a one-point-cover of $k[[t]]$ of conductor $\mathfrak{C}(\mathcal{D}, \psi, \bar{0})$. That is because $\text{sw}_{\psi|_{\mathcal{D}}}(\bar{0})$ coincides with $\text{sw}_{\phi}(\bar{0})$ (Remark 3.5 (i)), and the latter is equal to the ramification jump of ϕ at $\bar{0}$. Corollary 3.9 is thus equivalent to the different criterion (Proposition 2.22). We will utilize the former for the rest of the paper.

COROLLARY 3.11. *Let ψ be an admissible cyclic cover of C of order p , let $\mathcal{D} \subseteq C$ be an affinoid, and let \bar{x} be a point on the canonical reduction $\overline{\mathcal{D}}$ of \mathcal{D} . Then*

$$\text{sw}_{\psi|_{\mathcal{D}}}(\bar{x}) \leq \mathfrak{C}(\mathcal{D}, \psi, \bar{x}) - 1.$$

Moreover, if ψ has good reduction, then equality holds.

Proof. Parallel to the proof of [OW14, Corollary 5.13]. \square

The following result follows immediately from Corollary 3.11 and Remark 3.5.

COROLLARY 3.12. *In the same situation of Corollary 3.11, if $\delta(\psi|_{\mathcal{D}}) > 0$ and $\bar{x} \neq \infty$, we have*

$$\text{ord}_{\bar{x}}(\omega(\psi|_{\mathcal{D}})) \geq -\mathfrak{C}(\mathcal{D}, \psi, \bar{x})$$

with equality if ψ has good reduction.

Remark 3.13. Corollary 3.9 and Corollary 3.12 indicate that, for ψ to have good reduction, $\omega(\psi|_{\mathcal{D}})$ cannot have any zeros besides one at infinity for any affinoid $\mathcal{D} \subseteq C$.

Remark 3.14. The results in this section easily generalize to all cyclic covers. We will discuss it in the "forthcoming paper" mentioned in Remark 2.6

3.4.2 Refined Swan conductor of Artin-Schreier extensions In this section, we study the depth and differential Swan conductors of an Artin-Schreier cover of a closed disc $\text{Spec } R\{X\}$. Suppose ψ is a \mathbb{Z}/p -cover of the disc defined by

$$Y^p - Y = u,$$

where $u \in \mathbb{K} \setminus \wp\mathbb{K}$.

DEFINITION 3.15. Such an $u \in \mathbb{K} \setminus \mathcal{O}_{\mathbb{K}}$ is said to be *reduced* if $u = \tau\omega$, with $\tau \in \mathbb{K}$, $v(\tau) < 0$, $\omega \in \mathcal{O}_{\mathbb{K}}^{\times}$, and $\bar{\omega} \notin \kappa^p$. We say that u is *reducible* if there exists an $a \in \mathbb{K}^{\times}$ such that $u + a^p - a$ is reduced.

Remark 3.16. In our situation, as we can always replace R by its finite extensions, we may assume that u is reducible.

PROPOSITION 3.17. *Suppose ψ is a \mathbb{Z}/p -cover of $R\{X\}$ defined by*

$$Y^p - Y = u, \tag{8}$$

where $u \in \mathbb{K}$ is reduced when $v(u) < 0$. If $u = \pi^{-n}\omega$ (π is a fixed uniformizer of R), with $\omega \in \mathcal{O}_{\mathbb{K}}^{\times}$, $\bar{\omega} \notin \kappa^p$, and $n \in \mathbb{Q}_{\geq 0}$, then the depth of ψ is n/p and

- (i) *when $n = 0$, the degeneration is $\bar{\omega}$,*
- (ii) *when $n > 0$, its differential Swan conductor is $d\bar{\omega}$.*

Proof. The case $n = 0$ is immediate from the definition of the refined Swan conductors. When $n > 0$, we first describe a standard process to calculate $\text{rsw}(\psi)$ (see (7)).

PROPOSITION 3.18. *Suppose L/K , which we denote by ψ , is a $\mathbb{Z}/p = \langle \sigma \rangle$ -extension of complete discrete valuation field whose residue extension \bar{L}/\bar{K} is purely inseparable. Fix an element $x \in \mathcal{O}_{\bar{L}}^{\times}$ with $\bar{L} = \bar{K}[\bar{x}]$ and an element $\pi_K \in K$ with valuation one. Set $y := N_{L/K}(x)$ (hence $\bar{y} = \bar{x}^p \in \bar{K} \setminus \bar{K}^p$), $a := \sigma(x)/x - 1$ and $b := N_{L/K}(a)$. Then the refined Swan conductor of ψ with respect to the valuation of K is*

$$\text{rsw}(\psi) = b^{-1} \otimes \frac{d\bar{y}}{\bar{y}} \in \mathfrak{m}_K^{-\text{sw}(\psi)} \otimes_{\mathcal{O}_K} \Omega_{\bar{K}}^1. \tag{9}$$

where $\mathfrak{m}_K = (\pi_K)$ is an ideal of \mathcal{O}_K .

Proof. By [Kat80, §3.3, Lemma 15], with the notation as above, we have

$$\{\psi, 1 - bc\}_K = \{i_1(\bar{c}), y\}_K = i_2\left(\bar{c} \frac{d\bar{y}}{\bar{y}}\right)$$

for all $c \in \mathcal{O}_K$, where $\{_, _ \}_K$, i_1 , and i_2 are defined in [Kat87, §3]. The second equality follows from the definition of i_2 . Equation (9) then follows from the definition of rsw and [Kat87, Theorem 3.6 (2)]. \square

Let us now go back to the situation of Proposition 3.17. One may rewrite (8) as

$$Z^p - \pi^{n(p-1)/p} Z = \omega$$

simply by setting $Z = Y\pi^{n/p}$. Hence $\overline{Z}^p = \overline{\omega}$. The assumption $\overline{\omega} \notin \kappa^p$ suggests that the residue field upstairs is $\kappa[\overline{Z}]$. Set $x := Z$. Then $a = \sigma(x)/x - 1 = \pi^{n/p}/Z = 1/Y$, $y = N_{\mathbb{L}/\mathbb{K}}(x) = \omega$, and $b = N_{\mathbb{L}/\mathbb{K}}(a) = \pi^n/\omega$. Apply Proposition 3.18, we obtain

$$\text{rsw}(\psi) = \frac{\omega}{\pi^n} \otimes \frac{d\overline{\omega}}{\overline{\omega}} = (\pi^n)^{-1} \otimes d\overline{\omega}.$$


The rest then follows from the definition of the conductors in §3.4. \square

Remark 3.19. The proposition implies, if ψ is an Artin-Schreier cover where $\delta(\psi) > 0$, then its differential conductor $\omega(\psi)$ is an exact differential form. That is no longer true, however, for a cyclic G -cover (for instance, $G = \mathbb{Z}/p^2$) in general. More details will also be given in the "forthcoming paper" to which Remark 2.6 refers.

Example 3.20. Recall that, in example 2.24, the cover ψ is defined by

$$Y^5 - Y = \frac{-2X + t^{10}}{(-2)X^5(X - t^{10})^2(X - t^5)^5}. \quad (10)$$

Fix a valuation v_0 such that $v_0(t) = 1$. As the generic fiber is branched at three points $X = 0, X = t^5$, and $X = t^{10}$, we follow §3.2 to construct the dual graph of the associated semistable model as follows.



We set the orientation of the graph to be the direction from its leaves to its root v_0 .

At the initial vertex of e_1 , which corresponds to a punctured disc of radius 5^{-2} defined by $\text{Spec } \mathbb{R}\{t^{-10}X, t^{10}X^{-1}, t^{10}(X - t^{10})^{-1}\}$, we set $X_2 := Xt^{-10}$. Replacing X by X_2 equates to looking at the restriction of the cover to the sub-disc $\mathcal{D}[10] = \text{Spec } \mathbb{R}\{t^{-10}X\}$. We rewrite (10) as

$$Y^5 - Y = \frac{-2X_2 + 1}{(-2)X_2^5(X_2 - 1)^2(X_2 t^5 - 1)^5 t^{5 \cdot 17}} \quad (11)$$

Let x_2 the image of X_2 in the residue field of $\mathcal{D}[10]$. The derivative of the reduction modulo 5 of $\frac{-2X_2 + 1}{(-2)X_2^5(X_2 - 1)^2(X_2 t^5 - 1)^5}$ is $\omega(2) = \frac{dx_2}{x_2^4(x_2 - 1)^3}$. Hence, it follows from Proposition 3.17 that $\omega(2)$, after replacing x_2 by x , is the differential Swan conductor of ψ restricting to $\mathcal{D}[10]$, and its depth conductor is 17. Its boundary Swan conductors are $\text{sw}_{\psi|_{\mathcal{D}[10]}}(\infty) = -\text{ord}_\infty(\omega(2)) - 1 = -6$, $\text{sw}_{\psi|_{\mathcal{D}[10]}}(x_2) = -\text{ord}_{x_2}(\omega(2)) - 1 = 3$, and $\text{sw}_{\psi|_{\mathcal{D}[10]}}(x_2 - 1) = -\text{ord}_{x_2 - 1}(\omega(2)) - 1 = 2$.

At the initial vertex of e_0 , which resembles the punctured $\text{Spec } \mathbb{R}\{t^{-5}X, t^5X^{-1}, t^5(X - t^5)^{-1}\}$. Define $X_1 := Xt^{-5}$. We then rewrite (10) as

$$Y^5 - Y = \frac{-2X_1 + t^5}{(-2)X_1^5(X_1 - t^5)^2(X_1 - 1)^5 t^{5 \cdot 11}}.$$

Apply the same argument as above, we see that the degeneration type of ψ on $\text{Spec } \mathbb{R}\{X_1\}$ is $(11, \frac{dx}{x^7(x-1)^5})$.


Finally, when $r = 0$, then $X_0 = X$, the depth is 0, and the reduction ϕ is defined by $y^5 - y = \frac{1}{x^4}$. Therefore, we have $\mathfrak{C}(\psi, \bar{0}) = 5 + 3 + 4 = 12 = \text{sw}_\phi(\bar{0}) + 1$. Hence, ψ has good reduction by Corollary 3.9.

In the next example, we will go into detail of describing how the degeneration data vary along the dual graph.

Example 3.21. Consider the $\mathbb{Z}/2$ -cover ψ of $\text{Spec } \mathbb{R}\{X\}$ that is given by

$$Y^2 - Y = \frac{1}{X(X-t^2)} := f(X, t). \quad (12)$$

The associated dual graph of the semistable model is as follows.



Recall from §3.2 that the vertex v_1 represents the subdisc $\mathcal{D}[2] := \text{Spec } \mathbb{R}\{t^{-2}X\}$. Substitute $t^{-2}X$ by X_1 in (12), we obtain

$$f(X_1, t) = t^{-2 \cdot 2} \frac{1}{X_1(X_1 - 1)}.$$

As $1/(x_1(x_1 - 1)) \notin (k(x_1)^\times)^2$, we obtain $\delta(\psi|_{\mathcal{D}[2]}) = 2$ and $\omega(\psi|_{\mathcal{D}[2]}) = \frac{dx}{x^2(x-1)^2}$.

Set $X_r := Xt^{-2r}$. If $r > 1$, we can write $f(X, t)$ as

$$f(X_r, t) = t^{-2(r+1)} \frac{1}{X_r(X_r t^{2r-2} - 1)}.$$

Again, as $\frac{-1}{x_r} \notin (k(x_r)^\times)^2$, Proposition 3.17 shows $\delta(\psi|_{\mathcal{D}[2r]}) = r + 1$ and $\omega(\psi|_{\mathcal{D}[2r]}) = \frac{dx}{x^2}$.

Suppose $r < 1$. Then $[f]_{2r}$ is always a square. Set $a := 1/X^2$. We get

$$f(X, t) + a^2 - a = \frac{X^2 + t^2X + t^2}{X^2(X - t^2)} =: g(X, t).$$


Moreover, we can rewrite $g(X, t)$ as

$$g(X_r, t) = t^{-4r} \frac{t^{2-2r} - t^2X_r + X_r^2 t^{2r}}{X_r^2(X_r - t^{2-2r})}.$$

Note that $2 - 2r < 2r$ for $r \in (1/2, 1]$ and $2 - 2r > 2r$ otherwise. Hence, one can describe $\delta(\psi|_{\mathcal{D}[2r]})$ in terms of r as follows,

$$\delta(\psi|_{\mathcal{D}[2r]}) = \begin{cases} r & 0 \leq r \leq \frac{1}{2}, \\ 3r - 1 & \frac{1}{2} \leq r \leq 1, \\ r + 1 & 1 \leq r. \end{cases}$$

Below is the graph of $\delta(\psi|_{\mathcal{D}[2r]})$ with respect to r . Note that it is a piecewise linear function with respect to r . This phenomenon will be discussed further in §3.4.4.


 FIGURE 3. Graph of $\delta(\psi|_{\mathcal{D}[2r]})$

Moreover, we can write down the equation of $\omega(\psi|_{\mathcal{D}[2r]})$ with respect to r as follows.

$$\omega(\psi|_{\mathcal{D}[2r]}) = \begin{cases} \frac{dx}{x^2} & 0 < r < \frac{1}{2}, \\ \frac{(1+x^2)dx}{x^4} & r = \frac{1}{2}, \\ \frac{dx}{x^4} & \frac{1}{2} < r < 1, \\ \frac{dx}{x^2(x-1)^2} & r = 1, \\ \frac{dx}{x^2} & 1 < r. \end{cases}$$

When $r = 0$, the reduction ϕ of ψ is an Artin-Schreier cover given by $y^2 - y = \frac{1}{x^2}$. As $1/x^2$ is a square, we may replace the right-hand-side of the previous equation by $1/x^2 + (1/x)^2 - (1/x) = 1/x$. Thus, by Corollary 3.9, ψ does not have good reduction as $\sum_{x \in B(\psi)} h_x = 2 + 2 > \text{sw}_\psi(\bar{0}) + 1 = \text{sw}_\phi(\bar{0}) + 1 = 1 + 1$.

Remark 3.22. For future reference, we list down some important degeneration data of the cover ψ in Example 3.21 here. The degeneration type of ψ (resp. $\psi|_{\mathcal{D}[2]}$) is $(0, \frac{1}{x})$ (resp. $(2, \frac{dx}{x^2(x-1)^2})$).

Remark 3.23. If $f(X, t)$ in (12) is replaced by $\frac{1}{x^2(x-t)^2}$, then the deformation is flat of type [4] $\rightarrow [2, 2]^\top$. This type of deformation will be discussed further in § 6.1.1.

In the next two sections, we will study the degeneration of \mathbb{Z}/p -covers with good reduction on sub-discs and sub-annuli of the formal disc.

3.4.3 Degeneration data on the boundary The following is a result by Saïdi, which characterizes \mathbb{Z}/p -covers of a boundary of a disc.

PROPOSITION 3.24 c.f. [Saï07, Proposition 2.3.1]. *Let $A := \mathbb{R}[[X^{-1}]]\{X\}$, and let $\psi : \text{Spec } B \rightarrow \text{Spec } A$ be a nontrivial Galois cover of degree p . Assume that the ramification index of the corresponding extension of discrete valuation rings equals 1. Then ψ is a torsor under a finite flat \mathbb{R} -group scheme $G_{\mathbb{R}}$ of rank p and the following cases occur:*

- (i) *Suppose the depth is $\delta = 0$. Then ψ is a torsor under the étale group $(\mathbb{Z}/p\mathbb{Z})_{\mathbb{R}}$. Moreover, for a suitable choice of the parameter X of A , the torsor f is given by an equation $Y^p - Y = X^m$ for $m \in \mathbb{Z}$, prime to p . In addition, $Y^{1/m}$ is a parameter of B .*
- (ii) *Suppose the depth is $\delta > 0$. Then ψ is a torsor under the group scheme $\mathcal{M}_{\delta p, \mathbb{R}}$. Moreover, for a suitable choice of the parameter X , the torsor ψ is given by an equation $Y^p - Y = X^m/\pi^{p\delta}$ for $m \in \mathbb{Z}$, prime to p . In addition, $(Y\pi^\delta)^{1/m}$ is a parameter of B .*

The proposition thus shows, in the language of this paper, that an order p cover of $R[[X^{-1}]]\{X\}$ is determined by its depth δ and its boundary conductor $\text{sw}_{\psi|_{\mathcal{D}}}(\bar{0}) = m$ (where \mathcal{D} is the disc associated with $\text{Spec } R\{X\}$). We say it has *degeneration type* $(\delta, -m)$.

Remark 3.25. As $\text{Spec } A$ can be thought of as the boundary of the unit $\text{Spec } R[[X]]$, we will see in the next section that the number m may also be considered as the instantaneous rate of change of the depth on the centrifugal direction. In addition, it follows from Corollary 3.12 that, for ψ to be a restriction to a subdisc of a cover (of a disc) with good reduction, it is necessary that $m = -\mathfrak{C}(\mathcal{D}, \psi, \bar{0}) + 1 < 0$.

Remark 3.26. Saïdi's result is motivated by [Hen00, Corollaire 1.8] for the case R is of mixed characteristic. However, it is not true that a \mathbb{Z}/p^n -cover (where $n > 1$) of a boundary is determined by its depth and boundary conductor in both mixed and equal characteristic cases. See [HB09, §5.3] for a counterexample to the mixed characteristic case. We will discuss this phenomenon further in a forthcoming paper.

3.4.4 Good degeneration data on an annulus Fix a closed disc \mathcal{D} . For a K -point z on \mathcal{D} and $r \in \mathbb{Q}_{\geq 0}$, we denote by $\delta_{\psi}(r, z)$ (resp. $\delta_{\psi}(r)$) and $\omega_{\psi}(r, z)$ (resp. $\omega_{\psi}(r)$) the depth and the differential conductors of the restriction of ψ to $\mathcal{D}[r, z]$ (resp. $\mathcal{D}[r]$). When the point z is fixed, we may regard $\delta_{\psi}(r, z)$ and $\omega_{\psi}(r, z)$ as functions in terms of r . Suppose \bar{y} is a point on the canonical the reduction of $\mathcal{D}[r, z]$, or a point a infinity $\bar{y} = \infty$. Set $\text{sw}_{\psi}(r, z, \bar{y}) := \text{sw}_{\psi|_{\mathcal{D}[r, z]}}^K(\bar{y})$. The following results show how understanding the differential conductor can give a lot of information about the depth. They are the exact analogs of ones from [OW14, 5.3.2] in mixed characteristic.

PROPOSITION 3.27. *Suppose $z \in K$ is fixed. Then $\delta_{\psi}(_, z)$ extends to a continuous, piecewise linear function*

$$\delta_{\psi}(_, z) : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}.$$

Furthermore,

- (i) For $r \in \mathbb{Q}_{> 0}$, the left (resp. right) derivative of $\delta_{\psi}(_, z)$ at r is $-\text{sw}_{\psi}(r, z, \infty)$ (resp. $\text{sw}_{\psi}(r, z, \bar{0})$).
- (ii) If r is a kink of $\delta_{\psi}(_, z)$ (meaning the left and right derivatives do not agree), then $r \in \mathbb{Q}$.

Proof. See [Wew05, Proposition 2.9]. □

DEFINITION 3.28. Thanks to the above fact, it makes sense to define the refined Swan conductors of the restriction of ψ to an open disc $D[r, z]$ as $\delta(\psi|_{D[r, z]}) := \delta(\psi|_{\mathcal{D}[r, z]})$ and $\omega(\psi|_{D[r, z]}) := \omega(\psi|_{\mathcal{D}[r, z]})$. We set the boundary conductor of $\psi|_{D[r, z]}$ to be that of $\psi|_{\mathcal{D}[r, z]}$.

COROLLARY 3.29. *If $r \geq 0$ and $\delta_{\psi}(r, z) > 0$, then the left and right derivatives of δ_{ψ} at r are given by $\text{ord}_{\infty}(\omega_{\psi}(r)) + 1$ and $-\text{ord}_{\bar{0}}(\omega_{\psi}(r)) - 1$, respectively. In particular, the integer $-\text{ord}_{\bar{x}}(\omega_{\psi}(r, z)) - 1$ is the instantaneous rate of change of $\delta_{\psi}(r, z)$ on the direction with respect to \bar{x} (defined in Example 3.1).*

Proof. Immediate from Proposition 3.27 (i) and Remark 3.5 (ii). □

Remark 3.30. It follows from the above results and Corollary 3.11 that, if a cover of a disc has good reduction, then its depth varies from a point inside the disc to its boundary like a piecewise linear, weakly concave down function. Moreover, the rate of change of the depth at each place is determined by the sum of conductors of the branch points further away from the boundary (i.e. ones inside the corresponding closed disc). It is not the case in Example 3.21 because $\delta(\psi|_{\mathcal{D}[r]})$,

regarded as a function with respect to r , fails to concave down at $r = 1/2$ as showed in Figure 3.

We thus can describe the degeneration of a cover of an annulus as below.

PROPOSITION 3.31 c.f. [Sai07, Proposition 3.3.9]. *Suppose ψ is a \mathbb{Z}/p -cover of a unit disc D with good reduction. Let \mathcal{X} be an annulus of thickness $p\epsilon$ lying inside D that contains no branch points of ψ . Let S_1 and S_2 be the inside (one further from the boundary of D) and the outside boundary of \mathcal{X} , respectively. Then the reduction on S_1 is of type (d, q) (where $d \geq 0$) if and only if the degeneration type on S_2 is $(d - q\epsilon, -q)$.*

Proof. Let $S, U \in \text{Frac } R[[X]]$ be such that $\mathcal{X} \cong \text{Spec } R[[S, U]] / (SU - \pi^{p\epsilon})$. One may think of \mathcal{X} as the complement of $\text{Spec } R\{U^{-1}\}$ inside $\text{Spec } R[[S]]$. Hence $S_2 \cong \text{Spec } R\{S\}[[S^{-1}]]$ and $S_1 \cong \text{Spec } R\{U^{-1}\}[[U]]$. As in §3.2, we may associate with $\text{Spec } R[[S]]$ a unit disc D over K parameterized by S , and with $\text{Spec } R[[U^{-1}]]$ its sub-disc $D[p\epsilon]$. Set $\mathfrak{C} := \mathfrak{C}(D, \psi, \bar{0})$. For $r \in \mathbb{Q} \cap [0, p\epsilon]$, define $\delta(r) := \delta(\psi|_{\mathcal{D}[r]})$, $\omega(r) := \omega(\psi|_{\mathcal{D}[r]})$. Apply Corollary 3.12 and Corollary 3.29, we obtain

$$\text{ord}_{\bar{0}} \omega(r) = -\mathfrak{C} \text{ for } r \in \mathbb{Q} \cap [0, p\epsilon]. \quad (13)$$

Thus, it must also be true that $\text{ord}_{\infty} \omega(r) = \mathfrak{C} - 2$ for $r \in \mathbb{Q} \cap (0, p\epsilon]$. Therefore, the boundary Swan conductors of S_2 and S_1 are $q := -\mathfrak{C} + 1$ and $-q$, respectively. Finally, it follows from (13) and Proposition 3.27 that $\delta(_)$ is linear with slope $\mathfrak{C} - 1$ on $[0, p\epsilon]$. That requires the depth at the two boundaries to behave as in the statement. \square

Remark 3.32. [Sai07, Proposition 3.3.9] is actually much stronger than Proposition 3.31. It claims that the two boundaries' degeneration types completely determine such a cover of an annulus. However, the author does not give the proof but only cites [Sai04, Proposition 4.2.5], where the corresponding result in mixed characteristic is proved. We also do not fully understand that proof at the moment as the approach is quite different from the one above. Ours is still good enough for the purpose of this paper, though.

4. Hurwitz trees of Artin-Schreier covers

Suppose R is a complete discrete valuation ring whose residue field is of characteristic $p > 0$. When R is of mixed characteristic (e.g., $R = W(k)$), Henrio, Bouw, Wewers, and Brewis define a combinatorial object called *Hurwitz tree* from a G -cover of a formal disc over R (see [Hen00], [BW06], [BW09]). This construction gives an obstruction for the lifting of Galois covers in characteristic p . In this section, we introduce the notion of Hurwitz tree in equal characteristic, which basically formalizes what Maugeais and Saïdi did in [Mau03] and [Sai07], using the language from [Hen00] and [BW06]. Then, we will describe how a \mathbb{Z}/p -deformation gives rise to such a tree. Finally, we will provide an obstruction for the equal characteristic deformation of Artin-Schreier covers and compute the Hurwitz trees of the previous examples.

Throughout the rest of the paper, we assume $R = k[[t]]$, and $K := k((t))$ is the fraction field of R .

4.1 Hurwitz tree

The below definition and the two following paragraphs are almost identical to ones in [BW06, §3.1]. We recite here for the convenience of the reader.

DEFINITION 4.1. A *decorated tree* is given by the following data:

- a semistable curve C over k of genus 0,
- a family $(x_b)_{b \in B}$ of pairwise distinct smooth k -rational points of C , indexed by a finite nonempty set B ,
- a distinguished smooth k -rational point $x_0 \in C$, distinct from any of the point x_b .

We require that C is stably marked by the points $((x_b)_{b \in B}, x_0)$.

The *combinatorial tree* underlying a decorated tree C is the graph $T = (V, E)$, defined as follows. The vertex set V of T is the set of irreducible components of C , together with a distinguished element v_0 . The edge set E is the set of singular points of C , together with a distinguished elements e_0 . We write C_v for the component corresponding to a vertex $v \neq v_0$ and x_e for the singular point corresponding to an edge $e \neq e_0$. An edge e corresponding to a singular point x_e is adjacent to the vertices corresponding to the two components which intersect at x_e . The edge e_0 is adjacent to the root v_0 and the vertex v corresponding to the (unique) component C_v containing the distinguished point x_0 . For each edge $e \in E$, the *source* (resp. the *target*) of e is the unique vertex $s(e) \in V$ (resp. $t(e) \in V$) adjacent to e which lies in the direction of the root (resp. in the direction away from the root).

Note that, since $(C, (x_b), x_0)$ is stably marked of genus 0, the components C_v have genus zero, too, and the graph T is a tree. Moreover, we have $|B| \geq 1$. For a vertex $v \in V$, we write $U_v \subsetneq C_v$ for the complement in C_v of the set of singular and marked points.

DEFINITION 4.2. Let $\vec{E} = \{h_1, \dots, h_r\}$ an element of $\Omega_{\sum_{i=1}^r h_i}$ (defined in §2.3). A \mathbb{Z}/p -Hurwitz tree (or just *Hurwitz tree* in the context this paper) of type \vec{E} is given by the following data:

- A decorated tree $C = (C, (x_b), x_0)$ with underlying combinatorial tree $T = (V, E)$.
- For every $v \in V$, a rational number $\delta_v \geq 0$, called the *depth* of v .
- For each $v \in V \setminus \{v_0\}$, an exact differential form ω_v called the *differential conductor* at v .
- For every $e \in E$, a positive rational number ϵ_e , called the *thickness* of e .
- For every $b_i \in B = \{b_1, \dots, b_r\}$, the positive number h_i (from \vec{E}), called the *conductor* at b .
- For v_0 , a fraction $\frac{1}{x^d}$, where $d \not\equiv 0 \pmod{p}$, called the *degeneration* at v_0 .

These objects are required to satisfy the following conditions.

(H1) Let $v \in V$. We have $\delta_v \neq 0$ if $v \neq v_0$.

(H2) For each $v \in V \setminus \{v_0\}$, the differential form ω_v does not have zeros nor poles on $U_v \subsetneq C_v$.

(H3) For every edge $e \in E - \{e_0\}$, we have the equality

$$-\text{ord}_{x_e} \omega_{t(e)} - 1 = \text{ord}_{x_e} \omega_{s(e)} + 1 \not\equiv 0 \pmod{p}.$$

(H4) For v_0 , we have $d = \text{ord}_{x_{e_0}} \omega_{t(e_0)} + 1$.

(H5) For every edge $e \in E$, we have

$$\delta_{s(e)} + \epsilon_e d_e = \delta_{t(e)},$$

where

$$d_e := -\text{ord}_{x_e} \omega_{t(e)} - 1 \stackrel{\text{(H2)}}{=} \text{ord}_{x_e} \omega_{s(e)} + 1.$$

(H6) For $b \in B$, let C_v be the component containing the point x_b . Then the differential ω_v has a pole in x_b of order h_b .

For each $v \in V \setminus \{v_0\}$, we call (δ_v, ω_v) the *degeneration type* of v . For each $e \in E$, we call $(\delta_{s(e)}, -d_{s(e)})$ (resp. $(\delta_{t(e)}, d_{t(e)})$) the *initial degeneration type* (resp. the *final degeneration type*) of e . The integer $h := d + 1$ is called the *conductor* of the Hurwitz tree. The rational $\delta := \delta_{v_0}$ is the *depth*. We define the *height* of the tree to be the maximal number of edges on a direction from its root to its leaves.

One can easily obtain the following result. See [Hen00, §2] and [GM98] for the proofs of its mixed characteristic analogs.

LEMMA 4.3. *Let $(C, \omega_v, \delta_v, \epsilon_e, h_b, d)$ be a Hurwitz tree. Fix an edge $e \in E$ and let $C_e \subseteq C$ be the union of all components C_v corresponding to vertices v which are separated from the root v_0 by the edge e . Then:*

$$d_e = \sum_{\substack{b \in B \\ x_b \in C_e}} h_b - 1 > 0.$$

In particular, $h = \sum_{b \in B} h_b$.

4.2 The Hurwitz tree associated to a \mathbb{Z}/p -cover of a disc.

Let $D_K := \{X \mid |X|_K < 1\}$ be the rigid open unit disc over K . Suppose we are given a \mathbb{Z}/p -cover ψ of D_K , which is algebraically given by

$$\psi : R[[Z]] \rightarrow R[[X]]$$

with good reduction. Hence $\delta(\psi) = 0$. Obviously, ψ induces a \mathbb{Z}/p -cover of the boundary $\text{Spec } R[[X^{-1}]]\{X\}$ of the disc D_K . Let d be the boundary conductor of this cover. Following [BW09], we will now associate to ψ a Hurwitz tree.

Let $x_{b,K} \in D_K$ be the branch points of ψ , indexed by the finite set B . We assume that the points $x_{b,K}$ are all K -rational and the conductor at $x_{b,K}$ is h_b . By the different criterion (Proposition 2.22) and Proposition 3.24, we obtain $\sum_{b \in B} h_b = d + 1 =: h$. We assume that ψ has at least two branch points ($|B| \geq 2$), i.e. the deformation is non-trivial. Applying §3.2, one obtains D_R , the semistable model of the marked disc $(D_K, (x_{b,K})_{b \in B})$ with special fiber $(\overline{D}, (x_b), x_0)$. Note that it is a decorated tree, in the sense of Definition 4.1.

Let (V, E) be the combinatorial tree underlying $T := (\overline{D}, (x_b), x_0)$. For $v \in V$, let $U_v \subsetneq D_v$ be the complement of the singular and marked points and let $U_{v,K} \subsetneq D_K$ be the affinoid subdomain with reduction U_v . Let V_v (resp. $V_{v,K}$) denote the inverse image of U_v (resp. $U_{v,K}$) under ψ . By construction, $V_{v,K} \rightarrow U_{v,K}$ is a torsor under the constant K -group scheme G . We set $\delta_v \in \mathbb{Q}_{\geq 0}$ to be the depth conductor of the restriction of the cover to $V_{v,K}$. When $\psi|_{V_{v,K}}$ is radical, we set $\omega_v \in \Omega_{k(x)}^1(U_v)$ to be the differential conductor of the same restriction. When $\psi|_{V_{v,K}}$ is étale, which only happens when $v = v_0$, its reduction can be represented by a fraction $\frac{1}{x^d}$, where d is prime to p , by Proposition 2.15. We set the degeneration at v_0 to be that fraction. Finally, for $e \in E$ we let $A_e \subseteq D_K$ denote the subset of all points which specialize to the singular point $x_e \in \overline{D}$ corresponding to e . This is an open annulus. We define ϵ_e as the thickness of A_e divided by p , i.e., the positive rational number such that

$$A_e \cong \{x \mid |p|_K^{p\epsilon_e} < |x|_K < 1\}.$$

PROPOSITION 4.4. *The datum $(T, \omega_v, \delta_v, \epsilon_e, h_b, h - 1)$ defines a Hurwitz tree of type $\{h_1, \dots, h_r\}$. Moreover, h is the conductor, 0 is the depth.*

Proof. The proof is parallel to one for the mixed characteristic case [Hen00, §3.1]. The only

differences are: all differential conductors are exact, each branch point contributes more to the degree of the different (see Remark 2.5), and the depth can be arbitrarily large.

(H3) It follows from Proposition 3.31, as $-\text{ord}_{x_e} \omega_{t(e)} - 1$ is the inside boundary conductor of the associated annulus and $\text{ord}_{x_e} \omega_{t(e)} + 1$ is its outside boundary one.


(H5), (H4) Suppose e is an edge with initial depth $\delta_{s(e)}$, final depth $\delta_{t(e)}$, and thickness ϵ_e . Then, as the restriction of the cover to the corresponding annulus has good reduction, Proposition 3.31 shows that $\delta_{t(e)} = \delta_{s(e)} + \epsilon_e d_e$.

(H2), (H6) follow from Remark 3.13.

(H1) It follows immediately from the definition of the depth that $\delta_{v_0} \geq 0$. Moreover, by (H5), δ_v is a strictly increasing function, as v goes away from the root.

Finally, it is immediate from the construction and the good reduction assumption that the depth is 0 and the conductor is h , completing the proof. \square


Example 4.5. Let ψ be the $\mathbb{Z}/5$ -cover in Example 2.24. Then it follows from Example 3.20 that the Hurwitz tree associated with ψ has the following form.



At each vertex v of the tree, the first component of the pair is the depth conductor at the boundary of the disc associated to v . When the depth is positive, the second component is the differential Swan conductor. When the depth is 0, the second component represents the degeneration of the restriction. The rational number below each edge e is the thickness of the corresponding annulus divided by p . The integer on the right at each leaf denotes the conductor of the associated branch point, and inside $[\cdot]$ is the branch point. We often disregard this information as you can see in Example 4.6. One can easily read off from the leaves that ψ has type $[12] \rightarrow [5, 4, 3]^\top$.

By Proposition 4.4, the existence of a Hurwitz tree of type $\{h_1, \dots, h_r\}$ is necessary for the existence of a deformation of corresponding type. It gives us an obstruction for the deformation of \mathbb{Z}/p -covers, which we call the *Hurwitz tree obstruction*.

Example 4.6. Suppose again that $p = 5$. If there exists a deformation of type $[5] \rightarrow [3, 2]^\top$, then the associated Hurwitz tree must have the form below.



Thus, there exists an exact differential form $\omega = \frac{dx}{x^3(x-a)^2}$, where $a \neq 0$. A straightforward calculation shows that the residue of ω at 0 is $3/a^4$, which is a contradiction. Therefore, the Hurwitz tree obstruction does not vanish. Hence, there is no deformation of type $[5] \rightarrow [3, 2]^\top$.

PROPOSITION 4.7. *Suppose a \mathbb{Z}/p -cover ψ of a formal disc gives rise to a Hurwitz tree of type $\{h_1, \dots, h_r\}$ and with depth zero from the construction at the beginning of §4.2. Then ψ is a deformation of type $[h_1, \dots, h_r]^\top$.*

Proof. It follows from the depth zero assumption that ψ has étale reduction. Moreover, as the reduction has conductor $h = \sum_{i=1}^r h_i$ by Lemma 4.3, it is smooth by the different criterion (Proposition 2.22) or Corollary 3.9. \square

Example 4.8. The tree below arises from the deformation in Example 3.21 using the data from Remark 3.22.

$$\left(0, \frac{1}{x}\right) \xrightarrow{\frac{e_0}{1}} \left(2, \frac{dx}{x^2(x-1)^2}\right) \begin{array}{l} \nearrow 2[0] \\ \searrow 2[t^2] \end{array}$$

Note that $\left(0, \frac{1}{x}\right)$ (resp. $\left(2, \frac{dx}{x^2(x-1)^2}\right)$) is the degeneration type of ψ (resp. $\psi|_{\mathcal{D}[2]}$). It violates (H4) and (H5), as $2 = d \neq \text{ord}_{z_{e_0}} \omega_{t(e_0)} + 1 = 4$ and $2 = \delta_{t(e_0)} \neq \delta_{s(e_0)} + \epsilon_{e_0} d_{e_0} = 3$. Hence, it is not a flat deformation by Proposition 4.7.

5. Construction of \mathbb{Z}/p -covers from a Hurwitz tree

In the previous section, we have associated a Hurwitz tree to a \mathbb{Z}/p -cover $\psi : \text{Spec } R[[Z]] \rightarrow \text{Spec } R[[X]]$. The main result of this section is that this construction can be reversed.

THEOREM 5.1. *Every Hurwitz tree $\mathcal{T} = (T, \omega_v, \delta_v, \epsilon_e, h_b, h - 1)$ is associated to a \mathbb{Z}/p -cover ψ of $\text{Spec } R'[[X]]$ for some finite extension R' of R .*

The proof goes as follows. Using the geometry of the tree \mathcal{T} , we partition the open disc $\text{Spec } R[[X]]$ into sub-discs, sub-punctured-disc, and annuli as in §3.2. In sections 5.1, 5.2, 5.3, we construct explicitly a \mathbb{Z}/p -cover for each of these pieces that matches its associated datum on the tree. We then glue them together along their boundaries using the technique from §5.4. Finally, the details of the proof will be given in §5.5.

5.1 Realization of a vertex

Consider a vertex $v \neq v_0$ on a tree with datum $(\delta_v \in \mathbb{Q}_{>0}, \omega_v = df)$ where $f \in \text{Frac } k[x]$. Suppose ω_v has r poles $p_1, p_2, \dots, p_r \in \mathbb{A}_k^1$, and set $d_i := -\text{ord}_{p_i}(\omega) - 1$, $d := -\text{ord}_\infty(\omega) - 1$. Then it follows from (H2) that $d + 1 = \sum_{i=1}^r (d_i + 1)$. The points p_i (resp. the point ∞) correspond to the singular points x_{e_i} such that $s(e_i) = v$ (resp. to the unique singular point x_e with $t(e) = v$). As in the previous section, v corresponds to a complement of r open discs inside one closed disc

$$W_K \cong \{X \mid |X| \leq 1, |X - P_{i,K}| \geq 1, \}$$

where $P_{i,K}$ is a lift of p_i to K . W_K is the K -analytic space of $W = \text{Spec } R[X, (X - P_{i,K})^{-1}]_{i=1, \dots, r}$.

Define a \mathbb{Z}/p -cover $U \xrightarrow{\psi} W$ by

$$Y^p - Y = \frac{F}{t^{p\delta}}, \quad (14)$$

where F is an element of $R[X, (X - P_{i,K})^{-1}]_{i=1, \dots, r}$ that lifts f (hence only have the $P_{i,K}$'s as poles). Let $\text{Spec } S_i := R[[X - P_{i,K}]]\{(X - P_{i,K})^{-1}\}$ (resp. $\text{Spec } S_\infty := R[[X^{-1}]]\{X\}$) be the boundary of the missing open disc containing the point $P_{i,K}$ (resp. ∞). The first declaration of the below proposition follows easily from Proposition 3.17 and Remark 3.5.

PROPOSITION 5.2. *The cover ψ restricts to a \mathbb{Z}/p -cover on $\text{Spec } S_i$ (resp. on $\text{Spec } S_\infty$) which has*

boundary conductor d_i (resp. boundary conductor d) and depth δ . In fact, the covering space \mathcal{U} is itself a punctured disc.

Proof. Set $\mathcal{A} := \mathbb{R}\{X, (X - P_{i,K})^{-1}\}_{i=1,\dots,r}$ and $\mathcal{B} := \mathcal{A}[Z]/(Z^p - t^{(p-1)\delta}Z - F)$. The equation defining \mathcal{B} is exactly (14) with $Z = Y/t^\delta$. Hence, \mathcal{U} is the spectrum of \mathcal{B} . One may think of its reduction, $\overline{\psi} : \text{Spec } \overline{\mathcal{B}} \rightarrow \text{Spec } \overline{\mathcal{A}}$, as a degree- p -cover of $\mathbb{P}_k^1 \setminus \{\infty, p_i\}_{i=1,\dots,r}$ that is defined by the equation $\overline{Z}^p - f$. In addition, as $df = \omega_v$ has neither a zero nor a pole on $\mathbb{P}_k^1 \setminus \{\infty, p_i\}_{i=1,\dots,r}$ (by (H2)), the scheme $\text{Spec } \overline{\mathcal{B}}$ is a smooth superelliptic curve. Thus, we may write $\overline{\mathcal{B}} = k[v, (v - v_i)^{-1}]_{i=1,\dots,r}$. Therefore, there exists a V (resp. V_i) in \mathcal{B} that lifts v (resp. v_i) and so that $\mathcal{B} = \mathbb{R}\{V, (V - V_i)^{-1}\}_{i=1,\dots,r}$. \square

5.2 Realization of an edge

Suppose e is an edge of thickness ϵ with final degeneration type (δ_1, d_1) and initial degeneration type $(\delta_2, -d_2)$, where $d_1 = d_2 =: d$ (which is positive as discussed in §3.4.4) and $\delta_1 - d\epsilon = \delta_2$. The edge corresponds to an annulus \mathcal{X} of thickness $p\epsilon$, which can be identified with $\text{Spec } \mathbb{R}[[X, U]]/(XU - t^{p\epsilon})$. Consider a $G \cong \mathbb{Z}/p$ -cover of $\psi : \mathcal{Y} \rightarrow \mathcal{X}$ defined by

$$Y^p - Y = \frac{X^d}{t^{p\delta_1}}. \quad (15)$$

Call $S_1 := \text{Spec } \mathbb{R}[[X]][X^{-1}]$ the inner boundary of the annulus, and $S_2 := \text{Spec } \mathbb{R}[[U^{-1}]]\{U\}$ its outer boundary. Then it is immediate that the \mathbb{Z}/p -extension of S_1 has boundary conductor d and depth δ_1 . Replace X by $t^{p\epsilon}/U$ in (15), we get

$$Y^p - Y = \frac{1}{t^{p(\delta_1 - d\epsilon)}U^d}. \quad (16)$$

PROPOSITION 5.3. *The covering space \mathcal{Y} is itself an annulus and the followings hold.*

- (i) *The induced G -cover of S_1 has conductor d and depth δ_1 .*
- (ii) *The induced G -cover of S_2 has conductor $-d$ and depth $\delta_1 - d\epsilon$.*

Proof. Set $Y =: Y_1/t^{\delta_1}$ and $Y =: Y_2/t^{(\delta_1 - d\epsilon)}$. One may rewrite (15) (resp. (16)) as $Y_1^p - t^{(p-1)\delta_1}Y_1 = X^d$ (resp. $Y_2^p - t^{(p-1)(\delta_1 - d\epsilon)}Y_2 = 1/U^d$). Hence, the covering space of the inner disc $\text{Spec } \mathbb{R}[[X^{-1}]]$ (resp. the outer disc $\text{Spec } \mathbb{R}[[U]]$) is also a disc with parameter $Z_1 := Y_1^{1/d}$ (resp. $Z_2 := Y_2^{-1/d}$). A straightforward computation then confirms that \mathcal{Y} is the spectrum of the \mathbb{R} -algebra $\mathbb{R}[[Z_1, Z_2]]/(Z_1Z_2 - t^\epsilon)$. The rest then immediately follows from Proposition 3.24. \square

5.3 Realization of a leaf

Consider a leaf $b_i \in B$ on a tree with depth $\delta > 0$ and conductor $d_i \not\equiv 0 \pmod{p}$. It corresponds to a singular point x_i and can be associated with a closed disc

$$C_{i,K} = \{X \mid |X - x_i| \leq 1\}$$

which are K -points of $\text{Spec } \mathbb{R}\{X - x_i\}$. Let $S := \text{Spec } \mathbb{R}[[X - x_i]^{-1}]\{X - x_i\}$ be the boundary of $C_{i,K}$. Consider the \mathbb{Z}/p -cover ψ_i of $C_{i,K}$ given by the following equation

$$Y^p - Y = \frac{1}{t^{p\delta}(X - x_i)^{d_i}}.$$

Then, by applying Proposition 3.17, we obtain the following result.

PROPOSITION 5.4. *The cover ψ_i of $C_{i,K}$ has depth δ and boundary conductor d_i .*

5.4 Gluing the boundaries together

In this section, we glue covers of smaller pieces together to form one for a larger piece.

5.4.1 Filling in a punctured disc Suppose $\delta \in \mathbb{Q}_{>0}$, and d_1, d_2, \dots, d_r are not-divisible-by- p integers. Suppose $\psi_1, \psi_2, \dots, \psi_r$ are $G \cong \mathbb{Z}/p$ -covers of the open discs $\text{Spec } \mathbb{R}[[Y_1]], \dots, \text{Spec } \mathbb{R}[[Y_r]]$, respectively (whose covering spaces are not necessary discs). Suppose ψ_i has depth δ and boundary conductor d_i . It then follows from Proposition 3.24 that we may assume, after a change of variable, that ψ_i induces on $M_i := \text{Spec } \mathbb{R}[[Y_i^{-1}]]\{Y_i\}$ a \mathbb{Z}/p -cover

$$Y^p - Y = \frac{1}{t^{p\delta} Y_i^{d_i}}.$$

Suppose ψ is a \mathbb{Z}/p -cover of a closed punctured disc $\text{Spec } \mathbb{R}\{X, (X - \alpha_i)^{-1}\}_{i=1, \dots, r}$ as in §5.1. We may assume that the restriction of ψ to $S_i = \text{Spec } \mathbb{R}[[X - \alpha_i]]\{(X - \alpha_i)^{-1}\}$, after a change of variable, is given by

$$Y^p - Y = \frac{1}{t^{p\delta} (X - \alpha_i)^{d_i}}$$

(possible by Proposition 3.24). We define the \mathbb{R} -module isomorphism $\psi_i : \mathbb{R}[[X - \alpha_i]^{-1}]\{X - \alpha_i\} \rightarrow \mathbb{R}[[Y_i^{-1}]]\{Y_i\}$ by mapping $(X - \alpha_i)$ to Y_i . It is clear from the construction that ψ_i and ψ coincides on the glued boundary.

We would like to construct a \mathbb{Z}/p -cover of $\text{Spec } \mathbb{R}\{X\}$ whose restriction to the closed punctured disc W coincides with ψ , and whose restriction to each open disc $\text{Spec } \mathbb{R}[[X - \alpha_i]]$ is identical to ψ_i (when $\text{Spec } \mathbb{R}[[Y_i]]$ is identified with $\text{Spec } \mathbb{R}[[X - \alpha_i]]$). To do that, we fill in W by identifying the boundary of $\text{Spec } \mathbb{R}[[Y_i]]$ with S_i just like above.

The below lemma shows explicitly how to patch together the disks on the *bottom* and using the compatibilities on the top.

LEMMA 5.5 [Hen00, Lemma 3.7]. *Suppose the elements $\alpha_1, \dots, \alpha_r$ of \mathbb{R} are pairwise distinct modulo t . We denote, for each $1 \leq i \leq r$, α_i (resp. β_i) the canonical injection of the \mathbb{R} -algebras $\mathbb{R}\{X, (X - \alpha_j)^{-1}\}_{1 \leq j \leq r}$ (resp. $\mathbb{R}[[Y_i]]$) in $\mathbb{R}[[X - \alpha_i]]\{(X - \alpha_i)^{-1}\}$ (resp. $\mathbb{R}[[Y_i]]\{Y_i^{-1}\}$). Let $\psi_i : \mathbb{R}[[Y_i]]\{Y_i^{-1}\} \rightarrow \mathbb{R}[[X - \alpha_i]]\{(X - \alpha_i)^{-1}\}$ be an isomorphism of \mathbb{R} -algebras. If θ is the \mathbb{R} -module homomorphism*

$$\mathbb{R}\{X, (X - \alpha_j)^{-1}\}_{1 \leq j \leq r} \times \prod_{1 \leq i \leq r} \mathbb{R}[[Y_i]] \xrightarrow{\theta} \prod_{1 \leq i \leq r} \mathbb{R}[[X - \alpha_i]]\{(X - \alpha_i)^{-1}\}$$

$$\theta(f_0, f_1, \dots, f_r) = (\alpha_1(f_0) - \psi_1 \circ \beta_1(f_1), \dots, \alpha_r(f_0) - \psi_r \circ \beta_r(f_r)),$$

for $f_0 \in \mathbb{R}\{X, (X - \alpha_j)^{-1}\}_{1 \leq j \leq r}$ and $f_i \in \mathbb{R}[[Y_i]]$ for $1 \leq i \leq r$, then θ is surjective and its kernel N is an \mathbb{R} -algebra $\mathbb{R}\{Y_0\}$ as desired.

Recall that for each i , $\psi_i|_{M_i}$ and $\psi|_{S_i}$ coincides via the identification $X - \alpha_i \mapsto Y_i$. We thus can also glue together the covers ψ and ψ_i 's along the lifts of their boundaries M_i 's and S_i 's on the top (which are boundaries themselves) in a G -equivariant way to obtain a well-defined G -action on the whole lift. The result is a \mathbb{Z}/p -cover of $\mathbb{R}\{Y_0\}$ that restricts to ψ on $\text{Spec } \mathbb{R}\{X, (X - \alpha_i)^{-1}\}_{i=1, 2, \dots, r}$ and ψ_i on $\text{Spec } \mathbb{R}[[Y_i]]$ as desired. Note that, the preimage of $\mathbb{R}\{Y_0\}$ under ψ is not necessary a disc. To prove that the final cover has good reduction, we only need to show that it gives rise to a Hurwitz tree with depth zero (Proposition 4.7).

Remark 5.6. In [Hen00], Henrio pastes the discs and punctured discs on the *top* together G -equivariantly using his Lemma 3.7. We can do the exactly same thing as the covering space of

a punctured disc realized in §5.1 is also a punctured disc (by Proposition 5.2), and one may assume that the preimages of the ψ_i 's are likewise discs using induction.

5.4.2 Glueing a closed disc with an annulus Suppose we are given \mathbb{Z}/p -cover ψ of an annulus $\mathcal{X} = \text{Spec } \mathbb{R}[[X, U]] / (XU - t^{p \cdot e})$ as in §5.2 and a \mathbb{Z}/p -cover ψ' of $\text{Spec } \mathbb{R}\{Y\}$ whose restriction to $\text{Spec } \mathbb{R}[[Y^{-1}]]\{Y\}$ has depth δ_2 and conductor m_2 . As the previous subsection, we can use the following lemma to glue the boundary of the closed disc to the inside boundary of the annulus at the bottom to form an open disc.

LEMMA 5.7 [Hen00, Lemma 3.8]. *Suppose e is a strictly positive integer, β the canonical injection of $\mathbb{R}[[X, U]] / (XU - t^{p \cdot e})$ in $\mathbb{R}[[U]]\{U^{-1}\}$, α is the canonical injection of $\mathbb{R}\{Y\}$ to $\mathbb{R}[[Y^{-1}]]\{Y\}$ and ψ is an isomorphism of \mathbb{R} -algebras $\mathbb{R}[[X]]\{X^{-1}\}$ and $\mathbb{R}[[Y^{-1}]]\{Y\}$ (by mapping X to Y^{-1}). If θ is a homomorphism of \mathbb{R} -modules*

$$\mathbb{R}\{Y\} \times \frac{\mathbb{R}[[X, U]]}{XU - t^{p \cdot e}} \rightarrow \mathbb{R}[[Y^{-1}]]\{Y\}$$

defined by $\theta(f, g) := \alpha(f) - \psi \circ \beta(g)$, then θ is surjective, and its kernel \mathbb{N} is an algebra $\mathbb{R}[[Y_0]]$.

As before, by patching together the cover ψ and ψ' using the above gluing on the bottom, we also attain a \mathbb{Z}/p -cover of an open unit disc corresponding to $\text{Spec } \mathbb{R}[[U]]$ that restricts to ψ on $\text{Spec } \mathbb{R}[[X, U]] / (XU - t^{p \cdot e})$ and to ψ' on $\text{Spec } \mathbb{R}\{Y\}$.

Remark 5.8. The method of glueing covering spaces on the top (discussed in Remark 5.6) can also apply here as we show the preimage of \mathcal{X} is also an annulus (Proposition 5.3). As in the previous remark, we need the preimage of ψ' to be a closed disc.

5.5 Proof of Theorem 5.1

Let $\mathcal{T} = (C, \omega_v, \delta_v, \epsilon_e, h_b, h - 1)$ be a Hurwitz tree with conductor $h - 1$ and depth δ . We call a $G = \mathbb{Z}/p$ -cover of the open disc a *realization* of \mathcal{T} if \mathcal{T} is associated to this cover by the construction in §4.2. Theorem 5.1 claims that we can realize \mathcal{T} . We will prove this claim by induction on the height of the tree \mathcal{T} . Suppose first that the height of \mathcal{T} is one. Hence, the associated Hurwitz tree has the form like in Figure 4. Let e_0 be the trunk of \mathcal{T} , we set $v_1 = t(e_0)$. Then $C_0 := C_{v_1}$ is the unique component of C which contains the distinguished point $x_0 \in C$. Set $\delta_1 := \delta_{v_1}$, and suppose b_1, \dots, b_r are the leaves with source v , and the $x_i := x_{e_i}$ are the corresponding singular points associated to the leaves. Let $h_i := h_{b_i}$. By §5.3, there exists an open unit disc $C_{i,K}$ over some finite extension K of $k((t))$, together with a G -cover $\psi_{v_1, i}$, whose depth is δ_1 and whose boundary conductor is h_i . Let $\text{Spec } S_i$ be the boundary of $C_{i,K}$. By assumption, the differential form $\omega := \omega_{v_1}$ on C_0 has a zero of order $d_{t(e_0)} - 1$ at x_0 and poles of order $d_i + 1$ at x_i . Let $W_{v_1, K}$ be the punctured disc with the G -cover constructed in Section 5.1, starting from the datum (C_0, ω, δ_1) . By Proposition 3.24 and Section 5.2, we can identify the boundary of the missing open disc corresponding to the point x_i with $\text{Spec } S_i$ in a way which makes $\psi_{v_1, i}$ and ψ_{v_1} coincides on the boundary. We can now use Lemma 5.5 to patch together the punctured disc $W_{v_1, K}$ and the discs $C_{i,K}$, in a G -equivariant way. The result is a closed disc $C_{v_1, K} \cong \{C_{v_1} \mid |C_{v_1}| \leq 1\}$, together with a G -cover ψ_{v_1} . By Proposition 5.2 and the construction, the restriction of the cover to the boundary of $C_{v_1, K}$ has conductor $d_{t(e_0)}$ and depth δ_1 .

Let A_K be the open annulus with the G -cover constructed in Section 5.2, starting from the target datum $(\delta_1, d_{t(e_0)})$ and the initial datum $(\delta, -d_{s(e_0)} = -d)$. By §5.4.2, we can identify the boundary of $X_{v_1, K}$ with the "inner boundary" (associated with the target datum) of A_K , to-




FIGURE 4. A tree of height one

gether with the G -cover, along these boundaries. The result is a G -cover ψ of an open disc with boundary conductor d , depth δ . By construction ψ is a realization of the Hurwitz tree \mathcal{T} . That completes the base case of the induction.

Therefore, we may assume that we can realize all Hurwitz trees with lower height than C . Again, let e_0 be the trunk of \mathcal{T} , $v_1 = t(e_0)$, $\delta_1 := \delta_{v_1}$, and suppose e_1, \dots, e_r are the edges with source v , and $x_i := x_{e_i}$ the corresponding singular points. Let $d_i := d_{s(e_i)}$. Let $\mathcal{T}_i \subsetneq \mathcal{T}$ be the subtree of \mathcal{T} which contains the point x_i but not the component C_0 . It is clear that \mathcal{T}_i inherits from \mathcal{T} the structure of a Hurwitz tree, with conductor h_i , depth δ_1 , and height strictly less than \mathcal{T} . By our induction hypothesis, there exists an open unit disc $C_{i,K}$ over some finite inseparable extension K of $k((t))$, together with a G -cover $\psi_{v_1,i}$, whose associated Hurwitz tree is \mathcal{T}_i . We than can apply the same process as in the base case to construct a G -cover ψ that realizes the tree \mathcal{T} . This completes the proof of the theorem. \square

Remark 5.9. In [BW06], Bouw and Wewers generalize the notion of Hurwitz tree from [Hen00] to the case where $G = \mathbb{Z}/p \rtimes_{\chi} \mathbb{Z}/m$ (m is prime to p) by adding an extra piece of information, which they call the *tame inertia character*, to Henrio's tree. It comes from the group homomorphism $\chi : \mathbb{Z}/m \rightarrow \text{Aut}(\mathbb{Z}/p)$ that defines the semi-direct product. Furthermore, they prove an analog of [Hen00, Corollary 1.8] and Proposition 3.24, which says a G -action on a boundary of a disc is determined by its depth, its boundary conductor, and its tame inertia character [BW06, Proposition 2.3]. Using this fact, they utilize Henrio's technique to prove that every local D_p -cover in characteristic p lifts to characteristic zero [BW06, Theorem 4.4]. We would expect an analogous result for the equal characteristic case using a parallel theory.

5.6 Hurwitz trees and deformations

Combining Theorem 5.1 and Proposition 4.7, we acquire the following result.

COROLLARY 5.10. *Let $\phi : Z_k \rightarrow X_k$ be a local G -cover with conductor $h - 1$. Then there exists a deformation of ϕ over $k[[t]]$ of type $[h] \rightarrow [h_1, \dots, h_r]^{\top}$ if and only if there exists a Hurwitz tree of type $\{h_1, \dots, h_r\}$ and with depth zero.*

Remark 5.11. The only difference between Corollary 5.10 and Theorem 2.20 is that we still need the Hurwitz tree in the statement to have depth zero. That requirement will be abolished by Proposition 5.14.

DEFINITION 5.12. Suppose $\vec{A} \prec \vec{B}$. Then one can easily see that there exist a minimal submultiset \vec{A}' of \vec{A} and \vec{B}' of \vec{B} , such that $\vec{A}' \prec \vec{B}'$ and $\vec{A} \setminus \vec{A}'$ is the same as $\vec{B} \setminus \vec{B}'$. We call $\vec{A}' \prec \vec{B}'$ the *difference* of $\vec{A} \prec \vec{B}$.

Example 5.13. The difference of $\{3, 4, 5, 6\} \prec \{3, 2, 2, 3, 2, 6\}$ is $\{4, 5\} \prec \{2, 2, 3, 2\}$.

The next result shows that the existence of a Hurwitz tree equates to the existence of a chain of exact differential forms of certain type. This phenomenon does not generalize to any cyclic group G , though.

PROPOSITION 5.14. *There exists a Hurwitz tree of type $\{h_1, \dots, h_r\}$ if and only if there exists a chain of partitions of integers $\underline{e}_0 := \{e\} \prec \underline{e}_1 \prec \dots \prec \underline{e}_m := \{h_1, \dots, h_r\}$, where the difference between \underline{e}_{i-1} and \underline{e}_i is $\{l_i\} \prec \{l_{i,1}, \dots, l_{i,r_i}\}$, together with m exact differential forms*


$$\frac{dx}{\prod_{j=1}^{r_i} (x - P_{i,j})^{l_{i,j}}}. \quad (17)$$

for $1 \leq i \leq m$.

Proof. The " \Rightarrow " direction is easily deduced from Corollary 5.10 where each differential form corresponds to the differential conductor at a vertex $v \neq v_0$ of the associated tree. For instance, one can obtain from the tree in Example 4.5 a chain $\{12\} \prec \{5, 7\} \prec \{5, 4, 3\}$ and two exact differential forms $\frac{dx}{x^5(x-1)^7}, \frac{dx}{x^4(x-1)^3}$.

" \Leftarrow " We prove by induction on m . Suppose first that $m = 1$. Then, given an exact differential form of type (17), the tree in Figure 4 works. Suppose the statement is true for $m = n$. By induction, there exists a Hurwitz tree \mathcal{T}' with n non-root-vertices whose differential conductors are exact of type (17). Suppose that the difference between \underline{e}_n and \underline{e}_{n+1} is $\{l\} \prec \{l_1, \dots, l_r\}$. Suppose, moreover, that there exists an exact differential form $\omega = dx / (\prod_{i=1}^r (x - P_i)^{l_i})$, and the depth at the leaf corresponding to l is δ . Then one can "lengthen" \mathcal{T}' by "thickening" the leaf corresponding to l to an edge of thickness 1 and adding r leaves of conductors l_1, \dots, l_r to the end of this edge. Finally, we assign the exact differential form ω and the depth $\delta + l - 1$ to the end of the edge. One can easily check that the extended tree is Hurwitz. \square

Example 5.15. The tree below extends one from Figure 4,



where $l_2 = \sum_{j=1}^s l_{2,j}$, and the degeneration types at v_0, v_1 , and v_2 are $(0, \frac{1}{x^\epsilon})$, $(\epsilon(l-1), \frac{dx}{\prod_{i=1}^r (x-P_i)^{l_i}})$, and $(\epsilon(l-1) + l_2 - 1, \frac{dx}{\prod_{j=1}^s (x-P_{2,j})^{l_{2,j}}})$, respectively.

Remark 5.16. Proposition 5.14 implies that one can drop the "depth zero" condition in Corollary 5.10. Theorem 2.20 then easily follows.

We call an Artin-Schreier cover *equidistant* if the distance between any two branch points is the same. A deformation of a one point cover is equidistant if its generic fiber is equidistant, hence its Hurwitz tree has height one. Most of the known deformations have equidistant deformations as building blocks. The follows results then follows immediately from Proposition 5.14.

COROLLARY 5.17. *Suppose $\{h_1, h_2, \dots, h_r\}$ is a partition of $\{h\}$. Then there is a equidistant deformation over $k[[t]]$ of type $[h] \rightarrow [h_1, \dots, h_r]^T$ if and only if there exists an exact differential of*

the form

$$\frac{dx}{\prod_{i=1}^r (x - q_i)^{h_i}} \quad (18)$$

for some distinct elements q_i 's in k .

DEFINITION 5.18. We say the exact differential form in (18) has *type* $\{h_1, \dots, h_r\}$.

In the remainder of the paper, we will focus on differential forms of type (18).

Remark 5.19. Suppose we are given a rational function of the form $f(x) = g(x)^p h(x)$. Then $f'(x) = g(x)^p h'(x)$. Hence, it suffices to study differential forms of type (18), where $1 < h_i < p$.

Remark 5.20. In [BM00], Bertin and Mézard show that the infinitesimal deformation functor of a local Galois cover is represented by a *versal deformation ring*. Besides, they give some descriptions for the versal deformation rings of \mathbb{Z}/p -covers. Theorem 3.5 of [Hen00] and Proposition 5.14 imply that one can describe a deformation (over either a mixed or equal characteristic ring) of a \mathbb{Z}/p -cover by the associated Hurwitz tree, which, in turn, is determined by a sequence of exact differential forms in equal characteristic case, or exact and logarithmic differential forms in mixed characteristic case. We wonder if one can use a space of differential forms to give a full description of the versal deformation ring of an Artin-Schreier cover.

6. Applications

6.1 Some general exact differential form results

Recall from Proposition 5.14 that the deformation of an Artin-Schreier cover is determined by exact differential forms of type (17). Combining with Remark 5.19, we simplify Question 3.4 as follows.

QUESTION 6.1. Let k be an algebraically closed field of characteristic $p > 0$. Suppose $1 < h_i < p$ for $i = 1, 2, \dots, n$ are integers ($n \geq 2$). What are the conditions on the h_i 's so that the rational function

$$\frac{1}{\prod_{i=1}^n (x - P_i)^{h_i}} \quad (19)$$

is a derivative of some rational function in $k(x)$ for some P_i 's in k pairwise distinct?

When the multi-set $\{h_1, \dots, h_n\}$ is given, we can answer the question using Gröbner Basis techniques as follows. Equation (19) could be written as

$$\frac{\prod_{i=1}^n (x - P_i)^{p-h_i}}{\prod_{i=1}^n (x - P_i)^p} = \frac{\sum_j a_j x^j}{\prod_{i=1}^n (x - P_i)^p} \quad (20)$$

where a_j can be thought of as a polynomial in $k[P_1, \dots, P_n]$. As the denominator is a p -power, the fraction is exact if and only if the numerator $\sum_j a_j x^j$ is exact. That equates to there existing a choice of values for the P_i 's so that $a_j = 0$ for all $j \equiv -1 \pmod{p}$, or $1 \notin (a_j)_{j \equiv -1 \pmod{p}}$ by Hilbert's nullstellensatz. Moreover, all the P_i 's have to be distinct. That translates to $\prod_{j < k} (P_j - P_k)$ not lying in the radical of $(a_j)_{j \equiv -1 \pmod{p}}$, or the following ideal

$$\left(\{a_j\}_{j \equiv -1 \pmod{p}}, 1 - s \cdot \prod_{j < k} (P_j - P_k) \right)$$

is not the unit ideal of $k[P_1, \dots, P_n, s]$ by [DF04, § 15, Corollary 35]. We summarize by the below proposition.

PROPOSITION 6.2. *Suppose we are given a multi-set $\{h_1, \dots, h_n\} \in \Omega_h$. Then there exists a differential form*

$$\frac{dx}{\prod_{i=1}^n (x - P_i)^{h_i}}$$

if and only if $(\{\alpha_j\}_{j \equiv -1 \pmod{p}}, 1 - s \prod_{j < k} (P_j - P_k)) \in k[P_1, \dots, P_n, s]$ is not the unit ideal, where the α_j 's are defined in (20).

Remark 6.3. We can show whether the ideal in Proposition 6.2 is a unit one by checking whether 1 is its reduced Gröbner basis (with respect to any monomial ordering).

PROPOSITION 6.4. *Let k be an algebraically closed field of characteristic $p > 0$. Suppose $1 < h_i < p$ for $i = 1, 2, \dots, n$ are integers ($n \geq 2$). The rational differential form*

$$\omega = \frac{dx}{\prod_{i=1}^n (x - P_i)^{h_i}} \quad (21)$$

is exact for some distinct P_i 's in k only if $\sum_{i=1}^n h_i \geq p + n$. The converse is true when $n = 2$.

Proof. This argument is due to Fedor Petrov (see §1.1). Suppose that $\sum_{i=1}^n h_i < n + p$ and the differential form ω in (21) is exact. One may assume that the rational antiderivative of ω is of the form g/f , where $f = \prod_{i=1}^n (x - P_i)^{h_i - 1}$, and $\deg g < \deg f = \sum_{i=1}^n (h_i - 1) < p$. This may be seen from integrating the partial fraction decomposition of $\prod_{i=1}^n (x - P_i)^{-h_i}$ and convert it back to a single fraction. We have $(g/f)' = (g'f - f'g)/f^2$, and if $\deg f = a$, $\deg g = b$, the degree of the numerator equals $a + b - 1$, since the leading coefficient does not vanish (here we use that p can not divide $a - b$). Thus $1 = \prod_{i=1}^n (x - P_i)^{h_i} (g/f)'$ has degree $(n + a) + (a + b - 1) - 2a = n + b - 1 > 0$, a contradiction. Therefore, $\sum_{i=1}^n h_i$ has to be at least $n + p$ for (21) to be exact.

Suppose $n = 2$. One may assume that $P_1 = 0$. Consider the rational function

$$\omega = \frac{1}{x^{e_1} (x - Q)^{e_2}} = \frac{x^{p-e_1} (x - Q)^{p-e_2}}{x^p (x - Q)^p}.$$

Hence, ω is a derivative of some rational function if and only if $x^{p-e_1} (x - Q)^{p-e_2}$ is. The later is exact if and only if all the degree $kp - 1$ coefficients are equal to 0 as k varies. Suppose $e_1 + e_2 \geq p + 2$. Then the degree of the numerator is $2p - (e_1 + e_2) \leq p - 2$. Thus, it is a derivative. \square

Remark 6.5. The converse is not true in general for $n > 2$. See <https://mathoverflow.net/questions/310575> for some counterexamples to differential forms of type $\{2, 2, \dots, 2\}$ by Gjergji Zaimi. Using the Gröbner Basis technique (Proposition 6.2 and Remark 6.3), one can also show that there is no exact differential form of type $\{2, 2, 2, 6\}$ where $p = 7$, even though $2 + 2 + 2 + 6 > 7 + 4$.

6.1.1 *Oort-Sekiguchi-Suwa deformations* We first observe the following phenomenon.

PROPOSITION 6.6. *The differential form*

$$\frac{dx}{\prod_{i=1}^r (x - q_i)^{h_i}} \quad (22)$$

where $q_i \in k$'s are pairwise distinct, is exact if all but at most one of the h_i 's are divisible by p . In particular, there exists a flat deformation of type $[h] \rightarrow [h_1, \dots, h_r]^T$.

Proof. Suppose, without loss of generality, that h_1 is the only exponent that may not be divisible by p . Then we can easily see that differential form (22) is the derivative of the following rational

function

$$\frac{1}{(1 - h_1) \prod_{i=2}^r (x - q_i)^{h_i} (x - q_1)^{h_1 - 1}}.$$

The rest of the proposition follows from Proposition 5.14. \square

Remark 6.7. Suppose h_i 's are as in Proposition 6.6. Consider the \mathbb{Z}/p -cover of $\mathbb{P}_{k[[t_1, \dots, t_r]]}^1$ defined by

$$Y^p - Y = \frac{1}{(x - t_1)^{h_1 - 1} \prod_{i=2}^r (x - t_i)^{h_i}} \quad (23)$$

Replacing the t_i 's by any distinct elements P_i 's of $\text{mk}[[t]]$ (i.e. $v_i(P_i) > 0$), one can show that (23) defines a flat deformation (over $k[[t]]$) of type $[h] \rightarrow [h_1, h_2, \dots, h_r]^\top$.

Remark 6.8. Suppose Ψ is a \mathbb{Z}/p -cover over $k[[t]]$. In [BM00], Bertin and Mézard study the *versal deformation ring* R_Ψ whose spectrum is the formal deformation space of Ψ . They construct explicitly a family of deformations of Ψ over a polynomial ring over $R = \mathcal{W}(k)[\zeta_p]$. This family is parameterized by an irreducible component of the formal deformation space, called *the Oort-Sekiguchi-Suwa component*. Theorem 5.3.3 of the same paper shows that the dimension of this component is equal to the dimension of the versal deformation ring. Following [Dan20, §3.1.1], one can realize the characteristic p fibers of these deformations as special cases of one described in Proposition 6.6.

6.1.2 Non Oort-Sekiguchi-Suwa deformations In [Dan20], to prove that \mathcal{AS}_g is connected when g is large, we construct some equidistant deformations that do *not* lie in p -fibers of the Oort-Sekiguchi-Suwa component. In this section, we realize these deformations in term of exact differential forms.

PROPOSITION 6.9. *There exist exact differential forms of the following types:*

- (i) $\{h_1, h_2\}$ where $\bar{h}_1 + \bar{h}_2 \geq p + 2$,
- (ii) $\{h_1, h_2, h_3\}$ where $h_i \equiv (p + 1)/2 \pmod{p}$ and $p \geq 3$,
- (iii) $\{p - 1, h_1, h_2\}$ where $h_1, h_2 \not\equiv 1 \pmod{p}$,
- (iv) $\underbrace{\{n + 1, n + 1, \dots, n + 1\}}_{p-n+1}$ where $1 \leq n \leq p - 1$
- (v) $\{3, 2, 2, 2\}$ and $\{3, 3, 2, 2\}$ where $p = 5$

Proof. Item (i) follows immediately from 6.4.

Consider item (ii). Suppose the desired differential is of the form

$$\frac{dx}{(x - a_1)^{h_1} (x - a_2)^{h_2} (x - a_3)^{h_3}}. \quad (24)$$

As discussed in Remark 5.19, one may assume, without loss of generality, that $h_i = (p + 1)/2$. Set $a_1 = 0$, $a_2 = 1$, and rewrite the differential form (24) as

$$\frac{x^{(p-1)/2} (x - 1)^{(p-1)/2} (x - a_3)^{(p-1)/2} dx}{x^p (x - 1)^p (x - a_3)^p}.$$

As the denominator is a p -power, the differential form is exact if and only if all the terms of degree congruent to -1 modulo p in the numerator are zero. One can easily see the leading term has degree $3(p - 1)/2 < 2p - 1$. Hence, it suffices to make the term of degree $p - 1$, which is

$$\sum_{i+j=\frac{p-1}{2}} \binom{\frac{p-1}{2}}{i} \binom{\frac{p-1}{2}}{j} a_3^{(p-1)/2-j} = \sum_{i=0}^{(p-1)/2} \binom{\frac{p-1}{2}}{i}^2 a_3^{(p-1)/2-i}, \quad (25)$$

to be equal to zero. We thus want a_3 to be a root of (25) (which can be thought of as a polynomial in $k[a_3]$ that is different from 0 and 1. As the constant coefficient of the polynomial is nonzero, 0 cannot be a root. Moreover, since the polynomial $\sum_{i=0}^{(p-1)/2} \binom{\frac{p-1}{2}}{i}^2 x^{(p-1)/2-i}$ is separable by [Sil86, Theorem 4.1] and $(p-1)/2$ is at least 2 for $p \geq 5$, there are roots (in k) of (25) that are different from 1. Hence, if we pick a_3 to be one of these roots, the differential form is exact as desired.

By direct computation, one can show that the differential forms $\frac{dx}{x^{p-1}(x-1)^{h_1}(x-(1-h_2)/(h_1-1))^{h_2}}$, $\frac{dx}{x^{n+1}(x^{p-n}-1)^{n+1}}$, $\frac{dx}{x^3(x-1)^2(x^2+x+1)^2}$, and $\frac{dx}{x^3(x-1)^3(x^2+4x+2)^2}$ are solutions to (iii), (iv), and (v), respectively. \square

This gives an alternative proof for [Dan20, Theorem 3.7]. We paraphrase the statement of that theorem using the language in this paper.

THEOREM 6.10 c.f. [Dan20, Theorem 3.7]. *There exist equidistant deformations over $k[[t]]$ of the following type.*

- (i) $[h] \rightarrow [h_1, h_2]^\top$ where \bar{h}_1 and \bar{h}_2 are non-zero and $\bar{h}_1 + \bar{h}_2 \geq p + 2$.
- (ii) $[h] \rightarrow [h_1, h_2, h_3]^\top$, where $h_i \equiv (p+1)/2 \pmod{p}$ and $p \geq 3$.
- (iii) $[h] \rightarrow [p-1, h_2, h_3]^\top$ where \bar{h}_1 and \bar{h}_2 are non-zero.
- (iv) $[(n+1)(p-n+1)] \rightarrow \underbrace{[n+1, \dots, n+1]^\top}_{p-n+1}$, where $1 \leq n \leq p-1$.
- (v) $[9] \rightarrow [3, 2, 2, 2]^\top$ or $[10] \rightarrow [3, 3, 2, 2]^\top$ where $p = 5$.

Proof. The theorem follows immediately from Proposition 5.14 and Proposition 6.9. \square

6.2 Disconnectedness of \mathcal{AS}_g

Recall that the moduli space \mathcal{AS}_g can be partitioned by strata that are associated to partitions of $2g/(p-1) + 2$. Moreover, Pries and Zhu show that the dimension of the stratum indexed by $\vec{E} = \{h_1, \dots, h_r\}$ is given by

$$d - 1 - \sum_{j=1}^r \left(\left\lfloor \frac{h_j - 1}{p} \right\rfloor \right)$$

[PZ12, Corollary 3.11], recall that $d = \frac{2g}{p-1} + 2$. Therefore, the irreducible components of \mathcal{AS}_g are the closure of the strata indexed by partitions of the form $\{h_1, \dots, h_r\}$, where $h_i \leq p$ for all i . One key ingredient of our first connectedness result is that we prove, where the sum of conductors $d + 2$ is not congruent 1 modulo p , that all the strata of non-zero-codimension lie in the same connected component [Dan20, Corollary 4.3]. In general, \mathcal{AS}_g is connected only if none of the strata of codimension-zero is closed (unless there is only one stratum in \mathcal{AS}_g). Furthermore, the closedness of a stratum can be realized by the following result.

PROPOSITION 6.11. *A stratum of \mathcal{AS}_g indexed by $\vec{E} = \{h_1, \dots, h_r\}$ is not closed if and only if there exists a partition $\{g_1, \dots, g_s\} \subseteq \{h_1, \dots, h_r\}$, and an exact differential form of type $\{g_1, \dots, g_s\}$.*

Proof. " \Leftarrow ": Suppose, without loss of generality, that $\{g_1, \dots, g_s\} = \{h_1, \dots, h_s\}$. The " \Leftarrow " direction then follows immediately from Proposition 5.14 as the closure of $\Gamma_{\vec{E}}$ contains the stratum indexed by $\{\sum_{i=1}^s h_i, h_{s+1}, \dots, h_r\}$.

" \Rightarrow ": Suppose \vec{E} is not closed. Then, by Proposition 2.10, there exists a partition $\vec{E}' = \{l_1, \dots, l_m\} \prec \vec{E}$ such that $\Gamma_{\vec{E}'} \subsetneq \Gamma_{\vec{E}}$. It then follows from Proposition 2.16 that there exist $1 \leq i \leq m$ so that $\{l_i\} \prec \{h_{i,1}, \dots, h_{i,m}\} \subseteq \vec{E}$, and a deformation of type $[l_i] \rightarrow [h_{i,1}, \dots, h_{i,m}]^\top$. Finally, Proposition 5.14 shows that there exists an exact differential form of type $\{g_1, \dots, g_s\} \subseteq \{h_{i,1}, \dots, h_{i,m}\}$, where $\{\sum_{i=1}^s g_i\} \prec \{g_1, \dots, g_s\}$ is the difference between \vec{E} and the immediate one below in the chain of partitions from that proposition. That completes the proof. \square

We can finally give the proof of Theorem 1.3.

6.2.1 *Proof of Theorem 1.3* It is already known from Theorem 1.1 that, when $p = 5$, \mathcal{AS}_g is connected when $g > (p-1)(p-2) = 12$. The same result shows that, when $p \geq 5$, \mathcal{AS}_g is disconnected when $(p-1)/2 < g \leq (p-1)^2/2$. Hence, it suffices to show that \mathcal{AS}_g is disconnected if $(p-1)^2/2 < g \leq (p-1)(p-2)$, or the sum of conductors $d+2$ is between (and including) $p+3$ and $2p-2$.

Recall that the irreducible components of \mathcal{AS}_g are indexed by the closure of the strata corresponding to $\{h_1, \dots, h_r\}$ where $h_i \leq p$ for all i . Suppose $d+2$ is odd (resp. even). Then there exists a stratum indexed by $\vec{E}_1 := \{3, 2, \dots, 2\}$ (resp. $\vec{E}_2 := \{2, 2, \dots, 2\}$). It is straightforward to check that, for any $\{k_1, \dots, k_s\} \subseteq \vec{E}_1$ (resp. \vec{E}_2), $\sum_{j=1}^s k_j < p+s$. Here we use that fact that the sum of the entries of \vec{E}_1 (resp. \vec{E}_2), i.e., the number $d+2$, is smaller than $2p-2$. Therefore, it follows from Proposition 6.11 that the closure of the stratum corresponding to $\{3, 2, \dots, 2\}$ (resp. $\{2, 2, \dots, 2\}$) only contains itself. Thus, the moduli space \mathcal{AS}_g is disconnected. \square

REFERENCES

- BL93 Siegfried Bosch and Werner Lütkebohmert. Formal and rigid geometry. I. Rigid spaces. *Math. Ann.*, 295(2):291–317, 1993.
- BM00 José Bertin and Ariane Mézard. Déformations formelles des revêtements sauvagement ramifiés de courbes algébriques. *Invent. Math.*, 141(1):195–238, 2000.
- BW06 Irene I. Bouw and Stefan Wewers. The local lifting problem for dihedral groups. *Duke Math. J.*, 134(3):421–452, 2006.
- BW09 Louis Hugo Brewis and Stefan Wewers. Artin characters, Hurwitz trees and the lifting problem. *Math. Ann.*, 345(3):711–730, 2009.
- CTT16 Adina Cohen, Michael Temkin, and Dmitri Trushin. Morphisms of Berkovich curves and the different function. *Adv. Math.*, 303:800–858, 2016.
- Dan20 Huy Dang. Connectedness of the moduli space of Artin-Schreier curves of fixed genus. *Journal of Algebra*, 547:398 – 429, 2020.
- DF04 David S. Dummit and Richard M. Foote. *Abstract algebra*. John Wiley & Sons, Inc., Hoboken, NJ, third edition, 2004.
- Epp73 Helmut P. Epp. Eliminating wild ramification. *Invent. Math.*, 19:235–249, 1973.
- GM98 Barry Green and Michel Matignon. Liftings of Galois covers of smooth curves. *Compositio Math.*, 113(3):237–272, 1998.
- Har77 Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.

- HB09 Louis Hugo Brewis. *Ramification theory of the p -adic open disc and the lifting problem*. PhD thesis, Mathematik und Wirtschaftswissenschaften der Universität Ulm, 10 2009.
- Hen00 Y. Henrio. Arbres de Hurwitz et automorphismes d'ordre p des disques et des couronnes p -adiques formels. *ArXiv Mathematics e-prints*, November 2000.
- Kat80 Kazuya Kato. A generalization of local class field theory by using K -groups. II. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 27(3):603–683, 1980.
- Kat87 Kazuya Kato. Vanishing cycles, ramification of valuations, and class field theory. *Duke Math. J.*, 55(3):629–659, 1987.
- Lan02 Serge Lang. *Algebra*, volume 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2002.
- Lor08 Falko Lorenz. *Algebra. Vol. II*. Universitext. Springer, New York, 2008. Fields with structure, algebras and advanced topics, Translated from the German by Silvio Levy, With the collaboration of Levy.
- Mau03 Sylvain Maugeais. Relèvement des revêtements p -cycliques des courbes rationnelles semi-stables. *Math. Ann.*, 327(2):365–393, 2003.
- Obu12 Andrew Obus. The (local) lifting problem for curves. In *Galois-Teichmüller theory and arithmetic geometry*, volume 63 of *Adv. Stud. Pure Math.*, pages 359–412. Math. Soc. Japan, Tokyo, 2012.
- OW14 Andrew Obus and Stefan Wewers. Cyclic extensions and the local lifting problem. *Ann. of Math. (2)*, 180(1):233–284, 2014.
- OW16 Andrew Obus and Stefan Wewers. Wild ramification kinks. *Res. Math. Sci.*, 3:Paper No. 21, 27, 2016.
- Pop14 Florian Pop. The Oort conjecture on lifting covers of curves. *Ann. of Math. (2)*, 180(1):285–322, 2014.
- Pri03 Rachel J. Pries. Conductors of wildly ramified covers. III. *Pacific J. Math.*, 211(1):163–182, 2003.
- PZ12 Rachel Pries and Hui June Zhu. The p -rank stratification of Artin-Schreier curves. *Ann. Inst. Fourier (Grenoble)*, 62(2):707–726, 2012.
- Sai04 Mohamed Saïdi. Wild ramification and a vanishing cycles formula. *J. Algebra*, 273(1):108–128, 2004.
- Sai07 Mohamed Saïdi. Galois covers of degree p and semi-stable reduction of curves in equal characteristic $p > 0$. *Math. J. Okayama Univ.*, 49:113–138, 2007.
- Ser79 Jean-Pierre Serre. *Local fields*, volume 67 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1979. Translated from the French by Marvin Jay Greenberg.
- Sil86 Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1986.
- Wea18 Bradley Weaver. The local lifting problem for D_4 . *Israel J. Math.*, 228(2):587–626, 2018.
- Wew05 Stefan Wewers. Swan conductors on the boundary of Lubin-Tate spaces. *arXiv Mathematics e-prints*, page math/0511434, November 2005.
- Wew14 Stefan Wewers. Fiercely ramified cyclic extensions of p -adic fields with imperfect residue field. *Manuscripta Math.*, 143(3-4):445–472, 2014.

Huy Dang hqd4bz@virginia.edu

Institute of Mathematics, Vietnam Academy of Science and Technology, Hanoi, Vietnam