

# A remarkable congruence involving Appell polynomials

Abdelkader Benyattou

UNIV-DJELFA, Dep. of Mathematics and Informatics, RECITS Laboratory, Djelfa, Algeria.

a.benyattou@univ-djelfa.dz, abdelkaderbenyattou@gmail.com

## Abstract

In this paper, by the umbral calculus method, we give a remarkable congruence involving Appell polynomials. Some applications on derangement polynomials are also presented.

**Keywords.** Congruence, Appell polynomials, classical umbral calculus.

**2010 MSC.** 11B73, 11A07, 05A40.

## 1 Introduction

Let  $(\mathcal{A}_n(x))$  be a sequence of Appell polynomials with integer coefficients defined by [1]:

$$\sum_{n \geq 0} \mathcal{A}_n(x) \frac{t^n}{n!} = F(t) \exp(xt), \quad \mathcal{A}_n(0) = \mathcal{A}_n \in \mathbb{N},$$

where

$$F(t) = 1 + \sum_{n \geq 1} \mathcal{A}_n \frac{t^n}{n!},$$

Let  $\mathbf{A}$  be the Appell umbra defined by  $\mathbf{A}^n = \mathcal{A}_n$ , then we can define the generalized Appell umbra  $\mathbf{A}_x$  as follows

$$\mathbf{A}_x^n = \mathcal{A}_n(x) = (\mathbf{A} + x)^n$$

see [4]. For more information on the umbral calculus see [2, 3, 5, 6]. In the remainder of this paper, for any polynomials  $f$  and  $g$ , we denote by  $f(x) \equiv g(x)$  to mean  $f(x) \equiv g(x) \pmod{\mathbb{Z}[x]}$  and for any numbers  $a$  and  $b$  by  $a \equiv b$  we mean  $a \equiv b \pmod{p}$ .

## 2 Congruences involving Appell polynomials

Main result is given by the following theorem.

**Theorem 1** *Let  $f$  be a polynomial in  $\mathbb{Z}[x]$ ,  $m \geq 0, s \geq 1$  be integers and  $p$  be an odd prime number. Then if for any integer  $n \geq 0$ , there exists an integer  $t$  such that  $\mathcal{A}_{n+mp} \equiv t\mathcal{A}_n$ , we have the following congruence*

$$(\mathbf{A} + x)^{mp^s} f(\mathbf{A} + x) \equiv (x^{p^s} + t)^m f(\mathbf{A} + x).$$

*This congruence is equivalent when  $f(x) = x^n$  to*

$$\mathcal{A}_{n+mp^s}(x) \equiv (x^{p^s} + t)^m \mathcal{A}_n(x).$$

*In particular for  $n = 0$ , we have*

$$\mathcal{A}_{mp^s}(x) \equiv (x^{p^s} + t)^m.$$

**Proof.** It suffices to take  $f(x) = x^n$ . For  $m = 1$  we proceed by induction on  $s$ . Indeed, for  $s = 1$  we have

$$\begin{aligned}
(\mathbf{A}_x^p - t) \mathbf{A}_x^n &= \mathbf{A}_x^{n+p} - t \mathbf{A}_x^n \\
&= (\mathbf{A} + x)^p (\mathbf{A} + x)^n - t \mathbf{A}_x^n \\
&\equiv (\mathbf{A}^p + x^p) (\mathbf{A} + x)^n - t \mathbf{A}_x^n \\
&= x^p \mathbf{A}_x^n + \sum_{k=0}^n \binom{n}{k} \mathbf{A}^{n+p-k} x^k - t \mathbf{A}_x^n \\
&\equiv x^p \mathbf{A}_x^n + t \sum_{k=0}^n \binom{n}{k} \mathbf{A}^{n-k} x^k - t \mathbf{A}_x^n \\
&= x^p \mathbf{A}_x^n.
\end{aligned}$$

Assume it is true for  $s \geq 1$ . Then we have

$$\begin{aligned}
\mathbf{A}_x^n (\mathbf{A}_x^{p^{s+1}} - t) &= \mathbf{A}_x^n \left( (\mathbf{A}_x^{p^s} - t + t)^p - t \right) \\
&= \mathbf{A}_x^n \left( (\mathbf{A}_x^{p^s} - t)^p + t - t \right) \\
&\equiv \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^p \\
&= \left[ \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t) \right] (\mathbf{A}_x^{p^s} - t)^{p-1} \\
&\equiv x^{p^s} \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^{p-1} \\
&= x^{p^s} \left[ \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t) \right] (\mathbf{A}_x^{p^s} - t)^{p-2} \\
&\equiv x^{2p^s} \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^{p-2} \\
&\vdots \\
&\equiv (x^{p^s})^p \mathbf{A}_x^n \\
&= x^{p^{s+1}} \mathbf{A}_x^n.
\end{aligned}$$

So, we proved that  $(\mathbf{A} + x)^{p^s} f(\mathbf{A} + x) \equiv (x^{p^s} + t) f(\mathbf{A} + x)$ . For  $m \geq 1$  we use this last congruence to obtain

$$\begin{aligned}
(\mathbf{A} + x)^{m p^s} f(\mathbf{A} + x) &= (\mathbf{A} + x)^{p^s} (\mathbf{A} + x)^{(m-1)p^s} f(\mathbf{A} + x) \\
&\equiv (x^{p^s} + t) (\mathbf{A} + x)^{(m-1)p^s} f(\mathbf{A} + x) \\
&\vdots \\
&\equiv (x^{p^s} + t)^m f(\mathbf{A} + x).
\end{aligned}$$

■

**Corollary 2** Let  $f$  be a polynomial in  $\mathbb{Z}[x]$ ,  $m \geq 0$ ,  $s \geq 1$  be integers and  $p$  be an odd prime number. Then if for any integer  $n \geq 0$ , there exists an integer  $t$  such that  $\mathcal{A}_{n+p} \equiv t \mathcal{A}_n$ , there holds

$$(\mathbf{A} + x)^{m_1 p + \dots + m_s p^s} f(\mathbf{A} + x) \equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} f(\mathbf{A} + x).$$

This congruence is equivalent when  $f(x) = x^{m_0}$  to

$$\mathcal{A}_{m_0 + m_1 p + \dots + m_s p^s}(x) \equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} \mathcal{A}_{m_0}(x).$$

In particular, for  $m_0 = 0$ , we have

**Proof.** By the Theorem 1 we can write

$$\begin{aligned}
(\mathbf{A} + x)^{m_1 p + \dots + m_s p^s} f(\mathbf{A} + x) &= (\mathbf{A} + x)^{m_1 p} (\mathbf{A} + x)^{m_2 p^2 + \dots + m_s p^s} f(\mathbf{A} + x) \\
&\equiv (x^p + t)^{m_1} (\mathbf{A} + x)^{m_2 p^2 + \dots + m_s p^s} f(\mathbf{A} + x) \\
&\vdots \\
&\equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} f(\mathbf{A} + x).
\end{aligned}$$

■

### 3 Applications

In this section we give some applications of Theorem 1.

The derangement polynomials are defined by

$$\mathcal{D}_n(x) = \sum_{k=0}^n \binom{n}{k} \mathcal{D}_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} k! (x-1)^{n-k},$$

where  $\mathcal{D}_n = \mathcal{D}_n(0)$  is the  $n$ -th derangement number, denoted by  $\mathcal{D}_n$  counting the number of permutation of the set  $[n]$  without a fixed point. With exponential generating series

$$\sum_{i=0}^{\infty} \mathcal{D}_i(x) \frac{t^i}{i!} = \frac{e^{-t}}{1-t} e^{xt}.$$

Recall that for any integer  $n \geq 2$ , we have

$$\mathcal{D}_n(x) = n\mathcal{D}_{n-1}(x) + (x-1)^n \tag{1}$$

The derangement numbers satisfy the following congruence [7]. For any non-negative integer  $n$  and any prime number  $p$ ,

$$\mathcal{D}_{n+p} \equiv -\mathcal{D}_n.$$

Let  $D$  be the derangement umbra defined by  $D^n = \mathcal{D}_n$  and  $\mathbf{D}_x$  be the generalized derangement umbra defined by  $\mathbf{D}_x^n = (D+x)^n = \mathcal{D}_n(x)$ .

**Lemma 3** Let  $g(x) = \sum_{k=2}^n a_k x^k$ , be a polynomial with real coefficients. Then

$$g(\mathbf{D}_x) = g'(\mathbf{D}_x) + \sum_{k=2}^n a_k (x-1)^k.$$

where  $g'$  is the derivative polynomial.

**Proof.** We have

$$\begin{aligned}
g(\mathbf{D}_x) &= \sum_{k=2}^n a_k (\mathbf{D}_x)^k \\
&= \sum_{k=2}^n a_k \left( k\mathbf{D}_x^{k-1} + (x-1)^k \right) \\
&= \sum_{k=2}^n a_k k \mathbf{D}_x^{k-1} + \sum_{k=2}^n a_k (x-1)^k.
\end{aligned}$$

■ When we replace  $g(x)$  in lemma 3, by  $x^n$  we obtain the identity (1)

**Corollary 4** For any integers  $n \geq 1$ ,  $s \geq 1$ ,  $m \geq 0$  and for any prime number  $p \geq 3$ , there holds

$$\mathcal{D}_{n+mp^s}(x) \equiv (x^{p^s} - 1)^m \mathcal{D}_n(x). \tag{2}$$

For  $x = 0$ , we obtain

**Proof.** The congruence (2) follows by setting  $f(x) = x^n$  and  $t = -1$  in Theorem 1 ■

**Corollary 5** For any prime number  $p \geq 3$  and any integers  $s \geq 1, m_0, \dots, m_s \in \{0, \dots, p-1\}$ , there holds

$$\mathcal{D}_{m_0+m_1p+\dots+m_sp^s}(x) \equiv (x^p - 1)^{m_1} (x^{p^2} - 1)^{m_2} \dots (x^{p^s} - 1)^{m_s} \mathcal{D}_{m_0}(x).$$

In particular, we have

$$\begin{aligned} \mathcal{D}_{m_1p+\dots+m_sp^s}(x) &\equiv (x^p - 1)^{m_1} (x^{p^2} - 1)^{m_2} \dots (x^{p^s} - 1)^{m_s}, \\ \mathcal{D}_{m_1p+\dots+m_sp^s}(k) &\equiv (k - 1)^{m_1+m_2+\dots+m_s}. \end{aligned}$$

## References

- [1] P. Appell. Sur une classe de polynomes. *Ann Sci Ecole Norm Sup.* **9** (1880), 119–144.
- [2] A. Benyattou, M. Mihoubi, Curious congruences related to the Bell polynomials, *Quaest. Math.*, 41 (3), (2018), pp-437–448.
- [3] I. M. Gessel, Applications of the classical umbral calculus, *Algebra Universalis* **49** (2003), 397–434.
- [4] M. Mihoubi and S. Taharbouchet, Some identities involving Appell polynomials, *Quaest. Math.*, DOI: 10.2989/16073606.2018.1544944
- [5] G. C. Rota and B. D. Taylor, The classical umbral calculus, *SIAM J. Math. Anal.* **25** (1994), 694–711.
- [6] S. Roman, *The Umbral Calculus*, Academic Press, Orlando, FL, 1984.
- [7] Z.-W. Sun, D. Zagier, On a curious property of Bell numbers, *Bull. Aust. Math. Soc.* **84** (2011), 153–158.