

A remarkable congruence involving Appell polynomials

Abdelkader Benyattou

UNIV-DJELFA, Dep. of Mathematics and Informatics, RECITS Laboratory, Djelfa, Algeria.

a.benyattou@univ-djelfa.dz, abdelkaderbenyattou@gmail.com

Abstract

In this paper, by the umbral calculus method, we give a remarkable congruence involving Appell polynomials. Some applications on derangement polynomials are also presented.

Keywords. Congruence, Appell polynomials, classical umbral calculus.

2010 MSC. 11B73, 11A07, 05A40.

1 Introduction

Let $(\mathcal{A}_n(x))$ be a sequence of Appell polynomials with integer coefficients defined by [1]:

$$\sum_{n \geq 0} \mathcal{A}_n(x) \frac{t^n}{n!} = F(t) \exp(xt), \quad \mathcal{A}_n(0) = \mathcal{A}_n \in \mathbb{N},$$

where

$$F(t) = 1 + \sum_{n \geq 1} \mathcal{A}_n \frac{t^n}{n!},$$

Let \mathbf{A} be the Appell umbra defined by $\mathbf{A}^n = \mathcal{A}_n$, then we can define the generalized Appell umbra \mathbf{A}_x as follows

$$\mathbf{A}_x^n = \mathcal{A}_n(x) = (\mathbf{A} + x)^n$$

see [4]. For more information on the umbral calculus see [2, 3, 5, 6]. In the remainder of this paper, for any polynomials f and g , we denote by $f(x) \equiv g(x)$ to mean $f(x) \equiv g(x) \pmod{\mathbb{Z}[x]}$ and for any numbers a and b by $a \equiv b$ we mean $a \equiv b \pmod{p}$.

2 Congruences involving Appell polynomials

Main result is given by the following theorem.

Theorem 1 *Let f be a polynomial in $\mathbb{Z}[x]$, $m \geq 0, s \geq 1$ be integers and p be an odd prime number. Then if for any integer $n \geq 0$, there exists an integer t such that $\mathcal{A}_{n+mp} \equiv t\mathcal{A}_n$, we have the following congruence*

$$(\mathbf{A} + x)^{mp^s} f(\mathbf{A} + x) \equiv (x^{p^s} + t)^m f(\mathbf{A} + x).$$

This congruence is equivalent when $f(x) = x^n$ to

$$\mathcal{A}_{n+mp^s}(x) \equiv (x^{p^s} + t)^m \mathcal{A}_n(x).$$

In particular for $n = 0$, we have

$$\mathcal{A}_{mp^s}(x) \equiv (x^{p^s} + t)^m.$$

Proof. It suffices to take $f(x) = x^n$. For $m = 1$ we proceed by induction on s . Indeed, for $s = 1$ we have

$$\begin{aligned}
(\mathbf{A}_x^p - t) \mathbf{A}_x^n &= \mathbf{A}_x^{n+p} - t \mathbf{A}_x^n \\
&= (\mathbf{A} + x)^p (\mathbf{A} + x)^n - t \mathbf{A}_x^n \\
&\equiv (\mathbf{A}^p + x^p) (\mathbf{A} + x)^n - t \mathbf{A}_x^n \\
&= x^p \mathbf{A}_x^n + \sum_{k=0}^n \binom{n}{k} \mathbf{A}^{n+p-k} x^k - t \mathbf{A}_x^n \\
&\equiv x^p \mathbf{A}_x^n + t \sum_{k=0}^n \binom{n}{k} \mathbf{A}^{n-k} x^k - t \mathbf{A}_x^n \\
&= x^p \mathbf{A}_x^n.
\end{aligned}$$

Assume it is true for $s \geq 1$. Then we have

$$\begin{aligned}
\mathbf{A}_x^n (\mathbf{A}_x^{p^{s+1}} - t) &= \mathbf{A}_x^n \left((\mathbf{A}_x^{p^s} - t + t)^p - t \right) \\
&= \mathbf{A}_x^n \left((\mathbf{A}_x^{p^s} - t)^p + t - t \right) \\
&\equiv \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^p \\
&= \left[\mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t) \right] (\mathbf{A}_x^{p^s} - t)^{p-1} \\
&\equiv x^{p^s} \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^{p-1} \\
&= x^{p^s} \left[\mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t) \right] (\mathbf{A}_x^{p^s} - t)^{p-2} \\
&\equiv x^{2p^s} \mathbf{A}_x^n (\mathbf{A}_x^{p^s} - t)^{p-2} \\
&\vdots \\
&\equiv (x^{p^s})^p \mathbf{A}_x^n \\
&= x^{p^{s+1}} \mathbf{A}_x^n.
\end{aligned}$$

So, we proved that $(\mathbf{A} + x)^{p^s} f(\mathbf{A} + x) \equiv (x^{p^s} + t) f(\mathbf{A} + x)$. For $m \geq 1$ we use this last congruence to obtain

$$\begin{aligned}
(\mathbf{A} + x)^{m p^s} f(\mathbf{A} + x) &= (\mathbf{A} + x)^{p^s} (\mathbf{A} + x)^{(m-1)p^s} f(\mathbf{A} + x) \\
&\equiv (x^{p^s} + t) (\mathbf{A} + x)^{(m-1)p^s} f(\mathbf{A} + x) \\
&\vdots \\
&\equiv (x^{p^s} + t)^m f(\mathbf{A} + x).
\end{aligned}$$

■

Corollary 2 Let f be a polynomial in $\mathbb{Z}[x]$, $m \geq 0$, $s \geq 1$ be integers and p be an odd prime number. Then if for any integer $n \geq 0$, there exists an integer t such that $\mathcal{A}_{n+p} \equiv t \mathcal{A}_n$, there holds

$$(\mathbf{A} + x)^{m_1 p + \dots + m_s p^s} f(\mathbf{A} + x) \equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} f(\mathbf{A} + x).$$

This congruence is equivalent when $f(x) = x^{m_0}$ to

$$\mathcal{A}_{m_0 + m_1 p + \dots + m_s p^s}(x) \equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} \mathcal{A}_{m_0}(x).$$

In particular, for $m_0 = 0$, we have

Proof. By the Theorem 1 we can write

$$\begin{aligned}
(\mathbf{A} + x)^{m_1 p + \dots + m_s p^s} f(\mathbf{A} + x) &= (\mathbf{A} + x)^{m_1 p} (\mathbf{A} + x)^{m_2 p^2 + \dots + m_s p^s} f(\mathbf{A} + x) \\
&\equiv (x^p + t)^{m_1} (\mathbf{A} + x)^{m_2 p^2 + \dots + m_s p^s} f(\mathbf{A} + x) \\
&\vdots \\
&\equiv (x^p + t)^{m_1} \dots (x^{p^s} + t)^{m_s} f(\mathbf{A} + x).
\end{aligned}$$

■

3 Applications

In this section we give some applications of Theorem 1.

The derangement polynomials are defined by

$$\mathcal{D}_n(x) = \sum_{k=0}^n \binom{n}{k} \mathcal{D}_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} k! (x-1)^{n-k},$$

where $\mathcal{D}_n = \mathcal{D}_n(0)$ is the n -th derangement number, denoted by \mathcal{D}_n counting the number of permutation of the set $[n]$ without a fixed point. With exponential generating series

$$\sum_{i=0}^{\infty} \mathcal{D}_i(x) \frac{t^i}{i!} = \frac{e^{-t}}{1-t} e^{xt}.$$

Recall that for any integer $n \geq 2$, we have

$$\mathcal{D}_n(x) = n\mathcal{D}_{n-1}(x) + (x-1)^n \tag{1}$$

The derangement numbers satisfy the following congruence [7]. For any non-negative integer n and any prime number p ,

$$\mathcal{D}_{n+p} \equiv -\mathcal{D}_n.$$

Let D be the derangement umbra defined by $D^n = \mathcal{D}_n$ and \mathbf{D}_x be the generalized derangement umbra defined by $\mathbf{D}_x^n = (D+x)^n = \mathcal{D}_n(x)$.

Lemma 3 Let $g(x) = \sum_{k=2}^n a_k x^k$, be a polynomial with real coefficients. Then

$$g(\mathbf{D}_x) = g'(\mathbf{D}_x) + \sum_{k=2}^n a_k (x-1)^k.$$

where g' is the derivative polynomial.

Proof. We have

$$\begin{aligned}
g(\mathbf{D}_x) &= \sum_{k=2}^n a_k (\mathbf{D}_x)^k \\
&= \sum_{k=2}^n a_k \left(k\mathbf{D}_x^{k-1} + (x-1)^k \right) \\
&= \sum_{k=2}^n a_k k\mathbf{D}_x^{k-1} + \sum_{k=2}^n a_k (x-1)^k.
\end{aligned}$$

■ When we replace $g(x)$ in lemma 3, by x^n we obtain the identity (1)

Corollary 4 For any integers $n \geq 1$, $s \geq 1$, $m \geq 0$ and for any prime number $p \geq 3$, there holds

$$\mathcal{D}_{n+mp^s}(x) \equiv (x^{p^s} - 1)^m \mathcal{D}_n(x). \tag{2}$$

For $x = 0$, we obtain

Proof. The congruence (2) follows by setting $f(x) = x^n$ and $t = -1$ in Theorem 1 ■

Corollary 5 For any prime number $p \geq 3$ and any integers $s \geq 1, m_0, \dots, m_s \in \{0, \dots, p-1\}$, there holds

$$\mathcal{D}_{m_0+m_1p+\dots+m_sp^s}(x) \equiv (x^p - 1)^{m_1} (x^{p^2} - 1)^{m_2} \dots (x^{p^s} - 1)^{m_s} \mathcal{D}_{m_0}(x).$$

In particular, we have

$$\begin{aligned} \mathcal{D}_{m_1p+\dots+m_sp^s}(x) &\equiv (x^p - 1)^{m_1} (x^{p^2} - 1)^{m_2} \dots (x^{p^s} - 1)^{m_s}, \\ \mathcal{D}_{m_1p+\dots+m_sp^s}(k) &\equiv (k - 1)^{m_1+m_2+\dots+m_s}. \end{aligned}$$

References

- [1] P. Appell. Sur une classe de polynomes. *Ann Sci Ecole Norm Sup.* **9** (1880), 119–144.
- [2] A. Benyattou, M. Mihoubi, Curious congruences related to the Bell polynomials, *Quaest. Math.*, 41 (3), (2018), pp-437—448.
- [3] I. M. Gessel, Applications of the classical umbral calculus, *Algebra Universalis* **49** (2003), 397–434.
- [4] M. Mihoubi and S. Taharbouchet, Some identities involving Appell polynomials, *Quaest. Math.*, DOI: 10.2989/16073606.2018.1544944
- [5] G. C. Rota and B. D. Taylor, The classical umbral calculus, *SIAM J. Math. Anal.* **25** (1994), 694–711.
- [6] S. Roman, *The Umbral Calculus*, Academic Press, Orlando, FL, 1984.
- [7] Z.-W. Sun, D. Zagier, On a curious property of Bell numbers, *Bull. Aust. Math. Soc.* **84** (2011), 153–158.