

ON THE AREA BOUNDED BY THE CURVE

$$\prod_{k=1}^n \left| x \sin\left(\frac{k\pi}{n}\right) - y \cos\left(\frac{k\pi}{n}\right) \right| = 1$$

ANTON MOSUNOV

ABSTRACT. For a positive integer n , let

$$F_n^*(X, Y) = \prod_{k=1}^n \left(X \sin\left(\frac{k\pi}{n}\right) - Y \cos\left(\frac{k\pi}{n}\right) \right).$$

In 2000 Bean and Laugesen proved that for every $n \geq 3$ the area bounded by the curve $|F_n^*(x, y)| = 1$ is equal to $4^{1-1/n} B\left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2}\right)$, where $B(x, y)$ is the beta function. We provide an elementary proof of this fact based on the polar formula for the area calculation. We also prove that

$$F_n^*(X, Y) = 2^{1-n} \sum_{\substack{1 \leq k \leq n \\ k \text{ is odd}}} (-1)^{\frac{k-1}{2}} \binom{n}{k} X^{n-k} Y^k$$

and demonstrate that $\ell_n = 2^{n-1-\nu_2(n)}$ is the smallest positive integer such that the binary form $\ell_n F_n^*(X, Y)$ has integer coefficients. Here $\nu_2(n)$ denotes the 2-adic order of n .

1. Introduction. Let $F(X, Y) = a_0 X^n + a_1 X^{n-1} Y + \dots + a_n Y^n$ be a binary form of degree $n \geq 3$ with complex coefficients and nonzero discriminant D_F . In this article we evaluate the area A_F bounded by the curve

$$|F(x, y)| = 1, \quad (x, y) \in \mathbb{R},$$

associated with a particular family of binary forms F . The quantity A_F arises in the study of *Thue inequalities* $|F(x, y)| \leq h$ where the coefficients of F are integers and h is a fixed positive integer. In 1933 Mahler [5] proved that the number of integer solutions $Z_F(h)$ to the Diophantine inequality above satisfies

$$\left| Z_F(h) - A_F h^{2/n} \right| \ll_F h^{1/(n-1)},$$

2020 *Mathematics Subject Classification*. Primary 11D75, 51M25; Secondary 11J25, 33B15.

Keywords and phrases. Beta function, Diophantine inequality, polar coordinates, area, trigonometry.

Received by the editors February, 1, 2020.

provided that F is irreducible. More recently, Stewart and Xiao [6] proved that the number of integers of absolute value at most h which are represented by F is asymptotic to $C_F h^{2/n}$, where a positive number C_F depends solely on F and is a rational multiple of A_F .

In 1994 Bean [1] proved that the quantity $|D_F|^{1/n(n-1)} A_F$ is bounded by the absolute constant $3B\left(\frac{1}{3}, \frac{1}{3}\right) \approx 15.90$, where

$$(1) \quad B(x, y) = 2 \int_0^{\pi/2} (\sin \theta)^{2x-1} (\cos \theta)^{2y-1} d\theta$$

is the beta function. In his investigation of the quantity

$$M_n = \max \left\{ |D(F)|^{1/n(n-1)} A_F \right\},$$

where the maximum is taken over all binary forms F of degree n with nonzero discriminant, Bean [2] conjectured that the value of M_n is attainable by the binary form

$$F_n^*(X, Y) = \prod_{k=1}^n \left(X \sin\left(\frac{k\pi}{n}\right) - Y \cos\left(\frac{k\pi}{n}\right) \right).$$

The curve $|F_n^*(x, y)| = 1$ has the special property that it is invariant under rotation by any integer multiple of π/n . In 2000 Bean and Laugesen [3] proved that for every $n \geq 3$,

$$(2) \quad A_{F_n^*} = 4^{1-1/n} B\left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2}\right), \quad D_{F_n^*}^{1/n(n-1)} = \frac{1}{2} n^{1/(n-1)}.$$

In Section 3 we prove the first of these equalities via elementary methods. Apart from basic trigonometric identities, our proof involves the identity $\sin(n\theta) = 2^{n-1} \prod_{k=1}^n \sin\left(\frac{k\pi}{n} - \theta\right)$ (established in Section 2), the formula for the area bounded by a curve in polar form; and the trigonometric form of the beta function (1).

In Section 4 we explain what the coefficients of $F_n^*(X, Y)$ look like. More precisely, define the binary form

$$S_n(X, Y) = \ell_n F_n^*(X, Y),$$

where ℓ_n is the smallest positive integer such that S_n has integer coefficients. We prove the following result.

Proposition 1. For any positive integer n ,

$$(3) \quad F_n^*(X, Y) = 2^{1-n} \sum_{\substack{1 \leq k \leq n \\ k \text{ is odd}}} (-1)^{\frac{k-1}{2}} \binom{n}{k} X^{n-k} Y^k$$

and

$$(4) \quad \begin{aligned} S_n(X, Y) &= 2^{n-1-\nu_2(n)} F_n^*(X, Y) \\ &= 2^{-\nu_2(n)} \sum_{\substack{1 \leq k \leq n \\ k \text{ is odd}}} (-1)^{\frac{k-1}{2}} \binom{n}{k} X^{n-k} Y^k, \end{aligned}$$

where $\nu_2(n)$ is the 2-adic order of n .

Finally, in Section 5 we derive the formula for A_{S_n} :

$$A_{S_n} = 4^{\nu_2(n)/n} B\left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2}\right).$$

2. Proof of the Identity $\sin(n\theta) = 2^{n-1} \prod_{k=1}^n \sin\left(\frac{k\pi}{n} - \theta\right)$. Let $U_{n-1}(X)$ be the $(n-1)$ -st Chebyshev polynomial of the second kind:

$$U_{n-1}(X) = 2^{n-1} \prod_{k=1}^{n-1} \left(X - \cos\left(\frac{k\pi}{n}\right) \right).$$

Let $\theta \in \mathbb{R}$. In view of the identity $\sin(n\theta) = U_{n-1}(\cos \theta) \sin \theta$, we obtain

$$\begin{aligned} \sin(n\theta) &= \left[2^{n-1} \prod_{k=1}^{n-1} \left(\cos \theta - \cos\left(\frac{k\pi}{n}\right) \right) \right] \sin \theta \\ &= \left[2^{n-1} \prod_{k=1}^{n-1} \left(2 \sin\left(\frac{k\pi}{2n} - \frac{\theta}{2}\right) \sin\left(\frac{k\pi}{2n} + \frac{\theta}{2}\right) \right) \right] \sin \theta \\ &= \left[2^{n-1} \prod_{k=1}^{n-1} \left(2 \sin\left(\frac{k\pi}{2n} - \frac{\theta}{2}\right) \right) \prod_{k=1}^{n-1} \cos\left(\frac{(n-k)\pi}{2n} - \frac{\theta}{2}\right) \right] \sin \theta \\ &= \left[2^{n-1} \prod_{k=1}^{n-1} \left(2 \sin\left(\frac{k\pi}{2n} - \frac{\theta}{2}\right) \cos\left(\frac{k\pi}{2n} - \frac{\theta}{2}\right) \right) \right] \sin(\pi - \theta) \\ &= 2^{n-1} \prod_{k=1}^n \sin\left(\frac{k\pi}{n} - \theta\right). \end{aligned}$$

3. Proof of the Identity $A_{F_n^*} = 4^{1-1/n} B\left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2}\right)$. Consider the curve $|F_n^*(x, y)| = 1$. The change of variables

$$x = r(\theta) \cos \theta, \quad y = r(\theta) \sin \theta$$

enables us to transform this equation into polar coordinates:

$$\begin{aligned} r(\theta) &= |F_n^*(\cos \theta, \sin \theta)|^{-1/n} \\ &= \left| \prod_{k=1}^n \left(\cos \theta \sin \left(\frac{k\pi}{n} \right) - \sin \theta \cos \left(\frac{k\pi}{n} \right) \right) \right|^{-1/n} \\ &= \left| \prod_{k=1}^n \sin \left(\frac{k\pi}{n} - \theta \right) \right|^{-1/n} \\ &= |2^{1-n} \sin(n\theta)|^{-1/n}. \end{aligned}$$

We are now able to apply a well-known formula for the area bounded by a curve in polar form:

$$A_{F_n^*} = \frac{1}{2} \int_0^{2\pi} r(\theta)^2 d\theta = \frac{1}{2} \int_0^{2\pi} |2^{1-n} \sin(n\theta)|^{-2/n} d\theta = \frac{2^{1-2/n}}{n} \int_0^{2\pi} |\sin \theta|^{-2/n} d\theta.$$

Notice that the function $|\sin \theta|^{-2/n}$ has period π , and furthermore $\int_0^{\pi/2} (\sin \theta)^{-2/n} d\theta = \int_{\pi/2}^{\pi} (\sin \theta)^{-2/n} d\theta$. Hence

$$\begin{aligned} A_{F_n^*} &= \frac{2^{1-2/n}}{n} \cdot 4n \int_0^{\pi/2} (\sin \theta)^{-2/n} d\theta \\ &= 4^{1-1/n} \cdot 2 \int_0^{\pi/2} (\sin \theta)^{-2/n} d\theta \\ &= 4^{1-1/n} B\left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2}\right), \end{aligned}$$

where the last equality follows from (1).

4. Coefficients of $F_n^*(X, Y)$. In this section we prove Proposition 1. If $n = 1$ the result holds, so we assume that $n \geq 2$. Let $f_n^*(X)$ and

$h_n^*(X)$ denote the polynomials

$$f_n^*(X) = \prod_{k=1}^{n-1} \left(\sin\left(\frac{k\pi}{n}\right) X - \cos\left(\frac{k\pi}{n}\right) \right)$$

and

$$h_n^*(X) = 2^{1-n} \operatorname{Im}((X+i)^n),$$

where $\operatorname{Im}(z)$ denotes the imaginary part of a complex number z . It is a consequence of the Binomial Theorem that

$$\begin{aligned} h_n^*(X) &= 2^{1-n} \operatorname{Im} \left(\sum_{0 \leq k \leq n} \binom{n}{k} X^{n-k} i^k \right) \\ &= 2^{1-n} \sum_{\substack{1 \leq k \leq n \\ k \text{ is odd}}} (-1)^{\frac{k-1}{2}} \binom{n}{k} X^{n-k}. \end{aligned}$$

We claim that $f_n^* = h_n^*$. To prove this, we will demonstrate that these two polynomials have the same roots, namely $\cot\left(\frac{k\pi}{n}\right)$ for $k = 1, 2, \dots, n-1$, as well as equal leading coefficients.

First, notice that both f_n^* and h_n^* have degrees $n-1$. Second, if k is an integer between 1 and $n-1$, then $\sin\left(\frac{k\pi}{n}\right) \neq 0$, and so it immediately follows from the definition of f_n^* that $f_n^*\left(\cot\left(\frac{k\pi}{n}\right)\right) = 0$. In turn, De Moivre's formula tells us that

$$\begin{aligned} h_n^*\left(\cot\left(\frac{k\pi}{n}\right)\right) &= 2^{1-n} \operatorname{Im} \left(\left(\cot\left(\frac{k\pi}{n}\right) + i \right)^n \right) \\ &= 2^{1-n} \sin\left(\frac{k\pi}{n}\right)^{-n} \operatorname{Im} \left(\left(\cos\left(\frac{k\pi}{n}\right) + i \sin\left(\frac{k\pi}{n}\right) \right)^n \right) \\ &= 2^{1-n} \sin\left(\frac{k\pi}{n}\right)^{-n} \operatorname{Im}(\cos(k\pi) + \underbrace{i \sin(k\pi)}_0) \\ &= 0. \end{aligned}$$

Therefore, polynomials f_n^* and h_n^* have the same roots. Third, notice that the leading coefficient of $f_n^*(X)$ is $\prod_{k=1}^{n-1} \sin\left(\frac{k\pi}{n}\right)$, while the leading coefficient of $h_n^*(X)$ is $2^{1-n}n$. We use the result established in Section 2 to prove that these coefficients are equal to each other:

$$\prod_{k=1}^{n-1} \sin\left(\frac{k\pi}{n}\right) = \lim_{\theta \rightarrow 0} \prod_{k=1}^{n-1} \sin\left(\frac{k\pi}{n} - \theta\right) = \lim_{\theta \rightarrow 0} 2^{1-n} \frac{\sin(n\theta)}{\sin(\theta)} = 2^{1-n}n.$$

Therefore, $f_n^* = h_n^*$.

Now, let $f_n^*(X, Y)$ and $h_n^*(X, Y)$ denote the homogenizations of $f_n^*(X)$ and $h_n^*(X)$, respectively, so that $f_n^*(X, Y) = h_n^*(X, Y)$. Then $F_n^*(X, Y) = Y f_n^*(X, Y) = Y h_n^*(X, Y)$, which is the same as (3).

It remains to explain why the binary form $S_n = 2^{n-1-\nu_2(n)} F_n^*$ has integer coefficients and why the greatest common divisor of its coefficients is equal to 1. Let

$$g(n) = \gcd_{\substack{1 \leq k \leq n \\ k \text{ is odd}}} \left\{ \binom{n}{k} \right\}.$$

In view of (4), we need show that $g(n) = 2^{\nu_2(n)}$.

Assume for a contradiction that there exists an odd prime p such that $p \mid g(n)$. Let $k = p^r$, where r is the largest positive integer such that $p^r \mid n$. Since k is an odd integer between 1 and n , we conclude that $p \mid \binom{n}{k}$. Letting $\nu_p(m)$ denote the p -adic order of a positive integer m , we conclude that $\nu_p \left(\binom{n}{k} \right) > 0$. One can then use Legendre's formula $\nu_p(m!) = \sum_{j=1}^{\infty} \left\lfloor \frac{m}{p^j} \right\rfloor$ to show that $\nu_p \left(\binom{n}{k} \right) = \nu_p(n!) - \nu_p(k!) - \nu_p((n-k)!) = 0$, which contradicts our observation that the p -adic order of $\binom{n}{k}$ is positive. Therefore, $g(n)$ has no odd prime factors, or in other words $g(n) = 2^s$ for some non-negative integer s .

It remains to show that $s = \nu_2(n)$. Since $g(n) \mid \binom{n}{1} = n$ and $g(n) = 2^s$, we see that $s \leq \nu_2(n)$. To establish the reverse inequality $s \geq \nu_2(n)$, recall the well-known result

$$\frac{n}{\gcd(n, k)} \mid \binom{n}{k},$$

which was attributed by Dickson [4] to Hermite. Now, if k is an odd integer between 1 and n , we see that $\gcd(n, k)$ is odd, which means that $2^{\nu_2(n)} \mid \frac{n}{\gcd(n, k)}$. But then $2^{\nu_2(n)} \mid \binom{n}{k}$. Since k was chosen arbitrarily, we conclude that $2^{\nu_2(n)} \mid g(n)$. Since $g(n) = 2^s$, we see that $s \geq \nu_2(n)$, and so we come to the desired conclusion that $s = \nu_2(n)$.

5. Proof of the Identity $A_{S_n} = 4^{\nu_2(n)/n} B \left(\frac{1}{2} - \frac{1}{n}, \frac{1}{2} \right)$. The formula $A_F = \int_{-\infty}^{+\infty} |F(x, 1)|^{-2/n} dx$ implies that, for any nonzero complex

number c , $A_{cF} = |c|^{-2/n} A_F$. Combining this result with (2) and (4), we find that

$$A_{S_n} = A_{2^{n-1-\nu_2(n)} F_n^*} = 2^{-\frac{2(n-1-\nu_2(n))}{n}} A_{F_n^*} = 4^{\nu_2(n)/n} B\left(\frac{1}{2}, \frac{1}{n}, \frac{1}{2}\right).$$

REFERENCES

1. M.A. Bean, *An isoperimetric inequality for the area of plane regions defined by binary forms*, *Compositio Math.* **92** (1994), 115–131.
2. M.A. Bean, *Binary forms, hypergeometric functions and the Schwarz-Christoffel mapping formula*, *Trans. Amer. Math. Soc.* **347** (1995), 4959–4983.
3. M.A. Bean and R.S. Laugesen, *Binary forms, equiangular polygons and harmonic measure*, *Rocky Mt. J. Math.* **30** (2000), 15–62.
4. L. E. Dickson, *History of the Theory of Numbers*. Vol. 1. Washington, D.C.: Carnegie Institution (1919); rpt. New York: Chelsea (1952).
5. K. Mahler, *Zur Approximation algebraischer Zahlen III*, *Acta Math.* **62** (1933), 91–166.
6. C.L. Stewart and S.Y. Xiao, *On the representation of integers by binary forms*, *Math. Ann.* **375** (2019), 133–163.

UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVE W, WATERLOO ON, CANADA N2L 3G1

Email address: amosunov@uwaterloo.ca