

THE POSET OF COPIES FOR AUTOMORPHISM GROUPS OF COUNTABLE RELATIONAL STRUCTURES

CLAUDE LAFLAMME*, MAURICE POUZET**, NORBERT SAUER*,
AND ROBERT WOODROW*

†Dedicated to the memory of Ivo G. Rosenberg

ABSTRACT. Let G be a subgroup of the symmetric group $\mathfrak{S}(U)$ of all permutations of a countable set U . Let \overline{G} be the topological closure of G in the function topology on U^U . We initiate the study of the poset $\overline{G}[U] := \{f[U] \mid f \in \overline{G}\}$ of images of the functions in \overline{G} , being ordered under inclusion. This set $\overline{G}[U]$ of subsets of the set U will be called the *poset of copies* for the group G . A denomination being justified by the fact that for every subgroup G of the symmetric group $\mathfrak{S}(U)$ there exists a homogeneous relational structure R on U such that \overline{G} is the set of embeddings of the homogeneous structure R into itself and $\overline{G}[U]$ is the set of copies of R in R and that the set of bijections $\overline{G} \cap \mathfrak{S}(U)$ of U to U forms the group of automorphisms of R .

1. INTRODUCTION

Countable sets will be infinite. For a countable set U let $\mathfrak{S}(U)$ be the symmetric group of U and let U^U be the set of functions of U into U and let $\mathcal{P}(U)$ be the power set of U and let $\mathcal{P}_{<\omega}(U)$ be the set of finite subsets of U . If K is a subset of U^U then \overline{K} denotes the adherence

Date: August 15, 2021.

2000 *Mathematics Subject Classification.* Relational Structures, Partially ordered sets and lattices (06A, 06B).

Key words and phrases. homogeneous relational structures, automorphism groups, homogeneity, posets.

*Supported by NSERC of Canada Grant # 690404.

**Research started while the author visited the Mathematics and Statistics Department of the University of Calgary in 2010; the support provided is gratefully acknowledged.

***Supported by NSERC of Canada Grant # 691325.

The first and fourth author were supported by by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon within the program “Investissements d’Avenir (ANR-11-IDEX-0007)” operated by the French National Research Agency (ANR).

of K with respect to the function topology, the set U being equipped with the discrete topology. Then \overline{K} is the set of functions $f \in U^U$ so that for every set $F \in \mathcal{P}_{<\omega}(U)$ there exists a function $g \in K$ for which $f(x) = g(x)$ for all $x \in F$. That is $f \upharpoonright F = g \upharpoonright F$. For a function f let $f[A] = \{f(x) \mid x \in \text{dom}(f) \cap A\}$. For a set of functions K let $K[A] = \{f[A] \mid f \in K\}$. For two functions f and g let the *composition* of f and g be the function $f \circ g$ with $\text{dom}(f \circ g) = \{x \in \text{dom}(g) \mid g(x) \in \text{dom}(f)\}$ and so that if $x \in \text{dom}(f \circ g)$ then $f \circ g(x) = f(g(x))$.

Let G be a subgroup of the symmetric group $\mathfrak{S}(U)$ of a countable set U . Then \overline{G} is the set of functions adherent to G and $\overline{G}[U]$ is the set of images of the functions in \overline{G} . It follows that \overline{G} is a set of injections of U into U closed under composition and that $\overline{G}[U] \subseteq \mathcal{P}(U)$ and that $(\overline{G}[U]; \subseteq)$ is a partially ordered set under the inclusion order \subseteq . For a simple minded example, if $G := \mathfrak{S}(U)$ then \overline{G} is the set of one-to-one maps from U to U . In this case, $\overline{G}[U]$ is the set of subsets of U with the same cardinality as U . The set of functions $f \in \overline{G}$ with $f[U] = U$, that is with $f \in \mathfrak{S}(U)$ forms a subgroup of $\mathfrak{S}(U)$. The group G is *closed* if $G = \overline{G} \cap \mathfrak{S}(U)$. The set of images $\overline{G}[U] = \{g[U] \mid g \in \overline{G}\}$ of the functions in \overline{G} will be called the *set of copies* for the group G . For a more detailed background of this and towards some established notions discussed in the next paragraph and further on we refer the reader to the first about forty pages of [1].

The notions for subgroups of $\mathfrak{S}(U)$ investigated in this article are closely related to notions arising in the study of automorphism groups of relational structures. It is not difficult to verify that $\text{Aut}(\mathbb{R})$ the group of automorphisms of a relational structures \mathbb{R} with countable domain U is closed in $\mathfrak{S}(U)$ for the function topology. Let $\text{Emb}(\mathbb{R})$ be the set of isomorphisms of \mathbb{R} to induced substructures of \mathbb{R} . Define a *copy of* \mathbb{R} to be a subset C of U for which the structure $\mathbb{R} \upharpoonright C$ is isomorphic to \mathbb{R} , that is the range of a member of $\text{Emb}(\mathbb{R})$. Then every copy *for* the group $\text{Aut}(\mathbb{R})$ is a copy *of* \mathbb{R} . That is $\overline{\text{Aut}(\mathbb{R})}[U]$ is a subset of the set of copies of \mathbb{R} . (A function with domain U adherent to $\text{Aut}(\mathbb{R})$ is by definition an isomorphism.) Let for example $\mathbb{N} := (N; \leq)$ be the binary relational structure on the set of natural numbers with \leq the natural order. The automorphism group, say G , of \mathbb{N} contains only the identity map on N . Hence the set of copies for the group G is the set $\{N\}$. On the other hand the set of copies of \mathbb{N} has 2^{\aleph_0} elements.

A relational structure \mathbb{R} is *homogeneous*, see [4] or [5], if every isomorphism between finite induced substructures of \mathbb{R} extends to an automorphism of \mathbb{R} . It follows that if \mathbb{R} is homogeneous then $\overline{\text{Aut}(\mathbb{R})}[U]$ is equal to the set of copies of \mathbb{R} . It is well known, using the standard

back and forth argument, that the order structure \mathbb{Q} of the set of rational numbers Q is homogeneous. Implying that every copy of \mathbb{Q} is also a copy for the automorphism group of \mathbb{Q} . Let G be a subgroup of $\mathfrak{S}(U)$ for a countable set U . Then the canonical structure determined by G , that is the relational structure with domain U whose relations on n -tuples are determined by the orbits of the action of G on the set of n -tuples, is homogeneous. See [1]; or [5] for a more extensive exposition. In particular it follows along the discussion in Section 2.4 of [1] that $\overline{G} \cap \mathfrak{S}(U)$ is a Polish group, see [7], with respect to the function topology and composition as group operation.

A relational structure R with domain U is *prehomogeneous*, (see [13]), if for every finite set $F \subseteq U$ there is a finite set $F' \supseteq F$ such that every isomorphism f of $R \upharpoonright F$ to $R \upharpoonright f[F]$ does then extend to an automorphism, provided that it extends to F' . If the cardinality of F' can be bounded by some function θ depending on the cardinality of F then R is *uniformly prehomogeneous*. It is the case that if a relational structure R is homogeneous or prehomogeneous then $\text{Emb}(R) = \overline{\text{Aut}(R)}$ and the set of copies of R is equal to the set of copies $\overline{\text{Aut}(R)}[U]$ for the group $\text{Aut}(R)$. Actually, if $\text{Aut}(R)$ is oligomorphic and $\text{Emb}(R) = \overline{\text{Aut}(R)}$ then R is prehomogeneous, see [14] Theorem 2.2. A group $G \subseteq \mathfrak{S}(U)$ being *oligomorphic* if for every n the number of orbits of n -element subsets of U is finite. This amounts to the fact, see [16], that the group $\overline{G} \cap \mathfrak{S}(U)$ is the automorphism group of an \aleph_0 -categorical theory.

2. AN OUTLINE OF THE RESULTS

Results on the theme of copies of an homogeneous structure have been developed in a series of papers by Kurilić, Kuzeljević and Todorčević (e.g. [9, 10, 11, 12]). Some of the results presented here have been the subject of a lecture by the third author in March 2019 at the FG1 Seminar of the Technical University of Vienna [18].

For this section let U be a countable set and let G be a subgroup of the symmetric group $\mathfrak{S}(U)$.

Theorem 2.1. *[Theorem 10.2] If the group G is oligomorphic then there is an embedding of $(\mathcal{P}(\omega); \subseteq)$, the power set of ω ordered by inclusion, into $(\overline{G}[U]; \subseteq)$, the set of copies for the group G , ordered by inclusion.*

For F a finite subset of U let $G\langle F \rangle = \{g \in G \mid \forall x \in F (g(x) = x)\}$ and let $\text{ac}(F)$, the *algebraic closure* of F , be the union of all finite point orbits of the stabilizer group $G\langle F \rangle$. If $S \subseteq U$ is infinite let $\text{ac}(S) := \cup\{\text{ac}(F) \mid F \in \mathcal{P}_{<\omega}(S)\}$. Note that if G is oligomorphic then $\text{ac}(F)$ is

finite for every finite subset F of U . A group G of permutations of U is *algebraically finite* if the algebraic closure of every finite subset of U is finite. Oligomorphic groups are algebraically finite. There are many others.

Theorem 2.2. *[Theorem 8.2] If G is algebraically finite then for every finite subset F of U the algebraic closure of F , $\mathbf{ac}(F)$, is the intersection of two copies for the group G .*

Problem 2.1. *Is it true that $\mathbf{ac}(S)$ is an intersection of the set of copies containing S if S is infinite and the group G is oligomorphic?*

Problem 2.2. *If G is algebraically finite, is it true that $(\overline{G}[U]; \subseteq)$ embeds $(\mathcal{P}(\omega); \subseteq)$?*

Theorem 2.3. *[Theorem 5.1] A subset A of U is a copy, that is belongs to $\overline{G}[U]$, if and only if for every finite subset F of A and every $x \in U \setminus F$ there is a function $g \in G\langle F \rangle$ with $g(x) \in A$.*

Using Theorem 2.3 we obtain:

Corollary 2.1. *[Lemma 5.3, Lemma 5.2, Corollary 3.3] The set of copies $\overline{G}[U]$ is closed under the union of non-empty up-directed families. Topologically, the set of copies $\overline{G}[U]$ is a G_δ -set of $\mathcal{P}(U)$ equipped with the powerset topology. (See Definition 3.1.) Moreover, if $\overline{G}[U]$ contains more than one copy, then it has no isolated point with respect to the powerset topology.*

Corollary 2.1 will then be used to establish that:

Theorem 2.4. *[Theorem 5.2] The cardinality of $\overline{G}[U]$ is either 1 or 2^{\aleph_0} .*

Problem 2.3. *Is it true that if $\overline{G}[U]$ has more than one element then the partial ordered set $(\overline{G}[U]; \subseteq)$ is not totally ordered by inclusion? Is it true that it embeds $(\mathcal{P}(\omega); \subseteq)$?*

In Section 7 we will introduce the notions of being ranked or un-ranked for point orbits of the stabilizer subgroups $G\langle F \rangle$ for finite subsets F of U . Assigning an ordinal number rank to the ranked point orbits of those stabilizer subgroups. Leading to the notion of *ranked closure* $\mathbf{rc}(S)$ of S , for subsets S of U . It turns out, Lemma 8.2, that if the group G is algebraically finite then $\mathbf{rc}(S) = \mathbf{ac}(S)$ for every subset S of U .

Theorem 2.5. *[Theorem 7.1] The ranked closure of a finite set F is equal to the intersection of all copies containing F . There is just one*

copy, namely U , for a group $G \subseteq \mathfrak{S}(U)$, if and only if the ranked closure of the empty set is U if and only if every point orbit of the group G is ranked.

We do not know if a more general version of Theorem 2.2 holds for all groups G with $\overline{G}[U] \neq \{U\}$. The next Theorem, being just of an enough strengthening of Theorem 2.5 in order to establish Theorem 2.2, does not seem to be sufficiently strong to lead to a generalization of Theorem 2.2. Note that Theorem 2.2 implies that $\overline{G}[U] \neq \{U\}$ if the group G is algebraically finite.

Theorem 2.6. *[Theorem 8.1] Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$ and $\overline{G}[U] \neq \{U\}$. Then for every finite subset F of U and every copy $C \in \overline{G}[U]$ with $F \subseteq C$ (hence $\mathbf{rc}(F) \subseteq C$), there exists: An ω -sequence $C = C_0 \supset C_1 \supset C_2 \supset C_3 \supset \dots$ of copies $C_i \in \overline{G}[U]$ for which $\mathbf{rc}(F) = \bigcap_{i \in \omega} C_i$.*

For the connection between the Hausdorff rank of a linear order and the rank defined in this article, see Subsection 7.1.

Besides the set of copies $\overline{G}[U]$, there are three related sets of interest: the set of intersections of copies $\widehat{\overline{G}[U]}$, the set $\overline{G}^{\mathbf{rc}}[U]$ of ranked closures of arbitrary subsets of U and the topological closure $\overline{\overline{G}[U]}$ of $\overline{G}[U]$ in $\mathcal{P}(U)$.

Lemma 2.1. $\overline{G}[U] \subseteq \widehat{\overline{G}[U]} \subseteq \overline{G}^{\mathbf{rc}}[U] = \overline{\overline{G}[U]}$.

The content of this lemma is in the equality $\overline{G}^{\mathbf{rc}}[U] = \overline{\overline{G}[U]}$. Indeed, the first inclusion is obvious. The family $\overline{G}^{\mathbf{rc}}[U]$ defines an algebraic closure system. As a closure system, it is closed under intersection, since it contains $\overline{G}[U]$ it contains $\widehat{\overline{G}[U]}$. Being algebraic, it is closed in $\mathcal{P}(U)$ equipped with the product topology. Since it contains $\overline{G}[U]$, it contains its topological closure, that is $\overline{\overline{G}[U]}$. It remains to show that $\overline{G}^{\mathbf{rc}}[U] \subseteq \overline{\overline{G}[U]}$. Note that a subset S of U is in $\overline{G}[U]$ if for every finite subset $F \subseteq S$ and every finite subset E of $U \setminus S$ there exists a copy $C \in \overline{G}[U]$ for which $F \subseteq C$ and $E \cap C = \emptyset$. Thus, let $S \in \overline{G}^{\mathbf{rc}}[U]$. Let $F \subseteq S$ be a finite subset, and let E be a finite subset of $U \setminus S$. Since F is finite, $\mathbf{rc}(F)$ is equal to $\mathbf{ic}(F)$, the intersection of copies containing F (Theorem 2.5), hence for every $x \notin F$ there is a copy C containing F and not x . In fact, there is some copy C containing F and no element of E (see Lemma 3.2). Hence, $\mathbf{ic}(F) \in \overline{\overline{G}[U]}$.

The fact that $\overline{\overline{G}[U]} = \overline{G}^{\mathbf{rc}}[U]$ has the following significant consequence:

Corollary 2.2. $\overline{\overline{G[U]}}$, the topological closure of the set of copies, is closed under intersections.

As an other consequence, we have:

Theorem 2.7. [Theorem 9.1] Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then $\mathbf{rc}(S)$ is the least element of $\overline{\overline{G[U]}}$ for every subset S of U .

The intersection of the copies containing a finite set is equal to the ranked closure of this finite set. We do not know if this is also the case for infinite subsets of U . That is we cannot decide:

Problem 2.4. Is it true that $\widehat{\overline{G[U]}} = \overline{\overline{G[U]}}$?

But, Lemma 9.2 states that $\widehat{\overline{G[U]}}$ is an algebraic closure system if and only if $\widehat{\overline{G[U]}} = \overline{\overline{G[U]}}$.

3. ELEMENTARY PROPERTIES

Let U be a countable set and let G be a subgroup of the symmetric group $\mathfrak{S}(U)$.

Definition 3.1. For $\mathcal{Q} \subseteq \mathcal{P}(U)$ and $S \cup T \subseteq U$ let $\mathcal{Q}[[S]] = \{V \in \mathcal{Q} \mid S \subseteq V\}$ and let $\mathcal{Q}[[S, T]] = \{V \in \mathcal{Q} \mid S \subseteq V \text{ and } T \cap V = \emptyset\}$.

For E and F two finite subsets of U the set $\mathcal{P}(U)[[F, E]]$ is a basic neighbourhood for the powerset topology.

Of course $\overline{G[U]} \subseteq \mathcal{P}(U)$. Then $\overline{G[U]}[[F]]$, for $F \subseteq U$, denotes the set of copies for the group G which contain F as a subset. The set $\overline{G[U]}[[F, E]]$ is then a neighbourhood for the powerset topology relative to the set of copies $\overline{G[U]}$. We investigate the set of copies $\overline{G[U]}$ for the group G with respect to the order \subseteq on $\mathcal{P}(U)$ and with respect to the powerset topology on $\mathcal{P}(U)$. Then $\overline{\overline{G[U]}}$ is the set of subsets of U which are in the topological closure of $\overline{G[U]}$ the set of copies for the group G and $(\overline{G[U]}; \subseteq)$ is the poset of copies and $\widehat{\overline{G[U]}}$ is the set of intersections of copies. The hypergraph of copies is the hypergraph $H(\overline{G}) := (U, \overline{G[U]})$ whose hyperedges are the copies.

Definition 3.2. Let X be a subset of U^U and F be a subset of U . Then $X\langle F \rangle := \{f \in X \mid f \upharpoonright F = \text{Id} \upharpoonright F\}$ denotes the stabilizer of F for the set X of functions and $X_F = \{f \in X \mid f[F] = F\}$ denotes the setwise stabilizer of F for the set X of functions. Also, set \overline{X} the topological closure of X in U^U equipped with the product topology.

Lemma 3.1. Let $X \subseteq U^U$ and $F \in \mathcal{P}_{<\omega}(U)$. Then

- (1) $\overline{X}\langle F \rangle = \overline{X}\langle \overline{F} \rangle$.
(2) $\overline{X}_F = \overline{X}_F$.

Proof. The sets of functions $\overline{X}\langle F \rangle$ and \overline{X}_F are closed because \overline{X} is closed and F is finite. Hence $\overline{X}\langle \overline{F} \rangle \subseteq \overline{X}\langle F \rangle$ and $\overline{X}_F \subseteq \overline{X}_F$. Conversely, let $f \in \overline{X}\langle F \rangle$. Let $H \in \mathcal{P}_{<\omega}(U)$. Since $f \in \overline{X}$ there is some $g \in X$ such that $g \upharpoonright (F \cup H) = f \upharpoonright (F \cup H)$. Since $g \in X\langle F \rangle$ and f and g coincide on H this implies $f \in \overline{X}\langle \overline{F} \rangle$.

Let $f \in \overline{X}_F$ and let $H \in \mathcal{P}_{<\omega}(U)$. Since $f \in \overline{X}$ there is some $g \in X$ such that $g \upharpoonright (F \cup H) = f \upharpoonright (F \cup H)$. Since $g \in X_F$ and f and g coincide on H this implies $f \in \overline{X}_F$. \square

Lemma 3.2. *If F is a finite subset of a copy $C \in \overline{G}[U]$ then there exists a function $f \in \overline{G}$ with $f[U] = C$ and $f(a) = a$ for all $a \in F$. That is: $\overline{G}\langle F \rangle[U] = \overline{G}\langle F \rangle[U] = \overline{G}[U][F]$ for every $F \in \mathcal{P}_{<\omega}(U)$.*

Proof. The following inclusion for every subset F of U (finite or not) is obvious: $\overline{G}\langle F \rangle[U] \subseteq \overline{G}[U][F]$. We claim that the reverse inclusion holds for every $F \in \mathcal{P}_{<\omega}(U)$. Let $C \in \overline{G}[U][F]$. Let $g \in \overline{G}$ such that $g[U] = C$. Let F' be such that $g(F') = F$. Then there is an $h \in \overline{G}$ such that $h \upharpoonright F' = g \upharpoonright F'$. Set $f := g \circ h^{-1}$. Then $C = f[U]$ and $f \in \overline{G}\langle F \rangle$ implying $C \in (\overline{G}\langle F \rangle)[U]$. The claim follows. The equality follows too. \square

Corollary 3.1. $(\overline{G}\langle F \rangle)[U] = \overline{G}_F[U] = \overline{G}[U][F]$ for every $F \in \mathcal{P}_{<\omega}(U)$.

Corollary 3.2. *If F is a finite subset of a copy $C \in \overline{G}[U]$ then $\mathbf{ic}(F)$, the intersection of copies containing F is equal to $\mathbf{ic}_C(F)$ the intersection of copies containing F and included into C . Furthermore, if E is a finite set disjoint from $\mathbf{ic}(F)$, there is a copy containing F and disjoint from E .*

Proof. Clearly, $\mathbf{ic}(F) \subseteq \mathbf{ic}_C(F) \subseteq C$. Let $x \in C \setminus \mathbf{ic}(F)$. Let $C' \in \overline{G}[U]$. Set $F' := F \cup \{x\}$. Apply Lemma 3.2 to C and F' . There is some $f \in \overline{G}\langle F' \rangle$ such that $f[U] = C$. Then $C'' := f[C']$ is a copy containing F included in C and not containing x , hence $x \notin \mathbf{ic}(F)$. Now, we prove that there is some copy D containing F and disjoint from E . Indeed, pick $x \in E$, then some copy containing F , say D , avoids x . The property above ensures that $\mathbf{ic}(F)$, the intersection of copies containing F is equal to $\mathbf{ic}_D(F)$, the intersection of copies containing F that are included into D . If D contains some element y of E we may find a copy D' included into D and avoiding it. Repeating the process, we find a copy containing no element of E . \square

Lemma 3.3. *Let G be a subgroup of $\mathfrak{S}(U)$ and let $\{f, g\} \subseteq \overline{G}$ with $f[U] \subseteq g[U]$. Then $g^{-1} \circ f \in \overline{G}$. If $f[U] = g[U]$ then $g^{-1} \circ f \in \overline{G} \cap \mathfrak{S}(U)$.*

Proof. Let $F \in \mathfrak{P}_{<\omega}(U)$. There exists a function $k \in G$ with $k \upharpoonright F = f \upharpoonright F$ and there exists a function $h \in G$ with $h \upharpoonright (g^{-1}[k[F]]) = g \upharpoonright (g^{-1}[k[F]])$. Implying that $(h^{-1} \circ k) \upharpoonright F = (g^{-1} \circ f) \upharpoonright F$ with $h^{-1} \circ k \in G$. Let $f[U] = g[U]$. Then $g^{-1} \circ f$ maps U onto U . \square

Lemma 3.4. *Let G be a subgroup of $\mathfrak{S}(U)$. Then the following properties are equivalent:*

- (i) $\overline{G}[U] = \{U\}$;
- (ii) $\overline{G} \subseteq \mathfrak{S}(U)$;
- (iii) $\overline{G}\langle F \rangle \subseteq \mathfrak{S}(U)$ for every $F \in \mathfrak{P}_{<\omega}(U)$;
- (iv) $\overline{G}\langle F \rangle \subseteq \mathfrak{S}(U)$ for some $F \in \mathfrak{P}_{<\omega}(U)$.

Proof. The equivalence between (i) and (ii) and the implications (ii) \Rightarrow (iii) \Rightarrow (iv) are obvious. Suppose that (iv) holds. We will prove that (ii) holds. Let $f \in \overline{G}$. Let $F \in \mathfrak{P}_{<\omega}(U)$ such that $\overline{G}\langle F \rangle \subseteq \mathfrak{S}(U)$ and let $g \in G$ such that $g \upharpoonright F = f \upharpoonright F$. Let $k := g^{-1} \circ f$. Since $\{g^{-1}, f\} \subseteq \overline{G}$ the function $k \in \overline{G}$, actually $k \in \overline{G}\langle F \rangle$. Implying $k \in \mathfrak{S}(U)$ according to Lemma 3.1. Hence $k \in \mathfrak{S}(U)$. Consequently $f = g \circ k \in \mathfrak{S}(U)$. \square

It follows that if the stabilizer $G\langle F \rangle$ of G for some finite subset F of U contains only the identity function then the group G is closed in $\mathfrak{S}(U)$.

Lemma 3.5. *The copy U is isolated, for the powerset topology relative to the set of copies $\overline{G}[U]$ if and only if $\overline{G}[U] = \{U\}$.*

Proof. The set U is isolated if and only if $\overline{G}[U][F] = \{U\}$ for some $F \in \mathfrak{P}_{<\omega}(U)$. According to Lemma 3.2 and Lemma 3.1 $\{U\} = \overline{G}[U][F]$ if and only if $\{U\} = (\overline{G}\langle F \rangle)[U] = (\overline{G}\langle F \rangle)[U]$ if and only if $\mathfrak{S}(U) \supseteq \overline{G}\langle F \rangle$ if and only if $\overline{G}[U] = \{U\}$ according to Lemma 3.4. \square

Lemma 3.6. *For every copy $V \in \overline{G}[U] \setminus \{U\}$ and subset $F \in \mathfrak{P}_{<\omega}(V)$ there exists a copy $W \in \overline{G}[U]$ with $F \subseteq W \subset V$.*

Proof. Let $V \in \overline{G}[U] \setminus \{U\}$ and $F \in \mathfrak{P}_{<\omega}(V)$. According to Lemma 3.2 there exists a function $g \in \overline{G}\langle F \rangle$ with $g[U] = V$. Set $W := g[V]$. \square

Corollary 3.3. *The set of copies $\overline{G}[U]$ has no isolated point for the powerset topology if and only if $\overline{G}[U] \neq \{U\}$.*

Proof. Suppose that a copy V is isolated in $\overline{G}[U]$. Let $f \in \overline{G}$ be a function with $f[U] = V$. The set of copies $\overline{G}[U][[F, E]]$, containing F and being disjoint from E , is open for every $\{F, E\} \subseteq \wp_{<\omega}(U)$. Because V is isolated, we may select F, E so that $\overline{G}[U][[F, E]] = \{V\}$. Let $F' := f^{-1}(F)$. Then $\overline{G}[U][[F']] = \{U\}$. For otherwise there exists a copy W with $F' \subseteq W \subset U$. Then $F \subseteq f[W] \subset V$ with $f[W] \cap E \subseteq V \cap E = \emptyset$; in contradiction to $\overline{G}[U][[F, E]] = \{V\}$. The equality $\overline{G}[U][[F']] = \{U\}$ implies, according to Lemma 3.5, that $\overline{G}[U] = \{U\}$. \square

4. ORBITS AND TYPES AND TYPESETS

Let U be a set and G be a subgroup of $\mathfrak{S}(U)$.

In order to indicate an orbit of the action of the group G it suffices to choose any element in the orbit. Let $\text{Orb}(G)$ denote the set of different orbits of the action of G on U . The orbits of the action of the stabilizer subgroups of the group G will play a role and we will need some particular notation for them. According to Definition 3.2 for $F \subseteq U$ the F -stabilizer of G will be denoted by $G\langle F \rangle$, that is:

$$G\langle F \rangle = \{g \in G \mid \forall x \in F (g(x) = x)\}.$$

Then the set of orbits, denoted $\text{Orb}(G\langle F \rangle)$, of the subgroup $G\langle F \rangle$ of G acting on U forms a partition of the set U . We emphasize the fact that the orbit of any element x of F w.r.t. the action of $G\langle F \rangle$ is $\{x\}$. A *type* is a pair of the form $\langle F|p \rangle$ for which F is a finite subset of U and $p \in U$. The set F is the *socket* of the type $\langle F|p \rangle$. Two types $\langle F|p \rangle$ and $\langle E|q \rangle$ are *equal* if $F = E$ and if there exists a function $g \in G\langle F \rangle$ with $g(p) = q$. The *typeset*, denoted $G\langle F|p \rangle$, of the type $\langle F|p \rangle$, is the set in $\text{Orb}(G\langle F \rangle)$ given by:

$$G\langle F|p \rangle := \{x \in U \mid \exists g \in G\langle F \rangle (g(p) = x)\}.$$

Note that for all $\{x, y\} \subseteq G\langle F|p \rangle$ there exists a function $f \in G\langle F \rangle$ with $f(x) = y$. It follows that two types $\langle F|p \rangle$ and $\langle E|q \rangle$ are *equal* if and only if $F = E$ and if $G\langle F|p \rangle = G\langle E|q \rangle$. That is, the set of typesets of the types with socket F is the set $\text{Orb}(G\langle F \rangle)$. Two typesets having the same socket are either disjoint or equal. Note:

Fact 4.1. *Let $\langle F|p \rangle$ be a type. Then*

$$G\langle F|p \rangle = \{x \in U \mid \exists f \in \overline{G}\langle F \rangle (f(p) = x)\}.$$

The type $\langle E|q \rangle$ is a *continuation* of the type $\langle F|p \rangle$ if $F \subseteq E$ and if $q \in G\langle F|p \rangle$. It follows that the type $\langle E|q \rangle$ is a continuation of the type $\langle F|p \rangle$ if and only if $F \subseteq E$ and $G\langle F|p \rangle \supseteq G\langle E|q \rangle$. Note:

Fact 4.2. *Let $G \subseteq \mathfrak{S}(U)$. Let $\langle F|p \rangle$ be a type and let $E \in \mathcal{P}_{<\omega}(U)$ with $F \subseteq E$. Then the typesets of the types $\langle E|q \rangle$ which are continuations of the type $\langle F|p \rangle$ form a partition of the typeset $G\langle F|p \rangle$. That is:*

$$G\langle F|p \rangle = \bigcup_{q \in G\langle F|p \rangle} G\langle E|q \rangle.$$

Noting that either $G\langle E|q \rangle = G\langle E|q' \rangle$ or $G\langle E|q \rangle \cap G\langle E|q' \rangle = \emptyset$ and that a type $\langle E|q \rangle$ with $F \subseteq E$ is a continuation of the type $\langle F|p \rangle$ if and only if $q \in G\langle F|p \rangle$.

For $1 \leq n \in \omega$: Let U^n denote the set of n -tuples with entries in U . Let G_n be the subgroup of $\mathfrak{S}(U^n)$ induced by the action of the group G on the set U^n . Let $\mathcal{P}_n(U)$ denote the set of n -element subsets of U . Let $G_{(\mathcal{P}_n)}$ be the subgroup of $\mathfrak{S}(\mathcal{P}_n(U))$ induced by the action of the group G on the set $\mathcal{P}_n(U)$.

Lemma 4.1. *The following properties are equivalent:*

- (1) *The set $\text{Orb}(G\langle F \rangle)$ is finite for every $F \in \mathcal{P}_{<\omega}(U)$.*
- (2) *The set $\text{Orb}(G_n)$ is finite for every $1 \leq n \in \omega$.*

Proof. Note: $\text{Orb}(G\langle \emptyset \rangle) = \text{Orb}(G) = \text{Orb}(G_1)$. The equivalence of conditions (1) and (2) follows then via induction on n and the fact that two $n+1$ tuples $\vec{x} = (x_0, x_1, \dots, x_{n-1}, x_n) \in U^{n+1}$ and $\vec{y} = (y_0, y_1, \dots, y_{n-1}, y_n) \in U^{n+1}$ are in the same orbit of G_{n+1} if and only if there exists an n -tuple $(u_0, u_1, \dots, u_{n-1}) \in U^n$ with functions g and f in G so that $g(x_i) = u_i = f(y_i)$ for all $i \in n+1$ and so that the typesets $G\langle \{u_0, u_1, \dots, u_{n-1}\} | g(x_n) \rangle$ and $G\langle \{u_0, u_1, \dots, u_{n-1}\} | f(y_n) \rangle$ are equal. \square

Definition 4.1. *If the second condition of Lemma 4.1 is satisfied the group is said to be oligomorphic.*

Lemma 4.2. *Let $\langle F|x \rangle$ be a type. If $f \in \overline{G}$ then $G\langle f[F] | f(x) \rangle \supseteq f[G\langle F|x \rangle]$. If in addition the function f is onto U , that is $f \in \overline{G} \cap \mathfrak{S}(U)$, then $G\langle f[F] | f(x) \rangle = f[G\langle F|x \rangle]$.*

Proof. Let $y \in G\langle F|x \rangle$. There exists a function $g \in G\langle F \rangle$ with $g(x) = y$. Then $(f \circ g \circ f^{-1})(f(x)) = f(y)$ and $(f \circ g \circ f^{-1})(f(z)) = f(z)$ for all $z \in F$. Hence $f(y) \in G\langle f[F] | f(x) \rangle$ and $G\langle f[F] | f(x) \rangle \supseteq f[G\langle F|x \rangle]$.

Let $f \in \overline{G} \cap \mathfrak{S}(U)$. Let $y \in G\langle f[F] | f(x) \rangle$. There exists a function $h \in G\langle f[F] \rangle$ with $y = h(f(x))$. Then $(f^{-1} \circ h \circ f)(z) = z$ for all $z \in F$, and $f(f^{-1} \circ h \circ f)(x) = f \circ f^{-1} \circ h \circ f(x) = h \circ f(x) = y$. Hence $y \in f[G\langle F|x \rangle]$ and $G\langle f[F] | f(x) \rangle \subseteq f[G\langle F|x \rangle]$. \square

Corollary 4.1. *Let $\langle F|x \rangle$ be a type whose typeset $G\langle F|x \rangle$ is finite. Then $G\langle f[F]|f(x) \rangle = f[G\langle F|x \rangle]$ for all functions $f \in \overline{G}$. The set $G\langle F|x \rangle$ is a subset of all copies $C \in \overline{G}[U]$ with $F \subseteq C$.*

Proof. The first assertion of the Corollary follows from Lemma 4.2 because there exists a function $h \in G$ for which $h \upharpoonright (F \cup G\langle F|x \rangle) = f \upharpoonright (F \cup G\langle F|x \rangle)$. Let $C \in \overline{G}[U]$ with $F \subseteq C$. It follows from Lemma 3.2 that there exists a function $f \in \overline{G}\langle F \rangle$, hence with $f[F] = F$, for which $f[U] = C$. Then $G\langle F|x \rangle = G\langle f[F]|f(x) \rangle = f[G\langle F|x \rangle] \subseteq C$. The two types $\langle F|x \rangle$ and $\langle f[F]|f(x) \rangle$ are equal because there exist a function $g \in G$ with $g \upharpoonright (F \cup \{x\}) = f \upharpoonright (F \cup \{x\})$. That is $g \in G\langle F \rangle$ with $h(x) = f(x)$.

□

5. A CHARACTERIZATION OF MEMBERS OF $\overline{G}[U]$

Theorem 5.1. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. A subset C of U belongs to $\overline{G}[U]$ if and only if for every $F \in \mathcal{P}_{<\omega}(C)$ and every $x \in U \setminus F$ there is a function $g \in G\langle F \rangle$ such that $g(x) \in C$.*

Proof. Let $(*)$ be the following statement: $(*)$: For every $F \in \mathcal{P}_{<\omega}(C)$ and every $x \in U \setminus F$ there is a function $g \in G\langle F \rangle$ such that $g(x) \in C$.

Let $C \in \overline{G}[U]$ and let $F \in \mathcal{P}_{<\omega}(C)$. Let $x \in U \setminus F$. According to Lemma 3.2 there exists a function $h \in \overline{G}\langle F \rangle$ with $h[U] = C$. Hence $h(x) \in C$. Since $h \in \overline{G}\langle F \rangle$ there exists a function $g \in G\langle F \rangle$ such that $h \upharpoonright (F \cup \{x\}) = g \upharpoonright (F \cup \{x\})$. Hence $g(x) \in C$. We conclude that property $(*)$ holds.

Given property $(*)$ we make use of the standard back and forth method in order to prove that there is a function $f \in \overline{G}$ with $f[U] = C$. (See Fraïssé's characterization of homogeneous structures, [5].) Let $\{u_i \mid i \in \omega\}$ be an ω -enumeration of U and let $U_n = \{u_i \mid i \in n\}$. Let $\{c_i \mid i \in \omega\}$ be an ω -enumeration of C and let $C_n = \{c_i \mid i \in n\}$. For $n \in \omega$ we will construct recursively sets $V_n \subseteq U$ and functions $l_n \in G$ with $l_n[V_n] \subseteq C$ and with $f_n := l_n \upharpoonright V_n \subseteq l_{n+1} \upharpoonright V_{n+1} := f_{n+1}$. Furthermore $U_n \subseteq V_n$ and $C_n \subseteq l_n[V_n]$ for every $n \in \omega$. Then $f := \bigcup_n f_n \in \overline{G}$ with $f[U] = C$.

Let $V_0 = \emptyset$ and $l_0 = \emptyset$. Given the set V_n and the function $l_n \in G$ let $i \in \omega$ be the smallest number for which $u_i \notin V_n$. Let $j \in \omega$ be the smallest number for which $c_j \notin l_n[V_n]$. Using property $(*)$ let $g \in G\langle l_n[V_n] \rangle$ with $g(l_n(u_i)) \in C$. Let $V_{n+1} = V_n \cup \{u_i, l_n^{-1}(g^{-1}(c_j))\}$ and let $l_{n+1} = g \circ l_n$.

□

Corollary 5.1. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. A subset C of U belongs to $\overline{G}[U]$ if and only if $C \cap G\langle F|x \rangle \neq \emptyset$ for every type $\langle F|x \rangle$ of G with $F \subseteq C$.*

Lemma 5.1. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. If for every $F \in \mathfrak{P}_{<\omega}(U)$ every orbit of the action of $G\langle F \rangle$ on $U \setminus F$ is infinite then there is a coinfinite subset A of U such that $A' \in \overline{G}[U]$ for every subset A' of U which contains A . In particular, there is an embedding of $\mathfrak{P}(\omega)$ ordered by inclusion into $\overline{G}[U]$ ordered by inclusion.*

Proof. Let $Type(G) := \{Orb(x, G\langle F \rangle) : F \in [U \setminus \{x\}]^{<\omega} \text{ and } x \in U\}$. This is a countable set of subsets of U . According to our hypothesis, each one is infinite. Bernstein's property ensures that there is a partition of U into two infinite parts A and B such that each member of $Type(G)$ meets A and B (for this general property of sets, replace the continuum in Lemma 2, p.514, of [8] by \aleph_0). If A' is any subset of U containing A then each member of $Type(G)$ will meet A' . In particular, the condition of Corollary 5.1 holds, hence $A' \in \overline{G}[U]$. \square

Lemma 5.2. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then $\overline{G}[U]$ is a G_δ -set of $\mathfrak{P}(U)$ equipped with the powerset topology.*

Proof. A base for the powerset topology consists of a set of subsets of the form $\mathfrak{P}(U)[[F, E]] = \{A \subseteq U \mid F \subseteq A \text{ and } E \cap A = \emptyset\}$ with F and E ranging over finite subsets of U . Let $\langle F | p \rangle$ be a type of G . Then

$$O(\langle F | p \rangle) := \bigcup_{a \in F} \mathfrak{P}(U)[[\emptyset, \{a}]] \cup \bigcup_{x \in G\langle F | p \rangle} \mathfrak{P}(U)[[F \cup \{x\}, \emptyset]]$$

is an open subset of $\mathfrak{P}(U)$. Then the intersection of the subsets $O(\langle F | p \rangle)$ of $\mathfrak{P}(U)$ with $\langle F | p \rangle$ ranging over the countable set of types of G is a G_δ -set. It follows from Corollary 5.1 that

$$\bigcap_{\langle F | p \rangle \in \text{Types}(G)} O(\langle F | p \rangle) = \overline{G}[U].$$

\square

Theorem 5.2. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then the cardinality of $\overline{G}[U]$ is either 1 or 2^{\aleph_0} .*

Proof. According to Lemma 3.3 if $\overline{G}[U]$ has more than one element, it has no isolated point. Since $\overline{G}[U]$ is a G_δ -subset of the Polish space $\mathfrak{P}(U)$ it has the cardinality of the continuum. (See for example Theorem (6.2) of [7].) \square

Lemma 5.3. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then $\overline{G}[U]$ is closed under the union of non-empty up-directed families of members of $\overline{G}[U]$.*

Proof. Let $\mathcal{C} := (C_i; i \in I)$ with $I \neq \emptyset$ be an up-directed family of members of $\overline{G}[U]$. Let $C := \bigcup_{i \in I} C_i$. We claim that $C \in \overline{G}[U]$. For that

we apply Theorem 5.1. Let $F \in \mathcal{P}_{<\omega}(C)$ and $x \in U \setminus F$. We have to show that there is a $g \in G\langle F \rangle$ such that $g(x) \in C$. This is immediate. Since \mathcal{C} is up-directed, there is an $i \in I$ such that $F \subseteq C_i$. Since $C_i \in \overline{G}[U]$ there is a $g_i \in G\langle F \rangle$ such that $g_i(x) \in C_i$. Since $C_i \subseteq C$, $g_i(x) \in C$. It suffices to set $g := g_i$. \square

Corollary 5.2. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then for every subset F of U and for every copy $C \in \overline{G}[U]$ with $F \cap C = \emptyset$ there exists a maximal copy $C' \in \overline{G}[U]$ with $C \subseteq C'$ and with $F \cap C' = \emptyset$.*

Definition 5.1. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Let $f \in \overline{G}$ and $h \in \overline{G}$. Then $f(h) := f \circ h \circ f^{-1}$ and $f[\overline{G}] := \{f(h) \mid h \in \overline{G}\}$ and $f[G] := \{f(g) \mid g \in G\}$.*

Lemma 5.4. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Let $f \in \overline{G}$. Then:*

- (1) $f(k) \circ f(h) = f(k \circ h)$ for all $\{k, h\} \subseteq \overline{G}$.
- (2) The function $f(g)$ is a bijection of $f[U]$ onto $f[U]$ with inverse $f(g^{-1})$ and $f[G]$ is a subgroup of the symmetric group $\mathfrak{S}(f[U])$.
- (3) $f : U \rightarrow f[U]$ is an isomorphism of the group action of G mapping every $g \in G$ to $f(g)$. It is also an isomorphism of the monoid action of \overline{G} mapping every $h \in \overline{G}$ to $f(h)$.
- (4) $C \in (f[\overline{G}])(f[U])$ if and only if $C \in \overline{G}[U]$ and $C \subseteq f[U]$.

Proof.

(1): Let $\{k, h\} \subseteq \overline{G}$. Then $(f(k) \circ f(h)) = f \circ k \circ f^{-1} \circ f \circ h \circ f^{-1} = f \circ (k \circ h) \circ f^{-1} = f(k \circ h)$.

(2): Let $g \in G$ and $x \in f[U]$. Clearly $f(g) \in (f[U])^{f[U]}$. It follows from (1) that $f(g) \circ f(g^{-1}) = f(g \circ g^{-1}) = f(\text{id}_U) = \text{id}_{f[U]}$ and hence $f(g)(f(g^{-1})(x)) = x$. Implying that the function $f(g)$ is a bijection of $f[U]$ onto $f[U]$ with inverse $f(g^{-1})$. Hence $f[G]$ is a subgroup of the symmetric group $\mathfrak{S}(f[U])$.

(3): It remains to prove that if $\{k, h\} \subseteq \overline{G}$ with $k \neq h$ then $f(k) \neq f(h)$. Let $x \in U$ with $k(x) \neq h(x)$. If $(f(k))(f(x)) = (f(h))(f(x))$ then $(f \circ k \circ f^{-1})(f(x)) = (f \circ h \circ f^{-1})(f(x))$ and $f \circ k(x) = f \circ h(x)$. The function f is one-to-one on U and hence the contradiction $k(x) = h(x)$. It follows that f is one-to-one on \overline{G} .

(4): Let $V = f[U]$ and $C \in (f[\overline{G}])(V)$. Then $C \subseteq V$ and there exists a function $h \in \overline{G}$ with $f(h)[V] = C$. Hence $(f \circ h)[U] = (f \circ h \circ f^{-1})[V] = C$. Which in turn implies that $C \in \overline{G}[U]$. Let then $C \in \overline{G}[U]$ and $C \subseteq V$. Let $h \in \overline{G}$ with $h[U] = C$. Let F be a finite subset of U . Let $k = f^{-1} \circ h$. There exists a function $g \in G$ for which $g \upharpoonright F = h \upharpoonright F$. There

exists a function $l \in G$ with $f \uparrow (f^{-1}(h[F])) = l \uparrow (f^{-1}(h[F]))$. Then $k \uparrow F = (l^{-1} \circ g) \uparrow F$. We conclude that $k \in \overline{G}$ and that $f(k)[f[U]] = f(k)[V] = (f \circ k \circ f^{-1})[V] = (h \circ f^{-1})[V] = h[U] = C$. \square

Item (4) of Lemma 5.4 states that a subset S of a copy $C = f[U]$ for $f \in \overline{G}[U]$ is a copy in respect to the action of G on U if and only if it is also a copy in respect to the action of $f[G]$ on C . That is being a copy is "invariant" of that interpretation.

Lemma 5.5. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Let $f \in \overline{G}$ and $F \cup \{x, y\}$ be a finite subset of $f[U]$. Then:*

- (1) *The types $\langle F|x \rangle$ and $\langle F|y \rangle$ are equal with respect to the group action of G on U if and only if they are equal with respect to the group action of the group $f[G]$ on $f[U]$.*
- (2) *If $F \cup \{x\} \subseteq f[U] := C$ for $f \in \overline{G}$, then:*
 - (a) $G\langle F|x \rangle \cap C = (f[G])\langle F|x \rangle$.
 - (b) *If $G\langle F|x \rangle$ is finite then $G\langle F|x \rangle = (f[G])\langle F|x \rangle \subseteq C$.*

Proof.

(1): Let $F \cup \{x, y\} \subseteq f[U] := C$ for $f \in \overline{G}$. Assume $\langle F|x \rangle = \langle F|y \rangle$. Then there exists a function $g \in G\langle F \rangle$ with $g(x) = y$. Let the function $k \in G$ be such that $k \uparrow (f^{-1}[F \cup \{x, y\}]) = (f^{-1} \circ g \circ f) \uparrow (f^{-1}[F \cup \{x, y\}])$. Then $k \in G\langle f^{-1}[F] \rangle$ and $k(f^{-1}(x)) = f^{-1}(y)$ and $(f \circ k \circ f^{-1}) \uparrow (F \cup \{x, y\}) = g \uparrow (F \cup \{x, y\})$. Hence, the function $f(k)$ is a witness for $\langle F|x \rangle = \langle F|y \rangle$ in respect to the group action of $f[G]$ on $f[U]$. On the other hand let $k \in G$ for which the function $f(k)$ is a witness for $\langle F|x \rangle = \langle F|y \rangle$ in respect to the group action of $f[G]$ on $f[U]$. Then $f(k)(a) = a$ for all $a \in F$ and $f(k)(x) = y$. Let $g \in G$ be such that $g \uparrow (F \cup \{x, y\}) = f(k) \uparrow (F \cup \{x, y\})$. Then the function g is a witness for $\langle F|x \rangle = \langle F|y \rangle$ in respect to the group action of G on U .

Item (2)(a) follows from Item (1). Item (2)(b) follows from Item (2)(a) and Corollary 4.1. \square

6. ALGEBRAIC CLOSURE

Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. For F a finite subset of U we defined in Section 4 the set $\text{Orb}(G\langle F \rangle)$ to be the set of typesets with sockel F , that is the set of orbits of the group $G\langle F \rangle$. Let $F \subseteq U$, not necessarily finite. The *algebraic closure* of F is the set, denoted $\mathbf{ac}(F)$, and defined by setting:

$$\mathbf{ac}(F) = \begin{cases} \bigcup \{T \in \text{Orb}(G\langle F \rangle) \mid T \text{ is finite}\}, & \text{if } \bullet : F \text{ is finite;} \\ \bigcup \{\mathbf{ac}(F') \mid F' \in \mathcal{P}_{<\omega}(F)\}, & \bullet\bullet : \text{in general.} \end{cases}$$

Definition $\bullet\bullet$ is consistent with Definition \bullet because if a typeset T of a type is finite then every continuation of this type has a typeset which is a subset of T and hence finite. (Fact 4.2.) Finally, set $K_{\text{ac}} = \text{ac}(\emptyset)$. Note that $x \in K_{\text{ac}}$ if and only if the set $G\langle\emptyset|x\rangle$, that is the orbit of G containing x is finite.

Lemma 6.1. *If G is oligomorphic then $\text{ac}(F)$ is finite for every finite subset F of U .*

Proof. By Definition 4.1, if G is oligomorphic the set $\text{Orb}(G\langle F\rangle)$ of typesets with sockel F is finite. Hence $\text{ac}(F)$ is then a finite union of finite sets. \square

For R a relational structure let $\text{Age}(R)$, the *age* of R , denote the class of structures which are isomorphic to a finite induced substructure of R . A group G is *algebraically finite* if $\text{ac}(F)$ is finite for every finite subset F of U . There are many algebraically finite groups which are not oligomorphic. In fact there are many non oligomorphic groups such that $\text{ac}(F) = F$ for every finite F . If, in this case, a structure R is a countable homogeneous relational structure such that $\text{Aut}(R) = \overline{G} \cap \mathfrak{S}$ then this amounts to the fact that $\text{Age}(R)$ satisfies the *disjoint amalgamation property* (DAP). See [5].

Lemma 6.2. *Let $x \in U$. Then $x \in K_{\text{ac}}$ if and only if there exists a finite subset F of U such that $x \in g[F]$ for every function $g \in G$.*

Proof. Suppose that $x \in K_{\text{ac}}$. Set $F := G\langle\emptyset|x\rangle$, the orbit of x w.r.t. the action of G . By definition, F is finite and $g[F] = F$ for every $g \in G$.

Conversely, let $F \in \mathfrak{P}_{<\omega}(U)$ of minimal cardinality for which $x \in g[F]$ for every function $g \in G$. This means that $x \in F' := \bigcap \{g[F] : g \in G\}$. We have $h[F'] = \bigcap \{h \circ g[F] : g \in G\} \supseteq F'$ for every $h \in G$ and since F' is finite, $h[F'] = F'$. Since $x \in F'$, the orbit of x w.r.t. the action of G is included into F' . By minimality, this orbit, F' and F are equal. Hence, $x \in K_{\text{ac}}$. \square

Let R be a relational structure with domain U . The *kernel* of R is the set $\text{Ker}(R) := \{x \in U : \text{Age}(R_{-x}) \neq \text{Age}(R)\}$. ($R_{-x} := R \upharpoonright (U \setminus \{x\})$.)

Corollary 6.1. *Let R be a homogeneous relational structure and $G = \text{Aut}(R)$. Then $K_{\text{ac}} = \text{Ker}(R)$.*

Proof. Let $x \in \text{Ker}(R)$. By definition there exists a finite subset F of U containing x such that $R \upharpoonright F$ does not embed into R_{-x} . It follows that $x \in g[F]$ for every $g \in G$. According to Lemma 6.2, the element $x \in K_{\text{ac}}$. Conversely, suppose that $x \in K_{\text{ac}}$. Then $F := G\langle\emptyset|x\rangle$ is finite. Since F is preserved by every member of G , there is no embedding of $R \upharpoonright F$ into R_{-x} . \square

Lemma 6.3. *The map $F \rightarrow \mathbf{ac}(F)$ is an algebraic closure operator.*

Proof. We have trivially $F \subseteq \mathbf{ac}(F) \subseteq \mathbf{ac}(F')$ for every $F \subseteq F'$. In particular $\mathbf{ac}(F) \subseteq \mathbf{ac}(\mathbf{ac}(F))$. We have $\mathbf{ac}(F) = \mathbf{ac}(\mathbf{ac}(F))$. Indeed, let $x \in \mathbf{ac}(\mathbf{ac}(F))$. Then by definition there is some finite $Q \subseteq \mathbf{ac}(F)$ such that $x \in \mathbf{ac}(Q)$. Select for each $q \in Q$ some finite $Q_q \subseteq F$ such that $q \in \mathbf{ac}(Q_q)$. Set $S := \bigcup_{q \in Q} Q_q$. Then $x \in \mathbf{ac}(S) \subseteq \mathbf{ac}(F)$. Thus, \mathbf{ac} defines a closure operator. According to the second clause in the definition, it is algebraic. \square

This operator does not necessarily have the exchange property. Here is an example where the group is transitive. Let $U := [\mathbb{N}]^2$ and G be the group of permutations of U induced by permutations of \mathbb{N} . Let $u := \{x, y\} \in U$, $u' := \{x', y'\} \in U$ be two disjoint pairs of \mathbb{N} and $v := \{x, x'\}$. Then $v \notin \mathbf{ac}(\{u\}) = \{u\}$, and $v \in \mathbf{ac}(\{u, u'\})$ while $u' \notin \mathbf{ac}(\{u, v\})$.

7. RANKED CLOSURE

Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. A type $\langle F|p \rangle$ has *rank* 0 if $G\langle F|p \rangle$ is finite. Let \mathfrak{R}_0 be the set of types $\langle F|p \rangle$ having rank 0. We define recursively sets \mathfrak{R}_α of types for α an ordinal. If for every ordinal $\beta < \alpha$ the set \mathfrak{R}_β has been determined then the type $\langle F|p \rangle$ is an element of \mathfrak{R}_α if there exists a finite subset $F' \supseteq F$ of U so that every continuation $\langle F'|q \rangle$ of $\langle F|p \rangle$ is a type in \mathfrak{R}_β for some ordinal $\beta < \alpha$. Note that $\mathfrak{R}_\beta \subseteq \mathfrak{R}_\alpha$ for all $\beta < \alpha$; letting $F' = F$. A type $\langle F|p \rangle$ has *rank* α if $\langle F|p \rangle \in \mathfrak{R}_\alpha$ and $\langle F|p \rangle \notin \mathfrak{R}_\beta$ for any $\beta < \alpha$. Hence if a type $\langle E|x \rangle \in \mathfrak{R}_\gamma$, then the rank of the type $\langle E|x \rangle$ is less than or equal to γ . A type $\langle F|p \rangle$ is *unranked* if there is no ordinal α with $\langle F|p \rangle \in \mathfrak{R}_\alpha$.

Note that a type $\langle F|p \rangle$ is unranked if and only if:

- (p): For every finite set F' with $F \subseteq F' \subseteq U$ exists a continuation $\langle F'|q \rangle$ of $\langle F|p \rangle$ which is unranked.

Lemma 7.1. *If a type $\langle F|p \rangle$ is ranked and has rank α then every continuation $\langle E|q \rangle$ of the type $\langle F|p \rangle$ is ranked and has rank less than or equal to α .*

Proof. By induction on the rank of the type $\langle F|p \rangle$. If $G\langle F|p \rangle$ is finite then $G\langle E|q \rangle \subseteq G\langle F|p \rangle$ is finite. Let $\alpha > 0$. If the continuation $G\langle E|q \rangle$ is finite then $\langle E|q \rangle$ is ranked and has rank $0 < \alpha$. Otherwise we proceed as follows: There exists a finite subset $F' \supseteq F$ of U so that the rank $\beta(x)$ of every continuation $\langle F'|x \rangle$ of $\langle F|p \rangle$ is smaller than α . The set $E' = E \cup F'$ is finite. The set of typesets $G\langle E'|x \rangle$ for $x \in G\langle E|q \rangle$ forms a partition of the typeset $G\langle E|q \rangle$. Each one of the types $\langle E'|x \rangle$ is a continuation of the type $\langle F'|x \rangle$ having rank $\beta(x) < \alpha$. Implying

that the type $\langle E|q \rangle \in \mathfrak{R}_\alpha$ and hence that the rank of the type $\langle E|q \rangle$ is less than or equal to α . \square

Corollary 7.1. *Let $\langle F|p \rangle$ be a ranked type and $F \subseteq E \in \mathcal{P}_{<\omega}(U)$. Then the set $G\langle F|p \rangle$ is the union over the set of typesets of the ranked types $\langle E|x \rangle$ with $x \in G\langle F|p \rangle$.*

Lemma 7.2. *Let $\langle F|p \rangle$ be a type and $g \in G$. Then the type $\langle F|p \rangle$ is ranked if and only if the type $\langle g[F]|g(p) \rangle$ is ranked. The type $\langle F|p \rangle$ has rank α if and only if the type $\langle g[F]|g(p) \rangle$ has rank α .*

Proof. By induction on the rank of the type $\langle F|p \rangle$. It follows from Corollary 4.1 that the type $\langle F|p \rangle$ has rank 0 if and only if the type $\langle g[F]|g(p) \rangle$ has rank 0. Let the type $\langle F|p \rangle$ have rank $\alpha > 0$. Then there exists a finite $F' \supseteq F$ for which every continuation $\langle F'|q \rangle$ of the type $\langle F|p \rangle$ has a rank, say β_q , which is smaller than α . Also $g[F'] \supseteq g[F]$. Let $\langle g[F']|s \rangle$ be a continuation of the type $\langle g[F]|g(p) \rangle$. Then $s \in G\langle g[F]|g(p) \rangle$. It follows, using Lemma 4.2, that $g^{-1}(s) \in G\langle F|p \rangle$ and hence that the type $\langle F'|g^{-1}(s) \rangle$ is a continuation of the type $\langle F|p \rangle$. It has a rank $\beta_{g^{-1}(s)} < \alpha$. Then, via the induction, the rank of the type $\langle g[F']|s \rangle$ is also equal to $\beta_{g^{-1}(s)} < \alpha$. Hence the rank of the type $\langle g[F]|g(p) \rangle$ is equal to $\bar{\alpha}$ less than or equal to α . Applying g^{-1} to the type $\langle g[F]|g(p) \rangle$ we obtain $\alpha \leq \bar{\alpha}$. Hence $\langle F|p \rangle$ is ranked if and only if $\langle g[F]|g(p) \rangle$ is ranked and both types have the same rank. \square

Lemma 7.3. *Let $\langle F|p \rangle$ be a ranked type. Then $G\langle F|p \rangle \subseteq C$ for every $C \in \overline{G}[U]$ with $F \subseteq C$.*

Proof. Let $F \subseteq C \in \overline{G}[U]$. According to Lemma 3.2 there is a function $h \in \overline{G}\langle F \rangle$ with $h[U] = C$. We proceed via induction on the rank of the type $\langle F|p \rangle$. In the case that the typeset $G\langle F|p \rangle$ is finite the Lemma follows from Corollary 4.1.

Let $0 < \alpha$ be an ordinal for which $G\langle F''|q \rangle \subseteq C$ for all types $\langle F''|q \rangle$ of rank $\beta < \alpha$ and with $F'' \subseteq C$. Let $\langle F|p \rangle$ be a type of rank α . There exists then a finite subset $F' \subseteq U$ with $F' \supseteq F$ and so that every continuation $\langle F'|q \rangle$ of $\langle F|p \rangle$ has a rank, say $\beta(q)$, which is smaller than α . Let $Q := \{q \in G\langle F|p \rangle\}$ be the set of those elements q . It follows from Fact 4.2 that the sets $G\langle F'|q \rangle$ for $q \in Q$ form a partition of the set $G\langle F|p \rangle$. Let $g \in G$ be such that $h \upharpoonright F' = g \upharpoonright F'$. Then $g[F'] \subseteq C$ and $g \in G\langle F \rangle$. Consequently the sets $G\langle g[F']|g(q) \rangle$ for $q \in Q$ form a partition of the set $G\langle F|p \rangle$. Each of the types $\langle g[F']|g(q) \rangle$ is ranked and its rank is equal to $\beta(q)$ according to Lemma 7.2. Then $G\langle h[F']|h(q) \rangle \subseteq C$ using the induction assumption. Implying that $G\langle F|p \rangle \subseteq C$. \square

Let $\{f, h\} \subseteq \overline{G}$ and $C = f[U] = h[U]$. Then the group $f[G] \subseteq \mathfrak{S}(f[U])$ acts on the set $f[U]$. See Definition 5.1 and Lemma 5.4. It follows from Lemma 5.5 that for every finite $F \subseteq C$ and $x \in C \setminus F$ the type $\langle F|x \rangle$ with respect to the group action $f[G]$ is the same as with respect to the group action $h[G]$. Hence we can define:

Definition 7.1. *Let $\langle F|p \rangle$ be a type and $f \in \overline{G}$ with $F \cup \{p\} \subseteq f[U] := C$. Then $\langle F|p \rangle_C$ denotes this type with socket F in respect to the action of the group $f[G]$.*

Note that, for every $f \in \overline{G}$, the typeset of the type $\langle F|p \rangle_C$ is the set

$$G\langle F|p \rangle \cap C = (f[G])\langle F|p \rangle_C = \{(f(g))(x) \mid g \in G\}.$$

Lemma 7.4. *Let $f \in \overline{G}$ with $f[U] := C$. Then a type $\langle F|p \rangle$, with $f \cup \{p\} \subseteq C$, is ranked if and only if the type $\langle F|p \rangle_C$ is ranked with respect to the group action $f[G]$. The type $\langle F|p \rangle$ has rank α if and only if the type $\langle F|p \rangle_C$ has rank α with respect to the group action $f[G]$.*

Proof. Let $\text{rank}\langle F|p \rangle$ denote the rank of a type with respect to the group action G and $\text{rank}_f\langle F|p \rangle_C$ denote the rank of a type in C with respect to the group action $f(G)$. We proceed via induction on the ranks. In the case that the typeset $G\langle F|p \rangle$ is finite the Lemma follows from Lemma 5.5 Item (2)(b).

Let α be the smallest ordinal for which there exists a type $\langle F|p \rangle_C$ with $\nu := \text{rank}\langle F|p \rangle \neq \alpha := \text{rank}_f\langle F|p \rangle_C$. There exists a finite subset $F' \subseteq C$ with $F' \supseteq F$ and so that every continuation $\langle F'|q \rangle_C$ of $\langle F|p \rangle_C$ has a rank, say $\beta(q) = \text{rank}_f\langle F'|q \rangle_C$, which is smaller than α . The types $\langle F'|q \rangle$ are then continuations of the type $\langle F|p \rangle$ and via induction have the ranks β_q . Hence $\nu \leq \alpha$. There exists a finite subset $F' \subseteq U$ with $F' \supseteq F$ and so that every continuation $\langle F'|q \rangle$ of $\langle F|p \rangle$ has a rank, say $\beta(q) = \text{rank}\langle F'|q \rangle$, which is smaller than ν . Let $Q := q \in G\langle F|p \rangle \setminus F'$ be the set of those elements q . It follows from Fact 4.2 that the sets $G\langle F'|q \rangle$ for $q \in Q$ form a partition of the set $G\langle F|p \rangle$. According to Lemma 3.2 there is a function $h \in \overline{G}\langle F \rangle$ with $h[U] = C$. Let $g \in G$ be such that $h \upharpoonright F' = g \upharpoonright F'$. Then $g[F'] \subseteq C$ and $g \in G\langle F \rangle$. Consequently the sets $G\langle g[F']|g(q) \rangle$ for $q \in Q$ form a partition of the set $G\langle F|p \rangle$. Each of the types $\langle g[F']|g(q) \rangle$ is ranked and its rank is equal to $\beta(q)$ according to Lemma 7.2. Implying according to Lemma 7.3 that $G\langle h[F']|h(q) \rangle \subseteq C$. Via induction then $\text{rank}\langle g[F']|g(q) \rangle_C = \beta(q) < \nu$. Hence $\alpha \leq \nu$. \square

Lemma 7.5. *If $G\langle F|x \rangle$ is an unranked type, then there exists an element $C \in \overline{G}[U]$ with $F \subseteq C$ and with $x \notin C$.*

Proof. Let $(u_i \mid i \in \omega)$ be an enumeration of $U \setminus F$. Let $U_n = \{u_i \mid i \in n\}$. Note that $U_0 = \emptyset$. We will construct a sequence $\{x_n \in U \mid n \in \omega\}$ so that for all $n \in \omega$:

- (1) $x_0 = x$.
- (2) $x_n \in U \setminus (F \cup U_n)$ and the type $\langle F \cup U_n \mid x_n \rangle$ is unranked.
- (3) The type $\langle F \cup U_{n+1} \mid x_{n+1} \rangle$ is a continuation of the type $\langle F \cup U_n \mid x_n \rangle$.

The type $\langle F \cup U_n \mid x_n \rangle$ is unranked. Hence there exists, according to property **(p)**, an unranked type $\langle F \cup U_{n+1} \mid x_{n+1} \rangle$ which is a continuation of the type $\langle F \cup U_n \mid x_n \rangle$. Let $f_n \in \mathbf{G}\langle F \cup U_n \rangle$ be the function with $f_n(x_{n+1}) = x_n$.

Let g_0 be the identity function on U , that is $\{g_0\} = \mathbf{G}\langle U \rangle$. We will construct recursively functions $g_n \in \mathbf{G}\langle F \rangle$ for $n \in \omega$ for which $g_n(x_n) = x$ and for which $l_n := g_n \upharpoonright (F \cup U_n) \subseteq g_{n+1} \upharpoonright (F \cup U_{n+1}) = l_{n+1}$. Assume that g_n has been constructed. Let $g_{n+1} = g_n \circ f_n$. Note that $x \notin l_n[U_n]$ for all $n \in \omega$.

Then $l_0 \subset l_1 \subset l_2 \subset \dots$ and $l := \bigcup_{n \in \omega} l_n \in \overline{\mathbf{G}}$ with $x \notin l[U]$. Let $C = l[U]$.

□

Definition 7.2. For $F \subseteq U$ let the intersection closure:

$$\mathbf{ic}_{\mathbf{G}}(F) := \bigcap \{C \in \overline{\mathbf{G}}[U] \mid F \subseteq C\} = \bigcap \overline{\mathbf{G}}[U][F].$$

For F a finite subset of U let $\mathbf{rc}_{\mathbf{G}}(F)$, the ranked closure of F , be the union of F together with the union of the typesets $\mathbf{G}\langle F \mid p \rangle$ for which $\langle F \mid p \rangle$ is a ranked type. (In particular then $\mathbf{rc}_{\mathbf{G}}(\emptyset)$ is the union of the ranked orbits of \mathbf{G} .)

In general, for a subset S of U let:

$$(1) \quad \mathbf{rc}_{\mathbf{G}}(S) := \bigcup \{\mathbf{rc}(F) \mid F \in \wp_{<\omega}(S)\}.$$

If the group \mathbf{G} is understood we write \mathbf{ic} for $\mathbf{ic}_{\mathbf{G}}$ and \mathbf{rc} for $\mathbf{rc}_{\mathbf{G}}$

It follows from Corollary 7.1 that the ranked closure definition (1) is consistent with the ranked closure definition for finite subsets F of U .

Note 7.1. Let the group \mathbf{G} be oligomorphic, then: Every ranked type has rank 0. A type is unranked if and only if its typeset is infinite. The ranked closure of every finite set is finite. For every set $S \subseteq U$ is $\mathbf{ac}(S) = \mathbf{rc}(S)$.

Theorem 7.1. Let U be a countable set and \mathbf{G} be a subgroup of $\mathfrak{S}(U)$. Let $S \subseteq U$. Then $\mathbf{rc}_{\mathbf{G}}(S) \subseteq \mathbf{ic}_{\mathbf{G}}(S)$. If F is a finite subset of U then $\mathbf{rc}_{\mathbf{G}}(F) = \mathbf{ic}_{\mathbf{G}}(F)$.

Proof. The inclusion $\mathbf{rc}_{\mathbf{G}}(S) \subseteq \mathbf{ic}_{\mathbf{G}}(S)$ follows from Lemma 7.3. The inclusion $\mathbf{ic}_{\mathbf{G}}(F) \supseteq \mathbf{rc}_{\mathbf{G}}(F)$ for finite F follows from Lemma 7.5. □

According to the definition of typeset, a subset T of U with $t \in T$ is the point orbit of G containing t if and only if $T = G\langle \emptyset | t \rangle$. Since by Theorem 7.1, $\mathbf{rc}_G(\emptyset) = \mathbf{ic}_G(\emptyset)$, the intersection $\mathbf{ic}(\emptyset) = \bigcap \overline{G}(U)$ of all copies for the group G is equal to the union of the point orbits of G . In particular, we have a characterization in terms of ranked closure of the nonexistence of a proper copy (compare with Lemma 3.4).

Corollary 7.2. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then there is no proper copy (i.e., $\overline{G}[U] = \{U\}$, or equivalently, if every function $f \in \overline{G}$ is bijective) if and only if every point orbit of the group G is ranked.*

Corollary 7.3. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$ and F a finite subset of U for which the set $\mathbf{rc}(F)$ is finite. Then $\overline{G}[U][\mathbf{rc}(F)] = \overline{G}[U][F]$.*

Proof. Clearly $\overline{G}[U][\mathbf{rc}(F)] \subseteq \overline{G}[U][F]$. It follows from Theorem 7.1 that $\overline{G}[U][\mathbf{rc}(F)] \supseteq \overline{G}[U][F]$. □

Lemma 7.6. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. Then the map $\mathbf{rc} : \wp(U) \rightarrow \wp(U)$, which maps every subset S of U to the subset $\mathbf{rc}(S)$ of U , is an algebraic closure operator.*

Proof. If $F \subseteq F'$ are two finite subsets of U then $F \subseteq \mathbf{rc}(F) \subseteq \mathbf{rc}(F')$ follows trivially from Definition 7.2 and then from Corollary 7.1. Hence $S \subseteq \mathbf{rc}(S) \subseteq \mathbf{rc}(S')$ for every $S \subseteq S'$ using Definition 7.2 again. In particular $\mathbf{rc}(S) \subseteq \mathbf{rc}(\mathbf{rc}(S))$. We have to verify $\mathbf{rc}(S) = \mathbf{rc}(\mathbf{rc}(S))$. Let $x \in \mathbf{rc}(\mathbf{rc}(S))$. Then by definition there exists a finite set $F \subseteq \mathbf{rc}(S)$ such that $x \in \mathbf{rc}(F)$. Select for each $y \in F$ a finite set $F_y \subseteq S$ such that $y \in \mathbf{rc}(F_y)$. Then $y \in \mathbf{ic}(F_y)$ for every $y \in F$ according to Theorem 7.1. Set $E := \bigcup_{y \in F} F_y$. Then $y \in \mathbf{ic}(F_y) \subseteq \mathbf{ic}(E)$ for every $y \in F$, hence $F \subseteq \mathbf{ic}(E)$. Implying, using Theorem 7.1, that $F \cup \{x\} \subseteq \mathbf{ic}(E)$ and $x \in \mathbf{rc}(E) \subseteq \mathbf{rc}(S)$. Thus, the map \mathbf{rc} defines a closure operator. According to stipulation (1) of Definition 7.2, the map \mathbf{rc} is algebraic. □

However, there are examples for which the ranked closure of the empty set is empty and still there are no pairwise disjoint copies.

Example 7.2. *Let $T_{\mathbb{Z}}$ be the ordered tree on a countable set U in which each interval is finite, every element has countably many successors and a unique predecessor. Let $G := \text{Aut}(T_{\mathbb{Z}})$. Then:*

The powerset $\wp(\omega)$ is embeddable in $\overline{G}[U]$. Any two copies pairwise intersect. For $x \in U$ is the set $\downarrow x := \{y \in U \mid y \leq x\}$ the intersection of all copies containing x .

7.1. The case of linear orders. Let $Q = (Q; \leq)$ be the linear order of the rationals. For $S \subseteq \omega$ associate the copy S_Q of Q given by:

$$S_Q := ((-1, 0) \cap Q) \cup \bigcup_{s \in S} ((s, s+1) \cap Q).$$

Because the binary relational structure Q is homogeneous, see [5], it follows that S_Q is a copy for the group $\text{Aut}(Q)$ and hence that the partial order $(\overline{\text{Aut}(Q)}[Q]; \subset)$ of copies for the group $\overline{\text{Aut}(Q)}$ embeds the partial order $(\wp(\omega); \subset)$.

Let $U = (U; \leq)$ be a linear order on a countable set U and let G be the group of automorphisms of U . Clearly, every finite orbit of G is a singleton set, that is a linear order of order type $\mathbf{1}$. Let ζ be the order type of the integers and note that ζ^0 is the order type $\mathbf{1}$. If U is the natural linear order of the set of integers then all of the orbits of the group $G\langle\{0\}\rangle$ are singleton sets. That is $\text{Orb}(G\langle\{0\}\rangle) = \{\{i\} \mid i \in U\}$. Implying that if an orbit of the group of automorphisms G of a linear order U has order type ζ then the rank of this orbit is 1. If an orbit of the group of automorphisms G of a linear order U has order type ζ^2 then the rank of this orbit is 2. Consulting the discussion on powers of linear orders of order type ζ and in particular Corollary 8.6 of the Rosenstein book [17], we obtain using Lemma 3.2 and Theorem 7.1:

Theorem 7.2. *Let $U = (U; \leq)$ be a linear order on a countable set U and let G be the group of automorphisms of U , then:*

- (1) *Every orbit of G has order type ζ^γ or $\zeta^{\gamma \cdot \eta}$ for some ordinal number γ . (The ordinal γ is the Hausdorff rank of the scattered linear order ζ^γ .)*
- (2) *There exists only one copy for G if and only if every orbit of G has order type ζ^γ for some ordinal number γ . The rank of such an orbit is then the ordinal γ .*
- (3) *If the group G has an orbit of order type $\zeta^{\gamma \cdot \eta}$ then the partial order $(\overline{G}[U]; \subseteq)$ of copies for the group G embeds the partial order $(\wp(\omega); \subseteq)$.*

8. AN APPLICATION TO ALGEBRAICALLY FINITE GROUPS

Lemma 8.1. *Let U be a countable set and G be a subgroup of $\mathfrak{S}(U)$. If $\mathfrak{rc}(F) = U$ for some finite subset F of U then $\mathfrak{rc}(E) = U$ for all finite subsets E of U .*

Proof. Assume that there exists a finite subset F of U with $\mathfrak{rc}(F) = U$. Then according to Theorem 7.1 and Definition 7.2 and Lemma 3.2: $U = \mathfrak{ic}(F) = \bigcap \{C \in \overline{G}[U] \mid F \subseteq C\} = \bigcap \overline{G}\langle F \rangle[U]$. Hence $\overline{G}\langle F \rangle = \{U\}$. Implying according to Lemma 3.4 that $\overline{G}\langle E \rangle = \{U\}$ for all finite subsets

E of U . Hence $\mathbf{rc}(E) = \mathbf{ic}(E) = \bigcap \overline{\mathbf{G}\langle E \rangle}[U] = \bigcap \{U\} = U$ for all finite subsets E of U . \square

Corollary 8.1. *Let U be a countable set and \mathbf{G} be a subgroup of $\mathfrak{S}(U)$ with $\mathbf{rc}(\emptyset) \neq U$. Let F be a finite subset of U . Then $C := \mathbf{rc}(F) \notin \overline{\mathbf{G}}[U]$.*

Proof. If $C \in \overline{\mathbf{G}}[U]$ let $f \in \overline{\mathbf{G}}$ with $f[U] = C$. According to Lemma 7.4 a type $\langle F|p \rangle$ is ranked if and only if the type $\langle F|p \rangle_C$ is ranked with respect to the group action $f[\mathbf{G}]$ on C . It follows then from Lemma 8.1 that $\mathbf{rc}(\emptyset) = C$ under the group action $f[\mathbf{G}]$ on C . Hence, every orbit of the group $f[\mathbf{G}]$ is ranked. Which according to Lemma 7.4 implies that every orbit of \mathbf{G} is ranked and hence that $\mathbf{rc}(\emptyset) = U$. \square

Theorem 8.1. *Let U be a countable set and \mathbf{G} be a subgroup of $\mathfrak{S}(U)$ and $\mathbf{rc}(\emptyset) \neq U$. Then for every finite subset F of U and every copy $C \in \overline{\mathbf{G}}[U]$ with $F \subseteq C$ (hence $\mathbf{rc}(F) \subseteq C$), there exists:*

An ω -sequence $C = C_0 \supset C_1 \supset C_2 \supset C_3 \supset \dots$ of copies $C_i \in \overline{\mathbf{G}}[U]$ for which $\mathbf{rc}(F) = \bigcap_{i \in \omega} C_i$.

Proof. Let F be a finite subset of U and let $C_0 := C$. It follows from Lemma 8.1 that $\mathbf{rc}(F) \neq C$. Implying that there exists an unranked type with sockel F . Let $P := \{p_j \mid j \in \omega\}$ be an ω -enumeration of the union P of all the typesets $\mathbf{G}\langle F|p \rangle \cap C$ with sockel F for unranked types $\langle F|p \rangle$. If for a type $\langle F|p \rangle$ the typeset $\mathbf{G}\langle F|p \rangle \cap C$ is finite then the rank of the type $\langle F|p \rangle_C$ with respect to the copy C is 0. It follows from Lemma 7.4 that then the type $\langle F|p \rangle$ is ranked. Hence the set P is infinite.

Assume now that we have already constructed a sequence $C_0 \supset C_1 \supset C_2 \supset \dots \supset C_{n-1} \supseteq \mathbf{rc}(F)$ of copies. Then $C_{n-1} \neq \mathbf{rc}(F)$ according to Corollary 8.1. Let $f \in \overline{\mathbf{G}}$ with $f[U] = C_{n-1}$. Let m be the smallest index for which $p_m \in C_{n-1}$. The type $\langle F|p_m \rangle$ is unranked for the group action \mathbf{G} . Implying, according to Lemma 7.4 that the type $\langle F|p_m \rangle_{C_{n-1}}$ is unranked with respect to the group action $f[\mathbf{G}]$. Implying that there exists a copy $C_n \subset C_{n-1}$ with $F \subseteq C_n \in \overline{(f[\mathbf{G}])}[C_{n-1}]$. It follows from Lemma 5.4 Item (5) that $C_n \in \overline{\mathbf{G}}[U]$. Because $F \subseteq C_n$ it follows from Lemma 7.3 that $\mathbf{rc}(F) \subseteq C_n$. \square

Corollary 8.2. *Let U be a countable set and \mathbf{G} be a subgroup of $\mathfrak{S}(U)$. Then for all finite subsets F and E of U with $E \cap \mathbf{rc}(F) = \emptyset$ and for every copy C with $F \subseteq C$:*

- (1) *There exists a copy C' with $C' \cap E = \emptyset$ and with $F \subseteq C' \subseteq C$.*
- (2) *For every function $f \in \overline{\mathbf{G}}\langle F \rangle$:*

- (a) $\mathbf{rc}(F) = \mathbf{ic}(F) = \bigcap (\overline{\mathbf{G}}\langle F \rangle)[U]$.
 (b) $f[\mathbf{rc}(F)] = \mathbf{rc}(F)$.

Proof. According to Theorem 7.1, we have $\mathbf{ac}(F) = \mathbf{rc}(F)$ for every finite subset $F \subseteq U$. Hence, in our case, we have $\mathbf{ac}(F) \cap E = \emptyset$. Applying Corollary 3.2, we get Item (1).

Item (2a): $\mathbf{rc}(F) = \mathbf{ic}(F)$ according to Theorem 7.1. Since $\mathbf{ic}(F) = \bigcap \overline{\mathbf{G}}[U][F]$ according to Definition 7.2 and $\overline{\mathbf{G}}[U][F] = \overline{\mathbf{G}}[U]\langle F \rangle$ according to Lemma 3.2 we obtain $\mathbf{ic}(F) = \bigcap \overline{\mathbf{G}}[U]\langle F \rangle$.

(2b): It follows from Definition 7.2 and from Lemma 4.2 that $f[\mathbf{rc}(F)] \subseteq \mathbf{rc}(f[F]) = \mathbf{rc}(F)$. The set $f[U]$ is a copy containing $F = f[F] \subseteq f[\mathbf{rc}(F)]$. In particular it follows from Lemma 4.2 that f maps every typeset of the types with sockel F to a subset of this typeset. Implying, because U is the union of F together with the union of the typesets of the types with sockel F , that if f would map the typeset of a ranked type $\langle F|p \rangle$ to a proper subset then $\mathbf{rc}(F)$ would not be a subset of $f[U]$. But, it follows from Theorem 7.1 that $\mathbf{rc}(F) \subseteq f[U]$. \square

Lemma 8.2. *If \mathbf{G} is a algebraically finite subgroup of $\mathfrak{S}(U)$ then $\mathbf{rc}(S) = \mathbf{ac}(S)$ for every subset S of U .*

Proof. It suffices to prove the lemma for finite subsets F of U . If $\langle F|p \rangle$ is a type of rank 1 then there exists a finite $F' \supseteq F$ so that every continuation $\langle F'|q \rangle$ of $\langle F|p \rangle$ has rank 0. Leading to the contradiction that $\mathbf{G}\langle F|p \rangle \subseteq \bigcup_q \mathbf{G}\langle F'|q \rangle$ is finite and hence has rank 0. Hence, the ranked closure of a finite set is the union of typesets of types which have rank 0, that is the union of typesets which are finite and therefore equal to the algebraic closure. \square

Lemma 8.3. *Let U be a countably infinite set and \mathbf{G} be a algebraically finite subgroup of $\mathfrak{S}(U)$ with $\mathbf{ac}(\emptyset) = \emptyset$. Then there exist two copies $\{C, D\} \subseteq \overline{\mathbf{G}}[U]$ with $C \cap D = \emptyset$.*

Proof. Let $\{u_i \mid i \in \omega\}$ be an ω -enumeration of the set U . For $n \in \omega$ let $U_n = \{u_i \mid i \in n\}$. Let $C_0 = D_0 = U$ and $V_0 = W_0 = \emptyset$. Let g_1 be the identity map on U . For $V_1 := \{u_0\}$ the algebraic closure of V_1 is finite. Hence there exists, according to Corollary 8.2 Item (1) with E for $\mathbf{ac}(V_1)$ and F for \emptyset and C for U , a copy D_1 with $\mathbf{ac}(V_1) \cap D_1 = \emptyset$. The typeset $\mathbf{G}\langle \emptyset|u_0 \rangle$ of the type $\langle \emptyset|u_0 \rangle$ is infinite because $\mathbf{ac}(\emptyset) = \emptyset$. Implying according to Lemma 7.4 that the typeset $\mathbf{G}\langle \emptyset|u_0 \rangle \cap D_1$ of the type $\langle \emptyset|u_0 \rangle_{D_1}$ is infinite. Let $w_0 \in (\mathbf{G}\langle \emptyset|u_0 \rangle \cap D_1) \setminus \mathbf{ac}(V_1)$ and let $W_1 = \{w_0\}$. Note that there exists a function $h_1 \in \mathbf{G}$ with $h_1(u_0) = w_0$. It follows from Theorem 7.1 that $\mathbf{ac}(W_1) \subseteq D_1$. Hence $\mathbf{ac}(V_1) \cap (\mathbf{ac}(W_1)) = \emptyset$. It follows using Corollary 8.2 Item (1) that there

exists a copy C_1 with $\mathbf{ac}(V_1) \subseteq C_1 \subseteq U$ and with $\mathbf{ac}(W_1) \cap C_1 = \emptyset$. Then $\mathbf{ac}(V_1) \subseteq C_1 \setminus D_1$ and $\mathbf{ac}(W_1) \subseteq D_1 \setminus C_1$.

For $n \in \omega$ assume that there are copies C_n and D_n and sets $V_n = \{v_i \mid i \in n\}$ with $\mathbf{ac}(V_n) \subseteq C_n \setminus D_n$ and $W_n = \{w_i \mid i \in n\}$ with $\mathbf{ac}(W_n) \subseteq D_n \setminus C_n$. Assume further that there are functions g_n and h_n in \mathbb{G} with $g_n(u_i) = v_i$ for all $i \in n$ and $h_n(u_i) = w_i$ for all $i \in n$. Note that $\mathbf{ac}(V_n) \cap \mathbf{ac}(W_n) = \emptyset$ and that the set $\mathbf{ac}(V_n) \cup \mathbf{ac}(W_n)$ is finite. If the typeset of the type $\langle V_n \mid g(u_n) \rangle$ is finite then it is a subset of the algebraic closure of V_n and hence a subset of $C_n \setminus D_n$. In this case let $v_n = g_n(u_n)$ and $g_{n+1} = g_n$. If the typeset of the type $\langle V_n \mid g(u_n) \rangle$ is infinite then the typeset of the type $\langle V_n \mid g(u_n) \rangle_{C_n}$ is infinite. Implying that there exists an element $v_n \in (\mathbb{G}\langle V_n \mid g(u_n) \rangle \cap C_n) \setminus \mathbf{ac}(W_n)$. Let $V_{n+1} = V_n \cup \{v_n\}$. There exists a function $f \in \mathbb{G}\langle V_n \rangle$ with $f(g_n(u_n)) = v_n$. Let $g_{n+1} = f \circ g_n$. According to Corollary 8.2 Item (1) there exists a copy C_{n+1} with $\mathbf{ac}(V_{n+1}) \subseteq C_{n+1}$ and with $C_{n+1} \cap (\mathbf{ac}(W_n)) = \emptyset$.

If the typeset of the type $\langle W_n \mid h(u_n) \rangle$ is finite then it is a subset of the algebraic closure of W_n and hence a subset of $D_n \setminus C_{n+1}$. In this case let $w_n = h_n(u_n)$ and $h_{n+1} = h_n$. If the typeset of the type $\langle W_n \mid h(u_n) \rangle$ is infinite then the typeset of the type $\langle W_n \mid h(u_n) \rangle_{D_n}$ is infinite. Implying that there exists an element $w_n \in (\mathbb{G}\langle W_n \mid h(u_n) \rangle \cap D_n) \setminus \mathbf{ac}(C_{n+1})$. Let $W_{n+1} = W_n \cup \{w_n\}$. There exists a function $f' \in \mathbb{G}\langle W_n \rangle$ with $f'(h_n(u_n)) = w_n$. Let $h_{n+1} = f' \circ h_n$. According to Corollary 8.2 Item (1) there exists a copy D_{n+1} with $\mathbf{ac}(W_{n+1}) \subseteq D_{n+1}$ and with $D_{n+1} \cap \mathbf{ac}(V_{n+1}) = \emptyset$. Then $(\mathbf{ac}(V_{n+1})) \subseteq C_{n+1} \setminus D_{n+1}$ and $\mathbf{ac}(W_{n+1}) \subseteq D_{n+1} \setminus C_{n+1}$.

Recursively for every $n \in \omega$ there exists a set $V_n = \{v_i \mid i \in n\}$ and a set $W_n = \{w_i \mid i \in n\}$ for which $V_n \cap W_n = \emptyset$ and there exists a function $g_n \in \mathbb{G}$ and a function $h_n \in \mathbb{G}$ with $g_n(u_i) = v_i$ for all $i \in n$ and with $h_n(u_i) = w_i$ for all $i \in n$. Let $V = \bigcup_{n \in \omega} V_n$ and let $W = \bigcup_{n \in \omega} W_n$. Then $V \cap W = \emptyset$. Let $g : U \rightarrow V$ be the function with $g(u_i) = v_i$ for all $i \in \omega$. Then $g \in \overline{\mathbb{G}}[U]$ implying that V is a copy. Let $h : U \rightarrow W$ be the function with $h(u_i) = w_i$ for all $i \in \omega$. Then $h \in \overline{\mathbb{G}}[U]$ implying that W is a copy. \square

Theorem 8.2. *Let U be a countable set and \mathbb{G} be an algebraically finite subgroup of $\mathfrak{S}(U)$ and F a finite subset of U . Then there exist two copies $\{C, D\} \subseteq \overline{\mathbb{G}}[U]$ with $C \cap D = \mathbf{ac}(F)$.*

Proof. It follows from Lemma 3.2 that every copy containing the finite set $\mathbf{ac}(F)$ is a copy of the group $\mathbb{G}\langle \mathbf{ac}(F) \rangle$. Let $V = U \setminus (\mathbf{ac}(F))$. Every copy I of the group $\mathbb{G}\langle \mathbf{ac}(F) \rangle$ is the disjoint union of $\mathbf{ac}(F)$ and $I \cap V$. The set $\mathbb{H} = \{g \upharpoonright V \mid g \in \mathbb{G}\}$ of functions is a subgroup of the symmetric $\mathfrak{S}(V)$ and the function $\sigma : \mathbb{G} \rightarrow \mathbb{H}$ with $\sigma(g) = g \upharpoonright V$ is a

group action automorphism. A subset I of U is a copy for the group G if and only if $I \cap V$ is a copy for the group H . Because $\mathbf{ac}_H(\emptyset) = \emptyset$ the Theorem follows from Lemma 8.3. \square

9. THE LATTICE OF INTERSECTIONS OF COPIES

Let U be a set. A *Moore family* on U is a collection \mathcal{M} of subsets of U which is closed under intersections. The map $F \rightarrow \phi_{\mathcal{M}}(F) := \bigcap \{X \in \mathcal{M} : F \subseteq X\}$ is the *closure operator* associated with the Moore family. This closure operator is *algebraic* if it is closed under union of chains; equivalently, the Moore family is topologically closed in $\mathcal{P}(U)$ equipped with the product topology. If a closure operator is algebraic, the Moore family, once ordered by inclusion, forms an *algebraic lattice*, i.e. a complete lattice in which every element is a join(possibly infinite) of compact elements, an element x being *compact* if $x \leq \bigvee X$ in the lattice \mathcal{M} implies $x \leq \bigvee X'$ for some finite subset of X (but the converse is false in general)(see [6] for information about algebraic lattices). Let G be a subgroup of $\mathfrak{S}(U)$ and $\overline{G}[U]$ the set of copies. We denote by $\widehat{\overline{G}[U]}$ the set of intersections of members of $\overline{G}[U]$. Being closed under intersections this set is a *Moore family*; once ordered by inclusion, this family of intersections is a complete lattice. The map $F \rightarrow \mathbf{ic}(F)$ is the closure operator associated with the above Moore family. If the closure $F \rightarrow \mathbf{ic}(F)$ is algebraic then $\widehat{\overline{G}[U]}$ is an algebraic lattice. We do not know if the converse holds. In fact:

Problem 9.1. *Is $\widehat{\overline{G}[U]}$ an algebraic lattice? Is the closure $F \rightarrow \mathbf{ic}(F)$ algebraic?*

Let us recall that an element u of a poset P is *completely meet-irreducible* if the poset P contains an element, denoted u^+ , such that $u^+ > u$ and $y \geq u^+$ for all $y > u$, with $y \in P$.

Lemma 9.1. *If U is countable, the following properties are equivalent for every element $C \in \widehat{\overline{G}[U]}$:*

- (i) C is completely meet-irreducible;
- (ii) $C \in \overline{G}[U]$ and there is an element $x \in U \setminus C$ such that C is maximal among the set of copies $\overline{G}[U]$ not containing x .

Proof. Let C be a completely meet-irreducible element of $\widehat{\overline{G}[U]}$. Then trivially $C \in \overline{G}[U]$. Let $x \in U \setminus C$. According to Corollary 5.2 there is some maximal member of $\overline{G}[U]$ which contains C and not x . For each $x \in U \setminus C$, let C_x be such an element. Then clearly, $C = \bigcap \{C_x : x \in U \setminus C\}$. Since C is completely meet-irreducible then $C = C_x$ for

some $x \in U \setminus C$. Conversely, set $C^+ := \{J \in \overline{\mathbb{G}}[U] : \{x\} \cup C \subseteq J\}$. Since, by hypothesis every $J \in \overline{\mathbb{G}}[U]$ with $C \subseteq J$ will contain x , it follows that $C^+ \subseteq J$. Hence C is completely meet-irreducible. \square

Corollary 9.1. *If U is countable, every member of $\widehat{\overline{\mathbb{G}}[U]}$ is an intersection of completely meet-irreducible elements.*

Proof. Let $J \in \widehat{\overline{\mathbb{G}}[U]}$. For every $x \in U \setminus J$ there is some $C' \in \overline{\mathbb{G}}[U]$ such that $J \subseteq C'$ and $x \in U \setminus C'$. According to Lemma 9.1 above every C maximal among the members of $\overline{\mathbb{G}}[U]$ containing C' and not containing x is completely meet-irreducible. The intersection of those C is J . \square

Theorem 9.1. *Let U be a countable set and \mathbb{G} be a subgroup of $\mathfrak{S}(U)$. Then for every subset S of U , the ranked closure $\mathbf{rc}(S)$ of S , is equal to the least element of $\widehat{\overline{\mathbb{G}}[U]}$ containing S , where $\overline{\mathbb{G}}[U]$ is the topological closure of $\overline{\mathbb{G}}[U]$ in $\mathcal{P}(U)$ for the powerset topology .*

Proof. According to Lemma 2.1, $\overline{\mathbb{G}}[U] = \overline{\mathbb{G}^{\text{rc}}[U]}$, thus $\overline{\mathbb{G}}[U]$ is closed under intersection, and the closure of S w.r.t. this closure system is equal to the closure of S w.r.t. $\overline{\mathbb{G}^{\text{rc}}[U]}$. \square

Lemma 9.2. *Let U be a countable set and \mathbb{G} be a subgroup of $\mathfrak{S}(U)$. Then the closure operator associated with $\widehat{\overline{\mathbb{G}}[U]}$ is algebraic if and only if $\widehat{\overline{\mathbb{G}}[U]} = \overline{\overline{\mathbb{G}}[U]}$.*

Proof. The Moore family $\widehat{\overline{\mathbb{G}}[U]}$ defines an algebraic closure system if and only if it is closed in $\mathcal{P}(U)$ equipped with the product topology. Since $\overline{\mathbb{G}}[U] \subseteq \widehat{\overline{\mathbb{G}}[U]} \subseteq \overline{\overline{\mathbb{G}}[U]}$ by Lemma 2.1, it is closed if and only if it is equal to $\overline{\overline{\mathbb{G}}[U]}$. \square

Problem 9.2. *Is $\mathbf{rc}(S) = \mathbf{ic}(S)$ for all subsets S of U ?*

Problem 9.3. *Suppose that there is more than one copy. Does then U contain an infinite independent set w.r.t the closure system given by the collection of intersections of copies. That is an infinite subset S such that $x \notin \mathbf{ic}(S \setminus \{x\})$ for every $x \in S$? Does this conclusion hold under the additional assumption that the algebraic closure of every finite set is finite? (The group is then algebraically finite.)*

10. EMBEDDING THE POWERSSET OF \aleph_0

Theorem 10.1. *Let $\mathfrak{M} := (M, R_i)_{i \in I}$ be an infinite countable relational structure with language \mathcal{L} in which $T := \text{Th}(\mathfrak{M})$ is \aleph_0 -categorical. Let*

\mathbb{Q} be the set of rational numbers and let \mathcal{B} be the lattice of subsets of \mathbb{Q} ordered under set inclusion. Let \mathcal{C} be the family of subsets $N \subseteq M$ for which $\mathfrak{M} \upharpoonright N$, the restriction of \mathfrak{M} to N , is isomorphic to \mathfrak{M} . (i.e., \mathcal{C} is the set of copies.) Then for \mathcal{C} ordered by set inclusion there exists an order embedding of \mathcal{B} into \mathcal{C} .

Proof. Step 1: Let $\mathfrak{M}_0 := \mathfrak{M}$ with language $\mathcal{L}_0 := \mathcal{L}$. Add Skolem functions expanding \mathfrak{M} to a structure \mathfrak{M}' . This is done iteratively. Fix an element, say $\nu \in M$. For each L_0 formula $\exists x\varphi(x, y_1, \dots, y_n)$, with at most y_1, \dots, y_n occurring freely, add a function $f_{\exists x\varphi} : M^n \rightarrow M$ and axiom:

$$\exists x\varphi(x, y_1, \dots, y_n) \leftrightarrow \varphi[f_{\exists x\varphi}(y_1, \dots, y_n), y_1, \dots, y_n].$$

Expand \mathfrak{M}_0 to \mathfrak{M}_1 using the axiom of choice to pick a witness when $\exists x\varphi(x, y_1, \dots, y_n)$ holds for $f_{\exists x\varphi}(y_1, \dots, y_n)$ and define $f_{\exists x\varphi}(y_1, \dots, y_n) = \nu$ otherwise. This gives the language \mathcal{L}_1 and the expansion \mathfrak{M}_1 of \mathfrak{M}_0 . Iterate to form $\mathcal{L}_k \subseteq \mathcal{L}_{k+1}$ and \mathfrak{M}_{k+1} an expansion of \mathfrak{M}_k . Then \mathcal{L}' is the union of the \mathcal{L}_k and \mathfrak{M}' the common expansion of the structures \mathfrak{M}_k in the natural way. Let T' be the theory of \mathfrak{M}' .

It is a standard fact that if \mathcal{S} with domain S is a substructure, (in the expanded language,) of \mathfrak{M}' then it is an elementary substructure of \mathfrak{M}' . (It is easy to check that if $\{n_1, n_2, \dots, n_k\} \subseteq S$ and if $\exists x\varphi(x, n_1, n_2, \dots, n_k)$ is a sentence true in \mathfrak{M}' then $f_{\exists x\varphi}(n_1, \dots, n_k) \in S$ and $\mathfrak{M}' \models \varphi[f_{\exists x\varphi}(n_1, \dots, n_k), n_1, \dots, n_k]$.)

Step 2: We use Theorem 3.3.10 of [3]:

Theorem: Let T be a theory in \mathcal{L} with infinite models, and let $\langle X, < \rangle$ be a simply ordered set. Then there is a model \mathfrak{A} of T with X subset of the domain of \mathfrak{A} and such that X is a set of indiscernibles in \mathfrak{A} .

Indiscernibles with respect to Theorem 10 in [3] means order indiscernibles. Translated to our notation we obtain:

Theorem: There is a model \mathfrak{M}'' of \mathcal{L}' with \mathbb{Q} subset of the domain M'' of \mathfrak{M}'' and such that \mathbb{Q} is a set of indiscernibles in \mathfrak{M}'' .

Let \mathfrak{A} be the reduct of \mathfrak{M}'' to \mathcal{L} . Because the structure \mathfrak{M} is \aleph_0 -categorical we may assume without loss of generality that $\mathfrak{M} = \mathfrak{A}$.

For $X \subset \mathbb{Q}$ let \mathfrak{M}'_X be the closure of X under Skolem functions. Because \mathfrak{M}'_X is an elementary restriction of \mathfrak{M}' , the theory of \mathfrak{M}'_X is T' and hence the reduct to \mathcal{L} is a model whose theory is T . Implying, again due to categoricity, that \mathfrak{M}'_X is a copy of \mathfrak{M} . Note that $\nu \in \mathfrak{M}'_X$ only if there is a term $f_{\exists x\varphi}(x_1, \dots, x_k)$ for which $\exists x\varphi(x, x_1, \dots, x_k)$ holds. Also, it is clear that $X \subseteq X_1$ implies $\mathfrak{M}'_X \subseteq \mathfrak{M}'_{X_1}$. It remains to prove that if $X \neq X_1$ then $\mathfrak{M}'_X \neq \mathfrak{M}'_{X_1}$. This will follow from the following:

Claim: Let $X \subseteq \mathbb{Q}$ then $\mathfrak{M}'_X \cap X = X$.

Let $y \notin X$ but $y \in \mathfrak{M}'_X$, for a contradiction. Then there is a term t of \mathcal{L}' and rationals $x_1 < \dots < x_k$ in X so that $y = t(x_1, \dots, x_k)$. According to construction $\mathfrak{M}' \models \exists! x (x = t(x_1, \dots, x_k))$. But if a rational y is in the same interval of x_1, \dots, x_k as x then $\mathfrak{M}' \models y = t(x_1, \dots, x_k)$, because the rationals form a set of order indiscernibles. □

Theorem 10.2. *Let U be a countable set and G be an oligomorphic subgroup of $\mathfrak{S}(U)$. Then there exists an embedding of $(\mathcal{P}(\omega); \subseteq)$, the power set of ω ordered by inclusion, into $(\overline{G}[U]; \subseteq)$, the set of copies for the group G ordered by inclusion.*

Proof. Let \mathfrak{M} be the homogeneous canonical relational structure associated with the group G . (See [1] page 26.) Then $G = \text{Aut}(\mathfrak{M})$. It follows from [16] that the structure \mathfrak{M} is \aleph_0 -categorical. Let \mathcal{C} be the family of subsets $N \subseteq M$ for which $\mathfrak{M} \upharpoonright N$, the restriction of \mathfrak{M} to N , is isomorphic to \mathfrak{M} . Then \mathcal{C} is the set of copies of \mathfrak{M} . The structure \mathfrak{M} is homogeneous implying that \mathcal{C} is also the set of copies for the group G , that is $\mathcal{C} = \overline{G}[U]$. The Theorem follows from Theorem 10.1. □

REFERENCES

- [1] P.J. Cameron, *Oligomorphic permutation groups*. Cambridge University Press, Cambridge, 1990.
- [2] P.J. Cameron, Metric and Topological Aspects of the Symmetric Group of Countable Degree, *European J. Combinatorics* (1996) **17**, 135–142.
- [3] C.C. Chang, H. J. Keisler, *Model Theory*, third edition, Dover publications, (2012)
- [4] R. Fraïssé, Sur l'extension aux relations de quelques propriétés des ordres, *Ann. Sci. Ecole Norm. Sup.* **71** (1954), pp. 361–388.
- [5] R. Fraïssé, *Theory of Relations*, Revised Edition, in: *Studies in Logic and the Foundations of Mathematics*, **145**, North Holland 2000 ISBN 0-444-50542-3 CV 0.
- [6] G. Grätzer, *General lattice theory*. With appendices by B. A. Davey, R. Freese, B. Ganter, M. Greferath, P. Jipsen, H. A. Priestley, H. Rose, E. T. Schmidt, S. E. Schmidt, F. Wehrung and R. Wille. Reprint of the 1998 second edition, Birkhäuser Verlag, Basel, 2003
- [7] A. S. Kechris, *Classical descriptive set theory*, *Graduate Texts in Mathematics*, vol. 156, Springer-Verlag, New York, 1995.
- [8] K. Kuratowski, *Topology*, vol I, Academic Press, 1966.
- [9] M. Kurilić, Posets of copies of countable scattered linear orders. *Ann. Pure Appl. Logic* 165 (2014), no. 3, 895-912.
- [10] M. Kurilić, Posets of isomorphic substructures of relational structures. *Zb. Rad. (Beogr.)* 17(25) (2015), Selected topics in combinatorial analysis, 117-144.

- [11] M. Kurilić, S. Todorčević, The poset of all copies of the random graph has the 2-localization property, *Ann. Pure Appl. Logic* **167** (2016), no. 8, 649-662.
- [12] M. Kurilić, B. Kuzeljević, Antichains of Copies of Ultrahomogeneous Structures, arXiv:1904.00656, April 2019.
- [13] J-F. Pabion, Relations préhomogènes, *C.R. Acad. Sciences Paris, Sér. A-B t.274(7)*, 1972, 529-531.
- [14] C. Laflamme, M. Pouzet, N. Sauer, and R. Woodrow, Siblings of an \aleph_0 -categorical relational structure, arXiv preprint arXiv:1811.04185 (2018).
- [15] M. Pouzet, Modèle universel d'une théorie n -complète: modèle uniformément préhomogène, *C.R. Acad. Sci. Paris, Sér. A-B t.274(7)*, 1972, 695-698.
- [16] C. Ryll-Nardzewski, On categoricity in power $\leq \aleph_\alpha$. *Bull. Acad. Pol. Sci. Sér. Math. Astr. Phys.* **7** (1959), pp. 545-548.
- [17] J. G. Rosenstein, Linear orderings, *Academic Press* **98**, (1982).
- [18] N. Sauer, Copies and equimorphic structures, March 29, 2019. FG1 Seminar, TU Wien.

MATHEMATICS & STATISTICS DEPARTMENT, UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA T2N 1N4

E-mail address: laflamme@ucalgary.ca

ICJ, MATHÉMATIQUES, UNIVERSITÉ CLAUDE-BERNARD LYON1, 43 BD. 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE AND MATHEMATICS & STATISTICS DEPARTMENT, UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA T2N 1N4

E-mail address: pouzet@univ-lyon1.fr

MATHEMATICS & STATISTICS DEPARTMENT, UNIVERSITY OF CALGARY, CALGARY, T2N1N4, ALBERTA, CANADA T2N1N4

E-mail address: nsauer@math.ucalgary.ca

MATHEMATICS & STATISTICS DEPARTMENT, UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA T2N 1N4

E-mail address: woodrow@ucalgary.ca