

On étale hypercohomology of henselian regular local rings with values in p -adic étale Tate twists

Makoto Sakagaito

Indian Institute of Science Education and Research, Bhopal
E-mail address: sakagaito43@gmail.com

Abstract

In this paper, we prove that an étale hypercohomology of the henselization R of a local ring of a semistable family over a discrete valuation of mixed characteristic $(0, p)$ with values in p -adic étale twists and a logarithmic Hodge-Witt cohomology of the residue field of R are isomorphic. As one of applications, we prove the local-global principle for Galois cohomology groups over function fields of curves over a mixed characteristic excellent henselian discrete valuation ring.

1 Introduction

Let k be a field, X a *normal crossing variety* over k , i.e. a pure-dimensional scheme of finite type over $\text{Spec}(k)$ which is everywhere étale locally isomorphic to

$$\text{Spec}(k[T_0, \dots, T_d]/(T_0 T_1 \cdots T_r))$$

for some integer r with $0 \leq r \leq d = \dim X$ (cf. [21, p.707]).

Let us denote the set of points on X of codimension i by X^i . Then we have three kind of étale sheaves $W_r \omega_{X, \log}^n$ ([14]),

$$\lambda_{X, r}^n := \text{Im} \left(d \log : (\mathbb{G}_{m, X})^{\otimes n} \rightarrow \bigoplus_{x \in X^0} i_{x*} W_r \Omega_{x, \log}^n \right)$$

and

$$\nu_{X, r}^n := \text{Ker} \left(\partial : \bigoplus_{x \in X^0} i_{x*} W_r \Omega_{x, \log}^n \rightarrow \bigoplus_{x \in X^1} i_{x*} W_r \Omega_{x, \log}^{n-1} \right)$$

([22, p.527, 2.2]) which agree with the logarithmic Hodge-Witt sheaves $W_r \Omega_{X, \log}^n$ ([15]) in the case where X is a smooth variety (cf. [15, p.528, Théorème 2.4.2], [11]). Moreover, we have inclusions of étale sheaves

$$\lambda_{X, r}^n \subset W_r \omega_{X, \log}^n \subset \nu_{X, r}^n$$

([21, p.736, Proposition 4.2.1]) and these inclusions are not equalities in general (cf. [21, p.737, Remark 4.2.3]). In §2, we prove the followings:

THEOREM 1.1. (Theorem 2.6) Let A be a local ring of a normal crossing variety over a field of positive characteristic $p > 0$. Then the sequence

$$0 \rightarrow H_{\text{ét}}^s(A, v_r^n) \rightarrow H_{\text{ét}}^s(k(A), v_r^n) \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(1)}} H_x^{s+1}(A_{\text{ét}}, v_r^n) \rightarrow \dots$$

is exact for any integers s and $r > 0$.

THEOREM 1.2. (Theorem 2.7) Let X be a normal crossing variety over a field of characteristic $p > 0$, A the henselization of a local ring $\mathcal{O}_{X,x}$ of x in X and k the residue field of A . Let n be a non negative integer and r a positive integer. Then the homomorphism

$$H_{\text{ét}}^1(A, \lambda_{A,r}^n) \rightarrow H_{\text{ét}}^1(k, \lambda_{k,r}^n) \quad (1)$$

is an isomorphism.

In the case where X is a smooth variety over a field of characteristic $p > 0$, Theorem 1.2 is [20, Theorem 5.3]. The proof of Theorem 1.2 is reduced to [20, Theorem 5.3] by using the covariant functoriality of the sheaves $\lambda_{X,r}^n$ for normal crossing varieties X ([21, p.734, Corollary 3.5.3]). If we replace $\lambda_{A,r}^n$ with $v_{A,r}^n$ in the homomorphism (1), the homomorphism (1) is not an isomorphism unless A is smooth over a field of positive characteristic. See Remark 2.8.

Let B be a discrete valuation ring of mixed characteristic $(0, p)$ with the quotient field K . Let \mathfrak{X} be a *semistable family* over $\text{Spec}(B)$, i.e. a regular scheme of pure dimension which is flat of finite type over $\text{Spec}(B)$, $\mathfrak{X} \otimes_B K$ is smooth over $\text{Spec}(K)$, and the special fiber of \mathfrak{X} is a reduced divisor with normal crossings on \mathfrak{X} .

Then we prove the following as an application of Theorem 1.2:

THEOREM 1.3. (Theorem 3.5) Let \mathfrak{X} be the same as above, R the henselization of a local ring $\mathcal{O}_{\mathfrak{X},x}$ of x in \mathfrak{X} and k the residue field of R . Let n be a non negative integer and r be a positive integer. Then we have an isomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \simeq H_{\text{ét}}^1(k, \lambda_{k,r}^n)$$

where $\mathfrak{T}_r(n)$ is the p -adic Tate twist (cf.[22, p.537, Definition 4.2.4]).

Theorem 1.3 has several applications. Let us denote the Bloch's cycle complex for the Zariski topology ([2], [16]) by $\mathbb{Z}(n)$. For a positive integer l , $\mathbb{Z}(n) \otimes \mathbb{Z}/l$ denotes by $\mathbb{Z}/l(n)$. Thus, $\mathbb{Z}(n)_{\text{ét}}$ (resp. $\mathbb{Z}/l(n)_{\text{ét}}$) denotes the étale sheafification of $\mathbb{Z}(n)$ (resp. $\mathbb{Z}/l(n)$).

Then, the first application of Theorem 1.3 is as follows:

PROPOSITION 1.4. (Proposition 4.5) Let R be the same as in Theorem 1.3. Then we have

$$H_{\text{Zar}}^i(R, \mathbb{Z}/p^r(n)) = 0$$

for $i = n + 1$.

By Proposition 1.4 and [23, p.209, Remark 7.2], we have an isomorphism

$$\tau_{\leq n+1}(\mathbb{Z}/p^r(n)_{\text{ét}}) \simeq \mathfrak{T}_r(n) \quad (2)$$

in $D^b(\mathfrak{X}_{\text{ét}}, \mathbb{Z}/p^r\mathbb{Z})$ which is the derived category of bounded complexes of étale $\mathbb{Z}/p^r\mathbb{Z}$ -sheaves on \mathfrak{X} . If \mathfrak{X} is smooth over $\text{Spec}(B)$, then truncation is unnecessary by [8, p.786, Corollary 4.4].

Then, by using (2), the second application of Theorem 1.3 is expressed as follows:

THEOREM 1.5. (Theorem 4.8 and (2)) Let \mathfrak{X} be a semistable family over a discrete valuation ring of mixed characteristic $(0, p)$ and $R = \tilde{\mathcal{O}}_{\mathfrak{X}, x}$ the henselian local ring of x in \mathfrak{X} . Suppose that $\dim(R) = 2$. Then the sequence

$$0 \rightarrow \mathbf{H}_{\text{ét}}^{n+1}(R, \mathbb{Z}/l(n)) \rightarrow \mathbf{H}_{\text{ét}}^{n+1}(k(R), \mathbb{Z}/l(n)) \rightarrow \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1} \mathbf{H}_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathbb{Z}/l(n)) \quad (3)$$

is exact for $l = p^r$. Here $k(R)$ is the fraction field of R .

Let l be a positive integer which is invertible in B and μ_l the étale sheaf of l -th roots of unity. Then we have an isomorphism

$$\tau_{\leq n+1}(\mathbb{Z}/l(n)_{\text{ét}}) \simeq \mu_l^{\otimes n} \quad (4)$$

in $D^b(\mathfrak{X}_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$ (See Remark 4.7). If $\dim(B) \leq 1$ and \mathfrak{X} is smooth over $\text{Spec}(B)$, then truncation is unnecessary by [10, Theorem 1.5] and [8, p.786, Corollary 4.4]. In the case where R is equi-characteristic and $(l, \text{char}(R)) = 1$, Theorem 1.5 is a part of the Bloch-Ogus Theorem ([4]) by (4).

We are able to prove the following local-global principle as an application of Theorem 1.5 (and Theorem 1.3):

THEOREM 1.6. (Corollary 4.9 and Theorem 4.11) Let X be a regular scheme over a henselian discrete valuation ring B of mixed characteristic $(0, p)$ and Y the special fiber of X . Let $\tilde{\mathcal{O}}_{X, \mathfrak{p}}$ be the henselization of a local ring $\mathcal{O}_{X, \mathfrak{p}}$ of \mathfrak{p} in X , $\kappa(\mathfrak{p})$ the residue field of $\mathfrak{p} \in X$ and $k(X)$ (resp. $k(\tilde{\mathcal{O}}_{X, \mathfrak{p}})$) the ring of rational functions on X (resp. $\text{Spec}(\tilde{\mathcal{O}}_{X, \mathfrak{p}})$). Suppose that $\dim(X) = 2$.

Then the natural map

$$\mathbf{H}_{\text{ét}}^{n+1}(k(X), \mu_{p^r}^{\otimes n}) \rightarrow \bigoplus_{\mathfrak{p} \in X^1 \setminus Y^0} \mathbf{H}_{\text{ét}}^n(\kappa(\mathfrak{p}), \mu_{p^r}^{\otimes(n-1)}) \oplus \bigoplus_{\mathfrak{p} \in Y^0} \mathbf{H}_{\text{ét}}^{n+1}(k(\tilde{\mathcal{O}}_{X, \mathfrak{p}}), \mu_{p^r}^{\otimes n})$$

is injective for $n \geq 1$ in the following cases:

- (i) X is a scheme \mathfrak{X} which is a proper and semistable family over a henselian discrete valuation ring B of mixed characteristic $(0, p)$.
- (ii) $X = \tilde{\mathcal{O}}_{\mathfrak{X}, \mathfrak{p}}$. Here \mathfrak{X} is the same as in (i) and $\mathfrak{p} \in \mathfrak{X}^2$.

This implies that the local-global map

$$\mathbf{H}_{\text{ét}}^{n+1}(k(X), \mu_l^{\otimes n}) \rightarrow \prod_{\mathfrak{p} \in X^1} \mathbf{H}_{\text{ét}}^{n+1}(k(\tilde{\mathcal{O}}_{X, \mathfrak{p}}), \mu_l^{\otimes n}) \quad (5)$$

is injective for $n \geq 1$ and $l = p^r$.

Suppose that B is equi-characteristic and $(l, \text{char}(B)) = 1$. In the case (i), Colliot-Thélène raised a question whether the local-global map (5) is injective ([5]). In [12, Theorem 3.3.6], Harbater–Hartmann–Krashen provided an affirmative answer to this question in this case.

In the case where B is mixed characteristic $(0, p)$, due to Hu’s method (cf.[13, The proof of Theorem 2.5], [13, Remark 2.6 (2)]), Theorem 1.6 (i) follows from Theorem 1.5 (or Theorem 1.6 (ii)) without assuming $(l, p) = 1$. If $(l, p) = 1$, then Theorem 1.5 has been proved in [19, Theorem 1.1]. Theorem 1.6 (i) is an extension of [20, Theorem 6.3].

Acknowledgments

I thank Indian Institute of Science Education and Research, Bhopal where the main part of this work was completed.

Notations

For a scheme X , X^i denotes the set of points on X of codimension i . Moreover, $k(X)$ denotes the ring of rational functions on X and $\kappa(x)$ denotes the residue field of $x \in X$. If $X = \text{Spec}(A)$ for a ring A , $k(A)$ denotes $k(\text{Spec}(A))$. For a scheme X , X_{Zar} , X_{Nis} and $X_{\text{ét}}$ denote the category of étale schemes over X equipped with the Zariski, Nisnevich and étale topology, respectively. For a scheme X , $D(X)$ denotes the derived category of complexes of étale sheaves of abelian groups on X , and $D_+(X) \subset D(X)$ denotes the full subcategory of complexes that are bounded below.

2 Logarithmic Hodge-Witt sheaves

In this section, we study the logarithmic Hodge-Witt sheaves $v_{X,r}^n$ and $\lambda_{X,r}^n$ ([21]) for a normal crossing variety X .

2.1 $v_{X,r}^n$

Let X be a normal crossing variety over a field of characteristic $p > 0$. Let n be a non negative integer and r a positive integer. We show a relation between

$$v_{X,r}^n := \text{Ker} \left(\partial : \bigoplus_{x \in X^0} i_{x*} W_r \Omega_{x,\log}^n \rightarrow \bigoplus_{x \in X^1} i_{x*} W_r \Omega_{x,\log}^{n-1} \right)$$

([21]) and $\mathbb{Z}/p^r(n)$ of Zariski sheaves, where $\mathbb{Z}(n)$ is the Bloch’s cycle complex for the Zariski topology ([2],[16]) and $\mathbb{Z}/p^r(n) = \mathbb{Z}(n) \otimes \mathbb{Z}/p^r$.

PROPOSITION 2.1. Let X be a normal crossing variety over a field. Then we have

$$\mathcal{H}^i(\mathbb{Z}(n)) = 0$$

for $i \geq n + 1$.

Proof. It suffices to show the statement in the case where X is the spectrum of a local ring of a normal crossing variety. Hence it suffices to show

$$H_{\text{Zar}}^i(A, \mathbb{Z}(n)) = 0 \quad (6)$$

for $i \geq n + 1$ in the case where A is a local ring of a normal crossing variety by [8, p.779, Theorem 3.2 b)]. We prove the equation (6) by induction on $\#(\text{Spec}(A)^0)$.

In the case where $\#(\text{Spec}(A)^0) = 1$, the equation (6) follows from [2, p.299, Theorem (10.1)].

Assume that the equation (6) holds in the case where $\#(\text{Spec}(A)^0) \leq r$. Let

$$\text{Spec}(A_1), \dots, \text{Spec}(A_{r+1}) \quad (7)$$

be the irreducible components of $\text{Spec}(A)$. Then elements of the set (7) and

$$\text{Spec}(B_1) = \text{Spec}(A_1) \cap \text{Spec}(A_{r+1}), \dots, \text{Spec}(B_r) = \text{Spec}(A_r) \cap \text{Spec}(A_{r+1})$$

are smooth varieties by the definition of normal crossing variety. Let

$$\text{Spec}(B) = \bigcup_{j=1}^{j=r} \text{Spec}(B_j) = \left(\bigcup_{j=1}^{j=r} \text{Spec}(A_j) \right) \cap \text{Spec}(A_{r+1}).$$

Then we have

$$H_{\text{Zar}}^i(B, \mathbb{Z}(n)) = 0$$

for $i \geq n + 1$ by the inductive hypothesis. Since the homomorphism

$$H_{\text{Zar}}^n(\text{Spec}(A_{r+1}) \setminus \text{Spec}(B), \mathbb{Z}(n)) \rightarrow H_{\text{Zar}}^{n-1}(\text{Spec}(B), \mathbb{Z}(n-1))$$

is surjective by the localization theorem ([8, p.780, Corollary 3.3]) and we have

$$\text{Spec}(A) \setminus \text{Spec}(B) = \left(\bigcup_{j=1}^{j=r} \text{Spec}(A_j) \setminus \text{Spec}(B) \right) \bigoplus (\text{Spec}(A_{r+1}) \setminus \text{Spec}(B)),$$

the homomorphism

$$H_{\text{Zar}}^n(\text{Spec}(A) \setminus \text{Spec}(B), \mathbb{Z}(n)) \rightarrow H_{\text{Zar}}^{n-1}(\text{Spec}(B), \mathbb{Z}(n-1)) \quad (8)$$

is also surjective. For $i \geq n + 1$, we have

$$H_{\text{Zar}}^i\left(\bigcup_{j=1}^{j=r} \text{Spec}(A_j) \setminus \text{Spec}(B), \mathbb{Z}(n)\right) = H_{\text{Zar}}^i(\text{Spec}(A_{r+1}) \setminus \text{Spec}(B), \mathbb{Z}(n)) = 0$$

by the inductive hypothesis and the localization theorem. Hence we have

$$H_{\text{Zar}}^i(\text{Spec}(A) \setminus \text{Spec}(B), \mathbb{Z}(n)) = 0 \quad (9)$$

for $i \geq n + 1$. By using the localization theorem, the equation (6) follows from the equation (9) and the surjectivity of the homomorphism (8). This completes the proof. \square

PROPOSITION 2.2. Let X be a normal crossing variety over a field.

1. If $i : Z \rightarrow X$ is a closed subscheme of codimension c , then the canonical map

$$\tau_{\leq n+2}(\mathbb{Z}(n-c)_{\acute{e}t}[-2c]) \rightarrow \tau_{\leq n+2}Ri^!\mathbb{Z}(n)_{\acute{e}t}$$

is a quasi-isomorphism, where $\mathbb{Z}(n)_{\acute{e}t}$ is the étale sheafification of $\mathbb{Z}(n)$.

2. Let $\varepsilon : X_{\acute{e}t} \rightarrow X_{\text{Zar}}$ be the change of site map. Then the canonical map induces a quasi-isomorphism

$$\mathbb{Z}(n)_{\text{Zar}} \xrightarrow{\sim} \tau_{\leq n+1}R\varepsilon_*\mathbb{Z}(n)_{\acute{e}t}.$$

Proof. By the same argument as in the proof of [20, Proposition 2.1], Proposition 2.2.1 follows from Proposition and Proposition 2.2.2. So it suffices to show Proposition 2.2.2. Moreover, it suffices to show Proposition 2.2.2 in the case where X is the spectrum of a local ring A of a normal crossing variety.

We prove Proposition 2.2.2 by induction on $\#(\text{Spec}(A)^0)$. By, Proposition 2.2.2 holds in the case where $\#(\text{Spec}(A)^0) = 1$.

Assume that Proposition 2.2.2 holds in the case where $\#(\text{Spec}(A)^0) \leq r$. Suppose that $\#(\text{Spec}(A)^0) = r+1$ and $\text{Spec}(A_k)$ ($1 \leq k \leq r+1$) are the irreducible components of $\text{Spec}(A)$.

Let $\text{Spec}(B) = \text{Spec}(A) \cap \text{Spec}(A_{r+1})$ and $i : \text{Spec}(B) \rightarrow \text{Spec}(A)$ be the closed immersion. Let $U = \text{Spec}(A) \setminus \text{Spec}(B)$ and $j : U \rightarrow \text{Spec}(A)$. Then we have

$$U = \left(\bigcup_{k=1}^{k=r} \text{Spec}(A_k) \setminus \text{Spec}(B) \right) \bigoplus (\text{Spec}(A_{r+1}) \setminus \text{Spec}(B)).$$

Moreover, $\bigcup_{k=1}^{k=r} \text{Spec}(A_k) \setminus \text{Spec}(B)$ and $\text{Spec}(A^r) \setminus \text{Spec}(B)$ are normal crossing varieties,

$$\# \left(\bigcup_{k=1}^{k=r} \text{Spec}(A_k) \setminus \text{Spec}(B) \right) = r \text{ and } \#(\text{Spec}(A^r) \setminus \text{Spec}(B)) = 1.$$

Hence Proposition 2.2.2 holds for U by the assumption. Then Proposition 2.2.1 holds for $i : \text{Spec}(B) \rightarrow \text{Spec}(A)$ by the same argument as in the proof of [20, Proposition 2.1]. Moreover, $\#(\text{Spec}(B)^0) = r$ and Proposition 2.2.2 holds for $\text{Spec}(B)$ by the assumption. Therefore Proposition 2.2.2 follows by the same argument as in the proof of [8, Theorem 1.2.2]. This completes the proof. \square

PROPOSITION 2.3. Let X be a normal crossing variety over a field of characteristic $p > 0$. Then we have an isomorphism in $D^b(X_{\text{Zar}}, \mathbb{Z}/p^r\mathbb{Z})$

$$\mathbb{Z}/p^r(n) \simeq v_{X,r}^!\mathbb{Z}[-n],$$

where $D^b(X_{\text{Zar}}, \mathbb{Z}/p^r\mathbb{Z})$ is the derived category of bounded complexes of Zariski $\mathbb{Z}/p^r\mathbb{Z}$ -sheaves on X .

Proof. It suffices to show the statement in the case where X is the spectrum of a local ring of a normal crossing variety. We prove the statement by induction on $\dim X$. In the case where $\dim X = 0$, the statement follows from [9, p.491, Theorem 8.3].

Assume that the statement holds in the case where $\dim X \leq m$. Suppose that $\dim X = m + 1$. Let $i : Y \rightarrow X$ be a closed immersion of codimension 1, Y regular and U the complement of Y . Let Y be the spectrum of a regular local ring A' . Then A' is a local ring of a regular ring of finite type over a field. By Quillen's method (cf.[18, §7, The proof of Theorem 5.11]),

$$A' = \lim_{\rightarrow} A'_i \quad (10)$$

where A'_i is a local ring of a smooth algebra over \mathbb{F}_p and the maps $A'_i \rightarrow A'_j$ are flat. Since $\dim U = m$ and

$$\cdots \rightarrow i_* \mathbb{Z}/p^r(n-1)^Y[-2] \rightarrow \mathbb{Z}/p^r(n)^X \rightarrow Rj_* \mathbb{Z}/p^r(n)^U \rightarrow \cdots$$

is a distinguished triangle by [2, p.277, Theorem (3.1)], we have a quasi-isomorphism

$$\tau_{\leq n}(\mathbb{Z}/p^r(n)) \simeq v_{X,r}^n[-n]$$

by [9, p.491, Theorem 8.3]. Moreover, we have

$$\mathcal{H}^i(\mathbb{Z}/p^r(n)) = 0$$

for $i > n$ by Proposition 2.1. This completes the proof. \square

COROLLARY 2.4. Let A be a local ring at a point of a normal crossing variety over a field of positive characteristic $p > 0$. Then the sequence

$$\begin{aligned} 0 \rightarrow H^0(A, v_r^n) \rightarrow H^0(k(A), v_r^n) \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(1)}} H^0(\kappa(x), v_r^n) \\ \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(2)}} H^0(\kappa(x), v_r^n) \rightarrow \cdots \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(i)}} H^0(\kappa(x), v_r^n) \rightarrow \cdots \end{aligned}$$

is exact for any integer n .

Proof. We consider the spectral sequence

$$E_1^{s,t} = \bigoplus_{x \in \text{Spec}(A)^{(s)}} H_{\text{Zar}}^{2n-s+t}(\kappa(x), \mathbb{Z}/p^r(n-s)) \Rightarrow E^{s+t} = H_{\text{Zar}}^{2n+s+t}(A, \mathbb{Z}/p^r(n)) \quad (11)$$

(cf.[8, p.782]). By Proposition 2.1, we have $E_1^{s,t} = 0$ for $t \neq -n$ and $E^{s+t} = 0$ for $s+t \neq -n$. Hence the statement follows from the spectral sequence (11). \square

PROPOSITION 2.5. Let A be a local ring of characteristic $p > 0$ and F be an étale p -torsion sheaf.

Then we have

$$H_x^{r+j}(A_{\text{ét}}, F) = 0$$

for $r \geq 2$ and $x \in \text{Spec}(A)^{(j)}$.

Proof. We prove the statement by induction on $\text{codim}(\{\bar{x}\}, \text{Spec}(A)) = \dim(A_x)$.

In the case where $\dim(A_x) = 0$, the statement is true because a field k of characteristic $p > 0$ has p -cohomological dimension $\text{cd}_p(k)$ at most 1.

Assume that the statement is true for $\dim(A_x) \leq j$. Then we prove the statement in the case where $\dim(A_x) = j + 1$. By [17, p.93, III, Corollary 1.28], we have

$$H_x^s(A_{\text{ét}}, F) = H_x^s((A_x^h)_{\text{ét}}, F)$$

for any integer $s \geq 0$. So we can assume that A is henselian and $\dim(A) = j + 1$. In order to prove the statement, we use the spectral sequence

$$E_1^{s,t} = \bigoplus_{x \in \text{Spec}(A)^{(s)}} H_x^{s+t}(A_{\text{ét}}, F) \Rightarrow E^{s,t} = H_{\text{ét}}^{s+t}(A, F) \quad (12)$$

(cf.[6, Part 1, §1]). Since $\dim(A) = j + 1$, we have

$$E_2^{s,t} = 0 \quad (13)$$

for $s > j + 1$. Moreover, we have (13) for $s \leq j$ and $t \geq 2$ by the assumption. Hence we have

$$E_2^{j+1,r} = E_{\infty}^{j+1,r} \quad (14)$$

for $r \geq 2$. Since $\text{cd}_p(A) \leq 1$ by [1, Exposé X, Théorème 5.1], we have

$$E_{\infty}^{j+1,r} = E^{j+r+1} = 0 \quad (15)$$

for $r \geq 2$ by the spectral sequence (12). Hence the sequence

$$\bigoplus_{x \in \text{Spec}(A)^{(j)}} H_x^{r+j}(A_{\text{ét}}, F) \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(j+1)}} H_x^{r+j+1}(A_{\text{ét}}, F) \rightarrow 0$$

is exact for $r \geq 2$ by (14) and (15). Therefore we have

$$H_x^{r+j+1}(A_{\text{ét}}, F) = 0$$

for $r \geq 2$ by the assumption. This completes the proof. \square

THEOREM 2.6. Let A be a local ring of a normal crossing variety over a field of positive characteristic $p > 0$. Then the sequence

$$0 \rightarrow H_{\text{ét}}^s(A, \mathbf{v}_r^n) \rightarrow H_{\text{ét}}^s(k(A), \mathbf{v}_r^n) \rightarrow \bigoplus_{x \in \text{Spec}(A)^{(1)}} H_x^{s+1}(A_{\text{ét}}, \mathbf{v}_r^n) \rightarrow \dots$$

is exact for any integers s and $r > 0$.

Proof. In the case where $s < 0$, the statement follows from [21, p.718, Theorem 2.4.2]. In the case where $s = 0$, the statement follows from Corollary 2.4 and [21, p.718, Theorem 2.4.2]. In the case where $s \geq 2$, the statement follows from Proposition 2.5 and [1, Exposé X, Théorème 5.1]. So it suffices to show the statement in the case where $s = 1$. In order to prove the statement, we use the spectral sequence (12). By Corollary 2.4 and [21, p.718, Theorem 2.4.2], we have

$$E_2^{s,t} = 0 \quad (16)$$

for $s > 0$ and $t \leq 0$. By Proposition 2.5, we have the equation (16) for $t \geq 2$. Moreover, we have $E^{s+t} = 0$ for $s + t \geq 2$ by [1, Exposé X, Théorème 5.1]. Hence the statement follows from the spectral sequence (12). \square

2.2 $\lambda_{X,r}^n$

Let n be a non negative integer and r a positive integer. For a normal crossing variety X , we define a kind of generalized logarithmic Hodge-Witt sheaves by

$$\lambda_{X,r}^n := \text{Im} \left(d \log : (\mathbb{G}_{m,X})^{\otimes n} \rightarrow \bigoplus_{x \in X^0} i_{x*} \mathbf{W}_r \Omega_{x,\log}^n \right)$$

(cf. [21]). Then we show the following:

THEOREM 2.7. Let A be the henselization of a local ring of a normal crossing variety over a field of characteristic $p > 0$ and k the residue field of A . Then the homomorphism

$$\mathbf{H}_{\text{ét}}^1(A, \lambda_{A,r}^n) \rightarrow \mathbf{H}_{\text{ét}}^1(k, \lambda_{k,r}^n) \quad (17)$$

is an isomorphism.

Proof. Let C be a finitely generated k -algebra. Then there exists a surjective k -algebra homomorphism

$$k[T_1, \dots, T_N] \rightarrow C$$

for some integer N . Hence A is embedded into an equi-characteristic henselian regular local ring B . Let $i : \text{Spec}(A) \rightarrow \text{Spec}(B)$ be the closed immersion. Then the natural pull-back map

$$i^* : \mathbb{G}_{m,B}^{\otimes n} \rightarrow i_* \left(\mathbb{G}_{m,A}^{\otimes n} \right) \quad (18)$$

is surjective. Moreover the homomorphism (18) induces the pull-back map

$$i^* : \lambda_{B,r}^n \rightarrow i_* \lambda_{A,r}^n \quad (19)$$

(cf. [21, p.732, Theorem 3.5.1]) and the homomorphism (19) is also surjective. Since k has p -cohomological dimension at most 1, we have

$$\mathbf{H}_{\text{ét}}^2(B, \text{Ker}(i^*)) = 0$$

by [8, p.777, The proof of Proposition 2.2.b)]. Hence the homomorphism

$$\mathbf{H}_{\text{ét}}^1(B, \lambda_{B,r}^n) \rightarrow \mathbf{H}_{\text{ét}}^1(A, \lambda_{A,r}^n) \quad (20)$$

which is induced by the pull-back map (19) is surjective. Moreover a composition of the homomorphism (20) and the homomorphism

$$\mathbf{H}_{\text{ét}}^1(A, \lambda_{A,r}^n) \rightarrow \mathbf{H}_{\text{ét}}^1(k, \lambda_{k,r}^n) \quad (21)$$

which is induced by the pull-back map coincides with the homomorphism

$$\mathbf{H}_{\text{ét}}^1(B, \lambda_{A,r}^n) \rightarrow \mathbf{H}_{\text{ét}}^1(k, \lambda_{k,r}^n) \quad (22)$$

which is induced by the pull-back map by [21, p.734, Corollary 3.5.3]. Then the homomorphism (22) is an isomorphism by [20, Theorem 5.3]. Therefore the homomorphism (20) is an isomorphism and the homomorphism (21) is also an isomorphism. This completes the proof. \square

REMARK 2.8. If we replace $\lambda_{A,r}^n$ with $\nu_{A,r}^n$ in the homomorphism (17), the homomorphism (17) is not an isomorphism in general. In fact,

$$\lambda_{A,r}^0 = \mathbb{Z}/p^r \qquad \nu_{A,r}^0 = \underbrace{\mathbb{Z}/p^r \oplus \cdots \oplus \mathbb{Z}/p^r}_{\#(\text{Spec}(A)^0) \text{ times}}$$

and

$$H_{\text{ét}}^1(A, \lambda_{A,r}^0) \neq H_{\text{ét}}^1(A, \nu_{A,r}^0)$$

unless A is smooth over a field of positive characteristic.

3 p -adic Tate twist $\mathfrak{T}_r(n)$

Let B be a discrete valuation ring of mixed characteristic $(0, p)$ with the quotient field K .

Let \mathfrak{X} be a semistable family over $\text{Spec}(B)$, i.e. a regular scheme of pure dimension which is flat of finite type over $\text{Spec}(B)$, $\mathfrak{X}_K = \mathfrak{X} \otimes K$ is smooth over $\text{Spec}(K)$, and the special fiber Y of \mathfrak{X} is a reduced divisor with normal crossings on \mathfrak{X} .

Let j and i be as follows:

$$\mathfrak{X}_K \xrightarrow{j} \mathfrak{X} \xleftarrow{i} Y.$$

Let n be a non negative integer and r a positive integer. In this section, we study the p -adic Tate twist $\mathfrak{T}_r(n)$ (cf.[22, p.537, Definition 4.2.4]). In order to study it, the p -adic vanishing cycle

$$M_r^n := i^* R^n j_* \mu_{p^r}^{\otimes n}$$

plays important roles.

3.1 Review on the structure of the p -adic vanishing cycle

Let the notations be same as above. We define the étale sheaf $\mathcal{H}_{n, \mathfrak{X}_K/Y}^M$ on Y as

$$\mathcal{H}_{n, \mathfrak{X}_K/Y}^M := (i^* j_* \mathcal{O}_{\mathfrak{X}_K}^\times)^{\otimes n} / J,$$

where J denotes the subsheaf which is generated by local sections of the form

$$x_1 \otimes \cdots \otimes x_n \quad (x_i \in i^* j_* \mathcal{O}_{\mathfrak{X}_K}^\times) \text{ with } x_i + x_j = 0 \text{ or } 1$$

for some $1 \leq i < j \leq n$. By [3, (1.2)], there is a natural map

$$\mathcal{H}_{n, \mathfrak{X}_K/Y}^M \rightarrow M_r^n \tag{23}$$

and we define the filtrations U^* and V^* on the p -adic vanishing cycle M_r^n using this map (23), as follows:

DEFINITION 3.1. (1) Let π be a prime element of B . Let $U_{\mathfrak{X}_K}^0$ be the full sheaf $i^* j_* \mathcal{O}_{\mathfrak{X}_K}^\times$. For $q \geq 1$, let $U_{\mathfrak{X}_K}^q$ be the étale subsheaf of $i^* j_* \mathcal{O}_{\mathfrak{X}_K}^\times$ which is generated by local sections of the form $1 + \pi^q \cdot a$ with $a \in i^* \mathcal{O}_{\mathfrak{X}_K}$. We define the subsheaf $U^q \mathcal{H}_{n, \mathfrak{X}_K/Y}^M$ as the part which is generated by $U_{\mathfrak{X}_K}^q \otimes \{i^* j_* \mathcal{O}\}^{\otimes n-1}$.

- (2) We define the subsheaf $U^q M_r^n$ ($q \geq 0$) of M_r^n as the image of $U^q \mathcal{K}_{n, \mathfrak{X}_K/Y}^M$ under the map (23). We define the subsheaf $V^q M_r^n$ ($q \geq 0$) of M_r^n as the part which is generated by $U^{q+1} M_r^n$ and the image of $U^p \mathcal{K}_{n-1, \mathfrak{X}_K/Y}^M \otimes \langle \pi \rangle$ under the map (23).

Then we give a brief review of the structure of M_r^n . For $q \geq 0$,

$$\mathrm{gr}_{U/V}^q M_r^n := U^q M_r^n / V^q M_r^n \quad \text{and} \quad \mathrm{gr}_{V/U}^q M_r^n := V^q M_r^n / U^{q+1} M_r^n$$

are expressed by using subsheaves of the modified differential modules ω_Y^* as follows:

THEOREM 3.2. (Bloch-Kato [3, p.112, Corollary (1.4.1)] /Hyodo [14, p.548, (1.7) Corollary], Sato[23, pp.184–185, Theorem 3.3])

Let the notations be same as above. Then

- (1) The map (23) is surjective, that is, the subsheaf $U^0 M_r^n$ is the full sheaf M_r^n for any $n \geq 0$ and $r > 0$.
- (2) Let e be the absolute ramification index of K , and let $r = 1$. Then for q with $1 \leq q < e' := pe/(p-1)$, there are isomorphisms

$$\mathrm{gr}_{U/V}^q M_1^n \simeq \begin{cases} \omega_Y^{n-1} / \mathcal{B}_Y^{n-1} & ((p, q) = 1), \\ \omega_Y^{n-1} / \mathcal{Z}_Y^{n-1} & (p|q), \end{cases}$$

$$\mathrm{gr}_{V/U}^q M_1^n \xrightarrow{\sim} \omega_Y^{n-2} / \mathcal{Z}_Y^{n-2},$$

where $\omega_Y^n = W_1 \omega_Y^n$ and \mathcal{B}_Y^m (resp. \mathcal{Z}_Y^m) denotes the image of $d : \omega_Y^{m-1} \rightarrow \omega_Y^m$ (resp. the kernel of $d : \omega_Y^m \rightarrow \omega_Y^{m+1}$).

- (3) We have

$$U^q M_1^n = V^q M_1^n = 0$$

for any $q \geq e'$.

We define the étale subsheaf FM_r^n as the part which is generated by $U^1 M_r^n$ and the image of $(\iota^* \mathcal{O}_{\mathfrak{X}}^\times)^{\otimes n}$ under the map (23).

THEOREM 3.3. (Sato [22, p.533, Theorem 3.4.2], [23, p.186, Theorem 3.4]) Let \mathfrak{X} be same as in Theorem 3.2. Then there exists short exact sequence of sheaves on $Y_{\text{ét}}$

$$0 \rightarrow FM_r^n \rightarrow M_r^n \xrightarrow{\sigma_{\mathfrak{X}, r}^n} v_{Y, r}^{n-1} \rightarrow 0,$$

where $\sigma_{\mathfrak{X}, r}^n$ is induced by the boundary map of Galois cohomology groups (cf.[22, p.530, (3.2.5)]). Furthermore there is an isomorphism

$$FM_r^n / U^1 M_r^n \xrightarrow{\sim} \lambda_{Y, r}^n$$

sending a symbol

$$\{x_1, x_2, \dots, x_n\} \quad (x_i \in \iota^* \mathcal{O}_{\mathfrak{X}}^\times)$$

to

$$d \log(\bar{x}_1 \otimes \bar{x}_2 \otimes \dots \otimes \bar{x}_n).$$

Here for a section $x \in \iota^* \mathcal{O}_{\mathfrak{X}}^\times$, \bar{x} denotes its residue class in \mathcal{O}_Y^\times .

3.2 Rigidity

LEMMA 3.4. Let A be the henselization of a local ring of a normal crossing variety over a field of characteristic $p > 0$. Then

$$H_{\text{ét}}^i(A, \mathcal{B}_A^n) = H_{\text{ét}}^i(A, \mathcal{L}_A^n) = 0$$

for $i \geq 1$. Here \mathcal{B}_A^n (resp. \mathcal{L}_A^n) is the image of $d : \omega_A^{n-1} \rightarrow \omega_A^n$ (resp. the kernel of $d : \omega_A^n \rightarrow \omega_A^{n+1}$).

Proof. \mathcal{B}_A^n and \mathcal{L}_A^n are locally free $\mathcal{O}_{\text{Spec}(A)}$ -modules after twisting with Frobenius. Therefore the statement follows from [17, p.114, III, Remark 3.8] and [17, p.103, III, Lemma 2.15]. \square

THEOREM 3.5. Let B be a discrete valuation ring of mixed characteristic $(0, p)$, R the henselization of a local ring of a semistable family over B and k the residue field of R . Then we have an isomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \simeq H_{\text{ét}}^1(k, \lambda_{k,r}^n)$$

where $\mathfrak{T}_r(n)$ is the p -adic Tate twist.

Proof. Let $\iota : \text{Spec}(A) \hookrightarrow \text{Spec}(R)$ be the special fiber. Then it suffices to show that we have an isomorphism

$$H_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_r(n)) \simeq H_{\text{ét}}^1(A, \lambda_{A,r}^n) \quad (24)$$

by Theorem 2.7 and [8, p.777, The proof of Proposition 2.2.b)]. Since k has p -cohomological dimension at most 1, we have

$$H_{\text{ét}}^i(A, \iota^*(\tau_{\leq n-1} R j_* \mu_{p^r}^{\otimes n})) = 0 \quad (25)$$

for $i \geq n+1$ by the spectral sequence

$$H_{\text{ét}}^r(A, \mathcal{H}^s(\iota^*(\tau_{\leq n-1} R j_* \mu_{p^r}^{\otimes n}))) \Rightarrow H_{\text{ét}}^{r+s}(A, \iota^*(\tau_{\leq n-1} R j_* \mu_{p^r}^{\otimes n}))$$

and [8, p.777, The proof of Proposition 2.2.b)]. Hence the homomorphism

$$H_{\text{ét}}^n(A, \iota^*(\tau_{\leq n} R j_* \mu_{p^r}^{\otimes n})) \rightarrow H_{\text{ét}}^0(A, \iota^* R^n j_* \mu_{p^r}^{\otimes n})$$

is surjective and the homomorphism

$$H_{\text{ét}}^{n+1}(A, \iota^*(\tau_{\leq n} R j_* \mu_{p^r}^{\otimes n})) \rightarrow H_{\text{ét}}^1(A, \iota^* R^n j_* \mu_{p^r}^{\otimes n})$$

is an isomorphism by the distinguished triangle

$$\cdots \rightarrow \iota^*(\tau_{\leq n-1} R j_* \mu_{p^r}^{\otimes n}) \rightarrow \iota^*(\tau_{\leq n} R j_* \mu_{p^r}^{\otimes n}) \rightarrow \iota^* R^n j_* \mu_{p^r}^{\otimes n}[-n] \rightarrow \cdots$$

Since we have a commutative diagram with distinguished rows

$$\begin{array}{ccccc} \iota^* \mathfrak{T}_r(n) & \longrightarrow & \iota^*(\tau_{\leq n} R j_* \mu_{p^r}^{\otimes n}) & \longrightarrow & v_{A,r}^{n-1}[-n] \\ \downarrow & & \downarrow & & \parallel \\ FM_r^n[-n] & \longrightarrow & \iota^* R^n j_* \mu_{p^r}^{\otimes n}[-n] & \longrightarrow & v_{A,r}^{n-1}[-n] \end{array}$$

by the definition of $\mathfrak{T}_r(n)$ and FM_r^n , we have an isomorphism

$$\mathbf{H}_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_r(n)) \simeq \mathbf{H}_{\text{ét}}^1(A, FM_r^n)$$

by the five lemma. Moreover we have

$$\mathbf{H}_{\text{ét}}^i(A, U^1 M_1^n) = 0$$

for $i \geq 1$ by Theorem 3.2 and Lemma 3.4. Therefore we have an isomorphism

$$\mathbf{H}_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_1(n)) \simeq \mathbf{H}_{\text{ét}}^1(A, \lambda_{A,1}^n)$$

by Theorem 3.3. Since

$$\mathcal{H}^i(\mathfrak{T}_r(n)) = 0$$

for $i > n$, we can prove

$$\mathbf{H}_{\text{ét}}^{n+2}(A, \iota^* \mathfrak{T}_r(n)) = 0$$

in the same way as the proof of (25). Hence the top row in a commutative diagram

$$\begin{array}{ccccccc} \mathbf{H}_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_r(n)) & \longrightarrow & \mathbf{H}_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_{r+1}(n)) & \longrightarrow & \mathbf{H}_{\text{ét}}^{n+1}(A, \iota^* \mathfrak{T}_1(n)) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbf{H}_{\text{ét}}^1(A, \lambda_{A,r}^n) & \longrightarrow & \mathbf{H}_{\text{ét}}^1(A, \lambda_{A,r+1}^n) & \longrightarrow & \mathbf{H}_{\text{ét}}^1(A, \lambda_{A,1}^n) \end{array} \quad (26)$$

is exact. Moreover the bottom row in the commutative diagram (26) is exact by Theorem 2.7. If the homomorphism (24) is an isomorphism in the case where $r \leq j$, then the homomorphism (24) is an isomorphism in the case where $r = j + 1$ by applying the snake lemma to the commutative diagram (26). Therefore the statement follows by induction on r . \square

3.3 Gersten's conjecture

PROPOSITION 3.6. Let A be a discrete valuation ring of mixed characteristic $(0, p)$ and X a semi-stable family over $\text{Spec}(A)$. Let R be the local ring at a point of X and Z the closed fiber of $\text{Spec}(R)$. Then we have a spectral sequence

$$E_1^{s,t} = \bigoplus_{x \in \text{Spec}(R)^{(s)} \cap Z} \mathbf{H}_x^{s+t}(R_{\text{ét}}, \mathfrak{T}_r(n)) \Rightarrow E^{s+t} = \mathbf{H}_Z^{s+t}(R_{\text{ét}}, \mathfrak{T}_r(n))$$

for any integer n .

Proof. For a bounded below complex F^\bullet of étale sheaf on $\text{Spec}(R)$, we have a spectral sequence

$$\bigoplus_{x \in \text{Spec}(R)} \mathbf{H}_x^{s+t}(R_{\text{ét}}, F^\bullet) \Rightarrow E^{s+t} = \mathbf{H}^{s+t}(R_{\text{ét}}, F^\bullet) \quad (27)$$

(cf.[6, Part 1, §1]). Let $i : Z \rightarrow \text{Spec}(R)$ be the inclusion of the closed fiber of $\text{Spec}(R)$ and $F^\bullet = i_* Ri^! \mathfrak{T}_r(n)$. Then we have

$$\mathbf{H}_x^j(R, F^\bullet) = \begin{cases} \mathbf{H}_x^j(R, \mathfrak{T}_r(n)) & (\text{if } x \in Z) \\ 0 & (\text{if } x \notin Z) \end{cases} \quad (28)$$

for any integer $j > 0$. Hence the statement follows from the spectral sequence (27) and (28). \square

LEMMA 3.7. Let n, N and i be positive integers. Suppose that $i = 0$ or $i = 1$. Consider a spectral sequence $E_1^{s,t} \Rightarrow E^{s+t}$ which satisfies the following conditions:

- (a) If $s < 0$ or $s > N$, then $E_1^{s,t} = 0$;
- (b) If $s + t = n$ or $s + t = n + 1$, then $E_2^{s,t} = 0$ for $s \neq i$.

Then we have

$$E^n = E_2^{i,n-i}.$$

Proof. By the definition of the spectral sequence, we have a filtration

$$E^n = E_0^n \supset E_1^n \supset \cdots \supset E_N^n \supset 0$$

which satisfies

$$E_p^n / E_{p+1}^n \simeq E_\infty^{p,n-p}$$

for any integer p . So we have

$$E_{i+1}^n = 0 \text{ and } E_i^n = E^n$$

by the conditions. Moreover, we have

$$E_2^{i,n-i} = E_\infty^{i,n-i}$$

by the conditions. Hence the statement follows. \square

THEOREM 3.8. Let A be a discrete valuation ring of mixed characteristic $(0, p)$, X a semi-stable family over $\text{Spec}(A)$ and $R = \mathcal{O}_{X,x}$ the local ring at a point x of X . Let Z be the closed fiber of $\text{Spec}(R)$. Then the sequence

$$0 \rightarrow H_Z^{n+2}(R_{\text{ét}}, \mathfrak{T}_r(n)) \rightarrow \bigoplus_{x \in Z^{(0)}} H_x^{n+2}(R_{\text{ét}}, \mathfrak{T}_r(n)) \rightarrow \bigoplus_{x \in Z^{(1)}} H_x^{n+3}(R_{\text{ét}}, \mathfrak{T}_r(n))$$

is exact for any integer $r > 0$.

Proof. By Lemma 3.7, the statement of (i) follows from Proposition 3.6, Corollary 2.4 and [22, p.520, Theorem 4.4.7]. \square

REMARK 3.9. Let A be a discrete valuation ring of mixed characteristic $(0, p)$ and R a local ring of a smooth scheme over $\text{Spec}(A)$. Let r be a positive integer. Then the sequence

$$\bigoplus_{x \in \text{Spec}(R)^{(0)}} H_x^s(R_{\text{ét}}, \mathbb{Z}/p^r(n)) \rightarrow \bigoplus_{x \in \text{Spec}(R)^{(1)}} H_x^{s+1}(R_{\text{ét}}, \mathbb{Z}/p^r(n)) \rightarrow \cdots$$

is exact for $s \leq n$ by [8, p.774, Theorem 1.2] and [24]. Hence we can also give another proof of [20, Theorem 1.1] by Lemma 3.7 and the spectral sequence (27).

4 Applications

Throughout this section, n is a non negative integer and r is a positive integer.

LEMMA 4.1. Let R be a discrete valuation ring of mixed characteristic $(0, p)$. Then the homomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}) \quad (29)$$

is injective.

Proof. Let \mathfrak{m} be the residue field of R . We have a commutative diagram

$$\begin{array}{ccc} K_n^M(k(R))/p^r & \longrightarrow & K_{n-1}^M(\kappa(\mathfrak{m}))/p^r \\ \downarrow & & \downarrow \\ H_{\text{ét}}^n(k(R), \mu_{p^r}^{\otimes n}) & \longrightarrow & H_{\text{ét}}^0(\kappa(\mathfrak{m}), \nu_{\kappa(\mathfrak{m}), r}^{n-1}) \end{array}$$

where $K_n^M(F)$ is the n -th Milnor K -group for a field F . Then the top horizontal arrow is induced by the boundary map (cf.[22, p.529, §3.1]) and so is surjective. Moreover the right vertical map is an isomorphism by [3, p.113, Theorem (2.1)]. Hence the homomorphism

$$H_{\text{ét}}^n(k(R), \mu_{p^r}^{\otimes n}) \rightarrow H_{\text{ét}}^0(\kappa(\mathfrak{m}), \nu_{\kappa(\mathfrak{m}), r}^{n-1})$$

is surjective and the homomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow H_{\text{ét}}^{n+1}(R, \tau_{\leq n} R j_* \mu_{p^r}^{\otimes n})$$

is injective where $j : \text{Spec}(k(R)) \rightarrow \text{Spec}(R)$ is the generic fiber. Since we have a distinguished triangle

$$\cdots \rightarrow \tau_{\leq n} R j_* \mu_{p^r}^{\otimes n} \rightarrow \tau_{\leq n+1} R j_* \mu_{p^r}^{\otimes n} \rightarrow R^{n+1} j_* \mu_{p^r}^{\otimes n}[-(n+1)] \rightarrow \cdots$$

and

$$H_{\text{ét}}^{n+1}(R, \tau_{\leq n+1} R j_* \mu_{p^r}^{\otimes n}) = H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}),$$

the homomorphism

$$H_{\text{ét}}^{n+1}(R, \tau_{\leq n} R j_* \mu_{p^r}^{\otimes n}) \rightarrow H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n})$$

is injective. Therefore the statement follows. \square

PROPOSITION 4.2. Let \mathfrak{X} be a semistable family over the spectrum of a discrete valuation ring of mixed characteristic $(0, p)$ and $R = \mathcal{O}_{\mathfrak{X}, x}^h$ be the henselian local ring of x in \mathfrak{X} . Then the homomorphism (29) is injective.

Proof. Let \mathfrak{m} be the maximal ideal of R and $a \in \mathfrak{m} \setminus \mathfrak{m}^2$ and $\mathfrak{p} = (a)$. Suppose that $p \in \mathfrak{p}$. Then R/\mathfrak{p} is a regular local ring and we have a commutative diagram

$$\begin{array}{ccc} H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) & \longrightarrow & H_{\text{ét}}^{n+1}(R/\mathfrak{p}, \mathfrak{T}_r(n)) \\ \downarrow & & \downarrow \\ H_{\text{ét}}^{n+1}(R/\mathfrak{p}, \nu_{R/\mathfrak{p}, r}^n) & \longrightarrow & H_{\text{ét}}^{n+1}(\kappa(\mathfrak{p}), \nu_{\kappa(\mathfrak{p}), r}^n). \end{array} \quad (30)$$

Moreover the left vertical arrow in (30) is an isomorphism by Theorem 3.5 and the bottom horizontal arrow in (30) is injective by Theorem 2.6. Hence the top horizontal arrow in (30) is injective. Therefore the statement follows from Lemma 4.1. \square

LEMMA 4.3. Let R be a local ring of a semistable family over a discrete valuation ring of mixed characteristic $(0, p)$. Suppose that $\dim(R) = 2$. Let \mathfrak{p} be a prime ideal such that $\text{ht}(\mathfrak{p}) = 1$, R/\mathfrak{p} is regular and $(p) \subset \mathfrak{p}$. Then the homomorphism

$$H_{R/\mathfrak{p}}^{n+2}(R, \mathfrak{T}_r(n)) \rightarrow H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n))$$

is injective.

Proof. Let \mathfrak{m} be the maximal ideal of R . Since the sequence

$$H_{\mathfrak{p}}^{n+1}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)) \rightarrow H_{R/\mathfrak{m}}^{n+2}(R, \mathfrak{T}_r(n)) \rightarrow H_{R/\mathfrak{p}}^{n+2}(R, \mathfrak{T}_r(n)) \rightarrow H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n))$$

is exact by [17, p.92, III, Remark 1.26], it suffices to show that the homomorphism

$$H_{\mathfrak{p}}^{n+1}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)) \rightarrow H_{R/\mathfrak{m}}^{n+2}(R, \mathfrak{T}_r(n)) \quad (31)$$

is surjective. The homomorphism (31) coincides with the homomorphism

$$H_{\text{ét}}^0(k(R/\mathfrak{p}), v_r^{n-1}) \rightarrow H_{\text{ét}}^0(R/\mathfrak{m}, v_r^{n-2}) \quad (32)$$

by [22, p.540, Theorem 4.4.7] and [23, p.187, Remark 3.7]. Let $K_n^M(F)$ be the n -th Milnor K -group for a field F . Then the homomorphism (32) coincides with the boundary map

$$K_{n-1}^M(k(R/\mathfrak{p}))/p^r \rightarrow K_{n-2}^M(R/\mathfrak{m})/p^r$$

of Milnor K -groups (modulo p^r) by [3, p.113, Theorem (2.1)] and so is surjective. This completes the proof. \square

LEMMA 4.4. Let B be a discrete valuation ring of mixed characteristic $(0, p)$. Let \mathfrak{X} be a semistable family over $\text{Spec}(B)$. Then we have

$$\tau_{\leq n}(\mathbb{Z}/p^r(n)_{\text{Nis}}) \simeq \tau_{\leq n}(R\alpha_*\mathbb{Z}/p^r(n)_{\text{ét}}),$$

where $\alpha : \mathfrak{X}_{\text{ét}} \rightarrow \mathfrak{X}_{\text{Nis}}$ is the change of site map.

Proof. Let $\varepsilon : \mathfrak{X}_{\text{ét}} \rightarrow \mathfrak{X}_{\text{Zar}}$ and $\beta : \mathfrak{X}_{\text{Nis}} \rightarrow \mathfrak{X}_{\text{Zar}}$ be the change of site map. Let $j : U \rightarrow \mathfrak{X}$ be the generic fiber and $i : Z \rightarrow \mathfrak{X}$ be the closed fiber. Then the Beilinson-Lichtenbaum Conjecture holds for U by [10, Theorem 1.1] and the Bloch-Kato conjecture ([24]). Moreover, the Beilinson-Lichtenbaum Conjecture holds for Z by Proposition 2.2.2. Therefore we have a quasi-isomorphism

$$\tau_{\leq n}(\mathbb{Z}/p^r(n)_{\text{Zar}}) \simeq \tau_{\leq n}(R\mathcal{E}_*\mathbb{Z}/p^r(n)_{\text{ét}}) \quad (33)$$

by [20, Proposition 2.1] and the same argument as in the proof of [8, Theorem 1.2.2]. Moreover we have quasi-isomorphisms

$$\beta^*R\beta_* \simeq \text{id} \quad \text{and} \quad R\mathcal{E}_* \simeq R\beta_*R\alpha_*.$$

Since β^* is exact and $\beta^*(\mathbb{Z}(n)_{\text{Zar}}) \simeq \mathbb{Z}(n)_{\text{Nis}}$, the statement follows from (33). \square

PROPOSITION 4.5. Let A be a discrete valuation ring of mixed characteristic $(0, p)$ and R a local ring of a semi-stable family over $\text{Spec}(A)$. Then we have

$$H_{\text{Zar}}^{n+1}(R, \mathbb{Z}/p^r(n)) = 0$$

for any integer $r > 0$ where $\mathbb{Z}(n)$ is the Bloch's cycle complex for the Zariski topology ([2], [16]) and $\mathbb{Z}/p^r(n) = \mathbb{Z}(n) \otimes \mathbb{Z}/p^r$.

Proof. Let Z be the closed fiber of $\text{Spec}(R)$ and U the generic fiber of $\text{Spec}(R)$. By Proposition 4.2, the homomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow H_{\text{ét}}^{n+1}(U, \mu_{p^r}^{\otimes n})$$

is injective. So the homomorphism

$$H_{\text{Zar}}^n(U, \mathbb{Z}/p^r(n)) \rightarrow H_{\text{Zar}}^{n-1}(Z, \mathbb{Z}/p^r(n-1))$$

is surjective by [22, p.540, Theorem 4.4.7], [8, p.774, Theorem 1.2.2] and [24]. Hence the homomorphism

$$H_{\text{Zar}}^{n+1}(R, \mathbb{Z}/p^r(n)) \rightarrow H_{\text{Zar}}^{n+1}(U, \mathbb{Z}/p^r(n)) \quad (34)$$

is injective. Moreover, we have the commutative diagram

$$\begin{array}{ccccc} H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) & \longrightarrow & H_{\text{ét}}^{n+1}(U, \mu_{p^r}^{\otimes n}) & \longrightarrow & H_Z^{n+2}(R_{\text{ét}}, \mathfrak{T}_r(n)) & (35) \\ \downarrow & & \downarrow & & \downarrow & \\ 0 \longrightarrow \bigoplus_{x \in Z^{(0)}} H_{\text{ét}}^{n+1}(R_x, \mathfrak{T}_r(n)) & \longrightarrow & \bigoplus_{x \in Z^{(0)}} H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}) & \longrightarrow & \bigoplus_{x \in Z^{(0)}} H_x^{n+2}(R_{\text{ét}}, \mathfrak{T}_r(n)) \end{array}$$

where the sequences are exact. Then the left map in the diagram (35) is injective by Proposition 4.2 and the right map in the diagram (35) is injective by Theorem 3.8. So the middle map in the diagram (35) is injective and the homomorphism

$$H_{\text{ét}}^{n+1}(U, \mathbb{Z}/p^r(n)) \rightarrow H_{\text{ét}}^{n+1}(k(R), \mathbb{Z}/p^r(n))$$

is injective by [8, p.774, Theorem 1.2.4] and [24]. Moreover, the homomorphism

$$H_{\text{Zar}}^{n+1}(U, \mathbb{Z}/p^r(n)) \rightarrow H_{\text{ét}}^{n+1}(U, \mathbb{Z}/p^r(n))$$

is injective by [8, p.774, Theorem 1.2.2] and [24]. Hence the homomorphism

$$H_{\text{Zar}}^{n+1}(U, \mathbb{Z}/p^r(n)) \rightarrow H_{\text{Zar}}^{n+1}(k(R), \mathbb{Z}/p^r(n))$$

is injective and we have

$$H_{\text{Zar}}^{n+1}(U, \mathbb{Z}/p^r(n)) = 0 \quad (36)$$

by [8, p.786, Corollary 4.4]. Therefore the statement follows from the equation (36) and the injectivity of the homomorphism (34). \square

The following means that [22, p.524, Conjecture 1.4.1 (1)] holds in the case where \mathfrak{X}/B is semi-stable and $\dim(\mathfrak{X}) = 2$.

COROLLARY 4.6. Let \mathfrak{X} be a semistable family over a discrete valuation ring of mixed characteristic $(0, p)$. Suppose that $\dim(\mathfrak{X}) = 2$. Then we have an isomorphism in $D^b(\mathfrak{X}_{\text{ét}}, \mathbb{Z}/p^r\mathbb{Z})$

$$\mathbb{Z}/p^r(n)_{\text{ét}} \simeq \mathfrak{T}_r(n),$$

where $\mathbb{Z}/p^r(n)_{\text{ét}}$ is the étale sheafification of $\mathbb{Z}/p^r(n)$ and $D^b(\mathfrak{X}_{\text{ét}}, \mathbb{Z}/p^r\mathbb{Z})$ is the derived category of bounded complexes of étale $\mathbb{Z}/p^r\mathbb{Z}$ -sheaves on \mathfrak{X} .

Proof. We have a quasi-isomorphism

$$\tau_{\leq n}(\mathbb{Z}/p^r(n)_{\text{ét}}) \simeq \tau_{\leq n}(\mathfrak{T}_r(n))$$

by [23, p.209, Remark 7.2]. By the definition of $\mathfrak{T}_r(n)$, we have

$$\mathcal{H}^i(\mathfrak{T}_r(n)) = 0$$

for $i \geq n + 1$. Let $\alpha : \mathfrak{X}_{\text{ét}} \rightarrow \mathfrak{X}_{\text{Nis}}$ be the canonical map of sites. Then the sheafification α^* is exact and

$$\alpha^*\mathbb{Z}/p^r(n)_{\text{Nis}} = \mathbb{Z}/p^r(n)_{\text{ét}}.$$

Hence it suffices to show that

$$\mathcal{H}^i(\mathbb{Z}/p^r(n)) = 0$$

for $l \geq n + 1$. Since we have

$$\begin{aligned} \Gamma\left(\text{Spec}(\mathcal{O}_{\mathfrak{X},x}^h), \mathcal{H}^{n+1}(\mathbb{Z}/p^r(n)_{\text{Nis}})\right) &\simeq \mathbf{H}^{n+1}\left(\mathbb{Z}/p^r(n)_{\text{Zar}}(\text{Spec}(\mathcal{O}_{\mathfrak{X},x}^h))\right) \\ &\simeq \mathbf{H}_{\text{Zar}}^{n+1}(\mathcal{O}_{\mathfrak{X},x}^h, \mathbb{Z}/p^r(n)) \end{aligned}$$

by [8, p.779, Theorem 3.2 b)], it suffices to show that

$$\mathbf{H}_{\text{Zar}}^i(\mathcal{O}_{\mathfrak{X},x}^h, \mathbb{Z}/p^r(n)) = 0$$

for $i \geq n + 1$. Here $\mathcal{O}_{\mathfrak{X},x}^h$ is the henselian local ring at a point x of \mathfrak{X} .

Let $Z \hookrightarrow \mathfrak{X}$ be a closed immersion of codimension 1 with $U = \mathfrak{X} \setminus Z$. Suppose that Z is regular. Then we have

$$\mathbf{H}_{\text{Zar}}^i(Z, \mathbb{Z}/p^r(n-1)) = 0$$

for $i > n$ by [8, p.786, Corollary 4.4]. Since $\dim(U) \leq 1$, we have

$$\mathbf{H}^i(\mathbb{Z}(n)_{\text{Zar}}(U)) = 0$$

for $i > n + 1$ by the definition of $\mathbb{Z}(n)_{\text{Zar}}$. Moreover we have

$$\mathbf{H}_{\text{Zar}}^i(U, \mathbb{Z}(n)) = \mathbf{H}^i(\mathbb{Z}(n)_{\text{Zar}}(U)) = 0$$

for $i > n + 1$ by [8, p.779, Theorem 3.2 b)]. Hence we have

$$H_{\text{Zar}}^i(U, \mathbb{Z}/p^r(n)) = 0$$

for $i > n + 1$. Therefore we have

$$H_{\text{Zar}}^i(R, \mathbb{Z}/p^r(n)) = 0$$

for $i > n + 1$ by the localization theorem ([8, p.779, Theorem 3.2]). Moreover we have

$$H_{\text{Zar}}^{n+1}(R, \mathbb{Z}/p^r(n)) = 0$$

by Proposition 4.5. This completes the proof. \square

REMARK 4.7. Let \mathfrak{X} be a regular scheme which is finite type over a Dedekind domain. Let R be the henselization of a local ring $\mathcal{O}_{\mathfrak{X},x}$ and l a positive integer which is invertible in R . Then we can prove

$$H_{\text{Zar}}^{n+1}(R, \mathbb{Z}/l(n)) = 0$$

by the same argument as Proposition 4.5. Hence we have

$$\mathcal{H}^{n+1}(\mathbb{Z}/l(n)_{\text{ét}}) = 0$$

by [8, p.779, Theorem 3.2.b)]. Let $i : Z \hookrightarrow \text{Spec}(R)$ be a close immersion of regular subschemes of pure codimension 1. Then we have an isomorphism

$$Ri^! \mu_l^{\otimes n} \simeq \mu_l^{\otimes(n-1)}[-2]$$

in $D^b(Z_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$ by the absolute purity theorem ([7]). Here $D^b(Z_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$ is the derived category of bounded complexes of étale $\mathbb{Z}/l\mathbb{Z}$ -sheaves on Z . On the other hand, we have an isomorphism

$$\tau_{\leq n+1}(Ri^! \mathbb{Z}/l(n)_{\text{ét}}) \simeq \tau_{\leq n+1}(\mathbb{Z}/l(n-1)_{\text{ét}}[-2])$$

in $D^b(Z_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$ by [20, Proposition 2.1] and [24]. Hence we have an isomorphism

$$\tau_{\leq n}(\mathbb{Z}/l(n)_{\text{ét}}) \simeq \mu_l^{\otimes n}$$

in $D^b(\text{Spec}(R)_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$ by induction on $\dim(R)$. Therefore we have an isomorphism

$$\tau_{\leq n+1}(\mathbb{Z}/l(n)_{\text{ét}}) \simeq \mu_l^{\otimes n}$$

in $D^b(\mathfrak{X}_{\text{ét}}, \mathbb{Z}/l\mathbb{Z})$.

THEOREM 4.8. Let \mathfrak{X} be a semistable family over a discrete valuation ring of mixed characteristic $(0, p)$ and $R = \tilde{\mathcal{O}}_{\mathfrak{X},x}$ the henselian local ring of x in \mathfrak{X} . Suppose that $\dim(R) = 2$. Then the sequence

$$0 \rightarrow H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}) \xrightarrow{(*)} \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1} H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n))$$

is exact.

Proof. Let \mathfrak{m} be the maximal ideal of R . Let $a \in \mathfrak{m} \setminus \mathfrak{m}^2$. Let U be $\text{Spec}(R) \setminus \text{Spec}(R/(a))$. Suppose that $p \in (a)$. Since R/\mathfrak{m} has p -cohomological dimension at most one and

$$\mathcal{H}^i(\mathfrak{T}_r(n)) = 0$$

for $i \geq n+1$ by the definition of $\mathfrak{T}_r(n)$, we have

$$H_{\text{ét}}^{n+2}(R, \mathfrak{T}_r(n)) = 0$$

by the spectral sequence

$$H_{\text{ét}}^i(R, \mathcal{H}^j(\mathfrak{T}_r(n))) \Rightarrow H_{\text{ét}}^{i+j}(R, \mathfrak{T}_r(n)).$$

Hence we have a commutative diagram

$$\begin{array}{ccccccc} H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) & \longrightarrow & H_{\text{ét}}^{n+1}(U, \mathfrak{T}_r(n)) & \longrightarrow & H_{R/\mathfrak{p}}^{n+2}(R, \mathfrak{T}_r(n)) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \text{Ker}((*)) & \longrightarrow & H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n})' & \longrightarrow & H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)) \end{array} \quad (37)$$

where

$$H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n})' = \text{Ker} \left(H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}) \rightarrow \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1} H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)) \right)$$

and the rows are exact. Then the middle vertical arrow is surjective by [8, p.778, Lemma 2.4]. Moreover the left vertical arrow is injective by Lemma 4.3. Therefore the statement follows from the snake lemma. \square

COROLLARY 4.9. Let R be the same as in Theorem 4.8 and Y the special fiber of $\text{Spec}(R)$. Then the natural map

$$H_{\text{ét}}^{n+1}(k(R), \mu_{p^r}^{\otimes n}) \rightarrow \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1 \setminus Y^0} H_{\text{ét}}^n(\kappa(\mathfrak{p}), \mu_{p^r}^{\otimes(n-1)}) \oplus \bigoplus_{\mathfrak{p} \in Y^0} H_{\text{ét}}^{n+1}(k(\tilde{R}_{\mathfrak{p}}), \mu_{p^r}^{\otimes n})$$

is injective for non negative integer n . Here $\tilde{R}_{\mathfrak{p}}$ is the henselization of a local ring $R_{\mathfrak{p}}$.

Proof. Put

$$\text{RH}^{n+1}(R) := \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1 \setminus Y^0} H_{\text{ét}}^n(\kappa(\mathfrak{p}), \mu_{p^r}^{\otimes(n-1)}) \oplus \bigoplus_{\mathfrak{p} \in Y^0} H_{\text{ét}}^{n+1}(k(\tilde{R}_{\mathfrak{p}}), \mu_{p^r}^{\otimes n}).$$

Then we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) & \longrightarrow & H_{\text{ét}}^{n+1}(k(R), \mathfrak{T}_r(n)) & \longrightarrow & \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1} H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)) \\ \downarrow & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \bigoplus_{\mathfrak{p} \in Y^0} H_{\text{ét}}^{n+1}(\tilde{R}_{\mathfrak{p}}, \mathfrak{T}_r(n)) & \longrightarrow & \text{RH}^{n+1}(R) & \longrightarrow & \bigoplus_{\mathfrak{p} \in \text{Spec}(R)^1} H_{\mathfrak{p}}^{n+2}(R_{\mathfrak{p}}, \mathfrak{T}_r(n)), \end{array}$$

where the rows are exact by Theorem 4.8 and the absolute purity theorem (cf.[17, p.241, VI, Theorem 5.1]). Hence it suffices to show that the homomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow \bigoplus_{\mathfrak{p} \in Y^0} H_{\text{ét}}^{n+1}(\tilde{R}_{\mathfrak{p}}, \mathfrak{T}_r(n))$$

is injective. Since a diagram

$$\begin{array}{ccc} H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) & \longrightarrow & H_{\text{ét}}^{n+1}(\tilde{R}_{\mathfrak{p}}, \mathfrak{T}_r(n)) \\ \downarrow & & \downarrow \\ H_{\text{ét}}^1(R/\mathfrak{p}, \mathfrak{v}_r^n) & \longrightarrow & H_{\text{ét}}^1(\kappa(\mathfrak{p}), \mathfrak{v}_r^n) \end{array}$$

is commutative, the homomorphism

$$H_{\text{ét}}^{n+1}(R, \mathfrak{T}_r(n)) \rightarrow H_{\text{ét}}^{n+1}(\tilde{R}_{\mathfrak{p}}, \mathfrak{T}_r(n))$$

is injective by Theorem 3.5 and Theorem 2.6. This completes the proof. \square

LEMMA 4.10. Let v_i ($1 \leq i \leq m$) be a finite collection of independent discrete valuations on a field K of characteristic 0. Let K_i be the henselization of K at v_i and $\kappa(v_i)$ the residue field of v_i for each i . Suppose that $\text{char}(\kappa(v_i)) = p > 0$ for all i . Then the natural map

$$H_{\text{ét}}^n(K, \mu_{p^r}^{\otimes n}) \rightarrow \bigoplus_i H_{\text{ét}}^n(K_i, \mu_{p^r}^{\otimes n})$$

is surjective for $n \geq 1$.

Proof. The statement follows from [3, p.131, Theorem (5.12)] and [20, Lemma 6.2]. \square

THEOREM 4.11. Let \mathfrak{X} be a proper and semistable family over a henselian discrete valuation ring of mixed characteristic $(0, p)$ and Y the special fiber of \mathfrak{X} . Suppose that $\dim(\mathfrak{X}) = 2$. Then the map

$$H_{\text{ét}}^{n+1}(k(\mathfrak{X}), \mu_{p^r}^{\otimes n}) \rightarrow \bigoplus_{\mathfrak{p} \in \mathfrak{X}^1 \setminus Y^0} H_{\text{ét}}^n(\kappa(\mathfrak{p}), \mu_{p^r}^{\otimes(n-1)}) \oplus \bigoplus_{\mathfrak{p} \in Y^0} H_{\text{ét}}^{n+1}(k(\mathcal{O}_{\tilde{\mathfrak{X}, \mathfrak{p}}}), \mu_{p^r}^{\otimes n})$$

is injective for $n \geq 1$. Here $\mathcal{O}_{\tilde{\mathfrak{X}, \mathfrak{p}}}$ is the henselization of a local ring $\mathcal{O}_{\mathfrak{X}, \mathfrak{p}}$. This implies that the local-global map

$$H_{\text{ét}}^{n+1}(k(\mathfrak{X}), \mu_{p^r}^{\otimes n}) \rightarrow \prod_{\mathfrak{p} \in \mathfrak{X}^1} H_{\text{ét}}^{n+1}(k(\mathcal{O}_{\tilde{\mathfrak{X}, \mathfrak{p}}}), \mu_{p^r}^{\otimes n})$$

is injective for $n \geq 1$.

Proof. The proof of the statement is the same as [13, Theorem 2.5]. The statement follows from Corollary 4.9 and Lemma 4.10. We review the proof of [13, Theorem 2.5] for convenience.

Let

$$\begin{array}{ccc} X_1 & \xleftarrow{j'_2} & X_3 \\ j_1 \downarrow & & \downarrow j'_1 \\ X_0 & \xleftarrow{j_2} & X_2 \end{array}$$

be Cartesian. If j_1 and j_2 are étale and a complex $\mathcal{U} \in D_+(X_1)$, then we have an isomorphism

$$j_2^* Rj_{1*}(\mathcal{U}) \simeq Rj'_1{}^* j_2'^*(\mathcal{U}) \quad (38)$$

in $D_+(X_2)$ by [17, p.88, III, Theorem 1.15].

Write $\Lambda = \mu_{p^r}^{\otimes n}$ and $U := \mathfrak{X} \setminus Y$. The natural inclusion $j : U \hookrightarrow \mathfrak{X}$ is the complement of the closed immersion $i : Z \rightarrow \mathfrak{X}$. For each $x \in U$, $\{\bar{x}\}$ denotes its closure in \mathfrak{X} . Then we have an isomorphism

$$H_x^s(U, \Lambda) \simeq H_{\{\bar{x}\}}^s(\mathfrak{X}, Rj_*\Lambda)$$

for any integer s by (38). Thus the sequence

$$\bigoplus_{x \in U^1} H_x^{n+1}(U, \Lambda) \rightarrow H^{n+1}(U, \Lambda) \rightarrow H^{n+1}(k(U), \Lambda) \rightarrow \bigoplus_{x \in U^1} H_x^{n+2}(U, \Lambda)$$

corresponds to the sequence

$$\bigoplus_{x \in U^1} H_{\{\bar{x}\}}^{n+1}(\mathfrak{X}, Rj_*\Lambda) \rightarrow H^{n+1}(\mathfrak{X}, Rj_*\Lambda) \rightarrow H^{n+1}(k(\mathfrak{X}), \Lambda) \rightarrow \bigoplus_{x \in U^1} H_{\{\bar{x}\}}^{n+2}(\mathfrak{X}, Rj_*\Lambda).$$

Put $\mathcal{F} = i^* Rj_*\Lambda$. By the proper base change theorem ([17, pp.223–224, VI, Corollary 2.3]), we have an isomorphism

$$H_{\{\bar{x}\}}^s(\mathfrak{X}, i_*\mathcal{F}) \simeq H_{\{\bar{x}\} \cap Y}^s(Y, \mathcal{F})$$

for any integer s and a commutative diagram with exact rows

$$\begin{array}{ccccccc} \bigoplus_{x \in U^1} H_{\{\bar{x}\}}^{n+1}(\mathfrak{X}, Rj_*\Lambda) & \longrightarrow & H^{n+1}(\mathfrak{X}, Rj_*\Lambda) & \longrightarrow & H^{n+1}(k(\mathfrak{X}), \Lambda) & \longrightarrow & \bigoplus_{x \in U^1} H_{\{\bar{x}\}}^{n+2}(\mathfrak{X}, Rj_*\Lambda) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \bigoplus_{y \in Y^1} H_y^{n+1}(Y, \mathcal{F}) & \longrightarrow & H^{n+1}(Y, \mathcal{F}) & \longrightarrow & \bigoplus_{\eta \in Y^0} H^{n+1}(\kappa(\eta), \mathcal{F}) & \longrightarrow & \bigoplus_{y \in Y^1} H_y^{n+2}(Y, \mathcal{F}). \end{array}$$

By the proper base change theorem, (38) and [8, p.777, The proof of Proposition 2.2.b)], we have an isomorphism

$$H^{n+1}(\kappa(\eta), \mathcal{F}) \simeq H^{n+1}(K_{(\eta)}, \Lambda)$$

where $K_{(\eta)}$ is the fraction field of the henselian local ring $\mathcal{O}_{\mathfrak{X}, \eta}^{\sim}$. Moreover, we have an isomorphism

$$H^{s-2}(\kappa(x), \Lambda(-1)) \simeq H_x^s(U, \Lambda)$$

for any integer s by the absolute purity theorem ([17, p.241, VI, Theorem 5.1]). Hence we have a commutative diagram with exact rows

$$\begin{array}{ccccccc}
\bigoplus_{x \in U^1} \mathrm{H}^{n-1}(\kappa(x), \Lambda(-1)) & \longrightarrow & \mathrm{H}^{n+1}(\mathfrak{X}, R_{j_*} \Lambda) & \longrightarrow & \mathrm{H}^{n+1}(k(\mathfrak{X}), \Lambda)' & \longrightarrow & 0 \\
\downarrow \gamma & & \downarrow \simeq & & \downarrow & & \\
\bigoplus_{y \in Y^1} \mathrm{H}_y^{n+1}(Y, \mathcal{F}) & \longrightarrow & \mathrm{H}^{n+1}(Y, \mathcal{F}) & \longrightarrow & \bigoplus_{\eta \in Y^0} \mathrm{H}^{n+1}(K(\eta), \Lambda) & &
\end{array} \quad (39)$$

where

$$\mathrm{H}^{n+1}(k(\mathfrak{X}), \Lambda)' = \mathrm{Ker} \left(\mathrm{H}^{n+1}(k(\mathfrak{X}), \Lambda) \rightarrow \bigoplus_{x \in U^1} \mathrm{H}^n(\kappa(x), \Lambda(-1)) \right).$$

The statement means that the right map in the diagram (39) is injective. By the proper base change theorem, the middle map in the diagram (39) is an isomorphism. So it suffices to show that the homomorphism

$$\gamma: \bigoplus_{x \in U^1} \mathrm{H}^{n-1}(\kappa(x), \Lambda(-1)) \rightarrow \bigoplus_{y \in Y^1} \mathrm{H}_y^{n+1}(Y, \mathcal{F})$$

is surjective.

Let $x \in U^1$. Since $\{\bar{x}\} \rightarrow \mathfrak{X} \rightarrow \mathrm{Spec}(A)$ is proper and quasi-finite, so is finite by [17, p.6, I, Corollary 1.10]. Hence $\{\bar{x}\}$ is the spectrum of the henselian local domain $\mathcal{O}_{\bar{x}, y}/x$ by [17, pp.32–33, I, Theorem 4.2] and [17, p.34, I, Corollary 4.3]. Therefore $\{\bar{x}\}$ meets Y at one and only one point and

$$U^1 = \bigsqcup_{y \in Y^1} (\mathrm{Spec}(\mathcal{O}_{\bar{x}, y}) \times_{\mathfrak{X}} U)^1.$$

Moreover we have isomorphisms

$$\mathcal{O}_{\bar{x}, y}/x \cdot \tilde{\mathcal{O}}_{\bar{x}, y} \simeq \widetilde{\mathcal{O}_{\bar{x}, y}/x} \simeq \mathcal{O}_{\bar{x}, y}/x$$

and $x \cdot \tilde{\mathcal{O}}_{\bar{x}, y}$ is a prime ideal of $\tilde{\mathcal{O}}_{\bar{x}, y}$. So we have a bijection

$$P_y := (\mathrm{Spec}(\tilde{\mathcal{O}}_{\bar{x}, y}) \times_{\mathfrak{X}} U)^1 \rightarrow (\mathrm{Spec}(\mathcal{O}_{\bar{x}, y}) \times_{\mathfrak{X}} U)^1.$$

Hence the map γ decomposes into a direct sum

$$\gamma = \bigoplus_{y \in Y^1} \left(\gamma_y: \bigoplus_{x \in P_y} \mathrm{H}^{n-1}(\kappa(x), \Lambda(-1)) \rightarrow \mathrm{H}_y^{n+1}(Y, \mathcal{F}) \right)$$

and it suffices to show that the map γ_y is surjective.

Put

$$V_y = \mathrm{Spec}(\tilde{\mathcal{O}}_{\bar{x}, y})^1 \setminus (\mathrm{Spec}(\mathcal{O}_{\bar{x}, y}) \times_{\mathfrak{X}} U)^1.$$

Let $(\mathcal{O}_{\tilde{x},y})_\eta$ be the localization of the henselian local ring $\mathcal{O}_{\tilde{x},y}$ at $\eta \in V_y$ and $K_{(y,\eta)}$ the fraction field of the henselization of $(\mathcal{O}_{\tilde{x},y})_\eta$. Since

$$\mathcal{O}_{Y,y} = \mathcal{O}_{\tilde{x},y} \otimes_{\mathcal{O}_{x,y}} \mathcal{O}_{Y,y}$$

by [17, p.37, I, Examples 4.10 (c)], we have a commutative diagram with exact rows

$$\begin{array}{ccccccc} \mathrm{H}^n(K_{(y)}, \Lambda) & \longrightarrow & \bigoplus_{x \in P_y} \mathrm{H}^{n-1}(\kappa(x), \Lambda(-1)) & \longrightarrow & \mathrm{H}^{n+1}(\mathcal{O}_{\tilde{x},y} \times_X U, \Lambda) & \longrightarrow & \mathrm{H}^{n+1}(K_{(y)}, \Lambda)' & (40) \\ \downarrow & & \downarrow \gamma & & \downarrow \simeq \varphi & & \downarrow & \\ \bigoplus_{\eta \in V_y} \mathrm{H}^n(K_{(y,\eta)}, \Lambda) & \longrightarrow & \mathrm{H}_y^{n+1}(Y, \mathcal{F}) & \longrightarrow & \mathrm{H}^{n+1}(\mathcal{O}_{Y,y}, \mathcal{F}) & \longrightarrow & \bigoplus_{\eta \in V_y} \mathrm{H}^{n+1}(K_{(y,\eta)}, \Lambda) \end{array}$$

where

$$\mathrm{H}^{n+1}(K_{(y)}, \Lambda)' = \mathrm{Ker} \left(\mathrm{H}^{n+1}(K_{(y)}, \Lambda) \rightarrow \bigoplus_{x \in P_y} \mathrm{H}^n(\kappa(x), \Lambda) \right).$$

In the diagram (40), the left vertical map is surjective by Lemma 4.10, the map φ is an isomorphism by the proper base change theorem and the right vertical map is injective by Corollary 4.9. Therefore the map γ_y is surjective by chasing the diagram. This completes the proof. \square

REMARK 4.12. In the proof of Theorem 4.11, we use the Bloch-Kato conjecture ([24]). But it suffices to use [3, p.131, Theorem (5.12)] which is a special case of the Bloch-Kato conjecture (cf. The proof of Lemma 4.10).

References

- [1] ARTIN, M., A. GROTHENDIECK, AND J.-L. VERDIER., Théorie des Topos et Cohomologie Étale des Schémas (SGA 4), Tome 3. Lecture Notes in Mathematics, vol. 305. Heidelberg, Berlin: Springer-Verlag, 1973.
- [2] S.BLOCH, Algebraic cycles and Algebraic K-theory, Adv. in Math. 61 (1986), no. 3, 267–304.
- [3] S.BLOCH AND K.KATO, p -adic étale cohomology. Inst. Hautes Études Sci. Publ. Math. No. 63 (1986), 107–152.
- [4] S.BLOCH AND A.OGUS, Gersten’s conjecture and the homology of schemes. Ann. Sci. École Norm. Sup. (4) 7 (1974), 181–201 (1975).
- [5] J.-L.COLLIOT-THÉLÈNE, Quelques problèmes locaux-globaux, Personal Notes. (2011).
- [6] COLLIOT-THÉLÈNE, JEAN-LOUIS; HOOBLER, RAYMOND T.; KAHN, BRUNO, The Bloch-Ogus-Gabber theorem, Snaith, Victor P. (ed.), Algebraic K-theory. Papers from the 2nd Great Lakes conference, Canada, March 1996, in memory of Robert Wayne Thomason. Providence, RI: American Mathematical Society. Fields Inst. Commun. 16, 31–94 (1997).

- [7] K.FUJIWARA, A proof of the absolute purity conjecture (after Gabber), Algebraic geometry 2000, Azumino (Hotaka), 153–183, Adv. Stud. Pure Math., 36, Math. Soc. Japan, Tokyo, 2002.
- [8] T.GEISSER, Motivic cohomology over Dedekind rings, Math. Z. 248 (2004), no. 4, 773–794.
- [9] T.GEISSER AND M.LEVINE, The K -Theory of fields in characteristic p , Invent. Math. 139 (2000), no. 3, 459–493.
- [10] T.GEISSER AND M.LEVINE, The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky, J. Reine Angew. Math. 530 (2001), 55–103.
- [11] M.GROS AND N.SUWA, La conjecture de Gersten pour les faisceaux de Hodge-Witt logarithmique, Duke Math. J. 57 (1988), no. 2, 615–628.
- [12] D.HARBATER AND J.HARTMANN AND D.KRASHEN, Local-global principles for Galois cohomology, Comment. Math. Helv.89 (2014), no.1, 215–253.
- [13] Y.HU, A Cohomological Hasse Principle Over Two-dimensional Local Rings, Int. Math. Res. Not. IMRN 2017, no. 14, 4369–4397.
- [14] O.HYODO, A note on p -adic étale cohomology in the semistable reduction case, Invent. Math. 91 (1988), no.3, 543–557.
- [15] L.ILLUSIE, Complexe de de Rham-Witt et cohomologie cristalline, Ann. Sci. École Norm. Sup. (4) 12 (1979), no. 4, 501–661.
- [16] M.LEVINE, Techniques of localization in the theory of algebraic cycles, J. Algebraic Geom. 10 (2001), no.2, 299–363.
- [17] J.S.MILNE, Etale Cohomology, Princeton Math. Ser. Princeton University Press, Princeton, N.J., 1980.
- [18] D.QUILLEN, Higher algebraic K -theory: I. Higher K -theories (Proc.Conf., Battelle Memorial Inst., Seattle, Wash., 1972), pp. 85–147. Lecture Notes in Math., Vol. 341, Springer, 1973.
- [19] M.SAKAGAITO, A note on Gersten’s conjecture for étale cohomology over two-dimensional henselian regular local rings, Comptes Rendus - Mathématique 358 (2020), no.1, 33–39.
- [20] M.SAKAGAITO, On a generalized Brauer group in mixed characteristic cases, J. Math. Sci. Univ. Tokyo 27, No.1, 29–64 (2020).
- [21] K.SATO, Logarithmic Hodge–Witt sheaves on normal crossing varieties, Math. Z. 257 (2007), no. 4, 707–743.
- [22] K.SATO, p -adic étale Tate twists and arithmetic duality. With an appendix by Kei Hagihara, Ann. Sci. École Norm. Sup. (4) 40 (2007), no.4, 519–588.

- [23] K.SATO, Cycle classes for p -adic étale Tate twists and the image of p -adic regulators. Doc. Math. 18 (2013), 177–247.
- [24] V.VOEVOFSKY, On motivic cohomology with \mathbf{Z}/l -coefficients, Ann. of Math. (2) 174 (2011), no. 1, 401–438.