

A HIGHER RANK EULER SYSTEM FOR THE MULTIPLICATIVE GROUP OVER A TOTALLY REAL FIELD

RYOTARO SAKAMOTO

ABSTRACT. In this paper, we construct a higher rank Euler system for the multiplicative group over a totally real field by using the Iwasawa main conjecture proved by Wiles. A key ingredient of the construction is to generalize the notion of the characteristic ideal. Under certain technical assumptions, we prove that all higher Fitting ideals of a certain p -ramified Iwasawa module are described by analytic invariants canonically associated with Stickelberger elements.

CONTENTS

1. Introduction	1
1.1. Notation	4
1.2. Acknowledgments	5
2. Higher rank Euler systems	5
3. Higher rank Euler systems over totally real fields	9
3.1. Stickelberger elements and Iwasawa main conjecture	10
3.2. Iwasawa modules	12
3.3. Higher rank Euler systems for the multiplicative group	14
4. Stark systems	15
4.1. Review of the theory of Stark systems	15
4.2. Kolyvagin derivative homomorphism	20
4.3. Regulator homomorphism	22
5. On the Fitting ideals of X_K^χ	26
5.1. Proof of Theorem 5.3	27
Appendix A. Notes on Galois cohomology	31
A.1. Remarks on continuous cochain complexes	31
A.2. Selmer structures	32
Appendix B. Exterior bi-dual	33
B.1. Definition of the exterior bi-dual	34
B.2. Basic properties	34
B.3. Base change	36
Appendix C. Characteristic ideal	39
References	44

1. INTRODUCTION

Euler systems were introduced by Kolyvagin in order to study the structure of Selmer groups of elliptic curves. The theory of Euler systems has been playing a crucial role in studying the arithmetic of global Galois representations. However, there are very few known examples of Euler systems (see [18, 22, 23, 24, 32]).

In the paper [29], Perrin-Riou introduced the notion of higher rank Euler systems. However, at the present moment, we know an example of a higher rank Euler system only in the $\mathrm{GL}_2 \times \mathrm{GL}_2$ -case (see [9, 10, 11]). In this paper, we construct a higher rank Euler system for the multiplicative group over a totally real field by using the classical Iwasawa main conjecture for totally real fields proved by Wiles in [36].

Remark 1.1. The definition of higher rank Euler systems in this paper is not the same as the one introduced by Perrin-Riou in [29]. Perrin-Riou used the exterior power in order to define higher rank Euler systems, while we use the exterior bi-dual in this paper. Our definition of higher rank Euler systems is a generalization of Perrin-Riou's one and these definitions are essentially the same in the rank 1 case (see [1, 3, 34]).

To explain the result we obtain in this article more precisely, we introduce some notations. Let k be a totally real field with degree $r := [k: \mathbb{Q}]$ and

$$\chi: G_k \longrightarrow \overline{\mathbb{Q}}^\times$$

a non-trivial finite order even character. Here, for a field k' , we write $G_{k'}$ for the absolute Galois group of k' . Put $L := \overline{k}^{\ker(\chi)}$. Let p be an odd prime such that both the order of χ and the class number of k are coprime to p . Fix a finite abelian p -extension K/k satisfying $S_{\mathrm{ram}}(K/k) \cap (S_p(k) \cup S_{\mathrm{ram}}(L/k)) = \emptyset$. Here $S_p(k)$ denotes the set of all primes of k above p and, for an algebraic extension K'/k , we write $S_{\mathrm{ram}}(K'/k)$ for the set of all primes of k at which K'/k is ramified. Fix an embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ and put

$$\mathcal{O} := \mathbb{Z}_p[\mathrm{im}(\chi)] \quad \text{and} \quad T := \mathcal{O}(1) \otimes \chi^{-1}.$$

We write $M_{KL, \infty}$ for the maximal p -ramified pro- p abelian extension of $KL(\mu_{p^\infty})$. We note that $M_{KL, \infty}/k$ is a Galois extension. Since $M_{KL, \infty}/KL(\mu_{p^\infty})$ is an abelian extension, the Galois group $\mathrm{Gal}(KL(\mu_{p^\infty})/k)$ acts on the Galois group $\mathrm{Gal}(M_{KL, \infty}/KL(\mu_{p^\infty}))$. Furthermore, by using this Galois action, we see that the module

$$X_K := \mathcal{O} \otimes_{\mathbb{Z}_p} \mathrm{Gal}(M_{KL, \infty}/KL(\mu_{p^\infty}))$$

has an $\mathcal{O}[[\mathrm{Gal}(KL(\mu_{p^\infty})/k)]]$ -module structure. Let k_∞/k denote the cyclotomic \mathbb{Z}_p -extension. Then we have the canonical isomorphism

$$\mathrm{Gal}(KL(\mu_{p^\infty})/k) \xrightarrow{\sim} \mathrm{Gal}(k_\infty K/k) \times \mathrm{Gal}(L(\mu_p)/k)$$

since $p \nmid [L: k]$. Set

$$e_\chi := \frac{1}{[L(\mu_p): k]} \sum_{\sigma \in \mathrm{Gal}(L(\mu_p)/k)} \chi(\sigma)\sigma^{-1} = \frac{1}{[KL(\mu_{p^\infty}): k_\infty K]} \sum_{\sigma \in \mathrm{Gal}(KL(\mu_{p^\infty})/k_\infty K)} \chi(\sigma)\sigma^{-1}.$$

By the above isomorphism, we see that the Iwasawa module

$$X_K^\chi := e_\chi X_K$$

is a $\Lambda_K := \mathcal{O}[[\mathrm{Gal}(k_\infty K/k)]]$ -module.

Suppose that

- $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{p} \in S_p(k)$,

where \mathfrak{m} denotes the maximal ideal of \mathcal{O} . Then one can show that, for any finite abelian extension K'/k and prime $\mathfrak{p} \in S_p(k)$,

$$H^1(G_{k_{\mathfrak{p}}}, T \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\mathrm{Gal}(K'/k)]^c)$$

is a free $\mathcal{O}[\mathrm{Gal}(K'/k)]$ -module of rank $[k_{\mathfrak{p}} : \mathbb{Q}_p]$ (see Corollary A.4). Here $\mathbb{Z}_p[\mathrm{Gal}(K'/k)]^{\iota}$ is a free $\mathbb{Z}_p[\mathrm{Gal}(K'/k)]$ -module of rank 1 and the action of $\mathbb{Z}_p[\mathrm{Gal}(K'/k)]$ is defined by

$$\iota: \mathbb{Z}_p[\mathrm{Gal}(K'/k)] \longrightarrow \mathbb{Z}_p[\mathrm{Gal}(K'/k)]; \quad \sum_{g \in \mathrm{Gal}(K'/k)} a_g g \mapsto \sum_{g \in \mathrm{Gal}(K'/k)} a_g g^{-1}.$$

The action of G_k on $\mathbb{Z}_p[\mathrm{Gal}(K'/k)]^{\iota}$ is induced by its $\mathbb{Z}_p[\mathrm{Gal}(K'/k)]$ -action.

Fix an isomorphism

$$\varprojlim_{K'} \bigoplus_{\mathfrak{p}|p} H^1(G_{k_{\mathfrak{p}}}, T \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\mathrm{Gal}(K'/k)]^{\iota}) \xrightarrow{\sim} \varprojlim_{K'} \mathcal{O}[\mathrm{Gal}(K'/k)]^r,$$

where K' runs over all the finite abelian extensions of k .

For each integer $s \geq 0$, we denote by $\mathrm{ES}_s(T)$ the module of Euler systems of rank s for T (see Definition 2.6). The above fixed isomorphism induces an injective ‘rank reduction’ homomorphism

$$\mathrm{ES}_r(T) \hookrightarrow \mathrm{ES}_0(T).$$

Remark 1.2. A rank reduction homomorphism $\mathrm{ES}_r(T) \longrightarrow \mathrm{ES}_s(T)$ (for $0 \leq s \leq r$) was considered firstly by Rubin in [31] and Perrin-Riou in [29]. After that, in the papers [4, 5, 6], Büyükboduk developed a machinery for higher rank Euler systems by using the rank reduction homomorphism.

An Euler system of rank 0 is a collection of elements in group rings satisfying certain norm-relations, and, philosophically, it should be related to p -adic L -functions. We actually show in Lemma 3.5 that the collection of p -adic L -functions for the multiplicative group over totally real fields is an Euler system of rank 0. Let $\mathcal{L}_p^{\chi} \in \mathrm{ES}_0(T)$ denote this Euler system.

Under this setting, the following is the main result of this paper.

Theorem 1.3 (Theorem 3.13). *Suppose that*

- $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$ for each prime \mathfrak{p} of k above p , and
- the module X_k^{χ} is p -torsion-free.

Then there is a unique Euler system of rank r in $\mathrm{ES}_r(T)$ such that its image under the injection $\mathrm{ES}_r(T) \hookrightarrow \mathrm{ES}_0(T)$ is \mathcal{L}_p^{χ} .

Remark 1.4. When $k = \mathbb{Q}$, Theorem 1.3 follows from the well-known relation between cyclotomic units and Kubota–Leopoldt p -adic L -functions. In general, we expect that the higher rank Euler system constructed in Theorem 1.3 gives us the conjectural Rubin–Stark units (up to unit multiples). Under the Rubin–Stark conjecture, this expectation is equivalent to the certain relation between the conjectural Rubin–Stark units and p -adic L -functions which has already been considered by Büyükboduk in the papers [4, 5, 7].

A key ingredient of the proof of Theorem 1.3 is to generalize the notion of the characteristic ideal. In order to define the characteristic ideal of a finitely generated torsion module over a noetherian ring R , we need to impose that R is a normal ring since the definition relies on the structure theorem for finitely generated modules over a normal ring. In this paper, by using an exterior bi-dual instead of the structure theorem, we give a new definition of the characteristic ideal. This new definition is applicable to a finitely generated module over any noetherian ring, and we can compute the image of the injection $\mathrm{ES}_r(T) \hookrightarrow \mathrm{ES}_0(T)$ by using characteristic ideals of certain Iwasawa modules. Furthermore, by using the classical Iwasawa main conjecture for totally real fields proved by Wiles in [36], we show that the image contains the Euler system \mathcal{L}_p^{χ} .

As an application of Theorem 1.3, by using the theory of higher rank Euler, Kolyvagin and Stark systems (which have been developed by Rubin in [32], Mazur–Rubin in [25, 26], Büyükboduk in [4, 5, 6, 8], the author in [33], Burns–Sano in [3], and Burns–Sano and the

author in [1, 2]), we prove that all higher Fitting ideals of X_K^χ are described by analytic invariants canonically associated with Stickelberger elements. To be more precise, one can define an increasing sequence $\{\Theta_{K,\infty}^i\}_{i \geq 0}$ of ideals of the Iwasawa algebra $\Lambda_K = \mathcal{O}[[\text{Gal}(k_\infty K/k)]]$. The ideals $\Theta_{K,\infty}^i$ are generated by some Kolyvagin derivative classes associated with Stickelberger elements (see Definition 5.1).

Corollary 1.5 (Theorem 5.3). *Suppose that*

- *the same assumptions as in Theorem 1.3 hold, and*
- *$H^2(k_{\mathfrak{q}}, T/\mathfrak{m}T)$ vanishes for each prime $\mathfrak{q} \in S_{\text{ram}}(K/k)$.*

Then, for any integer $i \geq 0$, we have

$$\Theta_{K,\infty}^i = \text{Fitt}_{\Lambda_K}^i(X_K^\chi).$$

Remark 1.6. When $K = k$, Corollary 1.5 has already been proved by Kurihara in [19, 20]. Hence Corollary 1.5 can be viewed as an equivariant generalization of Kurihara's result. In the proof of [20, Theorem 2.1], one of the most important steps is the construction of a certain system of elements $\{\kappa_{n,\mathfrak{l}}\}$, and Kurihara proved that the system $\{\kappa_{n,\mathfrak{l}}\}$ has some important and beautiful properties.

To prove Corollary 1.5, we will also construct a similar system of elements $\{\kappa_{n,\mathfrak{l}}^\sigma\}$ in § 5, and prove that the system $\{\kappa_{n,\mathfrak{l}}^\sigma\}$ has the same kind of properties as $\{\kappa_{n,\mathfrak{l}}\}$ (see Proposition 5.6). However, the construction of $\{\kappa_{n,\mathfrak{l}}\}$ and that of $\{\kappa_{n,\mathfrak{l}}^\sigma\}$ are completely different. In this paper, we use Stark systems in order to construct the system $\{\kappa_{n,\mathfrak{l}}^\sigma\}$. It is assumed in [20] that $\mathfrak{n}\mathfrak{l}$ is well-ordered, while we do not need to impose any assumptions in the construction of the system $\{\kappa_{n,\mathfrak{l}}^\sigma\}$ and the proof of its properties.

We remark that the proof of the equality $\Theta_{k,\infty}^i = \text{Fitt}_{\Lambda}^i(X_k^\chi)$ is also different from that in [20]. In this paper, this equality is derived from the fact proved in [3, 33] that all higher Fitting ideals of a Selmer group are controlled by Stark systems. This allows us to control all higher Fitting ideals of the Iwasawa module X_K^χ even in the equivariant case.

1.1. Notation. Let p be an odd prime. For a field k , we fix a separable closure \bar{k} of k and denote by $G_k := \text{Gal}(\bar{k}/k)$ the absolute Galois group of k .

For a commutative ring R and an R -module M , we write

$$M^* := \text{Hom}_R(M, R)$$

for the R -dual of M . For any integer $r \geq 0$, we define an r -th exterior bi-dual $\bigcap_R^r M$ of M to be

$$\bigcap_R^r M := \left(\bigwedge_R^r (M^*) \right)^*.$$

Remark 1.7. If R is a reduced noetherian ring, then we have the canonical isomorphism

$$\left\{ x \in Q \otimes_R \bigwedge_R^r M \mid \Phi(a) \in R \text{ for all } \Phi \in \bigwedge_R^r M^* \right\} \xrightarrow{\sim} \bigcap_R^r M.$$

Here Q denotes the total ring of fractions of R . This fact tells us that the notion of exterior bi-dual $\bigcap_R^r M$ is a generalization of the notion of Rubin-lattice.

For a profinite group G and a topological G -module M , let $C^\bullet(G, M)$ denote the complex of inhomogeneous continuous cochains of G with values in M . We also denote the object in the derived category corresponding to the complex $C^\bullet(G, M)$ by $\mathbf{R}\Gamma(G, M)$. For each integer $i \geq 0$, we write $H^i(G, M)$ for its i -th cohomology group.

For any number field k , we denote by $S_p(k)$ and $S_\infty(k)$ the set of places of k above p and ∞ , respectively. For a finite set S of places of k containing $S_\infty(k)$, we denote by k_S the maximal extension of k contained in \bar{k} which is unramified outside S . Set

$$G_{k,S} := \text{Gal}(k_S/k).$$

For a prime \mathfrak{q} of k , we denote by $k_{\mathfrak{q}}$ the completion of k at \mathfrak{q} . For an algebraic extension K/k , we denote by $S_{\text{ram}}(K/k)$ the set of primes at which K/k is ramified.

For an abelian extension K/k , we write

$$\iota: \mathbb{Z}_p[[\text{Gal}(K/k)]] \longrightarrow \mathbb{Z}_p[[\text{Gal}(K/k)]]$$

for the involution induced by $\sigma \mapsto \sigma^{-1}$ for $\sigma \in \text{Gal}(K/k)$. For a $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ -module M , we set

$$M^\iota := M \otimes_{\mathbb{Z}_p[[\text{Gal}(K/k)]]} \mathbb{Z}_p[[\text{Gal}(K/k)]]^\iota.$$

In this paper, we regard $\mathbb{Z}_p[[\text{Gal}(K/k)]]^\iota$ as a Galois representation of G_k . The action of G_k on $\mathbb{Z}_p[[\text{Gal}(K/k)]]^\iota$ is defined by its $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ -action.

1.2. Acknowledgments. The author would like to express his gratitude to his supervisor Takeshi Tsuji for many helpful discussions. The author would also like to thank Büyükboduk Kâzım, Masato Kurihara and Hiroki Matsui for helpful advice and comments. This work was supported by the Program for Leading Graduate Schools, MEXT, Japan and JSPS KAKENHI Grant Number 17J02456.

2. HIGHER RANK EULER SYSTEMS

In this section, we will recall the definition of a higher rank Euler system and define a characteristic ideal for any finitely generated module over a noetherian ring.

First, let us introduce some notations and hypotheses. Let \mathcal{O} be a complete discrete valuation ring of mixed characteristic $(0, p)$ with maximal ideal \mathfrak{m} . Let k be a number field. In this section, we fix a finite set S of places of a number field k with $S_\infty(k) \cup S_p(k) \subseteq S$ and let T be a free \mathcal{O} -module of finite rank on which $G_{k,S}$ acts continuously. We also fix a \mathbb{Z}_p^s -extension $k_\infty \subseteq \bar{k}$ of k such that $s \geq 1$ and no prime of k splits completely in k_∞ . We then put

$$\Lambda := \mathcal{O}[[\text{Gal}(k_\infty/k)]] \quad \text{and} \quad \mathbb{T} := T \otimes_{\mathcal{O}} \mathcal{O}[[\text{Gal}(k_\infty/k)]]^\iota.$$

We take a pro- p abelian extension $\mathcal{K} \subseteq \bar{k}$ of k such that

- $S_{\text{ram}}(\mathcal{K}/k) \cap S = \emptyset$, and
- \mathcal{K} contains the maximal p -subextension of the ray class field modulo \mathfrak{q} for all but finitely many primes \mathfrak{q} of k .

We denote the set of finite extensions of k in \mathcal{K} by Ω ;

$$\Omega := \{K \mid k \subseteq K \subseteq \mathcal{K}, [K:k] < \infty\}.$$

We also fix a non-empty subset $\Sigma \subseteq S_p(k)$. For a field $K \in \Omega$, we put

- $S_K := S \cup S_{\text{ram}}(K/k)$,
- $\Lambda_K := \mathcal{O}[[\text{Gal}(k_\infty K/k)]]$,
- $\mathbb{T}_K := T \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\text{Gal}(k_\infty K/k)]]^\iota$, and
- $H_\Sigma^1(\mathbb{T}_K) := \bigoplus_{\mathfrak{p} \in \Sigma} H^1(G_{k_{\mathfrak{p}}}, \mathbb{T}_K)$.

For each field $K \in \Omega$, we fix a Λ_K -submodule $H_{f,\Sigma}^1(\mathbb{T}_K)$ of $H_\Sigma^1(\mathbb{T}_K)$. We take an integer $r > 0$. In this section, we assume the following hypotheses:

- (H.0) The class number of k is coprime to p .
- (H.1) The module $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T)$ vanishes for any prime $\mathfrak{p} \in \Sigma$.
- (H.2) For each field $K \in \Omega$, the Λ_K -module $H_{f,\Sigma}^1(\mathbb{T}_K)$ is free of rank r and the canonical map $H_\Sigma^1(\mathbb{T}_{K'}) \longrightarrow H_\Sigma^1(\mathbb{T}_K)$ induces an isomorphism

$$\Lambda_K \otimes_{\Lambda_{K'}} H_{f,\Sigma}^1(\mathbb{T}_{K'}) \xrightarrow{\sim} H_{f,\Sigma}^1(\mathbb{T}_K)$$

for any field $K' \in \Omega$ with $K \subseteq K'$.

- (H.3) The direct sum of localization maps $H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_\Sigma^1(\mathbb{T}_K)$ is injective.

Definition 2.1. For each field $K \in \Omega$, we set

$$H_f^1(G_{k,S_K}, \mathbb{T}_K) := \ker \left(H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_\Sigma^1(\mathbb{T}_K) / H_{f,\Sigma}^1(\mathbb{T}_K) \right).$$

Remark 2.2.

(i) Hypothesis (H.0) implies that, for any field $K \in \Omega$, we have the canonical isomorphism

$$\mathrm{Gal}(k_\infty K/k) \xrightarrow{\sim} \mathrm{Gal}(k_\infty/k) \times \mathrm{Gal}(K/k).$$

Hence we obtain an isomorphism $\Lambda_K \xrightarrow{\sim} \Lambda[\mathrm{Gal}(K/k)]$. In this paper, by using this isomorphism, we identify Λ_K with $\Lambda[\mathrm{Gal}(K/k)]$.

(ii) For any Λ_K -module M , we have the canonical isomorphism

$$\mathrm{Hom}_\Lambda(M, \Lambda) \xrightarrow{\sim} \mathrm{Hom}_{\Lambda_K}(M, \Lambda_K); \varphi \mapsto \left(m \mapsto \sum_{g \in \mathrm{Gal}(K/k)} \varphi(gm)g^{-1} \right).$$

In particular, $\mathrm{Ext}_\Lambda^i(M, \Lambda) \cong \mathrm{Ext}_{\Lambda_K}^i(M, \Lambda_K)$ for each integer $i \geq 0$. Since Λ is a regular local ring with residue field \mathcal{O}/\mathfrak{m} , we have

$$\mathrm{Ext}_{\Lambda_K}^i(\mathcal{O}/\mathfrak{m}, \Lambda_K) \cong \mathrm{Ext}_\Lambda^i(\mathcal{O}/\mathfrak{m}, \Lambda) \cong \begin{cases} 0 & \text{if } i < \dim(\Lambda) = \dim(\Lambda_K), \\ \mathcal{O}/\mathfrak{m} & \text{if } i = \dim(\Lambda) = \dim(\Lambda_K), \end{cases}$$

and we conclude that Λ_K is Gorenstein since the residue field of Λ_K is also \mathcal{O}/\mathfrak{m} .

Definition 2.3. For a field $K \in \Omega$, we define a Λ_K -module $R_{f,\Sigma}(\mathbb{T}_K)$ to be the cokernel of the canonical homomorphism $H_f^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_{f,\Sigma}^1(\mathbb{T}_K)$.

We note that, by the hypothesis (H.3), we have an exact sequence of Λ_K -modules

$$(1) \quad 0 \longrightarrow H_f^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_{f,\Sigma}^1(\mathbb{T}_K) \longrightarrow R_{f,\Sigma}(\mathbb{T}_K) \longrightarrow 0.$$

Definition 2.4.

- (i) Throughout this paper, we fix an arithmetic Frobenius element $\mathrm{Frob}_\mathfrak{q} \in \mathrm{Gal}(\bar{k}/k)$ for each prime \mathfrak{q} of k .
- (ii) For each prime $\mathfrak{q} \notin S$ of k , we define the Frobenius characteristic polynomial at \mathfrak{q} by

$$P_\mathfrak{q}(x) := \det(1 - x \cdot \mathrm{Frob}_\mathfrak{q} \mid T) \in \mathcal{O}[x].$$

- (iii) For a field $K \in \Omega$, take a Λ_K -module M_K . Suppose that $\{M_K\}_{K \in \Omega}$ is an inverse system of $\Lambda[[\mathrm{Gal}(\mathcal{K}/k)]]$ -modules with transition maps $\psi_{K',K}: M_{K'} \longrightarrow M_K$ for $K', K \in \Omega$ with $K \subseteq K'$. We then define a module $\mathrm{ES}(\{M_K\}_{K \in \Omega})$ by

$$\mathrm{ES}(\{M_K\}_{K \in \Omega}) := \left\{ (m_K)_{K \in \Omega} \in \prod_{K \in \Omega} M_K \mid \begin{array}{l} \psi_{K',K}(m_{K'}) = \left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} P_\mathfrak{q}(\mathrm{Frob}_\mathfrak{q}^{-1}) \right) \cdot m_K \\ \text{for any fields } K', K \in \Omega \text{ with } K \subseteq K' \end{array} \right\}.$$

- (iv) We set $\mathcal{E}(T) := \mathrm{ES}(\{\Lambda_K\}_{K \in \Omega})$.

Lemma 2.5. *We have an inverse system $\{\bigcap_{\Lambda_K}^r H^1(G_{k,S_K}, \mathbb{T}_K)\}_{K \in \Omega}$ of $\mathcal{O}[[k_\infty \mathcal{K}/k]]$ -modules, where the transition maps are induced by the canonical maps $\mathbb{T}_{K'} \longrightarrow \mathbb{T}_K$ for fields $K', K \in \Omega$ with $K \subseteq K'$.*

Proof. Let $K \in \Omega$ be a field. Take a field $K' \in \Omega$ containing K . Since no prime of k splits completely in k_∞ and $S_K \subseteq S_{K'}$, we have

$$H^1(G_{k,S_{K'}}, \mathbb{T}_K) = H^1(G_{k,S_K}, \mathbb{T}_K)$$

(see [32, Corollary B.3.6]). By hypothesis (H.1), the complex $\mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_{K'})$ satisfies the assumption in Corollary A.3. Hence $\mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_{K'})$ is a perfect complex of $\Lambda_{K'}$ -modules

having perfect amplitude contained in [1, 2] by Corollary A.3. Since Λ_K and $\Lambda_{K'}$ are Gorenstein by Remark 2.2 (ii), Lemma B.14 shows that we have the canonical homomorphism

$$\bigcap_{\Lambda_{K'}}^r H^1(G_{k,S_{K'}}, \mathbb{T}_{K'}) \longrightarrow \bigcap_{\Lambda_K}^r H^1(\mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_{K'}) \otimes_{\Lambda_{K'}}^{\mathbb{L}} \Lambda_K).$$

By Theorem A.2 (i), we have an isomorphism $\mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_{K'}) \otimes_{\Lambda_{K'}}^{\mathbb{L}} \Lambda_K \xrightarrow{\sim} \mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_K)$, and hence

$$H^1(\mathbf{R}\Gamma(G_{k,S_{K'}}, \mathbb{T}_{K'}) \otimes_{\Lambda_{K'}}^{\mathbb{L}} \Lambda_K) \xrightarrow{\sim} H^1(G_{k,S_{K'}}, \mathbb{T}_K) = H^1(G_{k,S_K}, \mathbb{T}_K).$$

□

Let us recall the definition of higher rank Euler systems for T .

Definition 2.6. For an integer $r \geq 0$, we define the module $\text{ES}_r(T)$ of Euler systems of rank r (for T) by

$$\text{ES}_r(T) := \text{ES} \left(\left\{ \bigcap_{\Lambda_K}^r H^1(G_{k,S_K}, \mathbb{T}_K) \right\}_{K \in \Omega} \right).$$

Note that $\text{ES}_0(T) = \mathcal{E}(T)$. Since the canonical map

$$\bigcap_{\Lambda_K}^r H_f^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow \bigcap_{\Lambda_K}^r H^1(G_{k,S_K}, \mathbb{T}_K)$$

is injective by Lemma C.1, we can also define

$$\text{ES}_{f,r}(T) := \text{ES}_r(T) \cap \prod_{K \in \Omega} \bigcap_{\Lambda_K}^r H_f^1(G_{k,S_K}, \mathbb{T}_K).$$

Remark 2.7. The definition of the polynomial $P_q(x)$ is the same as [25, Definition 1.2.2] and slightly different from the one appeared in [32, Definition 2.1.1] and [1, §6.1]. However, one can switch back between the two choices (see [32, §9.6] and [25, Remark 3.2.3]). Hence Lemma B.15 shows that the definition of higher rank Euler systems in this paper is essentially the same as the one occurred in [1, Definition 6.4].

Next, let us generalize the notion of characteristic ideals.

Definition 2.8. Let R be a noetherian ring and M a finitely generated R -module. Take an integer $r > 0$ and an exact sequence of R -modules

$$0 \longrightarrow N \longrightarrow R^r \longrightarrow M \longrightarrow 0.$$

We define a characteristic ideal of M by

$$\text{char}_R(M) := \text{im} \left(\bigcap_R^r N \longrightarrow \bigcap_R^r R^r = R \right).$$

We study basic properties of characteristic ideals in Appendix C. In particular, we show in Appendix C that the ideal $\text{char}_R(M)$ is independent of the choice of the exact sequence $0 \longrightarrow N \longrightarrow R^r \longrightarrow M \longrightarrow 0$ (Remark C.5) and that, when R is a normal ring, the ideal $\text{char}_R(M)$ coincides with the usual one (Remark C.14).

Since, by Remark 2.2 (ii), the ring Λ_K is Gorenstein for any field $K \in \Omega$, the following proposition follows from Theorem C.10.

Proposition 2.9. *Let $K \in \Omega$ be a field and M a finitely generated Λ_K -module. Then for any Λ_K -submodule N of M , we have*

$$\text{char}_{\Lambda_K}(M) \subseteq \text{char}_{\Lambda_K}(N).$$

Proposition 2.10. *Assume Hypotheses (H.0) – (H.3). Let us fix an isomorphism*

$$\varprojlim_{K \in \Omega} H_{f, \Sigma}^1(\mathbb{T}_K) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^r.$$

Then it naturally induces an injective homomorphism of $\Lambda[[\text{Gal}(\mathcal{K}/k)]]$ -modules

$$\text{ES}_{f,r}(T) \hookrightarrow \mathcal{E}(T).$$

Furthermore, we have

$$\mathcal{E}(T) \cap \prod_{K \in \Omega} \text{char}_{\Lambda_K}(R_{f, \Sigma}(\mathbb{T}_K)) = \text{im}(\text{ES}_{f,r}(T) \hookrightarrow \mathcal{E}(T)).$$

Proof. By the definition of $\mathcal{E}(T)$, the fixed isomorphism

$$\varprojlim_{K \in \Omega} H_{f, \Sigma}^1(\mathbb{T}_K) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^r$$

induces an isomorphism of $\Lambda[[\text{Gal}(\mathcal{K}/k)]]$ -modules

$$\text{ES} \left(\left\{ \bigcap_{\Lambda_K}^r H_{f, \Sigma}^1(\mathbb{T}_K) \right\}_{K \in \Omega} \right) \xrightarrow{\sim} \mathcal{E}(T).$$

By the same argument as in the proof of Lemma 2.5, we obtain an inverse system

$$\left\{ \bigcap_{\Lambda_K}^r H_{\Sigma}^1(\mathbb{T}_K) \right\}_{K \in \Omega},$$

and hence we get a $\Lambda[[\text{Gal}(\mathcal{K}/k)]]$ -module

$$\text{ES} \left(\left\{ \bigcap_{\Lambda_K}^r H_{\Sigma}^1(\mathbb{T}_K) \right\}_{K \in \Omega} \right).$$

Thus we obtain a diagram

$$\begin{array}{ccc} \text{ES}_{f,r}(T) & & \mathcal{E}(T) \\ \downarrow & & \downarrow \\ \text{ES}_r(T) & \longrightarrow & \text{ES} \left(\left\{ \bigcap_{\Lambda_K}^r H_{\Sigma}^1(\mathbb{T}_K) \right\}_{K \in \Omega} \right). \end{array}$$

Furthermore, for each field $K \in \Omega$, there is also a commutative diagram

$$\begin{array}{ccc} \bigcap_{\Lambda_K}^r H_f^1(G_{k, S_K}, \mathbb{T}_K) & \longrightarrow & \Lambda_K \\ \downarrow & & \downarrow \\ \bigcap_{\Lambda_K}^r H^1(G_{k, S_K}, \mathbb{T}_K) & \longrightarrow & \bigcap_{\Lambda_K}^r H_{\Sigma}^1(\mathbb{T}_K). \end{array}$$

These diagrams show the existence of the canonical homomorphism $\text{ES}_{f,r}(T) \longrightarrow \mathcal{E}(T)$, and the injectivity follows from Hypothesis (H.3) and Lemma C.1. Since $H_{f, \Sigma}^1(\mathbb{T}_K)$ is a free Λ_K -module of rank r by Hypothesis (H.2), the second assertion follows from the exact sequence (1) and the definition of characteristic ideal. \square

3. HIGHER RANK EULER SYSTEMS OVER TOTALLY REAL FIELDS

In this section, we will prove the main theorem of this paper.

Throughout this section, let k be a totally real field with degree r and

$$\chi: G_k \longrightarrow \overline{\mathbb{Q}}^\times$$

a non-trivial finite order even character. Let p be an odd prime, coprime to the class number of k and the order of χ . We put

$$L := \overline{\mathbb{Q}}^{\ker(\chi)}, \mathcal{O} := \mathbb{Z}_p[\mathrm{im}(\chi)], S := S_\infty(k) \cup S_p(k) \cup S_{\mathrm{ram}}(L/k), \text{ and } \Sigma := S_p(k).$$

Let

$$T := \mathcal{O}(1) \otimes \chi^{-1},$$

that is, $T \cong \mathcal{O}$ as \mathcal{O} -modules and the Galois group $G_{k,S}$ acts on T via the character $\chi_{\mathrm{cyc}}\chi^{-1}$, where χ_{cyc} denotes the cyclotomic character of k . We write k_∞/k for the cyclotomic \mathbb{Z}_p -extension of k . We also set

$$\Lambda := \mathcal{O}[[\mathrm{Gal}(k_\infty/k)]] \text{ and } \mathbb{T} := T \otimes_{\mathcal{O}} \mathcal{O}[[\mathrm{Gal}(k_\infty/k)]]^\iota.$$

Let \mathcal{K} denote the maximal pro- p abelian extension of k satisfying $S_{\mathrm{ram}}(\mathcal{K}/k) \cap S = \emptyset$. Let Ω , Λ_K , and \mathbb{T}_K be as in §2. Put

$$H_{f,\Sigma}^1(\mathbb{T}_K) := H_\Sigma^1(\mathbb{T}_K) := \bigoplus_{\mathfrak{p} \in S_p(k)} H^1(G_{k_{\mathfrak{p}}}, \mathbb{T}_K).$$

In this setting, the injectivity of the direct sum of localization maps (the weak Leopoldt conjecture for the multiplicative group) is proved by Iwasawa in [16]:

Proposition 3.1. *For a field $K \in \Omega$, the canonical homomorphism*

$$H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_\Sigma^1(\mathbb{T}_K)$$

is injective, i.e., Hypothesis (H.3) is satisfied.

Proof. Let \tilde{S}_K denote the set of primes of KL above S_K . Then, by Shapiro's lemma, we have the following commutative diagram:

$$\begin{array}{ccc} H^1(G_{k,S_K}, \mathbb{T}_K) & \longrightarrow & \bigoplus_{\mathfrak{p} \in S_p(k)} H^1(G_{k_{\mathfrak{p}}}, \mathbb{T}_K) \\ \downarrow & & \downarrow \\ H^1(G_{KL,\tilde{S}_K}, \mathbb{T}') & \longrightarrow & \bigoplus_{\mathfrak{p} \in S_p(KL)} H^1(G_{(KL)_{\mathfrak{p}}}, \mathbb{T}') \end{array}$$

Here $\mathbb{T}' := \mathcal{O}(1) \otimes_{\mathcal{O}} \mathcal{O}[[\mathrm{Gal}(k_\infty/k)]]^\iota$ and the vertical arrows are the restriction maps. The inflation-restriction exact sequence and [32, Lemma B.3.2] show that both of the vertical arrows are injective, and hence it suffices to prove that

$$H^1(G_{KL,\tilde{S}_K}, \mathbb{T}') \longrightarrow \bigoplus_{\mathfrak{p} \in S_p(KL)} H^1(G_{(KL)_{\mathfrak{p}}}, \mathbb{T}')$$

is injective.

We note that [32, Corollary B.3.6] implies

$$H^1(G_{KL,\tilde{S}_K}, \mathbb{T}') = H^1(G_{KL,S_p(KL) \cup S_\infty(KL)}, \mathbb{T}').$$

Hence, by the Poitou-Tate exact sequence, we have the exact sequence

$$H^2(G_{KL,S_p(KL) \cup S_\infty(KL)}, (\mathbb{T}')^\vee(1))^\vee \longrightarrow H^1(G_{KL,\tilde{S}_K}, \mathbb{T}') \longrightarrow \bigoplus_{\mathfrak{p} \in S_p(KL)} H^1(G_{(KL)_{\mathfrak{p}}}, \mathbb{T}'),$$

and the vanishing of $H^2(G_{KL, S_p(KL) \cup S_\infty(KL)}, (\mathbb{T}')^\vee(1))$ follows from [28, Theorem 11.3.2(ii)] since k_∞/k is the cyclotomic \mathbb{Z}_p -extension. \square

3.1. Stickelberger elements and Iwasawa main conjecture. We will recall the definition of Stickelberger elements and prove that the collection of Stickelberger elements is an Euler system of rank 0.

Let $K \in \Omega$ be a field and $n \geq 0$ an integer. Let μ_{p^n} denote the set of p^n -th roots of unity in $\overline{\mathbb{Q}}$ and $\mu_{p^\infty} := \bigcup_{n>0} \mu_{p^n}$. To simplify the notation, we set

$$G_{K,n} := \text{Gal}(KL(\mu_{p^n})/k) = \text{Gal}(K/k) \times \text{Gal}(L/k) \times \text{Gal}(k(\mu_{p^n})/k).$$

We have the complex conjugate $c \in \text{Gal}(L(\mu_p)/k) = G_{k,1}$ since L is a totally real field. For an $\mathcal{O}[G_{k,1}]$ -module M , we have a decomposition

$$M = M^{c=1} \oplus M^{c=-1}.$$

Here $M^{c=\pm 1}$ denotes the ± 1 -eigenspace with respect to the action of the complex conjugate $c \in G_{k,1}$. Since $p \nmid [L(\mu_p): k]$, we have a decomposition

$$M^{c=-1} = \bigoplus_{\psi: \text{ odd}} M^\psi,$$

where ψ runs over all characters of $G_{k,1}$ satisfying $\psi(c) = -1$ and M^ψ denotes the maximal submodule of M on which $G_{k,1}$ acts via the character ψ . Let

$$\omega: G_{k,1} \longrightarrow \text{Gal}(k(\mu_p)/k) \longrightarrow \mathbb{Z}_p^\times$$

denote the Teichmüller character. We write M^b for the component obtained from $M^{c=-1}$ by removing M^ω , that is,

$$M^{c=-1} = M^b \oplus M^\omega.$$

For an element x of M , we write x^b for the component of x in M^b .

Definition 3.2. Let $\zeta_{KL,n,S}(s, \sigma)$ denote the partial zeta function for $\sigma \in G_{K,n}$:

$$\zeta_{KL,n,S}(s, \sigma) := \sum_{(\mathfrak{a}, KL(\mu_{p^n})/k) = \sigma} N(\mathfrak{a})^{-s}$$

where \mathfrak{a} runs over all integral ideals of k coprime to all the primes in S_K such that the Artin symbol $(\mathfrak{a}, KL(\mu_{p^n})/k)$ is equal to σ and $N(\mathfrak{a})$ is the norm of \mathfrak{a} .

It is well-known that the function $\zeta_{KL,n,S}(s, \sigma)$ is continued to a holomorphic function on $\mathbb{C} \setminus \{1\}$. We put

$$\theta_{KL,n,S} := \sum_{\sigma \in G_{K,n}} \zeta_{KL,n,S}(0, \sigma) \sigma^{-1}$$

which is contained in $\mathbb{Q}[G_{K,n}]$ (see [13]).

Proposition 3.3 ([35, Proposition IV.1.8]). *Let $K' \in \Omega$ with $K \subseteq K'$ and n' an integer with $n \leq n'$. Then the image of $\theta_{K',n',S}$ in $\mathbb{Q}[G_{K,n}]$ is*

$$\left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} (1 - \text{Frob}_{\mathfrak{q}}^{-1}) \right) \cdot \theta_{KL,n,S}.$$

The elements $\{\theta_{KL,n,S}^b\}_{n>0}$ are norm-coherent by Proposition 3.3. In addition, Deligne–Ribet proved in [14] that the element $\theta_{KL,n,S}^b$ is contained in $\mathbb{Z}_p[G_{K,n}]^b$. Hence one can define an element

$$\theta_{KL,\infty,S}^b := \varprojlim_{n>0} \theta_{KL,n,S}^b \in \mathbb{Z}_p[[\text{Gal}(KL(\mu_{p^\infty})/k)]]^b.$$

Let

$$\mathrm{Tw}: \mathbb{Z}_p[[\mathrm{Gal}(KL(\mu_{p^\infty})/k)]] \longrightarrow \mathbb{Z}_p[[\mathrm{Gal}(KL(\mu_{p^\infty})/k)]]$$

denote the homomorphism induced by $\sigma \mapsto \chi_{\mathrm{cyc}}(\sigma)\sigma^{-1}$ for $\sigma \in \mathrm{Gal}(KL(\mu_{p^\infty})/k)$. We then get an element

$$\tilde{L}_{p,K}^\chi := e_\chi \mathrm{Tw}(\theta_{KL,\infty,S}^b) \in \Lambda_K,$$

where

$$e_\chi := \frac{1}{[L(\mu_p):k]} \sum_{\sigma \in \mathrm{Gal}(L(\mu_p)/k)} \chi(\sigma)\sigma^{-1}.$$

(Note that χ is an even character.)

Definition 3.4. For each prime $\mathfrak{q} \notin S$, we set

$$u_{\mathfrak{q}} := \chi(\mathrm{Frob}_{\mathfrak{q}})^{-1} \chi_{\mathrm{cyc}}(\mathrm{Frob}_{\mathfrak{q}}) \mathrm{Frob}_{\mathfrak{q}}^{-1} \in \mathcal{O}[[\mathrm{Gal}(k^{\mathrm{ab}}/k)]]^\times.$$

For a field $K \in \Omega$, we define a modified p -adic L -function $L_{p,K}^\chi \in \Lambda_K$ to be

$$L_{p,K}^\chi := \left(\prod_{\mathfrak{q} \in S_{\mathrm{ram}}(K/k)} (-u_{\mathfrak{q}}) \right) \cdot \tilde{L}_{p,K}^\chi.$$

Lemma 3.5. *The system $\{L_{p,K}^\chi\}_{K \in \Omega}$ is contained in $\mathcal{E}(T) = \mathrm{ES}_0(T)$.*

Proof. Let $K, K' \in \Omega$ with $K \subseteq K'$. We note that the diagram

$$\begin{array}{ccc} \mathcal{O}[[\mathrm{Gal}(K'L(\mu_{p^\infty})/k)]] & \xrightarrow{\mathrm{Tw}} & \mathcal{O}[[\mathrm{Gal}(K'L(\mu_{p^\infty})/k)]] \\ \downarrow \pi & & \downarrow \pi \\ \mathcal{O}[[\mathrm{Gal}(KL(\mu_{p^\infty})/k)]] & \xrightarrow{\mathrm{Tw}} & \mathcal{O}[[\mathrm{Gal}(KL(\mu_{p^\infty})/k)]] \end{array}$$

commutes, where π denotes the canonical projection. Hence Proposition 3.3 shows

$$\begin{aligned} \pi(\tilde{L}_{p,K'}^\chi) &= e_\chi(\mathrm{Tw}(\pi(\theta_{K',\infty,S}^b))) \\ &= e_\chi \left(\mathrm{Tw} \left(\left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} (1 - \mathrm{Frob}_{\mathfrak{q}}^{-1}) \right) \cdot (\theta_{K,\infty,S}^b) \right) \right) \\ &= e_\chi \left(\mathrm{Tw} \left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} (1 - \mathrm{Frob}_{\mathfrak{q}}^{-1}) \right) \right) \cdot \tilde{L}_{p,K}^\chi \\ &= \left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} (1 - u_{\mathfrak{q}}^{-1}) \right) \cdot \tilde{L}_{p,K}^\chi. \end{aligned}$$

Since $P_{\mathfrak{q}}(x) = 1 - \chi(\mathrm{Frob}_{\mathfrak{q}})^{-1} \chi_{\mathrm{cyc}}(\mathrm{Frob}_{\mathfrak{q}})x$ by definition, we have

$$P_{\mathfrak{q}}(\mathrm{Frob}_{\mathfrak{q}}^{-1}) = 1 - u_{\mathfrak{q}} = -u_{\mathfrak{q}}(1 - u_{\mathfrak{q}}^{-1}).$$

As $S_{K'} \setminus S_K = S_{\text{ram}}(K') \setminus S_{\text{ram}}(K)$, we conclude

$$\begin{aligned}
\pi(L_{p,K'}^\chi) &= \left(\prod_{\mathfrak{q} \in S_{\text{ram}}(K'/k)} (-u_{\mathfrak{q}}) \right) \cdot \pi(\tilde{L}_{p,K'}^\chi) \\
&= \left(\prod_{\mathfrak{q} \in S_{\text{ram}}(K'/k)} (-u_{\mathfrak{q}}) \prod_{\mathfrak{q} \in S_{K'} \setminus S_K} (1 - u_{\mathfrak{q}}^{-1}) \right) \cdot \tilde{L}_{p,K}^\chi \\
&= \left(\prod_{\mathfrak{q} \in S_{\text{ram}}(K/k)} (-u_{\mathfrak{q}}) \prod_{\mathfrak{q} \in S_{K'} \setminus S_K} P_{\mathfrak{q}}(\text{Frob}_{\mathfrak{q}}^{-1}) \right) \cdot \tilde{L}_{p,K}^\chi \\
&= \left(\prod_{\mathfrak{q} \in S_{K'} \setminus S_K} P_{\mathfrak{q}}(\text{Frob}_{\mathfrak{q}}^{-1}) \right) \cdot L_{p,K}^\chi.
\end{aligned}$$

□

3.2. Iwasawa modules. In this section, we will introduce several Iwasawa modules and recall their important properties. We will freely use the facts in Appendix A.2.

For a topological \mathbb{Z}_p -module M , let

$$M^\vee := \text{Hom}_{\text{cont}}(M, \mathbb{Q}_p/\mathbb{Z}_p)$$

denote the Pontryagin dual of M .

Definition 3.6. For a field $K \in \Omega$, we write $M_{KL,\infty}$ for the maximal p -ramified pro- p abelian extension of $KL(\mu_{p^\infty})$ and set

$$X_K^\chi := e_\chi(\mathcal{O} \otimes_{\mathbb{Z}_p} \text{Gal}(M_{KL,\infty}/KL(\mu_{p^\infty}))).$$

We note that we have the canonical isomorphism

$$H_{\mathcal{F}_{\text{str}}}^1(k, \mathbb{T}_K^\vee(1))^\vee \cong X_K^\chi$$

(see Example A.7 and Definition A.10 for the definition of $H_{\mathcal{F}_{\text{str}}}^1(k, \mathbb{T}_K^\vee(1))$).

Let $\mathfrak{q} \nmid p$ be a prime of k . By [32, Corollary B.3.6], the subgroup of unramified cohomology classes in $H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K)$ coincides with $H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K)$. In particular, we have

$$H_{\mathcal{F}_{\text{str}}}^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K) = H_{\mathcal{F}_{\text{can}}}^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K) = H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K).$$

Furthermore, Proposition 3.1 shows that $H_{\mathcal{F}_{\text{str}}}^1(k, \mathbb{T}_K)$ vanishes. Hence applying Theorem A.11 with $\mathcal{F}_1 = \mathcal{F}_{\text{str}}$ and $\mathcal{F}_2 = \mathcal{F}_{\text{can}}$, we obtain an exact sequence of Λ_K -modules:

$$(2) \quad 0 \longrightarrow H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_{\Sigma}^1(\mathbb{T}_K) \longrightarrow X_K^\chi \longrightarrow H_{\mathcal{F}_{\text{can}}}^1(k, \mathbb{T}_K^\vee(1))^\vee \longrightarrow 0.$$

Definition 3.7. We denote by $M_{KL,S,\infty}$ the maximal S_K -ramified pro- p abelian extension of $KL(\mu_{p^\infty})$ and put

$$X_{K,S}^\chi := e_\chi(\mathcal{O} \otimes_{\mathbb{Z}_p} \text{Gal}(M_{KL,S,\infty}/KL(\mu_{p^\infty}))).$$

We note that we have the canonical isomorphism

$$H^1(G_{k,S_K}, \mathbb{T}_K^\vee(1))^\vee = H_{\mathcal{F}_{S_K}}^1(k, \mathbb{T}_K^\vee(1))^\vee \cong X_{K,S}^\chi.$$

Applying Theorem A.11 with $\mathcal{F}_1 = \mathcal{F}_{S_K}$ and $\mathcal{F}_2 = \mathcal{F}_{\text{str}}$, we get an exact sequence of Λ_K -modules:

$$(3) \quad 0 \longrightarrow \bigoplus_{\mathfrak{q} \in S_K \setminus S} H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K) \longrightarrow X_{K,S}^\chi \longrightarrow X_K^\chi \longrightarrow 0.$$

Remark 3.8. If $\mathfrak{q} \in S_{\text{ram}}(L/k) \setminus S_p(k)$, then local duality shows that

$$H^2(G_{k_{\mathfrak{q}}}, T/\mathfrak{m}T) \cong H^0(G_{k_{\mathfrak{q}}}, (T/\mathfrak{m}T)^\vee(1)) = 0.$$

Furthermore, since $k(\mu_p)/k$ is unramified at \mathfrak{q} , we also see $H^0(G_{k_{\mathfrak{q}}}, T/\mathfrak{m}T) = 0$. Hence local Euler characteristic formula implies that $H^1(G_{k_{\mathfrak{q}}}, T/\mathfrak{m}T)$ vanishes. In particular, the module $H^1(G_{k_{\mathfrak{q}}}, \mathbb{T})$ vanishes, and so

$$X_{k,S}^\chi = X_k^\chi$$

by the exact sequence (3).

Lemma 3.9. *The μ -invariant of $X_{K,S}^\chi$ vanishes, i.e., the module $X_{K,S}^\chi$ is p -torsion-free if and only if the module $H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1))$ vanishes.*

Proof. Since $X_{K,S}^\chi$ is isomorphic to $H^1(G_{k,S_K}, \mathbb{T}_K^\vee(1))^\vee$, we have an exact sequence of Λ_K -modules

$$0 \longrightarrow H^2(G_{k,S_K}, \mathbb{T}_K^\vee(1))^\vee \otimes_{\mathbb{Z}} \mathbb{F}_p \longrightarrow H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1))^\vee \longrightarrow X_{K,S}^\chi[p] \longrightarrow 0.$$

By the Poitou–Tate exact sequence, we also have the exact sequence

$$\bigoplus_{\mathfrak{q} \in S_K} H^0(G_{k_{\mathfrak{q}}}, \mathbb{T}_K) \longrightarrow H^2(G_{k,S_K}, \mathbb{T}_K^\vee(1))^\vee \longrightarrow H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow \bigoplus_{\mathfrak{q} \in S_K} H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K).$$

[32, Lemma B.3.2] implies that the module $H^0(G_{k_{\mathfrak{q}}}, \mathbb{T}_K)$ vanishes for each prime $\mathfrak{q} \in S_K$. Hence Proposition 3.1 shows that $H^2(G_{k,S_K}, \mathbb{T}_K^\vee(1))$ vanishes. Therefore, we obtain an isomorphism

$$H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1))^\vee \xrightarrow{\sim} X_{K,S}^\chi[p],$$

which proves this lemma. \square

By using Lemma 3.9, we obtain the following well-known proposition:

Proposition 3.10 ([17]). *Let $K \in \Omega$ be a field. If the μ -invariant of X_k^χ vanishes, then so does $X_{K,S}^\chi$.*

Proof. By the Poitou–Tate exact sequence and [32, Lemma B.3.2], we have the following commutative diagram with exact rows:

$$\begin{array}{ccccc} 0 = H^2(G_{k,S_K}, (\mathbb{T}/p\mathbb{T})^\vee(1))^\vee & \hookrightarrow & H^1(G_{k,S_K}, \mathbb{T}/p\mathbb{T}) & \longrightarrow & \bigoplus_{\mathfrak{q} \in S_K} H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}/p\mathbb{T}) \\ \downarrow & & \downarrow & & \downarrow \\ H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1))^\vee & \hookrightarrow & H^1(G_{k,S_K}, \mathbb{T}_K/p\mathbb{T}_K) & \longrightarrow & \bigoplus_{\mathfrak{q} \in S_K} H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}_K/p\mathbb{T}_K). \end{array}$$

Let $\mathfrak{q} \in S_K$ be a prime and fix a prime \mathfrak{q}_K of K above \mathfrak{q} . The inflation–restriction sequence and [32, Lemma B.3.2] show

$$\ker \left(H^1(G_{k_{\mathfrak{q}}}, \mathbb{T}/p\mathbb{T}) \longrightarrow H^1(G_{K_{\mathfrak{q}_K}}, \mathbb{T}/p\mathbb{T}) \right) = H^1(\text{Gal}(K_{\mathfrak{q}_K}/k_{\mathfrak{q}}), (\mathbb{T}/p\mathbb{T})^{G_{K_{\mathfrak{q}_K}}}) = 0.$$

Hence the right vertical map in the diagram is injective. Furthermore, $H^2(G_{k,S_K}, (\mathbb{T}/p\mathbb{T})^\vee(1))^\vee$ vanishes by Lemma 3.9 and

$$H^1(G_{k,S_K}, \mathbb{T}/p\mathbb{T}) \xrightarrow{\sim} H^1(G_{K,S_K}, \mathbb{T}/p\mathbb{T})^{\text{Gal}(K/k)} = H^1(G_{k,S_K}, \mathbb{T}_K/p\mathbb{T}_K)^{\text{Gal}(K/k)}$$

by the inflation–restriction exact sequence and [32, Lemma B.3.2]. Thus we conclude that

$$(H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1))^\vee)^{\text{Gal}(K/k)} = 0,$$

which implies $H^2(G_{k,S_K}, (\mathbb{T}_K/p\mathbb{T}_K)^\vee(1)) = 0$ by Nakayama’s lemma. Therefore, this proposition follows from Lemmas 3.9. \square

Corollary 3.11. *If the μ -invariant of X_k^χ vanishes, then, for a field $K \in \Omega$, we have*

$$L_{p,K}^\chi \in \text{char}_{\Lambda_K}(X_K^\chi).$$

Proof. By Proposition 3.10, the μ -invariant of $X_{K,S}^\chi$ vanishes. In this case, Kurihara showed $\text{Fitt}_{\Lambda_K}^0(X_{K,S}^\chi) = L_{p,K}^\chi \Lambda_K$ (see [19, Proposition 2.1]). Hence we conclude that

$$L_{p,K}^\chi \in \text{Fitt}_{\Lambda_K}^0(X_{K,S}^\chi) \subseteq \text{Fitt}_{\Lambda_K}^0(X_K^\chi) \subseteq \text{char}_{\Lambda_K}(X_K^\chi).$$

□

3.3. Higher rank Euler systems for the multiplicative group. In this subsection, we will show the existence of a higher rank Euler system related to the (modified) Stickelberger elements $\{L_{p,K}^\chi\}_{K \in \Omega}$.

Recall that the prime $p > 2$ is coprime to the class number of k and the order of χ . Since $H_{f,\Sigma}^1(\mathbb{T}_K) = H_\Sigma^1(\mathbb{T}_K)$, by the exact sequence (2), we obtain an exact sequence of Λ_K -modules:

$$(4) \quad 0 \longrightarrow R_{f,\Sigma}(\mathbb{T}_K) \longrightarrow X_K^\chi \longrightarrow H_{\mathcal{F}_{\text{can}}^*}^1(k, \mathbb{T}_K^\vee(1))^\vee \longrightarrow 0.$$

Recall that \mathfrak{m} denotes the maximal ideal of \mathcal{O} .

Lemma 3.12. *Suppose that*

$$H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$$

for each prime $\mathfrak{p} \in S_p(k)$. Then Hypotheses (H.0) – (H.3) are satisfied.

Proof. Since we assume that p is coprime to the class number of k and $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{p} \in S_p(k)$, Hypotheses (H.0) and (H.1) are satisfied. Furthermore, we also assume that, for any prime $\mathfrak{p} \in S_p(k)$, the module $H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T)$ vanishes, Corollary A.4 implies Hypothesis (H.2). Hypothesis (H.3) follows from Proposition 3.1. □

Theorem 3.13. *Let us fix an isomorphism $\varprojlim_{K \in \Omega} H_\Sigma^1(\mathbb{T}_K) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^r$. Suppose that*

- $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{p} \in S_p(k)$, and
- the μ -invariant of X_k^χ vanishes.

Then there is a unique Euler system $c^{\text{DR}} = \{c_K^{\text{DR}}\}_{K \in \Omega} \in \text{ES}_r(T)$ of rank r such that the image of c^{DR} under the map $\text{ES}_r(T) \hookrightarrow \mathcal{E}(T)$ (defined in Proposition 2.10) is $\{L_{p,K}^\chi\}_{K \in \Omega}$.

Remark 3.14. By using the Kummer duality and the Ferrero–Washington theorem (proved in [15]), we see that the μ -invariant of X_k^χ vanishes if L/\mathbb{Q} is an abelian extension. Furthermore, by using the argument as the proof of Proposition 3.10, we also see that the μ -invariant of X_k^χ vanishes if there is a subfield $L' \subseteq L$ such that L'/\mathbb{Q} is an abelian extension and that L/L' is a Galois extension of p -power degree.

Proof. Let $K \in \Omega$ be a field. Note that

$$H_f^1(G_{k,S_K}, \mathbb{T}_K) = H^1(G_{k,S_K}, \mathbb{T}_K) \quad \text{and} \quad \text{ES}_{f,r}(T) = \text{ES}_r(T).$$

Since $R_{f,\Sigma}(\mathbb{T}_K)$ is a submodule of X_K^χ by the exact sequence (4), Proposition 2.9 shows that

$$\text{char}_{\Lambda_K}(X_K^\chi) \subseteq \text{char}_{\Lambda_K}(R_{f,\Sigma}(\mathbb{T}_K)).$$

Furthermore, Lemma 3.5 and Corollary 3.11 imply

$$\{L_{p,K}^\chi\}_{K \in \Omega} \in \mathcal{E}(T) \cap \prod_{K \in \Omega} \text{char}_{\Lambda_K}(X_K^\chi) \subseteq \mathcal{E}(T) \cap \prod_{K \in \Omega} \text{char}_{\Lambda_K}(R_{f,\Sigma}(\mathbb{T}_K)).$$

Thus, by Proposition 2.10 and Lemma 3.12, we have

$$\{L_{p,K}^\chi\}_{K \in \Omega} \in \mathcal{E}(T) \cap \prod_{K \in \Omega} \text{char}_{\Lambda_K}(R_{f,\Sigma}(\mathbb{T}_K)) = \text{im}(\text{ES}_r(T) \hookrightarrow \mathcal{E}(T)).$$

4. STARK SYSTEMS

In this section, we use the same notations as in §3. We will freely use the results in Appendix B. Throughout this section, we fix a field $K \in \Omega$ and suppose that

- p is coprime to the class number of k and the order of χ ,
- $H^0(G_{k_p}, T/\mathfrak{m}T) = H^2(G_{k_p}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{p} \in S_p(k)$, and
- $H^2(G_{k_q}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{q} \in S_{\text{ram}}(K/k)$.

Let $n > 0$ be an integer. For a field $K' \in \Omega$, put

$$R_{K',n} := \mathcal{O}/p^n \mathcal{O}[\text{Gal}(k_n K'/k)] \quad \text{and} \quad T_{K',n} := T \otimes_{\mathbb{Z}_p} \mathbb{Z}_p/p^n \mathbb{Z}_p[\text{Gal}(k_n K'/k)]^\iota,$$

where k_n denotes the n -th layer of the cyclotomic \mathbb{Z}_p -extension k_∞/k . Note that $R_{K',n}$ is a zero-dimensional Gorenstein local ring.

By using the second assumption and Corollary A.4, for each prime $\mathfrak{p} \in S_p(k)$, one can fix an isomorphism

$$\varprojlim_{K' \in \Omega} H^1(G_{k_p}, \mathbb{T}_{K'}) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^{[k_p: \mathbb{Q}_p]},$$

which induces an isomorphism

$$\varprojlim_{K' \in \Omega} H_\Sigma^1(\mathbb{T}_{K'}) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^r.$$

Remark 4.1. Let $\mathfrak{q} \in S_{\text{ram}}(K/k)$ be a prime. Since K/k is an abelian p -power degree extension and $\mathfrak{q} \notin S_p(k)$, we have

$$N(\mathfrak{q}) \equiv 1 \pmod{p}.$$

Hence local duality and the vanishing of $H^2(G_{k_q}, T/\mathfrak{m}T)$ imply the vanishing of $H^0(G_{k_q}, T/\mathfrak{m}T)$. Thus, for any integer $n > 0$, local Euler characteristic formula shows

$$H^1(G_{k_q}, T/\mathfrak{m}T) = 0.$$

By Remark 3.8, for a prime $\mathfrak{q} \in S_{\text{ram}}(L/k) \setminus S_p(k)$, we also have

$$H^0(G_{k_q}, T/\mathfrak{m}T) = H^1(G_{k_q}, T/\mathfrak{m}T) = H^2(G_{k_q}, T/\mathfrak{m}T) = 0.$$

In particular, the exact sequence (3) shows that

$$(5) \quad X_{K,S}^\chi = X_K^\chi.$$

4.1. Review of the theory of Stark systems. In this subsection, we recall the definition of Stark systems and some results proved in [1, 33].

We write \mathcal{H} for the Hilbert class field of k . Note that $p \nmid [\mathcal{H}: k]$ since p is coprime to the class number of k . Recall that $L = \overline{k}^{\ker(\chi)}$. Set $q := \#\mathcal{O}/\mathfrak{m}$ and $\ell_{K,n} := \text{length}_{R_{K,n}}(R_{K,n})$. Let

$$\mathcal{P}_n := \mathcal{P}_{K,n} := \{\mathfrak{q} \notin S_K \mid \mathfrak{q} \text{ splits completely in } \mathcal{H}KL(\mu_{q^{n+\ell_{K,n}}}, (\mathcal{O}_k^\times)^{1/q^{n+\ell_{K,n}}})\},$$

where $(\mathcal{O}_k^\times)^{1/q^{n+\ell_{K,n}}} := \{x \in \overline{k}^\times \mid x^{q^{n+\ell_{K,n}}} \in \mathcal{O}_k^\times\}$. We denote by \mathcal{N}_n the set of square-free products of primes in \mathcal{P}_n . For a prime $\mathfrak{q} \in \mathcal{P}_n$, let $k(\mathfrak{q})$ be the maximal p -extension inside the ray class field modulo \mathfrak{q} . For an ideal $\mathfrak{n} \in \mathcal{N}_n$ we set

$$k(\mathfrak{n}) := \prod_{\mathfrak{q} \mid \mathfrak{n}} k(\mathfrak{q}) \quad \text{and} \quad K(\mathfrak{n}) := k(\mathfrak{n})K.$$

We write $\nu(\mathfrak{n}) := \#\{\mathfrak{q} \mid \mathfrak{n}\}$ for the number of prime divisors of $\mathfrak{n} \in \mathcal{N}_n$.

By definition, for a prime $\mathfrak{q} \in \mathcal{P}_n$, the Frobenius element $\text{Fr}_{\mathfrak{q}}$ acts trivially on $T_{K,n}$. Hence the module

$$H_f^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := H^1(\text{Gal}(k_{\mathfrak{q}}^{\text{ur}}/k_{\mathfrak{q}}), T_{K,n}) \subseteq H^1(G_{k_{\mathfrak{q}}}, T_{K,n})$$

is a free $R_{K,n}$ -module of rank 1 (see [25, Lemma 1.2.1(i)]). Here $k_{\mathfrak{q}}^{\text{ur}}$ denotes the maximal unramified extension of $k_{\mathfrak{q}}$. Furthermore, [25, Lemma 1.2.3] implies that

$$H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := H^1(G_{k_{\mathfrak{q}}}, T_{K,n})/H_f^1(G_{k_{\mathfrak{q}}}, T_{K,n})$$

is also free of rank 1.

Definition 4.2. Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal and \mathcal{F} a Selmer structure on $T_{K,n}$. We then define a Selmer structure $\mathcal{F}^{\mathfrak{n}}$ on $T_{K,n}$ by the following data:

- $S(\mathcal{F}^{\mathfrak{n}}) := S(\mathcal{F}) \cup \{\mathfrak{q} \mid \mathfrak{n}\}$,
- we define a local condition at a prime $\mathfrak{q} \mid \mathfrak{n}$ by

$$H_{\mathcal{F}^{\mathfrak{n}}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := H^1(G_{k_{\mathfrak{q}}}, T_{K,n}),$$

- we define a local condition at a prime $\mathfrak{q} \nmid \mathfrak{n}$ by

$$H_{\mathcal{F}^{\mathfrak{n}}}^1(G_{k_{\mathfrak{q}}}, T) := H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T).$$

Let \mathcal{F} be a Selmer structure on $T_{K,n}$. Then, for any ideals $\mathfrak{n}, \mathfrak{m} \in \mathcal{N}_n$ with $\mathfrak{n} \mid \mathfrak{m}$, we have a cartesian diagram

$$(6) \quad \begin{array}{ccc} H_{\mathcal{F}^{\mathfrak{n}}}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}^{\mathfrak{m}}}^1(k, T_{K,n}) \\ \downarrow & & \downarrow \\ \bigoplus_{\mathfrak{q} \mid \mathfrak{n}} H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) & \hookrightarrow & \bigoplus_{\mathfrak{q} \mid \mathfrak{m}} H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}). \end{array}$$

Definition 4.3. For an ideal $\mathfrak{n} \in \mathcal{N}_n$, we put

$$W_{\mathfrak{n}} := \bigoplus_{\mathfrak{q} \mid \mathfrak{n}} H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n})^*.$$

By Definition B.4, the cartesian diagram (6) induces a homomorphism

$$(7) \quad \Phi_{\mathfrak{m}, \mathfrak{n}} : \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{m})} H_{\mathcal{F}^{\mathfrak{m}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{m}}) \longrightarrow \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{n})} H_{\mathcal{F}^{\mathfrak{n}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{n}})$$

for any integer $r' \geq 0$.

Definition 4.4. Let $r' \geq 0$ be an integer. The module $\text{SS}_{r'}(T_{K,n}, \mathcal{F})$ of Stark systems of rank r' is defined by the inverse limit with respect to $\Phi_{\mathfrak{m}, \mathfrak{n}}$:

$$\text{SS}_{r'}(T_{K,n}, \mathcal{F}) := \varprojlim_{\mathfrak{n} \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{n})} H_{\mathcal{F}^{\mathfrak{n}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{n}}).$$

Throughout this section, by fixing isomorphism $\det(W_{\mathfrak{n}}) \cong R_{K,n}$, we identify the homomorphism (7) with

$$\Phi_{\mathfrak{m}, \mathfrak{n}} : \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{m})} H_{\mathcal{F}^{\mathfrak{m}}}^1(k, T_{K,n}) \longrightarrow \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{n})} H_{\mathcal{F}^{\mathfrak{n}}}^1(k, T_{K,n}),$$

and we regard

$$\text{SS}_{r'}(T_{K,n}, \mathcal{F}) \subseteq \prod_{\mathfrak{n} \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'+\nu(\mathfrak{n})} H_{\mathcal{F}^{\mathfrak{n}}}^1(k, T_{K,n}).$$

To simplify the notation, we put $H_{\Sigma}^1(T_{K,n}) := \bigoplus_{\mathfrak{p} \in S_p(k)} H^1(G_{k_{\mathfrak{p}}}, T_{K,n})$. We note that, by Corollary A.4, the fixed isomorphism $\varprojlim_{K \in \Omega} H_{\Sigma}^1(\mathbb{T}_K) \xrightarrow{\sim} \Lambda[[\text{Gal}(\mathcal{K}/k)]]^r$ induces an $R_{K,n}$ -isomorphism

$$H_{\Sigma}^1(T_{K,n}) \xrightarrow{\sim} R_{K,n}^r.$$

Let us explain that we have the canonical homomorphism

$$(8) \quad \text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \longrightarrow \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}).$$

Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal. By the definition of the Selmer structures \mathcal{F}_{str} and \mathcal{F}_{can} (see Example A.7), we have an exact sequence of $R_{K,n}$ -modules

$$0 \longrightarrow H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow H_{\Sigma}^1(T_{K,n}).$$

Hence, by using the fixed isomorphism $H_{\Sigma}^1(T_{K,n}) \xrightarrow{\sim} R_{K,n}^r$ and Definition B.4, we obtain a homomorphism

$$l_{\mathfrak{n}}: \bigcap_{R_{K,n}}^{r+\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}).$$

Lemma 4.5. *Let $\mathfrak{n}, \mathfrak{m} \in \mathcal{N}_n$ be ideals with $\mathfrak{n} \mid \mathfrak{m}$. Then the diagram*

$$\begin{array}{ccc} \bigcap_{R_{K,n}}^{r+\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{can}}^{\mathfrak{m}}}^1(k, T_{K,n}) & \xrightarrow{(-1)^{r\nu(\mathfrak{m})} l_{\mathfrak{m}}} & \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n}) \\ \Phi_{\mathfrak{m}, \mathfrak{n}} \downarrow & & \Phi_{\mathfrak{m}, \mathfrak{n}} \downarrow \\ \bigcap_{R_{K,n}}^{r+\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) & \xrightarrow{(-1)^{r\nu(\mathfrak{n})} l_{\mathfrak{n}}} & \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) \end{array}$$

commutes. Thus we obtain the canonical homomorphism

$$\text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \longrightarrow \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}); (\epsilon_{\mathfrak{n}})_{\mathfrak{n} \in \mathcal{N}_n} \mapsto ((-1)^{r\nu(\mathfrak{n})} l_{\mathfrak{n}}(\epsilon_{\mathfrak{n}}))_{\mathfrak{n} \in \mathcal{N}_n}.$$

Proof. We have the following commutative diagrams of $R_{K,n}$ -modules:

$$\begin{array}{ccccc} H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}_{\text{can}}^{\mathfrak{m}}}^1(k, T_{K,n}) \\ \downarrow & & \downarrow & & \downarrow \\ W_{\mathfrak{n}}^* & \hookrightarrow & W_{\mathfrak{m}}^* & \hookrightarrow & W_{\mathfrak{m}}^* \oplus H_{\Sigma}^1(T_{K,n}), \\ \\ H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}_{\text{can}}^{\mathfrak{m}}}^1(k, T_{K,n}) \\ \downarrow & & \downarrow & & \downarrow \\ W_{\mathfrak{n}}^* & \hookrightarrow & W_{\mathfrak{n}}^* \oplus H_{\Sigma}^1(T_{K,n}) & \hookrightarrow & W_{\mathfrak{m}}^* \oplus H_{\Sigma}^1(T_{K,n}). \end{array}$$

Since any square of the above two diagrams is cartesian, by Proposition B.5, we get a commutative diagram

$$\begin{array}{ccc} \bigcap_{R_{K,n}}^{r+\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{can}}^{\mathfrak{m}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{m}} \oplus H_{\Sigma}^1(T_{K,n})^*) & \longrightarrow & \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{m}}) \\ \downarrow & & \downarrow \\ \bigcap_{R_{K,n}}^{r+\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{n}} \oplus H_{\Sigma}^1(T_{K,n})^*) & \longrightarrow & \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) \otimes_{R_{K,n}} \det(W_{\mathfrak{n}}). \end{array}$$

Hence this lemma follows from Definition B.4. \square

Proposition 4.6. *Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal with $H^1_{(\mathcal{F}_{\text{can}}^{\mathfrak{n}})^*}(k, T_{K,n}^{\vee}(1)) = 0$. Then the $R_{K,n}$ -module $H^1_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}(k, T_{K,n})$ is free of rank $r + \nu(\mathfrak{n})$ and the projection map*

$$\text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \longrightarrow \prod_{R_{K,n}}^{r+\nu(\mathfrak{n})} H^1_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}(k, T_{K,n})$$

is an isomorphism. In particular, the $R_{K,n}$ -module $\text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}})$ is free of rank 1.

Proof. We put $\mathcal{H}_{\infty} := \mathcal{H}(\mu_{p^{\infty}}, (\mathcal{O}_k^{\times})^{1/p^{\infty}})$. Here $(\mathcal{O}_k^{\times})^{1/p^{\infty}} := \bigcup_{n>0} (\mathcal{O}_k^{\times})^{1/p^n}$.

Since this proposition follows from [33, Lemma 4.6 and Theorem 4.7] (see also [25, Theorem 5.2.15] or [33, Example 3.20]), we only need to check that the following assumptions that appeared in [33, §3 and §4] are satisfied:

- (a) $(T/\mathfrak{m}T)^{G_k} = (T^{\vee}(1)[\mathfrak{m}])^{G_k} = 0$ and $T/\mathfrak{m}T$ is an irreducible $\mathcal{O}/\mathfrak{m}[G_k]$ -module,
- (b) There is a $\tau \in \text{Gal}(\bar{k}/\mathcal{H}_{\infty})$ such that $T_{K,n}/(\tau - 1)T_{K,n} \cong R_{K,n}$ as $R_{K,n}$ -modules,
- (c) $H^1(\text{Gal}(\mathcal{H}_{\infty}KL/k), T/\mathfrak{m}T) = H^1(\text{Gal}(\mathcal{H}_{\infty}KL/k), T^{\vee}(1)[\mathfrak{m}]) = 0$,
- (d) the Selmer structure \mathcal{F}_{can} is cartesian (see [33, Definition 3.8]).

Since χ is an even non-trivial character, we have $\chi \neq 1, \omega$. Here ω denotes the Teichmüller character. Hence Claim (a) is clear, and $\tau = 1$ satisfies Claim (b).

Let us show Claim (c). Since $p \nmid [L(\mu_p): k]$, we have

$$H^1(\text{Gal}(\mathcal{H}_{\infty}KL/k), T/\mathfrak{m}T) = \text{Hom}(\text{Gal}(\mathcal{H}_{\infty}KL/L(\mu_p)), \mathcal{O}/\mathfrak{m} \otimes \omega\chi^{-1})^{\text{Gal}(L(\mu_p)/k)}.$$

Hence if $H^1(\text{Gal}(\mathcal{H}_{\infty}KL/k), T/\mathfrak{m}T)$ does not vanish, there is an abelian extension $N/L(\mu_p)$ in $\mathcal{H}_{\infty}KL$ such that

- N/k is a Galois extension,
- $\text{Gal}(N/L(\mu_p)) \cong \mathcal{O}/\mathfrak{m}$ as abelian groups, and
- $\text{Gal}(L(\mu_p)/k)$ acts on $\text{Gal}(N/L(\mu_p))$ via the character $\omega\chi^{-1}$.

Put $A := \mathcal{H}KL(\mu_{p^{\infty}})$. Since A/k is an abelian extension, the action of $\text{Gal}(L(\mu_p)/k)$ on $\text{Gal}(A/L(\mu_p))$ is trivial. The fact that $\omega\chi^{-1}$ is not trivial implies that

$$N \cap A = L(\mu_p).$$

Since $AN/L(\mu_p)$ is an abelian extension, the action of $\text{Gal}(A/k)$ on $\text{Gal}(AN/A)$ factors through $\text{Gal}(L(\mu_p)/k)$. Hence we obtain an isomorphism of $\text{Gal}(L(\mu_p)/k)$ -modules

$$\text{Gal}(AN/A) \xrightarrow{\sim} \text{Gal}(N/L(\mu_p)).$$

Let Q denote the maximal abelian extension of $L(\mu_p)$ in $\mathcal{H}_{\infty}KL$. We note that Q/k is a Galois extension and $N \subseteq Q$. Since $Q/L(\mu_p)$ is an abelian extension, $\text{Gal}(L(\mu_p)/k)$ acts canonically on $\text{Gal}(Q/A)$. Hence the canonical surjection

$$\text{Gal}(Q/A) \longrightarrow \text{Gal}(AN/A) \xrightarrow{\sim} \text{Gal}(N/L(\mu_p))$$

is $\text{Gal}(L(\mu_p)/k)$ -equivariant. Kummer theory shows that $\text{Gal}(L(\mu_p)/k)$ acts on $\text{Gal}(Q/A)$ via ω . This contradicts the fact that $\omega \neq \omega\chi^{-1}$.

The similar argument shows that $H^1(\text{Gal}(\mathcal{H}_{\infty}KL/k)/k, T^{\vee}(1)[\mathfrak{m}])$ also vanishes.

Let us show Claim (d). By Remark 4.1, we only need to check that, for any prime $\mathfrak{p} \in S_p(k)$, the canonical map $H^1(G_{k_{\mathfrak{p}}}, \mathbb{T}_K) \longrightarrow H^1(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T)$ is surjective. Since we assume the vanishing of $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T)$ and $H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T)$, this claim follows from Corollary A.4. \square

We show in the proof of Proposition 4.6 that [33, Hypothesis 3.12] is satisfied. Furthermore, $T/\mathfrak{m}T$ and $(T/\mathfrak{m}T)^{\vee}(1)$ have no nonzero isomorphic $\mathbb{Z}_p[G_k]$ -subquotients. Hence the same argument as the proof of [25, Proposition 3.6.1] shows the following lemma:

Lemma 4.7 ([25, Proposition 3.6.1]). *Let m be an integer with $n \leq m$. For any non-zero elements $c_1, c_2 \in H^1(G_k, T_{K,n})$ and $c_3, c_4 \in H^1(G_k, T_{K,n}^\vee(1))$, there are infinitely many primes $\mathfrak{q} \in \mathcal{P}_m$ such that, for any $1 \leq i \leq 4$, the image of c_i under the localization map at \mathfrak{q} is non-zero.*

Remark 4.8. Let \mathcal{F} be a Selmer structure on $T_{K,n}$ and m a integer with $n \leq m$. By Lemma 4.7, for any integer $n \leq m$, there is an ideal $\mathfrak{n} \in \mathcal{N}_m$ such that

$$H_{(\mathcal{F}^{\mathfrak{n}})^*}^1(k, T_{K,n}^\vee(1)) = \ker \left(H_{\mathcal{F}^*}^1(k, T_{K,n}^\vee(1)) \longrightarrow \bigoplus_{\mathfrak{q}|\mathfrak{n}} H^1(G_{k,\mathfrak{q}}, T_{K,n}^\vee(1)) \right) = 0.$$

Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal with $H_{(\mathcal{F}_{\text{str}}^{\mathfrak{n}})^*}^1(k, T_{K,n}^\vee(1)) = 0$. Applying Theorem A.11 with $\mathcal{F}_1 = \mathcal{F}_{\text{str}}^{\mathfrak{n}}$ and $\mathcal{F}_1 = \mathcal{F}_{\text{can}}^{\mathfrak{n}}$, we get an exact sequence

$$0 \longrightarrow H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow H_{\Sigma}^1(T_{K,n}) \longrightarrow 0.$$

Hence the following proposition is obtained from this exact sequence and Proposition 4.6.

Proposition 4.9. *Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal with $H_{(\mathcal{F}_{\text{str}}^{\mathfrak{n}})^*}^1(k, T_{K,n}^\vee(1)) = 0$. Then the $R_{K,n}$ -module $H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n})$ is free of rank $\nu(\mathfrak{m})$ and the projection map*

$$\text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) \longrightarrow \bigcap_{R_{K,n}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n})$$

is an isomorphism. In particular, the $R_{K,n}$ -module $\text{SS}_0(T_{K,n}, \mathcal{F}_{\text{can}})$ is free of rank 1 and the natural homomorphism

$$\text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \longrightarrow \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$$

is an isomorphism.

Remark 4.10. Proposition 4.9 and [33, Lemma 4.6] show that the core rank $\chi(\mathcal{F}_{\text{str}})$ of \mathcal{F}_{str} is 0 (see [33, Definition 3.19] for the definition of the core rank).

Definition 4.11. Let $\epsilon = (\epsilon_{\mathfrak{n}})_{\mathfrak{n} \in \mathcal{N}_n} \in \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$ be a system. For each $\mathfrak{n} \in \mathcal{N}_n$, we regard $\epsilon_{\mathfrak{n}}$ as an $R_{K,n}$ -homomorphism

$$\epsilon_{\mathfrak{n}} : \bigwedge_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}^{\mathfrak{n}}}^1(k, T_{K,n})^* \longrightarrow R_{K,n}.$$

For an integer $i \geq 0$, we define an ideal $I_i(\epsilon) \subseteq R_{K,n}$ by

$$I_i(\epsilon) := \sum_{\mathfrak{n} \in \mathcal{N}_n, \nu(\mathfrak{n})=i} \text{im}(\epsilon_{\mathfrak{n}}).$$

Remark 4.12. Let $\epsilon \in \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$ be a system.

- (i) By [33, Theorem 4.10] or [1, Theorem 4.6(ii)], we have $I_i(\epsilon) \subseteq I_{i+1}(\epsilon)$ for an integer $i \geq 0$, and hence

$$I_i(\epsilon) = \sum_{\mathfrak{n} \in \mathcal{N}_n, \nu(\mathfrak{n}) \leq i} \text{im}(\epsilon_{\mathfrak{n}}).$$

- (ii) Let $\mathfrak{m} \in \mathcal{N}_n$ be an ideal such that $H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n})$ is free of rank $\nu(\mathfrak{m})$. Then, for each integer $0 \leq i \leq \nu(\mathfrak{m})$, we have

$$I_i(\epsilon) = \sum_{\mathfrak{n}|\mathfrak{m}, \nu(\mathfrak{n})=i} \text{im}(\epsilon_{\mathfrak{n}}) = \sum_{\mathfrak{n}|\mathfrak{m}, \nu(\mathfrak{n}) \leq i} \text{im}(\epsilon_{\mathfrak{n}})$$

(see the proof of [33, Theorem 4.10] or [1, Corollary 4.5]).

Take an ideal $\mathfrak{m} \in \mathcal{N}_{n+1}$ such that

- the $R_{K,n}$ -module $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n})$ is free of rank $\nu(\mathfrak{m})$ and the projection map

$$\text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) \longrightarrow \bigcap_{R_{K,n}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n})$$

is an isomorphism, and that

- the $R_{K,n+1}$ -module $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1})$ is free of rank $\nu(\mathfrak{m})$ and the projection map

$$\text{SS}_0(T_{K,n+1}, \mathcal{F}_{\text{str}}) \longrightarrow \bigcap_{R_{K,n+1}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1})$$

is an isomorphism.

Since the $R_{K,n+1}$ -module $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1})$ is free of rank $\nu(\mathfrak{m})$, the canonical homomorphism

$$\bigwedge_{R_{K,n+1}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1}) \longrightarrow \bigcap_{R_{K,n+1}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1})$$

is an isomorphism. Hence we obtain an $R_{K,n+1}$ -homomorphism

$$\bigcap_{R_{K,n+1}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n+1}) \longrightarrow \bigcap_{R_{K,n}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n}),$$

which induces an $R_{K,n+1}$ -homomorphism

$$\text{SS}_0(T_{K,n+1}, \mathcal{F}_{\text{str}}) \longrightarrow \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}).$$

This homomorphism is independent of the choice of the ideal $\mathfrak{m} \in \mathcal{N}_{n+1}$.

As in [33, §5], we define

$$\text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}}) := \varprojlim_{n>0} \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}).$$

For a system $\epsilon = (\epsilon_n) \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ and an integer $i \geq 0$, we also set

$$I_i(\epsilon) := \varprojlim_{n>0} I_i(\epsilon_n) \subseteq \Lambda_K.$$

We note that we have the canonical projection

$$\text{pr}_K: \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}}) \longrightarrow \varprojlim_{n>0} \bigcap_{R_{K,n}}^0 H_{\text{str}}^1(G_{k,S_K}, T_{K,n}) = \varprojlim_{n>0} R_{K,n} = \Lambda_K.$$

Remark 4.13. By definition, we have $I_0(\epsilon) = \text{pr}_K(\epsilon) \cdot \Lambda_K$ for a system $\epsilon \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$.

Theorem 4.14 ([33, Theorems 5.4 and 5.6]).

- (i) *The Λ_K -module $\text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ is free of rank 1.*
- (ii) *If $\epsilon \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ is a basis, then we have*

$$I_i(\epsilon) = \text{Fitt}_{\Lambda_K}^i(X_K^\chi).$$

In particular, we have $\text{pr}_K(\epsilon) \cdot \Lambda_K = \text{Fitt}_{\Lambda_K}^0(X_K^\chi)$.

4.2. Kolyvagin derivative homomorphism. Let $n > 0$ be an integer. In this subsection, we recall the definition of the Kolyvagin derivative homomorphism

$$\mathcal{D}_{r',n}: \text{ES}_{r'}(T) \longrightarrow \prod_{\mathfrak{n} \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'} H^1(G_{k,S_{K(\mathfrak{n})}}, T_{K,n}).$$

for an integer $r' \geq 0$. We note that $S_{K(\mathfrak{n})} = S_K \cup \{\mathfrak{q} \mid \mathfrak{n}\}$ and

$$H^1(G_{k,S_{K(\mathfrak{n})}}, T_{K,n}) = H_{\mathcal{F}_{\text{can}}}^1(k, T_{K,n})$$

for an ideal $\mathfrak{n} \in \mathcal{N}_n$.

Let $c = (c_{K'})_{K' \in \Omega} \in \mathcal{E}(T)$ be an Euler system of rank 0. Set

$$K_n := k_n K \text{ and } K_n(\mathfrak{n}) = k(\mathfrak{n})K_n$$

for an ideal $\mathfrak{n} \in \mathcal{N}_n$. We note that the field $K(\mathfrak{n})$ is contained in Ω and that there are canonical identifications

$$\mathrm{Gal}(K_n(\mathfrak{q})/K_n) = \mathrm{Gal}(k_n(\mathfrak{q})/k_n) =: G_{\mathfrak{q}} \text{ and } \mathrm{Gal}(K_n(\mathfrak{n})/K_n) = \prod_{\mathfrak{q}|\mathfrak{n}} G_{\mathfrak{q}}.$$

Let $\mathfrak{q} \in \mathcal{P}_n$ be a prime of k . Since p is coprime to the class number of k and $\mathfrak{q} \nmid p$, the field extension $k(\mathfrak{q})/k$ is cyclic, and hence $G_{\mathfrak{q}}$ is cyclic. We fix a generator $\sigma_{\mathfrak{q}}$ of $G_{\mathfrak{q}}$ and denote by $D_{\mathfrak{q}} \in \mathbb{Z}[G_{\mathfrak{q}}]$ the Kolyvagin's derivative operator:

$$D_{\mathfrak{q}} := \sum_{i=0}^{\#G_{\mathfrak{q}}-1} i\sigma_{\mathfrak{q}}^i.$$

We also set $D_{\mathfrak{n}} := \prod_{\mathfrak{q}|\mathfrak{n}} D_{\mathfrak{q}} \in \mathbb{Z}[\mathrm{Gal}(K_n(\mathfrak{n})/K_n)]$. Let

$$\pi_{K(\mathfrak{n}),n} : \Lambda_{K(\mathfrak{n})} \longrightarrow \mathcal{O}/p^n \mathcal{O}[\mathrm{Gal}(K_n(\mathfrak{n})/k)] = R_{K,n}[\mathrm{Gal}(K_n(\mathfrak{n})/K_n)]$$

be the canonical projection. Then it is well-known that Euler system relations imply

$$\kappa(c)_{K,n,n} := \pi_{K(\mathfrak{n}),n}(D_{\mathfrak{n}}c_{K(\mathfrak{n})}) \in R_{K,n}[\mathrm{Gal}(K(\mathfrak{n})/K)]^{\mathrm{Gal}(K(\mathfrak{n})/K)} \xleftarrow{\sim} R_{K,n}$$

(see, for example, [32, Lemma 4.4.2]).

Remark 4.15. Let $I_{\mathfrak{n}} := \ker(R_{K,n}[\mathrm{Gal}(K_n(\mathfrak{n})/K_n)] \longrightarrow R_{K,n})$ denote the augmentation ideal. The proof of [19, Lemma 4.4] shows that $\pi_{K(\mathfrak{n}),n}(c_{K(\mathfrak{n})}) \in I_{\mathfrak{n}}^{\nu(\mathfrak{n})}$ and

$$\pi_{K(\mathfrak{n}),n}(c_{K(\mathfrak{n})}) \equiv (-1)^{\nu(\mathfrak{n})} \kappa(c)_{K,n,n} \cdot \prod_{\mathfrak{q}|\mathfrak{n}} (\sigma_{\mathfrak{q}} - 1) \pmod{I_{\mathfrak{n}}^{\nu(\mathfrak{n})+1}}.$$

Hence one can regard $(-1)^{\nu(\mathfrak{n})} \kappa(c)_{K,n,n} \in R_{K,n}$ as the ‘‘leading coefficient’’ of $\pi_{K(\mathfrak{n}),n}(c_{K(\mathfrak{n})})$.

Definition 4.16. Let $\mathfrak{S}(\mathfrak{n})$ denote the set of permutations of prime divisors of \mathfrak{n} . For a permutation $\tau \in \mathfrak{S}(\mathfrak{n})$, we set $\mathfrak{d}_{\tau} = \prod_{\tau(\mathfrak{q})=\mathfrak{q}} \mathfrak{q}$. By using an isomorphism $\mathbb{Z}/(\#G_{\mathfrak{q}}) = G_{\mathfrak{q}} \xrightarrow{\sim} I_{\mathfrak{q}}/I_{\mathfrak{q}}^2$, we get an element $a_{\tau,\mathfrak{q}} \in \mathbb{Z}/(\#G_{\mathfrak{q}})$ which corresponds to the element $P_{\tau(\mathfrak{q})}(\mathrm{Frob}_{\tau(\mathfrak{q})}^{-1})$ for a prime $\mathfrak{q} \mid \mathfrak{n}/\mathfrak{d}_{\tau}$. Then we define an element $\tilde{\kappa}(c)_{n,n}$ by

$$\tilde{\kappa}(c)_{K,n,n} := \sum_{\tau \in \mathfrak{S}(\mathfrak{n})} \mathrm{sgn}(\tau) \left(\prod_{\mathfrak{q} \mid \frac{\mathfrak{n}}{\mathfrak{d}_{\tau}}} a_{\tau,\mathfrak{q}} \right) \kappa(c)_{K,n,\mathfrak{d}_{\tau}}.$$

Definition 4.17. We call the map

$$\mathcal{D}_{0,n} : \mathcal{E}(T) \longrightarrow \prod_{\mathfrak{n} \in \mathcal{N}_n} R_{K,n}; c \mapsto (\tilde{\kappa}(c)_{k,n,n})_{\mathfrak{n} \in \mathcal{N}_n}$$

a Kolyvagin derivative homomorphism.

Take a field $K' \in \Omega$ containing K . By Lemma B.14, we have the canonical homomorphism

$$\bigcap_{\Lambda_{K'}}^{r'} H^1(G_{k,S_{K'}}, \mathbb{T}_{K'}) \longrightarrow \bigcap_{R_{K',n}}^{r'} H^1(G_{k,S_{K'}}, T_{K',n}).$$

Furthermore, by [3, Proposition A.4], we have the canonical isomorphism

$$\begin{aligned} \left(\prod_{R_{K',n}}^{r'} H^1(G_{k,S_{K'}}, T_{K',n}) \right)^{\text{Gal}(K'/K)} &\cong \prod_{R_{K,n}}^{r'} H^1(G_{k,S_{K'}}, T_{K',n})^{\text{Gal}(K'/K)} \\ &= \prod_{R_{K,n}}^{r'} H^1(G_{k,S_{K'}}, T_{K,n}). \end{aligned}$$

Hence, in the same manner, one can also define a higher Kolyvagin derivative homomorphism

$$\mathcal{D}_{r',n} : \text{ES}_{r'}(T) \longrightarrow \prod_{\mathfrak{n} \in \mathcal{N}_n} \prod_{R_{K,n}}^{r'} H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n})$$

for any integer $r' \geq 0$ (see [3, §4.3] and [1, §§6.3 and 6.4] for the detail).

By using the localization map at p and the fixed isomorphism $H_{\Sigma}^1(T_{K,n}) \xrightarrow{\sim} R_{K,n}^r$, we obtain a homomorphism

$$H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow R_{K,n}^r$$

for each ideal $\mathfrak{n} \in \mathcal{N}_n$. Hence by (11), this map induces a homomorphism

$$\prod_{\mathfrak{n} \in \mathcal{N}_n} \prod_{R_{K,n}}^r H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \longrightarrow R_{K,n}^r.$$

Let $\text{ES}_r(T) = \text{ES}_{f,r}(T) \hookrightarrow \mathcal{E}(T)$ denote the homomorphism defined in Proposition 2.10. The following proposition follows from the construction of the Kolyvagin derivative homomorphisms $\mathcal{D}_{0,n}$ and $\mathcal{D}_{r,n}$.

Proposition 4.18. *The following diagram commutes:*

$$\begin{array}{ccc} \text{ES}_r(T) & \xrightarrow{\mathcal{D}_{r,n}} & \prod_{\mathfrak{n} \in \mathcal{N}_n} \prod_{R_{K,n}}^r H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n}) \\ \downarrow & & \downarrow \\ \mathcal{E}(T) & \xrightarrow{\mathcal{D}_{0,n}} & \prod_{\mathfrak{n} \in \mathcal{N}_n} R_{K,n}. \end{array}$$

4.3. Regulator homomorphism. Let $r' \geq 0$ be an integer and \mathcal{F} a Selmer structure on $T_{K,n}$. In this subsection, we recall the definition of regulator homomorphisms

$$\text{Reg}_{r'} : \text{SS}_{r'}(T_{K,n}, \mathcal{F}) \longrightarrow \prod_{\mathfrak{n} \in \mathcal{N}_n} H_{\mathcal{F}_{\text{can}}^{\mathfrak{n}}}^1(k, T_{K,n})$$

defined in [3, 26], and construct a Stark system $\epsilon_K^{\text{DR}} \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ from the p -adic L -functions $\{L_{p,K'}^{\chi}\}_{K' \in \Omega}$ by using the higher rank Euler system $c^{\text{DR}} \in \text{ES}_r(T)$ constructed in Theorem 3.13.

Lemma 4.19 ([25, Lemmas 1.2.3 and 1.2.4]). *Let $\mathfrak{q} \in \mathcal{P}_n$. We set*

$$H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := \ker \left(H^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \longrightarrow H^1(G_{k(\mathfrak{q})_{\mathfrak{q}}}, T_{K,n}) \right),$$

where $k(\mathfrak{q})_{\mathfrak{q}}$ is the closure of $k(\mathfrak{q})$ in $\bar{k}_{\mathfrak{q}}$. Put $H_{/ \text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := H^1(G_{k_{\mathfrak{q}}}, T_{K,n}) / H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$.

(i) *The composite maps*

$$H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \longrightarrow H^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \longrightarrow H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$$

and

$$H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \longrightarrow H^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \longrightarrow H_{/\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$$

are isomorphisms.

(ii) *The Frobenius $\text{Frob}_{\mathfrak{q}}$ induces an isomorphism*

$$H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \xrightarrow{\sim} T_{K,n}.$$

(iii) The fixed generator $\sigma_{\mathfrak{q}} \in G_{\mathfrak{q}}$ induces an isomorphism

$$H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) \xrightarrow{\sim} T_{K,n}.$$

We fix a basis e of \mathbb{T}_K . By using the isomorphisms in Lemma 4.19 and the fixed basis e of \mathbb{T}_K , we identify $H_f^1(G_{k_{\mathfrak{q}}}, T_{K,n})$, $H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$, $H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$, and $H_{/tr}^1(G_{k_{\mathfrak{q}}}, T_{K,n})$ with $R_{K,n}$. We note that these identifications are compatible with n , i.e., for $*$ \in $\{f, \text{tr}, /f, /tr\}$, the following diagram commutes:

$$\begin{array}{ccc} H_*^1(G_{k_{\mathfrak{q}}}, T_{K,n}) & \xrightarrow{=} & R_{K,n} \\ \downarrow & & \downarrow \\ H_*^1(G_{k_{\mathfrak{q}}}, T_{K,n-1}) & \xrightarrow{=} & R_{K,n-1}. \end{array}$$

Definition 4.20. For an ideal $\mathfrak{n} \in \mathcal{N}_n$ and a prime $\mathfrak{q} \in \mathcal{P}_n$ with $\mathfrak{q} \mid \mathfrak{n}$, we put

$$\varphi_{\mathfrak{q}}: H_{\mathcal{F}_{\text{can}}}^1(k, T_{K,n}) \longrightarrow H_{/tr}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) = R_{K,n}.$$

Definition 4.21. Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal and \mathcal{F} a Selmer structure on $T_{K,n}$. Then we define a Selmer structure $\mathcal{F}(\mathfrak{n})$ on $T_{K,n}$ by the following data:

- $S(\mathcal{F}(\mathfrak{n})) := S(\mathcal{F}) \cup \{\mathfrak{q} \mid \mathfrak{n}\}$,
- we define a local condition at a prime $\mathfrak{q} \mid \mathfrak{n}$ by

$$H_{\mathcal{F}(\mathfrak{n})}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) := H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}),$$

- we define a local condition at a prime $\mathfrak{q} \nmid \mathfrak{n}$ by

$$H_{\mathcal{F}(\mathfrak{n})}^1(G_{k_{\mathfrak{q}}}, T) := H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T).$$

Remark 4.22. By [26, Proposition 14.3], for any integer m with $n \leq m$ and ideal $\mathfrak{n} \in \mathcal{N}_m$, there are infinitely many ideals $\mathfrak{m} \in \mathcal{N}_m$ with $\mathfrak{n} \mid \mathfrak{m}$ such that $H_{\mathcal{F}_{\text{str}(\mathfrak{m})}^*}^1(k, (T/\mathfrak{m}T)^\vee(1))$ vanishes. Then, by [33, Corollary 3.21], the module $H_{\mathcal{F}_{\text{str}(\mathfrak{m})}}^1(k, T/\mathfrak{m}T)$ also vanishes since the core rank of $\chi(\mathcal{F}_{\text{str}})$ is zero by Remark 4.10. By [33, Lemmas 3.13 and 3.14, and Corollary 3.18] (see also [25, Lemma 3.5.3] and [1, § 3.2]), we see that

$$H_{\mathcal{F}_{\text{str}(\mathfrak{m})}}^1(k, T_{K,n}) = H_{\mathcal{F}_{\text{str}(\mathfrak{m})}^*}^1(k, T_{K,n}^\vee(1)) = 0.$$

Applying Theorem A.11 with $\mathcal{F}_1 = \mathcal{F}_{\text{str}(\mathfrak{m})}$ and $\mathcal{F}_2 = \mathcal{F}_{\text{str}}^{\mathfrak{m}}$, we conclude that the homomorphism

$$H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n}) \xrightarrow{\oplus_{\mathfrak{q} \mid \mathfrak{m}} \varphi_{\mathfrak{q}}} \bigoplus_{\mathfrak{q} \mid \mathfrak{m}} R_{K,n}$$

is an isomorphism. Hence $H_{\mathcal{F}_{\text{str}}^{\mathfrak{m}}}^1(k, T_{K,n})^*$ is generated by the set $\{\varphi_{\mathfrak{q}}\}_{\mathfrak{q} \mid \mathfrak{m}}$ and, for any ideal $\mathfrak{m}' \mid \mathfrak{m}$ and system $\epsilon \in \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$, the ideal $\text{im}(\epsilon_{\mathfrak{m}'})$ is generated by the set

$$\{\epsilon_{\mathfrak{m}'}(\wedge_{\mathfrak{q} \mid \mathfrak{n}} \varphi_{\mathfrak{q}}) \mid \nu(\mathfrak{n}) = \nu(\mathfrak{m}'), \mathfrak{n} \mid \mathfrak{m}\}.$$

We also note that, applying Theorem A.11 with $\mathcal{F}_1 = \mathcal{F}_{\text{str}(\mathfrak{m})}$ and $\mathcal{F}_2 = \mathcal{F}_{\mathbb{L}}(\mathfrak{m})$, we see that $H_{\mathcal{F}_{\mathbb{L}}(\mathfrak{m})}^1(k, T_{K,n})$ is a free $R_{K,n}$ -module of rank 1 and $H_{\mathcal{F}_{\mathbb{L}}(\mathfrak{m})}^1(k, T_{K,n}^\vee(1))$ vanishes.

Definition 4.23 ([25, Definition 3.1.3] and [3, Definition 4.1]). Let \mathcal{F} be a Selmer structure on $T_{K,n}$ and $r' > 0$ an integer. Let $\mathfrak{n} \in \mathcal{N}_n$ be an ideal and \mathfrak{q} a prime divisor of \mathfrak{n} . Then we have two cartesian diagrams

$$\begin{array}{ccc} H_{\mathcal{F}_{\mathfrak{q}}(\mathfrak{n})}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}(\mathfrak{n})}^1(k, T_{K,n}) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & H_{\text{tr}}^1(G_{k_{\mathfrak{q}}}, T_{K,n}) = R_{K,n}, \end{array}$$

$$\begin{array}{ccc} H_{\mathcal{F}_q(n)}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}(n/q)}^1(k, T_{K,n}) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & H_f^1(G_{k_q}, T_{K,n}) = R_{K,n}. \end{array}$$

Here we denote by $H_{\mathcal{F}_q(n)}^1(k, T_{K,n})$ the kernel of the map $H_{\mathcal{F}(n)}^1(k, T_{K,n}) \rightarrow H_{\text{tr}}^1(G_{k_q}, T_{K,n})$. Hence by Definition B.4, we obtain maps

$$\varphi_q : \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n)}^1(k, T_{K,n}) \longrightarrow \bigcap_{R_{K,n}}^{r'-1} H_{\mathcal{F}_q(n)}^1(k, T_{K,n})$$

and

$$\text{div}_q : \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n/q)}^1(k, T_{K,n}) \longrightarrow \bigcap_{R_{K,n}}^{r'-1} H_{\mathcal{F}_q(n)}^1(k, T_{K,n}).$$

We say that a family of elements $\{\kappa_n\}_{n \in \mathcal{N}_n} \in \prod_{n \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n)}^1(k, T_{K,n})$ is a Kolyvagin system of rank r' if $\{\kappa_n\}_{n \in \mathcal{N}_n}$ satisfies

$$\varphi_q(\kappa_n) = \text{div}_q(\kappa_{n/q})$$

for any ideal $n \in \mathcal{N}_n$ and prime divisor q of n . We denote by $\text{KS}_{r'}(T_{K,n}, \mathcal{F})$ the module of Kolyvagin systems of rank r' .

Let \mathcal{F} be a Selmer structure on $T_{K,n}$ and $r' \geq 0$ an integer. By definition, we have the following cartesian diagram

$$\begin{array}{ccc} H_{\mathcal{F}(n)}^1(k, T_{K,n}) & \hookrightarrow & H_{\mathcal{F}^n}^1(k, T_{K,n}) \\ \downarrow & & \downarrow \oplus_{q|n} \varphi_q \\ 0 & \longrightarrow & \bigoplus_{q|n} H_{\text{tr}}^1(G_{k_q}, T_{K,n}) = R_{K,n}^{\nu(n)}. \end{array}$$

Hence by applying Definition B.4 to the above cartesian diagram, we see that the homomorphisms $\{\varphi_q\}_{q|n}$ induce the following homomorphism defined by $\Phi \mapsto (-1)^{\nu(n)} \Phi(\wedge_{q|n} \varphi_q)$:

$$\text{Reg}_n : \bigcap_{R_{K,n}}^{r'+\nu(n)} H_{\mathcal{F}^n}^1(k, T_{K,n}) \longrightarrow \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n)}^1(k, T_{K,n}).$$

The regulator homomorphism $\text{Reg}_{r'}$ is defined by $(\epsilon_n)_{n \in \mathcal{N}_n} \mapsto (\text{Reg}_n(\epsilon_n))_{n \in \mathcal{N}_n}$:

$$\text{Reg}_{r'} : \text{SS}_{r'}(T_{K,n}, \mathcal{F}) \longrightarrow \prod_{n \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n)}^1(k, T_{K,n}).$$

Proposition 4.24 ([3, Proposition 4.3]). *Let \mathcal{F} be a Selmer structure. For an integer $r' > 0$, the image of the regulator homomorphism $\text{Reg}_{r'} : \text{SS}_{r'}(T_{K,n}, \mathcal{F}) \rightarrow \prod_{n \in \mathcal{N}_n} \bigcap_{R_{K,n}}^{r'} H_{\mathcal{F}(n)}^1(k, T_{K,n})$ is contained in $\text{KS}_{r'}(T_{K,n}, \mathcal{F})$.*

Lemma 4.25. *The regulator homomorphism*

$$\text{Reg}_0 : \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) \longrightarrow \prod_{n \in \mathcal{N}_n} R_{K,n}$$

is injective.

Proof. By Proposition 4.9 and Remark 4.22, one can take an ideal $m \in \mathcal{N}_n$ such that

- the projection $\text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) \rightarrow \bigcap_{R_{K,n}}^{\nu(m)} H_{\mathcal{F}_{\text{str}}^m}^1(k, T_{K,n})$ is an isomorphism, and

- the homomorphism

$$H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n}) \xrightarrow{\bigoplus_{\mathfrak{q}|\mathfrak{m}} \varphi_{\mathfrak{q}}} \bigoplus_{\mathfrak{q}|\mathfrak{m}} R_{K,n}$$

is an isomorphism.

Then we have a commutative diagram

$$\begin{array}{ccc} \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) & \xrightarrow{\cong} & \bigcap_{R_{K,n}}^{\nu(\mathfrak{m})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n}) \\ \downarrow \text{Reg}_0 & & \downarrow \text{Reg}_{\mathfrak{m}} \\ \prod_{\mathfrak{n} \in \mathcal{N}_n} R_{K,n} & \longrightarrow & R_{K,n}. \end{array}$$

Since $\bigoplus_{\mathfrak{q}|\mathfrak{m}} \varphi_{\mathfrak{q}} : H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n}) \rightarrow \bigoplus_{\mathfrak{q}|\mathfrak{m}} R_{K,n}$ is an isomorphism, the homomorphism $\text{Reg}_{\mathfrak{m}}$ is an isomorphism, and hence Reg_0 is injective. \square

Theorem 4.26 ([1, Theorem 5.2 (i)]). *The regulator homomorphism*

$$\text{Reg}_r : \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{can}}) \longrightarrow \text{KS}_r(T_{K,n}, \mathcal{F}_{\text{can}})$$

is an isomorphism.

Proof. This theorem follows from [1, Theorem 5.2 (i)], and hence we only need to check that Hypotheses 3.2, 3.3 and 4.2 in [1] are satisfied. In the proof of Proposition 4.6, we prove that Hypotheses 3.2 and 3.3 in [1] are satisfied. By Proposition 4.6, Hypotheses 4.2 is satisfied. \square

When $r' = 1$, the following theorem is proved by Mazur–Rubin in [25, Theorem 3.2.4]. When $r' > 1$, it is proved by Burns–Sano and the author in [1, Theorem 6.12].

Theorem 4.27. *For any integer $r' > 0$, the homomorphism $\mathcal{D}_{r',n}$ induces a homomorphism*

$$\mathcal{D}_{r',n} : \text{ES}_r(T) \longrightarrow \text{KS}_{r'}(T_{K,n}, \mathcal{F}_{\text{can}}).$$

By Proposition 4.18, we have the following commutative diagram:

$$(9) \quad \begin{array}{ccccc} \text{ES}_r(T) & \xrightarrow{\mathcal{D}_{r,n}} & \text{KS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) & \xleftarrow[\text{Reg}_r]{\cong} & \text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \\ \downarrow & & \downarrow & & \downarrow \cong \\ \mathcal{E}(T) & \xrightarrow{\mathcal{D}_{0,n}} & \prod_{\mathfrak{n} \in \mathcal{N}_n} R_{K,n} & \xleftarrow[\text{Reg}_0]{} & \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}). \end{array}$$

To simplify the notation, we put $\mathcal{L}_p^\chi := \{L_{p,K'}^\chi\}_{K' \in \Omega} \in \mathcal{E}(T)$.

Proposition 4.28. *Suppose that the same assumptions as in Theorem 3.13 hold. Then there is a unique system*

$$\epsilon_K^{\text{DR}} = (\epsilon_{K,n}^{\text{DR}})_{n>0} \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}}) = \varprojlim_{n>0} \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$$

such that, for any integer $n > 0$,

$$\text{Reg}_0(\epsilon_{K,n}^{\text{DR}}) = \mathcal{D}_{0,n}(\mathcal{L}_p^\chi),$$

that is,

$$(10) \quad \epsilon_{K,n,\mathfrak{n}}^{\text{DR}}(\wedge_{\mathfrak{q}|\mathfrak{n}} \varphi_{\mathfrak{q}}) = (-1)^{\nu(\mathfrak{n})} \tilde{\kappa}(\mathcal{L}_p^\chi)_{K,n,\mathfrak{n}}$$

for any ideal $\mathfrak{n} \in \mathcal{N}_n$. In particular, we have $\text{pr}_K(\epsilon_K^{\text{DR}}) = L_{p,K}^\chi$.

Proof. Let c^{DR} denote the Euler system of rank r constructed in Theorem 3.13 and $n > 0$ an integer. By Theorems 4.26 and 4.27, there exists a unique system

$$\epsilon_{\text{can},n} \in \text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}})$$

such that the image of $\epsilon_{\text{can},n}$ under the regulator map $\text{Reg}_r: \text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \rightarrow \text{KS}_r(T_{K,n}, \mathcal{F}_{\text{can}})$ is $\mathcal{D}_{r,n}(c^{\text{DR}})$. We define a system $\epsilon_{K,n}^{\text{DR}} \in \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$ to be the image of $\epsilon_{\text{can},n}$ under the isomorphism $\text{SS}_r(T_{K,n}, \mathcal{F}_{\text{can}}) \xrightarrow{\sim} \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}})$. Then the relation $\text{Reg}_0(\epsilon_{K,n}^{\text{DR}}) = \mathcal{D}_{0,n}(\mathcal{L}_p^X)$ follows from the commutative diagram (9).

Recall that the identification $H_{\text{tr}}^1(G_{k_q}, T_{K,n}) = R_{K,n}$ is compatible with n . Hence the regulator map $\text{Reg}_0: \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) \rightarrow \prod_{\mathbf{n} \in \mathcal{N}_n} R_{K,n}$ is also compatible with n . Thus, by the definition of $\mathcal{D}_{0,n}$, it is easy to see that the set $\{\epsilon_{K,n}^{\text{DR}}\}_{n>0}$ is an inverse system. Hence $\epsilon_K^{\text{DR}} := (\epsilon_{K,n}^{\text{DR}})_{n>0}$ is an element of $\varprojlim_{n>0} \text{SS}_0(T_{K,n}, \mathcal{F}_{\text{str}}) = \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$. \square

Proposition 4.29. *Suppose that the same assumptions as in Theorem 3.13 hold.*

- (i) *The system ϵ_K^{DR} constructed in Proposition 4.28 is a basis of $\text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$.*
- (ii) *For each integer $i \geq 0$, we have*

$$I_i(\epsilon_K^{\text{DR}}) = \text{Fitt}_{\Lambda_K}^i(X_K^X).$$

Proof. Theorem 4.14(i) shows that the Λ_K -module $\text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ is free of rank 1. Let $\epsilon_K \in \text{SS}_0(\mathbb{T}_K, \mathcal{F}_{\text{str}})$ be a basis. Then there is an element $a \in \Lambda_K$ such that $\epsilon_K^{\text{DR}} = a \cdot \epsilon_K$. Theorem 4.14(ii) and Proposition 4.28 imply

$$a \cdot \text{Fitt}_{\Lambda_K}^0(X_K^X) = I_0(\epsilon_K^{\text{DR}}) = \text{pr}_K(\epsilon_K^{\text{DR}}) \cdot \Lambda_K = L_{p,K}^X \cdot \Lambda_K.$$

Since $\text{Fitt}_{\Lambda_K}^0(X_K^X) = L_{p,K}^X \cdot \Lambda_K$ by [19, Proposition 2.1] and (5), we have

$$aL_{p,K}^X \cdot \Lambda_K = L_{p,K}^X \cdot \Lambda_K.$$

Hence we conclude that a is a unit, and ϵ_K^{DR} is a basis. Claim (ii) follows from Claim (i) and Theorem 4.14(ii). \square

5. ON THE FITTING IDEALS OF X_K^X

In this section, by using the system ϵ_K^{DR} constructed in Proposition 4.29, we prove that all higher Fitting ideals of X_K^X are described by analytic invariants canonically associated with Stickelberger elements.

Throughout this section, we use the same notations as in §§3 and 4. Suppose that the odd prime p is coprime to the class number of k and the order of χ . Fix a field $K \in \Omega$.

Definition 5.1. For each integer $i \geq 0$, we define an ideal $\Theta_{K,n}^i$ by

$$\Theta_{K,n}^i := \langle \tilde{\kappa}(\mathcal{L}_p^X)_{K,n,\mathbf{n}} \mid \nu(\mathbf{n}) \leq i, \mathbf{n} \in \mathcal{N}_n \rangle = \langle \kappa(\mathcal{L}_p^X)_{K,n,\mathbf{n}} \mid \nu(\mathbf{n}) \leq i, \mathbf{n} \in \mathcal{N}_n \rangle \subseteq R_{K,n}.$$

Since $\mathcal{N}_{n+1} \subseteq \mathcal{N}_n$, one can define an ideal

$$\Theta_{K,\infty}^i := \varprojlim_{n>0} \Theta_{K,n}^i \subseteq \Lambda_K.$$

Remark 5.2.

- (i) By Remark 4.15, the ideal $\Theta_{K,n}^i$ is generated by the leading coefficients of elements $\{\pi_{K(\mathbf{n}),n}(L_{p,K(\mathbf{n})}^X)\}_{\mathbf{n} \in \mathcal{N}_n}$, where

$$\pi_{K(\mathbf{n}),n}: \Lambda_{K(\mathbf{n})} \longrightarrow R_{K,n}[\text{Gal}(K_n(\mathbf{n})/K_n)]$$

is the canonical projection map.

- (ii) We have $\Theta_{K,\infty}^0 = L_{p,K}^X \cdot \Lambda_K = \text{Fitt}_{\Lambda_K}^0(X_K^X)$ by [19, Proposition 2.1].

We give a proof of the following theorem in the next subsection. As we mentioned in Remark 1.6, the following theorem is an equivariant generalization of the main theorem in [20].

Theorem 5.3. *Assume that*

- $H^0(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = H^2(G_{k_{\mathfrak{p}}}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{p} \in S_p(k)$, and
- $H^2(G_{k_{\mathfrak{q}}}, T/\mathfrak{m}T) = 0$ for each prime $\mathfrak{q} \in S_{\text{ram}}(K/k)$,
- the μ -invariant of X_k^χ vanishes.

Then, for an integer $i \geq 0$, we have

$$\Theta_{K,\infty}^i = \text{Fitt}_{\Lambda_K}^i(X_K^\chi).$$

5.1. Proof of Theorem 5.3. In this subsection, suppose that the same assumptions as in Theorem 5.3 hold. We fix an integer $n > 0$. Since

$$\varprojlim_{n' > 0} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n'}) = \ker(H^1(G_{k,S_K}, \mathbb{T}_K) \longrightarrow H_{\Sigma}^1(\mathbb{T}_K)) = 0$$

by Proposition 3.1 and $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n'})$ is finite for each integer $n' > 0$, one can take a sequence $\{n_i\}_{i=0}^\infty$ of positive integers such that $n = n_0 < n_1 < n_2 < \dots$ and a canonical map $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_{i+1}}) \longrightarrow H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_i})$ is zero.

Let $\text{End}(\mathcal{P}_{n_i})$ denote the set of maps from \mathcal{P}_{n_i} to \mathcal{P}_{n_i} . Fix an integer $i > 0$ and take an ideal $\mathfrak{n} \in \mathcal{N}_{n_i}$ and a prime $\mathfrak{q} \in \mathcal{P}_{n_i}$ with $\mathfrak{q} \nmid \mathfrak{n}$. For a map $\sigma \in \text{End}(\mathcal{P}_{n_i})$, we set

$$\varphi_{\mathfrak{n}}^\sigma := \wedge_{\tau|\mathfrak{n}} \varphi_{\sigma(\tau)} : \bigwedge_{R_{K,n}}^{\nu(\mathfrak{n})} H^1(G_k, T_{K,n}) \longrightarrow R_{K,n},$$

where the homomorphism φ_{τ} is defined in Definition 4.20 and we fix an order on the set of prime divisors of \mathfrak{n} .

Let

$$\epsilon_{K,n_i}^{\text{DR}} = (\epsilon_{K,n_i,\mathfrak{n}}^{\text{DR}})_{\mathfrak{n} \in \mathcal{N}_{n_i}} \in \text{SS}_0(T_{K,n_i}, \mathcal{F}_{\text{str}})$$

be the system constructed in Proposition 4.28. Let

$$\text{div}_{\mathfrak{q}} : H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n_j}) \xrightarrow{\sim} R_{K,n_j}$$

denote the fixed isomorphism (see the paragraph after Lemma 4.19). We also denote by $\text{div}_{\mathfrak{q}}$ the composite map

$$H^1(G_k, T_{K,n_j}) \longrightarrow H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n_j}) \xrightarrow{\sim} R_{K,n_j}.$$

Recall that

$$\epsilon_{K,n_i,\mathfrak{n}}^{\text{DR}} \in \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_i}) \otimes \det(W_{\mathfrak{n}}).$$

Here $W_{\mathfrak{n}} := \bigoplus_{\mathfrak{q}|\mathfrak{n}} H_{/f}^1(G_{k_{\mathfrak{q}}}, T_{K,n_i})^*$.

Definition 5.4. We define an element $\bar{\epsilon}_{K,n_i,\mathfrak{n},\mathfrak{q}} \in \bigcap_{R_{K,n}}^{\nu(\mathfrak{n}\mathfrak{q})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_i})$ by

$$\epsilon_{K,n_i,\mathfrak{n},\mathfrak{q}}^{\text{DR}} = \bar{\epsilon}_{K,n_i,\mathfrak{n},\mathfrak{q}} \otimes (\wedge_{\tau|\mathfrak{n}} \text{div}_{\tau} \wedge \text{div}_{\mathfrak{q}}),$$

where, in order to define $\wedge_{\tau|\mathfrak{n}} \text{div}_{\tau}$, we use the same order on the set of prime divisors of \mathfrak{n} as above. We also define an element $\bar{\epsilon}_{K,n_i,\mathfrak{n},1} \in \bigcap_{R_{K,n}}^{\nu(\mathfrak{n})} H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_i})$ by

$$\epsilon_{K,n_i,\mathfrak{n}}^{\text{DR}} = \bar{\epsilon}_{K,n_i,\mathfrak{n},1} \otimes \wedge_{\tau|\mathfrak{n}} \text{div}_{\tau}.$$

We set

$$\kappa_{n_i,\mathfrak{n},\mathfrak{q}}^\sigma := (-1)^{\nu(\mathfrak{n})} \bar{\epsilon}_{K,n_i,\mathfrak{n},\mathfrak{q}}(\varphi_{\mathfrak{n}}^\sigma) \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_i}).$$

Note that $\kappa_{n_i,\mathfrak{n},\mathfrak{q}}^\sigma$ is independent of the choice of the fixed order on the set of prime divisors of \mathfrak{n} . We denote by $\kappa_{n_j,\mathfrak{n},\mathfrak{q}}^\sigma$ the image of $\kappa_{n_i,\mathfrak{n},\mathfrak{q}}^\sigma$ in $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n_j})$ for $0 \leq j \leq i$.

Definition 5.5. We set

$$\delta_{n_i, \mathbf{n}}^\sigma := (-1)^{\nu(\mathbf{n})} \bar{\epsilon}_{K, n_i, \mathbf{n}, 1}(\varphi_{\mathbf{n}}^\sigma) \in R_{K, n_i}.$$

The following proposition follows from the construction of cohomology classes $\{\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma\}$.

Proposition 5.6. *The elements $\{\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma \mid \mathbf{n} \in \mathcal{N}_{n_i}, \mathbf{q} \in \mathcal{P}_{n_i}, \mathbf{q} \nmid \mathbf{n}, \sigma \in \text{End}(\mathcal{P}_{n_i})\}$ satisfies the following relations:*

- (i) $\text{div}_{\mathfrak{r}}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) = \varphi_{\sigma(\mathfrak{r})}(\kappa_{n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}}^\sigma)$ for each prime $\mathfrak{r} \mid \mathbf{n}$,
- (ii) $\text{div}_{\mathbf{q}}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) = \delta_{n_j, \mathbf{n}}^\sigma$,
- (iii) $\varphi_{\sigma(\mathfrak{r})}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) = 0$ for each prime $\mathfrak{r} \mid \mathbf{n}$,
- (iv) $\varphi_{\sigma(\mathbf{q})}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) = -\delta_{n_j, \mathbf{n}\mathbf{q}}^\sigma$.

Remark 5.7. The key of the proof of [20, Theorem 2.1] is the construction of certain cohomology classes which satisfy the same properties (i)–(iv) in Proposition 5.6 (see [20, §1]). However, Kurihara's construction is different from ours. We use a higher rank Euler system and a Stark system in order to construct cohomology classes $\{\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma\}$.

Proof. Let us show Claim (i). Since $\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma$ and $\kappa_{n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}}^\sigma$ are independent of the choice of the fixed order on the set of prime divisors of \mathbf{n} and of \mathbf{n}/\mathfrak{r} , respectively, we may assume that $\varphi_{\mathbf{n}}^\sigma = \varphi_{\sigma(\mathfrak{r})} \wedge \varphi_{\mathbf{n}/\mathfrak{r}}^\sigma$. By the definition of Stark systems, we have $\Phi_{\mathbf{n}\mathbf{q}, \mathbf{n}\mathbf{q}/\mathfrak{r}}^{\text{DR}}(\epsilon_{K, n_j, \mathbf{n}\mathbf{q}}^{\text{DR}}) = \epsilon_{K, n_j, \mathbf{n}\mathbf{q}/\mathfrak{r}}^{\text{DR}}$ (see the paragraph after Definition 4.3 for the definition of $\Phi_{\mathbf{n}\mathbf{q}, \mathbf{n}\mathbf{q}/\mathfrak{r}}^{\text{DR}}$). Hence by Definition B.4, we have

$$\text{div}_{\mathfrak{r}}(\bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}}) = \bar{\epsilon}_{K, n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}},$$

which implies

$$\begin{aligned} \text{div}_{\mathfrak{r}}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) &= (-1)^{\nu(\mathbf{n})} \bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}}(\varphi_{\mathbf{n}}^\sigma \wedge \text{div}_{\mathfrak{r}}) \\ &= \text{div}_{\mathfrak{r}}(\bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}})(\varphi_{\sigma(\mathfrak{r})} \wedge \varphi_{\mathbf{n}/\mathfrak{r}}^\sigma) \\ &= \bar{\epsilon}_{K, n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}}(\varphi_{\sigma(\mathfrak{r})} \wedge \varphi_{\mathbf{n}/\mathfrak{r}}^\sigma) \\ &= (-1)^{\nu(\mathbf{n}/\mathfrak{r})} \bar{\epsilon}_{K, n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}}(\varphi_{\mathbf{n}/\mathfrak{r}}^\sigma \wedge \varphi_{\sigma(\mathfrak{r})}) \\ &= \varphi_{\sigma(\mathfrak{r})}(\kappa_{n_j, \mathbf{n}/\mathfrak{r}, \mathbf{q}}^\sigma). \end{aligned}$$

Next, we will show Claim (ii). By the definition of Stark systems, we have $\text{div}_{\mathbf{q}}(\epsilon_{K, n_j, \mathbf{n}\mathbf{q}}^{\text{DR}}) = \epsilon_{K, n_j, \mathbf{n}}^{\text{DR}}$. Hence by Definition B.4, we have

$$\text{div}_{\mathbf{q}}(\bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}}) = (-1)^{\nu(\mathbf{n})} \bar{\epsilon}_{K, n_j, \mathbf{n}, 1},$$

which implies

$$\begin{aligned} \text{div}_{\mathbf{q}}(\kappa_{n_j, \mathbf{n}, \mathbf{q}}^\sigma) &= (-1)^{\nu(\mathbf{n})} \bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}}(\varphi_{\mathbf{n}}^\sigma \wedge \text{div}_{\mathbf{q}}) \\ &= \text{div}_{\mathbf{q}}(\bar{\epsilon}_{K, n_j, \mathbf{n}\mathbf{q}})(\varphi_{\mathbf{n}}^\sigma) \\ &= (-1)^{\nu(\mathbf{n})} \bar{\epsilon}_{K, n_j, \mathbf{n}, 1}(\varphi_{\mathbf{n}}^\sigma) \\ &= \delta_{n_j, \mathbf{n}}^\sigma. \end{aligned}$$

Claim (iii) follows from the fact that $\varphi_{\sigma(\mathfrak{r})} \wedge \varphi_{\mathbf{n}}^\sigma$ vanishes for each prime divisor $\mathfrak{r} \mid \mathbf{n}$.

By definition, the sign difference between $\bar{\epsilon}_{K, n_j, \mathbf{n}\mathbf{q}, 1}$ and $\bar{\epsilon}_{K, n_j, \mathbf{n}, \mathbf{q}}$ is the same as the one between $\varphi_{\mathbf{n}\mathbf{q}}^\sigma$ and $\varphi_{\mathbf{n}}^\sigma \wedge \varphi_{\sigma(\mathbf{q})}$. Since $(-1)^{\nu(\mathbf{n})} \times (-1)^{\nu(\mathbf{n}\mathbf{q})} = -1$, this fact implies Claim (iv). \square

When the map σ is the identity map, put $\delta_{n_i, \mathbf{n}} := \delta_{n_i, \mathbf{n}}^\sigma$ for simplicity.

Lemma 5.8. *We have*

$$I_i(\epsilon_{K, n}^{\text{DR}}) = \langle \delta_{n, \mathbf{n}}^\sigma \mid \mathbf{n} \in \mathcal{N}_{n_i}, \nu(\mathbf{n}) = i, \sigma \in \text{End}(\mathcal{P}_{n_i}) \rangle.$$

Proof. This lemma follows from Remark 4.22 and (10). \square

Lemma 5.9 ([21, Lemma 6.9]). *Let $0 < j \leq i$ and $c \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_j})$. If $\text{div}_{\mathfrak{r}}(c) = 0$ for any $\mathfrak{r} \mid \mathfrak{n}$, then $c = 0$ in $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_{j-1}})$.*

Proof. We have an exact sequence

$$0 \longrightarrow H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_j}) \longrightarrow H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_j}) \xrightarrow{\oplus_{\mathfrak{q} \mid \mathfrak{n}} \text{div}_{\mathfrak{q}}} \bigoplus_{\mathfrak{q} \mid \mathfrak{n}} R_{K, n_j}.$$

Hence $c \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_j})$. Since $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_j}) \longrightarrow H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_{j-1}})$ is zero, we have $c = 0$ in $H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_{j-1}})$. \square

Lemma 5.10 ([21, Proposition 7.14]). *Let $\mathfrak{r} \in \mathcal{P}_{n_i}$ be a prime with $\mathfrak{r} \nmid \mathfrak{nq}$ and $\sigma \in \text{End}(\mathcal{P}_{n_i})$. Take a homomorphism $f: H^1(G_k, T_{K, n}) \longrightarrow R_{K, n}$. Suppose that there is an element $z \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K, n_i})$ with $\text{div}_{\mathfrak{q}}(z) = 1$. If $\nu(\mathfrak{n}) = i - 1$, then we have*

$$f(\kappa_{n, \mathfrak{n}, \mathfrak{q}}^{\sigma}) \equiv -\text{div}_{\mathfrak{r}}(z) f(\kappa_{n, \mathfrak{n}, \mathfrak{r}}^{\sigma}) \pmod{I_{i-1}(\epsilon_{K, n}^{\text{DR}})}.$$

In particular, we have

$$\delta_{n, \mathfrak{n}, \mathfrak{q}}^{\sigma} \equiv \text{div}_{\mathfrak{r}}(z) \varphi_{\sigma(\mathfrak{q})}(\kappa_{n, \mathfrak{n}, \mathfrak{r}}^{\sigma}) \pmod{I_{i-1}(\epsilon_{K, n}^{\text{DR}})}.$$

Proof. The proof of this lemma is based on that of [21, Proposition 7.14]. For an ideal $\mathfrak{m} \mid \mathfrak{n}$, we set $j := \nu(\mathfrak{m})$ and

$$\tilde{\kappa}_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma} := -\text{div}_{\mathfrak{r}}(z) \kappa_{n_i, \mathfrak{m}, \mathfrak{r}}^{\sigma} + \delta_{n_i, \mathfrak{m}}^{\sigma} z + \sum_{\mathfrak{s} \mid \mathfrak{m}} \varphi_{\sigma(\mathfrak{s})}(z) \kappa_{n_i, \mathfrak{m}/\mathfrak{s}, \mathfrak{s}}^{\sigma},$$

where \mathfrak{s} runs over the set of prime divisors of \mathfrak{m} . Let us show

$$\kappa_{n_{i-1-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma} = \tilde{\kappa}_{n_{i-1-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma}$$

by induction on $j \leq i - 1$. Since $\text{div}_{\mathfrak{q}}(z) = 1$, Proposition 5.6 (ii) shows that

$$\begin{aligned} \text{div}_{\mathfrak{q}}(\tilde{\kappa}_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma}) &= \delta_{n_i, \mathfrak{m}}^{\sigma} = \text{div}_{\mathfrak{q}}(\kappa_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma}), \\ \text{div}_{\mathfrak{r}}(\tilde{\kappa}_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma}) &= -\text{div}_{\mathfrak{r}}(z) \delta_{n_i, \mathfrak{m}}^{\sigma} + \text{div}_{\mathfrak{r}}(z) \delta_{n_i, \mathfrak{m}}^{\sigma} = 0 = \text{div}_{\mathfrak{r}}(\kappa_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma}). \end{aligned}$$

Hence if $j = 0$, then $\mathfrak{m} = 1$, and this claim follows from Lemma 5.9.

Let us suppose that $j > 0$. Take a prime divisor \mathfrak{s} of \mathfrak{m} . Then Proposition 5.6 (i) and (ii) implies

$$\begin{aligned} \text{div}_{\mathfrak{s}}(\tilde{\kappa}_{n_i, \mathfrak{m}, \mathfrak{q}}^{\sigma}) &= -\text{div}_{\mathfrak{r}}(z) \text{div}_{\mathfrak{s}}(\kappa_{n_i, \mathfrak{m}, \mathfrak{r}}^{\sigma}) + \varphi_{\sigma(\mathfrak{s})}(z) \text{div}_{\mathfrak{s}}(\kappa_{n_i, \mathfrak{m}/\mathfrak{s}, \mathfrak{s}}^{\sigma}) + \sum_{\mathfrak{s}' \mid \mathfrak{m}/\mathfrak{s}} \varphi_{\sigma(\mathfrak{s}')} (z) \text{div}_{\mathfrak{s}}(\kappa_{n_i, \mathfrak{m}/\mathfrak{s}'\mathfrak{s}, \mathfrak{s}'}^{\sigma}) \\ &= \varphi_{\sigma(\mathfrak{s})}(-\text{div}_{\mathfrak{r}}(z) \kappa_{n_i, \mathfrak{m}/\mathfrak{s}, \mathfrak{r}}^{\sigma}) + \delta_{n_i, \mathfrak{m}/\mathfrak{s}}^{\sigma} \varphi_{\sigma(\mathfrak{s})}(z) + \sum_{\mathfrak{s}' \mid \mathfrak{m}/\mathfrak{s}} \varphi_{\sigma(\mathfrak{s}')} (z) \varphi_{\sigma(\mathfrak{s})}(\kappa_{n_i, \mathfrak{m}/\mathfrak{s}'\mathfrak{s}, \mathfrak{s}'}^{\sigma}) \\ &= \varphi_{\sigma(\mathfrak{s})}(\tilde{\kappa}_{n_i, \mathfrak{m}/\mathfrak{s}, \mathfrak{q}}^{\sigma}). \end{aligned}$$

Hence, the induction hypothesis and Proposition 5.6 (i) show that

$$\text{div}_{\mathfrak{s}}(\tilde{\kappa}_{n_{i-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma}) = \varphi_{\sigma(\mathfrak{s})}(\tilde{\kappa}_{n_{i-j}, \mathfrak{m}/\mathfrak{s}, \mathfrak{q}}^{\sigma}) = \varphi_{\sigma(\mathfrak{s})}(\kappa_{n_{i-j}, \mathfrak{m}/\mathfrak{s}, \mathfrak{q}}^{\sigma}) = \text{div}_{\mathfrak{s}}(\kappa_{n_{i-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma}).$$

Thus, by Lemma 5.9, we conclude that $\kappa_{n_{i-1-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma} = \tilde{\kappa}_{n_{i-1-j}, \mathfrak{m}, \mathfrak{q}}^{\sigma}$.

Take a homomorphism $f: H^1(G_k, T_{K,n}) \rightarrow R_{K,n}$. Since $\delta_{n,n}^\sigma$ and $f(\kappa_{n,n/\mathfrak{s},\mathfrak{s}}^\sigma)$ are contained in $I_{i-1}(\epsilon_{K,n}^{\text{DR}})$ by definition, we have

$$\begin{aligned} f(\kappa_{n,n,\mathfrak{q}}^\sigma) &= f(\tilde{\kappa}_{n,n,\mathfrak{q}}^\sigma) \\ &= -\text{div}_{\mathfrak{r}}(z)f(\kappa_{n,n,\mathfrak{r}}^\sigma) + \delta_{n,n}^\sigma f(z) - \sum_{\mathfrak{s}|\mathfrak{n}} \varphi_{\sigma(\mathfrak{s})}(z)f(\kappa_{n,n/\mathfrak{s},\mathfrak{s}}^\sigma) \\ &\equiv -\text{div}_{\mathfrak{r}}(z)f(\kappa_{n,n,\mathfrak{r}}^\sigma) \pmod{I_{i-1}(\epsilon_{K,n}^{\text{DR}})}. \end{aligned}$$

□

The following lemma is proved by Kurihara in [21].

Lemma 5.11 ([21, Proposition 9.3]). *Let $\mathfrak{m} \in \mathcal{N}_n$ be an ideal, $x \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n})$, and $\mathfrak{q} \mid \mathfrak{m}$. Then there are infinitely many primes $\mathfrak{r} \in \mathcal{P}_n$ with $\mathfrak{r} \nmid \mathfrak{m}$ such that*

- $\varphi_{\sigma(\mathfrak{q})}(x) \in R_{K,n} \cdot \varphi_{\mathfrak{r}}(x)$, and
- there is an element $z \in H_{\mathcal{F}_{\text{str}}}^1(k, T_{K,n})$ with $\text{div}_{\mathfrak{q}}(z) = 1$ and $\text{div}_{\mathfrak{r}}(z) \in R_{K,n}^\times$.

The following corollary follows from Lemmas 5.10 and 5.11.

Corollary 5.12. *Let $\sigma \in \text{End}(\mathcal{P}_{n_i})$ be a map and $\mathfrak{n} \in \mathcal{N}_{n_i}$ an ideal with $\nu(\mathfrak{n}) = i$. Then there are infinitely many primes $\mathfrak{r} \in \mathcal{P}_{n_i}$ such that*

$$\delta_{n,n}^\sigma \in R_{K,n} \cdot \varphi_{\mathfrak{r}}(\kappa_{n,n/\mathfrak{q},\mathfrak{r}}^\sigma) + I_{i-1}(\epsilon_{K,n}^{\text{DR}}).$$

Proof of Theorem 5.3. Let $\mathfrak{n} \in \mathcal{N}_{n_i}$ be an ideal satisfying $\nu(\mathfrak{n}) = i$. Take a map $\sigma \in \text{End}(\mathcal{P}_{n_i})$. Let us prove

$$\delta_{n,n}^\sigma \in \Theta_{K,n}^i + I_{i-1}(\epsilon_{K,n}^{\text{DR}})$$

by induction on $d := i - \#\{\mathfrak{q} \mid \mathfrak{q} \text{ is a prime divisor of } \mathfrak{n} \text{ and } \sigma(\mathfrak{q}) = \mathfrak{q}\}$.

If $d = 0$, then the claim follows from Lemma 5.8. Suppose that $d > 0$. Take a prime divisor \mathfrak{q} of \mathfrak{n} satisfying $\sigma(\mathfrak{q}) \neq \mathfrak{q}$. By Corollary 5.12, there is a prime $\mathfrak{r} \in \mathcal{P}_{n_i}$ such that

$$\delta_{n,n}^\sigma \in R_{K,n} \cdot \varphi_{\mathfrak{r}}(\kappa_{n,n/\mathfrak{q},\mathfrak{r}}^\sigma) + I_{i-1}(\epsilon_{K,n}^{\text{DR}}).$$

We define a map $\sigma' \in \text{End}(\mathcal{P}_{n_i})$ to be $\sigma'(\mathfrak{r}) = \mathfrak{r}$ and $\sigma'(\mathfrak{s}) = \sigma(\mathfrak{s})$ for $\mathfrak{s} \in \mathcal{P}_{n_i} \setminus \{\mathfrak{r}\}$. Then Proposition 5.6 (iv) shows

$$\delta_{n,n}^\sigma \in R_{K,n} \cdot \delta_{n,n\mathfrak{r}/\mathfrak{q}}^{\sigma'} + I_{i-1}(\epsilon_{K,n}^{\text{DR}}).$$

Since $\sigma(\mathfrak{q}) \neq \mathfrak{q}$ and $\sigma'(\mathfrak{r}) = \mathfrak{r}$, we obtain

$$i - \#\{\mathfrak{s} \mid \mathfrak{s} \text{ is a prime divisor of } n\mathfrak{r}/\mathfrak{q} \text{ and } \sigma'(\mathfrak{s}) = \mathfrak{s}\} = d - 1.$$

Hence applying the induction hypothesis for σ' and $n\mathfrak{r}/\mathfrak{q}$, we have $\delta_{n,n\mathfrak{r}/\mathfrak{q}}^{\sigma'} \in \Theta_{K,n}^i + I_{i-1}(\epsilon_{K,n}^{\text{DR}})$ and hence $\delta_{n,n}^\sigma \in \Theta_{K,n}^i + I_{i-1}(\epsilon_{K,n}^{\text{DR}})$.

In particular, Lemma 5.8 shows $I_i(\epsilon_{K,n}^{\text{DR}}) = \Theta_{K,n}^i + I_{i-1}(\epsilon_{K,n}^{\text{DR}})$ for any integer $i > 0$. Hence Proposition 4.29 implies

$$\text{Fitt}_{\Lambda_K}^i(X_K^\times) = I_i(\epsilon_K^{\text{DR}}) = \Theta_{K,\infty}^i + I_{i-1}(\epsilon_K^{\text{DR}}) = \Theta_{K,\infty}^i + \text{Fitt}_{\Lambda_K}^{i-1}(X_K^\times).$$

Since $\text{Fitt}_{\Lambda_K}^0(X_K^\times) \subseteq \Theta_{K,\infty}^1$ by Remark 5.2(ii), we conclude that $\text{Fitt}_{\Lambda_K}^i(X_K^\times) = \Theta_{K,\infty}^i$ for any integer $i > 0$. □

APPENDIX A. NOTES ON GALOIS COHOMOLOGY

A.1. Remarks on continuous cochain complexes. The contents of this subsection are based on [12, Appendix A].

Let R be a complete noetherian local ring with maximal ideal \mathfrak{m} , and k a number field. Fix a finite set S of places of k satisfying $S_\infty(k) \cup S_p(k) \subseteq S$. Let T be a free R -module of finite rank on which $G_{k,S}$ acts continuously; namely, the composite map

$$G_{k,S} \longrightarrow R[G_{k,S}] \longrightarrow \text{End}_R(T)$$

is continuous. Let G be one of the profinite groups in $\{G_{k_p} \mid \mathfrak{p} \in S_p(k)\} \cup \{G_{k,S}\}$. Note that the pair (T, G) satisfies [30, Hypothesis A].

Definition A.1. Let A be a noetherian ring and M^\bullet a complex of A -modules.

- (i) We say that M^\bullet is a perfect complex if M^\bullet is quasi-isomorphic to a bounded complex P^\bullet of projective A -modules of finite type. Furthermore, we say that M^\bullet has perfect amplitude contained in $[a, b]$ if $P^i = 0$ for every $i < a$ and $i > b$.
- (ii) We write $D_{\text{perf}}^{[a,b]}(A)$ for the full subcategory consisting of perfect complexes having perfect amplitude contained in $[a, b]$ in the derived category of the abelian category of A -modules.

Theorem A.2 ([30, Theorem 1.4 (1) and Corollary 1.2]).

- (i) For an ideal I of R , we have the canonical isomorphism

$$\mathbf{R}\Gamma(G, T) \otimes_R^{\mathbb{L}} R/I \xrightarrow{\sim} \mathbf{R}\Gamma(G, T/IT).$$

- (ii) We have $\mathbf{R}\Gamma(G, T) \in D_{\text{perf}}^{[0,2]}(R)$.

Corollary A.3 ([12, Corollary A.5]). If $H^0(G, T/\mathfrak{m}T) = 0$, then we have $\mathbf{R}\Gamma(G, T) \in D_{\text{perf}}^{[1,2]}(R)$.

Proof. By Theorem A.2(ii), there is a complex

$$P^\bullet = [\cdots \longrightarrow 0 \longrightarrow P^0 \longrightarrow P^1 \longrightarrow P^2 \longrightarrow 0 \longrightarrow \cdots]$$

of finitely generated free R -modules such that P^\bullet is quasi-isomorphic to $\mathbf{R}\Gamma(G, T)$. The vanishing of $H^0(G, T/\mathfrak{m}T)$ implies $(\mathfrak{m}^i/\mathfrak{m}^{i+1} \otimes_R T)^G = 0$. Hence the exact sequence

$$0 \longrightarrow \mathfrak{m}^i/\mathfrak{m}^{i+1} \longrightarrow R/\mathfrak{m}^{i+1} \longrightarrow R/\mathfrak{m}^i \longrightarrow 0$$

shows $(T/\mathfrak{m}^i T)^G = 0$ for any integer $i \geq 1$ and hence $H^0(P^\bullet) = T^G = 0$, which means that the map $P^0 \longrightarrow P^1$ is injective. Thus it suffices to show that the cokernel

$$X := \text{coker}(P^0 \longrightarrow P^1)$$

is a projective R -module. Since R is a noetherian local ring, we only need to check

$$\text{Tor}_1^R(X, R/\mathfrak{m}) = 0$$

by the local criterion for flatness. Since P^1 is a flat R -module, we have

$$\begin{aligned} \text{Tor}_1^R(X, R/\mathfrak{m}) &= \ker(P^0 \otimes_R R/\mathfrak{m} \longrightarrow P^1 \otimes_R R/\mathfrak{m}) \\ &= H^0(\mathbf{R}\Gamma(G, T) \otimes_R^{\mathbb{L}} R/\mathfrak{m}). \end{aligned}$$

By Theorem A.2(i), we conclude

$$\text{Tor}_1^R(X, R/\mathfrak{m}) = H^0(\mathbf{R}\Gamma(G, T/\mathfrak{m}T)) = (T/\mathfrak{m}T)^G = 0.$$

□

Corollary A.4 ([12, Corollary A.7]). *If $H^0(G, T/\mathfrak{m}T) = H^2(G, T/\mathfrak{m}T) = 0$, then the R -module $H^1(G, T)$ is free and, for any ideal J of R , the canonical homomorphism*

$$H^1(G, T) \otimes_R R/J \longrightarrow H^1(G, T/JT)$$

is an isomorphism.

Proof. Since $H^2(G, T/\mathfrak{m}T) = 0$, by Theorem A.2(ii), the module $H^2(G, T/JT)$ also vanishes for any ideal J of R , and hence $\mathbf{R}\Gamma(G, T/JT) \in D_{\text{parf}}^{[1,1]}(R/J)$ by Corollary A.3. In particular, $H^1(G, T)$ is free. The second assertion follows from Theorem A.2(i). \square

A.2. Selmer structures. In this subsection, we recall the definition of Selmer structures. The contents of this subsection are based on [25]. In this subsection, we assume that the residue field of R is finite.

Definition A.5. A Selmer structure \mathcal{F} on T is a collection of the following data:

- a finite set $S(\mathcal{F})$ of places of k , including the set S ,
- a choice of R -submodule $H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T) \subseteq H^1(G_{k_{\mathfrak{q}}}, T)$ for each $\mathfrak{q} \in S(\mathcal{F})$.

Put

$$H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T) := H_f^1(G_{k_{\mathfrak{q}}}, T) := \ker(H^1(G_{k_{\mathfrak{q}}}, T) \longrightarrow H^1(\mathcal{I}_{\mathfrak{q}}, T))$$

for each prime $\mathfrak{q} \notin S(\mathcal{F})$. Here $k_{\mathfrak{q}}^{\text{ur}}$ denotes the maximal unramified extension of $k_{\mathfrak{q}}$ and $\mathcal{I}_{\mathfrak{q}} := \text{Gal}(\overline{k}_{\mathfrak{q}}/k_{\mathfrak{q}}^{\text{ur}})$. We call $H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T)$ the local condition of \mathcal{F} at a prime \mathfrak{q} of k .

Remark A.6. Since p is an odd prime, we have $H^1(k_v, T) = 0$ for any infinite place v of k . Thus we ignore local condition at any infinite place of k .

Example A.7. Suppose that R is p -torsion-free.

(i) We define a canonical Selmer structure \mathcal{F}_{can} on T by the following data:

- $S(\mathcal{F}_{\text{can}}) := S$,
- we define a local condition at a prime $\mathfrak{q} \nmid p$ by

$$H_{\mathcal{F}_{\text{can}}}^1(G_{k_{\mathfrak{q}}}, T) := \ker(H^1(G_{k_{\mathfrak{q}}}, T) \longrightarrow H^1(\mathcal{I}_{\mathfrak{q}}, T \otimes_{\mathbb{Z}_p} \mathbb{Q}_p)),$$

- we define a local condition at a prime $\mathfrak{p} \mid p$ by

$$H_{\mathcal{F}_{\text{can}}}^1(G_{k_{\mathfrak{p}}}, T) := H^1(G_{k_{\mathfrak{p}}}, T).$$

(ii) We define a strict Selmer structure \mathcal{F}_{str} on T by the following data:

- $S(\mathcal{F}_{\text{str}}) := S$,
- we define a local condition at a prime $\mathfrak{q} \nmid p$ by

$$H_{\mathcal{F}_{\text{str}}}^1(G_{k_{\mathfrak{q}}}, T) := \ker(H^1(G_{k_{\mathfrak{q}}}, T) \longrightarrow H^1(\mathcal{I}_{\mathfrak{q}}, T \otimes_{\mathbb{Z}_p} \mathbb{Q}_p)),$$

- we define a local condition at a prime $\mathfrak{p} \mid p$ by

$$H_{\mathcal{F}_{\text{str}}}^1(G_{k_{\mathfrak{p}}}, T) := 0.$$

(iii) We define a Selmer structure \mathcal{F}_S on T by the following data:

- $S(\mathcal{F}_S) := S$,
- we define a local condition at a prime $\mathfrak{q} \in S$ by

$$H_{\mathcal{F}_S}^1(G_{k_{\mathfrak{q}}}, T) := 0.$$

Remark A.8. Let \mathcal{F} be a Selmer structure on T and $R \longrightarrow R'$ a surjective ring homomorphism. Then \mathcal{F} induces a Selmer structure on the R' -module $T \otimes_R R'$, that we will denote by $\mathcal{F}_{R'}$. If there is no risk of confusion, then we write \mathcal{F} instead of $\mathcal{F}_{R'}$.

Definition A.9. Let \mathcal{F} be a Selmer structure on T . Set

$$H_{/\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T) := H^1(G_{k_{\mathfrak{q}}}, T) / H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T)$$

for any prime \mathfrak{q} of k . We define the Selmer group $H_{\mathcal{F}}^1(k, T) \subseteq H^1(G_{k, S(\mathcal{F})}, T)$ associated with \mathcal{F} to be the kernel of the direct sum of localization maps:

$$H_{\mathcal{F}}^1(k, T) := \ker \left(H^1(G_{k, S(\mathcal{F})}, T) \longrightarrow \bigoplus_{\mathfrak{q} \in S(\mathcal{F})} H_{/\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T) \right),$$

where \mathfrak{q} runs through all the primes of k .

For a topological \mathbb{Z}_p -module M , let $M^\vee := \text{Hom}_{\text{cont}}(M, \mathbb{Q}_p/\mathbb{Z}_p)$ denote the Pontryagin duality of M . For any integer $n > 0$, let μ_{p^n} denote the group of all p^n -th roots of unity. The Cartier dual of T is defined by

$$T^\vee(1) := T^\vee \otimes_{\mathbb{Z}_p} \varprojlim_n \mu_{p^n} = \text{Hom}_{\text{cont}}(T, \mathbb{Q}_p/\mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \varprojlim_n \mu_{p^n}.$$

Then we have the local Tate paring

$$\langle \cdot, \cdot \rangle_{\mathfrak{q}}: H^1(G_{k_{\mathfrak{q}}}, T) \times H^1(G_{k_{\mathfrak{q}}}, T^\vee(1)) \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p$$

for each prime \mathfrak{q} of k .

Definition A.10. Let \mathcal{F} be a Selmer structure on T . Put

$$H_{\mathcal{F}^*}^1(G_{k_{\mathfrak{q}}}, T^\vee(1)) := \{x \in H^1(G_{k_{\mathfrak{q}}}, T^\vee(1)) \mid \langle y, x \rangle_{\mathfrak{q}} = 0 \text{ for any } y \in H_{\mathcal{F}}^1(G_{k_{\mathfrak{q}}}, T)\}.$$

In this manner, the Selmer structure \mathcal{F} on T gives rise to a Selmer structure \mathcal{F}^* on $T^\vee(1)$. Hence we obtain the dual Selmer group $H_{\mathcal{F}^*}^1(k, T^\vee(1))$ associated with \mathcal{F} ;

$$H_{\mathcal{F}^*}^1(k, T^\vee(1)) := \ker \left(H^1(G_{k, S(\mathcal{F})}, T^\vee(1)) \longrightarrow \bigoplus_{\mathfrak{q} \in S(\mathcal{F})} H_{\mathcal{F}^*}^1(G_{k_{\mathfrak{q}}}, T^\vee(1)) \right).$$

In this paper, we refer to the following theorem as global duality.

Theorem A.11 ([25, Theorem 2.3.4]). *Let \mathcal{F}_1 and \mathcal{F}_2 be Selmer structures on T satisfying $H_{\mathcal{F}_1}^1(k_{\mathfrak{q}}, T) \subseteq H_{\mathcal{F}_2}^1(k_{\mathfrak{q}}, T)$ for any prime \mathfrak{q} of k ,*

Then we have the following exact sequence:

$$\begin{aligned} 0 \longrightarrow H_{\mathcal{F}_1}^1(k, T) \longrightarrow H_{\mathcal{F}_2}^1(k, T) \longrightarrow \bigoplus_{\mathfrak{q}} H_{\mathcal{F}_2}^1(G_{k_{\mathfrak{q}}}, T) / H_{\mathcal{F}_1}^1(G_{k_{\mathfrak{q}}}, T) \\ \longrightarrow H_{\mathcal{F}_1^*}^1(k, T^\vee(1))^\vee \longrightarrow H_{\mathcal{F}_2^*}^1(k, T^\vee(1))^\vee \longrightarrow 0, \end{aligned}$$

where \mathfrak{q} runs through all the primes of k which satisfies $H_{\mathcal{F}_1}^1(G_{k_{\mathfrak{q}}}, T) \neq H_{\mathcal{F}_2}^1(G_{k_{\mathfrak{q}}}, T)$.

APPENDIX B. EXTERIOR BI-DUAL

We fix a noetherian commutative ring R . For each R -module M we set

$$M^* := \text{Hom}_R(M, R).$$

B.1. Definition of the exterior bi-dual. Let M and N be R -modules, and $r \geq 0$ an integer. For a map $\varphi: M \rightarrow N$, there exists a unique homomorphism of R -modules

$$\bigwedge_R^{r+1} M \rightarrow N \otimes_R \bigwedge_R^r M$$

with the property that

$$m_1 \wedge \cdots \wedge m_{r+1} \mapsto \sum_{i=1}^{r+1} (-1)^{i+1} \varphi(m_i) \otimes m_1 \wedge \cdots \wedge m_{i-1} \wedge m_{i+1} \wedge \cdots \wedge m_{r+1}$$

for each subset $\{m_i\}_{1 \leq i \leq r+1}$ of M . We will also denote this map by φ . In particular, for integers $0 \leq r \leq s$, we have the canonical homomorphism

$$\bigwedge_R^r M^* \rightarrow \text{Hom}_R \left(\bigwedge_R^s M, \bigwedge_R^{s-r} M \right); \varphi_1 \wedge \cdots \wedge \varphi_r \mapsto \varphi_r \circ \cdots \circ \varphi_1.$$

Definition B.1. We define the r -th exterior bi-dual $\bigcap_R^r M$ of M by

$$\bigcap_R^r M := \left(\bigwedge_R^r M^* \right)^*.$$

Note that there is a canonical homomorphism

$$\xi_M^r : \bigwedge_R^r M \rightarrow \bigcap_R^r M; m \mapsto (\Phi \mapsto \Phi(m)),$$

which is neither injective nor surjective in general. However, if M is a finitely generated projective R -module, then ξ_M^r is an isomorphism.

Definition B.2. For integers $0 \leq r \leq s$ and $\Phi \in \bigwedge_R^r M^*$, define an R -homomorphism

$$(11) \quad \bigcap_R^s M \rightarrow \bigcap_R^{s-r} M$$

as the R -dual of the R -homomorphism

$$\bigwedge_R^{s-r} M^* \rightarrow \bigwedge_R^s M^*; \Psi \mapsto \Phi \wedge \Psi.$$

We denote the map (11) also by Φ .

One can check that the diagram

$$\begin{array}{ccc} \bigwedge_R^s M & \xrightarrow{\Phi} & \bigwedge_R^{s-r} M \\ \xi_M^s \downarrow & & \downarrow \xi_M^{s-r} \\ \bigcap_R^s M & \xrightarrow{\Phi} & \bigcap_R^{s-r} M \end{array}$$

is commutative.

B.2. Basic properties. In this subsection, suppose that

- R is a zero dimensional Gorenstein local ring.

Hence all free R -modules are injective by the definition of R . In particular, the functor $(-)^* = \text{Hom}_R(-, R)$ is exact. Furthermore, Matlis duality tells us that the canonical map

$$\xi_M^1 : M \rightarrow \bigcap_R^1 M$$

is an isomorphism for any finitely generated R -module M .

Lemma B.3 ([33, Lemma 2.1]). *Let $F \xrightarrow{h} M \xrightarrow{g} N \rightarrow 0$ be an exact sequence of R -modules and $r \geq 0$ an integer. If F is a free R -module of rank $s \leq r$, then there is a unique R -homomorphism*

$$\varphi: \bigwedge_R^{r-s} N \otimes_R \det(F) \longrightarrow \bigwedge_R^r M$$

such that $\varphi(\wedge^{r-s} g(b) \otimes a) = (\wedge^s h(a)) \wedge b$ for any $a \in \det(F)$ and $b \in \bigwedge_R^{r-s} M$.

For $i \in \{1, 2\}$, let M_i be an R -module and F_i a free R -module of rank s_i . Suppose that we have the following cartesian diagram:

$$\begin{array}{ccc} M_1 & \hookrightarrow & M_2 \\ \downarrow & & \downarrow \\ F_1 & \hookrightarrow & F_2, \end{array}$$

where the horizontal arrows are injective. The R -module F_2/F_1 is free of rank $s_2 - s_1$ since F_1 is an injective R -module. Applying Lemma B.3 to the exact sequence $(F_2/F_1)^* \rightarrow M_2^* \rightarrow M_1^* \rightarrow 0$, we obtain an R -homomorphism

$$\tilde{\Phi}: \bigcap_R^r M_2 \otimes_R \det((F_2/F_1)^*) \longrightarrow \bigcap_R^{r-s_2+s_1} M_1$$

for any integer $r \geq s_2 - s_1$.

Definition B.4 ([33, Definition 2.3]). For any cartesian diagram as above and an integer $r \geq s_2 - s_1$, we have an R -homomorphism

$$\begin{aligned} \Phi: \bigcap_R^r M_2 \otimes_R \det(F_2^*) &\xrightarrow{\sim} \bigcap_R^r M_2 \otimes_R \det((F_2/F_1)^*) \otimes_R \det(F_1^*) \\ &\longrightarrow \bigcap_R^{r-s_2+s_1} M_1 \otimes_R \det(F_1^*), \end{aligned}$$

where the first map is induced by the isomorphism

$$\det((F_2/F_1)^*) \otimes_R \det(F_1^*) \xrightarrow{\sim} \det(F_2^*); a \otimes b \longrightarrow a \wedge \tilde{b},$$

where \tilde{b} is a lift of b in $\bigwedge_R^{s_1} F_2^*$ and the second map is induced by $\tilde{\Phi}$.

Proposition B.5 ([33, Proposition 2.4]). *Suppose that we have the following commutative diagram of R -modules*

$$\begin{array}{ccccc} M_1 & \hookrightarrow & M_2 & \hookrightarrow & M_3 \\ \downarrow & & \downarrow & & \downarrow \\ F_1 & \hookrightarrow & F_2 & \hookrightarrow & F_3 \end{array}$$

such that F_1 , F_2 , and F_3 are free of finite rank and the two squares are cartesian. Let $s_i = \text{rank}_R(F_i)$ and $r \geq s_3 - s_1$. For $i, j \in \{1, 2, 3\}$ with $j < i$, we denote by

$$\Phi_{ij}: \bigcap_R^{r-s_3+s_i} M_i \otimes_R \det(F_i^*) \longrightarrow \bigcap_R^{r-s_3+s_j} M_j \otimes_R \det(F_j^*)$$

the map given by Definition B.4. Then we have

$$\Phi_{31} = \Phi_{21} \circ \Phi_{32}.$$

B.3. Base change. In this subsection, we slightly generalize the results proved in [12, Appendix B]. Recall that R is a noetherian commutative ring. Let M be a finitely generated R -module.

Definition B.6.

- (i) The ring R is said to satisfy the condition (G_1) if the local ring $R_{\mathfrak{r}}$ is Gorenstein for any prime \mathfrak{r} of R with $\text{ht}(\mathfrak{r}) \leq 1$.
- (ii) For an integer $n \geq 0$, we say that the R -module M satisfies the Serre's condition (S_n) if the inequality $\text{depth}(M_{\mathfrak{r}}) \geq \min\{n, \dim \text{Supp}(M_{\mathfrak{r}})\}$ holds for all prime ideal \mathfrak{r} of R . Here $M_{\mathfrak{r}} := R_{\mathfrak{r}} \otimes_R M$.
- (iii) The R -module M is said to be 2-syzygy if there exists an exact sequence $0 \rightarrow M \rightarrow P^0 \rightarrow P^1$ of R -modules with P^i projective.

Remark B.7. A noetherian ring satisfying the conditions (G_1) and (S_2) has the following characterization (proved by Evans–Griffith and Matsui–Takahashi–Tsuchiya in [27]).

The noetherian ring R satisfies (G_1) and (S_2) if and only if, for any finitely generated R -module M , the following assertions are equivalent:

- (i) M satisfies (S_2) ;
- (ii) M is 2-syzygy;
- (iii) M is reflexive i.e., $M \rightarrow M^{**}$ is an isomorphism.

Remark B.8. Normal rings and Gorenstein rings satisfy (G_1) and (S_2) .

Remark B.9. If R' is a flat R -algebra, then we have the canonical isomorphism

$$R' \otimes_R \bigcap_R^r M \xrightarrow{\sim} \bigcap_{R'}^r (R' \otimes_R M)$$

for any integer $r \geq 0$. However, in general, there dose not exist a canonical homomorphism from $R' \otimes_R \bigcap_R^r M$ to $\bigcap_{R'}^r (R' \otimes_R M)$. We will show the existence of such a homomorphism if R and R' satisfy (G_1) and (S_2) (see Lemma B.14).

Definition B.10. We say that the finitely generated R -module M is pseudo-null if $M_{\mathfrak{r}}$ vanishes for any prime \mathfrak{r} of R with $\text{ht}(\mathfrak{r}) \leq 1$. When $\dim(R) \leq 1$, the module M is pseudo-null if and only if $M = 0$.

Lemma B.11. *If R satisfies (S_2) and M is a pseudo-null R -module, then we have*

$$\text{Hom}_R(M, R) = \text{Ext}_R^1(M, R) = 0.$$

Proof. We may assume $\dim(R) > 1$. Since R is a noetherian ring, there is a sequence of submodules $0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_n = M$ such that M_{i+1}/M_i is isomorphic to R/\mathfrak{r} for some prime ideal \mathfrak{r} of R for each integer $0 \leq i < n$. Hence we may assume that $M = R/\mathfrak{r}$ for some prime ideal \mathfrak{r} of R . Since R/\mathfrak{r} is pseudo-null, we have $\text{ht}(\mathfrak{r}) \geq 2$.

There is an exact sequence

$$0 \rightarrow \text{Hom}_R(R/\mathfrak{r}, R) \rightarrow R \rightarrow \text{Hom}_R(\mathfrak{r}, R) \rightarrow \text{Ext}_R^1(R/\mathfrak{r}, R) \rightarrow 0.$$

The ring R has no embedded primes since R satisfies (S_1) . Let Q denote the total ring of fractions of R . we have $\mathfrak{r} \cdot Q = Q$ since $\text{ht}(\mathfrak{r}) \geq 2$ and R has no embedded primes. Hence we conclude that $\text{Hom}_R(R/\mathfrak{r}, R)$ vanishes and there is a canonical identification

$$\text{Hom}_R(\mathfrak{r}, R) = \{f \in Q \mid f \cdot \mathfrak{r} \subseteq R\}.$$

It suffices to show that

$$R = \{f \in Q \mid f \cdot \mathfrak{r} \subseteq R\}.$$

Let $f \in Q$ satisfying $f \cdot \mathfrak{r} \subseteq R$. There is a regular element $x \in R$ with $y := xf \in R$. Let us show $y \in xR$. Let $xR = \bigcap_{i=1}^n I_i$ be a minimal primary decomposition of xR . Since x is a regular

element and R satisfies (S_2) , the ring R/xR satisfies (S_1) and hence R/xR has no embedded primes. This fact implies the prime ideal $\sqrt{I_i}$ has height one. Since $\text{ht}(\mathfrak{r}) \geq 2$, we have

$$y \in yR_{\sqrt{I_i}} = y\mathfrak{r}R_{\sqrt{I_i}} = xR_{\sqrt{I_i}} \subseteq I_iR_{\sqrt{I_i}}.$$

As I_i is a $\sqrt{I_i}$ -primary ideal, we see $y \in I_i$. Thus we conclude that $y \in \bigcap_{i=1}^n I_i = xR$. \square

Lemma B.12. *Let $0 \rightarrow M \rightarrow P^1 \xrightarrow{\alpha} P^2$ be an exact sequence of R -modules, where P^1 and P^2 are free R -modules of finite rank. Put $I = \text{im}(P^1 \rightarrow P^2)$. If R satisfies (G_1) and (S_2) , then the canonical map*

$$\bigcap_R^r M \rightarrow \bigcap_R^r P^1 \xleftarrow{\sim} \bigwedge_R^r P^1$$

induces an isomorphism

$$\bigcap_R^r M \xrightarrow{\sim} \ker \left(\bigwedge_R^r P^1 \xrightarrow{\alpha} I \otimes_R \bigwedge_R^{r-1} P^1 \right).$$

for any integer $r \geq 1$.

Remark B.13. When R is Gorenstein, this lemma has already been proved in [12, Lemma B.2].

Proof. Since P^2 is a projective R -module, we have an isomorphism

$$\text{Ext}_R^1(I, R) \xrightarrow{\sim} \text{Ext}_R^2(P^2/I, R).$$

Hence, since P^1 is a projective R -module, by taking R -duals to the exact sequence $0 \rightarrow M \rightarrow P^1 \rightarrow I \rightarrow 0$, we obtain an exact sequence of R -modules

$$0 \rightarrow I^* \rightarrow (P^1)^* \rightarrow M^* \rightarrow \text{Ext}_R^2(P^2/I, R) \rightarrow 0.$$

Put $Y := \text{im}((P^1)^* \rightarrow M^*)$. This exact sequence gives rise to the exact sequence

$$I^* \otimes_R \bigwedge_R^{r-1} (P^1)^* \rightarrow \bigwedge_R^r (P^1)^* \rightarrow \bigwedge_R^r Y \rightarrow 0.$$

Passing to R -duals once again and using the canonical identification $\bigwedge_R^r P^1 = \bigcap_R^r P^1$, we get an exact sequence

$$0 \rightarrow \left(\bigwedge_R^r Y \right)^* \rightarrow \bigwedge_R^r P^1 \rightarrow \left(I^* \otimes_R \bigwedge_R^{r-1} (P^1)^* \right)^*.$$

Since I is a submodule of P^2 , the canonical map $I \rightarrow I^{**}$ is injective, and hence we have the commutative diagram

$$\begin{array}{ccc} \bigwedge_R^r P^1 & \xrightarrow{\alpha} & I \otimes_R \bigwedge_R^{r-1} P^1 \\ \downarrow & & \downarrow \\ \left(I^* \otimes_R \bigwedge_R^{r-1} (P^1)^* \right)^* & \xleftarrow{\cong} & I^{**} \otimes_R \bigcap_R^{r-1} P^1, \end{array}$$

where the bottom arrow is the composite map

$$\begin{aligned} I^{**} \otimes_R \bigcap_R^{r-1} P^1 &= \text{Hom}_R(I^*, R) \otimes_R \bigcap_R^{r-1} P^1 \\ &\xrightarrow{\sim} \text{Hom}_R \left(I^*, \bigcap_R^{r-1} P^1 \right) \\ &= \left(I^* \otimes_R \bigwedge_R^{r-1} (P^1)^* \right)^*. \end{aligned}$$

Thus we conclude

$$\left(\bigwedge_R^r Y \right)^* = \ker \left(\bigwedge_R^r P^1 \rightarrow I \otimes_R \bigwedge_R^{r-1} P^1 \right).$$

We remark that $\text{Ext}_R^2(P^2/I, R)$ is pseudo-null since R satisfies (G_1) . Hence the kernel and cokernel of the homomorphism

$$\bigwedge_R^r Y \longrightarrow \bigwedge_R^r M^*$$

are also pseudo-null. By taking R -duals to the homomorphism $\bigwedge_R^r Y \longrightarrow \bigwedge_R^r M^*$, Lemma B.11 shows that the map

$$\bigcap_R^r M \xrightarrow{\sim} \left(\bigwedge_R^r Y \right)^*$$

is an isomorphism, which completes the proof. \square

Lemma B.14. *Let $R \longrightarrow R'$ be a ring homomorphism of noetherian rings satisfying (G_1) and (S_2) . Let $C \in \bigcup_{n>0} D_{\text{perf}}^{[1, n]}(R)$. Then for any integer $r \geq 0$, there is a canonical homomorphism*

$$\bigcap_R^r H^1(C) \longrightarrow \bigcap_{R'}^r H^1(C \otimes_R^{\mathbb{L}} R')$$

Proof. This lemma follows from Lemma B.12. \square

Lemma B.15. *Let J be a directed poset and $\{R_j\}_{j \in J}$ an inverse system of noetherian rings satisfying (G_1) and (S_2) . Set $R := \varprojlim_{j \in J} R_j$ and let*

$$C = [\cdots \longrightarrow 0 \longrightarrow P^1 \longrightarrow P^2 \longrightarrow P^3 \longrightarrow \cdots] \in \bigcup_{n>0} D_{\text{perf}}^{[1, n]}(R)$$

such that P^i is a finitely generated projective R -module. If R is a noetherian ring satisfying (G_1) and (S_2) , then, for any integer $r \geq 0$, the canonical maps

$$\bigcap_R^r H^1(C) \longrightarrow \bigcap_{R_j}^r H^1(C \otimes_R^{\mathbb{L}} R_j)$$

for $j \in J$ induce an isomorphism

$$\bigcap_R^r H^1(C) \xrightarrow{\sim} \varprojlim_{j \in J} \bigcap_{R_j}^r H^1(C \otimes_R^{\mathbb{L}} R_j).$$

Proof. For $j \in J$ and $i \in \{1, 2\}$, put $P_j^i := P^i \otimes_R R_j$ and $I_j := \text{im}(P_j^1 \longrightarrow P_j^2)$. We also set $I := \text{im}(P^1 \longrightarrow P^2)$. Since P_j^1 is a free R_j -module, by Lemma B.12, we have

$$\begin{aligned} \varprojlim_{j \in J} \bigcap_R^r H^1(C \otimes_R^{\mathbb{L}} R_j) &= \varprojlim_{j \in J} \ker \left(\bigwedge_{R_j}^r P_j^1 \longrightarrow I_j \otimes_{R_j} \bigwedge_{R_j}^{r-1} P_j^1 \right) \\ &= \ker \left(\bigwedge_R^r P^1 \longrightarrow \left(\varprojlim_{j \in J} I_j \right) \otimes_R \bigwedge_R^{r-1} P^1 \right) \end{aligned}$$

Since I_j is a submodule of P_j^2 , the canonical map $I \longrightarrow \varprojlim_{j \in J} I_j$ is injective. This shows that

$$\ker \left(\bigwedge_R^r P^1 \longrightarrow I \otimes_R \bigwedge_R^{r-1} P^1 \right) = \ker \left(\bigwedge_R^r P^1 \longrightarrow \left(\varprojlim_{j \in J} I_j \right) \otimes_R \bigwedge_R^{r-1} P^1 \right)$$

and completes the proof by Lemma B.12. \square

APPENDIX C. CHARACTERISTIC IDEAL

Let R be a noetherian ring and M a finitely generated R -module. We write Q for the total ring of fractions of R .

Lemma C.1. *Suppose that R satisfies (G_0) and (S_1) . For an R -submodule N of M and an integer $r > 0$, the canonical homomorphism*

$$\bigcap_R^r N \longrightarrow \bigcap_R^r M$$

is injective.

Proof. By taking R -duals to the tautological exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$, we obtain an exact sequence of R -modules

$$0 \rightarrow (M/N)^* \rightarrow M^* \rightarrow N^* \rightarrow \text{Ext}_R^1(M/N, R).$$

Since R satisfies (G_0) and (S_1) , the ring Q is Gorenstein with $\dim(Q) = 0$. Hence the module $Q \otimes_R \text{Ext}_R^1(M/N, R) = \text{Ext}_Q^1(Q \otimes_R M/N, Q)$ vanishes, and we have a commutative diagram

$$\begin{array}{ccc} \bigcap_R^r N & \longrightarrow & \bigcap_R^r M \\ \downarrow & & \downarrow \\ Q \otimes_R \bigcap_R^r N & \longrightarrow & Q \otimes_R \bigcap_R^r M. \end{array}$$

As $\bigcap_R^r N = (\bigwedge_R^r N^*)^*$, any R -regular element is $\bigcap_R^r N$ -regular. Hence the left vertical arrow is injective, which completes the proof. \square

Remark C.2. When R satisfies (G_0) and (S_1) , for an ideal I of R , the canonical homomorphism $I^{**} \rightarrow R^{**} = R$ is injective by Lemma C.1. Therefore, in this case, one can regard I^{**} as an ideal of R and I as a submodule of I^{**} .

Definition C.3. Take an integer $r > 0$ and an exact sequence of R -modules

$$0 \rightarrow N \rightarrow R^r \rightarrow M \rightarrow 0.$$

We then define the characteristic ideal $\text{char}_R(M)$ of M by

$$\text{char}_R(M) := \text{im} \left(\bigcap_R^r N \rightarrow \bigcap_R^r R^r = R \right).$$

Remark C.4. If R' is a flat R -algebra, then we have $\text{char}_R(M)R' = \text{char}_{R'}(R' \otimes_R M)$.

Remark C.5. The ideal $\text{char}_R(M)$ is independent of the choice of the exact sequence $0 \rightarrow N \rightarrow R^r \rightarrow M \rightarrow 0$.

In fact, by Remark C.4, we may assume that R is a local noetherian ring. Then any free resolution of M is isomorphic to the direct sum of the minimal free resolution of M and a trivial complex. Hence it suffices to show that

$$\text{im} \left(\bigcap_R^r N \rightarrow \bigcap_R^r R^r = R \right) = \text{im} \left(\bigcap_R^{r+s} (N \oplus R^s) \rightarrow \bigcap_R^{r+s} R^{r+s} = R \right).$$

The homomorphism $(R^{r+s})^* = (R^r)^* \oplus (R^s)^* \rightarrow N^* \oplus (R^s)^* = (N \oplus R^s)^*$ induces a commutative diagram

$$\begin{array}{ccc} \bigwedge_R^r (R^r)^* \otimes \bigwedge_R^s (R^s)^* & \longrightarrow & \bigwedge_R^r (N)^* \otimes \bigwedge_R^s (R^s)^* \\ \downarrow = & & \downarrow \\ \bigwedge_R^{r+s} (R^{r+s})^* & \longrightarrow & \bigwedge_R^{r+s} (N \oplus R^s)^*. \end{array}$$

By taking R -duals to this commutative diagram, we obtain the following commutative diagram:

$$\begin{array}{ccc} \bigcap_R^{r+s}(N \oplus R^s) & \longrightarrow & \bigcap_R^{r+s}R^{r+s} \\ \downarrow & & \downarrow = \\ \bigcap_R^r N \otimes \bigcap_R^s R^s & \longrightarrow & \bigcap_R^r R^r \otimes \bigcap_R^s R^s. \end{array}$$

Hence $\text{im}(\bigcap_R^r N \longrightarrow \bigcap_R^r R^r = R) = \text{im}(\bigcap_R^{r+s}(N \oplus R^s) \longrightarrow \bigcap_R^{r+s} R^{r+s} = R)$.

Example C.6. For an ideal I of R , we have $\text{char}_R(R/I) = \text{im}(I^{**} \longrightarrow R^{**} = R)$.

Proposition C.7.

- (i) We have $\text{Fitt}_R^0(M) \subseteq \text{char}_R(M)$.
- (ii) If the projective dimension of M is at most 1, then $\text{Fitt}_R^0(M) = \text{char}_R(M)$.

Proof. Take an integer $r > 0$, an exact sequence of R -modules

$$0 \longrightarrow N \longrightarrow R^r \longrightarrow M \longrightarrow 0,$$

and a surjection $R^s \longrightarrow N$. Then we have a commutative diagram

$$\begin{array}{ccc} \bigwedge_R^r R^s & \longrightarrow & \bigwedge_R^r R^r = R \\ \downarrow & & \downarrow = \\ \bigcap_R^r N & \longrightarrow & \bigcap_R^r R^r = R. \end{array}$$

Hence we conclude

$$\text{Fitt}_R^0(M) = \text{im}\left(\bigwedge_R^r R^s \longrightarrow \bigwedge_R^r R^r = R\right) \subseteq \text{im}\left(\bigcap_R^r N \longrightarrow \bigcap_R^r R^r = R\right) = \text{char}_R(M).$$

Furthermore, if the projective dimension of M is at most 1, then N is a projective R -module. In this case, the canonical map $\bigwedge_R^r R^s \longrightarrow \bigwedge_R^r N = \bigcap_R^r N$ is surjective, and hence $\text{Fitt}_R^0(M) = \text{char}_R(M)$. \square

Proposition C.8. If R is a zero dimensional Gorenstein ring, then we have

$$\text{char}_R(M) = \text{Ann}_R(M).$$

Proof. Take an integer $r > 0$ and an exact sequence of R -modules

$$0 \longrightarrow N \longrightarrow R^r \longrightarrow M \longrightarrow 0.$$

Since the functor $(-)^*$ is exact, we have an exact sequence

$$M^* \otimes_R \bigwedge_R^{r-1}(R^r)^* \longrightarrow \bigwedge_R^r (R^r)^* \longrightarrow \bigwedge_R^r N^* \longrightarrow 0.$$

Note that $M = M^{**}$ by Matlis duality. By taking R -duals to this exact sequence, we obtain an exact sequence

$$0 \longrightarrow \bigcap_R^r N \longrightarrow R \longrightarrow \left(M^* \otimes_R \bigwedge_R^{r-1}(R^r)^*\right)^* = M^r.$$

The homomorphism $R \longrightarrow M^r$ corresponds to the homomorphism $R^r \longrightarrow M$ under the canonical isomorphism $\text{Hom}_R(R, M^r) \xrightarrow{\sim} \text{Hom}_R(R^r, M)$. Hence we conclude

$$\text{char}_R(M) = \text{Ann}_R(\text{im}(R \longrightarrow M^r)) = \text{Ann}_R(M).$$

\square

Remark C.9. In general, the characteristic ideal is not additive on the (split) short exact sequence of finitely generated R -modules. We give two counter-examples.

(i) Suppose that R is a zero dimensional Gorenstein ring. We then have

$$\text{char}_R(R/I \oplus R/J) = I \cap J$$

for any ideals I and J of R by Proposition C.8. Hence if $IJ \neq I \cap J$, we have

$$\text{char}_R(R/I \oplus R/J) \neq \text{char}_R(R/I)\text{char}_R(R/J).$$

(ii) Suppose that R is a noetherian local ring. Take a regular element $r \in R$ and an ideal I of R with $r \in I$. We then have an exact sequence of R -modules

$$0 \longrightarrow I/rR \longrightarrow R/rR \longrightarrow R/I \longrightarrow 0.$$

We note that $\text{char}_R(R/rR) = rR \cong R$ since r is a regular element. Therefore, if we have

$$\text{char}_R(R/I)\text{char}_R(I/rR) = \text{char}_R(R/rR) \cong R,$$

the ideal $\text{char}_R(R/I)$ is invertible. Since an invertible ideal is projective, the ideal $\text{char}_R(R/I)$ is principal. Hence we conclude that if $\text{char}_R(R/I)$ is not principal, then we have

$$\text{char}_R(R/rR) \neq \text{char}_R(R/I)\text{char}_R(I/rR).$$

We devote the rest of this appendix to proving the following important property of characteristic ideals.

Theorem C.10. *Let R be a noetherian ring satisfying (G_1) and (S_2) . Let M be a finitely generated R -module. Then for any R -submodule N of M , we have*

$$\text{char}_R(M) \subseteq \text{char}_R(N).$$

Lemma C.11. *Let R be a noetherian ring satisfying (G_0) and (S_1) . Let M be a finitely generated R -module. Then the homomorphisms*

$$\xi_{M^*}^1 : M^* \longrightarrow M^{***} \quad \text{and} \quad (\xi_M^1)^* : M^{***} \longrightarrow M^*$$

are isomorphisms.

Proof. The composite map

$$M^* \xrightarrow{\xi_{M^*}^1} M^{***} \xrightarrow{(\xi_M^1)^*} M^*$$

is the identity map. Hence it suffices to prove that the homomorphism $(\xi_M^1)^* : M^{***} \longrightarrow M^*$ is injective. To see this, we take an exact sequence $R^a \longrightarrow R^b \longrightarrow M \longrightarrow 0$ and set $C := \text{coker}(M^* \hookrightarrow (R^b)^*)$. We then have the commutative diagram with exact rows:

$$\begin{array}{ccccccc} R^a & \longrightarrow & R^b & \longrightarrow & M & \longrightarrow & 0 \\ \downarrow & & \downarrow \cong & & \downarrow & & \\ 0 & \longrightarrow & C^* & \longrightarrow & (R^b)^{**} & \longrightarrow & M^{**} \longrightarrow \text{Ext}_R^1(C, R) \longrightarrow 0, \end{array}$$

and it induces an exact sequence of R -modules

$$M \longrightarrow M^{**} \longrightarrow \text{Ext}_R^1(C, R) \longrightarrow 0.$$

Since $Q \otimes_R \text{Ext}_R^1(C, R) = \text{Ext}_Q^1(Q \otimes_R C, Q) = 0$, the module $\text{Ext}_R^1(C, R)^*$ also vanishes. In particular, the homomorphism $(\xi_M^1)^* : M^{***} \longrightarrow M^*$ is injective. \square

Proposition C.12. *Let R be a noetherian ring satisfying (G_0) and (S_1) . Let M be a finitely generated R -module. Then the characteristic ideal $\text{char}_R(M)$ is reflexive, that is, $\text{char}_R(M) = \text{char}_R(M)^{**}$.*

Proof. Take an exact sequence of R -modules

$$0 \longrightarrow N \longrightarrow R^r \longrightarrow M \longrightarrow 0$$

with $r > 0$. Then Lemmas C.1 shows that we have the canonical isomorphism

$$\bigcap_R^r N \xrightarrow{\sim} \text{char}_R(M).$$

Hence this proposition follows from Lemma C.11. \square

Lemma C.13. *Let R be a noetherian ring satisfying (G_0) and (S_2) . Let I and J be ideals of R . If $I_{\mathfrak{r}} \subseteq J_{\mathfrak{r}}$ for any prime \mathfrak{r} of R with $\text{ht}(\mathfrak{r}) \leq 1$, then we have $I^{**} \subseteq J^{**}$.*

Proof. By assumption, the R -module $(I + J)/J$ is pseudo-null. Hence $(I + J)^* \longrightarrow J^*$ is an isomorphism by Lemma B.11. This implies $J^{**} = (I + J)^{**} \supseteq I^{**}$. \square

Remark C.14. Let R be a noetherian ring satisfying (G_0) and (S_2) . By Proposition C.12 and Lemma C.13, we see that the ideal $\text{char}_R(M)$ is determined by ideals $\text{char}_{R_{\mathfrak{r}}}(M_{\mathfrak{r}})$ for any primes \mathfrak{r} of R with $\text{ht}(\mathfrak{r}) \leq 1$. Hence one can easily show the following properties of characteristic ideals:

- (i) If there is a pseudo-isomorphism $M \longrightarrow N$ of finitely generated R -modules, then we have $\text{char}_R(M) = \text{char}_R(N)$.
- (ii) When R is a normal ring, we have $\text{Fitt}_R^0(M)^{**} = \text{char}_R(M)$ for any finitely generated R -module, and hence, in this case, the notion of the characteristic ideal coincides with the usual one.

Proof of Theorem C.10. By Proposition C.12, the characteristic ideals $\text{char}_R(M)$ and $\text{char}_R(N)$ are reflexive. Hence, by Lemma C.13 and the condition (G_1) , we may assume that R is a Gorenstein local ring with $\dim(R) \leq 1$.

By the horseshoe lemma, a projective resolution of M can be built up inductively with the n -th item in the resolution of M equal to the direct sum of the n -th items in projective resolutions of N and M/N . Hence one can take the following commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & Y & \xrightarrow{\beta} & R^r & \longrightarrow & N & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & X & \xrightarrow{\alpha} & R^{r+s} & \longrightarrow & M & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & X/Y & \longrightarrow & R^s & \longrightarrow & M/N & \longrightarrow & 0. \end{array}$$

Here r is a positive integer and the homomorphism $R^r \hookrightarrow R^{r+s}$ is a split injection.

Since R is Gorenstein and $\dim(R) \leq 1$, the module $\text{Ext}_R^1(X/Y, R) \xrightarrow{\sim} \text{Ext}_R^2(M/N, R)$ vanishes. Hence by taking R -duals to the above commutative diagram, we get exact sequences of R -modules

$$\begin{array}{ccccccccc} 0 & \longrightarrow & (R^*)^s & \longrightarrow & (R^*)^{r+s} & \longrightarrow & (R^*)^r & \longrightarrow & 0 \\ & & \downarrow & & \downarrow \alpha^* & & \downarrow \beta^* & & \\ 0 & \longrightarrow & (X/Y)^* & \longrightarrow & X^* & \longrightarrow & Y^* & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ & & \text{Ext}_R^1(M/N, R) & \longrightarrow & \text{Ext}_R^1(M, R) & \longrightarrow & \text{Ext}_R^1(N, R) & \longrightarrow & 0. \end{array}$$

We fix a basis $\omega \in \bigwedge_R^s (R^*)^s$. We also denote by ω the images of ω in $\bigwedge_R^s (R^*)^{r+s}$ and $\bigwedge_R^s X^*$. Then the homomorphism

$$\bigwedge_R^r (R^*)^{r+s} \longrightarrow \bigwedge_R^{r+s} (R^*)^{r+s}; x \mapsto x \wedge \omega$$

induces an R -isomorphism

$$\bigwedge_R^r (R^*)^r \xrightarrow{\sim} \bigwedge_R^{r+s} (R^*)^{r+s}.$$

We also have the composite map

$$(12) \quad \bigwedge_R^r X^* \xrightarrow{x \mapsto x \wedge \omega} \bigwedge_R^{r+s} X^* \longrightarrow \left(\bigwedge_R^{r+s} X^* \right)^{**}.$$

The homomorphism $Q \otimes_R (R^*)^s \longrightarrow Q \otimes_R (X/Y)^*$ is surjective since $Q \otimes_R \text{Ext}_R^1(M/N, R)$ vanishes. Hence the composite map

$$(X/Y)^* \otimes_R \bigwedge_R^{r-1} X^* \longrightarrow \bigwedge_R^r X^* \longrightarrow Q \otimes_R \bigwedge_R^{r+s} X^*$$

is zero. The canonical homomorphism $\left(\bigwedge_R^{r+s} X^* \right)^{**} \longrightarrow Q \otimes_R \bigwedge_R^{r+s} X^*$ is injective, and hence the composite map

$$(X/Y)^* \otimes_R \bigwedge_R^{r-1} X^* \longrightarrow \bigwedge_R^r X^* \longrightarrow \left(\bigwedge_R^{r+s} X^* \right)^{**}$$

is also zero. Since the kernel of the surjection $\bigwedge_R^r X^* \longrightarrow \bigwedge_R^r Y^*$ coincides with the image of the homomorphism $(X/Y)^* \otimes_R \bigwedge_R^{r-1} X^* \longrightarrow \bigwedge_R^r X^*$, the homomorphism (12) induces an R -homomorphism

$$\bigwedge_R^r Y^* \longrightarrow \left(\bigwedge_R^{r+s} X^* \right)^{**}.$$

Hence, we have the following commutative diagram:

$$\begin{array}{ccccccc} & & \cong & & & & \\ & \swarrow & & \searrow & & & \\ \bigwedge_R^r (R^r)^* & \longleftarrow & \bigwedge_R^r (R^*)^{r+s} & \xrightarrow{x \mapsto x \wedge \omega} & \bigwedge_R^{r+s} (R^*)^{r+s} & \xrightarrow{\cong} & \left(\bigwedge_R^{r+s} (R^*)^{r+s} \right)^{**} \\ \downarrow \wedge^r \beta^* & & \downarrow \wedge^r \alpha^* & & \downarrow \wedge^{r+s} \alpha^* & & \downarrow (\wedge^{r+s} \alpha^*)^{**} \\ \bigwedge_R^r Y^* & \longleftarrow & \bigwedge_R^r X^* & \xrightarrow{x \mapsto x \wedge \omega} & \bigwedge_R^{r+s} X^* & \longrightarrow & \left(\bigwedge_R^{r+s} X^* \right)^{**} \\ & \searrow & & \swarrow & & & \end{array}$$

Therefore, we obtain a commutative diagram

$$\begin{array}{ccccc} \bigcap_R^{r+s} X & \xleftarrow{(\xi^1)^*} & \left(\bigwedge_R^{r+s} X^* \right)^{***} & \longrightarrow & \bigcap_R^r Y \\ \downarrow \bigcap^{r+s} \alpha & & \downarrow (\wedge^{r+s} \alpha^*)^{***} & & \downarrow \bigcap^{r+s} \beta \\ \bigcap_R^{r+s} R^{r+s} & \xleftarrow{\cong} & \left(\bigwedge_R^{r+s} (R^{r+s})^* \right)^{***} & \xrightarrow{\cong} & \bigcap_R^r R^r. \end{array}$$

By Lemma C.11, the homomorphism $(\bigwedge_R^{r+s} X^*)^{***} \rightarrow \bigcap_R^{r+s} X$ is an isomorphism, and hence this commutative diagram shows that

$$\begin{aligned} \text{char}_R(M) &= \text{im} \left(\bigcap_R^{r+s} X \longrightarrow \bigcap_R^{r+s} R^{r+s} = R \right) \\ &= \text{im} \left(\left(\bigwedge_R^{r+s} X^* \right)^{***} \longrightarrow \left(\bigwedge_R^{r+s} (R^{r+s})^* \right)^{***} = R \right) \\ &\subseteq \text{im} \left(\bigcap_R^r Y \longrightarrow \bigcap_R^r R^r = R \right) \\ &= \text{char}_R(N). \end{aligned}$$

□

REFERENCES

- [1] D. Burns, R. Sakamoto, T. Sano, On the theory of higher rank Euler, Kolyvagin and Stark systems, II, submitted for publication, arXiv:1805.08448.
- [2] D. Burns, R. Sakamoto, T. Sano, On the theory of higher rank Euler, Kolyvagin and Stark systems, III: applications, arXiv:1902.07002.
- [3] D. Burns, T. Sano, On the Theory of Higher Rank Euler, Kolyvagin and Stark Systems, to appear in Int. Math. Res. Not. <https://doi.org/10.1093/imrn/rnz103>
- [4] K. Büyükboduk, Kolyvagin systems of Stark units. J. Reine Angew. Math. 631 (2009), 85–107.
- [5] K. Büyükboduk, Stark units and the main conjectures for totally real fields. Compos. Math. 145 (2009), no. 5, 1163–1195.
- [6] K. Büyükboduk, On Euler systems of rank r and their Kolyvagin systems. Indiana Univ. Math. J. 59 (2010), no. 4, 1277–1332.
- [7] K. Büyükboduk, Stickelberger elements and Kolyvagin systems. Nagoya Math. J. 203 (2011), 123–173.
- [8] K. Büyükboduk, Λ -adic Kolyvagin systems. Int. Math. Res. Not. IMRN 2011, no. 14, 3141–3206.
- [9] K. Büyükboduk, A. Lei, Rank-two Euler systems for symmetric squares. Trans. Amer. Math. Soc. 372 (2019), no. 12, 8605–8619.
- [10] K. Büyükboduk, A. Lei, D. Loeffler, G. Venkat, Iwasawa theory for Rankin-Selberg products of p -nonordinary eigenforms, Algebra Number Theory 13 (2019), no. 4, 901–941.
- [11] K. Büyükboduk, A. Lei, G. Venkat, Iwasawa theory for Symmetric Square of non- p -ordinary eigenforms, preprint, arXiv:1807.11517v2.
- [12] K. Büyükboduk, R. Sakamoto, On the non-critical exceptional zeros of Katz p -adic L -functions for CM fields. arXiv:1904.01644.
- [13] C. L. Siegel, Über die Fourierschen Koeffizienten von Modulformen, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II 1970 1970 15–56.
- [14] P. Deligne, K. A. Ribet, Values of abelian L -functions at negative integers over totally real fields, Invent. Math. 59 (1980), no. 3, 227–286.
- [15] B. Ferrero, L. Washington, The Iwasawa invariant μ_p vanishes for abelian number fields, Ann. Math. 109 (1979), 377–395.
- [16] K. Iwasawa, On \mathbb{Z}_l -extensions of algebraic number fields, Ann. of Math. (2) 98 (1973), 246–326.
- [17] K. Iwasawa, On the μ -invariants of \mathbb{Z}_l -extensions, Number theory, algebraic geometry and commutative algebra, in honor of Yasuo Akizuki, pp. 1–11. Kinokuniya, Tokyo, 1973.
- [18] K. Kato, p -adic Hodge theory and values of zeta functions of modular forms. Cohomologies p -adiques et applications arithmétiques. III. Astérisque No. 295 (2004), ix, 117–290.
- [19] M. Kurihara, On the structure of ideal class groups of CM-fields, Kazuya Kato’s fiftieth birthday. Doc. Math. 2003, Extra Vol., 539–563.
- [20] M. Kurihara, Refined Iwasawa theory and Kolyvagin systems of Gauss sum type, Proc. Lond. Math. Soc. (3) 104 (2012), no. 4, 728–769.
- [21] M. Kurihara, Refined Iwasawa theory for p -adic representations and the structure of Selmer groups, Münster J. Math. 7 (2014), no. 1, 149–223.
- [22] A. Lei, D. Loeffler, S. Zerbes, Euler systems for Rankin-Selberg convolutions of modular forms. Ann. of Math. (2) 180 (2014), no. 2, 653–771.
- [23] D. Loeffler, C. Skinner, S. Zerbes, Euler systems for $\text{GSp}(4)$, arXiv:1706.00201v3.
- [24] D. Loeffler, S. Zerbes, Euler systems with local conditions, arXiv:1710.04956v2, to appear in proceedings of the Iwasawa 2017 conference.

- [25] B. Mazur, K. Rubin, Kolyvagin systems, *Mem. Amer. Math. Soc.* 799 (2004).
- [26] B. Mazur, K. Rubin, Controlling Selmer groups in the higher core rank case, *J. Th. Nombres Bordeaux* 28 (2016) 145–183.
- [27] H. Matsui, R. Takahashi, Y. Tsuchiya, When are n -syzygy modules n -torsionfree? *Arch. Math. (Basel)* 108 (2017), no. 4, 351–355.
- [28] J. Neukirch, A. Schmidt, K. Wingberg, *Cohomology of number fields*, Second edition, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 323. Springer-Verlag, Berlin, 2008. xvi+825 pp.
- [29] B. Perrin-Riou, Systèmes d’Euler p -adiques et théorie d’Iwasawa, *Ann. Inst. Fourier (Grenoble)* 48 (1998) 1231–1307.
- [30] J. Pottharst. Analytic families of finite-slope Selmer groups, *Algebra Number Theory* 7 (2013), no. 7, 1571–1612.
- [31] K. Rubin, A Stark Conjecture ‘over \mathbb{Z} ’ for abelian L -functions with multiple zeros, *Ann. Inst. Fourier* 46 (1996) 33–62.
- [32] K. Rubin, *Euler systems*, *Annals of Math. Studies* 147, Princeton Univ. Press, 2000.
- [33] R. Sakamoto, Stark systems over Gorenstein local rings, *Algebra Number Theory* 12 (2018), no. 10, 2295–2326.
- [34] R. Sakamoto, On the theory of higher rank Euler, Kolyvagin and Stark systems: a research announcement, to appear in *Algebraic Number Theory and Related Topics 2017*, RIMS Kôkyûroku Bessatsu.
- [35] J. Tate, *Les Conjectures de Stark sur les Fonctions L d’Artin en $s = 0$* (notes par D. Bernardi et N. Schappacher), *Progress in Math.*, 47, Birkhäuser, Boston, 1984.
- [36] A. Wiles, The Iwasawa conjecture for totally real fields, *Ann. of Math.* 131 (1990) 493–540.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN

E-mail address: sakamoto@ms.u-tokyo.ac.jp