

THE HODGE REALIZATION OF THE POLYLOGARITHM ON THE PRODUCT OF MULTIPLICATIVE GROUPS

KENICHI BANNAI^{*◊}, KEI HAGIHARA^{◊*}, KAZUKI YAMADA^{*}, AND SHUJI YAMAMOTO^{*◊}

ABSTRACT. The purpose of this article is to describe explicitly the polylogarithm class in absolute Hodge cohomology of a product of multiplicative groups, in terms of the Bloch-Wigner-Ramakrishnan polylogarithm functions. We will use the logarithmic Dolbeault complex defined by Burgos to calculate the corresponding absolute Hodge cohomology groups.

1. INTRODUCTION

The classical polylogarithm was constructed by Beilinson and Deligne as a class in the motivic cohomology of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$. The Hodge realization of this class gives an element in the corresponding absolute Hodge cohomology and may be described in terms of a unipotent variation of mixed \mathbb{R} -Hodge structures on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ whose periods are given by the classical polylogarithm functions (see [HW98]). Since $\mathbb{P}^1 \setminus \{0, 1, \infty\} = \mathbb{G}_m \setminus \{1\}$, we may view the classical polylogarithm as an object associated to the multiplicative group \mathbb{G}_m . In fact, the construction of the polylogarithm was extended to elliptic curves by Beilinson-Levin [BL94], to abelian schemes by Wildeshaus [Wil97] and Kings [Kin09], and more recently to general commutative group schemes by Huber and Kings [HK18]. The explicit description of the Hodge realization of the polylogarithm was given for elliptic curves by Beilinson-Levin [BL94] (see also [BKT10, Appendix]), and the description of the topological sheaf underlying the Hodge realization of the polylogarithm for general abelian varieties was given by Levin [Lev97] and Blottière [Blo09]. The purpose of this article is to describe directly the Hodge realization of the polylogarithm as an explicit class in absolute Hodge cohomology for the product of multiplicative groups.

For an integer $g > 0$, let $X = \mathbb{G}_m^g$ be the g -fold product of the multiplicative group defined over the field of complex numbers \mathbb{C} . We denote by $\mathbb{L}og$ the logarithm sheaf (see Definition 3.1), which is a certain admissible pro-unipotent variation of mixed \mathbb{R} -Hodge structures on X . The Hodge realization of the polylogarithm is defined as a class

$$\mathbf{pol} \in H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}og(g))$$

in the absolute Hodge cohomology $H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}og(g))$ of $U := X \setminus \{1\}$ with coefficients in $\mathbb{L}og(g)$, the g -fold Tate twist of $\mathbb{L}og$. The polylogarithm class is characterized as the class which maps to 1 through the residue map. In this article, we describe the polylogarithm class in terms of the Bloch-Wigner-Ramakrishnan polylogarithm functions defined by Ramakrishnan [Ram86] (see also [Zag90, §1]), which are single-valued real analytic variants of the classical polylogarithm functions.

Date: July 22, 2019 (Version 2.05).

2010 Mathematics Subject Classification. 14C30, 11G55.

This research was conducted as part of the KiPAS program FY2014–2018 of the Faculty of Science and Technology at Keio University. This research was supported in part by KAKENHI 26247004, 16J01911, 16K13742, 18H05233 as well as the JSPS Core-to-Core program “Foundation of a Global Research Cooperative Center in Mathematics focused on Number Theory and Geometry”.

Contrary to the case when $g = 1$, the classes in absolute Hodge cohomology of degree > 1 cannot be interpreted in terms of variations of mixed \mathbb{R} -Hodge structures on U . We will use the logarithmic Dolbeault complex defined by Burgos [Bur94] to construct an explicit complex yielding absolute Hodge cohomology. Then in our main result, Theorem 5.11, we will describe the polylogarithm class in terms of cocycles. We admit that our result is not very surprising, but our method illustrates the strength of the logarithmic Dolbeault complex in elegantly describing directly the polylogarithm class in absolute Hodge cohomology for higher dimensional cases.

Our method also applies to the case $g = 1$ and then recovers the following well-known result.

Corollary 1.1 (=Corollary 5.10). *Let $d > 1$ be an integer and let ζ be a primitive d -th root of unity. The inclusion $i_\zeta : \text{Spec } \mathbb{C} \rightarrow U := \text{Spec } \mathbb{C}[t, t^{-1}, (t-1)^{-1}]$ defined by $t \mapsto \zeta$ induces the restriction map*

$$i_\zeta^* : H_{\mathcal{A}}^1(U, \text{Log}(1)) \rightarrow H_{\mathcal{A}}^1(\text{Spec } \mathbb{C}, i_\zeta^* \text{Log}(1)) \cong \prod_{k=0}^{\infty} \mathbb{C}/(2\pi i)^{k+1} \mathbb{R}.$$

Then we have

$$i_\zeta^* \mathbf{pol} = ((-1)^k \text{Li}_{k+1}(\zeta))_{k=0}^{\infty} \in \prod_{k=0}^{\infty} \mathbb{C}/(2\pi i)^{k+1} \mathbb{R}.$$

Here, $\text{Li}_k(t)$ denotes the classical polylogarithm function, defined as the complex analytic continuation of the series

$$\text{Li}_k(t) := \sum_{n=1}^{\infty} \frac{t^n}{n^k}$$

on the open unit disc in the complex plane.

The arithmetic significance of our result for the case $g > 1$ will be treated in subsequent research.

CONTENTS

1. Introduction	1
2. Mixed Hodge Structure on Cohomology	3
3. Geometric Polylogarithm	6
4. Logarithmic Dolbeault Complex	15
5. Absolute Polylogarithm	21
References	27

The content of this article is as follows. In §2, we will review the theory of mixed \mathbb{R} -Hodge structures and the construction of mixed \mathbb{R} -Hodge structure on cohomology with coefficients in an admissible unipotent variation of mixed \mathbb{R} -Hodge structures. In §3, we will define the logarithm sheaf Log on $X = \mathbb{G}_m^g$, and describe the geometric polylogarithm class in terms of differential forms. In §4, we will introduce the logarithmic Dolbeault complex defined by Burgos [Bur94], which will be used to describe the absolute Hodge cohomology of $U = X \setminus \{1\}$. In §5, we will define the absolute polylogarithm class and prove our main result.

Acknowledgement. The authors express their sincere gratitude to the referee for carefully reading the manuscript and giving helpful suggestions. The authors would also like to thank the KiPAS program FY2014–2018 of the Faculty of Science and Technology at Keio University for providing an excellent environment making this research possible.

2. MIXED HODGE STRUCTURE ON COHOMOLOGY

In what follows, for a smooth scheme X of finite type over \mathbb{C} , we denote by \mathcal{O}_X the sheaf of holomorphic functions on $X(\mathbb{C})$, and by Ω_X^\bullet the complex of sheaves of holomorphic differential forms. In this section, we review the construction by Deligne [Del71] and Hain-Zucker [HZ87] of mixed \mathbb{R} -Hodge structures on cohomology of a complex variety with coefficients in an admissible unipotent variation of mixed \mathbb{R} -Hodge structures.

2.1. Construction of Mixed Hodge Structures on Cohomology. We first start with the definition of pure and mixed \mathbb{R} -Hodge structures.

Definition 2.1 (pure \mathbb{R} -Hodge structure). Let $V = (V, F^\bullet)$ be a pair consisting of a finite dimensional \mathbb{R} -vector space and a finite descending Hodge filtration by \mathbb{C} -linear subspaces F^\bullet on $V_{\mathbb{C}} := V \otimes \mathbb{C}$. We say that V is a *pure \mathbb{R} -Hodge structure of weight n* , if

$$V_{\mathbb{C}} = \bigoplus_{p+q=n} F^p V_{\mathbb{C}} \cap \overline{F^q} V_{\mathbb{C}},$$

where $\overline{F^q} V_{\mathbb{C}} := \overline{F^q V_{\mathbb{C}}}$ is the complex conjugate of $F^q V_{\mathbb{C}}$.

Example 2.2. For any integer $n \in \mathbb{Z}$, the Tate object $\mathbb{R}(n) := (V, F^\bullet)$, where $V_{\mathbb{C}} := \mathbb{C}\underline{\omega}^n$ for a basis $\underline{\omega}^n$ and $V := \mathbb{R}\underline{u}^n \subset V_{\mathbb{C}}$ for $\underline{u}^n := (2\pi i)^n \underline{\omega}^n$, with a filtration F^\bullet satisfying $F^{-n} V_{\mathbb{C}} = V_{\mathbb{C}}$ and $F^{-n+1} V_{\mathbb{C}} = 0$ is a pure \mathbb{R} -Hodge structure of weight $-2n$.

Definition 2.3 (mixed \mathbb{R} -Hodge structure). Let $V = (V, W_\bullet, F^\bullet)$ be a triple consisting of a finite dimensional \mathbb{R} -vector space V , a finite ascending filtration W_\bullet on V by \mathbb{R} -linear subspaces, and a finite descending filtration F^\bullet on $V_{\mathbb{C}}$ by \mathbb{C} -linear subspaces. We say that V is a *mixed \mathbb{R} -Hodge structure*, if the pair $\text{Gr}_n^W V := (\text{Gr}_n^W V, F^\bullet)$ for each $n \in \mathbb{Z}$ is a pure \mathbb{R} -Hodge structure of weight n , where $\text{Gr}_n^W V := W_n V / W_{n-1} V$ and F^\bullet is the filtration on $\text{Gr}_n^W V$ induced by F^\bullet .

We denote by $\text{MHS}_{\mathbb{R}}$ the category of mixed \mathbb{R} -Hodge structures, whose morphisms are \mathbb{R} -linear homomorphisms of the underlying \mathbb{R} -vector spaces compatible with the filtrations W_\bullet and F^\bullet .

Let X be a smooth algebraic variety of dimension g over \mathbb{C} , which we view as a complex manifold. The purpose of this section is to give the construction of a mixed \mathbb{R} -Hodge structure on the cohomology $H^m(X, \mathbb{R}) := H^m(X(\mathbb{C}), \mathbb{R})$. We first start with the construction of the weight filtration on $H^m(X, \mathbb{R})$. Let $j : X \hookrightarrow \overline{X}$ be a smooth and proper compactification of X such that the complement $D := \overline{X} \setminus X$ is a simple normal crossing divisor. Note that we have $H^m(X, \mathbb{R}) = \mathbb{H}^m(\overline{X}, Rj_* \mathbb{R})$. For any complex K^\bullet in an abelian category, let $\tau_{\leq \bullet}$ be the canonical filtration which is defined by

$$\tau_{\leq n}(K^m) = \begin{cases} 0 & n < m, \\ \text{Ker}(d^n) & n = m, \\ K^m & n > m. \end{cases}$$

We define the filtration \widetilde{W}_\bullet on $H^m(X, \mathbb{R})$ to be the filtration induced from the canonical filtration $\tau_{\leq \bullet}$ on $Rj_* \mathbb{R}$. In other words, we let

$$\widetilde{W}_n H^m(X, \mathbb{R}) := \text{Im}(\mathbb{H}^m(\overline{X}, \tau_{\leq n} Rj_* \mathbb{R}) \rightarrow H^m(X, \mathbb{R})).$$

We define the filtration $W_\bullet := \widetilde{W}_\bullet[m]$ on $H^m(X, \mathbb{R})$ by $W_n H^m(X, \mathbb{R}) := \widetilde{W}_{n-m} H^m(X, \mathbb{R})$.

In order to define the Hodge filtration F^\bullet on cohomology, we introduce the de Rham cohomology. For the complex of differentials Ω_X^\bullet on X , we define the de Rham cohomology of X by

$$H_{\text{dR}}^m(X) := \mathbb{H}^m(X, \Omega_X^\bullet).$$

Note that we have canonical isomorphisms $H^m(X, \mathbb{R}) \otimes \mathbb{C} = H^m(X, \mathbb{C}) \cong H_{\text{dR}}^m(X)$. As in [Del71, (3.1.2)], we let $\Omega_{\overline{X}}^\bullet(D)$ be the complex of sheaves of logarithmic differentials around D . By [Del71, (3.1.7)] (see Proposition 2.4 below), we have an isomorphism

$$(2.1) \quad H_{\text{dR}}^m(X) \cong \mathbb{H}^m(\overline{X}, \Omega_{\overline{X}}^\bullet(D)).$$

We define the weight and Hodge filtrations on de Rham cohomology. For any $x \in \overline{X}$, we say that a coordinate neighborhood $U \subset \overline{X}$ of x is *adapted to D* , if x has coordinates $(0, \dots, 0)$ and $U \cap D$ is defined by the equation $t_{\mu_1} \cdots t_{\mu_h} = 0$ for local coordinates t_1, \dots, t_g on U . For a fixed U adapted to D , we let $I := \{\mu_1, \dots, \mu_h\}$. Then following [Del71, (3.1.5)], we define the weight filtration W_\bullet on $\Omega_{\overline{X}}^\bullet(D)$ to be the multiplicative ascending filtration defined on each open neighborhood $U \subset \overline{X}$ adapted to D by giving weight 0 to the sections of $\Omega_{\overline{X}}^\bullet$ and weight 1 to the sections dt_μ/t_μ for $\mu \in I$. The Hodge filtration F^\bullet on $\Omega_{\overline{X}}^\bullet(D)$ is defined to be the stupid filtration

$$F^p \Omega_{\overline{X}}^m(D) := \begin{cases} \Omega_{\overline{X}}^m(D) & p \leq m, \\ 0 & p > m. \end{cases}$$

We define the filtrations \widetilde{W}_\bullet and F^\bullet on the de Rham cohomology $H_{\text{dR}}^m(X)$ to be the filtrations induced through the isomorphism (2.1) from the filtrations W_\bullet and F^\bullet on $\Omega_{\overline{X}}^\bullet(D)$. In other words, we let

$$\begin{aligned} \widetilde{W}_n H_{\text{dR}}^m(X) &:= \text{Im}(\mathbb{H}^m(\overline{X}, W_n \Omega_{\overline{X}}^\bullet(D))) \rightarrow H_{\text{dR}}^m(X) \\ F^p H_{\text{dR}}^m(X) &:= \text{Im}(\mathbb{H}^m(\overline{X}, F^p \Omega_{\overline{X}}^\bullet(D))) \rightarrow H_{\text{dR}}^m(X). \end{aligned}$$

By [Del71, (3.1.7)], we have the following result.

Proposition 2.4. *The resolution $\mathbb{C} \hookrightarrow \Omega_{\overline{X}}^\bullet$ and the quasi-isomorphism $\Omega_{\overline{X}}^\bullet(D) \hookrightarrow j_* \Omega_X^\bullet$ induce filtered quasi-isomorphisms*

$$(Rj_* \mathbb{C}, \tau_{\leq \bullet}) \rightarrow (j_* \Omega_X^\bullet, \tau_{\leq \bullet}) \leftarrow (\Omega_{\overline{X}}^\bullet(D), \tau_{\leq \bullet}) \rightarrow (\Omega_{\overline{X}}^\bullet(D), W_\bullet),$$

hence a quasi-isomorphism $\alpha : (Rj_* \mathbb{C}, \tau_{\leq \bullet}) \xrightarrow{\cong} (\Omega_{\overline{X}}^\bullet(D), W_\bullet)$ in the filtered derived category.

Proposition 2.4 implies that the filtration \widetilde{W}_\bullet on $H_{\text{dR}}^m(X)$ coincides with the filtration \widetilde{W}_\bullet on $H^m(X, \mathbb{R})$ through the isomorphism $H^m(X, \mathbb{R}) \otimes \mathbb{C} \cong H_{\text{dR}}^m(X)$. Again, we denote by W_\bullet the filtration $W_\bullet := \widetilde{W}_\bullet[m]$ on $H_{\text{dR}}^m(X)$. The filtrations W_\bullet and F^\bullet on $H_{\text{dR}}^m(X)$ induces the filtrations W_\bullet and F^\bullet on $H^m(X, \mathbb{R}) \otimes \mathbb{C}$.

Theorem 2.5 ([Del71, Théorème 3.2.5]). *For any $m \in \mathbb{Z}$, the triple*

$$H^m(X, \mathbb{R}) := (H^m(X, \mathbb{R}), W_\bullet, F^\bullet)$$

defined above is a mixed \mathbb{R} -Hodge structure.

2.2. Admissible Unipotent Variation of Mixed Hodge Structures. In this subsection, we review the definition of an admissible unipotent variation of mixed \mathbb{R} -Hodge structures on X , and give the construction by Hain and Zucker of the mixed \mathbb{R} -Hodge structure on its cohomology.

Definition 2.6. Let $\mathbb{V} = (\mathbb{V}, W_\bullet, F^\bullet)$ be a triple, where \mathbb{V} is an \mathbb{R} -local system on X , W_\bullet is a finite ascending filtration of \mathbb{V} by \mathbb{R} -local subsystems, and F^\bullet is a finite descending filtration on $\mathcal{V} := \mathbb{V} \otimes \mathcal{O}_X$ by coherent \mathcal{O}_X -submodules, satisfying the Griffiths transversality

$$\nabla(F^p \mathcal{V}) \subset F^{p-1} \mathcal{V} \otimes \Omega_X^1,$$

where ∇ is the integrable connection on \mathcal{V} defined by $\nabla := 1 \otimes d$. We say that \mathbb{V} is a *variation of mixed \mathbb{R} -Hodge structures* on X , if for any point $x \in X$ giving an inclusion $i_x : \text{Spec } \mathbb{C} \hookrightarrow X$, the triple $i_x^* \mathbb{V} := (i_x^* \mathbb{V}, W_\bullet, F^\bullet)$ is a mixed \mathbb{R} -Hodge structure, where W_\bullet is the ascending filtration by \mathbb{R} -linear subspaces on $i_x^* \mathbb{V}$ given by $W_n(i_x^* \mathbb{V}) := i_x^*(W_n \mathbb{V})$ for any $n \in \mathbb{Z}$, the filtration F^\bullet is the descending filtration by \mathbb{C} -linear subspaces on $i_x^* \mathbb{V} \otimes \mathbb{C}$ given by $F^p(i_x^* \mathbb{V} \otimes \mathbb{C}) := i_x^*(F^p \mathcal{V}) \otimes_{\mathcal{O}_x} \mathbb{C}$ for any $p \in \mathbb{Z}$.

Example 2.7. Suppose $V = (V, W_\bullet, F^\bullet)$ is a mixed \mathbb{R} -Hodge structure. The constant variation of mixed \mathbb{R} -Hodge structure V on X is defined as the triple

$$V_X := (V_X, W_\bullet, F^\bullet),$$

where V_X and W_\bullet are the constant \mathbb{R} -local systems on X given by V_X and W_\bullet , and the filtration F^\bullet on $\mathcal{V} := V_X \otimes \mathcal{O}_X$ is given by $F^p \mathcal{V} := (F^p V_{\mathbb{C}})_X \otimes \mathcal{O}_X$. In particular, if V is a pure \mathbb{R} -Hodge structure of weight n , then we say that V_X is a constant variation of pure \mathbb{R} -Hodge structure of weight n . We will often denote V_X simply by V if there is no fear of confusion.

Example 2.7 shows that the Tate object $\mathbb{R}(n)$ of Example 2.2 in $\text{MHS}_{\mathbb{R}}$ defines a variation of pure \mathbb{R} -Hodge structure of weight $-2n$ on X . We next define the notion of a unipotent variation of mixed \mathbb{R} -Hodge structures.

Definition 2.8. Let $\mathbb{V} = (\mathbb{V}, W_\bullet, F^\bullet)$ be a variation of mixed \mathbb{R} -Hodge structures on X . We say that \mathbb{V} is *unipotent*, if $\text{Gr}_n^W \mathbb{V} := (\text{Gr}_n^W \mathbb{V}, F^\bullet)$ is a constant variation of pure \mathbb{R} -Hodge structure of weight n , where F^\bullet is the filtration on $\text{Gr}_n^W \mathcal{V} = \text{Gr}_n^W \mathbb{V} \otimes \mathcal{O}_X$ induced by F^\bullet .

We next review the notion of admissibility of mixed Hodge structures defined in [HZ87, (1.5)], for the case of unipotent variation of mixed \mathbb{R} -Hodge structures. Let $X \hookrightarrow \bar{X}$ be a smooth compactification of X such that the complement $D := \bar{X} \setminus X$ is a normal crossing divisor. Let $\mathbb{V} = (\mathbb{V}, W_\bullet, F^\bullet)$ be a variation of mixed \mathbb{R} -Hodge structures on X . We let $\mathcal{V} := \mathbb{V} \otimes \mathcal{O}_X$, with integrable connection $\nabla := 1 \otimes d$. By [Del70, Proposition 5.2], there exists a canonical extension $\mathcal{V}(D)$ of \mathcal{V} , which is a certain locally free $\mathcal{O}_{\bar{X}}$ -module of finite type with integrable logarithmic connection along D . Since $W_n \mathcal{V} := W_n \mathbb{V} \otimes \mathcal{O}_X$ for $n \in \mathbb{Z}$ is an \mathcal{O}_X -submodule of \mathcal{V} compatible with the connection, the canonical extension of W_\bullet on \mathcal{V} defines a finite ascending filtration W_\bullet on $\mathcal{V}(D)$.

Definition 2.9. Let $\mathbb{V} = (\mathbb{V}, W_\bullet, F^\bullet)$ be a variation of mixed \mathbb{R} -Hodge structures on X . Let F^\bullet be a finite descending filtration on $\mathcal{V}(D)$ extending the filtration F^\bullet on \mathcal{V} , again satisfying the Griffiths transversality with respect to ∇ . This implies in particular that $F^p \mathcal{V} = F^p \mathcal{V}(D)|_X$ for any $p \in \mathbb{Z}$. We say that F^\bullet is *compatible with W_\bullet* , if for any $n \in \mathbb{Z}$, the filtration F^\bullet induced on $\text{Gr}_n^W \mathcal{V}(D)$ extends the filtration F^\bullet on $\text{Gr}_n^W \mathcal{V}$.

The admissibility of \mathbb{V} is defined as follows.

Definition 2.10. Suppose \mathbb{V} is a unipotent variation of mixed \mathbb{R} -Hodge structures on X . We say that \mathbb{V} is *admissible*, if it satisfies the following conditions.

- (1) Let $\mathcal{V}(D)$ be the canonical extension of \mathcal{V} to \bar{X} . The filtration F^\bullet of \mathcal{V} extends to a filtration F^\bullet of $\mathcal{V}(D)$ compatible with W_\bullet .
- (2) For any component D_i of D , the residue Res_{D_i} of ∇ along D_i defined in [Del70, (3.8.3)] satisfies $\text{Res}_{D_i}(W_n \mathcal{V}(D)|_{D_i}) \subset W_{n-2} \mathcal{V}(D)|_{D_i}$ for any $n \in \mathbb{Z}$.

In what follows, let $\mathbb{V} = (\mathbb{V}, W_\bullet, F^\bullet)$ be an admissible unipotent variation of mixed \mathbb{R} -Hodge structures on X . We let $\mathcal{V} := \mathbb{V} \otimes \mathcal{O}_X$ with connection $\nabla := 1 \otimes d$. Then the de Rham cohomology of X with coefficients in \mathcal{V} is defined by

$$H_{\text{dR}}^m(X, \mathcal{V}) := \mathbb{H}^m(X, \mathcal{V} \otimes \Omega_X^\bullet).$$

Note that we have canonical isomorphisms $H^m(X, \mathbb{V}) \otimes \mathbb{C} = H^m(X, \mathbb{V}_{\mathbb{C}}) \cong H_{\text{dR}}^m(X, \mathcal{V})$. If $\mathcal{V}(D)$ is the canonical extension of \mathcal{V} , then by [Del70, Théorème 6.2] and the definition of the canonical extension, we have an isomorphism

$$H_{\text{dR}}^m(X, \mathcal{V}) \cong \mathbb{H}^m(\bar{X}, \mathcal{V}(D) \otimes \Omega_{\bar{X}}^{\bullet}(D)).$$

We define the filtrations W_{\bullet} and F^{\bullet} on $\mathcal{V}(D) \otimes \Omega_{\bar{X}}^{\bullet}(D)$ by

$$\begin{aligned} W_n(\mathcal{V}(D) \otimes \Omega_{\bar{X}}^{\bullet}(D)) &:= \sum_{k \in \mathbb{Z}} W_{n-k} \mathcal{V}(D) \otimes W_k \Omega_{\bar{X}}^{\bullet}(D), \\ F^p(\mathcal{V}(D) \otimes \Omega_{\bar{X}}^{\bullet}(D)) &:= \sum_{q \in \mathbb{Z}} F^{p-q} \mathcal{V}(D) \otimes F^q \Omega_{\bar{X}}^{\bullet}(D). \end{aligned}$$

The following result is due to Hain and Zucker.

Proposition 2.11 ([HZ87, (8.6) Proposition]). *We denote by \tilde{W}_{\bullet} and F^{\bullet} the filtrations induced on $H_{\text{dR}}^m(X, \mathcal{V})$ from the filtrations W_{\bullet} and F^{\bullet} of $\mathcal{V}(D) \otimes \Omega_{\bar{X}}^{\bullet}(D)$, and we denote again by \tilde{W}_{\bullet} the filtration on $H^m(X, \mathbb{V}) \subset H_{\text{dR}}^m(X, \mathcal{V})$ induced from \tilde{W}_{\bullet} . Then the triple*

$$H^m(X, \mathbb{V}) := (H^m(X, \mathbb{V}), \tilde{W}_{\bullet}[m], F^{\bullet})$$

is a mixed \mathbb{R} -Hodge structure.

If $\mathbb{V} = (\mathbb{V}, W_{\bullet}, F^{\bullet})$ is an admissible unipotent variation of mixed \mathbb{R} -Hodge structures on X , then we let $\mathbb{V}(n) := \mathbb{V} \otimes \mathbb{R}(n)$ for any $n \in \mathbb{Z}$, that is, $\mathbb{V}(n) = \mathbb{V} \otimes \mathbb{R}\underline{u}^n$ with the filtrations given by $W_m(\mathbb{V}(n)) = (W_{m+2n} \mathbb{V}) \otimes \mathbb{R}\underline{u}^n$ and $F^p(\mathbb{V}(n)) = (F^{p+n} \mathbb{V}) \otimes \mathbb{C}\underline{u}^n$. Then we have $H^m(X, \mathbb{V}(n)) = H^m(X, \mathbb{V})(n)$ as mixed \mathbb{R} -Hodge structures.

3. GEOMETRIC POLYLOGARITHM

In what follows, we let g be a positive integer, and we let $X := \mathbb{G}_m^g = \text{Spec } \mathbb{C}[t_1^{\pm 1}, \dots, t_g^{\pm 1}]$ be the g -fold product of the multiplicative group over \mathbb{C} . The purpose of this section is to give the definition and an explicit construction of the geometric polylogarithm class of X . We start with the definition of the logarithm sheaf on X .

3.1. The Logarithm Sheaf. In this subsection, we define the logarithm sheaf $\mathcal{L}\text{og}$ of X . In what follows, we let $\mathbb{R}(1) := \mathbb{R}(1)^{\oplus g}$, which is a pure \mathbb{R} -Hodge structure of weight -2 . We denote by \mathbb{N} the set of natural numbers, and for any $\mathbf{k} = (k_{\mu}) \in \mathbb{N}^g$, we let $|\mathbf{k}| := k_1 + \dots + k_g$.

For an integer $N > 0$, consider the free \mathcal{O}_X -module

$$\mathcal{L}\text{og}^N := \prod_{\mathbf{k} \in \mathbb{N}^g, |\mathbf{k}| \leq N} \mathcal{O}_X \underline{\omega}^{\mathbf{k}}$$

with connection $\nabla: \mathcal{L}\text{og}^N \rightarrow \mathcal{L}\text{og}^N \otimes \Omega_X^1$ given by

$$(3.1) \quad \nabla(\underline{\omega}^{\mathbf{k}}) = \sum_{\mu=1}^g \underline{\omega}^{\mathbf{k}+1_{\mu}} \otimes \frac{dt_{\mu}}{t_{\mu}},$$

where 1_{μ} denotes the element in \mathbb{N}^g whose μ -th component is 1 and the other components are 0, and we let $\underline{\omega}^{\mathbf{k}} = 0$ if $|\mathbf{k}| > N$. We define the filtrations W_{\bullet} and F^{\bullet} on $\mathcal{L}\text{og}^N$ by

$$W_{-2m} \mathcal{L}\text{og}^N := W_{-2m+1} \mathcal{L}\text{og}^N := \prod_{m \leq |\mathbf{k}| \leq N} \mathcal{O}_X \underline{\omega}^{\mathbf{k}}, \quad F^{-p} \mathcal{L}\text{og}^N := \prod_{|\mathbf{k}| \leq p} \mathcal{O}_X \underline{\omega}^{\mathbf{k}}.$$

By choosing local branches of $\log t_\mu$, we define sections

$$(3.2) \quad \underline{u}^k := (2\pi i)^{|k|} \exp\left(\sum_{\mu=1}^g (-\log t_\mu)\omega_\mu\right) \cdot \underline{\omega}^k$$

of $\mathcal{L}\text{og}^N$, where ω_μ is an operator which acts on $\underline{\omega}^k$ by $\omega_\mu \cdot \underline{\omega}^k = \underline{\omega}^{k+1_\mu}$. We have

$$\begin{aligned} \nabla(\underline{u}^k) &= (2\pi i)^{|k|} \exp\left(\sum_{\nu=1}^g (-\log t_\nu)\omega_\nu\right) \cdot \sum_{\mu=1}^g \underline{\omega}^{k+1_\mu} \otimes \frac{dt_\mu}{t_\mu} \\ &\quad - (2\pi i)^{|k|} \sum_{\mu=1}^g \exp\left(\sum_{\nu=1}^g (-\log t_\nu)\omega_\nu\right) \omega_\mu \cdot \underline{\omega}^k \otimes \frac{dt_\mu}{t_\mu} = 0, \end{aligned}$$

hence \underline{u}^k gives a horizontal section of $\mathcal{L}\text{og}^N$ with respect to the connection ∇ . Since the transformation (3.2) is invertible, the sections \underline{u}^k are linearly independent over \mathbb{C} , hence span the space of horizontal sections of $\mathcal{L}\text{og}^N$.

We define the \mathbb{R} -local system $\mathbb{L}\text{og}^N$ by

$$\mathbb{L}\text{og}^N := \prod_{k \in \mathbb{N}^g, |k| \leq N} \mathbb{R}\underline{u}^k,$$

which gives an \mathbb{R} -structure on the \mathbb{C} -local system of horizontal sections of $\mathcal{L}\text{og}^N$. Although the basis \underline{u}^k depends on the branches of $\log t_\nu$ and does not give a global basis of $\mathbb{L}\text{og}^N$, the sheaf $\mathbb{L}\text{og}^N$ is independent of the branches of $\log t_\mu$.

We define the weight filtration on $\mathbb{L}\text{og}^N$ by

$$W_{-2m}\mathbb{L}\text{og}^N = W_{-2m+1}\mathbb{L}\text{og}^N := \prod_{m \leq |k| \leq N} \mathbb{R}\underline{u}^k,$$

which is compatible with the weight filtration W_\bullet on $\mathcal{L}\text{og}^N$ through the natural isomorphism

$$\mathbb{L}\text{og}^N \otimes_{\mathcal{O}_X} \cong \mathcal{L}\text{og}^N.$$

Then the triple $\mathbb{L}\text{og}^N = (\mathbb{L}\text{og}^N, W_\bullet, F^\bullet)$ is a variation of mixed \mathbb{R} -Hodge structures on X . By mapping the sections \underline{u}^k of $\mathbb{L}\text{og}^N$ to $\underline{u}_1^{k_1} \cdots \underline{u}_g^{k_g}$, where $(\underline{u}_1, \dots, \underline{u}_g)$ is the standard basis of $\mathbb{R}(1)$, we obtain an isomorphism

$$\text{Gr}_\bullet^W \mathbb{L}\text{og}^N \cong \prod_{k=0}^N \text{Sym}^k \mathbb{R}(1)$$

of variations of \mathbb{R} -Hodge structures on X , where we view $\text{Sym}^k \mathbb{R}(1)$ as a constant variation of pure \mathbb{R} -Hodge structures on X . We see from this fact that $\mathbb{L}\text{og}^N$ is a unipotent variation of mixed \mathbb{R} -Hodge structures.

The unipotent variation of mixed \mathbb{R} -Hodge structures $\mathbb{L}\text{og}^N$ is known to satisfy the *splitting principle*, given as follows. Let $d \geq 1$ and let $\zeta = (\zeta_1, \dots, \zeta_g)$ be an element in X whose components are d -th roots of unity, and we denote by $i_\zeta : \text{Spec } \mathbb{C} \hookrightarrow X$ the inclusion corresponding to ζ . By mapping the basis $\underline{\omega}^k$ of $i_\zeta^* \mathcal{L}\text{og}$ to $\underline{\omega}_1^{k_1} \cdots \underline{\omega}_g^{k_g}$, we obtain an isomorphism

$$(3.3) \quad i_\zeta^* \mathbb{L}\text{og}^N \cong \prod_{k=0}^N \text{Sym}^k \mathbb{R}(1)$$

of mixed \mathbb{R} -Hodge structures.

We let $\overline{X} := \mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ be the g -fold product of \mathbb{P}^1 . We have a natural inclusion $X \hookrightarrow \overline{X}$ with complement $D := \overline{X} \setminus X$. We denote by $\Omega_{\overline{X}}^{\bullet}(D)$ the complex of differentials on \overline{X} with logarithmic singularities along D . Note that $\mathcal{L}\text{og}^N$ extends naturally to a coherent $\mathcal{O}_{\overline{X}}$ -module

$$\mathcal{L}\text{og}^N(D) := \prod_{|k| \leq N} \mathcal{O}_{\overline{X}} \underline{\omega}^k,$$

and (3.1) defines a connection

$$\nabla: \mathcal{L}\text{og}^N(D) \rightarrow \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^1(D)$$

with logarithmic singularities along D . Then $\mathcal{L}\text{og}^N(D)$ is the canonical extension of $\mathcal{L}\text{og}^N$ to \overline{X} , and the filtrations W_{\bullet} and F^{\bullet} on $\mathcal{L}\text{og}^N$ naturally extend to $\mathcal{L}\text{og}^N(D)$. We see from the existence of such $\mathcal{L}\text{og}^N(D)$ that $\mathbb{L}\text{og}^N$ is an admissible unipotent variation of mixed \mathbb{R} -Hodge structures.

For integers $N > 0$, we have natural projections

$$(3.4) \quad \mathbb{L}\text{og}^N \rightarrow \mathbb{L}\text{og}^{N-1}.$$

This gives an exact sequence

$$(3.5) \quad 0 \rightarrow \text{Sym}^N \mathbb{R}(\mathbb{1}) \rightarrow \mathbb{L}\text{og}^N \rightarrow \mathbb{L}\text{og}^{N-1} \rightarrow 0$$

of variations of mixed \mathbb{R} -Hodge structures.

Definition 3.1. We define the *logarithm sheaf* $\mathbb{L}\text{og}$ to be the projective system with respect to (3.4) of admissible unipotent variations of mixed \mathbb{R} -Hodge structures $\mathbb{L}\text{og}^N$ on X .

We define the cohomology of X with coefficients in $\mathbb{L}\text{og}$ by

$$H^m(X, \mathbb{L}\text{og}) := \varprojlim_N H^m(X, \mathbb{L}\text{og}^N)$$

for any $m \in \mathbb{Z}$. We will calculate this cohomology in the next subsection.

3.2. Cohomology of the Logarithm Sheaf. In [HK18, Corollary 7.1.6], Huber and Kings calculated the cohomology of the logarithm sheaf for a general commutative group scheme given as an extension of an abelian variety by a torus. Applied to X , this gives the following.

Proposition 3.2. *We have*

$$H^m(X, \mathbb{L}\text{og}(g)) := \varprojlim_N H^m(X, \mathbb{L}\text{og}^N(g)) = \begin{cases} \mathbb{R} & m = g, \\ 0 & m \neq g. \end{cases}$$

Proposition 3.2 follows immediately from the following Lemma 3.3 concerning the cohomology of X with coefficients in $\mathbb{L}\text{og}^N$, a proof of which we give for the convenience of the reader.

Lemma 3.3. *For integers m and $N > 0$, the natural map*

$$H^m(X, \mathbb{L}\text{og}^N) \rightarrow H^m(X, \mathbb{L}\text{og}^{N-1})$$

is the zero map if $m \neq g$, and is an isomorphism if $m = g$. Moreover, we have

$$H^g(X, \mathbb{L}\text{og}^N) \xrightarrow{\cong} \cdots \xrightarrow{\cong} H^g(X, \mathbb{L}\text{og}^0) \cong \mathbb{R}(-g).$$

Proof. For $N > 0$, the exact sequence (3.5) gives rise to the long exact sequence

$$(3.6) \quad \begin{aligned} 0 &\rightarrow H^0(X, \mathrm{Sym}^N \mathbb{R}(1)) \rightarrow H^0(X, \mathbb{L}\mathrm{og}^N) \rightarrow H^0(X, \mathbb{L}\mathrm{og}^{N-1}) \\ &\rightarrow H^1(X, \mathrm{Sym}^N \mathbb{R}(1)) \rightarrow H^1(X, \mathbb{L}\mathrm{og}^N) \rightarrow H^1(X, \mathbb{L}\mathrm{og}^{N-1}) \\ &\rightarrow \dots \\ \dots &\rightarrow H^g(X, \mathrm{Sym}^N \mathbb{R}(1)) \rightarrow H^g(X, \mathbb{L}\mathrm{og}^N) \rightarrow H^g(X, \mathbb{L}\mathrm{og}^{N-1}) \rightarrow 0. \end{aligned}$$

In order to prove the statement for $m \neq g$, it suffices to show that the morphism

$$(3.7) \quad H^m(X, \mathrm{Sym}^N \mathbb{R}(1)) \rightarrow H^m(X, \mathbb{L}\mathrm{og}^N)$$

is surjective. We prove that the corresponding morphism for de Rham cohomology is surjective. We first let $\mathcal{H} := \mathbb{R}(1) \otimes \mathcal{O}_{\overline{X}}$ be the coherent $\mathcal{O}_{\overline{X}}$ -module on \overline{X} with connection $\nabla := 1 \otimes d$. Let $\mathbb{A}^g := \mathrm{Spec} \mathbb{C}[t_1, \dots, t_g] \subset \overline{X} = \mathbb{P}^1 \times \dots \times \mathbb{P}^1$, and we denote by Y the normal crossing divisor of \mathbb{A}^g defined by $t_1 \cdots t_g = 0$. Then the restriction to \mathbb{A}^g of \mathcal{H} and $\mathcal{L}\mathrm{og}^N(D)$ give coherent $\mathcal{O}_{\mathbb{A}^g}$ -modules with logarithmic connection along Y , and we have

$$\begin{aligned} H_{\mathrm{dR}}^m(X, \mathrm{Sym}^N \mathcal{H}) &\cong \mathbb{H}^m(\mathbb{A}^g, \mathrm{Sym}^N \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^\bullet(Y)) = H^m(\Gamma(\mathbb{A}^g, \mathrm{Sym}^N \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^\bullet(Y))) \\ H_{\mathrm{dR}}^m(X, \mathcal{L}\mathrm{og}^N) &\cong \mathbb{H}^m(\mathbb{A}^g, \mathcal{L}\mathrm{og}^N(D) \otimes \Omega_{\mathbb{A}^g}^\bullet(Y)) = H^m(\Gamma(\mathbb{A}^g, \mathcal{L}\mathrm{og}^N(D) \otimes \Omega_{\mathbb{A}^g}^\bullet(Y))), \end{aligned}$$

where the equalities on the right follow from the fact that \mathbb{A}^g is affine. This proves in particular that the de Rham cohomology is zero for $m > g$, hence it is sufficient to prove our assertion for non-negative integers $m < g$. Let $[\xi]$ be a class in $H_{\mathrm{dR}}^m(X, \mathcal{L}\mathrm{og}^N)$. Since \mathbb{A}^g is affine, the class $[\xi]$ is represented by a cocycle $\xi \in \Gamma(\mathbb{A}^g, \mathcal{L}\mathrm{og}^N(D) \otimes \Omega_{\mathbb{A}^g}^m(Y))$. Since $\mathcal{L}\mathrm{og}^N(D) \cong \bigoplus_{k=0}^N \mathrm{Sym}^k \mathcal{H}$ as $\mathcal{O}_{\mathbb{A}^g}$ -modules, we may write

$$\xi = \sum_{k=0}^N \xi_k$$

for $\xi_k \in \Gamma(\mathbb{A}^g, \mathrm{Sym}^k \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^m(Y))$. Then the cocycle condition $\nabla(\xi) = 0$ gives

$$\nabla(\xi) = \sum_{k=0}^N \nabla(\xi_k) = \sum_{k=0}^N \left((1 \otimes d)(\xi_k) + \sum_{\mu=1}^g \omega_\mu \cdot \xi_k \wedge \frac{dt_\mu}{t_\mu} \right) = 0.$$

This gives the differential equations

$$(1 \otimes d)(\xi_0) = 0, \quad (1 \otimes d)(\xi_k) + \sum_{\mu=1}^g \omega_\mu \cdot \xi_{k-1} \wedge \frac{dt_\mu}{t_\mu} = 0$$

for $k = 1, \dots, N$. We prove by induction on k that the class of ξ may be represented by an element satisfying $\xi_i = 0$ for $i < N$. Let $k \geq 0$ be an integer $< N$ and suppose $\xi_i = 0$ for $i < k$. Then the above differential equation becomes

$$(1 \otimes d)(\xi_k) = 0, \quad (1 \otimes d)(\xi_{k+1}) + \sum_{\mu=1}^g \omega_\mu \cdot \xi_k \wedge \frac{dt_\mu}{t_\mu} = 0.$$

The first differential equation shows that ξ_k is a cocycle representing an element in

$$H^m(\Gamma(\mathbb{A}^g, \mathrm{Sym}^k \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^\bullet(Y))) \cong H_{\mathrm{dR}}^m(X, \mathrm{Sym}^k \mathcal{H}).$$

The second differential equation shows that if we write

$$\xi_k = \sum_{\mu=\{\mu_1, \dots, \mu_m\} \subseteq \{1, \dots, g\}} u_\mu \frac{dt_{\mu_1}}{t_{\mu_1}} \wedge \dots \wedge \frac{dt_{\mu_m}}{t_{\mu_m}}$$

for $u_\mu \in \Gamma(\mathbb{A}^g, \text{Sym}^k \mathcal{H})$, the constant term of u_μ is zero. This shows that the class

$$[\xi_k] \in \mathbb{H}^m(\mathbb{A}^g, \text{Sym}^k \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^\bullet(Y))$$

is zero, which implies that there exists an element $\eta_k \in \Gamma(\mathbb{A}^g, \text{Sym}^k \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^{m-1}(Y))$ such that $(1 \otimes d)(\eta_k) = \xi_k$. If we let

$$\tilde{\xi}_{k+1} := \xi_{k+1} - \sum_{\mu=1}^g \omega_\mu \cdot \eta_k \wedge \frac{dt_\mu}{t_\mu} \in \Gamma(\mathbb{A}^g, \text{Sym}^{k+1} \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^m(Y)),$$

where ω_μ acts on $\underline{\omega}^m$ by $\omega_\mu \cdot \underline{\omega}^m = \underline{\omega}^{m+1\mu}$, and $\tilde{\xi}_j := \xi_j$ for $j = k+2, \dots, N$, then the class of

$$\tilde{\xi} = \sum_{j=k+1}^N \tilde{\xi}_j \in \Gamma(\mathbb{A}^g, \mathcal{L} \log^N \otimes \Omega_{\mathbb{A}^g}^m(Y))$$

coincides with that of ξ . By induction on k , we see that the class of ξ coincides with the class of a certain element $\xi_N \in \Gamma(\mathbb{A}^g, \text{Sym}^N \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^m(Y))$. Since ξ_N satisfies $(1 \otimes d)(\xi_N) = 0$, this element represents a class

$$[\xi_N] \in H^m(\Gamma(\mathbb{A}^g, \text{Sym}^N \mathcal{H} \otimes \Omega_{\mathbb{A}^g}^\bullet(Y))) \cong H_{\text{dR}}^m(X, \text{Sym}^N \mathcal{H}).$$

By construction, $[\xi_N]$ maps to the class $[\xi]$ through the homomorphism (3.7). This gives our statement for the case $m \neq g$. As a consequence, the exact sequence (3.6) gives an isomorphism

$$(3.8) \quad H^0(X, \text{Sym}^N \mathbb{R}(1)) \xrightarrow{\cong} H^0(X, \mathbb{L} \log^N),$$

short exact sequences

$$(3.9) \quad 0 \rightarrow H^{m-1}(X, \mathbb{L} \log^{N-1}) \rightarrow H^m(X, \text{Sym}^N \mathbb{R}(1)) \rightarrow H^m(X, \mathbb{L} \log^N) \rightarrow 0$$

for $m = 1, \dots, g-1$, and the exact sequence

$$(3.10) \quad 0 \rightarrow H^{g-1}(X, \mathbb{L} \log^{N-1}) \rightarrow H^g(X, \text{Sym}^N \mathbb{R}(1)) \rightarrow H^g(X, \mathbb{L} \log^N) \rightarrow H^g(X, \mathbb{L} \log^{N-1}) \rightarrow 0.$$

Next, we prove our assertion for the case $m = g$. From (3.10), it is sufficient to prove that the dimensions of $H^{g-1}(X, \mathbb{L} \log^{N-1})$ and $H^g(X, \text{Sym}^N \mathbb{R}(1))$ coincide. Note that since $H^g(X, \mathbb{R}) = H^g(\mathbb{G}_m^g, \mathbb{R}) \cong \mathbb{R}(-g)$, we have

$$\dim_{\mathbb{R}} H^g(X, \text{Sym}^N \mathbb{R}(1)) = \dim_{\mathbb{R}}(\text{Sym}^N \mathbb{R}(1)) = \binom{N+g-1}{g-1}.$$

We prove the equality

$$(3.11) \quad \dim_{\mathbb{R}} H^m(X, \mathbb{L} \log^N) = \binom{N+g-1-m}{g-1-m} \binom{N+g}{m}$$

for any integers $m = 0, \dots, g-1$ and $N \geq 0$.

If $N = 0$, then $\mathbb{L} \log^0$ is the constant \mathbb{R} -Hodge structure $\mathbb{R}(0)$ on X , hence

$$(3.12) \quad H^m(X, \mathbb{L} \log^0) = H^m(\mathbb{G}_m^g, \mathbb{R}) = \bigwedge^m H^1(\mathbb{G}_m^g, \mathbb{R}) \cong \bigwedge^m \mathbb{R}(-1)^{\oplus g} \cong \mathbb{R}(-m)^{\oplus \binom{g}{m}},$$

which gives (3.11) in this case. If $m = 0$, then by (3.8), we have

$$\dim_{\mathbb{R}} H^0(X, \mathbb{L} \log^N) = \dim_{\mathbb{R}} H^0(X, \text{Sym}^N \mathbb{R}(1)) = \dim_{\mathbb{R}} \text{Sym}^N \mathbb{R}(1) = \binom{N+g-1}{g-1},$$

which gives (3.11) in this case. Finally, suppose $m, N > 0$ and (3.11) is true for the pair $(m-1, N-1)$. Then the exact sequence (3.9) shows that we have

$$\begin{aligned} \dim_{\mathbb{R}} H^m(X, \mathbb{L}\text{og}^N) &= \dim_{\mathbb{R}} H^m(X, \text{Sym}^N \mathbb{R}(\mathbb{1})) - \dim_{\mathbb{R}} H^{m-1}(X, \mathbb{L}\text{og}^{N-1}) \\ &= \dim_{\mathbb{R}} \left(\left(\bigwedge^m H^1(X) \right) \otimes \text{Sym}^N \mathbb{R}(\mathbb{1}) \right) - \dim_{\mathbb{R}} H^{m-1}(X, \mathbb{L}\text{og}^{N-1}) \\ &= \binom{g}{m} \binom{N+g-1}{g-1} - \binom{N+g-m-1}{g-m} \binom{N+g-1}{m-1} = \binom{N+g-m-1}{g-m-1} \binom{N+g}{m}, \end{aligned}$$

which proves (3.11) for (m, N) as desired.

The last statement of our lemma follows from (3.12) for the case $m = g$. \square

3.3. The Geometric Polylogarithm Class. In this subsection, we give the definition of the geometric polylogarithm class of X . We let $Z := \{1\} \subset X$ and $U := X \setminus Z$. Then we have the following.

Proposition 3.4. *For $g > 1$, we have canonical isomorphisms*

$$H^m(U, \mathbb{L}\text{og}(g)) := \varprojlim_N H^m(U, \mathbb{L}\text{og}^N(g)) \cong \begin{cases} \mathbb{R}(0) & m = g, \\ \prod_{k=0}^{\infty} \text{Sym}^k \mathbb{R}(\mathbb{1}) & m = 2g-1, \\ 0 & \text{otherwise.} \end{cases}$$

For $g = 1$, we have canonical isomorphisms

$$H^m(U, \mathbb{L}\text{og}(1)) \cong \begin{cases} \mathbb{R}(0) \oplus \prod_{k=0}^{\infty} \mathbb{R}(k) & m = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 3.4 follows immediately from the following Lemma 3.5 concerning the cohomology of U with coefficients in $\mathbb{L}\text{og}^N$ and Lemma 3.3.

Lemma 3.5. *For $g > 1$, we have canonical isomorphisms*

$$H^m(U, \mathbb{L}\text{og}^N(g)) \cong \begin{cases} H^m(X, \mathbb{L}\text{og}^N(g)) & m = 0, \dots, g, \\ \prod_{k=0}^N \text{Sym}^k \mathbb{R}(\mathbb{1}) & m = 2g-1, \\ 0 & \text{otherwise.} \end{cases}$$

For $g = 1$, we have canonical isomorphisms

$$H^m(U, \mathbb{L}\text{og}^N(1)) \cong \begin{cases} \mathbb{R}(0) \oplus \prod_{k=0}^N \mathbb{R}(k) & m = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If we denote by $i : Z \hookrightarrow X$ the natural inclusion. then for any integer $N \geq 0$, we have a long exact sequence of mixed \mathbb{R} -Hodge structures

$$(3.13) \quad \cdots \rightarrow H^m(X, \mathbb{L}\text{og}^N(g)) \rightarrow H^m(U, \mathbb{L}\text{og}^N(g)) \xrightarrow{\text{res}_1} H^{m+1-2g}(Z, i^* \mathbb{L}\text{og}^N) \rightarrow \cdots .$$

Since $i^* \mathbb{L}\text{og}^N = \prod_{k=0}^N \text{Sym}^k \mathbb{R}(\mathbb{1})$, we have

$$H^m(Z, i^* \mathbb{L}\text{og}^N) \cong \begin{cases} \prod_{k=0}^N \text{Sym}^k \mathbb{R}(\mathbb{1}) & m = 0, \\ 0 & m \neq 0. \end{cases}$$

Since the m -th cohomology of $\mathbb{L}\text{og}^N(g)$ on X is zero for $m > g$, the long exact sequence (3.13) gives the desired result. For the case $g = 1$, we have the short exact sequence

$$0 \rightarrow H^1(X, \mathbb{L}\text{og}^N(1)) \rightarrow H^1(U, \mathbb{L}\text{og}^N(1)) \rightarrow H^0(Z, i^*\mathbb{L}\text{og}^N) \rightarrow 0.$$

By the diagram

$$\begin{array}{ccc} H^1(X, \mathbb{L}\text{og}^N(1)) & \longrightarrow & H^1(U, \mathbb{L}\text{og}^N(1)) \\ \downarrow \cong & & \downarrow \\ H^1(X, \mathbb{L}\text{og}^0(1)) & \longrightarrow & H^1(U, \mathbb{L}\text{og}^0(1)) \\ & \searrow \text{res}_0 \cong & \downarrow \text{res}_0 \\ & & \mathbb{R}(0), \end{array}$$

where the left vertical isomorphism follows from Lemma 3.3, we see that there exists a natural splitting $H^1(U, \mathbb{L}\text{og}^N(1)) \rightarrow H^1(X, \mathbb{L}\text{og}^N(1))$ of the first injection. Our assertion follows from Proposition 3.2. \square

Definition 3.6. We define the *geometric polylogarithm class* to be the class

$$\xi = (\xi_N) \in H^{2g-1}(U, \mathbb{L}\text{og}(g))$$

which maps to the basis \underline{u}^0 of $\text{Sym}^0\mathbb{R}(\mathbb{1})$ through the isomorphism of Proposition 3.4.

Remark 3.7. The geometric polylogarithm class $\xi = (\xi_N)$ satisfies

$$\xi_N \in W_0 H^{2g-1}(U, \mathbb{L}\text{og}^N(g)) \cap (F^0 \cap W_0) H^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)),$$

since the residue map is a morphism of mixed \mathbb{R} -Hodge structures. Hence we may view this class as an element in

$$\text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^{2g-1}(U, \mathbb{L}\text{og}(g))) := \varprojlim_N \left(\text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^{2g-1}(U, \mathbb{L}\text{og}^N(g))) \right).$$

3.4. Explicit Construction of the Geometric Polylogarithm. In this subsection, we give an explicit construction of the geometric polylogarithm class. We first define certain complexes to explicitly calculate the mixed \mathbb{R} -Hodge structure on the cohomology of $U := X \setminus Z$ for $Z = \{1\} \subset X$, and we give an explicit description of the residue map res_1 of (3.13). We let $[g] := \{1, \dots, g\}$, and for any $\mu \in [g]$, we let Z_{μ} be the divisor

$$Z_{\mu} := \mathbb{P}^1 \times \cdots \times \mathbb{P}^1 \times \{1\} \times \mathbb{P}^1 \times \cdots \times \mathbb{P}^1 \subset \overline{X}$$

with $\{1\}$ in the μ -th component and \mathbb{P}^1 in the other components. For any $I \subset [g]$, we let $D^I := D \cup \bigcup_{\mu \in I} Z_{\mu}$. In what follows, let N be an integer ≥ 0 .

Definition 3.8. We define the Čech-de Rham complex $\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^{\bullet}(D^{\bullet})$ on \overline{X} to be the simple complex associated to the double complex

$$\bigoplus_{|I|=1} \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^{\bullet}(D^I) \xrightarrow{\partial} \cdots \xrightarrow{\partial} \bigoplus_{|I|=g} \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^{\bullet}(D^I),$$

with $\bigoplus_{|I|=1} \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^0(D^I)$ in degree 0, where the horizontal differentials are the standard alternating sum $(\partial u)_{\mu_1 \cdots \mu_h} = \sum_{\nu=1}^h (-1)^{\nu} u_{\mu_1 \cdots \check{\mu}_{\nu} \cdots \mu_h}$.

Proposition 3.9. *Denote by $j : U \hookrightarrow \overline{X}$ be the natural inclusion, and consider the \mathbb{C} -local system $\mathbb{L}\text{og}_{\mathbb{C}}^N := \mathbb{L}\text{og}^N \otimes \mathbb{C}$ on U . Then we have canonical isomorphisms*

$$Rj_* \mathbb{L}\text{og}_{\mathbb{C}}^N \cong Rj_*(\mathcal{L}\text{og}^N \otimes \Omega_U^\bullet) \cong \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)$$

in the derived category of abelian sheaves on \overline{X} . In particular, by taking the functor $R\Gamma(\overline{X}, -)$, we have canonical isomorphisms

$$(3.14) \quad H^m(U, \mathbb{L}\text{og}_{\mathbb{C}}^N) \otimes \mathbb{C} = H^m(U, \mathbb{L}\text{og}_{\mathbb{C}}^N) \cong H_{\text{dR}}^m(U, \mathcal{L}\text{og}^N) \cong H^m(\overline{X}, \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)).$$

Proof. Since $\mathcal{L}\text{og}^N = \mathbb{L}\text{og}_{\mathbb{C}}^N \otimes \mathcal{O}_X$, the complex $\mathcal{L}\text{og}^N \otimes \Omega_U^\bullet$ on U gives a resolution

$$(3.15) \quad 0 \rightarrow \mathbb{L}\text{og}_{\mathbb{C}}^N \rightarrow \mathcal{L}\text{og}^N \otimes \mathcal{O}_U \xrightarrow{\nabla} \mathcal{L}\text{og}^N \otimes \Omega_U^1 \rightarrow \cdots$$

of $\mathbb{L}\text{og}_{\mathbb{C}}^N$, which proves the first isomorphism of our assertion. For any $\mu \in [g]$, we let $U_\mu := X \setminus Z_\mu = \overline{X} \setminus (D \cup Z_\mu)$. Then $\{U_\mu\}$ is an open covering of U . For any $I \subset [g]$, we let $U_I := \bigcap_{\mu \in I} U_\mu = \overline{X} \setminus D^I$ and we denote by $j_I^\circ : U_I \hookrightarrow U$ and $j_I := j \circ j_I^\circ : U_I \hookrightarrow \overline{X}$ the natural inclusions. If we let $j_{\bullet*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet)$ be the Čech complex defined as the simple complex associated to the double complex

$$\prod_{|I|=1} j_{I*}^\circ(\mathcal{L}\text{og}^N \otimes \Omega_{U_I}^\bullet) \rightarrow \cdots \rightarrow \prod_{|I|=g} j_{I*}^\circ(\mathcal{L}\text{og}^N \otimes \Omega_{U_I}^\bullet),$$

then we have a quasi-isomorphism

$$\mathcal{L}\text{og}^N \otimes \Omega_U^\bullet \xrightarrow{\cong} j_{\bullet*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet).$$

This quasi-isomorphism combined with (3.15) gives the isomorphism

$$(3.16) \quad Rj_* \mathbb{L}\text{og}_{\mathbb{C}}^N \xrightarrow{\cong} Rj_{\bullet*} j_{\bullet*}^\circ(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet) \cong j_{\bullet*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet)$$

in the derived category, where we let $j_{\bullet*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet) := j_* j_{\bullet*}^\circ(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet)$. The second isomorphism of (3.16) follows from the fact that j_I is an affine morphism for any $I \subset [g]$. Note that since $\mathcal{L}\text{og}^N(D)$ is the canonical extension of $\mathcal{L}\text{og}^N$, we have quasi-isomorphisms

$$\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I) \xrightarrow{\cong} j_{I*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_I}^\bullet)$$

for any $I \subset [g]$, which induce the quasi-isomorphism

$$(3.17) \quad \mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet) \xrightarrow{\cong} j_{\bullet*}(\mathcal{L}\text{og}^N \otimes \Omega_{U_\bullet}^\bullet).$$

Our assertion follows by combining the quasi-isomorphisms (3.16) and (3.17). \square

We next define the filtrations W_\bullet and F^\bullet on $\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)$ which gives the weight and Hodge filtrations of the mixed \mathbb{R} -Hodge structure $H^m(U, \mathbb{L}\text{og}^N)$. For any integer $n \in \mathbb{Z}$, we let $W_n(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet))$ be the subcomplex of $\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)$ defined as the simple complex associated to the double complex

$$\bigoplus_{|I|=1} W_n(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)) \rightarrow \cdots \rightarrow \bigoplus_{|I|=g} W_{n+g-1}(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)),$$

and for any integer $p \in \mathbb{Z}$, we let $F^p(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet))$ be the subcomplex of $\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)$ defined as the simple complex associated to the double complex

$$\bigoplus_{|I|=1} F^p(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)) \rightarrow \cdots \rightarrow \bigoplus_{|I|=g} F^p(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)).$$

Then the filtrations \widetilde{W}_\bullet and F^\bullet induced on $\mathbb{H}^m(\overline{X}, \mathcal{L}og^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet))$ from the filtrations W_\bullet and F^\bullet on $\mathcal{L}og^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)$ coincide through the isomorphism (3.14) with the weight and Hodge filtrations on the mixed \mathbb{R} -Hodge structure $H^m(U, \mathbb{L}og^N)$ of Proposition 2.11.

We now describe the residue map res_1 of (3.13). Consider $\mathbb{L}og^N(g) = \mathbb{L}og^N \otimes \mathbb{R}(g)$ and $\mathcal{L}og^N(D)(g) = \mathcal{L}og^N(D) \otimes \mathbb{C}(g)$, and let $\underline{\omega}^k(g) := \underline{\omega}^k \otimes \underline{\omega}^g$. Note that $Z = Z_1 \cap \cdots \cap Z_g$ and we let $i : Z \hookrightarrow \overline{X}$ be the natural inclusion. Consider the map

$$\mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^g(D^{[g]}) \rightarrow i_*(i^* \mathcal{L}og^N(D) \otimes \Omega_Z^0)$$

given locally in a neighborhood of Z by

$$f \otimes \underline{\omega}^g \otimes \frac{dt_1}{t_1 - 1} \wedge \cdots \wedge \frac{dt_g}{t_g - 1} \mapsto f|_Z \otimes 1,$$

where t_1, \dots, t_g is the standard parameter on $\overline{X} = \mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ so that the divisor Z_μ is defined by $t_\mu = 1$. This map induces a map of complexes

$$\mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet) \rightarrow i_*(i^* \mathcal{L}og^N(D) \otimes \Omega_Z^\bullet)[-2g + 1],$$

which induces a map on cohomology

$$\mathbb{H}^{2g-1}(\overline{X}, \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)) \rightarrow \mathbb{H}^0(Z, i^* \mathcal{L}og^N(D) \otimes \Omega_Z^\bullet).$$

Then through the isomorphism of Proposition 3.9, we obtain the map

$$\text{res}_1 : H^{2g-1}(U, \mathbb{L}og_{\mathbb{C}}^N(g)) \rightarrow H^0(Z, i^* \mathbb{L}og_{\mathbb{C}}^N).$$

The map coincides with the residue map res_1 of (3.13).

Note that the inclusion induces a natural morphism of complexes

$$\mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^g(D^{[g]})[-2g + 1] \rightarrow \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet),$$

which induces a map

$$\Gamma(\overline{X}, \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^g(D^{[g]})) \rightarrow \mathbb{H}^{2g-1}(\overline{X}, \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)).$$

Proposition 3.10. *The sections*

$$(3.18) \quad \xi_N := \underline{\omega}^0(g) \otimes \frac{dt_1}{t_1 - 1} \wedge \cdots \wedge \frac{dt_g}{t_g - 1} \in \Gamma(\overline{X}, \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^g(D^{[g]}))$$

for integers $N \geq 0$ define classes

$$[\xi_N] \in \mathbb{H}^{2g-1}(\overline{X}, \mathcal{L}og^N(D)(g) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet)) \cong H^{2g-1}(U, \mathbb{L}og_{\mathbb{C}}^N)$$

which are mapped by res_1 to the elements $\underline{\omega}^0 = \underline{u}^0$ of $H^0(Z, i^* \mathbb{L}og^N) = \prod_{k=0}^N \text{Sym}^k \mathbb{R}(\mathbb{1})$, and are mapped to zero by res_0 in the case $g = 1$. In other words, $\xi := ([\xi_N])_N$ is the geometric polylogarithm class.

The fact that the residue maps preserve the real structure of mixed \mathbb{R} -Hodge structures implies that $[\xi_N]$ is a class in $H^{2g-1}(U, \mathbb{L}og^N(g)) \subset H^{2g-1}(U, \mathbb{L}og_{\mathbb{C}}^N(g))$.

4. LOGARITHMIC DOLBEAULT COMPLEX

In this section, we will use the theory of logarithmic Dolbeault complex defined by Burgos to construct an explicit complex calculating the mixed \mathbb{R} -Hodge structure on the cohomology of U with coefficients in $\mathbb{L}\text{og}$. For a general complex manifold X , we denote by $\mathcal{A}_{X,\mathbb{R}}$ (resp. $\mathcal{E}_{X,\mathbb{R}}$) the sheaf of real-valued real analytic (resp. C^∞) functions on X , and by $\mathcal{A}_{X,\mathbb{R}}^\bullet$ (resp. $\mathcal{E}_{X,\mathbb{R}}^\bullet$) the corresponding complex of sheaves of differential forms.

4.1. Review of Logarithmic Dolbeault Complex. We first review the theory of logarithmic Dolbeault complex defined by Burgos. In what follows, let X be a smooth algebraic variety over \mathbb{C} . The C^∞ -de Rham complex gives a resolution

$$(4.1) \quad 0 \rightarrow \mathbb{R} \rightarrow \mathcal{E}_{X,\mathbb{R}} \rightarrow \mathcal{E}_{X,\mathbb{R}}^1 \rightarrow \mathcal{E}_{X,\mathbb{R}}^2 \rightarrow \cdots$$

of the constant sheaf \mathbb{R} on X in terms of the fine sheaves $\mathcal{E}_{X,\mathbb{R}}^\bullet$. Burgos defined a logarithmic version of $\mathcal{E}_{X,\mathbb{R}}^\bullet$ on \overline{X} amenable for defining the weight and Hodge filtrations. We first define the complex $\mathcal{A}_{\overline{X},\mathbb{R}}^\bullet(D)$ to be the $\mathcal{A}_{\overline{X},\mathbb{R}}$ -subalgebra of $j_*\mathcal{A}_{X,\mathbb{R}}^\bullet$ generated locally on coordinate neighborhoods adapted to D by $\mathcal{A}_{\overline{X},\mathbb{R}}^\bullet$ and the sections

$$(4.2) \quad \log |t_\mu|, \quad \operatorname{Re} \frac{dt_\mu}{t_\mu}, \quad \operatorname{Im} \frac{dt_\mu}{t_\mu}$$

for $\mu \in I$ and $\operatorname{Re} dt_\mu, \operatorname{Im} dt_\mu$ for $\mu \notin I$, where the set $I := \{\mu_1, \dots, \mu_h\} \subset [g]$ is such that D is defined locally by $t_{\mu_1} \cdots t_{\mu_h} = 0$. We define the weight filtration W_\bullet on $\mathcal{A}_{\overline{X},\mathbb{R}}^\bullet$ to be the multiplicative ascending filtration obtained by assigning weight 0 to the sections of $\mathcal{A}_{\overline{X},\mathbb{R}}$ and weight 1 to the local sections given in (4.2).

Following Burgos [Bur94, §2], we define the complex $\mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D)$ to be the image of the natural map

$$\mathcal{A}_{\overline{X},\mathbb{R}}^\bullet(D) \otimes_{\mathcal{A}_{\overline{X},\mathbb{R}}} \mathcal{E}_{\overline{X},\mathbb{R}} \rightarrow j_*\mathcal{E}_{X,\mathbb{R}}^\bullet.$$

We let $\mathcal{E}_{\overline{X},\mathbb{R}}^0(D) := \mathcal{E}_{\overline{X},\mathbb{R}}^0(D)$. The complex $\mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D)$ has a weight filtration W_\bullet induced from the weight filtration W_\bullet on $\mathcal{A}_{\overline{X},\mathbb{R}}^\bullet(D)$. The complex $\mathcal{E}_X^\bullet := \mathcal{E}_{X,\mathbb{R}}^\bullet \otimes \mathbb{C}$ has a bigrading

$$\mathcal{E}_X^m = \bigoplus_{p,q \in \mathbb{Z}, p+q=m} \mathcal{E}_X^{p,q}$$

induced from the complex structure of X , and this bigrading induces a bigrading

$$\mathcal{E}_{\overline{X}}^m(D) = \bigoplus_{p,q \in \mathbb{Z}, p+q=m} \mathcal{E}_{\overline{X}}^{p,q}(D), \quad \mathcal{E}_{\overline{X}}^{p,q}(D) := \mathcal{E}_{\overline{X}}^{p+q}(D) \cap j_*\mathcal{E}_X^{p,q}$$

on $\mathcal{E}_{\overline{X}}^\bullet(D) := \mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D) \otimes \mathbb{C}$. We define the Hodge filtration on the complex $\mathcal{E}_{\overline{X}}^\bullet(D)$ by

$$F^p \mathcal{E}_{\overline{X}}^\bullet(D) = \bigoplus_{r \geq p} \mathcal{E}_{\overline{X}}^{r,s}(D).$$

Then we have the following.

Theorem 4.1 ([Bur94, Theorem 1.2, Theorem 2.1]). *There exist filtered quasi-isomorphisms*

$$(Rj_*\mathbb{R}, \tau_{\leq \bullet}) \xrightarrow{\cong} (j_*\mathcal{E}_{X,\mathbb{R}}^\bullet, \tau_{\leq \bullet}) \xleftarrow{\cong} (\mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D), \tau_{\leq \bullet}) \xrightarrow{\cong} (\mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D), W_\bullet)$$

which induces an isomorphism $(Rj_*\mathbb{R}, \tau_{\leq \bullet}) \xrightarrow{\cong} (\mathcal{E}_{\overline{X}, \mathbb{R}}^{\bullet}(D), W_{\bullet})$ in the filtered derived category, and a bifiltered quasi-isomorphism

$$(4.3) \quad (\Omega_{\overline{X}}^{\bullet}(D), W_{\bullet}, F^{\bullet}) \xrightarrow{\cong} (\mathcal{E}_{\overline{X}}^{\bullet}(D), W_{\bullet}, F^{\bullet}).$$

The proof of (4.3) is based on the following result, noting that $\mathcal{E}_{\overline{X}}^{\bullet}(D)$ is the simple complex associated to the double complex $\mathcal{E}_{\overline{X}}^{\bullet, \bullet}(D)$.

Proposition 4.2 ([Bur94] Proposition 2.3). *For any $n, p \in \mathbb{Z}$, we have the exact sequence*

$$0 \longrightarrow \mathrm{Gr}_n^W \Omega_{\overline{X}}^p(D) \longrightarrow \mathrm{Gr}_n^W \mathcal{E}_{\overline{X}}^{p,0}(D) \xrightarrow{\bar{\partial}} \mathrm{Gr}_n^W \mathcal{E}_{\overline{X}}^{p,1}(D) \xrightarrow{\bar{\partial}} \cdots$$

The advantage of using this complex is that the sheaf $\mathcal{E}_{\overline{X}, \mathbb{R}}$ is fine, which gives the following.

Lemma 4.3. *For integers $m, n, p \in \mathbb{Z}$, the sheaves $\mathrm{Gr}_n^W \mathcal{E}_{\overline{X}, \mathbb{R}}^m(D)$ and $\mathrm{Gr}_n^W \mathrm{Gr}_F^p \mathcal{E}_{\overline{X}}^m(D)$ are acyclic with respect to the global section functor $\Gamma(\overline{X}, -)$.*

Proof. Since the sheaf $\mathcal{E}_{\overline{X}, \mathbb{R}}$ has a partition of unity, the $\mathcal{E}_{\overline{X}, \mathbb{R}}$ -modules $\mathrm{Gr}_n^W \mathcal{E}_{\overline{X}, \mathbb{R}}^m(D)$ and $\mathrm{Gr}_n^W \mathrm{Gr}_F^p \mathcal{E}_{\overline{X}}^m(D)$ for $m, n, p \in \mathbb{Z}$ are fine, hence is acyclic with respect to the functor $\Gamma(\overline{X}, -)$. \square

4.2. Cohomology of the Logarithm Sheaf. In this subsection, we first define the $\mathcal{E}_{\overline{X}, \mathbb{R}}$ -module with connection $\mathcal{L}\mathrm{og}^N(D)$. We will apply Theorem 4.1 to this module to calculate the cohomology of U with coefficients in $\mathbb{L}\mathrm{og}^N$.

The module $\mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}}(D)$ has a connection $\nabla : \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}}(D) \rightarrow \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^1(D)$ induced from the connection on $\mathcal{L}\mathrm{og}^N(D)$. If we let

$$(4.4) \quad \underline{e}^k := i^{|\mathbf{k}|} \exp\left(\sum_{\nu=1}^g (-\log |t_{\nu}|) \omega_{\nu}\right) \cdot \underline{\omega}^k$$

for $\mathbf{k} \in \mathbb{N}^g$, where the operator ω_{μ} acts on $\underline{\omega}^k$ by $\omega_{\mu} \cdot \underline{\omega}^k = \underline{\omega}^{k+1_{\mu}}$, then we have

$$\begin{aligned} \nabla(\underline{e}^k) &= i^{|\mathbf{k}|} \exp\left(\sum_{\nu=1}^g (-\log |t_{\nu}|) \omega_{\nu}\right) \cdot \sum_{\mu=1}^g \underline{\omega}^{k+1_{\mu}} \otimes \frac{dt_{\mu}}{t_{\mu}} \\ &\quad - i^{|\mathbf{k}|} \sum_{\mu=1}^g \exp\left(\sum_{\nu=1}^g (-\log |t_{\nu}|) \omega_{\nu}\right) \omega_{\mu} \cdot \underline{\omega}^k \otimes \frac{1}{2} \left(\frac{dt_{\mu}}{t_{\mu}} + \frac{d\bar{t}_{\mu}}{\bar{t}_{\mu}} \right) \\ &= \sum_{\mu=1}^g i^{|\mathbf{k}|} \exp\left(\sum_{\nu=1}^g (-\log |t_{\nu}|) \omega_{\nu}\right) \cdot \underline{\omega}^{k+1_{\mu}} \otimes i \mathrm{Im} \frac{dt_{\mu}}{t_{\mu}} = \sum_{\mu=1}^g \underline{e}^{k+1_{\mu}} \otimes \mathrm{Im} \frac{dt_{\mu}}{t_{\mu}}, \end{aligned}$$

that is

$$(4.5) \quad \nabla(\underline{e}^k) = \sum_{\mu=1}^g \underline{e}^{k+1_{\mu}} \otimes \mathrm{Im} \frac{dt_{\mu}}{t_{\mu}}.$$

This calculation shows that we may define an \mathbb{R} -structure $\mathcal{L}\mathrm{og}^N(D)$ on $\mathcal{L}\mathrm{og}^N(D)$ as follows.

Definition 4.4. We let $\mathcal{L}\mathrm{og}^N(D)$ be the free $\mathcal{E}_{\overline{X}, \mathbb{R}}$ -module generated by \underline{e}^k of (4.4) for $\mathbf{k} \in \mathbb{N}^g$ such that $|\mathbf{k}| \leq N$, with connection

$$\nabla : \mathcal{L}\mathrm{og}^N(D) \rightarrow \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^1(D)$$

given by (4.5). By definition, $\mathcal{L}og^N(D)$ is an $\mathcal{E}_{\overline{X},\mathbb{R}}$ -submodule of $\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}}(D)$.

We define the filtration W_\bullet on $\mathcal{L}og^N(D)$ such that \underline{e}^k is of weight $-2|k|$. The basis $\underline{\omega}^k$ of $\mathcal{L}og^N(D)$ in terms of \underline{e}^k is given by

$$(4.6) \quad \underline{\omega}^k = (-i)^{|k|} \exp\left(\sum_{\nu=1}^g \log |t_\nu| \omega_\nu\right) \cdot \underline{e}^k,$$

where ω_μ acts on \underline{e}^k as

$$\begin{aligned} \omega_\mu \cdot \underline{e}^k &= i^{|k|} \exp\left(\sum_{\nu=1}^g (-\log |t_\nu|) \omega_\nu\right) \omega_\mu \cdot \underline{e}^k = i^{|k|} \exp\left(\sum_{\nu=1}^g (-\log |t_\nu|) \omega_\nu\right) \cdot \underline{\omega}^{k+1_\mu} \\ &= (-i) i^{|k|+1} \exp\left(\sum_{\nu=1}^g (-\log |t_\nu|) \omega_\nu\right) \cdot \underline{\omega}^{k+1_\mu} = (-i) \underline{e}^{k+1_\mu}. \end{aligned}$$

Note that we have an exact sequence

$$(4.7) \quad 0 \rightarrow \text{Sym}^N \mathbb{R}(1) \otimes \mathcal{E}_{\overline{X},\mathbb{R}} \rightarrow \mathcal{L}og^N(D) \rightarrow \mathcal{L}og^{N-1}(D) \rightarrow 0.$$

compatible with the connection and W_\bullet . The natural inclusion induces an isomorphism of projective systems

$$(4.8) \quad \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}}(D) = \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}}(D),$$

which is compatible with the connection and the filtration W_\bullet . The relation between the basis $(\underline{u}^k)_{k \in \mathbb{N}^g}$ of $\mathbb{L}og^N$ and $(\underline{e}^k)_{k \in \mathbb{N}^g}$ of $\mathcal{L}og^N(D)$ is given by

$$\begin{aligned} \underline{u}^k &= (2\pi i)^{|k|} \exp\left(\sum_{\nu=1}^g (-\log t_\nu) \omega_\nu\right) \cdot \underline{\omega}^k = (2\pi)^{|k|} \exp\left(\sum_{\nu=1}^g (-i \text{Im}(\log t_\nu)) \omega_\nu\right) \cdot \underline{e}^k \\ &= (2\pi)^{|k|} \exp\left(\sum_{\nu=1}^g (-\text{Im}(\log t_\nu)) e_\nu\right) \cdot \underline{e}^k, \end{aligned}$$

where e_ν acts on \underline{e}^k by $e_\nu \cdot \underline{e}^k = \underline{e}^{k+1_\nu}$, the equality given locally for each choice of a branch of $\log t_\nu$. Since both $(2\pi)^{|k|}$ and $\text{Im}(\log t_\nu)$ are real-valued functions on X , this gives an isomorphism

$$(4.9) \quad \mathbb{L}og^N \otimes \mathcal{E}_{X,\mathbb{R}} \cong \mathcal{L}og^N := \mathcal{L}og^N(D)|_X$$

which is compatible with the connection and the filtration W_\bullet . Moreover, since $\text{Gr}_n^W \mathbb{L}og^N$ is a constant variation of pure Hodge structure on X , this naturally extends to a constant variation of pure Hodge structures on \overline{X} , and the isomorphism induced on Gr_n^W by (4.9) extends to an isomorphism

$$(4.10) \quad \text{Gr}_n^W \mathcal{L}og^N(D) \cong \text{Gr}_n^W \mathbb{L}og^N \otimes \mathcal{E}_{\overline{X},\mathbb{R}}$$

on \overline{X} .

Definition 4.5. We define the logarithmic Čech-Dolbeault complex $\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D^\bullet)$ on \overline{X} to be the simple complex associated to the double complex

$$\bigoplus_{|I|=1} \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D^I) \xrightarrow{\partial} \dots \xrightarrow{\partial} \bigoplus_{|I|=g} \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D^I)$$

with $\bigoplus_{|I|=1} \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X},\mathbb{R}}^\bullet(D^I)$ in degree 0, where the horizontal differentials are the standard alternating sums.

Proposition 4.6. *Let $j : U \hookrightarrow \overline{X}$ be the natural inclusion. Then we have canonical isomorphisms*

$$(4.11) \quad Rj_* \mathbb{L}\mathrm{og}^N \cong Rj_*(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet) \cong \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)$$

in the derived category of abelian sheaves on \overline{X} . In particular, by taking the functor $R\Gamma(\overline{X}, -)$, we obtain a canonical isomorphism

$$H^m(U, \mathbb{L}\mathrm{og}^N) \cong H^m(\Gamma(\overline{X}, \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet))).$$

Proof. Since $\mathbb{L}\mathrm{og}^N$ is an \mathbb{R} -local system on U , the resolution (4.1) gives a resolution

$$0 \rightarrow \mathbb{L}\mathrm{og}^N \rightarrow \mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}} \rightarrow \mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^1 \rightarrow \cdots$$

of $\mathbb{L}\mathrm{og}^N$ on U . By taking Rj_* , we obtain the first isomorphism of (4.11). As in the proof of Proposition 3.9, for $\mu \in [g]$, we let $U_\mu = X \setminus Z_\mu = \overline{X} \setminus (D \cup Z_\mu)$, which gives an open covering $\{U_\mu\}$ of U . For $I \subset [g]$, we let $U_I = \overline{X} \setminus D^I = \bigcap_{\mu \in I} U_\mu$, and we let $j_I : U_I \hookrightarrow \overline{X}$ be the natural inclusion. If we let $j_{\bullet*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet)$ be the Čech complex defined as the simple complex associated to the double complex

$$\prod_{|I|=1} j_{I*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet) \rightarrow \cdots \rightarrow \prod_{|I|=g} j_{I*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet).$$

Following a similar argument to that in (3.16), noting that $\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet$ are complexes of fine sheaves, we have a quasi-isomorphism

$$(4.12) \quad Rj_*(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet) \xrightarrow{\cong} j_{\bullet*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet).$$

In order to complete our proof, we prove by induction on $N \geq 0$ that for any $I \subset [g]$, the natural morphism

$$(4.13) \quad \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I) \rightarrow j_{I*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet)$$

is a quasi-isomorphism. If $N = 0$, then $\mathbb{L}\mathrm{og}^0 \cong \mathbb{R}$ and $\mathcal{L}\mathrm{og}^0(D) \cong \mathcal{E}_{\overline{X}, \mathbb{R}}$, hence (4.13) is a quasi-isomorphism by Theorem 4.1. Suppose (4.13) is a quasi-isomorphism for $N - 1 \geq 0$. The short exact sequence of (4.7) gives the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Sym}^N \mathbb{R}(\mathbb{1}) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I) & \longrightarrow & \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I) & \longrightarrow & \mathcal{L}\mathrm{og}^{N-1}(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & j_{I*}(\mathrm{Sym}^N \mathbb{R}(\mathbb{1}) \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet) & \longrightarrow & j_{I*}(\mathcal{L}\mathrm{og}^N \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet) & \longrightarrow & j_{I*}(\mathcal{L}\mathrm{og}^{N-1} \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet) \longrightarrow 0. \end{array}$$

The first vertical morphism is a quasi-isomorphism by Theorem 4.1, noting that

$$j_{I*}(\mathrm{Sym}^N \mathbb{R}(\mathbb{1}) \otimes \mathcal{E}_{U_I, \mathbb{R}}^\bullet) = \mathrm{Sym}^N \mathbb{R}(\mathbb{1}) \otimes j_{I*} \mathcal{E}_{U_I, \mathbb{R}}^\bullet,$$

and the third vertical morphism is a quasi-isomorphism by the induction hypothesis. Hence the middle vertical morphism is a quasi-isomorphism, which by induction proves that (4.13) is a quasi-isomorphism for any $N \geq 0$. Combining this result with (4.12), we obtain isomorphisms

$$Rj_*(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet) \cong j_{\bullet*}(\mathbb{L}\mathrm{og}^N \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet) \cong \mathcal{L}\mathrm{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)$$

in the derived category, which proves the second isomorphism of (4.11). The statement for cohomology follows from (4.11) and the fact that $\mathcal{L}\mathrm{og}^N \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D)$ is a complex of fine sheaves. \square

4.3. Mixed Hodge Structure of the Logarithm Sheaf. In this subsection, we show that the weight and Hodge filtrations on $\mathcal{L}\text{og}^N(D)$ induce the weight and Hodge filtrations of the mixed \mathbb{R} -Hodge structure on the cohomology of U with coefficients in $\mathbb{L}\text{og}^N$.

We define the filtration W_\bullet on $\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I)$ and F^\bullet on $\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)$ by

$$\begin{aligned} W_n(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I)) &:= \sum_{k \in \mathbb{Z}} W_{n-k} \mathcal{L}\text{og}^N(D) \otimes W_k \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I), \\ F^p(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)) &:= \sum_{q \in \mathbb{Z}} F^{p-q} \mathcal{L}\text{og}^N(D) \otimes F^q \mathcal{E}_{\overline{X}}^\bullet(D^I), \end{aligned}$$

where we use the equality $\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I) = \mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)$ induced from (4.8) for the definition of F^\bullet . For any integer $n \in \mathbb{Z}$, we let $W_n(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet))$ be the subcomplex of $\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)$ defined as the simple complex associated to the double complex

$$\bigoplus_{|I|=1} W_n(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I)) \rightarrow \cdots \rightarrow \bigoplus_{|I|=g} W_{n+g-1}(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^I)),$$

and for any integer $p \in \mathbb{Z}$, we let $F^p(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet))$ be the subcomplex of $\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet)$ defined as the simple complex associated to the double complex

$$\bigoplus_{|I|=1} F^p(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)) \rightarrow \cdots \rightarrow \bigoplus_{|I|=g} F^p(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)).$$

Theorem 4.7. *The natural inclusion*

$$\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet) \rightarrow \mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet) = \mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet)$$

induces a bifiltered quasi-isomorphism

$$(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^\bullet), W_\bullet, F^\bullet) \xrightarrow{\cong} (\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet), W_\bullet, F^\bullet).$$

Proof. It is sufficient to prove that we have a bifiltered quasi-isomorphism

$$(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I), W_\bullet, F^\bullet) \xrightarrow{\cong} (\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I), W_\bullet, F^\bullet)$$

for any $I \subset [g]$. Since the filtration W_\bullet on $\mathcal{L}\text{og}^N(D)$ and $\mathcal{L}\text{og}^N(D)$ are filtrations by free submodules, we have

$$\begin{aligned} \text{Gr}_n^W(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)) &= \bigoplus_{k \in \mathbb{Z}} \text{Gr}_{n-k}^W \mathcal{L}\text{og}^N(D) \otimes \text{Gr}_k^W \Omega_{\overline{X}}^\bullet(D^I) = \bigoplus_{k \in \mathbb{Z}} \text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N \otimes \text{Gr}_k^W \Omega_{\overline{X}}^\bullet(D^I) \\ \text{Gr}_n^W(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)) &= \bigoplus_{k \in \mathbb{Z}} \text{Gr}_{n-k}^W \mathcal{L}\text{og}^N(D) \otimes \text{Gr}_k^W \mathcal{E}_{\overline{X}}^\bullet(D^I) = \bigoplus_{k \in \mathbb{Z}} \text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N \otimes \text{Gr}_k^W \mathcal{E}_{\overline{X}}^\bullet(D^I). \end{aligned}$$

Moreover, since the filtration F^\bullet on $\text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N$ is a filtration by \mathbb{C} -local systems, we see that

$$\begin{aligned} \text{Gr}_F^p \text{Gr}_n^W(\mathcal{L}\text{og}^N(D) \otimes \Omega_{\overline{X}}^\bullet(D^I)) &= \bigoplus_{k, q \in \mathbb{Z}} \text{Gr}_F^{p-q} \text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N \otimes \text{Gr}_k^W \Omega_{\overline{X}}^{q, \bullet}(D^I) \\ \text{Gr}_F^p \text{Gr}_n^W(\mathcal{L}\text{og}^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^I)) &= \bigoplus_{k, q \in \mathbb{Z}} \text{Gr}_F^{p-q} \text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N \otimes \text{Gr}_k^W \mathcal{E}_{\overline{X}}^{q, \bullet}(D^I). \end{aligned}$$

Since $\text{Gr}_F^{p-q} \text{Gr}_{n-k}^W \mathbb{L}\text{og}_{\mathbb{C}}^N$ are constant \mathbb{C} -local systems on \overline{X} , our assertion follows from Proposition 4.2. \square

By Theorem 4.7, we see that the complex $\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)$ may be used to calculate the mixed \mathbb{R} -Hodge structure $H^m(U, \mathbb{L}og^N)$. This implies in particular that

$$((\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet), W_\bullet), (\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet), W_\bullet, F^\bullet), \text{id})$$

is a cohomological mixed \mathbb{R} -Hodge complex on \overline{X} in the sense of [Del74, (8.1.6)]. Furthermore, by Lemma 4.3, $\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)$ as well as $\text{Gr}_F^p \text{Gr}_n^W$ are complexes of sheaves acyclic with respect to the global section functor, we may simply take the global section to calculate the corresponding cohomology groups and their filtrations.

Definition 4.8. We let

$$R^\bullet(U, \mathbb{L}og^N) := \Gamma(\overline{X}, \mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)), \quad R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}}) := R^\bullet(U, \mathbb{L}og^N) \otimes \mathbb{C}$$

with filtrations \widetilde{W}_\bullet and F^\bullet defined by

$$\begin{aligned} \widetilde{W}_n R^\bullet(U, \mathbb{L}og^N) &:= \Gamma(\overline{X}, W_n(\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet))), \\ F^p R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}}) &:= \Gamma(\overline{X}, F^p(\mathcal{L}og^N(D) \otimes \mathcal{E}_{\overline{X}}^\bullet(D^\bullet))). \end{aligned}$$

Furthermore, we let $W_\bullet := \text{Dec}(\widetilde{W})_\bullet$ be the décalage of \widetilde{W}_\bullet in the sense of [Del71, (1.3.3)], given by

$$W_n R^m(U, \mathbb{L}og^N) := \text{Ker}(d^m : \widetilde{W}_{n-m} R^m(U, \mathbb{L}og^N) \rightarrow R^{m+1}(U, \mathbb{L}og^N) / \widetilde{W}_{n-m-1} R^{m+1}(U, \mathbb{L}og^N)).$$

By [Del74, (8.1.9)], the spectral sequences for the filtrations W_\bullet and F^\bullet degenerates at E_1 . This implies in particular that the filtrations W_\bullet and F^\bullet are strictly compatible with the differential maps, hence

$$(4.14) \quad \begin{aligned} W_n H^m(R^\bullet(U, \mathbb{L}og^N)) &= H^m(W_n R^\bullet(U, \mathbb{L}og^N)), \\ F^q H^m(R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}})) &= H^m(F^q R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}})). \end{aligned}$$

By the property of the décalage, we have

$$W_n H^m(R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}})) = \widetilde{W}_{n-m} H^m(R^\bullet(U, \mathbb{L}og^N)).$$

Corollary 4.9. *The complex $(R^\bullet(U, \mathbb{L}og^N), W_\bullet, F^\bullet)$ given above calculates the mixed \mathbb{R} -Hodge structure $H^m(U, \mathbb{L}og^N)$ given in Proposition 2.11.*

Proof. By Proposition 4.6, we have an isomorphism

$$H^m(R^\bullet(U, \mathbb{L}og^N)) \cong H^m(U, \mathbb{L}og^N),$$

and by Theorem 4.7, the filtrations induced from \widetilde{W}_\bullet and F^\bullet on $R^\bullet(U, \mathbb{L}og^N)$ corresponds through this isomorphism to the filtrations \widetilde{W}_\bullet and F^\bullet on $H^m(U, \mathbb{L}og^N)$. Our assertion follows from $W_\bullet = \widetilde{W}_\bullet[m]$. \square

As a variant of $R^\bullet(U, \mathbb{L}og^N)$, we let

$$R^\bullet(U, \mathbb{L}og^N(n)) := \Gamma(\overline{X}, \mathcal{L}og^N(D)(n) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^\bullet(D^\bullet)), \quad R^\bullet(U, \mathbb{L}og^N_{\mathbb{C}}(n)) := R^\bullet(U, \mathbb{L}og^N(n)) \otimes \mathbb{C}$$

for any integer $n \in \mathbb{Z}$, where $\mathcal{L}og^N(D)(n) := \mathcal{L}og^N(D) \otimes \mathbb{R}(n)$ with the corresponding weight and Hodge filtrations. This filtered complex calculates the mixed \mathbb{R} -Hodge structure $H^m(U, \mathbb{L}og^N(n))$.

The geometric polylogarithm class is given explicitly as follows. Through the map

$$\mathcal{L}og^N(D)(g) \otimes \Omega_X^g(D^{[g]}) \rightarrow \mathcal{L}og^N(D)(g) \otimes \mathcal{E}_X^g(D^{[g]}),$$

the differential form

$$\xi_N = \underline{\omega}^0(g) \otimes \frac{dt_1}{t_1 - 1} \wedge \cdots \wedge \frac{dt_g}{t_g - 1}$$

of (3.18) giving the geometric polylogarithm class $[\xi_N]$ in $H^1(U, \mathbb{L}\text{og}^N(g))$ defines an element

$$\xi_N \in R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) = \Gamma(\overline{X}, \mathcal{L}\text{og}^N(D)(g) \otimes \mathcal{E}_{\overline{X}}^g(D^{[g]})).$$

Note that the weight and Hodge filtrations on $R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g))$ are given by

$$W_n R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) = \widetilde{W}_{n-2g+1} R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) = \Gamma(\overline{X}, W_{n-g}(\mathcal{L}\text{og}^N(D)(g) \otimes \mathcal{E}_{\overline{X}}^g(D^{[g]}))),$$

$$F^p R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) = \Gamma(\overline{X}, F^p(\mathcal{L}\text{og}^N(D)(g) \otimes \mathcal{E}_{\overline{X}}^g(D^{[g]}))).$$

Since ξ_N is an element in both

$$\Gamma(\overline{X}, W_{-2g} \mathcal{L}\text{og}^N(D)(g) \otimes W_g \mathcal{E}_{\overline{X}}^g(D^{[g]})) \subset W_0 R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)),$$

$$\Gamma(\overline{X}, F^{-g} \mathcal{L}\text{og}^N(D)(g) \otimes F^g \mathcal{E}_{\overline{X}}^g(D^{[g]})) \subset F^0 R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)),$$

we see that $\xi_N \in (F^0 \cap W_0) R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g))$, hence the class $[\xi_N]$ satisfies

$$[\xi_N] \in (F^0 \cap W_0) H^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)).$$

5. ABSOLUTE POLYLOGARITHM

The purpose of this section is to define and give an explicit construction of the absolute polylogarithm class

$$\mathbf{pol} \in H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g)).$$

We start with the definition of the absolute Hodge cohomology $H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g))$.

5.1. The Absolute Polylogarithm Class. We let

$$R^{\bullet}(U, \mathbb{L}\text{og}^N(g)) = \Gamma(\overline{X}, \mathcal{L}\text{og}^N(D)(g) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^{\bullet}(D^{\bullet}))$$

be the logarithmic Čech-Dolbeault complex of §4.3.

Definition 5.1. We define $R_{\mathcal{A}}^{\bullet}(U, \mathbb{L}\text{og}^N(g))$ to be the complex

$$\text{Cone}\left(W_0 R^{\bullet}(U, \mathbb{L}\text{og}^N(g)) \oplus (F^0 \cap W_0) R^{\bullet}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) \rightarrow W_0 R^{\bullet}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g))\right)[-1],$$

where the map is defined by $(x, y) \mapsto x - y$. Then we define the absolute Hodge cohomology $H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g))$ of U with coefficients in $\mathbb{L}\text{og}^N(g)$ by

$$(5.1) \quad H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g)) := H^m(R_{\mathcal{A}}^{\bullet}(U, \mathbb{L}\text{og}^N(g))).$$

In addition, we let

$$H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}(g)) := \varprojlim_N H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g)).$$

Remark 5.2. By Corollary 4.9, the complex $R^{\bullet}(U, \mathbb{L}\text{og}^N(g))$ is a complex of filtered modules which gives the mixed \mathbb{R} -Hodge structure on the cohomology of U with coefficients in $\mathbb{L}\text{og}^N(g)$. In fact, the complex $R^{\bullet}(U, \mathbb{L}\text{og}^N(g))$ gives a Hodge complex, and hence an object in the derived category $D^b(\text{MHS}_{\mathbb{R}})$ via Beilinson's equivalence of categories [Beĭ86, Theorem 3.4]. Then the absolute Hodge cohomology (5.1) may be interpreted as

$$H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g)) = \text{Hom}_{D^b(\text{MHS}_{\mathbb{R}})}(\mathbb{R}(0), R^{\bullet}(U, \mathbb{L}\text{og}^N(g))[m]).$$

By definition of the cone, we have a long exact sequence

$$\begin{aligned} \cdots &\rightarrow W_0 H^{m-1}(U, \mathbb{L}\text{og}^N(g)) \oplus (F^0 \cap W_0) H^{m-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) \rightarrow W_0 H^{m-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) \\ &\rightarrow H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g)) \\ &\rightarrow W_0 H^m(U, \mathbb{L}\text{og}^N(g)) \oplus (F^0 \cap W_0) H^m(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) \rightarrow W_0 H^m(U, \mathbb{L}\text{og}_{\mathbb{C}}^N(g)) \rightarrow \cdots \end{aligned}$$

Note that for any mixed \mathbb{R} -Hodge structure $V := (V, W_{\bullet}, F^{\bullet})$, the extension groups in $\text{MHS}_{\mathbb{R}}$ are calculated as

$$(5.2) \quad \text{Ext}_{\text{MHS}_{\mathbb{R}}}^m(\mathbb{R}(0), V) = \begin{cases} \text{Ker}(W_0 V \oplus (F^0 \cap W_0) V_{\mathbb{C}} \rightarrow W_0 V_{\mathbb{C}}) & m = 0, \\ \text{Coker}(W_0 V \oplus (F^0 \cap W_0) V_{\mathbb{C}} \rightarrow W_0 V_{\mathbb{C}}) & m = 1, \\ 0 & \text{otherwise,} \end{cases}$$

where the map is defined by $(x, y) \mapsto x - y$. Thus we obtain the short exact sequence

$$(5.3) \quad 0 \rightarrow \text{Ext}_{\text{MHS}_{\mathbb{R}}}^1(\mathbb{R}(0), H^{m-1}(U, \mathbb{L}\text{og}^N(g))) \rightarrow H_{\mathcal{A}}^m(U, \mathbb{L}\text{og}^N(g)) \rightarrow \text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^m(U, \mathbb{L}\text{og}^N(g))) \rightarrow 0$$

for any $m \in \mathbb{Z}$.

The following proposition is crucial in defining the polylogarithm class.

Proposition 5.3. *We have a canonical isomorphism*

$$H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g)) \cong \text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^{2g-1}(U, \mathbb{L}\text{og}(g))).$$

Moreover, we have

$$H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g)) \cong \begin{cases} \mathbb{R} & g > 1 \\ \mathbb{R} \oplus \mathbb{R} & g = 1. \end{cases}$$

Proof. By Lemma 3.5, we have

$$H^{2g-2}(U, \mathbb{L}\text{og}^N(g)) = \begin{cases} 0 & g \neq 2 \\ \mathbb{R}(0) & g = 2. \end{cases}, \quad H^{2g-1}(U, \mathbb{L}\text{og}^N(g)) = \begin{cases} \prod_{k=0}^N \text{Sym}^k \mathbb{R}(1) & g \neq 1 \\ \mathbb{R}(0) \oplus \prod_{k=0}^N \text{Sym}^k \mathbb{R}(1) & g = 1. \end{cases}$$

On the other hand, by (5.2), we have

$$\text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), \mathbb{R}(n)) \cong \begin{cases} \mathbb{R} & n = 0, \\ 0 & n \neq 0, \end{cases} \quad \text{Ext}_{\text{MHS}_{\mathbb{R}}}^1(\mathbb{R}(0), \mathbb{R}(n)) \cong \begin{cases} \mathbb{C}/(2\pi i)^n \mathbb{R} & n > 0, \\ 0 & n \leq 0, \end{cases}$$

and $\text{Ext}_{\text{MHS}_{\mathbb{R}}}^m(\mathbb{R}(0), \mathbb{R}(n)) = 0$ if $m \neq 0, 1$. Our assertion follows from (5.3) and by passing to the limit. \square

The absolute polylogarithm class for X is defined as follows.

Definition 5.4. We define the *absolute polylogarithm class* \mathbf{pol} to be the class

$$\mathbf{pol} = (\mathbf{pol}_N) \in H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g)) = \varprojlim_N H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}^N(g))$$

which maps to the geometric polylogarithm class in $\text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^{2g-1}(U, \mathbb{L}\text{og}(g)))$ of Remark 3.7 through the isomorphism of Proposition 5.3.

Remark 5.5. When $g > 1$, then $\mathbf{pol}_N \in H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}^N(g))$ is characterized as the class which maps to 1 through the isomorphism

$$H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}^N(g)) \cong \text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^{2g-1}(U, \mathbb{L}\text{og}^N(g))) \stackrel{\text{res}_1}{\cong} \text{Hom}_{\text{MHS}_{\mathbb{R}}}(\mathbb{R}(0), H^0(Z, \mathbb{L}\text{og}^N)) \cong \mathbb{R}.$$

In what follows, let

$$R^\bullet(U, \mathbb{L}\text{og}(g)) := \varprojlim_N R^\bullet(U, \mathbb{L}\text{og}^N(g)), \quad R_{\mathcal{A}}^\bullet(U, \mathbb{L}\text{og}(g)) := \varprojlim_N R_{\mathcal{A}}^\bullet(U, \mathbb{L}\text{og}^N(g)).$$

The filtrations W_\bullet and F^\bullet on $R^\bullet(U, \mathbb{L}\text{og}^N(g))$ induce filtrations W_\bullet and F^\bullet on $R^\bullet(U, \mathbb{L}\text{og}(g))$.

Proposition 5.6. *For any integer $m \geq 0$, we have*

$$H^m(R^\bullet(U, \mathbb{L}\text{og}(g))) \cong H^m(U, \mathbb{L}\text{og}(g)).$$

Moreover, for any integer $n \in \mathbb{Z}$, we have

$$H^m\left(\varprojlim_N W_n R^\bullet(U, \mathbb{L}\text{og}^N(g))\right) \cong \varprojlim_N H^m(W_n R^\bullet(U, \mathbb{L}\text{og}^N(g))).$$

Proof. We start by proving the first isomorphism, that is

$$H^m\left(\varprojlim_N R^\bullet(U, \mathbb{L}\text{og}^N(g))\right) \cong \varprojlim_N H^m(R^\bullet(U, \mathbb{L}\text{og}^N(g))).$$

By [KS94, Proposition 1.12.4], it is sufficient to check that the systems $(R^m(U, \mathbb{L}\text{og}^N(g)))_N$ and $(H^m(R^\bullet(U, \mathbb{L}\text{og}^N(g))))_N$ satisfy the Mittag-Leffler condition for any $m \in \mathbb{Z}$. For $(R^m(U, \mathbb{L}\text{og}^N(g)))_N$, recall the definition of $\mathcal{L}\text{og}^N(D)$ given in Definition 4.4. As an $\mathcal{E}_{\overline{X}, \mathbb{R}}^P$ -module, we have a splitting

$$(5.4) \quad \mathcal{L}\text{og}^{N+1}(D) = \mathcal{L}\text{og}^N(D) \oplus W_{-2N-2} \mathcal{L}\text{og}^{N+1}(D),$$

and the projection $\mathcal{L}\text{og}^{N+1}(D) \rightarrow \mathcal{L}\text{og}^N(D)$ is simply the projection to the first component. This implies that

$$\mathcal{L}\text{og}^{N+1}(D)(g) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^P(D^q) \rightarrow \mathcal{L}\text{og}^N(D)(g) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}^P(D^q)$$

is split surjective, hence the induced morphism $R^m(U, \mathbb{L}\text{og}^{N+1}(g)) \rightarrow R^m(U, \mathbb{L}\text{og}^N(g))$ is also split surjective. The Mittag-Leffler condition for $(H^m(R^\bullet(U, \mathbb{L}\text{og}^N(g))))_N$ follows from Lemmas 3.3 and 3.5.

We prove the second isomorphism in a similar way. Since the splitting (5.4) is compatible with the filtration W_\bullet , the system $(W_n R^m(U, \mathbb{L}\text{og}^N(g)))_N$ satisfies the Mittag-Leffler condition. On the other hand, the Mittag-Leffler condition for $(H^m(R^\bullet(U, \mathbb{L}\text{og}^N(g))))_N$ shown above implies that for $(W_n H^m(R^\bullet(U, \mathbb{L}\text{og}^N(g))))_N$, since the morphisms of mixed \mathbb{R} -Hodge structures are strictly compatible with the filtrations. By (4.14), the latter system is equal to $(H^m(W_n R^\bullet(U, \mathbb{L}\text{og}^N(g))))_N$, hence we obtain our assertion. \square

In the following subsections, we will construct an explicit cocycle giving the polylogarithm class. More precisely, we will define a triple

$$(\alpha, \eta, \xi) \in W_0 R^{2g-2}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g)) \oplus W_0 R^{2g-1}(U, \mathbb{L}\text{og}(g)) \oplus (F^0 \cap W_0) R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g)),$$

which satisfies:

- (1) The cocycle conditions $d\alpha = \eta - \xi$, $d\eta = 0$ and $d\xi = 0$.
- (2) $\xi = (\xi_N)$, where ξ_N are the differential forms of (3.18) giving the geometric polylogarithm class.

The first condition insures that (α, η, ξ) defines a class in $H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g))$, and the second implies that this class gives the absolute polylogarithm class **pol**.

5.2. Polylogarithm Function and the Case $g = 1$. We first review the definition of the classical polylogarithm function. The polylogarithm function, defined by the power series

$$\mathrm{Li}_k(t) := \sum_{n=1}^{\infty} \frac{t^n}{n^k}, \quad (t \in \mathbb{C}, |t| < 1, k \in \mathbb{N}),$$

may be extended analytically to $\mathbb{C} \setminus [1, \infty)$ using the integration

$$\mathrm{Li}_{k+1}(t) = \int_0^t \mathrm{Li}_k(u) \frac{du}{u}$$

for any integer $k \geq 0$.

For $m \geq 0$ and $|t| < 1$, let

$$(5.5) \quad L_{m+1}(t) := \sum_{k=0}^m \frac{(-\log |t|)^{m-k}}{(m-k)!} \mathrm{Li}_{k+1}(t).$$

In [Zag90, §1], Zagier defined the Bloch-Wigner-Ramakrishnan polylogarithm functions $D_m^\circ(t)$ by

$$D_m^\circ(t) := \begin{cases} \mathrm{Im}(L_m(t)) & (m: \text{even}), \\ \mathrm{Re}(L_m(t)) + \frac{(\log |t|)^m}{(2m)!} & (m: \text{odd}). \end{cases}$$

In this article, we take a slightly different normalization as follows:

Definition 5.7. For any $m \in \mathbb{N}$, we define the *Bloch-Wigner-Ramakrishnan polylogarithm function* $D_m(t)$ by

$$(5.6) \quad D_m(t) := \mathrm{Im}(i^m L_m(t)).$$

The relations between the normalizations are

$$D_m(t) = \mathrm{Im}(i^m L_m(t)) = (-1)^n D_m^\circ(t)$$

if $m = 2n$ is even, and

$$D_m(t) = \mathrm{Im}(i^m L_m(t)) = (-1)^n \mathrm{Re}(L_m(t)) = (-1)^n D_m^\circ(t) - (-1)^n \frac{(\log |t|)^m}{(2m)!}$$

if $m = 2n + 1$ is odd. Then the functions $D_m(t)$ satisfy the following functional equations.

Proposition 5.8 ([Zag90, Proposition 1]). *The function $D_m(t)$ of (5.6) can be continued to a single-valued real analytic function on $U := \mathbb{P}^1 \setminus \{0, 1, \infty\}$. In particular, it is a function in $\mathcal{E}_{U, \mathbb{R}}$. The function $D_m(t)$ satisfies the functional equation*

$$(5.7) \quad D_m(1/t) = \begin{cases} D_m(t) & (m: \text{even}), \\ D_m(t) + (\log |t|)^m / m! & (m: \text{odd}). \end{cases}$$

We will use the functions $D_m(t)$ given above to explicitly describe the polylogarithm class. In this subsection, we consider the case $g = 1$. Let $\bar{X} = \mathbb{P}^1$, $D = \{0, \infty\}$ and $Z = \{1\}$.

Theorem 5.9 (Case $g = 1$). *The polylogarithm class \mathbf{pol} is given by the triple (α, η, ξ) , where*

$$\begin{aligned} \xi &= \underline{\omega}^0(1) \otimes \frac{dt}{t-1} \in (F^0 \cap W_0)R^1(U, \mathbb{L}\mathrm{og}_{\mathbb{C}}(1)), \\ \eta &:= \mathrm{Re}(\xi) = (\xi + \bar{\xi})/2 \in W_0R^1(U, \mathbb{L}\mathrm{og}(1)), \end{aligned}$$

and

$$\alpha := \sum_{m=0}^{\infty} D_{m+1}(t) \underline{e}^m(1) \in W_0 R^0(U, \mathbb{L}\text{og}(1)).$$

Proof. Let us consider

$$\tilde{\alpha} := \sum_{k=0}^{\infty} (-1)^{k+1} \text{Li}_{k+1}(t) \underline{\omega}^k(1)$$

which is a multi-valued section of $\mathcal{L}\text{og}(1)$ on U . Then we have

$$\begin{aligned} \nabla(\tilde{\alpha}) &= \sum_{k=0}^{\infty} (-1)^{k+1} \underline{\omega}^k(1) \otimes d\text{Li}_{k+1}(t) + \sum_{k=0}^{\infty} (-1)^{k+1} \text{Li}_{k+1}(t) \underline{\omega}^{k+1}(1) \otimes \frac{dt}{t} \\ &= \sum_{k=0}^{\infty} (-1)^{k+1} \text{Li}_k(t) \underline{\omega}^k(1) \otimes \frac{dt}{t} + \sum_{k=1}^{\infty} (-1)^k \text{Li}_k(t) \underline{\omega}^k(1) \otimes \frac{dt}{t} = -\text{Li}_0(t) \underline{\omega}^0(1) \otimes \frac{dt}{t} = \xi. \end{aligned}$$

The section $\tilde{\alpha}$ only gives a section of

$$\mathcal{L}\text{og}(1) \otimes \mathcal{E}_{\overline{X}}(D) := \lim_{\leftarrow N} \left(\mathcal{L}\text{og}^N(1) \otimes \mathcal{E}_{\overline{X}}(D) \right)$$

on $\mathbb{C} \setminus [1, \infty)$ and does not extend to a global section of \overline{X} , since the functions $\text{Li}_k(t)$ extend to multi-valued functions and do not extend to single-valued functions on $\mathbb{C} \setminus \{1\}$. Note that we have $\nabla(\text{Re}(\tilde{\alpha})) = \text{Re}(\xi) = \eta$, hence if we let

$$\alpha := \text{Re}(\tilde{\alpha}) - \tilde{\alpha} = -i \text{Im}(\tilde{\alpha}),$$

then we have

$$(5.8) \quad \nabla(\alpha) = \eta - \xi = -i \text{Im}(\xi).$$

By (4.6), we have

$$\begin{aligned} \tilde{\alpha} &= \sum_{k=0}^{\infty} (-1)^{k+1} \text{Li}_{k+1}(t) (-i)^k \exp(\log |t| \omega) \cdot \underline{e}^k(1) = \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} (-1)^{k+1} (-i)^{n+k} \frac{(\log |t|)^n}{n!} \text{Li}_{k+1}(t) \underline{e}^{n+k}(1) \\ &= - \sum_{m=0}^{\infty} i^m \left(\sum_{k=0}^m \frac{(-\log |t|)^{m-k}}{(m-k)!} \text{Li}_{k+1}(t) \right) \underline{e}^m(1) = \sum_{m=0}^{\infty} i^{m+1} L_{m+1}(t) \cdot i \underline{e}^m(1). \end{aligned}$$

Since the basis $\underline{e}^m(1)$ is purely imaginary, we see that α is given by

$$\alpha = -i \text{Im}(\tilde{\alpha}) = -i \sum_{m=0}^{\infty} \text{Im}(i^{m+1} L_{m+1}(t)) \cdot i \underline{e}^m(1) = \sum_{m=0}^{\infty} D_{m+1}(t) \underline{e}^m(1),$$

which by Proposition 5.8 is an element in $W_0 \Gamma(U, \mathcal{L}\text{og}(1) \otimes \mathcal{E}_{U, \mathbb{R}})$.

It remains to show that α defines an element of $W_0 R^0(U, \mathbb{L}\text{og}(1)) = W_0 \Gamma(\overline{X}, \mathcal{L}\text{og}(D)(1) \otimes \mathcal{E}_{\overline{X}, \mathbb{R}}(D \cup Z))$. From the fact that the class $[\xi]$ is real, we have $[\xi] = [\eta]$. Hence by Proposition 5.6, there exists $\alpha' \in W_0 R^0(U, \mathbb{L}\text{og}(1))$ such that $\nabla(\alpha') = \eta - \xi$. Since

$$\nabla(\alpha - \alpha'|_U) = \nabla(\alpha) - \nabla(\alpha'|_U) = 0,$$

the section $w := \alpha - \alpha'|_U \in \Gamma(U, \mathcal{L}\text{og}(1) \otimes \mathcal{E}_{U, \mathbb{R}})$ defines a class in $\mathbb{H}^0(U, \mathcal{L}\text{og}(1) \otimes \mathcal{E}_{U, \mathbb{R}}^\bullet) = H^0(U, \mathbb{L}\text{og}(1))$.

By Proposition 3.4, we have $H^0(U, \mathbb{L}\text{og}(1)) = 0$, hence this shows that $w = 0$, which implies that $\alpha'|_U = \alpha$. This implies that α extends to an element of $W_0 R^0(U, \mathbb{L}\text{og}(1))$ as desired. \square

Using this result, we may recover the well-known result concerning the specialization of the polylogarithm class to torsion points of \mathbb{G}_m . Let $d > 1$ be an integer and let ζ be a primitive d -th root of unity. Let $\mathbb{R}(n) = (V_{\mathbb{R}}, W_{\bullet}, F^{\bullet})$ with $V_{\mathbb{R}} := \mathbb{R}\underline{u}^n$, $W_0 V_{\mathbb{R}} = V_{\mathbb{R}}$ if $n \geq 0$ and $F^0 V_{\mathbb{C}} = 0$ if $n > 0$ be the Tate object of Example 2.2. The inclusion $i_{\zeta} : \text{Spec } \mathbb{C} \rightarrow U$ defines the restriction map

$$i_{\zeta}^* : H_{\mathcal{A}}^1(U, \text{Log}(1)) \rightarrow H_{\mathcal{A}}^1(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}(1)) \cong \prod_{k=0}^{\infty} H_{\mathcal{A}}^1(\text{Spec } \mathbb{C}, \mathbb{R}(k+1)),$$

where the last equality follows from the splitting principle (3.3). For any mixed \mathbb{R} -Hodge structure $V = (V_{\mathbb{R}}, W_{\bullet}, F^{\bullet})$, we have by definition

$$H_{\mathcal{A}}^m(\text{Spec } \mathbb{C}, V) := \text{Ext}_{\text{MHS}_{\mathbb{R}}}^m(\mathbb{R}(0), V) = H^m \left(\text{Cone} \left(W_0 V_{\mathbb{R}} \oplus (F^0 \cap W_0) V_{\mathbb{C}} \rightarrow W_0 V_{\mathbb{C}} \right) [-1] \right).$$

For $\mathbb{R}(n) = (V_{\mathbb{R}}, W_{\bullet}, F^{\bullet})$, if $n > 0$ then we have

$$H_{\mathcal{A}}^1(\text{Spec } \mathbb{C}, \mathbb{R}(n)) \cong W_0 V_{\mathbb{C}} / W_0 V_{\mathbb{R}} = \mathbb{C}\underline{\omega}^n / \mathbb{R}\underline{u}^n \cong \mathbb{C} / (2\pi i)^n \mathbb{R}.$$

Our calculation gives the following:

Corollary 5.10. *Let $d > 1$ be an integer and let ζ be a primitive d -th root of unity. Then we have*

$$i_{\zeta}^* \mathbf{pol} = ((-1)^k \text{Li}_{k+1}(\zeta))_{k=0}^{\infty} \in \prod_{k=0}^{\infty} \mathbb{C} / (2\pi i)^{k+1} \mathbb{R}.$$

Proof. By Theorem 5.9, the class $i_{\zeta}^* \mathbf{pol}$ is represented by the triple

$$\begin{aligned} (i_{\zeta}^* \alpha, i_{\zeta}^* \eta, i_{\zeta}^* \xi) &\in W_0 R^0(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{C}}(1)) \oplus W_0 R^1(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{R}}(1)) \\ &\quad \oplus (F^0 \cap W_0) R^1(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{C}}(1)). \end{aligned}$$

However, since the dimension of $\text{Spec } \mathbb{C}$ is zero, we have

$$R^1(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{R}}(1)) = R^1(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{C}}(1)) = 0.$$

Hence $i_{\zeta}^* \mathbf{pol}$ is represented by the section

$$i_{\zeta}^* \alpha = \sum_{m=0}^{\infty} D_{m+1}(\zeta) \underline{e}^m(1) \in W_0 R^0(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{C}}(1)) := \Gamma(\text{Spec } \mathbb{C}, i_{\zeta}^* \text{Log}_{\mathbb{C}}(1)) = \prod_{m=0}^{\infty} \mathbb{C} \underline{e}^m(1).$$

Note that since $\log |\zeta| = 0$, we have $L_{m+1}(\zeta) = \text{Li}_{m+1}(\zeta)$ for any $m \geq 0$. By definition,

$$\begin{aligned} D_{m+1}(\zeta) \underline{e}^m(1) &= \text{Im}(i^{m+1} L_{m+1}(\zeta)) \underline{e}^m(1) = \text{Im}(i^{m+1} \text{Li}_{m+1}(\zeta)) \underline{e}^m(1) \\ &= \text{Re}(i^m \text{Li}_{m+1}(\zeta)) \underline{e}^m(1). \end{aligned}$$

hence we have

$$D_{m+1}(\zeta) \underline{e}^m(1) \equiv i^m \text{Li}_{m+1}(\zeta) \underline{e}^m(1) \pmod{\mathbb{R} i \underline{e}^m(1)}.$$

Noting that $i^m \underline{e}^m(1) = (-1)^m \underline{\omega}^{m+1}$ and $\mathbb{R} i \underline{e}^m(1) = \mathbb{R} \underline{u}^{m+1}$, we see that $i_{\zeta}^* \alpha$ coincides with the class of

$$\sum_{m=0}^{\infty} (-1)^m \text{Li}_{m+1}(\zeta) \underline{\omega}^{m+1}$$

in $\prod_{m=0}^{\infty} \mathbb{C} \underline{\omega}^{m+1} / \mathbb{R} \underline{u}^{m+1}$ as desired. \square

5.3. The Polylogarithm Class for the Case $g > 1$. We consider the case when $X = \mathbb{G}_m^g$ for an integer $g > 1$. For $\mu \in [g]$, we let α_μ and ξ_μ be the elements given by

$$\alpha_\mu := \sum_{m=0}^{\infty} D_{m+1}(t_\mu) \underline{e}^{m_\mu}(1), \quad \xi_\mu := \underline{\omega}^0(1) \otimes \frac{dt_\mu}{t_\mu - 1},$$

where m_μ is the element in \mathbb{N}^g with m in the μ -th component and 0 in the other components. Then the differential form ξ of (3.18) is given by

$$\xi = \xi_1 \wedge \cdots \wedge \xi_g \in (F^0 \cap W_0)R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g)).$$

We let

$$\eta := \text{Re}(\xi) = (\xi + \bar{\xi})/2 \in W_0R^{2g-1}(U, \mathbb{L}\text{og}(g)).$$

Then we have $\eta - \xi = (\bar{\xi} - \xi)/2$. If we let

$$\alpha := \sum_{\mu=1}^g (-1)^{\mu-1} \alpha_\mu \xi_1 \wedge \cdots \wedge \xi_{\mu-1} \wedge \overline{\xi_{\mu+1}} \wedge \cdots \wedge \overline{\xi_g} \in W_0R^{2g-2}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g)),$$

then we have

$$\begin{aligned} \nabla(\alpha) &= \sum_{\mu=1}^g (-1)^{\mu-1} \frac{\bar{\xi}_\mu - \xi_\mu}{2} \wedge \xi_1 \wedge \cdots \wedge \xi_{\mu-1} \wedge \overline{\xi_{\mu+1}} \wedge \cdots \wedge \overline{\xi_g} \\ &= \frac{1}{2} \sum_{\mu=1}^g \left(\xi_1 \wedge \cdots \wedge \xi_{\mu-1} \wedge \overline{\xi_\mu} \wedge \overline{\xi_{\mu+1}} \wedge \cdots \wedge \overline{\xi_g} - \xi_1 \wedge \cdots \wedge \xi_{\mu-1} \wedge \xi_\mu \wedge \overline{\xi_{\mu+1}} \wedge \cdots \wedge \overline{\xi_g} \right) \\ &= \frac{1}{2} \left(\overline{\xi_1} \wedge \cdots \wedge \overline{\xi_g} - \xi_1 \wedge \cdots \wedge \xi_g \right) = \frac{1}{2} (\bar{\xi} - \xi) = \eta - \xi, \end{aligned}$$

where the calculation of $\nabla(\alpha_\mu)$ follows from (5.8).

This proves our main result.

Theorem 5.11 (Case $g > 1$). *We let α , η and ξ be as above. Then the triple*

$$(\alpha, \eta, \xi) \in W_0R^{2g-2}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g)) \oplus W_0R^{2g-1}(U, \mathbb{L}\text{og}(g)) \oplus (F^0 \cap W_0)R^{2g-1}(U, \mathbb{L}\text{og}_{\mathbb{C}}(g))$$

represents the polylogarithm class \mathbf{pol} in $H_{\mathcal{A}}^{2g-1}(U, \mathbb{L}\text{og}(g))$.

REFERENCES

- [BKT10] Kenichi Bannai, Shinichi Kobayashi, and Takeshi Tsuji, *On the de Rham and p -adic realizations of the elliptic polylogarithm for CM elliptic curves*, Ann. Sci. Éc. Norm. Supér. (4) **43** (2010), no. 2, 185–234, DOI 10.24033/asens.2119 (English, with English and French summaries). MR2662664
- [Bei86] A. A. Beilinson, *Notes on absolute Hodge cohomology*, Applications of algebraic K -theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), Contemp. Math., vol. 55, Amer. Math. Soc., Providence, RI, 1986, pp. 35–68. MR862628
- [Blo09] David Blottière, *Réalisation de Hodge du polylogarithme d'un schéma abélien*, J. Inst. Math. Jussieu **8** (2009), no. 1, 1–38, DOI 10.1017/S1474748008000315 (French, with English and French summaries). MR2461901
- [BL94] A. A. Beilinson and A. Levin, *The elliptic polylogarithm*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 123–190. MR1265553
- [Bur94] José Ignacio Burgos, *A \mathcal{C}^∞ logarithmic Dolbeault complex*, Compositio Math. **92** (1994), no. 1, 61–86. MR1275721
- [Del70] Pierre Deligne, *Équations différentielles à points singuliers réguliers*, Lecture Notes in Mathematics, Vol. 163, Springer-Verlag, Berlin-New York, 1970 (French). MR0417174
- [Del71] ———, *Théorie de Hodge. II*, Inst. Hautes Études Sci. Publ. Math. **40** (1971), 5–57 (French). MR0498551
- [Del74] ———, *Théorie de Hodge. III*, Inst. Hautes Études Sci. Publ. Math. **44** (1974), 5–77 (French). MR0498552

- [HZ87] Richard M. Hain and Steven Zucker, *Unipotent variations of mixed Hodge structure*, Invent. Math. **88** (1987), no. 1, 83–124. MR877008
- [HW98] Annette Huber and Jörg Wildeshaus, *Classical motivic polylogarithm according to Beilinson and Deligne*, Doc. Math. **3** (1998), 27–133. MR1643974
- [HK18] Annette Huber and Guido Kings, *Polylogarithm for families of commutative group schemes*, J. Algebraic Geom. **27** (2018), 449–495.
- [KS94] Masaki Kashiwara and Pierre Schapira, *Sheaves on manifolds*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 292, Springer-Verlag, Berlin, 1994. With a chapter in French by Christian Houzel; Corrected reprint of the 1990 original. MR1299726
- [Kin09] Guido Kings, *A note on polylogarithms on curves and abelian schemes*, Math. Z. **262** (2009), no. 3, 527–537, DOI 10.1007/s00209-008-0387-5. MR2506305
- [Lev97] A. Levin, *Elliptic polylogarithms: an analytic theory*, Compositio Math. **106** (1997), no. 3, 267–282, DOI 10.1023/A:1000193320513. MR1457106
- [Ram86] Dinakar Ramakrishnan, *Analogues of the Bloch-Wigner function for higher polylogarithms*, Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), Contemp. Math., vol. 55, Amer. Math. Soc., Providence, RI, 1986, pp. 371–376, DOI 10.1090/conm/055.1/862642. MR862642
- [Wil97] Jörg Wildeshaus, *Realizations of polylogarithms*, Lecture Notes in Mathematics, vol. 1650, Springer-Verlag, Berlin, 1997. MR1482233
- [Zag90] Don Zagier, *The Bloch-Wigner-Ramakrishnan polylogarithm function*, Math. Ann. **286** (1990), no. 1-3, 613–624. MR1032949

E-mail address: bannai@math.keio.ac.jp

*DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, KEIO UNIVERSITY, 3-14-1 HIYOSHI, KOUHOKU-KU, YOKOHAMA 223-8522, JAPAN

◇MATHEMATICAL SCIENCE TEAM, RIKEN CENTER FOR ADVANCED INTELLIGENCE PROJECT (AIP), 1-4-1 NIHONBASHI, CHUO-KU, TOKYO 103-0027, JAPAN