

AN INVARIANT SUBBUNDLE OF THE KZ CONNECTION MOD p AND REDUCIBILITY OF $\widehat{\mathfrak{sl}}_2$ VERMA MODULES MOD p

ALEXANDER VARCHENKO*

*Department of Mathematics, University of North Carolina at Chapel Hill
Chapel Hill, NC 27599-3250, USA*

*Faculty of Mathematics and Mechanics, Lomonosov Moscow State University
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia*

Key words: KZ equations, reduction to characteristic p , arithmetic solutions

2010 Mathematics Subject Classification: 13A35 (33C60, 32G20)

ABSTRACT. We consider the KZ differential equations over \mathbb{C} in the case, when its multi-dimensional hypergeometric solutions are one-dimensional integrals. We also consider the same differential equations over a finite field \mathbb{F}_p . We study the space of polynomial solutions of these differential equations over \mathbb{F}_p , constructed in a previous work by V. Schechtman and the author. The module of these polynomial solutions defines an invariant subbundle of the associated KZ connection modulo p . We describe the algebraic equations for that subbundle and argue that the equations correspond to highest weight vectors of the associated $\widehat{\mathfrak{sl}}_2$ Verma modules over the field \mathbb{F}_p .

CONTENTS

1. Introduction	2
2. KZ equations	2
2.1. Description of equations	2
2.2. Solutions over \mathbb{C}	3
2.3. Solutions over \mathbb{F}_p	3
3. Module of arithmetic solutions	4
3.1. Auxiliary fields and rings	4
3.2. Auxiliary lemma	5
3.3. Vector subspace $\mathcal{M}_{\mathbb{F}_p(z)}$	5
3.4. Annihilator of $\mathcal{M}_{\mathbb{F}_p(z)}$	6
4. Spanning elements of $\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq M_n})$	7
4.1. Formula (3.11)	10
4.2. Corollary of Theorems 3.6, 4.3, 4.4	10
5. The space $\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq M_n})$ and coinvariants mod p	11
5.1. Complex (Ω, ∂) over \mathbb{C}	11
5.2. Lie algebra $\widehat{\mathfrak{sl}}_2$	11
5.3. Verma modules	11
5.4. Lie algebra $\mathfrak{sl}_2(z)$	12
5.5. Complexes (5.2) and (5.8)	12

* *E-mail:* anv@email.unc.edu, supported in part by NSF grant DMS-1665239

1. INTRODUCTION

The KZ connection is a flat connection on a trivial complex vector bundle, whose fiber is the tensor product of Lie algebra modules, see [KZ, EFK]. The connection has an important invariant subbundle of conformal blocks defined in conformal field theory. The algebraic equations for this subbundle are useful for applications. The algebraic equations were described in [FSV1, FSV2, FSV3]. In [SV1, SV2] flat sections of the KZ connection were constructed in the form of multidimensional hypergeometric integrals. It was shown in [FSV1, FSV2, FSV3] that these hypergeometric flat sections always are sections of the subbundle of conformal blocks.

Recently in [SV4] the KZ connection was considered over a finite field \mathbb{F}_p and polynomial flat sections of the KZ connection were constructed as p -analogs of the hypergeometric flat sections of the KZ connection over the field \mathbb{C} , see also [V3, V4, V5, Sliv2]. We call these sections the *arithmetic flat sections* of the KZ connection over \mathbb{F}_p .

The problem is to describe the algebraic equations for the subbundle spanned by the arithmetic flat sections. We consider the particular case of the KZ connection in which the hypergeometric flat sections over the field \mathbb{C} are given by one-dimensional integrals. We identify the annihilator of the subbundle of arithmetic flat sections with a certain space of rational functions in one variable x over the field $\mathbb{F}_p(z)$. That space is related to the De Rham complex of rational differential forms on the curve of genus zero with punctures. That De Rham complex over the field \mathbb{C} was studied in [SV3] to relate it to the highest weight vectors in reduced Verma modules over the complex affine Lie algebra $\widehat{\mathfrak{sl}}_2$.

In the end of the paper we argue that the annihilator of the module of arithmetic flat sections is related to highest weight vectors in the Verma modules of $\widehat{\mathfrak{sl}}_2$ considered over the field \mathbb{F}_p .

The paper is organized as follows. In Section 2 we describe our example of the KZ differential equations, construct its solutions over \mathbb{C} and \mathbb{F}_p . In Section 3 we discuss the annihilator of the module of arithmetic flat sections and identify it with a certain space of rational functions in one variable x . In Section 4 we describe a set of generators in the annihilator, which gives us a set of explicit algebraic equations satisfied by the arithmetic flat sections. In Section 5 we argue that the algebraic equations for arithmetic flat sections are related to highest weight vectors in the Verma modules of the affine Lie algebra $\widehat{\mathfrak{sl}}_2$ considered over \mathbb{F}_p .

The author thanks Vadim Schechtman and Alexey Slinkin for useful discussions.

2. KZ EQUATIONS

2.1. Description of equations. In this paper the numbers p, q are prime numbers, n a positive integer, $p > n$, $n = kq + 1$ for some $k \in \mathbb{Z}_{>0}$. We study the following system of

equations for a column vector $I(z) = (I_1(z), \dots, I_n(z))$:

$$(2.1) \quad \frac{\partial I}{\partial z_i} = \frac{1}{q} \sum_{j \neq i} \frac{\Omega_{ij}}{z_i - z_j} I, \quad i = 1, \dots, n, \quad I_1(z) + \dots + I_n(z) = 0,$$

where $z = (z_1, \dots, z_n)$,

$$\Omega_{ij} = \begin{pmatrix} & \vdots & & \vdots & \\ i \cdots & -1 & \cdots & 1 & \cdots \\ & \vdots & & \vdots & \\ j \cdots & 1 & \cdots & -1 & \cdots \\ & \vdots & & \vdots & \end{pmatrix},$$

and all other entries of Ω_{ij} are zero. This joint system of *differential and algebraic equations* is called the *system of KZ equations* in this paper.

System (2.1) is the system of the classical KZ differential equations associated with the Lie algebra \mathfrak{sl}_2 and the subspace of highest weight vectors of weight $n - 2$ of the tensor product $V^{\otimes n}$, where V is the irreducible two-dimensional \mathfrak{sl}_2 -module, up to a gauge transformation, see this example in [V2, Section 1.1]. See also [V1, SliV2].

We consider system (2.1) over the field \mathbb{C} and the field \mathbb{F}_p with p elements.

2.2. Solutions over \mathbb{C} . All solutions of (2.1) have the form

$$(2.2) \quad I^{(\gamma)}(z) = \left(\int_{\gamma} \frac{\Phi_{\mathbb{C}}(x, z)}{x - z_1} dx, \dots, \int_{\gamma} \frac{\Phi_{\mathbb{C}}(x, z)}{x - z_n} dx \right),$$

where

$$(2.3) \quad \Phi_{\mathbb{C}}(x, z) = \prod_{a=1}^n (x - z_a)^{-1/q}$$

and γ is an element of the first homology group of the algebraic curve with affine equation

$$y^q = (x - z_1) \dots (x - z_n).$$

Starting from such γ , chosen for given $\{z_1, \dots, z_n\} \subset \mathbb{C}$, the vector $I^{(\gamma)}(z)$ can be analytically continued, as a multivalued holomorphic function of z , to the complement in \mathbb{C}^n to the union of the diagonal hyperplanes $z_i = z_j$.

The complex vector space of such integral solutions is the $n - 1$ -dimensional vector space of all solutions of system (2.1). See these statements in the example in [V2, Section 1.1], also in [SliV2].

2.3. Solutions over \mathbb{F}_p . Polynomial solutions of the general KZ differential equations over \mathbb{F}_p were constructed in [SV4]. The particular case of system (2.1) was studied in [SliV2]. For $q = 2$ system (2.1) was studied in [V5].

Denote by a the unique integer such that

$$(2.4) \quad 1 \leq a < q \quad \text{and} \quad q \mid (ap - 1).$$

Denote

$$(2.5) \quad M = (ap - 1)/q \in \mathbb{Z}_{>0}.$$

Let

$$(2.6) \quad \Phi(x, z) := \prod_{i=1}^n (x - z_i)^M,$$

$$(2.7) \quad P(x, z) := \left(\frac{\Phi(x, z)}{x - z_1}, \dots, \frac{\Phi(x, z)}{x - z_n} \right) = \sum_i P^i(z) x^i,$$

where $P^i(z)$ are n -vectors of polynomials in z_1, \dots, z_n with coefficients in \mathbb{F}_p .

Theorem 2.1 ([SV4, Theorem 1.2]). *For any positive integer l , the vector of polynomials $P^{lp-1}(z)$ is a solution of system (2.1).*

Theorem 2.1 is a particular case of [SV4, Theorem 2.4]. Cf. Theorem 2.1 in [K]. See also [V3, V4, V5, Sliv2].

The solutions $P^{lp-1}(z)$ given by this construction are called the *arithmetic solutions* of system (2.1). They are p -analogs of the hypergeometric solutions (2.2), in which the integration over a cycle γ in the integrals of formula (2.2) is replaced by taking the coefficient of x^{lp-1} in the Taylor expansion of the integrands.

Denote $\mathbb{F}_p[z^p] := \mathbb{F}_p[z_1^p, \dots, z_n^p]$. The set of all polynomial solutions of system (2.1) with coefficients in \mathbb{F}_p is a module over the ring $\mathbb{F}_p[z^p]$ since equations (2.1) are linear and $\frac{\partial z_i^p}{\partial z_j} = 0$ in $\mathbb{F}_p[z]$ for all i, j .

The $\mathbb{F}_p[z^p]$ -module

$$(2.8) \quad \mathcal{M} = \left\{ \sum_l c_l(z) P^{lp-1}(z) \mid c_l(z) \in \mathbb{F}_p[z^p] \right\},$$

spanned by arithmetic solutions, is called the *module of arithmetic solutions*.

The range for the index l is defined by the inequalities $0 < lp - 1 \leq nM - 1$. This implies that $l = 1, \dots, ak$, see [Sliv2, Lemma 5.1].

Theorem 2.2 ([Sliv2, Theorem 3.2]). *The $\mathbb{F}_p[z^p]$ -module \mathcal{M} of arithmetic solutions is free of rank ak , that is, if $\sum_{l=1}^{ak} c_l(z) P^{lp-1}(z) = 0$ for some $c_l(z) \in \mathbb{F}_p[z^p]$, then $c_l(z) = 0$ for all l .*

3. MODULE OF ARITHMETIC SOLUTIONS

3.1. Auxiliary fields and rings. For variables $u = (u_1, \dots, u_r)$, denote by $\mathbb{F}_p(u) = \mathbb{F}_p(u_1, \dots, u_r)$ the field of rational functions in u_1, \dots, u_r with coefficients in \mathbb{F}_p . Denote by $\mathbb{F}_p(u)[x]$ the ring of polynomials in x with coefficients in $\mathbb{F}_p(u)$. Denote by $\mathbb{F}_p(u^p) \subset \mathbb{F}_p(u)$ the subfield of rational functions in u_1^p, \dots, u_r^p .

3.2. Auxiliary lemma.

Lemma 3.1. *Let $f(t) \in \mathbb{F}_p(t)$ be a rational function in one variable t such that $f'(t) := \frac{df}{dt}(t) = 0$, then $f(t) \in \mathbb{F}_p(t^p)$.*

Proof. For $g, h \in \mathbb{F}_p(t)$ denote

$$\text{Wr}(g, h) = gh' - g'h,$$

the Wronskian of g, h . If $a, b, c, d \in \mathbb{F}_p(t^p)$, then

$$(3.1) \quad \text{Wr}(ag + bh, cg + dh) = (ad - bc) \text{Wr}(g, h).$$

If $f = g/h$ with $g, h \in \mathbb{F}_p[t]$, then $f' = \text{Wr}(g, h)/g^2$ and $\text{Wr}(g, h) = 0$.

Using (3.1) we show that there exists $j(t) \in \mathbb{F}_p[t]$ and $a, b \in \mathbb{F}_p(t^p)$ such that $g = aj$, $h = bj$ and hence $g/h = a/b \in \mathbb{F}_p(t^p)$.

Indeed, let $g(t) = gt^l + \cdots + g_0$, $g_i \in \mathbb{F}_p$, $g_l \neq 0$, and $h(t) = h_mt^m + \cdots + h_0$, $h_i \in \mathbb{F}_p$, $h_m \neq 0$. Equation $\text{Wr}(g, h) = 0$ implies that $p \mid (l - m)$.

If $l \geq m$, consider the polynomial $\tilde{g}(t) = g(t) - t^{l-m} \frac{g_l}{h_m} h(t)$. Then $\deg \tilde{g}(t) < \deg g(t)$ and $\text{Wr}(\tilde{g}, h) = \text{Wr}(g, h)$.

If $m \geq l$, consider the polynomial $\tilde{h}(t) = h(t) - t^{m-l} \frac{h_m}{g_l} g(t)$. Then $\deg \tilde{h}(t) < \deg h(t)$ and $\text{Wr}(g, \tilde{h}) = \text{Wr}(g, h)$.

We iterate this procedure, which decreases the degree of one of the two polynomials h, g by a multiple of p while keeping the Wronskian of the two polynomials equal to zero. The procedure will stop when one of the polynomials becomes zero. That means that the last nonzero polynomial $j(t) \in \mathbb{F}_p[t]$ divides the initial polynomials $g(t), h(t)$ and the ratios $a(t) := g(t)/j(t)$ and $b(t) := h(t)/j(t)$ belong to $\mathbb{F}_p[t^p]$. \square

Corollary 3.2. *Let $f(u_1, \dots, u_r) \in \mathbb{F}_p(u_1, \dots, u_r)$ be such that $\frac{\partial f}{\partial u_i} = 0$, $i = 1, \dots, r$. Then $f(u_1, \dots, u_r) \in \mathbb{F}_p(u_1^p, \dots, u_r^p)$.* \square

3.3. Vector subspace $\mathcal{M}_{\mathbb{F}_p(z)}$. Let

$$\mathcal{M}_{\mathbb{F}_p(z)} := \mathcal{M} \otimes_{\mathbb{F}_p[z^p]} \mathbb{F}_p(z) \subset \mathbb{F}_p(z)^n,$$

be the $\mathbb{F}_p(z)$ -vector subspace of $\mathbb{F}_p(z)^n$ spanned by arithmetic solutions.

Recall that the *KZ connection* is defined by the commuting differential operators ∇_i , $i = 1, \dots, n$,

$$(3.2) \quad \nabla_i := \frac{\partial}{\partial z_i} - \frac{1}{q} \sum_{j \neq i} \frac{\Omega_{ij}}{z_i - z_j} : \mathbb{F}_p(z)^n \rightarrow \mathbb{F}_p(z)^n.$$

The arithmetic solutions $P^{lp-1}(z)$ are flat sections of the KZ connection.

Lemma 3.3. *The subspace $\mathcal{M}_{\mathbb{F}_p(z)}$ is invariant with respect to the KZ connection, namely, for any element $I = \sum_{l=1}^{ak} c_l(z) P^{lp-1}(z)$, $c_l(z) \in \mathbb{F}_p(z)$, of $\mathcal{M}_{\mathbb{F}_p(z)}$ we have*

$$\nabla_i I = \sum_{l=1}^{ak} \frac{\partial c_l}{\partial z_i}(z) P^{lp-1}(z) \in \mathcal{M}_{\mathbb{F}_p(z)}, \quad i = 1, \dots, n.$$

\square

Lemma 3.4. *Let $I^1(z), \dots, I^l(z)$ be flat sections of the KZ connection linearly independent over the field $\mathbb{F}_p(z^p)$. Then the $\mathbb{F}_p(z)$ -vector subspace of $\mathbb{F}_p(z)^n$ spanned by $I^1(z), \dots, I^l(z)$ is of dimension l .*

Proof. The proof is by induction. The statement is true for $l = 1$. Assume that it is true for all $l < m$ and prove it for $l = m$. Assume that $I^1(z), \dots, I^m(z)$ are linearly independent over $\mathbb{F}_p(z^p)$ but dependent over $\mathbb{F}_p(z)$:

$$(3.3) \quad c_1(z)I^1(z) + \dots + c_m(z)I^m(z) = 0$$

for some $c_j(z) \in \mathbb{F}_p(z)$ not all equal to zero. If at least one of the coefficients $c_j(z)$ is zero, then (3.3) contradicts to the induction assumption. Hence all $c_j(z)$ are nonzero. Divide (3.3) by $c_1(z)$ and obtain the new relation of the form

$$(3.4) \quad I^1(z) + c_2(z)I^2(z) + \dots + c_m(z)I^m(z) = 0.$$

Apply ∇_i to (3.4) and obtain

$$(3.5) \quad \frac{\partial c_2}{\partial z_i} I^2(z) + \dots + \frac{\partial c_m}{\partial z_i} I^m(z) = 0.$$

If $\frac{\partial c_j}{\partial z_i} \neq 0$ for some i and j , then equation (3.5) for that i contradicts to the induction assumption. If $\frac{\partial c_j}{\partial z_i} = 0$ for all i, j , then $c_j(z) \in \mathbb{F}_p(z^p)$ for all j by Lemma 3.1. Hence (3.3) gives a linear dependence of $I^1(z), \dots, I^m(z)$ over $\mathbb{F}_p(z^p)$, which contradicts to the assumptions of the lemma. \square

Corollary 3.5. *We have $\dim_{\mathbb{F}_p(z)} \mathcal{M}_{\mathbb{F}_p(z)} = \text{rank}_{\mathbb{F}_p[z^p]} \mathcal{M} = ak$.*

3.4. Annihilator of $\mathcal{M}_{\mathbb{F}_p(z)}$. Consider the $\mathbb{F}_p(z)$ -vector space $(\mathbb{F}_p(z)^n)^*$ dual to $\mathbb{F}_p(z)^n$ and the subspace

$$\text{Ann}(\mathcal{M}_{\mathbb{F}_p(z)}) = \{c \in (\mathbb{F}_p(z)^n)^* \mid \langle c, I \rangle = 0 \ \forall I \in \mathcal{M}_{\mathbb{F}_p(z)}\}.$$

This subspace is identified with the subspace

$$(3.6) \quad \left\{ (c_1(z), \dots, c_n(z)) \in \mathbb{F}_p(z)^n \mid \sum_{j=1}^n c_j(z) P_j^{lp-1}(z) = 0, \ l = 1, \dots, ak \right\},$$

where $P^{lp-1}(z) = (P_1^{lp-1}(z), \dots, P_n^{lp-1}(z))$, $l = 1, \dots, ak$, are arithmetic solutions.

We have

$$(3.7) \quad \dim_{\mathbb{F}_p(z)} \text{Ann}(\mathcal{M}_{\mathbb{F}_p(z)}) = n - ak = (q - a)k + 1,$$

by Corollary 3.5.

Theorem 3.6. *We have $(c_1(z), \dots, c_n(z)) \in \text{Ann}(\mathcal{M}_{\mathbb{F}_p(z)})$ if and only if there exists a polynomial $Q(x, z) \in \mathbb{F}_p(z)[x]$ such that*

$$(3.8) \quad \frac{\partial Q}{\partial x}(x, z) = \sum_{j=1}^n c_j(z) \frac{\Phi(x, z)}{x - z_j}.$$

Proof. The polynomial $Q(x, z)$ with property (3.8) exists if and only if the Taylor expansion of the polynomial $\sum_{j=1}^n c_j(z) \frac{\Phi(x, z)}{x-z_j}$ with respect to x has zero coefficients for all of the monomials x^{lp-1} , $l = 1, \dots, ak$. That property is equivalent to the property that $(c_1(z), \dots, c_n(z)) \in \text{Ann}(\mathcal{M}_{\mathbb{F}_p(z)})$. \square

For $m \in \mathbb{Z}_{\geq 0}$ denote by $\mathbb{F}_p(z)[x]_{\leq m}$ the $\mathbb{F}_p(z)$ -vector space of polynomials in $\mathbb{F}_p(z)[x]$ of degree $\leq m$ in x .

Clearly, $\deg_x \frac{\Phi(x, z)}{x-z_j} = Mn - 1$, where M is defined in (2.5). Hence a polynomial $Q(x, z)$ in Theorem 3.6, if exists, can be chosen to be in $\mathbb{F}_p(z)[x]_{\leq Mn}$.

Define the map

$$(3.9) \quad \partial : \mathbb{F}_p(z)[x]_{\leq Mn} \rightarrow \mathbb{F}_p(z)[x]_{\leq Mn-1}, \quad Q(x, z) \mapsto \frac{\partial Q}{\partial x}(x, z).$$

Define the n -dimensional vector space

$$\Omega_{\log} = \left\{ \sum_{j=1}^n c_j(z) \frac{\Phi(x, z)}{x-z_j} \mid (c_1(z), \dots, c_n(z)) \in \mathbb{F}_p(z)^n \right\},$$

the span of the polynomials $\frac{\Phi(x, z)}{x-z_j}$, $j = 1, \dots, n$.

Corollary 3.7. *Theorem (3.6) gives an isomorphism*

$$(3.10) \quad \text{Ann}(\mathcal{M}_{\mathbb{F}_p(z)}) \simeq \Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq Mn}).$$

\square

Corollary 3.8. *We have*

$$(3.11) \quad \dim_{\mathbb{F}_p(z)} \left(\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq Mn}) \right) = (q - a)k + 1.$$

Corollary 3.8 follows from Corollaries 3.7, 3.5, and Theorem 2.2.

Remark. The map $\partial : \mathbb{F}_p(z)[x]_{\leq Mn} \rightarrow \mathbb{F}_p(z)[x]_{\leq Mn-1}$ in (3.9) is a part of the larger De Rham complex

$$(3.12) \quad \Omega \xrightarrow{\partial} \Omega,$$

where Ω is the $\mathbb{F}_p(z)$ -vector space with basis

$$(3.13) \quad \frac{\Phi(x, z)}{(x - z_i)^m} \text{ for } m \in \mathbb{Z}_{>0}, i = 1, \dots, n, \quad \text{and} \quad \Phi(x, z)x^m \text{ for } m \in \mathbb{Z}_{\geq 0},$$

and $\partial = \frac{\partial}{\partial x}$.

4. SPANNING ELEMENTS OF $\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq Mn})$

Lemma 4.1. *The polynomials*

$$(4.1) \quad \frac{\Phi(x, z)}{(x - z_j)^m}, \quad j = 1, \dots, n, \quad 0 < m \leq M,$$

form a basis of $\mathbb{F}_p(z)[x]_{\leq Mn-1}$. The polynomials in (4.1) together with the polynomial $\Phi(x, z)$ form a basis of $\mathbb{F}_p(z)[x]_{\leq Mn}$.

Proof. Each of the polynomials in (4.1) lies in $\mathbb{F}_p(z)[x]_{\leq Mn-1}$. They are linearly independent and their number Mn equals $\dim \mathbb{F}_p(z)[x]_{\leq Mn-1}$. Hence they form a basis of $\mathbb{F}_p(z)[x]_{\leq Mn-1}$. Similarly for the second part of the lemma. \square

In these bases the map ∂ defined in (3.9) is given by the formulas:

$$(4.2) \quad \frac{\partial}{\partial x} \left(\Phi(x, z) \right) = M \sum_{j=1}^n \frac{\Phi(x, z)}{x - z_j},$$

$$(4.3) \quad \frac{\partial}{\partial x} \left(\frac{\Phi(x, z)}{(x - z_i)^m} \right) = (M - m) \frac{\Phi(x, z)}{(x - z_i)^{m+1}} - M \sum_{l=1}^m \left(\sum_{j \neq i} \frac{1}{(z_j - z_i)^l} \right) \frac{\Phi(x, z)}{(x - z_i)^{m+1-l}} \\ + M \sum_{j \neq i} \frac{1}{(z_j - z_i)^m} \frac{\Phi(x, z)}{x - z_j},$$

for $i = 1, \dots, n$ and $m \geq 1$. In particular for $m = M$ we have

$$(4.4) \quad \frac{\partial}{\partial x} \left(\frac{\Phi(x, z)}{(x - z_i)^M} \right) = -M \sum_{l=1}^M \left(\sum_{j \neq i} \frac{1}{(z_j - z_i)^l} \right) \frac{\Phi(x, z)}{(x - z_i)^{M+1-l}} \\ + M \sum_{j \neq i} \frac{1}{(z_j - z_i)^M} \frac{\Phi(x, z)}{x - z_j}.$$

Lemma 4.2. *For $i = 1, \dots, n$, there exist unique $A_{i,1}(z), \dots, A_{i,M-1}(z) \in \mathbb{F}_p(z)$ such that the polynomial*

$$(4.5) \quad Q_i(x, z) = \frac{\Phi(x, z)}{(x - z_i)^M} + \sum_{m=1}^{M-1} A_{i,m}(z) \frac{\Phi(x, z)}{(x - z_i)^m}$$

in x has the property: $\frac{\partial Q_i}{\partial x} \in \Omega_{\log}$, that is,

$$(4.6) \quad \frac{\partial Q_i}{\partial x}(x, z) = \sum_{j=1}^n B_{i,j}(z) \frac{\Phi(x, z)}{x - z_j},$$

for suitable $B_{i,j}(z) \in \mathbb{F}_p(z)$.

Proof. Let $Q_i(x, z)$ be an expression as in (4.5) for some $(A_{i,m}(z))_{m=1}^{M-1}$. Applying formula (4.3) we obtain

$$\frac{\partial Q_i}{\partial x}(x, z) = \sum_{m=1}^M C_m(z) \frac{\Phi(x, z)}{(x - z_i)^m} + \sum_{j \neq i} B_{ij}(z) \frac{\Phi(x, z)}{x - z_j},$$

where $C_m(z), B_{ij}(z)$ are linear expressions in $A_{i,1}(z), \dots, A_{i,M-1}(z)$ with coefficients in $\mathbb{F}_p(z)$. We have

$$(4.7) \quad C_M(z) = A_{i,M-1}(z) - \sum_{j \neq i} \frac{M}{z_i - z_j},$$

and for $m = 2, \dots, M - 1$,

$$(4.8) \quad C_m(z) = (M - m + 1)A_{i,m-1}(z) + \dots,$$

where the dots denote a linear expression in $A_{i,m}(z), \dots, A_{i,M-1}(z)$. To obtain property (4.6) it is necessary and sufficient to choose the coefficients $A_{i,1}(z), \dots, A_{i,M-1}(z)$ so that $C_m(z) = 0$ for $m = 2, \dots, M$. But this can be done uniquely by formulas (4.7) and (4.8). \square

Denote

$$(4.9) \quad C_{i,m}(z) = \sum_{\substack{j=1, \\ j \neq i}}^n \frac{M}{(z_j - z_i)^m}, \quad D_{i,m}(x, z) = \sum_{\substack{j=1, \\ j \neq i}}^n \frac{M}{(z_j - z_i)^m} \frac{\Phi(x, z)}{x - z_j}.$$

Theorem 4.3. *For $i = 1, \dots, n$, let $Q_i(x, z)$ be the polynomial determined in Lemma 4.2. Then*

$$(4.10) \quad Q_i(x, z) = \sum_{m \geq 0} \sum_{\substack{l_0 + \dots + l_m = M, \\ l_0, \dots, l_m > 0}} \frac{\Phi(x, z)}{(x - z_i)^{l_0}} \prod_{r=1}^m \frac{C_{i,l_r}(z)}{l_1 + \dots + l_r}.$$

$$(4.11) \quad \frac{\partial Q_i}{\partial x}(x, z) = \sum_{m \geq 0} \sum_{\substack{l_0 + \dots + l_m = M, \\ l_0, \dots, l_m > 0}} \left(D_{i,l_0}(x, z) - C_{i,l_0}(z) \frac{\Phi(x, z)}{x - z_i} \right) \prod_{r=1}^m \frac{C_{i,l_r}(z)}{l_1 + \dots + l_r}.$$

This theorem is a modification of [SV3, Corollary 6.4], where formula (4.11) is considered over \mathbb{C} . Notice that formula (40) in [SV3] has misprints, cf. formulas (40) and (4.11).

Proof. Using formula (4.3) we eliminate from formula (4.4) all the terms $\frac{\Phi(x, z)}{(x - z_i)^l}$ with $l > 1$. This leads to formulas (4.10) and (4.11).

For example, if $q = 2$, $p = 5$, then $M = (p - 1)/2 = 2$. Formulas (4.4) and (4.3) take the form

$$\begin{aligned} \frac{\partial}{\partial x} \left(\frac{\Phi}{(x - z_i)^2} \right) &= -C_{i,1} \frac{\Phi}{(x - z_i)^2} - C_{i,2} \frac{\Phi}{x - z_i} + D_{i,2}, \\ \frac{\partial}{\partial x} \left(\frac{\Phi}{x - z_i} \right) &= \frac{\Phi}{(x - z_i)^2} - C_{i,1} \frac{\Phi}{x - z_i} + D_{i,1}, \end{aligned}$$

and imply the formula

$$\frac{\partial}{\partial x} \left(\frac{\Phi}{(x - z_i)^2} + C_{i,1} \frac{\Phi}{x - z_i} \right) = \left(D_{i,2} - C_{i,2} \frac{\Phi}{x - z_i} \right) + \left(D_{i,1} - C_{i,1} \frac{\Phi}{x - z_i} \right) C_{i,1}.$$

\square

Theorem 4.4. *Assume that $F(x, z) \in \mathbb{F}_p(z)[x]_{\leq Mn}$ is such that*

$$(4.12) \quad \frac{\partial F}{\partial x}(x, z) = \sum_{j=1}^n A_j(z) \frac{\Phi(x, z)}{x - z_j}$$

for some $A_j(z) \in \mathbb{F}_p(z)$. Then there exist unique $D_0(z), D_1(z), \dots, D_n(z) \in \mathbb{F}_p(z)$ such that

$$(4.13) \quad F(x, z) = D_0(z) \Phi(x, z) + \sum_{j=1}^n D_j(z) Q_j(x, z).$$

Proof. We have

$$F(x, z) = c_0(z)\Phi(x, z) + \sum_{j=1}^n \sum_{m=1}^M c_{j,m}(z) \frac{\Phi(x, z)}{(x - z_j)^m}$$

for suitable coefficients $c_0(z), c_{j,m}(z)$ in $\mathbb{F}_p(z)$. Then

$$F(x, z) - \sum_{j=1}^n c_{j,M}(z) Q_j(x, z) = c_0(z) \Phi(x, z) + \sum_{j=1}^n \sum_{m=1}^{M-1} \tilde{c}_{j,m}(z) \frac{\Phi(x, z)}{(x - z_j)^m}$$

and

$$\frac{\partial}{\partial x} \left(F(x, z) - \sum_{j=1}^n c_{j,M}(z) Q_j(x, z) \right) = \sum_{j=1}^n \tilde{A}_j(z) \frac{\Phi(x, z)}{x - z_j},$$

for suitable $\tilde{A}_j(z)$. Then formula (4.3) implies that

$$F(x, z) - \sum_{j=1}^n c_{j,M} Q_j(x, z) = c_0(z) \Phi(x, z).$$

The theorem is proved. □

4.1. Formula (3.11). Here is a proof of formula (3.11) independent of Theorem 2.2.

Denote by \mathcal{Q} the $n + 1$ -dimensional $\mathbb{F}_p(z)$ -vector space of linear combinations of the polynomials $\Phi(x, z)$ and $Q_i(x, z)$, $i = 1, \dots, n$.

By Theorem 4.4

$$\mathcal{Q} = \partial^{-1} \left(\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq Mn}) \right).$$

The kernel of the map

$$(4.14) \quad \partial|_{\mathcal{Q}} : \mathcal{Q} \rightarrow \Omega_{\log}$$

is the subspace generated by monomials x^{lp} with $0 \leq lp \leq Mn$. We have

$$0 \leq l \leq \frac{Mn}{p} = \frac{ap - 1}{pq} (kq + 1) = ak + \frac{a}{q} - \frac{kq + 1}{pq}.$$

Using the fact that $p > q$, $p > n = kq + 1$, $q > a > 0$, we conclude that $l = 0, 1, \dots, ak$. Hence the kernel of the map in (4.14) is $ak + 1$ -dimensional and its image $\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq Mn})$ is of dimension $n + 1 - (ak + 1) = (q - a)k + 1$. This statement is the statement of formula (3.11).

It is easy to see that this independent proof of formula (3.11) implies Theorem 2.2.

4.2. Corollary of Theorems 3.6, 4.3, 4.4.

Corollary 4.5. *For any arithmetic solution $P^{mp-1}(z) = (P_1^{mp-1}(z), \dots, P_n^{mp-1}(z))$, $m = 1, \dots, ak$, we have the relation*

$$(4.15) \quad P_1^{mp-1}(z) + \dots + P_n^{mp-1}(z) = 0,$$

coming from formula (4.2) and for any $i = 1, \dots, n$ the relation

$$(4.16) \quad \sum_{m \geq 0} \sum_{\substack{l_0 + \dots + l_m = M, \\ l_0, \dots, l_m > 0}} \left(\sum_{\substack{j=1, \\ j \neq i}}^n \frac{M}{(z_j - z_i)^{l_0}} P_j^{mp-1}(z) - C_{i,l_0}(z) P_i^{mp-1}(z) \right) \prod_{r=1}^m \frac{C_{i,l_r}(z)}{l_1 + \dots + l_r} = 0,$$

coming from formula (4.11). Moreover, these relations generate the space $\text{Ann}(\mathcal{M}_{\mathbb{F}_p}(z))$. \square

5. THE SPACE $\Omega_{\log} \cap \partial(\mathbb{F}_p(z)[x]_{\leq M_n})$ AND COINVARIANTS MOD p

5.1. **Complex (Ω, ∂) over \mathbb{C} .** Let $\Omega_{\mathbb{C}}$ be the $\mathbb{C}(z)$ -vector space with basis

$$(5.1) \quad \frac{\Phi(x, z)}{(x - z_i)^m} \text{ for } m \in \mathbb{Z}_{>0}, i = 1, \dots, n, \quad \text{and} \quad \Phi(x, z)x^m \text{ for } m \in \mathbb{Z}_{\geq 0},$$

where the polynomial $\Phi(x, z)$ is defined in (2.6). Consider the new complex

$$(5.2) \quad \Omega_{\mathbb{C}} \xrightarrow{\partial} \Omega_{\mathbb{C}},$$

where $\partial = \frac{\partial}{\partial x}$. Complexes (3.12) and (5.1) are given by the same formulas but are defined over different fields $\mathbb{F}_p(z)$ and $\mathbb{C}(z)$, respectively.

5.2. **Lie algebra $\widehat{\mathfrak{sl}}_2$.** Let \mathfrak{sl}_2 be the Lie algebra over $\mathbb{C}(z)$ of the 2×2 -matrices with zero trace. Let e, f, h be standard generators subject to the relations

$$[e, f] = h, \quad [h, e] = 2e, \quad [h, f] = -2f.$$

Let $\widehat{\mathfrak{sl}}_2$ be the affine Lie algebra $\widehat{\mathfrak{sl}}_2 = \mathfrak{sl}_2[T, T^{-1}] \oplus \mathbb{C}(z)c$ with the bracket

$$[aT^i, bT^j] = [a, b]T^{i+j} + i \langle a, b \rangle \delta_{i+j,0} c,$$

where c is central element, $\langle a, b \rangle = \text{tr}(ab)$. Set

$$\begin{aligned} e_1 &= e, & f_1 &= f, & h_1 &= h, \\ e_2 &= fT, & f_2 &= eT^{-1}, & h_2 &= c - h. \end{aligned}$$

These are the standard Chevalley generators defining $\widehat{\mathfrak{sl}}_2$ as the Kac-Moody algebra corresponding to the Cartan matrix $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$.

The Lie algebra $\widehat{\mathfrak{sl}}_2$ has an automorphism π ,

$$(5.3) \quad \pi : c \mapsto c, \quad eT^i \mapsto fT^i, \quad fT^i \mapsto eT^i, \quad hT^i \mapsto -hT^i.$$

5.3. **Verma modules.** We fix $K \in \mathbb{C}$ and assume that the central element c acts on all our representations by multiplication by K .

For $L \in \mathbb{C}$, let $V(L, K - L)$ be the $\widehat{\mathfrak{sl}}_2$ Verma module with generating vector v . It is the $\mathbb{C}(z)$ -vector space generated by v subject to the relations

$$e_1 v = 0, \quad e_2 v = 0, \quad h_1 v = L, \quad h_2 v = (K - L)v.$$

The Verma module $V(L, K - L)$ is reducible if and only if at least one of the following relations holds:

$$(5.4) \quad L - l + 1 + (m - 1)(K + 2) = 0,$$

$$(5.5) \quad L + l + 1 - m(K + 2) = 0,$$

$$(5.6) \quad K + 2 = 0,$$

where $l, m \in \mathbb{Z}_{>0}$, see [KK, MFF].

Let $V(L, K - L)^*$ be the $\mathbb{C}(z)$ -vector space dual to $V(L, K - L)$. The space $V(L, K - L)^*$ is an $\widehat{\mathfrak{sl}}_2$ -module with the $\widehat{\mathfrak{sl}}_2$ -action defined by the formulas:

$$\langle f_i \varphi, x \rangle = \langle \varphi, e_i x \rangle, \quad \langle e_i \varphi, x \rangle = \langle \varphi, f_i x \rangle,$$

where $\varphi \in V(L, K - L)^*$, $x \in V(L, K - L)$, $i = 1, 2$.

5.4. Lie algebra $\mathfrak{sl}_2(z)$. Let $\mathfrak{sl}_2(z)$ be the Lie algebra over $\mathbb{C}(z)$ of the \mathfrak{sl}_2 -valued rational functions in x of the form $e \otimes u_1 + h \otimes u_2 + f \otimes u_3$ with $u_i \in \Omega_{\mathbb{C}}$, and the bracket is defined by the formula $[x \otimes u_1, y \otimes u_2] = [x, y] \otimes (u_1 u_2)$.

Let W_1, \dots, W_{n+1} be $\widehat{\mathfrak{sl}}_2$ -modules. Then $\mathfrak{sl}_2(z)$ acts on $W_1 \otimes \dots \otimes W_{n+1}$ by the formula

$$(5.7) \quad \begin{aligned} y \otimes u(x) \cdot (w_1 \otimes \dots \otimes w_{n+1}) &= ([y \otimes u(x)]^{(z_1)} w_1) \otimes w_2 \otimes \dots \otimes w_{n+1} + \dots \\ &+ w_1 \otimes \dots \otimes w_{n-1} \otimes ([y \otimes u(x)]^{(z_n)} w_n) \otimes w_{n+1} \\ &+ w_1 \otimes \dots \otimes w_n \otimes (\pi([y \otimes u(x)]^{(\infty)} w_{n+1})), \end{aligned}$$

where for $y \otimes u(x) \in \mathfrak{sl}_2(z)$ the symbol $[y \otimes u(x)]^{(z_j)}$ denotes the Laurent expansion of $y \otimes u(x)$ with respect to x at $x = z_j$ for $j = 1, \dots, n$, and $[y \otimes u(x)]^{(\infty)}$ denotes the Laurent expansion of $y \otimes u(x)$ with respect to x at $x = \infty$; the symbol π in the last term denotes the $\widehat{\mathfrak{sl}}_2$ -automorphism defined in (5.3).

The $\widehat{\mathfrak{sl}}_2$ -action is the map

$$(5.8) \quad \mathfrak{sl}_2(z) \otimes \left(\otimes_{j=1}^{n+1} W_j \right) \xrightarrow{\mu} \otimes_{j=1}^{n+1} W_j,$$

denoted by μ . The quotient space $\left(\otimes_{j=1}^{n+1} W_j \right) / \text{Im}(\mu)$ is called the space of *coinvariants* or *conformal blocks* at genus 0 with marked points $(x = z_j, W_j)$, $j = 1, \dots, n$, $(x = \infty, W_{n+1})$. See this construction, for example, in [FSV1, FSV2, FSV3, SV3, Sliv1]. One defines the KZ connection on the bundle of coinvariants with respect to changing z_1, \dots, z_n . Since the objects, considered in the previous sections of this paper, are related to the KZ connection mod p it is not surprising that they are related to the space of coinvariants.

5.5. Complexes (5.2) and (5.8). Define

$$(5.9) \quad K := q - 2,$$

$$(5.10) \quad \otimes_{j=1}^{n+1} W_j := \left(V(1 - ap, q + ap - 3)^* \right)^{\otimes n} \otimes V(n - nap - 2, q + nap - n)^*.$$

In [SV3, SliV1] the commutative diagram

$$(5.11) \quad \begin{array}{ccc} \Omega_{\mathbb{C}} & \xrightarrow{\partial} & \Omega_{\mathbb{C}} \\ \nu^0 \downarrow & & \downarrow \nu^1 \\ \mathfrak{sl}_2(z) \otimes \left(\otimes_{j=1}^{n+1} W_j \right) & \xrightarrow{\mu} & \otimes_{j=1}^{n+1} W_j \end{array}$$

was defined, see there the definition of ν^0, ν^1 .

The Verma module $V(1 - ap, q + ap - 3)$, whose dual is used in the definition (5.10), is reducible, due to equation (5.4) for

$$(5.12) \quad L = 1 - ap, \quad l = 1, \quad K = q - 2, \quad m - 1 = (ap - 1)/q.$$

For $i = 1, \dots, n$, let

$$(5.13) \quad Q_i(x, z) = \sum_{m \geq 0} \sum_{\substack{l_0 + \dots + l_m = M, \\ l_0, \dots, l_m > 0}} \frac{\Phi(x, z)}{(x - z_i)^{l_0}} \prod_{r=1}^m \frac{C_{i, l_r}(z)}{l_1 + \dots + l_r} \in \mathbb{C}(z)[x].$$

This is the same polynomial as in formula (4.10) but considered over a different field. This polynomial $Q_i(x, z)$ and its derivative produce elements

$$\nu^0(Q_i(x, z)) \in \mathfrak{sl}_2(z) \otimes \left(\otimes_{j=1}^{n+1} W_j \right), \quad \nu^1\left(\frac{\partial Q_i}{\partial x}(x, z)\right) \in \otimes_{j=1}^{n+1} W_j.$$

It was explained in [SV3, SliV1] that these elements can be constructed purely in terms of action of $\widehat{\mathfrak{sl}}_2(z)$ on $\otimes_{j=1}^{n+1} W_j$ by using the fact that the module $V(1 - ap, q + ap - 3)$, corresponding to the i -th factor of this tensor product, is reducible with the reducibility condition (5.14).

Summarizing these remarks, we may conclude that the linear relations (4.16) for the arithmetic solutions of our KZ equations (2.1) correspond to diagram (5.11), in which the affine Lie algebra $\widehat{\mathfrak{sl}}_2(z)$ is reduced modulo p as well as its Verma modules.

Remark. Notice that the Verma module $V(n - nap - 2, q + nap - n)$, whose dual is used in the definition (5.10), is also reducible, due to equation (5.5) for

$$(5.14) \quad L = n - nap - 2, \quad l = 1, \quad K = q - 2, \quad m = Mn,$$

cf. [SV3, Formula (41)].

REFERENCES

- [FSV1] B.A. Feigin, V.V. Schechtman, and A. Varchenko, *On Algebraic Equations Satisfied by Correlators in WZW Models*, Letters in Math. Physics **20** (1990), 291–297
- [FSV2] B. Feigin, V. Schechtman, and A. Varchenko, *On algebraic equations satisfied by hypergeometric correlators in WZW models, I*, Comm. Math. Phys. **163** (1994), 173–184
- [FSV3] B. Feigin, V. Schechtman, and A. Varchenko, *On algebraic equations satisfied by hypergeometric correlators in WZW models, II*, Comm. In Math. Phys. **70** (1995), 219–247

- [EFK] P. Etingof, I. Frenkel, A. Kirillov, *Lectures on representation theory and Knizhnik-Zamolodchikov equations*, Mathematical Surveys and Monographs, 58, AMS, Providence, RI, 1998. xiv+198 pp. ISBN: 0-8218-0496-0
- [K] N. Katz, *Expansion-coefficients as approximate solutions of differential equations*, *Asterisque* **119-120** (1984), 183–189
- [KK] V. Kac, D. Kazhdan, *Structure of representations with highest weight of infinite dimensional Lie algebras*, *Adv. Math.* **34**(1979), 97–108
- [KZ] V. Knizhnik and A. Zamolodchikov, *Current algebra and the Wess-Zumino model in two dimensions*, *Nucl. Phys.* **B247** (1984), 83–103
- [MFF] F. Malikov, B. Feigin, D. Fuchs, *Singular vectors in Verma modules over Kac-Moody algebras*, *Funct. Anal. and Appl.* **20**, no. 2(1986), 103–113
- [SV1] V. Schechtman and A. Varchenko, *Integral Representations of N-Point Conformal Correlators in the WZW Model*, Bonn, Max-Planck Institute, 1989, 1–22
- [SV2] V. Schechtman, A. Varchenko, *Arrangements of Hyperplanes and Lie Algebra Homology*, *Invent. Math.* **106** (1991), 139–194
- [SV3] V. Schechtman, A. Varchenko, *Rational differential forms on line and singular vectors in Verma modules over $\widehat{\mathfrak{sl}}_2$* , [arXiv:1511.09014](https://arxiv.org/abs/1511.09014), *Mosc. Math. J.* **17** (2017), no. 4, 787–802
- [SV4] V. Schechtman, A. Varchenko, *Solutions of KZ differential equations modulo p* , *The Ramanujan Journal*, **48** (3), 2019, 655–683, <https://doi.org/10.1007/s11139-018-0068-x>, [arXiv:1707.02615](https://arxiv.org/abs/1707.02615)
- [SliV1] A. Slinkin, A. Varchenko, *Twisted De Rham complex on line and $\widehat{\mathfrak{sl}}_2$ -singular vectors*, *SIGMA* **15** (2019), 075, 26 pages
- [SliV2] A. Slinkin, A. Varchenko, *Hypergeometric Integrals Modulo p and Hasse–Witt matrices*, [arXiv:2001.06869](https://arxiv.org/abs/2001.06869), 1–36
- [V1] A. Varchenko, *Multidimensional Hypergeometric Functions and Representation Theory of Lie Algebras and Quantum Groups*, Advanced Series in Mathematical Physics, Vol. 21, World Scientific, 1995
- [V2] A. Varchenko, *Special functions, KZ type equations, and Representation theory*, CBMS, Regional Conference Series in Math., n. 98, AMS (2003)
- [V3] A. Varchenko, *Remarks on the Gaudin model modulo p* , [arXiv:1708.06264](https://arxiv.org/abs/1708.06264), *Journal of Singularities*, **18** (2018), 486–499
- [V4] A. Varchenko, *Solutions modulo p of Gauss-Manin differential equations for multidimensional hypergeometric integrals and associated Bethe ansatz*, [arXiv:1709.06189](https://arxiv.org/abs/1709.06189), *Mathematics* 2017, 5(4), 52; [doi:10.3390/math5040052](https://doi.org/10.3390/math5040052), 1–18
- [V5] A. Varchenko, *Hyperelliptic integrals modulo p and Cartier-Manin matrices*, [arXiv:1806.03289](https://arxiv.org/abs/1806.03289), 1–16