

# ODD-ODD CONTINUED FRACTION ALGORITHM

DONG HAN KIM, SEUL BEE LEE, AND LINGMIN LIAO

ABSTRACT. By using an accelerated algorithm of Romik’s dynamical system, we define a new continued fraction algorithm called odd-odd continued fraction, whose principal convergents are rational numbers of odd denominators and odd numerators. Among others, it is proved that all the best approximating rationals of odd denominators and odd numerators of an irrational number are given by the principal convergents of the odd-odd continued fraction algorithm and vice versa.

## 1. INTRODUCTION

An important property of the regular continued fraction algorithm is that it gives us the best approximating rational numbers of a given real number. The regular continued fraction (RCF) algorithm can be attained by an acceleration of *the Farey map* defined by

$$F(x) = \begin{cases} \frac{x}{1-x}, & 0 \leq x \leq \frac{1}{2}, \\ \frac{1-x}{x}, & \frac{1}{2} \leq x \leq 1. \end{cases}$$

Let  $n(x)$  be *the first return time* to  $[\frac{1}{2}, 1]$  defined by  $n(x) = \min \{j \geq 0 : F^j(x) \in [\frac{1}{2}, 1]\}$ . *The regular continued fraction map* or *the Gauss map* is obtained by

$$G(0) = 0, \quad \text{and} \quad G(x) = F^{n(x)+1}(x), \quad \text{for all } x \in (0, 1]$$

Equivalently, we have the following continued fraction map

$$G(0) = 0, \quad \text{and} \quad G(x) = \left\{ \frac{1}{x} \right\}, \quad \text{for all } x \in (0, 1],$$

that is the left shift map of the continued fraction expansions. We note that the first return time  $n(x)$  is exactly the first partial quotient of the RCF expansion of  $x$ .

Schweiger investigated the continued fraction with even partial quotients which we call *the even integer continued fraction (EICF)* in [15, 16]. The EICF expansion of  $x \in \mathbb{R}$  is of the form

$$x = b_0 + \frac{\eta_0}{b_1 + \frac{\eta_1}{b_2 + \ddots}}$$

where  $b_0 \in 2\mathbb{Z}$ ,  $b_i \in 2\mathbb{N}$  for  $i \geq 1$  and  $\eta_i \in \{-1, +1\}$ . The left shift map of EICF expansions is the EICF map  $T_{\text{EICF}}$  on the unit interval  $[0, 1]$  defined by

$$T_{\text{EICF}}(0) = 0, \quad \text{and} \quad T_{\text{EICF}}(x) = \begin{cases} \frac{1}{x} - 2k, & \text{if } \frac{1}{2k+1} \leq x \leq \frac{1}{2k}, \\ 2k - \frac{1}{x}, & \text{if } \frac{1}{2k} \leq x \leq \frac{1}{2k-1}, \end{cases} \quad \text{for all } k \in \mathbb{N}.$$

---

2010 *Mathematics Subject Classification*. Primary 11J70; Secondary 37E05.  
 Research partially supported by the National Research Foundation of Korea (NRF-2018R1A2B6001624).

Note that an invariant measure of  $T_{\text{EICF}}$  is  $\frac{dx}{1-x^2}$ . Similar to RCF, for the principal convergents  $p_n^E/q_n^E$  of EICF, the following matrix relation holds (see Lemma 3.1 for the details):

$$(1.1) \quad \begin{pmatrix} a_0 & \varepsilon_1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & \varepsilon_1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_n & \varepsilon_n \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} q_n & \varepsilon_n q_{n-1} \\ p_n & \varepsilon_n p_{n-1} \end{pmatrix}.$$

The group  $\text{SL}_2(\mathbb{R})$  acts on the extended complex plane  $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$  as Möbius transformations defined by

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} : z \mapsto g(z) = \frac{az + b}{cz + d}.$$

The EICF has a connection with an index 3 subgroup  $\Theta$  of  $\text{SL}_2(\mathbb{Z})$  defined by

$$\Theta = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ or } \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \pmod{2} \right\}.$$

Kraaikamp-Lopes and Boca-Merriman explored relations between EICF and geodesics on  $\Theta \backslash \mathbb{H}$  (see [12] and [4]). Since  $\Theta \backslash \mathbb{H}$  has two cusps corresponding to  $\infty$  and 1, we classify rationals into two classes by the orbits of the cusps. We call a rational in the orbit  $\Theta \cdot \infty$  an  $\infty$ -rational and a rational in the orbit  $\Theta \cdot 1$  a 1-rational. Since each matrix in (1.1) belongs to  $\Theta$ , we can easily conclude that each EICF principal convergent  $p_n^E/q_n^E$  is an  $\infty$ -rational. In this paper, we will find a continued fraction which has 1-rational principal convergents.

As we obtain the Gauss map by accelerating the Farey map, we can obtain  $T_{\text{EICF}}$  by accelerating the Romik map. In [14], Romik introduced a dynamical system on the unit interval  $[0, 1]$  as follows

$$(1.2) \quad R(x) = \begin{cases} \frac{x}{1-2x}, & 0 \leq x \leq \frac{1}{3}, \\ \frac{1}{x} - 2, & \frac{1}{3} \leq x \leq \frac{1}{2}, \\ 2 - \frac{1}{x}, & \frac{1}{2} \leq x \leq 1. \end{cases}$$

Note that the Farey map sends all rational points in  $[0, 1]$  to 0 after iterations, i.e.,  $F^n(x) = 0$  for some  $n \geq 0$  if and only if  $x$  is rational in  $[0, 1]$ . In contrary, the Romik map has two neutral fixed points and we have

$$R^n(x) = 0 \text{ for some } n \geq 0 \iff x \text{ is an } \infty\text{-rational}$$

and

$$R^n(x) = 1 \text{ for some } n \geq 0 \iff x \text{ is a 1-rational.}$$

Some number theoretical properties of the Romik map were shown in [5]. See also [6].

The EICF map is the acceleration of  $R$  to return to  $E := [\frac{1}{3}, 1]$ . In fact, we have

$$T_{\text{EICF}}(x) = R^{n_E(x)+1}(x), \quad \text{where } n_E(x) = \min\{j \geq 0 : R^j(x) \in E\}.$$

Instead of  $E$ , we choose a neighborhood  $E' = [0, \frac{1}{2}]$  of 0 which is corresponding to a cusp  $\infty$  of  $\Theta \backslash \mathbb{H}$ . Then by the acceleration of the Romik dynamical system to return to  $E'$ , we induce the odd-odd continued fraction (OOCF) map  $T_{\text{OOCF}}(x)$  on  $[0, 1]$ .

We organize our paper as follows. In Section 2, we give definitions and properties of the OOCF map  $T_{\text{OOCF}}$ . In Section 3, we study the principal convergents of the OOCF algorithms. In Section 4, we prove that all the best approximating 1-rationals are obtained by the principal convergents of the OOCF algorithm and vice versa. Finally, in Section 5, we describe relations between the expansions of OOCF and the expansions of RCF.




FIGURE 1. Continued fraction maps  $T_{\text{OOCF}}$  and  $T_{\text{EICF}}$  are conjugate via  $f(t) = \frac{1-t}{1+t}$ .

## 2. THE ODD-ODD CONTINUED FRACTION MAP

Romik originally introduced the Romik dynamical system  $\widehat{R}$  as a dynamical system on a quadrant  $\mathcal{Q}$  to investigate an algorithm generating the Pythagorean triples by multiplying matrices (see [3] and also [1], [2], [8], [7]). More precisely, the Romik map  $\widehat{R}$  on  $\mathcal{Q} = \{(x, y) : x^2 + y^2 = 1, x \geq 0, y \geq 0\}$ , is defined by

$$\widehat{R}(x, y) = \left( \frac{|2-x-2y|}{3-2x-2y}, \frac{|2-2x-y|}{3-2x-2y} \right).$$

Let us define  $D : (0, 1) \rightarrow \mathcal{Q}$  and  $\tilde{D} : (0, 1) \rightarrow \mathcal{Q}$  by

$$D(t) = \left( \frac{1-t^2}{1+t^2}, \frac{2t}{1+t^2} \right) \quad \text{and} \quad \tilde{D}(t) = \left( \frac{2t}{1+t^2}, \frac{1-t^2}{1+t^2} \right).$$

The maps  $D$  and  $\tilde{D}$  are the inverses of the stereographic projections through  $(-1, 0)$  and  $(0, -1)$ , respectively. Then the Romik map  $R$  as in (1.2) is a conjugation of  $\widehat{R}$  such that

$$(2.1) \quad R = D^{-1} \circ \widehat{R} \circ D = \tilde{D}^{-1} \circ \widehat{R} \circ \tilde{D}.$$

See Theorem 4 in [14] for more details.

Let us consider the first return time  $n_{E'}(x)$  to  $E' = [0, \frac{1}{2}]$  defined by  $n_{E'}(x) = \min\{j \geq 0 : R^j(x) \in E'\}$ . The odd-odd continued fraction map  $T_{\text{OOCF}}$  from  $[0, 1]$  to itself is defined by the following acceleration

$$T_{\text{OOCF}}(1) = 1, \quad \text{and} \quad T_{\text{OOCF}}(x) = R^{n_{E'}(x)+1}(x).$$

By simple calculation, we have

$$(2.2) \quad T_{\text{OOCF}}(1) = 1, \quad \text{and} \quad T_{\text{OOCF}}(x) = \begin{cases} \frac{kx-(k-1)}{k-(k+1)x}, & \frac{k-1}{k} \leq x \leq \frac{2k-1}{2k+1}, \\ \frac{k-(k+1)x}{kx-(k-1)}, & \frac{2k-1}{2k+1} \leq x \leq \frac{k}{k+1}, \end{cases} \quad \text{for all } k \geq 1.$$

Evidently, the map  $T_{\text{OOCF}}$  has a neutral fixed point at 0.

Let  $f(x) = \tilde{D}^{-1} \circ D(x) = \frac{1-x}{1+x}$ . We note that  $f = f^{-1}$ . As in Figure 1,  $D(E) = \{(x, y) \in \mathcal{Q} : 0 < x < \frac{4}{5}\} = \tilde{D}(E')$ . Since the conditions  $R^j(f^{-1}(t)) \in E$  and  $R^j(t) \in E'$  are equivalent to  $\widehat{R}^j \tilde{D}(t) \in D(E) = \tilde{D}(E')$  by




FIGURE 2. The graph of the Romik map  $R$  (left) and the graph of the OOCF map  $T_{\text{OOCF}}$  (right)

(2.1), we can show that  $n_E(f^{-1}(t))$  and  $n_{E'}(t)$  are the same. Thus,  $T_{\text{OOCF}}$  is conjugate to  $T_{\text{EICF}}$  since

$$f \circ T_{\text{OOCF}} \circ f^{-1}(s) = \tilde{D}^{-1} \circ \widehat{R}^{n_E(f^{-1}(s))+1} \circ \tilde{D}(s) = R^{n_{E'}(s)+1}(s) = T_{\text{EICF}}(s).$$

The above arguments are summarized as the following theorem.

**Theorem 2.1.** *The OOCF map  $T_{\text{OOCF}}$  is conjugate to  $T_{\text{EICF}}$ . More precisely, let  $f$  be the function on the interval  $[0, 1]$  defined by  $x \mapsto \frac{1-x}{1+x}$ . Then*

$$f \circ T_{\text{OOCF}} \circ f^{-1} = T_{\text{EICF}}.$$

To calculate an invariant measure, let  $y = f(x)$ . Then  $dy = |f'(x)|dx = \frac{2dx}{(1+x)^2}$ , thus

$$\frac{dx}{1-x^2} = \frac{(1+x)^2 dy}{2(1-x^2)} = \frac{(1+x)dy}{2(1-x)} = \frac{dy}{2y}.$$

Hence, we have the absolutely continuous invariant measure as follows:

**Proposition 2.2.** *The map  $T_{\text{OOCF}} : [0, 1] \rightarrow [0, 1]$  preserves the measure*

$$\frac{1}{x} dx.$$

**Remark 2.3.** In [15], Schweiger proved that  $T_{\text{EICF}}$  admits an ergodic invariant measure  $d\mu := \frac{dx}{1-x^2}$ . Thus, by Theorem 2.1, we have the measure  $f_*^{-1}\mu$  is ergodic invariant with respect to  $T_{\text{OOCF}}$ . By simple calculation, one can find that this measure  $f_*^{-1}\mu$  is nothing but the one in Proposition 2.2.

Using the above map  $T_{\text{OOCF}}$ , we can induce an OOCF expansion of  $x \in (0, 1)$ . For the convenience, let  $T := T_{\text{OOCF}}$ . Denote

$$(2.3) \quad \mathcal{U} := \bigcup_{k \geq 1} \left\{ \frac{2k-1}{2k+1} \right\}, \quad \mathcal{V} := \bigcup_{k \geq 1} \left\{ \frac{k}{k+1} \right\}.$$

By (2.2), for  $k \geq 1$ , for  $x \in (\frac{k-1}{k}, \frac{2k-1}{2k+1})$ ,

$$(2.4) \quad 1-x = \frac{Tx+1}{(k+1)Tx+k} = \frac{1}{(k+1) - \frac{1}{Tx+1}} = \frac{1}{(k+1) + \frac{-1}{2-(1-Tx)}}$$

and, for  $x \in (\frac{2k-1}{2k+1}, \frac{k}{k+1})$ ,

$$(2.5) \quad 1 - x = \frac{Tx + 1}{(k+1) + kTx} = \frac{1}{k + \frac{1}{Tx+1}} = \frac{1}{k + \frac{1}{2 - (1 - Tx)}}.$$

Thus, for all  $x \in [0, 1] \setminus \bigcup_{n \geq 0} T^{-n}(\mathcal{U} \cup \mathcal{V} \cup \{0\})$ ,

$$1 - x = \frac{1}{a_1 + \frac{\varepsilon_1}{2 - \frac{1}{\ddots \frac{\varepsilon_n}{a_n + \frac{\varepsilon_n}{2 - (1 - T^n x)}}}}},$$

where

$$(2.6) \quad (a_n, \varepsilon_n) = \begin{cases} (k+1, -1), & \text{if } T_{\text{OOCF}}^{n-1}(x) \in (\frac{k-1}{k}, \frac{2k-1}{2k+1}), \\ (k, 1), & \text{if } T_{\text{OOCF}}^{n-1}(x) \in (\frac{2k-1}{2k+1}, \frac{k}{k+1}). \end{cases}$$

Hence, we have the OOCF expansion of any  $x \in [0, 1] \setminus \bigcup_{n \geq 0} T^{-n}(\mathcal{U} \cup \mathcal{V} \cup \{0\})$ :

$$x = 1 - \frac{1}{a_1 + \frac{\varepsilon_1}{2 - \frac{1}{a_2 + \frac{\varepsilon_2}{2 - \frac{1}{a_3 + \frac{\varepsilon_3}{2 - \ddots}}}}}}},$$

where  $a_n \in \mathbb{N}$ ; and  $\varepsilon_n \in \{1, -1\}$  for  $a_n \geq 2$  and  $\varepsilon_n = 1$  for  $a_n = 1$ . For simplicity, we denote the OOCF expansion of  $x$  by  $x = \llbracket (a_1, \varepsilon_1), (a_2, \varepsilon_2), \dots, (a_n, \varepsilon_n), \dots \rrbracket$ . We call  $(a_n, \varepsilon_n)$  the  $n$ -th digit of  $x$  in its OOCF expansion.

While for  $x \in \bigcup_{n \geq 0} T^{-n}(\mathcal{U} \cup \mathcal{V} \cup \{0\})$ , the situation is more complicated. First, note that for any  $x \in \mathcal{U}$ ,  $T(x) = 1$  and if  $x = \frac{2k-1}{2k+1}$ , we have

$$1 - x = \frac{1}{(k+1) + \frac{-1}{2 - (1 - Tx)}} = \frac{1}{(k+1) + \frac{-1}{2}}, \quad \text{or} \quad 1 - x = \frac{1}{k + \frac{1}{2 - (1 - Tx)}} = \frac{1}{k + \frac{1}{2}}.$$

Thus  $x = \frac{2k-1}{2k+1}$  has two finite OOCF expansions:  $x = \llbracket (k+1, -1) \rrbracket$  and  $x = \llbracket (k, 1) \rrbracket$ . Further, for  $x \in [0, 1]$  such that  $T^N(x) \in \mathcal{U}$ , for some  $N \geq 1$ , we can apply the iteration (2.4) or (2.5)  $N$  times, and then write  $1 - T^N(x)$  in two different ways. Therefore, any  $x \in \bigcup_{n \geq 0} T^{-n}\mathcal{U}$  has two finite OOCF expansions which differ at the last digit.

If  $x = 0$ , then  $T(0) = 0$  and we have

$$1 - x = \frac{1}{2 + \frac{-1}{2 - (1 - Tx)}}$$

Thus 0 has a unique infinite OOCF expansion:  $0 = \llbracket (2, -1), (2, -1), \dots \rrbracket = \llbracket (2, -1)^\infty \rrbracket$ .

For  $x \in \mathcal{V}$ ,  $T(x) = 0$  and if  $x = \frac{k}{k+1}$ , we have

$$1 - x = \frac{1}{(k+2) + \frac{-1}{2 - (1 - Tx)}} = \frac{1}{(k+2) + \frac{-1}{2 - (1 - 0)}}$$

or

$$1 - x = \frac{1}{(k+1) + \frac{1}{2 - (1 - Tx)}} = \frac{1}{(k+1) + \frac{1}{2 - (1 - 0)}}$$

Hence  $x = \frac{k}{k+1}$  has two infinite OOCF expansions:  $x = \llbracket (k+2, -1), (2, -1)^\infty \rrbracket$  and  $x = \llbracket (k+1, 1), (2, -1)^\infty \rrbracket$ . Similarly, any  $x \in \bigcup_{n \geq 0} T^{-n}\mathcal{V}$  has two infinite OOCF expansions.

### 3. CONVERGENTS OF THE ODD-ODD CONTINUED FRACTION ALGORITHM

In this section, we consider three forms of truncated continued fractions which give us three types of convergents. We investigate basic properties of such convergents and give geometrical interpretations of the hyperbolic plane.

For  $n \geq 1$ , the  $n$ -th *principal convergent* of OOCF is defined by

$$\frac{p_n}{q_n} = \llbracket (a_1, \varepsilon_1), (a_2, \varepsilon_2), \dots, (a_n, \varepsilon_n) \rrbracket = 1 - \frac{1}{a_1 + \frac{\varepsilon_1}{2 - \frac{1}{\dots \frac{\varepsilon_{n-1}}{2 - \frac{1}{a_n + \frac{\varepsilon_n}{2}}}}}}$$

We denote

$$\frac{p'_n}{q'_n} := 1 - \frac{1}{a_1 + \frac{\varepsilon_1}{2 - \frac{1}{\dots \frac{\varepsilon_{n-1}}{2 - \frac{1}{a_n}}}}}$$

and

$$\frac{p''_n}{q''_n} := 1 - \frac{1}{a_1 + \frac{\varepsilon_1}{2 - \frac{1}{\dots \frac{\varepsilon_{n-1}}{2 - \frac{1}{a_n + \varepsilon_n}}}}}$$

and call them the  $n$ -th *sub-convergent* and  $n$ -th *pseudo-convergent*, respectively. To study the convergents of a continued fraction, we have the following general lemma by induction (see [11, p. 3] for details).

**Lemma 3.1.** *Let us consider a general continued fraction of the form*

$$x = a_0 + \frac{b_0}{a_1 + \frac{b_1}{a_2 + \frac{b_2}{a_3 + \ddots}}},$$

where  $a_n, b_n$  are integers. Let  $\tau_n/s_n$  be the finite continued fraction of the form

$$a_0 + \frac{b_0}{a_1 + \frac{b_1}{a_2 + \frac{b_2}{\ddots + \frac{b_{n-1}}{a_n}}}}.$$

Then the following matrix relation holds:

$$\begin{pmatrix} \tau_n & b_n \tau_{n-1} \\ s_n & b_n s_{n-1} \end{pmatrix} = \begin{pmatrix} a_0 & b_0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_n & b_n \\ 1 & 0 \end{pmatrix}.$$

Consequently, we have the following recursive formulas:

$$\begin{cases} \tau_n = a_n \tau_{n-1} + b_{n-1} \tau_{n-2}, \\ s_n = a_n s_{n-1} + b_{n-1} s_{n-2}, \end{cases}$$

where  $\tau_{-1} = 1, s_{-1} = 0, \tau_0 = a_0$  and  $s_0 = 1$ .

Applying Lemma 3.1, we have recursive relations of three types of convergents.

**Lemma 3.2.** *Let  $p'_0 = 1, q'_0 = 0, p_0 = 1$  and  $q_0 = 1$ . We have the following recursive formulas:*

$$\begin{cases} p'_n = a_n p_{n-1} - p'_{n-1} \\ q'_n = a_n q_{n-1} - q'_{n-1} \end{cases}, \quad \begin{cases} p''_n = p'_n + \varepsilon_n p_{n-1} \\ q''_n = q'_n + \varepsilon_n q_{n-1} \end{cases} \quad \text{and} \quad \begin{cases} p_n = 2p'_n + \varepsilon_n p_{n-1} \\ q_n = 2q'_n + \varepsilon_n q_{n-1} \end{cases} \quad \text{for } n \geq 1.$$

By the above recursive formulas, we further have

$$(3.1) \quad \begin{cases} p_n = p'_n + p''_n, \\ q_n = q'_n + q''_n, \end{cases} \quad \text{and} \quad \begin{cases} p_{n-1} = \varepsilon_n (p''_n - p'_n), \\ q_{n-1} = \varepsilon_n (q''_n - q'_n). \end{cases}$$

*Proof.* The lemma is obvious if we plug  $a_0 = 1, b_0 = -1, a_{2n} = 2, a_{2n-1} = a_n, b_{2n} = -1, b_{2n-1} = \varepsilon_n$  in the general continued fraction form in Lemma 3.1.  $\square$

The recursive formulas for the principle convergents are given by the following lemma.

**Lemma 3.3.** *We have*

$$\begin{cases} p_n = (2a_n + \varepsilon_n - 1)p_{n-1} + \varepsilon_{n-1}p_{n-2}, \\ q_n = (2a_n + \varepsilon_n - 1)q_{n-1} + \varepsilon_{n-1}q_{n-2}, \end{cases}$$

where  $p_{-1}/q_{-1} = -1/1, p_0/q_0 = 1/1$  and  $\varepsilon_0 = 1$ .

*Proof.* By Lemma 3.2,

$$p_n = p'_n + p''_n = 2(a_n p_{n-1} - p'_{n-1}) + \varepsilon_n p_{n-1} = (2a_n + \varepsilon_n - 1)p_{n-1} + (p_{n-1} - 2p'_{n-1}).$$

Since  $p_{n-1} - 2p'_{n-1} = \varepsilon_{n-1}p_{n-2}$ , we arrive at the conclusion.  $\square$




FIGURE 3. Farey graph

Now, let us consider the hyperbolic plane  $\mathbb{H}$  as the upper half-plane model. The boundary of  $\mathbb{H}$  is  $\mathbb{R}_\infty = \mathbb{R} \cup \{\infty\}$ . Let  $\ell$  be the vertical line whose endpoints are 0 and  $\infty$ . We denote the union of the images of  $\ell$  under all  $\gamma \in \text{SL}_2(\mathbb{Z})$  by  $\mathcal{G}$  and call it *the Farey graph* (see Figure 3). The Farey graph  $\mathcal{G}$  is a graph on  $\mathbb{H} \cup \mathbb{R}_\infty$  and the vertices in  $\mathbb{R}_\infty$  are all rationals. Recall that the group  $\text{SL}_2(\mathbb{R})$  acts on  $\mathbb{H}$  as Möbius transformations

$$\gamma(z) = \frac{az + b}{cz + d} \quad \text{for } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Since  $\gamma(\infty) = a/c$  and  $\gamma(0) = b/d$ , two rationals  $a/b$  and  $c/d$  are adjacent to each other in  $\mathcal{G}$  if and only if  $|ad - bc| = 1$ . The RCF principal convergents  $p_n^R/q_n^R$  of  $x$  satisfy  $|p_{n+1}^R q_n^R - p_n^R q_{n+1}^R| = 1$ . Thus, we can find a corresponding path to  $x$  on  $\mathcal{G}$  starting from  $[x]$ , passing through all  $p_n^R/q_n^R$  consecutively and converging to  $x$ .

*Ford circles* are horocycles of  $\mathbb{H}$  whose base points are rational or  $\infty$ . Denote by  $C_{a/b}$  the horocycle based at  $a/b$  whose Euclidean radius is  $(2b^2)^{-1}$  and  $C_\infty$  as the line  $y = 1$  (see Figure 4). The collection of Ford circles is a dual of Farey graph in the sense that two Ford circles are tangent to each other if and only if their base points are adjacent to each other in the Farey graph.

Short and Walker [17] examined a similar relation between EICF and a subtree of the Farey graph and the Ford circles. The *Farey tree*  $\mathcal{F}$  is defined as the union of the images of  $\ell$  under  $\gamma \in \Theta$ . The shaded lines in Figure 3 represent the Farey tree. Every vertex of  $\mathcal{F}$  on  $\mathbb{R}_\infty$  is in the orbit of  $\infty$  under  $\Theta$  and its numerator and denominator have different parity. Other rationals have odd numerators and odd denominators.

We will see that each OOCF corresponds to a path on  $\mathcal{G} - \mathcal{F}$ . From now on, let

$$x = \llbracket (a_1, \varepsilon_1), (a_2, \varepsilon_2), \dots, (a_n, \varepsilon_n), \dots \rrbracket$$

be the OOCF expansion of  $x \in (0, 1)$ . The following lemma is on relations of the three distinct convergents of OOCF.

**Lemma 3.4.** *Each  $p_{n-1}/q_{n-1}$  (and also each  $p_n/q_n$ ) is adjacent to  $p'_n/q'_n$  and  $p''_n/q''_n$  in Farey graph. Moreover,  $p'_n/q'_n$  and  $p''_n/q''_n$  are adjacent to each other.*

*Proof.* Recall that  $p'_0/q'_0 = 0/1$ ,  $p''_0/q''_0 = 1/0$  and  $p_0/q_0 = 1/1$ . We have

$$\frac{p'_1}{q'_1} = \frac{a_1 - 1}{a_1}, \quad \frac{p''_1}{q''_1} = \frac{a_1 + \varepsilon_1 - 1}{a_1 + \varepsilon_1} \quad \text{and} \quad \frac{p_1}{q_1} = \frac{2a_1 + \varepsilon_1 - 2}{2a_1 + \varepsilon_1}.$$






FIGURE 4. Ford circles: white circles are based at  $\infty$ -rationals and gray circles are based at 1-rationals

Clearly,  $p_0/q_0$  is adjacent to  $p'_1/q'_1$  and  $p''_1/q''_1$ . So is  $p_1/q_1$ . Let

$$(3.2) \quad A_{(a_n, \varepsilon_n)} = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & a_n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & \varepsilon_n \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} a_n - 1 & a_n + \varepsilon_n - 1 \\ a_n & a_n + \varepsilon_n \end{pmatrix}.$$

Then

$$p_0/q_0 = A_{(a_1, \varepsilon_1)} \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \quad p'_1/q'_1 = A_{(a_1, \varepsilon_1)} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad p''_1/q''_1 = A_{(a_1, \varepsilon_1)} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \text{ and } p_1/q_1 = A_{(a_1, \varepsilon_1)} \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

By Lemma 3.2,

$$\begin{cases} p'_n = (a_n - 1)p'_{n-1} + a_n p''_{n-1} & \text{and} & p''_n = (a_n - 1 + \varepsilon_n)p'_{n-1} + (a_n + \varepsilon_n)p''_{n-1}, \\ q'_n = (a_n - 1)q'_{n-1} + a_n q''_{n-1} & \text{and} & q''_n = (a_n - 1 + \varepsilon_n)q'_{n-1} + (a_n + \varepsilon_n)q''_{n-1}. \end{cases}$$

Thus we have

$$(3.3) \quad \begin{pmatrix} p_{n-1} & p'_n & p''_n & p_n \\ q_{n-1} & q'_n & q''_n & q_n \end{pmatrix} = A_{(a_1, \varepsilon_1)} \cdots A_{(a_n, \varepsilon_n)} \begin{pmatrix} -1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix}.$$

Since  $|\det(A_{(a_n, \varepsilon_n)})| = |-\varepsilon_n| = 1$ , the matrix  $A_{(a_n, \varepsilon_n)}$  acting as the linear fractional map or the anti-linear fractional map preserves adjacency in the Farey graph.  $\square$

**Example 3.5.** Let  $x$  be in the subinterval  $[3/7, 1/2]$  which is the blue segment in Figure 5 and Figure 6. The principal convergents and sub-convergents of  $x$  are

$$\frac{p'_0}{q'_0} = \frac{1}{0}, \quad \frac{p_0}{q_0} = \frac{1}{1}, \quad \frac{p'_1}{q'_1} = \frac{0}{1}, \quad \frac{p_1}{q_1} = \frac{1}{3}, \quad \frac{p'_2}{q'_2} = \frac{2}{5}, \quad \frac{p_2}{q_2} = \frac{3}{7}, \quad \dots$$

The corresponding path of the OOCF expansion of  $x$  is the branch starting from  $\infty$ , going down to 1 and then going along with the arcs connecting  $p'_n/q'_n$  and  $p_n/q_n$ , then the arcs connecting  $p_n/q_n$  and  $p'_{n+1}/q'_{n+1}$  repeatedly. In Figure 5, the red arrows follow the convergents  $p_{n-1}/q_{n-1} \rightarrow p'_n/q'_n \rightarrow p_n/q_n \rightarrow \dots$ . From the duality between Farey graph and Ford circles, there are Ford circles tangent to each other when their




FIGURE 5. The ray and semicircles represent the corresponding path of  $x$  on the Farey graph.




FIGURE 6. The Ford circles numbered from 1 to 5 are Ford circles based at principal convergents and sub-convergents of  $x$  consecutively.

base points are  $p'_n/q'_n$ ,  $p_n/q_n$  or  $p_n/q_n$ ,  $p'_{n+1}/q'_{n+1}$ . The Ford circles numbered from 1 to 5 in Figure 6 are the first five Ford circles corresponding to the OOCF expansion of  $x$ .

For each OOCF digit  $(a, \varepsilon)$ , we partition  $[0, 1]$  into the subintervals  $B(a, \varepsilon)$  defined by

$$(3.4) \quad B(k+1, -1) = \left[ \frac{k-1}{k}, \frac{2k-1}{2k+1} \right], \quad B(k, 1) = \left[ \frac{2k-1}{2k+1}, \frac{k}{k+1} \right].$$

Note that the set of endpoints of  $B(a, \varepsilon)$  is  $\mathcal{U} \cup \mathcal{V}$  as in (2.3). The first OOCF digit of  $x \in B(a, \varepsilon)$  is  $(a, \varepsilon)$  and the restriction of  $T_{\text{OOCF}}$  to  $B(a, \varepsilon)$  is monotone (see Figure 2). We denote the inverse of  $T_{\text{OOCF}}|_{B(a, \varepsilon)}$  by  $f_{(a, \varepsilon)}$ , then by calculation, we have

$$(3.5) \quad f_{(a, \varepsilon)}(t) = 1 - \frac{1}{a + \frac{\varepsilon}{1+t}}.$$

**Lemma 3.6.** For all  $x \in (0, 1)$ ,  $x$  is between  $p_n/q_n$  and  $p'_n/q'_n$ .

*Proof.* By (2.6), we deduce that  $1 - T_{\text{OOCF}}^{n-1}(x)$  is between  $\frac{1}{a_n + (\varepsilon_n/2)}$  and  $\frac{1}{a_n + \varepsilon_n}$ . Let  $g = f_{(a_1, \varepsilon_1)} \circ \dots \circ f_{(a_n, \varepsilon_n)}$ . Since  $g$  is monotone,  $g$  does not change the relative positions of points. Thus  $g(1 - T_{\text{OOCF}}^{n-1}(x)) = x$  is between  $g(\frac{1}{a_n + (\varepsilon_n/2)}) = p_n/q_n$  and  $g(\frac{1}{a_n + \varepsilon_n}) = p'_n/q'_n$ .  $x$  is between  $p_n/q_n$  and  $p''_n/q''_n$ .  $\square$

By (3.2), it is easy to check that either  $A_{(a_n, \varepsilon_n)}$  or  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} A_{(a_n, \varepsilon_n)}$  belongs to the  $\Theta$ -group according to  $\det(A_{(a_n, \varepsilon_n)})$ . With (3.3) we obtain the following theorem.

**Theorem 3.7.** *All principal convergents  $p_n/q_n$  are 1-rationals. Moreover, all sub-convergents  $p'_n/q'_n$  and all pseudo-convergents  $p''_n/q''_n$  are  $\infty$ -rationals.*

Now, we will discuss the relative positions of the different convergents. We define the  $n$ -th complete quotient by  $\zeta_n = T_{\text{OOCF}}^{n-1}(x)$ , i.e.,

$$\zeta_n = 1 - \frac{1}{a_n + \frac{\varepsilon_n}{2 - \frac{1}{a_{n+1} + \frac{\varepsilon_{n+1}}{\ddots}}}}$$

We can show inductively,

$$(3.6) \quad x = \frac{p''_{n-1} + p'_{n-1}\zeta_n}{q''_{n-1} + q'_{n-1}\zeta_n}.$$

By (3.1), we have

$$\begin{cases} p''_{n-1} = \frac{1}{2}(p_{n-1} + \varepsilon_{n-1}p_{n-2}), \\ p'_{n-1} = \frac{1}{2}(p_{n-1} - \varepsilon_{n-1}p_{n-2}). \end{cases} \quad \text{and} \quad \begin{cases} q''_{n-1} = \frac{1}{2}(q_{n-1} + \varepsilon_{n-1}q_{n-2}), \\ q'_{n-1} = \frac{1}{2}(q_{n-1} - \varepsilon_{n-1}q_{n-2}). \end{cases}$$

Thus,

$$x = \frac{p_{n-1}(1 + \zeta_n) + \varepsilon_{n-1}p_{n-2}(1 - \zeta_n)}{q_{n-1}(1 + \zeta_n) + \varepsilon_{n-1}q_{n-2}(1 - \zeta_n)} \quad \text{and} \quad \zeta_n = -\frac{(q_{n-1}x - p_{n-1}) + \varepsilon_{n-1}(q_{n-2}x - p_{n-2})}{(q_{n-1}x - p_{n-1}) - \varepsilon_{n-1}(q_{n-2}x - p_{n-2})}.$$

With the above preparations, we can now show the following lemma.

**Lemma 3.8.** *The following statements hold.*

- (1) *The  $n$ -th principal convergent  $p_n/q_n$  is between  $p'_n/q'_n$  and  $p''_n/q''_n$ .*
- (2) *The  $(n-1)$ -th principal convergent  $p_{n-1}/q_{n-1}$  is not between  $p'_n/q'_n$  and  $p''_n/q''_n$ .*
- (3) *The three distinct convergents  $p_n/q_n$ ,  $p'_n/q'_n$  and  $p''_n/q''_n$  are in the half closed interval  $I_{n-1}$  of  $p_{n-1}/q_{n-1}$  and  $p''_{n-1}/q''_{n-1}$  which contains  $p'_{n-1}/q'_{n-1}$  but does not contain  $p_{n-1}/q_{n-1}$ .*

*Proof.* (1) By Lemma 3.2,  $p_n = p'_n + p''_n$  and  $q_n = q'_n + q''_n$ . Hence,  $p_n/q_n$  is between  $p'_n/q'_n$  and  $p''_n/q''_n$ .

(2) On the Farey graph, there are only two triangles which share the arc connecting  $p'_n/q'_n$  and  $p''_n/q''_n$ . By Lemma 3.4, the other end points of the two triangles are  $p_{n-1}/q_{n-1}$  and  $p_n/q_n$ . By (1),  $p_{n-1}/q_{n-1}$  can not be between  $p'_n/q'_n$  and  $p''_n/q''_n$  since two ideal triangles of the Farey graph do not overlap each other.

(3) By Lemma 3.2, we have

$$\begin{cases} p'_n = (a_n - 1)p_{n-1} + p''_{n-1}, \\ q'_n = (a_n - 1)q_{n-1} + q''_{n-1}, \end{cases} \quad \text{and} \quad \begin{cases} p''_n = (a_n - 1 + \varepsilon_n)p_{n-1} + p''_{n-1}, \\ q''_n = (a_n - 1 + \varepsilon_n)q_{n-1} + q''_{n-1}. \end{cases}$$

Since  $a_n - 1 \geq 0$  and  $a_n + \varepsilon_n - 1 \geq 0$ , both  $p'_n/q'_n$  and  $p''_n/q''_n$  are in  $I_{n-1}$ . By (1),  $p_n/q_n$  is also in  $I_{n-1}$ .  $\square$

**Theorem 3.9.** *The infinite OOCF expansions converge.*

*Proof.* Let  $x$  be a number whose OOCF expansion is infinite. By Lemma 3.8 (3), the intervals  $[p_n/q_n, p''_n/q''_n]$  are shrinking. Thus by Lemma 3.6,

$$\left| x - \frac{p_n}{q_n} \right| < \left| \frac{p_n}{q_n} - \frac{p''_n}{q''_n} \right| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad \square$$

**Lemma 3.10.** *The denominators of principal convergents are increasing.*

*Proof.* Since  $2a_{n+1} + \varepsilon_{n+1} \geq 3$ , we have

$$q_{n+1} = (2a_{n+1} + \varepsilon_{n+1} - 1)q_n + \varepsilon_n q_{n-1} > q_n + (q_n - q_{n-1}).$$

Since  $q_0 - q_{-1} = 1 > 0$ , inductively,  $q_n$  is increasing.  $\square$

As discussed in Section 2, any non-zero rational has exactly two OOCF expansions. For example, the 1-rational  $1/3$  has such two finite OOCF expansions:

$$\frac{1}{3} = 1 - \frac{1}{1 + \frac{1}{2}} = 1 - \frac{1}{2 + \frac{-1}{2}}.$$

For the  $\infty$ -rational  $2/3$ , there are two infinite OOCF expansions:

$$\frac{2}{3} = 1 - \frac{1}{4 + \frac{-1}{2 - \frac{1}{2 + \frac{-1}{2 - \ddots}}}} = 1 - \frac{1}{2 + \frac{1}{2 - \frac{1}{2 + \frac{-1}{2 - \ddots}}}}.$$

In general, we have the following theorem.

**Theorem 3.11.** *The following properties hold.*

- (1) Any finite OOCF is a 1-rational.
- (2) Each 1-rational has exactly two finite OOCF expansions which differ only in the last digit.
- (3) Any non-zero  $\infty$ -rational has exactly two infinite OOCF expansions ending with  $(2, -1)^\infty$ .
- (4) An eventually periodic OOCF expansion converges to an  $\infty$ -rational or a quadratic irrational.
- (5) A quadratic irrational has an eventually periodic OOCF expansion.
- (6) Every irrational has a unique infinite OOCF expansion.

*Proof.* (1): It follows directly from Theorem 3.7.

(2) and (3): For an irreducible rational  $m/n$ , the denominator of  $T_{\text{OOCF}}(m/n)$  is strictly less than  $n$ . Note that the continued fraction map  $T_{\text{OOCF}}$  sends a 1-rational to a 1-rational and an  $\infty$ -rational to an  $\infty$ -rational. Thus if  $m/n$  is a 1-rational, then  $T_{\text{OOCF}}^N(m/n) = 1$  for some  $N$ . Note that

$$T_{\text{OOCF}}^{N-1}(m/n) \in T_{\text{OOCF}}^{-1}(1) = \left\{ \frac{2k-1}{2k+1} : k \geq 1 \right\}.$$

Thus, by the discussion in Section 2,  $m/n$  has exactly two finite OOCF expansions which differ only at last digit.

If  $m/n$  is a non-zero  $\infty$ -rational, then  $T_{\text{OOCF}}^N(m/n) = 0$  for some  $N \geq 1$ . Since the OOCF expansion of 0 is  $[(2, -1)^\infty]$ , the OOCF expansion of  $x$  ends with  $(2, -1)^\infty$ . Similar to the case of 1-rational, note that

$$T_{\text{OOCF}}^{N-1}(m/n) \in \left\{ \frac{k}{k+1} : k \geq 1 \right\}.$$

Then by the discussion in Section 2,  $m/n$  has exactly two infinite OOCF expansions ending with  $(2, -1)^\infty$ .

(4): If  $x$  has an eventually periodic OOCF, then there exist  $i$  and  $j$  such that  $\zeta_{i+1} = \zeta_{j+1}$ . By (3.6),

$$x = \frac{p_i'' + p_i' \zeta_{i+1}}{q_i'' + q_i' \zeta_{i+1}} = \frac{p_j'' + p_j' \zeta_{j+1}}{q_j'' + q_j' \zeta_{j+1}}.$$

Thus we have  $x = p_i''/q_i''$  or

$$\frac{1}{\zeta_{i+1}} = \frac{p_i' - q_i' x}{q_i'' x - p_i''} = \frac{p_j' - q_j' x}{q_j'' x - p_j''} = \frac{1}{\zeta_{j+1}}.$$

The first case corresponds to the case that  $x$  is an  $\infty$ -rational and  $\zeta_{i+1} = 0$ . While the second case implies

$$(q_i' q_j'' - q_j' q_i'') x^2 + (q_j' p_i'' + p_j' q_i'' - q_i' p_j'' - p_i' q_j'') x + (p_i' p_j'' - p_i'' p_j') = 0,$$

which means  $x$  is a quadratic irrational (remark that  $q_i' q_j'' - q_j' q_i'' \neq 0$ ).

(5): Let  $x$  be a quadratic irrational between 0 and 1 such that  $a_1 x^2 + b_1 x + c_1 = 0$ . By (3.6), we have

$$a_1 (p_i'' + p_i' \zeta_{i+1})^2 + b_1 (p_i'' + p_i' \zeta_{i+1})(q_i'' + q_i' \zeta_{i+1}) + c_1 (q_i'' + q_i' \zeta_{i+1})^2 = 0.$$

Let

$$\begin{aligned} a_{i+1} &= a_1 (p_i')^2 + b_1 p_i' q_i' + c_1 (q_i')^2, \\ (3.7) \quad b_{i+1} &= 2a_1 p_i'' p_i' + b_1 (p_i'' q_i' + p_i' q_i'') + 2c_1 q_i'' q_i' \text{ and} \\ c_{i+1} &= a_1 (p_i'')^2 + b_1 p_i'' q_i'' + c_1 (q_i'')^2. \end{aligned}$$

Then  $a_{i+1} \zeta_{i+1}^2 + b_{i+1} \zeta_{i+1} + c_{i+1} = 0$ . Since  $|q_i' p_i'' - q_i'' p_i'| = 1$ , by direct calculation, we have

$$(3.8) \quad b_{i+1}^2 - 4a_{i+1} c_{i+1} = b_1^2 - 4a_1 c_1$$

and also we have  $|(x - p_i'/q_i') + (p_i''/q_i'' - x)| = 1/q_i' q_i''$ . By Lemma 3.6 and the fact that  $p_i/q_i$  is between  $p_i'/q_i'$  and  $p_i''/q_i''$ , two distances  $x - p_i'/q_i'$  and  $x - p_i''/q_i''$  have opposite sign. Thus,  $|q_i' x - p_i'| < 1/q_i''$  and  $|p_i'' - q_i'' x| < 1/q_i'$ . There are  $|\alpha| < 1$  and  $|\beta| < 1$  such that

$$(3.9) \quad p_i' = q_i' x + \alpha/q_i'' \text{ and } p_i'' = q_i'' x + \beta/q_i'.$$

We derive the following expressions by plugging (3.9) in (3.7):

$$a_{i+1} = \alpha \left( \frac{q_i'}{q_i''} (2a_1 x + b_1) + a_1 \frac{\alpha}{(q_i'')^2} \right), \quad c_{i+1} = \beta \left( \frac{q_i''}{q_i'} (2a_1 x + b_1) + a_1 \frac{\beta}{(q_i')^2} \right) \text{ and}$$

$$b_{i+1} = (2a_1 x + b_1)(\alpha + \beta) + 2a_1 \frac{\alpha\beta}{q_i' q_i''}.$$

Since  $|b_{i+1}| \leq 2(|2a_1| + |b_1|) + |2a_1|$ , the coefficient  $b_{i+1}$  is bounded. If  $q_i'' \geq q_i'$ , then  $a_{i+1}$  is bounded since  $|a_{i+1}| < 2|a_1| + |b_1| + |a_1|$ . By (3.8),  $c_{i+1}$  is bounded. Similarly, if  $q_i'' < q_i'$ , then  $c_{i+1}$  is bounded since  $|c_{i+1}| < 2|a_1| + |b_1| + |a_1|$ , then,  $a_{i+1}$  is also bounded. Thus the sequence of the complete quotients  $\{\zeta_i\}$  has only finitely many values. It means that  $\zeta_n = \zeta_m$  for some  $m$  and  $n$ . Therefore, the OOCF expansion of  $x$  is eventually periodic.




FIGURE 7. Two possible relative locations of  $x$ ,  $p/q$ ,  $a/b$  and  $r$  in the proof of Theorem 4.2. The dashed circles are the horocycles based at  $x$  tangent to  $C_{p/q}$  and  $C_{a/b}$ .

(6): Note that

$$\bigcup_{n=0}^{\infty} T_{\text{OOCF}}^{-n}(\mathcal{U} \cup \mathcal{V} \cup \{0\}) = \bigcup_{n=1}^{\infty} T_{\text{OOCF}}^{-n}(\{0, 1\}) = \mathbb{Q}.$$

By the discussion in Section 2, every irrational has a unique infinite OOCF expansion.  $\square$

**Remark 3.12.** From the proof of the point (4) of the Theorem 3.11, we see that if  $x$  is an  $\infty$ -rational, then its OOCF ends with  $(2, -1)^\infty$  and thus there exists  $n_0 \geq 0$  such that  $\zeta_{n+1} = 0$  for all  $n \geq n_0$ . Hence, by (3.6), we have  $x = p_n''/q_n''$  for all  $n \geq n_0$ .

#### 4. BEST 1-RATIONAL APPROXIMATION

Short and Walker defined “the best  $\infty$ -rational approximations” and showed that the EICF gives “the best  $\infty$ -rational approximations”. For  $x \notin \mathbb{Q}$ ,  $p/q$  is a *best  $\infty$ -rational approximation* of  $x$  if  $|qx - p| \leq |bx - a|$  for each  $\infty$ -rational  $a/b$  such that  $b \leq q$  and  $a/b \neq p/q$  (see [17] for more details). Similarly, we prove that the OOCF gives “the best 1-rational approximations”.

**Definition 4.1.** For  $x \notin \mathbb{Q}$ ,  $p/q$  is a *best 1-rational approximation* of  $x$  if for each 1-rational  $a/b$  such that  $b \leq q$  and  $a/b \neq p/q$ , we have

$$|qx - p| \leq |bx - a|.$$

**Theorem 4.2.** *All principal convergents  $p_n/q_n$  are best 1-rational approximations, and vice versa.*

*Proof.* Let  $R_{a/b}(x)$  be the Euclidean radius of the horocycle based at  $x$  which is externally tangent to  $C_{a/b}$ . We can check that  $R_{a/b}(x) := \frac{1}{2}|bx - a|^2$ . Let  $x$  be an irrational number and  $p/q$  be an OOCF convergent of  $x$ . Fix a 1-rational  $a/b$  such that  $1 \leq b \leq q$ . By Lemma 3.4, Lemma 3.6 and Theorem 3.7, there is an  $\infty$ -rational  $r$  which is adjacent to  $p/q$  and  $x$  is between  $p/q$  and  $r$ . The radius  $R_{p/q}(x)$  is at most the radius of  $C_r$ . Since the radius of  $C_{a/b}$  is at least the radius of  $C_{p/q}$ , the 1-rational  $a/b$  is outside of the interval  $[p/q, r]$  (as shown in Figure 7). Thus  $R_{a/b}(x)$  is at least the radius of  $C_r$ . (The equality holds if and only if  $a/b = \infty$ .) We conclude that  $R_{p/q}(x) \leq R_{a/b}(x)$ .

Conversely, let us assume that  $a/b$  is a 1-rational which is not a principal convergent of the OOCF expansion of  $x$ . There are consecutive principal convergents  $p_n/q_n$  and  $p_{n+1}/q_{n+1}$  such that  $q_n \leq b < q_{n+1}$ . Both are adjacent to  $p'_{n+1}/q'_{n+1}$  and  $p''_{n+1}/q''_{n+1}$  in the Farey graph, i.e.,  $C_{p_n/q_n}$  and  $C_{p_{n+1}/q_{n+1}}$  are tangent to  $C_{p'_{n+1}/q'_{n+1}}$  and  $C_{p''_{n+1}/q''_{n+1}}$ . Since the radius of  $C_{a/b}$  is larger than the radius of  $C_{p_{n+1}/q_{n+1}}$ ,  $a/b$  is not between  $p'_{n+1}/q'_{n+1}$  and  $p''_{n+1}/q''_{n+1}$ . Without loss of generality, we assume that  $p_n/q_n < p'_{n+1}/q'_{n+1} < p''_{n+1}/q''_{n+1}$  (see Figure 8).




FIGURE 8. A possible relative position of  $x$ ,  $a/b$  and the convergents. The dashed circles are horocycles based at  $x$  tangent to  $C_{p_n/q_n}$  and  $C_{a/b}$ .

If  $a/b < p_n/q_n$ , then obviously  $R_{a/b}(x) > R_{p_n/q_n}(x)$ . Let us consider the case of  $a/b > p''_{n+1}/q''_{n+1}$ . We note that if  $|x - t| > |x' - t|$ , then  $R_t(x) > R_t(x')$ . Thus,  $R_{a/b}(x) > R_{a/b}(p''_{n+1}/q''_{n+1})$  and  $R_{p_n/q_n}(x)$  is proportional to the distance of  $x$  and  $p_n/q_n$ . The radius  $R_{a/b}(p''_{n+1}/q''_{n+1})$  is at least the radius of  $C_{p''_{n+1}/q''_{n+1}}$  (they are the same when  $C_{p''_{n+1}/q''_{n+1}}$  is tangent to  $C_{a/b}$ ). Since  $x$  is between  $p'_{n+1}/q'_{n+1}$  and  $p''_{n+1}/q''_{n+1}$ , the radius  $R_{p_n/q_n}(x)$  is between  $R_{p_n/q_n}(p'_{n+1}/q'_{n+1}) = 1/2(q'_{n+1})^2$  and  $R_{p_n/q_n}(p''_{n+1}/q''_{n+1}) = 1/2(q''_{n+1})^2$ . Thus,  $R_{a/b}(x) > R_{p_n/q_n}(x)$ .

The last case is of  $a/b \in (p_n/q_n, p'_{n+1}/q'_{n+1})$ . Let us consider a horocycle  $C$  based at  $x$  which is tangent to  $C_{a/b}$ . (The thick dashed arc in Figure 8 represents a part of  $C$ .) The circle  $C$  intersects  $C_{p_n/q_n}$  since the tangent point of  $C$  and  $C_{a/b}$  is an interior point of the shape bounded by  $C_{p_n/q_n}$ ,  $C_{p'_{n+1}/q'_{n+1}}$  and the real line. Thus, for all the cases,  $R_{a/b}(x) > R_{p_n/q_n}(x)$ , i.e.,  $a/b$  is not a 1-rational best approximation.  $\square$

Now, we discuss the relation between the OOCF convergents of a number  $x$  and the EICF convergents of  $1 - x$ . We denote by  $p_n^E(x)/q_n^E(x)$  an EICF convergent of  $x$ . Observe that if  $p_n^E(x)/q_n^E(x)$  is of type even/odd, then  $1 - p_n^E(x)/q_n^E(x)$  is 1-rational. If  $p_n^E(x)/q_n^E(x)$  is of type odd/even, then  $1 - p_n^E(x)/q_n^E(x)$  is still of type odd/even.

**Theorem 4.3.** *Let  $x \in (0, 1)$ . All rationals of type odd/odd in  $1 - p_n^E(1 - x)/q_n^E(1 - x)$  are best 1-rational approximations of  $x$ , and hence are OOCF principal convergents of  $x$ .*

*Proof.* For each  $n \geq 0$ , denote  $P_n/Q_n := 1 - p_n^E(1 - x)/q_n^E(1 - x)$ . Then,  $P_n = q_n^E(1 - x) - p_n^E(1 - x)$  and  $Q_n = q_n^E(1 - x)$ . Thus,  $|P_n - Q_n x| = |q_n^E(1 - x) - p_n^E(1 - x) - x \cdot q_n^E(1 - x)| = |(1 - x) \cdot q_n^E(1 - x) - p_n^E(1 - x)| < |(1 - x)b - a|$  for any  $\infty$ -rational  $a/b$  such that  $1 \leq b \leq q_n^E(1 - x)$  and  $a/b \neq p_n^E(1 - x)/q_n^E(1 - x)$ .

For any 1-rational  $c/d$  such that  $1 \leq d \leq Q_n$  and  $c/d \neq P_n/Q_n$ , we have  $\frac{d-c}{d} = 1 - \frac{c}{d}$  is  $\infty$ -rational and  $\frac{d-c}{d} \neq p_n^E(1 - x)/q_n^E(1 - x)$ . Thus  $|P_n - Q_n x| < |(1 - x)d - (d - c)| = |c - dx|$ , which means that  $P_n/Q_n$  is a best 1-rational approximation of  $x$ .  $\square$

**Example 4.4.** Let  $x = 10 - \pi^2 = 0.1303955989 \dots$ . The OOCF convergents of  $x$  are:

$$(4.1) \quad \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \frac{3}{23}, \frac{287}{2201}, \frac{577}{4425}, \frac{867}{6649}, \frac{1157}{8873}, \frac{1447}{11097}, \frac{6945}{53261}, \frac{40223}{308469}, \dots$$

The sequence of  $1 - p_n^E/q_n^E$  where  $p_n^E/q_n^E$  is the EICF convergent of  $1 - x$  is:

$$(4.2) \quad \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \frac{1}{7}, \frac{1}{8}, \frac{3}{23}, \frac{145}{1112}, \frac{1447}{11097}, \frac{2749}{21082}, \frac{6945}{53261}, \frac{16639}{127604}, \dots$$

The subsequence of odd/odd in (4.2) is a (strict) subsequence of (4.1). However, the rational  $287/2201$  in (4.1) is not in (4.2) since

$$|1112x - 145| = 0.0000940113 \dots < |2201x - 287| = 0.00071320 \dots < |23x - 3| = 0.00090122 \dots$$

**Theorem 4.5.** *Let  $f(x) = \frac{1-x}{1+x}$ . Then, we have  $p_n^E(f(x))/q_n^E(f(x)) = f(p_n(x)/q_n(x))$ .*

*Proof.* From now on, we write an EICF expansion as a sequence in a double angle bracket:

$$\frac{1}{b_1 + \frac{\eta_1}{b_2 + \frac{\eta_2}{b_3 + \frac{\eta_3}{\ddots}}}} = \langle\langle (b_1, \eta_1), (b_2, \eta_2), \dots, (b_n, \eta_n), \dots \rangle\rangle = \langle\langle (b_n, \eta_n)_{n \in \mathbb{N}} \rangle\rangle.$$

Denote by  $\mathcal{A}_{\text{OOCF}}$  and  $\mathcal{A}_{\text{EICF}}$ , the sets of digits of OOCF and EICF respectively. Explicitly, we have

$$\mathcal{A}_{\text{OOCF}} = \{(a, \varepsilon) : a \in \mathbb{N}, \varepsilon = \pm 1\} \setminus \{(1, -1)\}, \quad \mathcal{A}_{\text{EICF}} = \{(b, \eta) : b' \in 2\mathbb{N}, \eta = \pm 1\}.$$

Recall the partition  $\{B(a, \varepsilon)\}_{(a, \varepsilon) \in \mathcal{A}_{\text{OOCF}}}$  in (3.4). For the measure preserving map  $f(x) = \frac{1-x}{1+x}$  on the unit interval in Theorem 2.1, by calculation, we have

$$f(B(k+1, -1)) = \left[ \frac{1}{2k}, \frac{1}{2k-1} \right], \quad f(B(k, 1)) = \left[ \frac{1}{2k+1}, \frac{1}{2k} \right].$$

Thus,  $f(B(k+1, -1))$ ,  $f(B(k, 1))$  are subintervals which correspond to  $(2k, 1), (2k, -1) \in \mathcal{A}_{\text{EICF}}$ . In other words, there is a natural correspondence  $\phi$  between  $\mathcal{A}_{\text{OOCF}}$  and  $\mathcal{A}_{\text{EICF}}$  as follows:

$$\phi : (k+1, -1) \mapsto (2k, 1), \quad (k, 1) \mapsto (2k, -1).$$

Inductively, we define a subinterval corresponding to an OOCF expansion  $\langle\langle (a_1, \varepsilon_1), \dots, (a_n, \varepsilon_n) \rangle\rangle$  by

$$B(\langle\langle (a_1, \varepsilon_1), \dots, (a_n, \varepsilon_n) \rangle\rangle) = B(a_1, \varepsilon_1) \cap T_{\text{OOCF}}^{-1} B(a_2, \varepsilon_2) \cap T_{\text{OOCF}}^{-2} B(a_3, \varepsilon_3) \dots \cap T_{\text{OOCF}}^{-n+1} B(a_n, \varepsilon_n).$$

Since  $f$  is 1-1 and onto,

$$(4.3) \quad \begin{aligned} & f(B(\langle\langle (a_1, \varepsilon_1), \dots, (a_n, \varepsilon_n) \rangle\rangle)) \\ &= f(B(a_1, \varepsilon_1)) \cap f(T_{\text{OOCF}}^{-1} B(a_2, \varepsilon_2)) \cap f(T_{\text{OOCF}}^{-2} B(a_3, \varepsilon_3)) \dots \cap f(T_{\text{OOCF}}^{-n+1} B(a_n, \varepsilon_n)). \end{aligned}$$

From Theorem 2.1, the interval in (4.3) is the subinterval corresponding to  $\langle\langle \phi(a_i, \varepsilon_i)_{i=1}^n \rangle\rangle$ . Thus, for  $x = \langle\langle (a_n, \varepsilon_n)_{n \in \mathbb{N}} \rangle\rangle$ , the EICF expansion of  $f(x)$  is  $\langle\langle \phi(a_n, \varepsilon_n)_{n \in \mathbb{N}} \rangle\rangle$  and  $p_n^E(f(x))/q_n^E(f(x)) = f(p_n(x)/q_n(x))$ .  $\square$

## 5. RELATION WITH THE REGULAR CONTINUED FRACTION

In their monograph [13], Rockett and Szűs introduced “the best approximation of the first kind” and “the best approximation of the second kind”. Our definition of the best approximation is the best approximation of the second kind. The best approximation of the first kind of  $x$  is a rational  $p/q$  such that  $|x - p/q| < |x - a/b|$  for any  $a/b \neq p/q$  and  $q \leq b$ . For  $n \in \mathbb{N}$ , Denote by  $p_n^R/q_n^R$  the principal convergents of RCF. Every best



approximation of the first kind is a form of

$$\frac{p_{n,j}^R}{q_{n,j}^R} = \frac{p_{n-2}^R + jp_{n-1}^R}{q_{n-2}^R + jq_{n-1}^R}$$

for  $1 \leq j \leq a_n$  and for  $n \geq 1$  (see [10, Section 6] and [13, p.36] for the details). The rationals are called *the intermediate convergents*. Kraaikamp and Lopes showed that the convergents of EICF are intermediate convergents of RCF (see [12] for the proof). We show that the OOCF principal convergents are also intermediate convergents of RCF. Now we denote an RCF expansion by

$$d_0 + \frac{1}{d_1 + \frac{1}{d_2 + \frac{1}{\ddots}}} = [d_0; d_1, d_2, \dots].$$

For  $(a, \varepsilon) \in \mathcal{A}_{\text{OOCF}}$ , let  $f_{(a,\varepsilon)}$  be as in (3.5). The following lemma tells us how the peicewise inverses of OOCF acts on RCF expansions.

**Lemma 5.1.** *Let  $x = [0; d_1, d_2, \dots]$ . Then, the RCF expansion of  $f_{(a,\varepsilon)}(x)$  is as follows:*

$$f_{(a,\varepsilon)}(x) = \begin{cases} [2, d_1, d_2, \dots] & \text{if } \varepsilon = 1, a = 1 \\ [1, (a-1), 1, d_1, d_2, \dots] & \text{if } \varepsilon = 1, a \geq 2 \\ [(d_1+2), d_2, \dots] & \text{if } \varepsilon = -1, a = 2 \\ [1, (a-1), (d_1+1), d_2, \dots] & \text{if } \varepsilon = -1, a \geq 3. \end{cases}$$

*Proof.* If  $\varepsilon = 1$ , then

$$f_{(a,\varepsilon)}(x) = 1 - [a, 1, d_1, d_2, \dots] = \begin{cases} [2, d_1, d_2, \dots] & \text{if } a = 1, \\ [1, a-1, 1, d_1, \dots] & \text{if } a \geq 2. \end{cases}$$

If  $\varepsilon = -1$ , then

$$\begin{aligned} f_{(a,\varepsilon)}(x) &= 1 - \frac{1}{a - [1, d_1, d_2, \dots]} = 1 - \frac{1}{(a-1) + 1 - [1, d_1, d_2, \dots]} \\ &= 1 - [a-1, d_1+1, d_2, \dots] = \begin{cases} [d_1+2, d_2, \dots] & \text{if } a = 2, \\ [1, a-1, 1, d_1+1, d_2, \dots] & \text{if } a \geq 3. \end{cases} \end{aligned}$$

□

**Theorem 5.2.** *The OOCF principal convergents of  $x$  are intermediate convergents of  $x$ .*

*Proof.* Let  $x = [d_1, d_2, \dots] = \llbracket (a_1, \varepsilon_1), (a_2, \varepsilon_2), \dots \rrbracket$ . Note that

$$x = f_{(a_1, \varepsilon_1)} \circ f_{(a_2, \varepsilon_2)} \circ \dots \circ f_{(a_k, \varepsilon_k)}(\llbracket (a_{k+1}, \varepsilon_{k+1}), \dots \rrbracket) \quad \text{and} \quad \frac{p_k}{q_k} = f_{(a_1, \varepsilon_1)} \circ f_{(a_2, \varepsilon_2)} \circ \dots \circ f_{(a_k, \varepsilon_k)}(1).$$

By Lemma 5.1,  $x$  and  $p_k/q_k$  have the same prefix in their RCF expansions, except for the last digit of  $p_k/q_k$ . Thus,  $p_k/q_k$  is an intermediate convergent of  $x$ . □

We use the following notations:

$$\begin{cases} x = [0; d_1, \dots, d_n, \alpha] & \text{if } G^n(x) = \alpha, \\ x = \llbracket (a_1, \varepsilon_1), (a_2, \varepsilon_2), \dots, (a_n, \varepsilon_n), \gamma \rrbracket & \text{if } T_{\text{OOCF}}^n(x) = \gamma. \end{cases}$$

Note that

$$(5.1) \quad \frac{1}{1+\alpha} = 1 - \frac{1}{1+\frac{1}{\alpha}}.$$

**Theorem 5.3.** *We can convert RCF expansions into OOCF expansions by the following relations:*

$$x = [0; d_1, d_2, \alpha] = \begin{cases} \llbracket (2, -1)^{\frac{d_1-1}{2}}, (d_2+1, 1), F(\alpha) \rrbracket & \text{if } d_1 \text{ is odd and } \alpha \in [\frac{1}{2}, 1), \\ \llbracket (2, -1)^{\frac{d_1-1}{2}}, (d_2+2, -1), F(\alpha) \rrbracket & \text{if } d_1 \text{ is odd and } \alpha \in [0, \frac{1}{2}), \\ \llbracket (2, -1)^{\frac{d_1}{2}-1}, (1, 1), G(x) \rrbracket & \text{if } d_1 \text{ is even.} \end{cases}$$

*Proof.* By (5.1), we have

$$[0; 1, d_2, \alpha] = \frac{1}{1 + \frac{1}{d_2 + \alpha}} = 1 - \frac{1}{(d_2 + 1) + \alpha}.$$

Since  $\alpha = \frac{1}{[\alpha^{-1}] + G(\alpha)}$ , if  $\alpha \in [\frac{1}{2}, 1)$ , then  $\alpha = \frac{1}{1+G(\alpha)}$ . If  $\alpha \in [0, \frac{1}{2})$ , then

$$\alpha = 1 - \frac{1}{1 + \frac{1}{[\alpha^{-1}] - 1 + G(\alpha)}}.$$

Using the fact that  $F(\alpha) = \frac{1}{[\alpha^{-1}] - 1 + G(\alpha)}$  if  $\alpha \in [0, \frac{1}{2})$  and otherwise,  $F(\alpha) = G(\alpha)$ , we have

$$[0; d_1, d_2, \alpha] = \begin{cases} \llbracket (d_2 + 1, 1), F(\alpha) \rrbracket, & \text{if } \alpha \in [\frac{1}{2}, 1), \\ \llbracket (d_2 + 2, -1), F(\alpha) \rrbracket, & \text{if } \alpha \in [0, \frac{1}{2}). \end{cases}$$

Similarly,

$$x = [0; 2, G(x)] = \frac{1}{2 + G(x)} = 1 - \frac{1}{1 + \frac{1}{1+G(x)}} = \llbracket (1, 1), G(x) \rrbracket.$$

If  $d_1 \geq 3$ , then

$$x = [0; d_1, G(x)] = \frac{1}{d_1 + G(x)} = 1 - \frac{1}{1 + \frac{1}{1+(d_1-2)+G(x)}} = 1 - \frac{1}{2 - \frac{1}{1+(d_1-2)+G(x)}} = \llbracket (2, -1), \gamma \rrbracket.$$

where  $\gamma = [0; d_1 - 2, G(x)] = R(x)$ . Thus, by induction, we complete the proof.  $\square$

Since  $p_n^R/q_n^R$  is a best approximation, if  $p_n^R/q_n^R$  is a 1-rational, then  $p_n^R/q_n^R$  is a 1-rational best approximation, thus,  $p_n^R/q_n^R$  is an OOCF convergent by Theorem 4.2. Now we check when an intermediate convergent is an OOCF convergent. Keita [9] proved the following proposition.

**Proposition 5.4** (Proposition 1.2 in [9]). *We have*

$$q_{n,0}^R = q_{n-2}^R < q_{n-1}^R \leq q_{n,1}^R < \dots < q_{n,d_n}^R = q_n^R,$$

$$|q_{n,d_n}^R x - p_{n,d_n}^R| = |q_n^R x - p_n^R| < |q_{n-1}^R x - p_{n-1}^R| \leq |q_{n,d_n-1}^R x - p_{n,d_n-1}^R| < \dots < |q_{n,0}^R x - p_{n,0}^R| = |q_{n-2}^R x - p_{n-2}^R|.$$

By the above proposition and Theorem 5.2, if  $p_{n-1}^R/q_{n-1}^R$  is a 1-rational, then  $p_{n,j}^R/q_{n,j}^R$  is not an OOCF principal convergent for any  $1 \leq j < d_n$ . If  $p_{n-1}^R/q_{n-1}^R$  is an  $\infty$ -rational and  $p_{n,j}^R/q_{n,j}^R$  is a 1-rational, then  $p_{n,j}^R/q_{n,j}^R$  is an OOCF principal convergent.

## REFERENCES

- [1] R. C. Alperin, *The modular tree of Pythagoras*, Amer. Math. Monthly **112** (2005), no. 9, 807–816, DOI 10.2307/30037602. MR2179860
- [2] F. J. M. Barning, *On Pythagorean and quasi-Pythagorean triangles and a generation process with the help of unimodular matrices*, Math. Centrum Amsterdam Afd. Zuivere Wisk. **1963** (1963), no. ZW-011, 37 (Dutch). MR0190077
- [3] B. Berggren, *Pytagoreiska trianglar*, Tidskrift för elementär matematik, fysik och kemi **17** (1934), 129–139.
- [4] F. P. Boca and C. Merriman, *Coding of geodesics on some modular surfaces and applications to odd and even continued fractions*, Indagationes Mathematicae **29** (2018), no. 5, 1214–1234.
- [5] B. Cha and D. H. Kim, *Number theoretical properties of Romik’s dynamical system*, Bull. Korean Math. Soc. **57** (2020), 251–274, DOI <https://doi.org/10.4134/BKMS.b190163>.
- [6] ———, *Lagrange spectrum of Romik’s dynamical system*, available at <https://arxiv.org/abs/1903.02882>.
- [7] B. Cha, E. Nguyen, and B. Tauber, *Quadratic forms and their Berggren trees*, J. Number Theory **185** (2018), 218–256, DOI <https://doi.org/10.1016/j.jnt.2017.09.003>.
- [8] K. Conrad, *Pythagorean descent*, available at <http://www.math.uconn.edu/~kconrad/blurbs/linmultialg/descentPythag.pdf>.
- [9] A. Keita, *Continued fractions and parametric geometry of numbers*, Journal de Théorie des Nombres de Bordeaux **29** (2017), no. 1, 129–135.
- [10] A. I. Khinchin, *Continued fractions*, P. Noordhoff, 1963.
- [11] C. Kraaikamp, *A new class of continued fraction expansions*, Acta Arithmetica **57** (1991), no. 1, 1–39.
- [12] C. Kraaikamp and A. Lopes, *The theta group and the continued fraction expansion with even partial quotients*, Geometriae Dedicata **59** (1996), no. 3, 293–333.
- [13] A. Rockett and P. Szűsz, *Continued fractions*, World Sci, Pub., Singapore, 1992.
- [14] D. Romik, *The dynamics of Pythagorean triples*, Trans. Amer. Math. Soc. **360** (2008), no. 11, 6045–6064, DOI 10.1090/S0002-9947-08-04467-X. MR2425702 (2009i:37101)
- [15] F. Schweiger, *Continued fractions with odd and even partial quotients*, Arbeitsber. Math. Inst. Univ. Salzburg **4** (1982), 59–70.
- [16] ———, *On the approximation by continued fractions with odd and even partial quotients*, Arbeitsber. Math. Inst. Univ. Salzburg **1** (1984), no. 2, 105–114.
- [17] I. Short and M. Walker, *Even-integer continued fractions and the Farey tree*, Symmetries in Graphs, Maps, and Polytopes Workshop (2014), 287–300.

DEPARTMENT OF MATHEMATICS EDUCATION, DONGGUK UNIVERSITY—SEOUL, 30 PILDONG-RO 1-GIL, JUNG-GU, SEOUL, 04620 KOREA  
*E-mail address:* kim2010@dongguk.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, KWANAK-RO 1, KWANAK-GU, SEOUL, 08826 KOREA  
*E-mail address:* seulbee.lee@snu.ac.kr

UNIVERSITÉ PARIS-EST, LAMA (UMR 8050), UPEMLV, UPEC, CNRS, F-94010, CRETEIL, FRANCE  
*E-mail address:* lingmin.liao@u-pec.fr