

On the residual of a factorized group with widely supersoluble factors

Victor S. Monakhov and Alexander A. Trofimuk

Department of Mathematics and Programming Technologies,

Francisk Skorina Gomel State University,

Gomel 246019, Belarus

e-mail: victor.monakhov@gmail.com

e-mail: alexander.trofimuk@gmail.com[†]

Abstract. Let \mathbb{P} be the set of all primes. A subgroup H of a group G is called \mathbb{P} -subnormal in G , if either $H = G$, or there exists a chain of subgroups $H = H_0 \leq H_1 \leq \dots \leq H_n = G$, $|H_i : H_{i-1}| \in \mathbb{P}$, $\forall i$. A group G is called *widely supersoluble*, *w-supersoluble* for short, if every Sylow subgroup of G is \mathbb{P} -subnormal in G . A group $G = AB$ with \mathbb{P} -subnormal *w-supersoluble* subgroups A and B is studied. The structure of its *w-supersoluble* residual is obtained. In particular, it coincides with the nilpotent residual of the \mathcal{A} -residual of G . Here \mathcal{A} is the formation of all groups with abelian Sylow subgroups. Besides, we obtain new sufficient conditions for the *w-supersolubility* of such group G .

Keywords. widely supersoluble groups, mutually *sn*-permutable subgroups, \mathbb{P} -subnormal subgroup, the \mathfrak{X} -residual.

Mathematics Subject Classification. 20D10, 20D20.

Introduction

Throughout this paper, all groups are finite and G always denotes a finite group. We use the standard notations and terminology of [6]. The formations of all nilpotent, supersoluble groups and groups with abelian Sylow subgroups are denoted by \mathfrak{N} , \mathfrak{U} and \mathcal{A} , respectively. The notation $Y \leq X$ means that Y is a subgroup of a group X and \mathbb{P} be the set of all primes. Let \mathfrak{X} be a formation. Then $G^{\mathfrak{X}}$ denotes the \mathfrak{X} -residual of G .

By Huppert's Theorem [6, VI.9.5], a group G is supersoluble if and only if for every proper subgroup H of G there exists a chain of subgroups

$$H = H_0 \leq H_1 \leq \dots \leq H_n = G, |H_i : H_{i-1}| \in \mathbb{P}, \forall i. \quad (1)$$

So naturally the following definition.

A subgroup H of a group G is called \mathbb{P} -subnormal in G , if either $H = G$, or there is a chain subgroups (1). We use the notation $H \text{ Psn } G$. This definition was proposed in [13] and besides, in this paper w -supersoluble (widely supersoluble) groups, i.e. groups with \mathbb{P} -subnormal Sylow subgroups, were investigated. Denote by $w\mathfrak{U}$ the class of all w -supersoluble groups.

The factorizable groups $G = AB$ with w -supersoluble factors A and B were investigated in [8], [10], [11], [14]. There are many other papers devoted to study factorizable groups, and the reader is referred to the book [1] and the bibliography therein. A criteria for w -supersolvability was obtained by A. F. Vasil'ev, T. I. Vasil'eva and V. N. Tyutyanyanov [14].

Theorem A. [14, Theorem 4.7] *Let $G = AB$ be a group which is the product of two w -supersoluble subgroups A and B . If A and B are \mathbb{P} -subnormal in G and G^A is nilpotent, then G is w -supersoluble.*

We recall that two subgroups A and B of a group G are said to be *mutually sn-permutable* if A permutes with all subnormal subgroups of B and B permutes with all subnormal subgroups of A . If A and B are mutually sn -permutable subgroups of a group $G = AB$, then we say that G is a *mutually sn-permutable product* of A and B , see [4]. In soluble groups, mutually sn -permutable factors are \mathbb{P} -subnormal [14, Lemma 4.5]. The converse is not true, see the example 3.1 below.

A. Ballester-Bolinches, W. M. Fakieh and M. C. Pedraza-Aguilera [3] obtained the following results for the sn -permutable product of the w -supersoluble subgroups.

Theorem B. *Let $G = AB$ be the mutually sn -permutable product of subgroups A and B . Then the following hold:*

- (1) *if A and B are w -supersoluble and N is a minimal normal subgroup of G , then both AN and BN are w -supersoluble, [3, Theorem 3];*
- (2) *if A and B are w -supersoluble and $(|A/A^A|, |B/B^A|) = 1$, then G is w -supersoluble, [3, Theorem 5].*

Present paper extends the Theorems A and B. We prove the following result.

Theorem 1. *Let A and B be w -supersoluble \mathbb{P} -subnormal subgroups of G and $G = AB$. Then the following hold:*

- (1) $G^{w\mathfrak{U}} = (G^A)^{\mathfrak{U}}$;
- (2) *if N is a nilpotent normal subgroup of G , then both AN and BN are w -supersoluble;*

(3) if $(|A/A^A|, |B/B^A|) = 1$, then G is w -supersoluble.

Theorem A follows from assertion (1) of Theorem 1. Theorem B follows from assertions (2) and (3) of Theorem 1 since the group G in Theorem B is soluble.

1 Preliminaries

In this section, we give some definitions and basic results which are essential in the sequel. A group whose chief factors have prime orders is called *supersoluble*. Recall that a p -closed group is a group with a normal Sylow p -subgroup and a p -nilpotent group is a group with a normal Hall p' -subgroup.

Denote by G' , $Z(G)$, $F(G)$ and $\Phi(G)$ the derived subgroup, centre, Fitting and Frattini subgroups of G respectively. We use E_{p^t} to denote an elementary abelian group of order p^t and Z_m to denote a cyclic group of order m . The semidirect product of a normal subgroup A and a subgroup B is written as follows: $A \rtimes B$.

Let \mathfrak{F} be a formation. Recall that the \mathfrak{F} -residual of G , that is the intersection of all those normal subgroups N of G for which $G/N \in \mathfrak{F}$. We define $\mathfrak{X}\mathfrak{Y} = \{G \in \mathfrak{E} \mid G^{\mathfrak{Y}} \in \mathfrak{X}\}$ and call $\mathfrak{X}\mathfrak{Y}$ the *formation product* of \mathfrak{X} and \mathfrak{Y} . Here \mathfrak{E} is the class of all finite groups.

If H is a subgroup of G , then $H_G = \bigcap_{x \in G} H^x$ is called *the core* of H in G . If a group G contains a maximal subgroup M with trivial core, then G is said to be *primitive* and M is its *primitivator*.

A simple check proves the following lemma.

Lemma 1.1. *Let \mathfrak{F} be a saturated formation and G be a group. Assume that $G \notin \mathfrak{F}$, but $G/N \in \mathfrak{F}$ for all non-trivial normal subgroups N of G . Then G is a primitive group.*

Lemma 1.2. ([6, Theorem II.3.2]) *Let G be a soluble primitive group and M is a primitivator of G . Then the following statements hold:*

- (1) $\Phi(G) = 1$;
- (2) $F(G) = C_G(F(G)) = O_p(G)$ and $F(G)$ is an elementary abelian subgroup of order p^n for some prime p and some positive integer n ;
- (3) G contains a unique minimal normal subgroup N and moreover, $N = F(G)$;
- (4) $G = F(G) \rtimes M$ and $O_p(M) = 1$.

Lemma 1.3. ([2, Proposition 2.2.8, Proposition 2.2.11]) *Let \mathfrak{F} and \mathfrak{H} be formations, K be normal in G . Then the following hold:*

- (1) $(G/K)^{\mathfrak{F}} = G^{\mathfrak{F}}K/K$;

- (2) $G^{\mathfrak{H}\mathfrak{H}} = (G^{\mathfrak{H}})^{\mathfrak{H}}$;
- (3) if $\mathfrak{H} \subseteq \mathfrak{F}$, then $G^{\mathfrak{H}} \leq G^{\mathfrak{F}}$;
- (4) if $G = HK$, then $H^{\mathfrak{H}}K = G^{\mathfrak{H}}K$.

Recall that a group G is said to be *siding* if every subgroup of the derived subgroup G' is normal in G , see [12, Definition 2.1]. Metacyclic groups, t-groups (groups in which every subnormal subgroup is normal) are siding. The group $G = (Z_6 \times Z_2) \rtimes Z_2$ ($\text{IdGroup}(G)=[24,8]$) [5] is siding, but not metacyclic and a t-group.

Lemma 1.4. *Let G be siding. Then the following hold:*

- (1) if N is normal in G , then G/N is siding;
- (2) if H is a subgroup of G , then H is siding;
- (3) G is supersoluble

Proof. 1. By [6, Lemma I.8.3], $(G/N)' = G'N/N$. Let A/N be an arbitrary subgroup of $(G/N)'$. Then

$$A \leq G'N, \quad A = A \cap G'N = (A \cap G')N.$$

Since $A \cap G' \leq G'$, we have $A \cap G'$ is normal in G . Hence $(A \cap G')N/N$ is normal in G/N .

2. Since $H \leq G$, it follows that $H' \leq G'$. Let A be an arbitrary subgroup of H' . Then $A \leq G'$ and A is normal in G . Therefore A is normal in H .

3. We proceed by induction on the order of G . Let $N \leq G'$ and $|N| = p$, where p is prime. By the hypothesis, N is normal in G . By induction, G/N is supersoluble and G is supersoluble. \square

Lemma 1.5. ([9, Lemma 3]) *Let H be a subgroup of G , and N be a normal subgroup of G . Then the following hold:*

- (1) if $N \leq H$ and $H/N \mathbb{P}sn G/N$, then $H \mathbb{P}sn G$;
- (2) if $H \mathbb{P}sn G$, then $(H \cap N) \mathbb{P}sn N$, $HN/N \mathbb{P}sn G/N$ and $HN \mathbb{P}sn G$;
- (3) if $H \leq K \leq G$, $H \mathbb{P}sn K$ and $K \mathbb{P}sn G$, then $H \mathbb{P}sn G$;
- (4) if $H \mathbb{P}sn G$, then $H^g \mathbb{P}sn G$ for any $g \in G$.

Lemma 1.6. ([9, Lemma 4]) *Let G be a soluble group, and H be a subgroup of G . Then the following hold:*

- (1) if $H \mathbb{P}sn G$ and $K \leq G$, then $(H \cap K) \mathbb{P}sn K$;
- (2) if $H_i \mathbb{P}sn G$, $i = 1, 2$, then $(H_1 \cap H_2) \mathbb{P}sn G$.

Lemma 1.7. ([9, Lemma 5]) *If H is a subnormal subgroup of a soluble group G , then H is \mathbb{P} -subnormal in G .*

Lemma 1.8. ([13, Theorem 2.7]) *The class $w\mathfrak{U}$ is a hereditary saturated formation.*

Lemma 1.9. (1) *If $G \in w\mathfrak{U}$, then G^A is nilpotent, [13, Theorem 2.13].*

(2) *$G \in w\mathfrak{U}$ if and only if every metanilpotent subgroup of G is supersoluble, [7, Theorem 2.6].*

(3) *$G \in w\mathfrak{U}$ if and only if G has a Sylow tower of supersoluble type and every biprimary subgroup of G is supersoluble, [9, Theorem B].*

2 Factorizable groups with \mathbb{P} -subnormal w -supersoluble subgroups

Lemma 2.1. ([14, Theorem 4.4]) *Let A and B be \mathbb{P} -subnormal subgroups of G , and $G = AB$. If A and B have an ordered Sylow tower of supersoluble type, then G has an ordered Sylow tower of supersoluble type.*

Proof of Theorem 1 (1). If G is w -supersoluble, then $G^{w\mathfrak{U}} = 1$ and G^A is nilpotent by Lemma 1.9 (1). Consequently $G^{w\mathfrak{U}} = 1 = (G^A)^{\mathfrak{N}}$ and the statement is true. Further, we assume that G is non- w -supersoluble. Since $w\mathfrak{U} \subseteq \mathfrak{NA}$, it follows that

$$G^{(\mathfrak{NA})} = (G^A)^{\mathfrak{N}} \leq G^{w\mathfrak{U}}$$

by Lemma 1.3 (2-3). Next we check the converse inclusion. For this we prove that $G/(G^A)^{\mathfrak{N}}$ is w -supersoluble. By Lemma 1.3 (1),

$$(G/(G^A)^{\mathfrak{N}})^A = G^A(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}} = G^A/(G^A)^{\mathfrak{N}}$$

and $(G/(G^A)^{\mathfrak{N}})^A$ is nilpotent. The quotients

$$G/(G^A)^{\mathfrak{N}} = (A(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}})(B(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}}),$$

$$A(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}} \simeq A/A \cap (G^A)^{\mathfrak{N}},$$

$$B(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}} \simeq B/B \cap (G^A)^{\mathfrak{N}},$$

hence the subgroups $A(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}}$ and $B(G^A)^{\mathfrak{N}}/(G^A)^{\mathfrak{N}}$ are w -supersoluble by Lemma 1.8 and by Lemma 1.5 (2), they are \mathbb{P} -subnormal in $G/(G^A)^{\mathfrak{N}}$. By Theorem A, $G/(G^A)^{\mathfrak{N}}$ is w -supersoluble. \square

Lemma 2.2. *Let G be a group, and A be a subgroup of G such that $|G : A| = p^\alpha$, where $p \in \pi(G)$ and $\alpha \in \mathbb{N}$. Suppose that A is w -supersoluble and \mathbb{P} -subnormal in G . If G is p -closed, then G is w -supersoluble.*

Proof. Let P be a Sylow p -subgroup of G . Since P is normal in G and $G = AP$, we have $G/P \simeq A/A \cap P \in \mathfrak{w}\mathfrak{U}$, in particular, G is soluble. Because G is soluble, it follows that P is \mathbb{P} -subnormal in G by Lemma 1.7. Let Q be a Sylow q -subgroup of G , $q \neq p$. Then $Q \leq A^x$ for some $x \in G$. By Lemma 1.5 (4), A^x is \mathbb{P} -subnormal in G . Since $A^x \in \mathfrak{w}\mathfrak{U}$, it follows that Q is \mathbb{P} -subnormal in A^x and Q is \mathbb{P} -subnormal in G by Lemma 1.5 (3). So, G is w-supersoluble. \square

Lemma 2.3. *Let A and B be w-supersoluble \mathbb{P} -subnormal subgroups of G , and $G = AB$. Suppose that $|G : A| = p^\alpha$, where $p \in \pi(G)$. If p is the greatest in $\pi(G)$, then G is w-supersoluble.*

Proof. Since every w-supersoluble group has an ordered Sylow tower of supersoluble type, then by Lemma 2.1, G has an ordered Sylow tower of supersoluble type. Hence G is p -closed. By Lemma 2.2, we have that G is w-supersoluble. \square

Theorem 2.1. *Let A be a w-supersoluble \mathbb{P} -subnormal subgroup of G , and $G = AB$. Then G is w-supersoluble in each of the following cases:*

- (1) B is nilpotent and normal in G ;
- (2) B is nilpotent and $|G : B|$ is prime;
- (3) B is normal in G and is a siding group.

Proof. We prove all three statements at the same time using induction on the order of G . Note that G is soluble in any case. By Lemma 1.7, B is \mathbb{P} -subnormal in G and G has an ordered Sylow tower of supersoluble type by Lemma 2.1. If N is a non-trivial normal subgroup of G , then AN/N is \mathbb{P} -subnormal in G/N by Lemma 1.5 (2) and $AN/N \simeq A/A \cap N$ is w-supersoluble by Lemma 1.8. The subgroup $BN/N \simeq B/B \cap N$ is nilpotent or a siding group by Lemma 1.4 (1). Hence $G/N = (AN/N)(BN/N)$ is w-supersoluble by induction. Since the formation of all w-supersoluble groups is saturated by Lemma 1.8, we have G is a primitive group by Lemma 1.1. By Lemma 1.2, $F(G) = N = G_p$ is a unique minimal normal subgroup of G and $N = C_G(N)$, where p is the greatest in $\pi(G)$.

Since A is \mathbb{P} -subnormal in G , it follows that G has a subgroup M such that $A \leq M$ and $|G : M|$ is prime. By Dedekind's identity, $M = A(M \cap B)$. The subgroup A is \mathbb{P} -subnormal in M . The subgroup $M \cap B$ satisfies the requirements (1)–(3). By induction, M is w-supersoluble.

1. If B is nilpotent and normal in G , then $B = N$. Hence $G = AN$ and A is a maximal subgroup of G . Since A is \mathbb{P} -subnormal in G , we have $|G : A| = p = |N|$ and G is supersoluble. Therefore G is w-supersoluble. So, in (1), the theorem is proved.

2. Let B be nilpotent and $|G : B| = q$, where q is prime. Besides, let $|G : M| = r$, where r is prime. If $q \neq r$, then $(|G : M|, |G : B|) = 1$. Since $G = MB$, M and B are \mathbb{P} -subnormal in G and w-supersoluble, it follows obviously that G is w-supersoluble. Hence $q = r$. If $q = p$, then N is not contained in M . Thus $G = N \rtimes M$ and $|N|$ is prime. Consequently G is supersoluble and therefore G is w-supersoluble. So, $q \neq p$. Then $G_p = N \leq M \cap B$. Since B is nilpotent, $G_p = B \leq M$. Because $G = MB$, we have $G = M$, a contradiction. So, in (2), the theorem is proved.

3. Let B is normal in G and is a siding group. If B is nilpotent, then G is w-supersoluble by (1). Hence $B' \neq 1$. Because B' is normal in G and nilpotent, we have $N = B'$. If N is not contained in M , then $G = N \rtimes M$ and $|N|$ is prime. Consequently G is supersoluble and therefore G is w-supersoluble. Let N be contained in M and N_1 be a subgroup of prime order of N such that N_1 is normal in M . Then N_1 is normal in B by definition of siding group. Hence N_1 is normal in G . Consequently G is w-supersoluble. So, in (3), the theorem is proved. \square

Proof of Theorem 1 (2).

Note that by the Lemma 2.1, G is soluble. By Theorem 2.1 (1), Theorem 1 (2) is true. \square

Proof of Theorem 1 (3). Assume that the claim is false and let G be a minimal counterexample. By Lemma 2.1, G has an ordered Sylow tower of supersoluble type. If N is a non-trivial normal subgroup of G , then AN/N and BN/N are \mathbb{P} -subnormal in G/N by Lemma 1.5 (2). Besides, $AN/N \simeq A/A \cap N$ and $BN/N \simeq B/B \cap N$ are w-supersoluble by Lemma 1.8. By Lemma 1.3, we have

$$\begin{aligned} & (|(AN/N)/(AN/N)^A|, |(BN/N)/(BN/N)^A|) = \\ & = (|AN/(AN)^A N|, |BN/(BN)^A N|) = \\ & = (|AN/A^A N|, |BN/B^A N|) = \left(\frac{|A/A^A|}{|S_1|}, \frac{|B/B^A|}{|S_2|} \right), \\ & S_1 = (A \cap N)/(A^A \cap N), \quad S_2 = (B \cap N)/(B^A \cap N). \end{aligned}$$

Since $(|A/A^A|, |B/B^A|) = 1$, it follows that

$$(|(AN/N)/(AN/N)^A|, |(BN/N)/(BN/N)^A|) = 1.$$

The quotient $G/N = (AN/N)(BN/N)$ is w-supersoluble by induction.

Since the formation of all w-supersoluble groups is saturated by Lemma 1.8, we have G is a primitive group by Lemma 1.1. By Lemma 1.2, $F(G) = N =$

G_p is a unique minimal normal subgroup of G and $N = C_G(N)$, where p is the greatest in $\pi(G)$.

By Lemma 2.2, AN is w-supersoluble. If $AN = G$, then G is w-supersoluble, a contradiction. Hence in the future we consider that AN and BN are proper subgroups of G .

By Lemma 1.9(1), $(AN)^A$ is nilpotent. Since $N = C_G(N)$, we have $(AN)^A$ is a p -group. Because $AN/(AN)^A \in \mathcal{A}$, it follows that all Sylow r -subgroups of A are abelian, $r \neq p$. Since $A_p \leq G_p$, where A_p is a Sylow p -subgroup of A , we have $A \in \mathcal{A}$. Similarly, $B \in \mathcal{A}$. Hence $A^A = 1 = B^A$ and $(|A|, |B|) = (|A/A^A|, |B/B^A|) = 1$. It is clear that G is w-supersoluble, a contradiction. \square

3 Examples

The following example shows that for a soluble group $G = AB$ the mutually sn -permutability of subgroups A and B doesn't follow from \mathbb{P} -subnormality of these factors.

Example 3.1. The group $G = S_3 \rtimes Z_3$ (IdGroup=[18,3]) has \mathbb{P} -subnormal subgroups $A \simeq E_{3^2}$ and $B \simeq Z_2$. However A and B are not mutually sn -permutable.

The following example shows that we cannot omit the condition « G is p -closed» in Lemma 2.2.

Example 3.2. The group $G = (S_3 \times S_3) \rtimes Z_2$ (IdGroup=[72,40]) has a \mathbb{P} -subnormal supersoluble subgroups $A \simeq Z_3 \times S_3$. Besides $|G : A| = 2^2$ and Sylow 2-subgroup is maximal in G . Hence G is non-w-supersoluble.

The following example shows that in Theorem 2.1(1) the normality of subgroup B cannot be weakened to \mathbb{P} -subnormality.

Example 3.3. The group $G = (Z_2 \times (E_{3^2} \rtimes Z_4)) \rtimes Z_2$ (IdGroup=[144,115]) is non-w-supersoluble and factorized by subgroups $A = D_{12}$ and $B = Z_{12}$. The subgroup A has the chain of subgroups $A < S_3 \times S_3 < Z_2 \times S_3 \times S_3 < G$ and B has the chain of subgroups $B < Z_3 \times (Z_3 \rtimes Z_4) < (Z_3 \times (Z_3 \rtimes Z_4)) \rtimes Z_2 < G$. Therefore A and B are \mathbb{P} -subnormal in G .

The following example shows that in Theorem 2.1(2) it is impossible to weak the restrictions on the index of subgroup B .

Example 3.4. The alternating group $G = A_4$ is non-w-supersoluble and factorized by subgroups $A = E_{2^2}$ and $B = Z_3$. It is clear that A is supersoluble and \mathbb{P} -subnormal in G , and B is nilpotent and $|G : B| = 2^2$. The group $G = E_{5^2} \rtimes Z_3$ is non-w-supersoluble and has a nilpotent subgroup Z_3 of index 5^2 . Therefore even for the greatest p of $\pi(G)$, the index of B cannot be equal p^α , $\alpha \geq 2$.

The following example shows that in Theorem 2.1 (3) the normality of subgroup B cannot be weakened to subnormality.

Example 3.5. The group $G = Z_3 \times ((S_3 \times S_3) \rtimes Z_2)$ (IdGroup=[216,157]) is non-w-supersoluble and factorized by \mathbb{P} -subnormal supersoluble subgroup $A \simeq S_3 \times S_3$ and subnormal siding subgroup $B \simeq Z_3 \times Z_3 \times S_3$.

References

- [1] Ballester-Bolinches, A., Esteban-Romero, R., Asaad, M. (2010). *Products of finite groups*. Berlin; New York: Walter de Gruyter.
- [2] Ballester-Bolinches, A., Ezquerro, L.M. (2006). *Classes of Finite Groups*. Dordrecht: Springer.
- [3] Ballester-Bolinches, A., Fakieh, W. M., Pedraza-Aguilera, M. C. (2019). On Products of Generalised Supersoluble Finite Groups. *Mediterr. J. Math.* 16:46.
- [4] Carocca, A. (1998). On factorized finite groups in which certain subgroups of the factors permute. *Arch. Math.* 71:257–262.
- [5] GAP (2019) Groups, Algorithms, and Programming, Version 4.10.2. www.gap-system.org.
- [6] Huppert, B. (1967). *Endliche Gruppen*. Berlin: Springer-Verlag.
- [7] Monakhov, V. S. (2016). Finite groups with abnormal and \mathfrak{U} -subnormal subgroups. *Siberian Math. J.* 57:352–363.
- [8] Monakhov, V. S., Chirik, I. K. (2017). On the supersoluble residual of a product of subnormal supersoluble subgroups. *Siberian Math. J.* 58:271–280.
- [9] Monakhov, V. S., Kniahina, V. N. (2013). Finite group with \mathbb{P} -subnormal subgroups. *Ricerche Mat.* 62:307–323.

- [10] Monakhov, V. S., Trofimuk, A. A. (2019). Finite groups with two supersoluble subgroups. *J. Group Theory*. 22:297–312.
- [11] Monakhov, V. S., Trofimuk, A. A. (2020). On supersolubility of a group with seminormal subgroups. *Siberian Math. J.* 61:118–126.
- [12] Perez, E. R. (1999). On products of normal supersoluble subgroups. *Algebra Colloq.* 6:341–347.
- [13] Vasil'ev, A. F., Vasil'eva, T. I., Tyutyaynov, V. N. (2010). On the finite groups of supersoluble type. *Siberian Math. J.* 51:1004–1012.
- [14] Vasil'ev, A. F., Vasil'eva, T. I., Tyutyaynov, V. N. (2012). On the products of \mathbb{P} -subnormal subgroups of finite groups. *Siberian Math. J.* 53:47–54.