

A Generation Criterion for Subsets of $SL_n(\mathbb{F}_q)$

Ziv Greenhut
Tel Aviv University

January 11, 2022

Abstract

Let G_0 be either $SL_n(\mathbb{F}_q)$, the special linear group over the finite field with q elements, or $\mathbb{P}SL_n(\mathbb{F}_q)$, its projective quotient, and let Σ be a symmetric subset of G_0 , namely, if $x \in \Sigma$ then $x^{-1} \in \Sigma$. We find a certain set $\mathcal{R}(G_0)$ of irreducible representations of G_0 whose size is at most 5, such that Σ generates G_0 if and only if $|\Sigma|$ is not an eigenvalue of $\sum_{\sigma \in \Sigma} \rho(\sigma)$ for every $\rho \in \mathcal{R}(G_0)$.

To achieve this result, let G be either $GL_n(\mathbb{F}_q)$ or $\mathbb{P}GL_n(\mathbb{F}_q)$. We consider $\mathcal{X}(G)$, some set of irreducible nontrivial characters of G , whose size is at most 5. We show that for every subgroup $K \leq G$ that does not contain G_0 , the restriction to K of at least one of the characters in $\mathcal{X}(G)$ contains the trivial character as an irreducible summand. We then restrict the characters to G_0 and use standard arguments about the Cayley graph of G to imply the result. In addition, we obtain slightly weaker results about the generation of symmetric subsets of G .

We finish by considering S_n , the symmetric group on n elements, and presenting $\mathcal{R}(S_n)$, a set of eight irreducible nontrivial representations of S_n , such that a symmetric subset $\Sigma \subseteq S_n$ generates S_n if and only if $|\Sigma|$ is not an eigenvalue of $\sum_{\sigma \in \Sigma} \rho(\sigma)$ for every $\rho \in \mathcal{R}(S_n)$, which is an improvement upon the previously known set of 12 irreducible nontrivial representations of S_n that satisfies this condition.

Contents

1	Introduction	2
1.1	Main Results	5
1.2	Proof Strategy	6
1.3	Paper Organization	7
1.4	Notations	8
2	Induced Characters of Action Stabilizers	8
3	Characters of $GL_n(\mathbb{F}_q)$	13
4	The Maximal Subgroups of $GL_n(\mathbb{F}_q)$	15
4.1	The Geometric Maximal Subgroups	16
4.2	The Restriction of the Characters to the Geometric Subgroups	17
4.3	The Non-Geometric Subgroups for $n = 3$	20
5	Proofs of the Main Theorems for $n \geq 3$	21
5.1	The Special Linear Group	21
5.2	The Projective Groups	22
5.3	$GL_n(\mathbb{F}_q)$	23
5.4	$GL_4(\mathbb{F}_2)$	23
6	The Main Theorems for $n = 2$	24
6.1	$\mathcal{C}_3(GL_2(\mathbb{F}_q))$	25
6.2	$\mathcal{C}_6(GL_2(\mathbb{F}_q))$ and $\mathcal{S}(GL_2(\mathbb{F}_q))$	27

7	Open Questions	28
A	Characters and Subgroups of S_n	29
B	Group Extensions	31
	B.1 $c_{\min}(G)$ and $c_{\min}(G/N)$	31
	B.2 Abelian Groups	33
	B.3 $c_{\min}(G)$ and $c_{\min}(N)$	34
	References	36

1 Introduction

Let G be a finite group, and let Σ be a symmetric subset of G , namely, if $x \in \Sigma$ then $x^{-1} \in \Sigma$. We look for criteria to determine whether Σ generates G .

For a representation ρ of G , define

$$A_\rho(\Sigma) := \sum_{s \in \Sigma} \rho(s). \tag{1}$$

Then, the following proposition presents one such criterion.

Proposition 1.1. *Let G be a finite group, and let Σ be a symmetric subset of G . Then, Σ generates G if and only if for every nontrivial irreducible representation ρ of G , the maximal eigenvalue of $A_\rho(\Sigma)$ is strictly less than $|\Sigma|$.*

We prove this proposition using Cayley graphs. Before we prove the proposition, we define and state several facts about Cayley graphs.

Define $\text{Cay}(G, \Sigma)$, the Cayley graph of G with respect to Σ , as the $|\Sigma|$ -regular graph whose vertices are the elements of G , and edges are (g, gs) for all $g \in G$ and $s \in \Sigma$ (with a single edge between g and gs if $s^2 = 1$). By the definition of the Cayley graph, Σ generates G if and only if $\text{Cay}(G, \Sigma)$ is connected.

For a graph Γ with n vertices, let

$$\lambda_1(\Gamma) \geq \lambda_2(\Gamma) \geq \dots \lambda_n(\Gamma)$$

be the eigenvalues of the adjacency matrix of the graph. The graph is connected if and only if $\lambda_1(\Gamma)$ has multiplicity 1, namely, if $\lambda_1(\Gamma) \not\geq \lambda_2(\Gamma)$. In this situation, we say that the graph admits a spectral gap.

Applying it to the case of Cayley graphs, we deduce that the graph $\text{Cay}(G, \Sigma)$ admits a spectral gap if and only if Σ generates G . Therefore, we consider the adjacency matrix of the Cayley graph, and study its spectrum.

The adjacency matrix of $\text{Cay}(G, \Sigma)$ is equal to $A_{\rho_{\text{Reg}}}(\Sigma)$, where ρ_{Reg} is the right regular representation of G , that is, the permutation representation of the action of G on itself, defined by right multiplication.

A known fact in representation theory¹ states that the regular representation ρ_{Reg} decomposes to the sum of all of the irreducible representations of G , each with multiplicity equal to its dimension. Therefore, the eigenvalues of $A_{\rho_{\text{Reg}}}(\Sigma)$ are (up to multiplicities) the union of the eigenvalues of $A_\rho(\Sigma)$, for every irreducible representation ρ of G . As such, every eigenvalue of $A_{\rho_{\text{Reg}}}(\Sigma)$ can be associated with some irreducible representation ρ of G . For example, the eigenvalue associated with the trivial representation is $\lambda_1(\text{Cay}(G, \Sigma)) = |\Sigma|$.

We can now prove the proposition.

¹Throughout this paper we use standard facts about representation theory. [FH13] is a good reference for this subject.

Proof of Proposition 1.1. The set Σ generates G if and only if the Cayley graph $\text{Cay}(G, \Sigma)$ is connected. This graph is connected if and only if it admits a spectral gap, namely that $\lambda_2(\text{Cay}(G, \Sigma)) \not\leq |\Sigma|$.

The eigenvalues of $A_{\rho_{\text{Reg}}}(\Sigma)$ are (up to multiplicities) the union of the eigenvalues of $A_\rho(\Sigma)$, for every irreducible representation ρ of G . This implies that Σ generates G if and only if for every nontrivial irreducible representation ρ of G , the maximal eigenvalue of $A_\rho(\Sigma)$ is strictly less than $|\Sigma|$. \square

When $n \in \mathbb{Z}_{\geq 1}$ is large enough, consider the symmetric group S_n . It is known that for several types of sets Σ , there exists some set \mathcal{R} of irreducible representations of S_n with $|\mathcal{R}| \leq 8$, such that the second largest eigenvalue of the Cayley graph is always associated with some representation in \mathcal{R} . Then, it is much easier to determine the spectral gap $\lambda_1(\text{Cay}(S_n, \Sigma)) - \lambda_2(\text{Cay}(S_n, \Sigma))$. In particular, it is much easier to determine whether $\text{Cay}(S_n, \Sigma)$ admits a spectral gap, as the matrices of these representations are of a dimension much smaller than $n! = |S_n|$, the dimension of the regular representation. See [CLR10] and [PP18].

Given a finite group G , this problem can be naturally generalized to the problem of finding a relatively small set \mathcal{R} of irreducible representations of G , such that the second largest eigenvalue of $A_{\rho_{\text{Reg}}}(\Sigma)$ is associated with some representation in \mathcal{R} . Then, the second largest eigenvalue of $A_{\rho_{\text{Reg}}}(\Sigma)$ is the largest eigenvalue of one of the matrices $\{A_\rho(\Sigma) : \rho \in \mathcal{R}\}$.

In this paper, we consider the following relaxation of this generalized problem. Given some finite group G , we wish to find some relatively small set \mathcal{R} of irreducible representations of G , such that for every symmetric subset Σ , $\text{Cay}(G, \Sigma)$ admits a spectral gap (equivalently, Σ generates G) if and only if the eigenvalues of $A_\rho(\Sigma)$ are strictly less than $|\Sigma|$ for all $\rho \in \mathcal{R}$.

To study this problem, we introduce the following definitions. We use the notation $\langle f_1, f_2 \rangle_G := \frac{1}{|G|} \sum_{g \in G} f_1(g) \overline{f_2(g)}$, for $f_1, f_2: G \rightarrow \mathbb{C}$.

Definition 1.2. Let G be a finite group, and let ϕ and ψ be some characters of G . We write $\phi \supseteq \psi$ if there exists some character ϕ' of G such that $\phi = \psi + \phi'$. $\phi \supseteq \psi$

If ψ is an irreducible character, $\phi \supseteq \psi$ if and only if ψ is a summand of ϕ in its decomposition to irreducible characters, namely, if and only if $\langle \phi, \psi \rangle_G > 0$.

Definition 1.3. Let G be a finite group, let \mathcal{X} be a set of characters of G and let K be a subgroup of G . We write \mathcal{X} marks K if there exists some $\chi \in \mathcal{X}$ such that $\chi \supseteq \mathbb{1}_K$. \mathcal{X} marks K

If $\mathcal{X} = \{\chi\}$, we may omit the curly brackets and write that χ marks K .

The following proposition shows the connection between these definitions and the eigenvalues of $A_\rho(\Sigma)$. Note that the proposition considers only a subset of the nontrivial irreducible representations of G , unlike Proposition 1.1 that considers every nontrivial irreducible representation.

Proposition 1.4. *Let G be a finite group, and let \mathcal{X} be a set of nontrivial irreducible characters of G . For a character χ , let ρ_χ be the corresponding representation. Let $\mathcal{R} := \{\rho_\chi : \chi \in \mathcal{X}\}$, and assume that \mathcal{X} marks every proper subgroup of G . Let $\Sigma \subseteq G$ be a symmetric subset. Then the following are equivalent:*

1. The subset Σ generates G .
2. The graph $\text{Cay}(G, \Sigma)$ admits a spectral gap.
3. The maximal eigenvalue of $A_\rho(\Sigma)$ is strictly less than $|\Sigma|$, for all $\rho \in \mathcal{R}$.

Proof. The implications (1) \iff (2) \implies (3) follow from the discussion above. Thus, to finish the proof, we only need to prove the implication (3) \implies (1). We prove that if Σ does not generate G , there exists some $\rho \in \mathcal{R}$ for which the largest eigenvalue of $A_\rho(\Sigma)$ is equal to $|\Sigma|$.

Let $K := \langle \Sigma \rangle$, and assume $K \not\cong G$. Then, there exists some $\chi \in \mathcal{X}$ for which $\chi|_K \supseteq \mathbb{1}_K$. Let ρ be its representation.

Since $\Sigma \subseteq K$, $A_\rho(\Sigma) = A_{\rho|_K}(\Sigma)$, we deduce that the eigenvalues of $A_\rho(\Sigma)$ contain the single eigenvalue of $A_{\mathbb{1}_K}(\Sigma)$, which is precisely $|\Sigma|$. \square

Remark The condition that \mathcal{X} contains only nontrivial characters is important, since for every group G the character $\mathbb{1}_G$ marks every subgroup of G .

Example 1.5. Using this proposition, we see that the classification of finite multiply transitive groups (e.g. [DM96, Chapter 7.7] or [Cam81, Theorem 5.3]) gives a subset $\overline{\mathcal{X}}(S_n)$ of characters of S_n , for $n \geq 25$, such that $\overline{\mathcal{X}}(S_n)$ marks every proper subgroup of S_n .

Let $n \geq 25$, and let χ' be the character of the permutation representation associated with the action of S_n on tuples of 4 distinct elements of $\{1, \dots, n\}$. Denote $\chi = \chi' - \mathbb{1}_{S_n}$. It is easy to see (with the details given in Proposition 2.4) that for all $K \not\cong G$, $\chi|_K \not\supseteq \mathbb{1}_K$ if and only if K acts transitively on these 4-tuples. This is equivalent to saying that K is 4-transitive with respect to this action. By the classification, this implies that K is either S_n or the alternating group A_n .

Therefore, we define $\overline{\mathcal{X}}(S_n)$ to be the collection of irreducible summands of χ , along with the sign character. Then, $\overline{\mathcal{X}}(S_n)$ marks every proper subgroup of S_n . By calculating the decomposition of χ to irreducible characters, we deduce that the characters in $\overline{\mathcal{X}}(S_n)$ are parametrized² by the following³ 12 partitions of n :

$$\left\{ \begin{array}{cccc} (n-1, 1), & (n-2, 2), & (n-2, 1^2), & (n-3, 3), \\ (n-3, 2, 1), & (n-3, 1^3), & (n-4, 4), & (n-4, 3, 1), \\ (n-4, 2^2), & (n-4, 2, 1^2), & (n-4, 1^4), & (1^n) \end{array} \right\}.$$

For a prime power q , let \mathbb{F}_q be the field with q elements. In this paper, we prove similar results about the general linear group over a finite field $GL_n(\mathbb{F}_q)$ and its relatives – $\mathbb{P}GL_n(\mathbb{F}_q)$, $SL_n(\mathbb{F}_q)$ and $\mathbb{P}SL_n(\mathbb{F}_q)$. For $n \geq 3$, the group $GL_n(\mathbb{F}_q)$ acts 2-transitively on the projective space $\mathbb{P}^{n-1}\mathbb{F}_q$, and every subgroup of $GL_n(\mathbb{F}_q)$ with a 2-transitive action on the projective space contains $SL_n(\mathbb{F}_q)$ ([CK79]), except in the exceptional case where $n = 4$ and $q = 2$. Thus, we may attempt to apply the methods we used for S_n on these groups, along with the classification of multiply transitive groups. However, two differences with the case of S_n arise.

First, in the case of S_n , we define a set of characters $\mathcal{X}(S_n)$ that marks every subgroup of S_n that does not contain A_n , and then we add the sign character, which satisfies $\text{sgn}|_{A_n} \supseteq \mathbb{1}_{A_n}$, to obtain $\overline{\mathcal{X}}(S_n)$.

In the case of $GL_n(\mathbb{F}_q)$, the same arguments would give a set of characters $\mathcal{X}(GL_n(\mathbb{F}_q))$ that marks every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$. In general, there exists no single character we can add to $\mathcal{X}(GL_n(\mathbb{F}_q))$ to obtain some set $\overline{\mathcal{X}}(GL_n(\mathbb{F}_q))$ that marks every proper subgroup of $GL_n(\mathbb{F}_q)$ (see Corollary B.9), and the number of characters we need to add can be very large for values of q where $q - 1$ has a large number of prime factors.

Thus, we only find a set of characters $\mathcal{X}(GL_n(\mathbb{F}_q))$ that marks every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$ in this paper. Then, by restricting the characters in $\mathcal{X}(GL_n(\mathbb{F}_q))$ to $SL_n(\mathbb{F}_q)$ (all of the characters in this set reduce irreducibly), we obtain a set of characters $\mathcal{X}(SL_n(\mathbb{F}_q))$ of $SL_n(\mathbb{F}_q)$ that marks every proper subgroup of $SL_n(\mathbb{F}_q)$.

Second, the character of the permutation representation associated with the action of $GL_n(\mathbb{F}_q)$ on pairs of projective vectors in $\mathbb{P}^{n-1}\mathbb{F}_q$ decomposes to $q + 3$ distinct irreducible characters for every $n \geq 4$. Our aim is to obtain sets $\mathcal{X}(GL_n(\mathbb{F}_q))$ of nontrivial irreducible characters whose sizes are bounded uniformly in n and q .

²[Ful97] is a good reference for this connection between partitions and characters of S_n , along with the representation theory of S_n in general.

³We present a smaller set of characters with 8 elements in Appendix A.

Therefore, we take a slightly different approach. Then, we obtain sets $\mathcal{X}(GL_n(\mathbb{F}_q))$ with $|\mathcal{X}(GL_n(\mathbb{F}_q))| \leq 5$.

1.1 Main Results

In the representation theory of $GL_n(\mathbb{F}_q)$, several classes of characters arise. One of these classes is the class of unipotent characters, which are irreducible characters parametrized by partitions of n (for more information about the representation theory of $GL_n(\mathbb{F}_q)$, see [Section 3](#)).

The unipotent characters are closely related to the action of $GL_n(\mathbb{F}_q)$ on the projective space $\mathbb{P}^{n-1}\mathbb{F}_q$, and have a fundamental role in the group's representation theory. For example, $\chi^{(n-1,1)}(g)$ counts the number of fixed points of g in this action, minus 1. The character $\chi^{(n-2,2)}(g)$ counts the number of 2-dimensional projective subspaces W with $gW = W$, minus the number of fixed points of g .

It turns out that the unipotent characters can be defined in a natural way for the projective and special groups as well – see [Section 5.1](#) and [Section 5.2](#). Thus, by abuse of notation, we define

$$\mathcal{X}(G) := \{\chi^{(n-3,1^3)}, \chi^{(n-3,2,1)}, \chi^{(n-2,1^2)}, \chi^{(n-2,2)}, \chi^{(n-1,1)}\} \subseteq \widehat{G}, \quad \mathcal{X}(G)$$

for G either $GL_n(\mathbb{F}_q)$ or one of its relatives, when $n \geq 3$ and $(n, q) \neq (4, 2)$. The character χ^α is omitted if α is not a partition⁴.

Theorem 1.6. *Let $n \geq 3$ and q a prime power, such that $(n, q) \neq (4, 2)$. Let G be either $GL_n(\mathbb{F}_q)$ or $\mathbb{P}GL_n(\mathbb{F}_q)$, and let $G' \leq G$ be $SL_n(\mathbb{F}_q)$ or $\mathbb{P}SL_n(\mathbb{F}_q)$ respectively. Then, $\mathcal{X}(G)$ marks every subgroup of G that does not contain G' .*

Theorem 1.7. *Let $n \geq 3$ and q a prime power, such that $(n, q) \neq (4, 2)$. Let G be either $SL_n(\mathbb{F}_q)$ or $\mathbb{P}SL_n(\mathbb{F}_q)$. Then, $\mathcal{X}(G)$ marks every proper subgroup of G .*

Note that we do not claim that these sets $\mathcal{X}(G)$ are unique, nor that they are minimal.

GL₄(F₂) For completeness, we present the following theorem that deals with the exceptional case of the previous theorems. For this case, let ψ be the single irreducible character of degree 7 of $GL_4(\mathbb{F}_2) \cong A_8$ (the character of the permutation representation associated with the action on 8 elements, minus the trivial). Here, $q = 2$ implies that $GL_4(\mathbb{F}_2)$ is equal to all of its relatives. Denote

$$\mathcal{X}(GL_4(\mathbb{F}_2)) := \{\chi^{(1^4)}, \chi^{(2,1^2)}, \chi^{(2^2)}, \chi^{(3,1)}, \psi\} \subseteq \widehat{GL_4(\mathbb{F}_2)}. \quad \mathcal{X}(GL_4(\mathbb{F}_2))$$

Theorem 1.8. *$\mathcal{X}(GL_4(\mathbb{F}_2))$ marks every proper subgroup of $GL_4(\mathbb{F}_2)$.*

GL₂(F_q) We present here the case of $n = 2$ as well. In this case, there exists a single nontrivial unipotent character, $\chi^{(1,1)}$, which does not encode enough information about subgroups of $GL_2(\mathbb{F}_q)$, since $\chi^{(1,1)}|_K \not\cong \mathbb{1}_K$ for every $K \leq GL_2(\mathbb{F}_q)$ that acts transitively on $\mathbb{P}^1\mathbb{F}_q$. There are such subgroups that do not contain $SL_2(\mathbb{F}_q)$. For example, $GL_1(\mathbb{F}_{q^2})$ acts transitively on $\mathbb{F}_{q^2} \setminus 0$. By the isomorphism $\mathbb{F}_{q^2} \cong \mathbb{F}_q^2$ of vector spaces over \mathbb{F}_q . We get an inclusion $GL_1(\mathbb{F}_{q^2}) \not\leq GL_2(\mathbb{F}_q)$. Thus, $GL_2(\mathbb{F}_q)$ has transitive subgroups that do not contain $SL_2(\mathbb{F}_q)$, and an additional character is required.

For G either $GL_2(\mathbb{F}_q)$ or one of its relatives, consider the irreducible character ψ_1 , defined in [Definition 6.1](#). Denote

$$\mathcal{X}(G) := \{\chi^{(1,1)}, \psi_1\} \subseteq \widehat{G}.$$

⁴For example, when $n = 3$, $(n - 2, 2) = (1, 2)$ which is not a partition, because the numbers are not non-increasing.

Theorem 1.9. *Let q be a prime power, let G be either $GL_2(\mathbb{F}_q)$ or $\mathbb{P}GL_2(\mathbb{F}_q)$, and let $G' \leq G$ be $SL_2(\mathbb{F}_q)$ or $\mathbb{P}SL_2(\mathbb{F}_q)$ respectively. Then, $\mathcal{X}(G)$ marks every subgroup of G that does not contain G' .*

Theorem 1.10. *Let q be a prime power, and let G be either $SL_2(\mathbb{F}_q)$ or $\mathbb{P}SL_2(\mathbb{F}_q)$. Then, $\mathcal{X}(G)$ marks every proper subgroup of G .*

1.2 Proof Strategy

Consider the family of subgroups $GL_n(\mathbb{F}_q)$ with $n \geq 3$, along with their special subgroups $SL_n(\mathbb{F}_q) \leq GL_n(\mathbb{F}_q)$. Denote by $\widehat{GL_n(\mathbb{F}_q)}$ the collection of irreducible characters of $GL_n(\mathbb{F}_q)$. We wish to find sets of nontrivial characters $\mathcal{X}(GL_n(\mathbb{F}_q)) \subseteq \widehat{GL_n(\mathbb{F}_q)}$ of bounded size (with respect to both n and q), that mark every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$.

The following notion of transitivity plays an important role in the proof:

Definition 1.11. Let G be a finite group acting on some set X , let $K \leq G$, let $\ell \in \mathbb{Z}_{\geq 1}$, and let x_1, \dots, x_ℓ be distinct elements of X . We say that K is (x_1, \dots, x_ℓ) - G -transitive if for every $g \in G$, there exists $k \in K$ such that $\forall i: kx_i = gx_i$. (x_1, \dots, x_ℓ) - G -transitive

Example 1.12. 1. (x_1, \dots, x_ℓ) - G -transitivity is equivalent to transitivity on the orbit $O_G((x_1, \dots, x_\ell))$ in the diagonal action of G on X^ℓ .

2. For $\ell \leq n$, a subgroup $K \leq S_n$ is $(1, \dots, \ell)$ - S_n -transitive if and only if the action of K on $\{1, \dots, n\}$ is ℓ -transitive.

3. If the action of G on X is ℓ -transitive, $K \leq G$ is (x_1, \dots, x_ℓ) - G -transitive if and only if the action of K is ℓ -transitive.

4. Let \bar{e}_i be the point in $\mathbb{P}^{n-1}\mathbb{F}_q$ represented by the i th element of the standard basis of \mathbb{F}_q^n . For $\ell \leq n$, $K \leq GL_n(\mathbb{F}_q)$ is $(\bar{e}_1, \dots, \bar{e}_\ell)$ - $GL_n(\mathbb{F}_q)$ -transitive if and only if for every pair of ℓ -tuples of linearly independent projective vectors – (x_1, \dots, x_ℓ) and (y_1, \dots, y_ℓ) – there exists an element $k \in K$ such that $\forall i: kx_i = y_i$.

5. $SL_n(\mathbb{F}_q)$ is $(\bar{e}_1, \dots, \bar{e}_n)$ - $GL_n(\mathbb{F}_q)$ -transitive.

We can now describe the framework for the proof. Let $G_n = GL_n(\mathbb{F}_q)$ for $n \geq 3$, acting on the sets $X_n = \mathbb{P}^{n-1}\mathbb{F}_q$. The sets X_n are increasing with respect to inclusion $i_n: \mathbb{P}^{n-1}\mathbb{F}_q \hookrightarrow \mathbb{P}^n\mathbb{F}_q$, $i_n(x_1, \dots, x_n) = (x_1, \dots, x_n, 0)$. Denote $Y := X_3 = \mathbb{P}^2\mathbb{F}_q$, and consider the stabilizer $(G_n)_Y = \{g \in G_n: gY = Y\}$, which can be realized as the group of upper diagonal block matrices

$$P_{(3,n-3)} := \begin{bmatrix} GL_3(\mathbb{F}_q) & * \\ 0 & GL_{n-3}(\mathbb{F}_q) \end{bmatrix} \leq GL_n(\mathbb{F}_q).$$

It projects to the symmetry group of Y , and we denote its image by H . Note that $H \cong \mathbb{P}GL_3(\mathbb{F}_q)$.

Given a character of H , we pull it back using the projection from the stabilizer. Here we take a character of a small group, with a relatively small degree, and lift it to a much larger group without increasing its degree, nor the number of its irreducible summands. We get a character of $(G_n)_Y$ which is then induced to the group G_n .

This allows us to study H and its characters, and then use the described process to lift them to the sequence of groups G_n and obtain results on these characters. Thus, assume that we have some subset $\mathcal{X}(H) \subseteq \widehat{H} = \widehat{\mathbb{P}GL_3(\mathbb{F}_q)}$, such that $\mathcal{X}(H)$ marks every subgroup of H that is not $(\bar{e}_1, \bar{e}_2, \bar{e}_3)$ - H -transitive. Lift the characters to characters of $(G_n)_Y$, then induce them to characters of G_n . Denote the induced characters by $\mathcal{X}(G_n)'$.

We prove that since H , $\mathcal{X}(H)$, G_n and Y satisfy several properties, which are described below (with $\ell = 3$), $\mathcal{X}(G_n)'$ marks every subgroup of G_n that is not (\bar{e}_1, \bar{e}_2) - G_n -transitive.

It turns out that this implies that $\mathcal{X}(GL_n(\mathbb{F}_q))'$ marks every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$, outside the exceptional case of $q = 2, n = 4$. Therefore, to finish the proof of the main theorems, it remains to decompose the characters in $\mathcal{X}(G_n)'$ to irreducible characters, and deduce the theorems.

The method described above can be used for every sequence of groups G_n acting on sets X_n , which are increasing with respect to inclusion. In the general case, we need the following three conditions to hold in order to apply this method:

1. Y contains some ℓ element subset $Y_0 = \{a_1, \dots, a_\ell\}$, which can be thought of as the basis of Y .
2. The translations of Y by elements of G_n cover all ℓ -element subsets of X_n .
3. If we can translate less than ℓ of the elements of Y_0 , with some element of G_n , while keeping them in Y , we can translate these same elements in the same way with some element of $(G_n)_Y$.

Then, we do the following to finish the proof:

1. Find some subset of characters $\mathcal{X}(H) \subseteq \widehat{H}$ that marks every subgroup of H that is not (a_1, \dots, a_ℓ) - H -transitive.
2. Decompose the characters of $\mathcal{X}(G_n)'$ to irreducibles, to obtain some set $\mathcal{X}(G_n)$.
3. Classify⁵ all $(a_1, \dots, a_{\ell-1})$ - G_n -transitive subgroups of G_n .

To finish the proof of the main theorems, we need to make sure the decomposed characters $\mathcal{X}(GL_n(\mathbb{F}_q))$ restrict to irreducible characters of $SL_n(\mathbb{F}_q)$ and project to irreducible characters of $\mathbb{P}GL_n(\mathbb{F}_q)$ and $\mathbb{P}SL_n(\mathbb{F}_q)$ for almost all pairs $n \in \mathbb{Z}_{\geq 1}$, q a prime power. Then we deal with the exceptional cases that arise.

1.3 Paper Organization

In [Section 2](#), we study characters induced by action stabilizers, which are the characters of permutation representations. The main result of this section is [Corollary 2.9](#). To prove it, we prove several technical lemmas. In general, we prove that there exist certain characters such that if their restriction to some $K \leq G$ does not contain the trivial representation, the action of K has certain transitivity properties.

In [Section 3](#), we introduce general facts about the characters of $GL_n(\mathbb{F}_q)$, and study certain characters that are important to the rest of the paper.

In [Section 4](#), we study maximal subgroups of $GL_n(\mathbb{F}_q)$. [Section 4.1](#) describes the families of a class of maximal subgroups called the geometric subgroups, [Section 4.2](#) restricts certain characters to the geometric subgroups and proves these do not contain the trivial representation, and [Section 4.3](#) handles the remaining case of the non-geometric maximal subgroups.

In [Section 5](#), we prove the main theorems for $n \geq 3$: for the general case ([Section 5.3](#)), the special cases ([Section 5.1](#)), the projective cases ([Section 5.2](#)), and the exceptional case of $GL_4(\mathbb{F}_2)$ ([Section 5.4](#)).

In [Section 6](#), we prove the main theorems for $n = 2$.

In [Section 7](#), we present several open questions raised in this paper.

In [Appendix A](#), we apply the method described in the introduction on S_n and state the results.

⁵The classification in the case of $GL_n(\mathbb{F}_q)$ is actually implied by the classification of multiply transitive groups, and is not done in this paper

In [Appendix B](#), we consider the situation where G is some finite group, with $N \trianglelefteq G$. We show how sets of nontrivial irreducible characters $\mathcal{X}(G)$, $\mathcal{X}(G/N)$ and $\mathcal{X}(N)$ behave with respect to one another, where these sets mark proper subgroups of G , G/N and N respectively.

1.4 Notations

Symbol	Definition
$\text{Tr}_{\mathbb{E}/\mathbb{F}}$	The trace of the field extension \mathbb{E}/\mathbb{F}
$\mathbb{P}V$	The projective space of V
\bar{v}	The point in the projective space represented by v
$\text{Ch}(G)$	The collection of characters of G (non-virtual)
\widehat{G}	The collection of irreducible characters of G
$\langle u, v \rangle$	The inner product of u and v
$\langle \psi, \phi \rangle_G$	The inner product of ψ and ϕ in the character ring of G
$\mathbb{1}_G$	The trivial character of G
$\text{Span}(v_1, \dots, v_k)$	The linear span of v_1, \dots, v_k
$e_i \in \mathbb{F}_q^n$	The i th element of the standard basis of \mathbb{F}_q^n
$\langle g_1, \dots, g_k \rangle$	The subgroup generated by $g_1, \dots, g_k \in G$
$\text{Ind}_K^G(\chi)$	The induced character from K to G
$Z(G)$	The center of G
S_n	The symmetric group on n elements
A_n	The alternating group on n elements
$G \circlearrowleft X$	G acts on X
G_Y	The stabilizer of $Y = \{g \in G : gY = Y\}$, where $G \circlearrowleft X$ and $Y \subseteq X$
$O_G(Y)$	The orbit of $Y = \{gy : g \in G, y \in Y\}$ where $G \circlearrowleft X$ and $Y \subseteq X$
$N_G(K)$	The normalizer of $K \leq G$
$\lambda \vdash n$	λ is a partition of n
λ'	The partition conjugate to $\lambda \vdash n$

2 Induced Characters of Action Stabilizers

In this section, we relate the notion of (y_1, \dots, y_ℓ) - G -transitivity and the properties of characters of groups. We start by stating several standard results in representation theory that we need.

Theorem 2.1 (Frobenius Reciprocity). *Let G be a group and let $K \leq G$. Let $\psi \in \text{Ch}(K)$ and $\phi \in \text{Ch}(G)$. Then*

$$\langle \text{Ind}_K^G(\psi), \phi \rangle_G = \langle \psi, \phi|_K \rangle_K.$$

Proposition 2.2 (e.g. [\[FH13, Equation 3.18\]](#)). *Let G be a finite group, let $K \leq G$, and let $\chi \in \text{Ch}(K)$. Then*

$$\text{Ind}_K^G(\chi)(g) = \frac{1}{|K|} \sum_{t \in G \text{ s.t. } t^{-1}gt \in K} \chi(t^{-1}gt).$$

Lemma 2.3 (Burnside's Lemma). *Let G be a finite group acting on some set X . For each $g \in G$, denote by $X^g := \{x \in X : gx = x\}$. Let $|G \backslash X|$ denote the number of orbits of the action. Then*

$$|G \backslash X| = \frac{1}{|G|} \sum_{g \in G} |X^g|.$$

Proposition 2.4. *Let G be a finite group acting transitively on some set X , let $K \leq G$, and let χ be the character of the permutation representation associated with the action, namely, $\chi(g)$ is the number of $x \in X$ fixed by g . Then, $\langle \chi|_K, \mathbb{1}_K \rangle_K = 1 \iff K$ acts transitively on X .*

Proof. By [Burnside's Lemma](#),

$$\langle \chi|_K, \mathbb{1}_K \rangle_K = \frac{1}{|K|} \sum_{g \in K} \chi(g) = \frac{1}{|K|} \sum_{g \in K} |X^g| = \left| \frac{K \backslash X}{K} \right|.$$

The right hand side is 1 if and only if K acts transitively on X . \square

The equality $\langle \chi|_K, \mathbb{1}_K \rangle_K = 1$ means that $\chi|_K \supseteq \mathbb{1}_K$ and $\mathbb{1}_K$ has multiplicity 1 in $\chi|_K$.

Remark 2.5. *The character of the permutation representation associated with a transitive group action is equal to the character $\text{Ind}_{G_x}^G(\mathbb{1}_{G_x})$, induced from the stabilizer G_x for every $x \in X$.*

Corollary 2.6. *Let G be a finite group acting transitively on some set X , and let χ be the character of the permutation representation associated with the action. Denote $\chi' := \chi - \mathbb{1}_G$. Then, χ' marks every subgroup of G that does not act transitively on X .*

Notation for the central example Here, we introduce notation useful for the remainder of this section. We start by an example and then define the notation in general.

Let $G = GL_n(\mathbb{F}_q)$ acting on $X = \mathbb{P}^{n-1}\mathbb{F}_q$, and let $\ell \leq n$. Consider the projective subspace $Y = \mathbb{P}(\text{Span}(e_1, \dots, e_\ell)) \subseteq X$, whose translations by elements of G are precisely the ℓ -dimensional projective subspaces of X . The stabilizer G_Y is

$$P_{(\ell, n-\ell)} := \left[\begin{array}{c|c} GL_\ell(\mathbb{F}_q) & * \\ \hline 0 & GL_{n-\ell}(\mathbb{F}_q) \end{array} \right] \leq GL_n(\mathbb{F}_q).$$

Let $p: P_{(\ell, n-\ell)} \rightarrow \mathbb{P}GL_\ell(\mathbb{F}_q)$ be the projection to the upper left block, composed with the projection to the projective group. This corresponds to the restriction of the action of G_Y to Y , with p being the projection from G_Y to $\text{Sym}(Y)$, the symmetric group on the set Y . We denote by H the image of this morphism – $\mathbb{P}GL_\ell(\mathbb{F}_q)$.

For every $t \in GL_n(\mathbb{F}_q)$, we define p_t and H_t in a similar way – by projecting from G_{tY} to $\text{Sym}(tY)$. Note that the translations tY of Y are the ℓ -dimensional projective subspaces of X .

For every $t \in GL_n(\mathbb{F}_q)$, the group H_t is isomorphic to $\mathbb{P}GL_\ell(\mathbb{F}_q)$, as this is the group of linear morphisms of the ℓ -dimensional projective space tY . Let $f_t: H \rightarrow H_t$ be the natural isomorphisms, defined by lifting an element of H to some element of G_Y , conjugating by t and projecting to H_t . The resulting element of H_t is independent of the choice of the element of G_Y .

Let χ be a character of $H = \mathbb{P}GL_\ell(\mathbb{F}_q)$. The pull-back of χ to G_Y through the projection then defines a new character $\text{P}_H^{G_Y}(\chi) \in \text{Ch}(G_Y)$, which is applied on an element of G_Y as follows:

$$\text{P}_H^{G_Y}(\chi) \left(\begin{bmatrix} A & B \\ 0 & C \end{bmatrix} \right) = \chi(A).$$

The character $\text{P}_H^{G_Y}(\chi)$ is then induced to G to define a character $\text{R}_H^G(\chi) \in \text{Ch}(G)$.

Before we move to the general case, we define the pull-back of a character. Let G and H be finite groups. For a character χ of H and a morphism $f: G \rightarrow H$, we denote by $f^*\chi$ the pull-back of χ by f . The pull-back is a character of G , with $f^*\chi(g) = \chi(f(g))$.

Notation 2.7. Given a finite group G acting on some set X , a subset $Y \subseteq X$ and some $t \in G$, define the following.

1. We denote by $H_t \leq \text{Sym}(tY)$ the restriction of the action of G_{tY} on tY (formally – H_t the quotient of G_{tY} by the kernel of its action on tY).

2. The corresponding epimorphism is denoted by $p_t: G_{tY} \twoheadrightarrow H_t$. p_t
3. We specialize the case when $t = e$, the identity of G , and denote $H := H_e$ and $p := p_e$. H, p
4. Denote by $f_t: H \rightarrow H_t$ – the "conjugation maps"⁶ $f_t(g) := p_t(tp^{-1}(g)t^{-1})$, which make f_t the following diagram commute. Note that f_t is an isomorphism.

$$\begin{array}{ccc}
G_Y^g & \xrightarrow{tgt^{-1}} & G_{tY} \\
p \downarrow & & \downarrow p_t \\
Y \circ H & \xrightarrow{f_t} & H_t \circ tY
\end{array}$$

5. For a character χ of H :

- 5.1. Define $\chi_t := (f_t^{-1})^*\chi \in \text{Ch}(H_t)$, that is, $\chi_t(g) = \chi(f_t^{-1}(g))$.
- 5.2. Define $P_H^{G_Y}(\chi) := p^*\chi$, that is, $P_H^{G_Y}(\chi)(g) := \chi(p(g))$.
- 5.3. Define⁷ $R_H^G(\chi) \in \text{Ch}(G)$ as the induction of $P_H^{G_Y}(\chi)$ to G , namely, $R_H^G(\chi) := \text{Ind}_{G_Y}^G(P_H^{G_Y}(\chi))$.

In the rest of this section, we study the induced characters $R_H^G(\chi)$. Our end goal is the following lemma:

Lemma 2.8. *Let G be a finite group acting on some set X and let $Y \subseteq X$. Using Notation 2.7 let $\chi \in \text{Ch}(H)$. Assume:*

1. $Y \supseteq Y_0 = \{a_1 \dots a_\ell\}$
2. χ marks every subgroup of H that is not (a_1, \dots, a_ℓ) - H -transitive.
3. The translations of Y cover all ℓ -element subsets of X , that is, for every $B \subseteq X$ with $|B| \leq \ell$, there exists some $t \in G$ such that $B \subseteq tY$.
4. For every $t \in G$ and $i < \ell$, if $ta_j \in Y$ for all $j \leq i$, then there exists some $s \in G_Y$ such that $\forall j \leq i: sa_j = ta_j$.

Then $R_H^G(\chi)$ marks every subgroup of G that is not $(a_1, \dots, a_{\ell-1})$ - G -transitive.

We apply the lemma on $G = GL_n(\mathbb{F}_q)$ acting on $X = \mathbb{P}^{n-1}\mathbb{F}_q$, along with $Y = \mathbb{P}(\text{Span}(e_1, \dots, e_\ell)) \subseteq X$ and $Y_0 = \{\bar{e}_1, \dots, \bar{e}_\ell\}$. Since $\mathbb{P}SL_\ell(\mathbb{F}_q)$ is $(\bar{e}_1, \dots, \bar{e}_\ell)$ - $\mathbb{P}GL_\ell(\mathbb{F}_q)$ -transitive, if χ is a character of $\mathbb{P}GL_\ell(\mathbb{F}_q)$ that marks every subgroup of $\mathbb{P}GL_\ell(\mathbb{F}_q)$ that does not contain $\mathbb{P}SL_\ell(\mathbb{F}_q)$, χ also marks every subgroup of $\mathbb{P}GL_\ell(\mathbb{F}_q)$ that is not $(\bar{e}_1, \dots, \bar{e}_\ell)$ - $\mathbb{P}GL_\ell(\mathbb{F}_q)$ -transitive.

Corollary 2.9. *Let $\ell \leq n$ be integers, let $G = GL_n(\mathbb{F}_q)$, $H = \mathbb{P}GL_\ell(\mathbb{F}_q)$, and let $K \leq G$. Let $\chi \in \text{Ch}(H)$ be a character that marks every subgroup of H that does not contain $\mathbb{P}SL_\ell(\mathbb{F}_q)$. Then, $R_H^G(\chi)$ marks every subgroup of G that is not $(\bar{e}_1, \dots, \bar{e}_{\ell-1})$ - G -transitive.*

Lemma 2.8 is not limited to the case of $GL_n(\mathbb{F}_q)$. For example, by applying the lemma for $G = S_n, Y = Y_0 = \{1, \dots, \ell\}$, we deduce:

Corollary 2.10. *Let $\ell \leq n$ be integers, and let $K \leq S_n$. Assume there exists some $\chi \in \text{Ch}(S_\ell)$ such that χ marks every proper subgroup of S_ℓ . Then, $R_{S_\ell}^{S_n}(\chi)$ marks every subgroup of S_n that is not $(\ell - 1)$ -transitive.*

⁶The morphisms f_t are well defined, as $a \in \ker(p) \Rightarrow tat^{-1} \in \ker(p_t)$.

⁷This definition is similar to the definition of \circ , described in Section 3.

Remark [Corollary 2.10](#) gives an example where we "lose a transitivity degree", that is, the resulting character $\mathbf{R}_H^G(\chi)$ does not mark all subgroups of G that are not $(a_1, \dots, a_{\ell-1}, a_\ell)$ - G -transitive, but only those that are not $(a_1, \dots, a_{\ell-1})$ - G -transitive. For example, choose $G = S_4$ with the usual action on $\{1, 2, 3, 4\}$, $Y = \{1, 2\}$, $\ell = 2$, $H = S_2$ and χ the single nontrivial irreducible character of H . Then, $\mathbf{R}_{S_2}^{S_4}(\chi)$ decomposes as the sum of the characters parametrized by $(3, 1)$ and $(2, 1, 1)$. These are not enough to mark all subgroups of S_4 that are not 2-transitive, as neither marks the subgroups of S_4 of index 3, which are conjugate to $K = \langle (1, 2, 3, 4), (1, 3) \rangle$ (with (x_1, \dots, x_a) denoting the cyclic permutation of x_1, \dots, x_a).

We prove [Lemma 2.8](#) in several steps. Let $K \leq G$ and $t \in G$. We consider the action of $K \cap G_{tY}$ on tY . To understand this action, we can consider the projection to H_t – the group $J_t := p_t(K \cap G_{tY})$, which is the quotient of $K \cap G_{tY}$ by the kernel of this action.

In [Lemma 2.11](#), we deduce that if $\mathbf{R}_H^G(\chi)|_K \not\supseteq \mathbb{1}_K$, then $\chi_t|_{J_t} \not\supseteq \mathbb{1}_{J_t}$ for all $t \in G$. This allows us to reduce the question about the subgroups and characters of G to a question about the subgroups and characters of H .

In [Lemma 2.12](#), we assume there exist some $a_1, \dots, a_\ell \in Y$ and some character χ of H that marks every subgroup of H that is not (a_1, \dots, a_ℓ) - H -transitive. Then, we show that this χ can be "pulled-back" to show that $\mathbf{R}_H^G(\chi)|_K \not\supseteq \mathbb{1}_K$ implies that $K \cap G_{tY}$ is (a_1, \dots, a_ℓ) - G_{tY} -transitive for all $t \in G$. We proceed to proving the lemmas.

Lemma 2.11. *Continuing with [Notation 2.7](#), let G be a finite group acting on some set X , let $Y \subseteq X$ be some subset, and let $\chi \in \text{Ch}(H)$. For every $K \leq G$, we have $\mathbf{R}_H^G(\chi)|_K \not\supseteq \mathbb{1}_K$ if and only if for every $t \in G$, the group $J_t := p_t(K \cap G_{tY})$ satisfies $\chi_t|_{J_t} \not\supseteq \mathbb{1}_{J_t}$.*

Proof. Let $K \leq G$. For every $t \in G$, $J_t = f_t(p((t^{-1}Kt) \cap G_Y))$. Now,

$$\begin{aligned}
\langle \mathbf{R}_H^G(\chi)|_K, \mathbb{1}_K \rangle_K &= \frac{1}{|K|} \sum_{g \in K} \mathbf{R}_H^G(\chi)(g) \stackrel{(1)}{=} \frac{1}{|K|} \sum_{g \in K} \frac{1}{|G_Y|} \sum_{t \in G \text{ s.t. } t^{-1}gt \in G_Y} \mathbf{P}_H^{G_Y}(\chi)(t^{-1}gt) \\
&= \frac{1}{|K| \cdot |G_Y|} \sum_{t \in G} \left(\sum_{g \in K \text{ s.t. } t^{-1}gt \in G_Y} \mathbf{P}_H^{G_Y}(\chi)(t^{-1}gt) \right) \\
&= \frac{1}{|K| \cdot |G_Y|} \sum_{t \in G} \left(\sum_{g' \in G_Y \text{ s.t. } tg't^{-1} \in K} \mathbf{P}_H^{G_Y}(\chi)(g') \right) \\
&= \frac{1}{|K| \cdot |G_Y|} \sum_{t \in G} \left(\sum_{g' \in (t^{-1}Kt) \cap G_Y} \chi(p(g')) \right) \\
&= \frac{1}{|K| \cdot |G_Y|} \sum_{t \in G} |\ker(p|_{(t^{-1}Kt) \cap G_Y})| \left(\sum_{h \in p((t^{-1}Kt) \cap G_Y)} \chi(h) \right) \\
&= \frac{1}{|K| \cdot |G_Y|} \sum_{t \in G} |\ker(p|_{(t^{-1}Kt) \cap G_Y})| \left(\sum_{h' \in J_t} \chi_t(h') \right) \\
&= \sum_{t \in G} c_t \langle \chi_t|_{J_t}, \mathbb{1}_{J_t} \rangle_{J_t},
\end{aligned}$$

where (1) follows from [Proposition 2.2](#), and

$$c_t = \frac{|\ker(p|_{(t^{-1}Kt) \cap G_Y})| \cdot |p((t^{-1}Kt) \cap G_Y)|}{|K| \cdot |G_Y|} = \frac{|(t^{-1}Kt) \cap G_Y|}{|K| \cdot |G_Y|} > 0.$$

Therefore, since $\langle \chi'|_{J_t}, \mathbb{1}_{J_t} \rangle_{J_t} \geq 0$ for every $t \in G$,

$$\langle \mathbf{R}_H^G(\chi)|_K, \mathbb{1}_K \rangle_K = 0 \iff \forall t \in G : \langle \chi|_{J_t}, \mathbb{1}_{J_t} \rangle_{J_t} = 0.$$

□

Lemma 2.12. *Continuing with Notation 2.7, let G be a finite group G acting on some set X , let $Y \subseteq X$ be some subset, and let $\chi \in \text{Ch}(H)$. Assume:*

1. $Y \supseteq Y_0 = \{a_1 \dots a_\ell\}$
2. χ marks every subgroup of H that is not (a_1, \dots, a_ℓ) - H -transitive.

Let $K \leq G$ such that $R_H^G(\chi)|_K \not\supseteq \mathbb{1}_K$. Then for every $t \in G$, $K \cap G_{tY}$ is (ta_1, \dots, ta_ℓ) - G_{tY} -transitive.

Proof. Assume that $R_H^G(\chi)|_K \not\supseteq \mathbb{1}_K$. By Lemma 2.11, for every $t \in G$, $J_t = p_t(K \cap G_{tY})$ satisfies $\chi_t \not\supseteq \mathbb{1}_{J_t}$.

Let $t \in G$. It is easy to see that for every $g \in H$, $f_t(g)tx = tgx$. For every $h \in H_t$ there exists some $g \in H$ such that $f_t(g) = h$. Since $\chi|_{J_e} \not\supseteq \mathbb{1}_{J_e}$, by assumption 2 of the lemma, J_e is (a_1, \dots, a_ℓ) - H -transitive. As such, there exists some $g' \in J_e$ such that $ga_i = g'a_i$ for every $1 \leq i \leq \ell$.

If we denote $h' := f_t(g')$, we get

$$h'(ta_i) = f_t(g')t(a_i) = tg'(a_i) = tg(a_i) = f_t(g)t(a_i) = h(ta_i)$$

for every $1 \leq i \leq \ell$. As such, J_t is (ta_1, \dots, ta_ℓ) - H_t -transitive. We pull it back by p_t (since p_t is the quotient by the kernel of the action) to show that $K \cap G_{tY}$ is (ta_1, \dots, ta_ℓ) - G_{tY} -transitive. \square

Proof of Lemma 2.8. Let $t_0 \in G$, and let $b_i := t_0 a_i$ for all $1 \leq i \leq \ell - 1$. Define $g_i \in K$ by induction, for $0 \leq i \leq \ell - 1$, so that $\forall j \leq i: g_i(a_j) = b_j$.

Base case – $i = 0$: define $g_0 = id$.

Induction step: assume that $g_j \in K$ are defined for all $j < i$. We define the following elements of G :

t' By assumption 3 of Lemma 2.8, with $i + 1 \leq \ell$, there exists some $t' \in G$ such that

$$g_{i-1}(a_1) = b_1, \dots, g_{i-1}(a_{i-1}) = b_{i-1}, g_{i-1}(a_i), b_i$$

all belong to $t'Y$.

s_1 Since $t'^{-1}g_{i-1}(a_j) \in Y$ for all $j \leq i$, assumption 4 of Lemma 2.8 implies that there exists some $s_1 \in G_Y$, such that $s_1(a_j) = t'^{-1}g_{i-1}(a_j)$ for all $j \leq i$. Equivalently, $t's_1(a_j) = g_{i-1}(a_j)$, for all $j \leq i$.

t Denote $t := t's_1$. Since $tY = t'Y$, $t^{-1}t'Y = Y$.

s_2 By assumption 4 of Lemma 2.8 with $t^{-1}(b_j) \in Y$ for $j = 1, \dots, i$, we obtain some $s_2 \in G_Y$, such that $s_2(a_j) = t^{-1}(b_j)$ for all $j \leq i$.

s_3 Denote $s_3 := ts_2t^{-1} \in G_{tY}$. For every $j \leq i$,

$$s_3(g_{i-1}(a_j)) = ts_2t^{-1}t(a_j) = ts_2(a_j) = b_j.$$

h Since, by Lemma 2.12, $K \cap G_{tY}$ is (ta_1, \dots, ta_ℓ) - G_{tY} -transitive and $t(a_j) = g_{i-1}(a_j)$ for all $j \leq i$, there exists some $h \in K$ such that for every $j \leq i$, $h(g_{i-1}(a_j)) = s_3(g_{i-1}(a_j)) = b_j$.

We can now define $g_i := hg_{i-1} \in K$. Then: $\forall j \leq i: g_i(a_j) = b_j$. By letting $g := g_{\ell-1} \in K$, we deduce that $g(a_i) = b_i$ for all $1 \leq i \leq \ell - 1$. \square

3 Characters of $GL_n(\mathbb{F}_q)$

This section introduces several facts about the characters of $GL_n(\mathbb{F}_q)$, and uses them to define and understand the main characters we need in the paper. We follow the terminology of [AT17].

Definition 3.1. Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_s)$, such that $\alpha_1, \dots, \alpha_s \geq 1$ and $\sum_{i=1}^s \alpha_i = n$. Define the parabolic subgroup

$$P_\alpha := \begin{bmatrix} GL_{\alpha_1}(\mathbb{F}_q) & * & \dots & * \\ 0 & GL_{\alpha_2}(\mathbb{F}_q) & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & GL_{\alpha_s}(\mathbb{F}_q) \end{bmatrix} \leq GL_n(\mathbb{F}_q).$$

Let $\chi_i \in \text{Ch}(GL_{\alpha_i}(\mathbb{F}_q))$. Define the characters

$$(\chi_1 \times \chi_2 \times \dots \times \chi_s) \left(\begin{bmatrix} A_1 & * & \dots & * \\ 0 & A_2 & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & A_s \end{bmatrix} \right) := \prod_{i=1}^s (\chi_i(A_i)),$$

$$\chi_1 \circ \chi_2 \circ \dots \circ \chi_s := \text{Ind}_{P_\alpha}^{GL_n(\mathbb{F}_q)} (\chi_1 \times \chi_2 \times \dots \times \chi_s) \in \text{Ch}(GL_n(\mathbb{F}_q)).$$

$$\chi_1 \circ \chi_2 \circ \dots \circ \chi_s$$

Proposition 3.2 ([Gre55, Lemma 2.5]). *The operator \circ is bilinear, associative and commutative.*

Remark Let $1 \leq \ell \leq n$, and let q be a prime power. Then, in the notation of Section 2, for every character χ of $\mathbb{P}GL_\ell(\mathbb{F}_q)$ we have $\text{R}_{\mathbb{P}GL_\ell(\mathbb{F}_q)}^{GL_n(\mathbb{F}_q)}(\chi) = \chi' \circ \mathbb{1}_{n-\ell}$ where χ' is the pull-back of χ to $GL_\ell(\mathbb{F}_q)$.

Using the operator \circ , we can define a family of characters.

Definition 3.3. Let $\lambda = (\lambda_1, \dots, \lambda_\ell) \vdash n$ be a partition of n . Define

$$\mathbb{1}_\lambda := \mathbb{1}_{\lambda_1} \circ \dots \circ \mathbb{1}_{\lambda_\ell} \in \text{Ch}(GL_n(\mathbb{F}_q)).$$

$$\mathbb{1}_\lambda$$

This family of characters is related to the action of $GL_n(\mathbb{F}_q)$ on \mathbb{F}_q^n . This can be seen as follows.

For a sequence of integers $0 = d_0 \leq d_1 \leq \dots \leq d_\ell = n$, we say that the sequence of subspaces $0 = V_0 \subseteq V_1 \subseteq \dots \subseteq V_\ell = \mathbb{F}_q^n$ is a *flag of signature* (d_1, \dots, d_ℓ) if $\dim(V_i) = d_i$ for all $i \leq \ell$.

flag of signature

For a partition $\lambda = (\lambda_1, \dots, \lambda_\ell) \vdash n$ we associate the increasing sequence $d_i(\lambda) := \sum_{j=1}^i \lambda_j$.

(d_1, \dots, d_ℓ)

Example 3.4. We present examples for the characters $\mathbb{1}_\lambda$. Since \circ is commutative, we may permute the entries of λ . For example, $\mathbb{1}_{(1, n-1)} = \mathbb{1}_{(n-1, 1)}$.

1. The character $\mathbb{1}_{(n)}$ is simply the trivial character, since the only flag of signature (n) is $0 \subseteq \mathbb{F}_q^n$.
2. The value $\mathbb{1}_{(1, n-1)}(g)$ is the number of 1-dimensional subspaces of \mathbb{F}_q^n that g fixes, or equivalently, the number of fixed points of g in the action of $GL_n(\mathbb{F}_q)$ on $\mathbb{P}^{n-1}\mathbb{F}_q$.
3. The value $\mathbb{1}_{(1, 1, n-2)}(g)$ is the number of flags $0 \subseteq V_1 \subseteq V_2 \subseteq \mathbb{F}_q^n$ that g fixes, where $\dim(V_1) = 1$ and $\dim(V_2) = 2$.

4. By [Remark 2.5](#), $\mathbb{1}_\lambda$ is the character of the permutation representation associated with the action of $GL_n(\mathbb{F}_q)$ on flags of signature $(d_1(\lambda), \dots, d_\ell(\lambda))$. In other words, $\mathbb{1}_\lambda(g)$ is the number of flags $0 = V_0 \subseteq V_1 \subseteq \dots \subseteq V_\ell = \mathbb{F}_q^n$ of signature $(d_1(\lambda), \dots, d_\ell(\lambda))$ such that $gV_i = V_i$ for all $i \leq \ell$.

The characters $\mathbb{1}_\lambda$ are not irreducible in general. The following lemma decomposes them to irreducible characters.

Lemma 3.5 ([\[AT17, Lemma 2.4\]](#)). *Let $\lambda = (\lambda_1, \dots, \lambda_\ell) \vdash n$. Then*

$$\mathbb{1}_\lambda = \sum_{\mu \vdash n} K_{\mu\lambda} \chi^\mu,$$

where $K_{\mu\lambda}$ are the Kostka numbers, and χ^μ are some irreducible characters⁸.

We define the unipotent characters $\{\chi^\mu: \mu \vdash n\}$ as these irreducible summands of χ^μ $\{\mathbb{1}_\lambda: \lambda \vdash n\}$.

In the remainder of the section, we study the characters θ_i of $GL_n(\mathbb{F}_q)$, defined below. We decompose them to irreducible characters, and characterize the properties of subgroups of $GL_n(\mathbb{F}_q)$ that they mark.

Definition 3.6. Define the following characters:

$$\theta_1 := \text{Ind}_{P_{(n-1,1)}}^{GL_n(\mathbb{F}_q)}(\mathbb{1}_{P_{(n-1,1)}}) - \mathbb{1}_{GL_n(\mathbb{F}_q)} = \mathbb{1}_{(n-1,1)} - \mathbb{1}_{(n)} \quad (n \geq 2) \quad \theta_1$$

$$\theta_2 := \text{Ind}_{P_{(n-2,1,1)}}^{GL_n(\mathbb{F}_q)}(\mathbb{1}_{P_{(n-2,1,1)}}) - \mathbb{1}_{GL_n(\mathbb{F}_q)} = \mathbb{1}_{(n-2,1,1)} - \mathbb{1}_{(n)} \quad (n \geq 3) \quad \theta_2$$

Using [Lemma 3.5](#), we deduce:

Corollary 3.7. *For the values of n where the characters are defined:*

1. $\mathbb{1}_{GL_n(\mathbb{F}_q)} = \mathbb{1}_{(n)} = \chi^{(n)}$
2. $\theta_1 = \chi^{(n-1,1)}$
3. $\theta_2 = 2\chi^{(n-1,1)} + \chi^{(n-2,1,1)} + \chi^{(n-2,2)}$

In 3, the term $\chi^{(n-2,2)}$ is omitted if $n = 3$.

Proposition 3.8. *Let $K \leq GL_n(\mathbb{F}_q)$, and let $V \cong \mathbb{F}_q^n$ be the vector space on which $GL_n(\mathbb{F}_q)$ acts. Consider the action of $GL_n(\mathbb{F}_q)$ on $\mathbb{P}(V)$, and denote by $\Phi(V)$ the collection of flags of signature $(1, 2, n)$, that is,*

$$\Phi(V) := \{(0, V_1, V_2, V) : 0 \subseteq V_1 \subseteq V_2 \subseteq V, \dim(V_1) = 1, \dim(V_2) = 2\}. \quad \Phi(V)$$

Then:

1. $\theta_1 \not\geq \mathbb{1}_K \iff K$ acts transitively on $\mathbb{P}(V)$
2. $\theta_2 \not\geq \mathbb{1}_K \iff K$ acts transitively on $\Phi(V) \iff \forall (u, v), (u', v') \in V \times V$, both pairs linearly independent, there exist $A \in K$ and elements of $\mathbb{F}_q - \alpha \neq 0, \beta, \gamma \neq 0$ - such that $Au = \alpha u', Av = \beta u' + \gamma v'$.

Proof. The proposition follows from [Corollary 2.6](#) and [Example 3.4](#). □

⁸In [\[AT17\]](#), these characters are labeled $\chi^{\mu^{(1)}}$. We define: $\chi^\mu = \chi^{\mu^{(1)}}$, which is more similar to the notation of [\[Gre55\]](#).

4 The Maximal Subgroups of $GL_n(\mathbb{F}_q)$

In this section we study the maximal subgroups of $GL_n(\mathbb{F}_q)$ that do not contain $SL_n(\mathbb{F}_q)$, and find a character χ such that $\chi|_K \supseteq \mathbb{1}_K$ for every maximal subgroup K .

It is enough to study maximal subgroups of $GL_n(\mathbb{F}_q)$, as $\chi|_K \supseteq \mathbb{1}_K$ implies $\chi|_{K'} \supseteq \mathbb{1}_{K'}$ for every $K' \leq K$. This can be easily seen, as $\chi|_K \supseteq \mathbb{1}_K$ implies that $\chi|_K = \mathbb{1}_K + \chi'$ for some character χ' of K , which implies that $\chi|_{K'} = \mathbb{1}_{K'} + \chi'|_{K'}$.

[Section 4.1](#) describes the geometric maximal subgroups, [Section 4.2](#) checks whether $\theta_i|_K \supseteq \mathbb{1}_K$ for K which is one of the geometric subgroups, and [Section 4.3](#) deals with the rest of the maximal subgroups, for the case of $n = 3$. For the main results, we only need these results for $n = 2$ and $n = 3$, but we add the general case since it is very similar to our specialized case.

Before we proceed to describing the geometric groups, we make several definitions.

Let G be a finite group, and let $d \in \mathbb{Z}_{\geq 1}$. We denote by dG some group extension of G with $[dG: G] = d$. There may exist many non-isomorphic groups that satisfy this condition, but we only consider properties that are common to all of the extensions.

For $K \leq GL_m(\mathbb{F}_q)$, the wreath product $K \wr S_\ell$, can be realized the group of matrices with $\ell \times \ell$ blocks, where:

1. Each block is of size $m \times m$.
2. The non-zero blocks are elements of K .
3. There is exactly one non-zero block in every row.
4. There is exactly one non-zero block in every column.

We denote by $\text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ the Galois group of the extension $\mathbb{F}_q/\mathbb{F}_{q_0}$ where q is a power of q_0 .

We use the nonstandard notation of $\Gamma L_d\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ to denote the group of semi-linear morphisms on \mathbb{F}_q^d over \mathbb{F}_{q_0} , that is, morphisms $f: \mathbb{F}_q^d \rightarrow \mathbb{F}_q^d$ that satisfy:

1. For every two vectors $u, v \in \mathbb{F}_q^d$, $f(u + v) = f(u) + f(v)$.
2. There exists some automorphism $\sigma \in \text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ such that for every $v \in \mathbb{F}_q^d$, $f(\lambda v) = \sigma(\lambda)f(v)$.

The group $\Gamma L_d\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ can be realized as the semidirect product $GL_d(\mathbb{F}_q) \rtimes \text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$, which is the set

$$\{(A, \sigma): A \in GL_d(\mathbb{F}_q), \sigma \in \text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)\}.$$

with multiplication defined by $(A, \sigma)(A', \sigma') = (A\sigma(A'), \sigma\sigma')$ where $\text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ acts on every matrix entry by entry. The action of $\Gamma L_d\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ on \mathbb{F}_q^d is defined by $(A, \sigma)(x) = A\sigma(x)$ where $\text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$ acts on every vector coordinate by coordinate.

When q_0 is prime, we write $\Gamma L_d(\mathbb{F}_q) := \Gamma L_d\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$. Similarly, we denote $\text{Aut}(\mathbb{F}_q) := \text{Gal}\left(\mathbb{F}_q/\mathbb{F}_{q_0}\right)$.

Table 1: The families $\mathcal{C}_i(GL_n(\mathbb{F}_q))$

\mathcal{C}_i	Type	Description	Conditions
\mathcal{C}_1	$P_{(m, n-m)}$	Stabilizer of $W \subseteq V$ with $\dim(W) = m$	$1 \leq m \leq n - 1$
\mathcal{C}_2	$GL_m(\mathbb{F}_q) \wr S_t$	Permutations of decompositions of V to t m -dimensional spaces	$n = mt$ $t \geq 2$
\mathcal{C}_3	$\Gamma L_m(\mathbb{F}_{q^r}/\mathbb{F}_q)$	Stabilizers of the structure of V as a vector space over \mathbb{F}_{q^r}	$n = mr$
\mathcal{C}_4	$GL_{n_1}(\mathbb{F}_q) \otimes GL_{n_2}(\mathbb{F}_q)$	Stabilizers of the structure of V as $V \cong V_1 \otimes V_2$	$n = n_1 n_2$ $2 \leq n_1 < n_2$
\mathcal{C}_5	$GL_n(\mathbb{F}_{q'})$	Matrices with coefficients in $\mathbb{F}_{q'}$	$\mathbb{F}_{q'} \subseteq \mathbb{F}_q$
\mathcal{C}_6	$N_{GL_n(\mathbb{F}_q)}(R)$	See Definition 4.2 .	r is prime $n = r^m, r q-1$ if $r = 2, 4 q-1$
\mathcal{C}_7	$GL_m(\mathbb{F}_q) \wr S_t$	Permutations of decompositions of $V = V_1 \otimes \cdots \otimes V_t$ where $\dim V_i = m$	$n = m^t$ $m \geq 3$
\mathcal{C}_8	$Sp_n(\mathbb{F}_q)$	See Definition 4.3 .	$4 \leq n$ is even
\mathcal{C}_8	$U_n(q^{\frac{1}{2}})$	See Definition 4.3 .	q is a square $n \geq 3$
\mathcal{C}_8	$O_n^\varepsilon(q)$	See Definition 4.3 .	q is odd $n \geq 3$

4.1 The Geometric Maximal Subgroups

Aschbacher's Theorem To understand the structure of subgroups of $GL_n(\mathbb{F}_q)$, we use a variation of a theorem due to Aschbacher⁹. Roughly speaking, given a finite simple classical group G_0 and an extension G of G_0 which satisfies $G_0 \leq G \leq \text{Aut}(G_0)$, Aschbacher's Theorem describes classes of subgroups $\mathcal{C}_1(G), \dots, \mathcal{C}_8(G)$ called "geometric classes" and an exceptional class $\mathcal{S}(G)$. The theorem states that every subgroup of G which does not contain G_0 and is maximal among all such subgroups is a member of one of the classes $\mathcal{C}_1(G), \dots, \mathcal{C}_8(G), \mathcal{S}(G)$.

The theorem can be generalized to several other families of groups, such as $GL_n(\mathbb{F}_q)$. To simplify the definitions and theorems we do not state the general theorem, but restrict to the case of $G = GL_n(\mathbb{F}_q)$. We use [KL90, Theorem 1.2.1] and the description of the geometric maximal subgroups in [KL90, Chapter 4] for the case of $GL_n(\mathbb{F}_q)$. While the classes are defined by their geometric meaning, we divide them further by their structure. We present the classes in the following theorem, followed by brief descriptions of the classes.

We note that while these classes may intersect and may contain subgroups that are not maximal among the subgroups of G not containing G_0 .

Theorem 4.1. *Let $G = GL_n(\mathbb{F}_q)$, and let $V \cong \mathbb{F}_q^n$ be the space upon which G acts with the usual action. Consider the families $\mathcal{C}_i(GL_n(\mathbb{F}_q))$ given in [Table 1](#). Then, every $K \leq G$ such that $K \not\cong SL_n(\mathbb{F}_q)$ is either contained in a member of $\mathcal{C}_i(G)$ for some $1 \leq i \leq 8$, or $K \in \mathcal{S}(G)$.*

Remark For subgroups $SL_n(\mathbb{F}_q) \leq G' \leq GL_n(\mathbb{F}_q)$, the families $\mathcal{C}_i(G')$ are defined as $\mathcal{C}_i(G') = \{G' \cap C : C \in \mathcal{C}_i(GL_n(\mathbb{F}_q))\}$, and likewise for \mathcal{S} .

⁹See [Asc84] for Aschbacher's original paper, or [KL90] for a more detailed introduction.

Description of the geometric classes Let $G = GL_n(\mathbb{F}_q)$, and let $V \cong \mathbb{F}_q^n$ be the space upon which G acts with the usual action.

$\mathcal{C}_1(\mathbf{G})$ This class contains G_W , which are the stabilizers of subspaces $W \subseteq V$. Up to conjugation, these are the groups $P_{(m,n-m)}$ defined in [Definition 3.1](#), with $m = \dim(W)$.

$\mathcal{C}_2(\mathbf{G})$ Write $V = \bigoplus_{i=1}^t V_i$, where each V_i is of dimension m . Consider the subgroup of G that stabilizes each of the vector spaces. It can be extended by allowing permutations of the vector spaces, to get a group conjugate to $GL_m(\mathbb{F}_q) \wr S_t$.

$\mathcal{C}_3(\mathbf{G})$ If $n = rm$, we can consider V as a vector space of dimension m over \mathbb{F}_{q^r} . Then, the semi-linear group $\Gamma L_m\left(\frac{\mathbb{F}_{q^r}}{\mathbb{F}_q}\right)$ naturally acts linearly (over \mathbb{F}_q) on V and embeds as a proper¹⁰ subgroup of $GL_n(\mathbb{F}_q)$.

$\mathcal{C}_4(\mathbf{G})$ This class contains the stabilizers of decompositions of the form $V \cong V_1 \otimes V_2$.

$\mathcal{C}_5(\mathbf{G})$ Considering G as a group of matrices, this class contains the subgroups of matrices $GL_n(\mathbb{F}_{\mathbb{F}_{q'}})$, where we restrict the coefficients to some subfield $\mathbb{F}_{q'}$ of \mathbb{F}_q .

$\mathcal{C}_6(\mathbf{G})$ See [Definition 4.2](#).

$\mathcal{C}_7(\mathbf{G})$ If we write $V = \bigotimes_{i=1}^t V_i$, where each V_i is of dimension m , consider the subgroup of G that stabilizes each of the vector spaces. It can be extended by allowing permutations of the vector spaces, to get a group isomorphic to $GL_m(\mathbb{F}_q) \wr S_t$.

$\mathcal{C}_8(\mathbf{G})$ See [Definition 4.3](#).

The class $\mathcal{S}(\mathbf{G})$ We only describe the groups in this class for $n = 2, 3$, in [Section 6.2](#) and [Section 4.3](#). For more information about the general case, see [[KL90](#), Section 1.2].

Definition 4.2. (A full definition is given in [[KL90](#), Section 4.6]) We define the groups in $\mathcal{C}_6(GL_n(\mathbb{F}_q))$. Assume that $n = r^m$, for a prime r that does not divide q , and let

$$\mathbb{F}_q^\times \ni \omega = \begin{cases} \text{primitive } r\text{-root of unity} & \text{if } r \neq 2 \\ \text{primitive 4-root of unity} & \text{if } r = 2 \end{cases}.$$

For some $R_i \leq \langle \omega \rangle \wr S_r$, $1 \leq i \leq m$, such that R_i is not contained in the group of diagonal matrices (these are given explicitly in [[KL90](#), Section 4.6]), let $R' = R_1 \otimes R_2 \cdots \otimes R_m$ as in \mathcal{C}_7 . Let ωI be the diagonal matrix with ω on the diagonal, and let $R = \langle \omega I, R' \rangle$ with its action on $\mathbb{F}_q^r \otimes \mathbb{F}_q^r \otimes \cdots \otimes \mathbb{F}_q^r \cong \mathbb{F}_q^n$. We define $\mathcal{C}_6(GL_n(\mathbb{F}_q))$ to be the collection of subgroups of $GL_n(\mathbb{F}_q)$ that are conjugate to the normalizers of such groups R .

Definition 4.3. (A full definition is given in [[KL90](#), Section 2]) The class $\mathcal{C}_8(GL_n(\mathbb{F}_q))$ consists of the groups listed in the last three rows of [Table 1](#). Each group has some bilinear form, such that the group is the subgroup of $GL_n(\mathbb{F}_q)$ that preserves the bilinear form. We describe the bilinear form by some standard basis, defining the group up to conjugation. These are described in [Table 2](#).

4.2 The Restriction of the Characters to the Geometric Subgroups

Recall [Definition 3.6](#) where θ_1 and θ_2 were introduced. In this section, we prove that for every subgroup $K \leq GL_n(\mathbb{F}_q)$ which is a member of one of the geometric families, $\chi|_K \supseteq \mathbb{1}_K$ for at least one of the characters $\chi = \theta_i$. While θ_1 is a summand of θ_2 for $n \geq 3$, θ_2 is not defined for $n = 2$. Therefore, whenever possible, we prove the result for θ_1 .

Throughout the section, we heavily use [Proposition 3.8](#) without explicitly referring to it.

¹⁰A single exceptional case occurs when $G = GL_2(\mathbb{F}_2)$. Then, the subgroup $\Gamma L_1\left(\frac{\mathbb{F}_4}{\mathbb{F}_2}\right)$ is not proper, so we define $\mathcal{C}_3(G)$ as the singleton $\{GL_1(\mathbb{F}_4)\}$.

Table 2: The bilinear forms defining the subgroups in the class $\mathcal{C}_8(GL_n(\mathbb{F}_q))$

K	Basis type	$Basis$	$Relations$
$Sp_n(\mathbb{F}_q)$ $n = 2m$	Symplectic	e_1, \dots, e_m f_1, \dots, f_m	$\langle e_i, e_j \rangle = 0$ $\langle f_i, f_j \rangle = 0$ $\langle e_i, f_j \rangle = \delta_{ij}$
$U_n(q^{\frac{1}{2}})$ $n = 2m$	Unitary	e_1, \dots, e_m f_1, \dots, f_m	$\langle e_i, e_j \rangle = 0$ $\langle f_i, f_j \rangle = 0$ $\langle e_i, f_j \rangle = \delta_{ij}$
$U_n(q^{\frac{1}{2}})$ $n = 2m + 1$	Unitary	e_1, \dots, e_m f_1, \dots, f_m x	$\langle e_i, e_j \rangle = 0, \langle e_i, x \rangle = 0$ $\langle f_i, f_j \rangle = 0, \langle f_i, x \rangle = 0$ $\langle x, x \rangle = 1, \langle e_i, f_j \rangle = \delta_{ij}$
$O_n^+(q)$ $n = 2m$	Orthogonal	e_1, \dots, e_m f_1, \dots, f_m	$\langle e_i, e_i \rangle = 0$ $\langle f_i, f_i \rangle = 0$ $\langle e_i, f_j \rangle = \delta_{ij}$
$O_n^-(q)$ $n = 2m$	Orthogonal	e_1, \dots, e_{m-1} f_1, \dots, f_{m-1} x, y	$\langle e_i, e_i \rangle = 0, \langle e_i, x \rangle = 0, \langle e_i, y \rangle = 0$ $\langle f_i, f_i \rangle = 0, \langle f_i, x \rangle = 0, \langle f_i, y \rangle = 0$ $\langle e_i, f_j \rangle = \delta_{ij}, \langle x, x \rangle = 1$ $\langle x, y \rangle = 1, \langle y, y \rangle \neq 0$
$O_n^{\circ}(q)$ $n = 2m + 1$	Orthogonal	e_1, \dots, e_m f_1, \dots, f_m x	$\langle e_i, e_i \rangle = 0, \langle e_i, x \rangle = 0$ $\langle f_i, f_i \rangle = 0, \langle f_i, x \rangle = 0$ $\langle x, x \rangle \neq 0, \langle e_i, f_j \rangle = \delta_{ij}$

(θ_1) In this part of the section, we prove that $\theta_1|_K \supseteq \mathbb{1}_K$ if K is a member of one of the families $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_4, \mathcal{C}_5$ or \mathcal{C}_7 . It is sufficient to find $v, w \in \mathbb{P}(V)$ such that no $k \in K$ satisfies $kv = w$.

Proposition 4.4. *If $K \in \mathcal{C}_1(GL_n(\mathbb{F}_q))$ then $\theta_1|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_1(GL_n(\mathbb{F}_q))$. The group K stabilizes some proper non-zero subspace W . Choose $v \in \mathbb{P}W$ and $w \notin \mathbb{P}W$. Then, for every $k \in K$, $kv \neq w$. \square

Proposition 4.5. *If $K \in \mathcal{C}_2(GL_n(\mathbb{F}_q))$ then $\theta_1|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_2(GL_n(\mathbb{F}_q))$. Consider the basis $\{e_1, \dots, e_n\}$ of V such that the elements of K are realized as block matrices (with respect to the decomposition $V = \bigoplus_{i=1}^t V_i$) over this basis. As a matrix with this basis, the first column of every element of K is all zeros but a single block of length $m < n$. Therefore, there is no element $k \in K$ with $k\bar{e}_1 = \overline{\sum_{i=1}^n e_i}$. \square

Proposition 4.6. *If $K \in \mathcal{C}_4(GL_n(\mathbb{F}_q))$ then $\theta_1|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_4(GL_n(\mathbb{F}_q))$. Recall that $K \cong GL_{n_1}(\mathbb{F}_q) \otimes GL_{n_2}(\mathbb{F}_q)$, and consider the decomposition $V \cong V_1 \otimes V_2$. The action of K is defined as follows: $(A \otimes B)(u \otimes v) = Au \otimes Bv$. By denoting $\{e_1, e_2, \dots\}$ and $\{e'_1, e'_2, \dots\}$ as bases of V_1 and V_2 respectively, we see that no element in K satisfies: $(A \otimes B)(e_1 \otimes e'_1) = e_1 \otimes e'_1 + e_2 \otimes e'_2$. \square

Proposition 4.7. *If $K \in \mathcal{C}_5(GL_n(\mathbb{F}_q))$ then $\theta_1|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_5(GL_n(\mathbb{F}_q))$. Choose $v \in \mathbb{P}(\mathbb{F}_{q'}^n)$, $w \in \mathbb{P}(\mathbb{F}_{q'} \times (\mathbb{F}_q - \mathbb{F}_{q'})^{n-1})$. Then $kv \neq w$ for all $k \in K$. \square

Proposition 4.8. *If $K \in \mathcal{C}_7(GL_n(\mathbb{F}_q))$ then $\theta_1|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_7(GL_n(\mathbb{F}_q))$. Then, an element of K permutes the subspaces and acts on each of the subspaces in the decomposition: $V \cong V_1 \otimes \dots \otimes V_t$. Let $e_1^{(i)}, \dots, e_m^{(i)}$ the basis of V_i .

Then, for every $k \in K$ there exist $v^{(i)} = \sum_{j=1}^m \lambda_j^{(i)} e_j^{(i)} \in V_i$ such that

$$k \left(\overline{\bigotimes_{i=1}^t e_1^{(i)}} \right) = \overline{\bigotimes_{i=1}^t v^{(i)}} = \overline{\bigotimes_{i=1}^t \sum_{j=1}^m \lambda_j^{(i)} e_j^{(i)}} = \sum_{j_1, \dots, j_t=1}^m \left(\prod_{i=1}^t \lambda_{j_i}^{(i)} \right) \cdot \left(\overline{\bigotimes_{i=1}^t e_{j_i}^{(i)}} \right).$$

By comparing the coefficients, we see that no $k \in K$ can satisfy

$$k \left(\overline{\bigotimes_{i=1}^t e_1^{(i)}} \right) = \overline{\bigotimes_{i=1}^t e_1^{(i)}} + \overline{\bigotimes_{i=1}^t e_2^{(i)}},$$

since this implies that both $\prod_{i=1}^t \lambda_1^{(i)} = \prod_{i=1}^t \lambda_2^{(i)} \neq 0$ and $\lambda_1^{(1)} \prod_{i=2}^t \lambda_2^{(i)} = 0$, which leads to a contradiction. \square

(θ_2) In this part of the section, we prove that $\theta_2|_K \supseteq \mathbb{1}_K$, for K in one of the families $\mathcal{C}_3, \mathcal{C}_6, \mathcal{C}_8$. Since θ_2 is only defined for $n \geq 3$, we assume this inequality holds.

Proposition 4.9. *If $K \in \mathcal{C}_3(GL_n(\mathbb{F}_q))$ then $\theta_2|_K \supseteq \mathbb{1}_K$.*

Proof. Assume otherwise. Then, for every three vectors $u, v, w \in \mathbb{F}_q^n$ such that u, v and u, w are linearly independent, there exists some $k \in K$ and $\lambda, \mu, \nu \in \mathbb{F}_q$ with $\lambda, \nu \neq 0$ such that $ku = \lambda u$ and $kv = \mu u + \nu w$.

Let $\alpha \in \mathbb{F}_{q^r}$ be such that $\mathbb{F}_{q^r} = \mathbb{F}_q(\alpha)$, and let $\{b_1, \dots, b_m\}$ be a basis of $V \cong \mathbb{F}_{q^r}^m$. Define $e_{i+r(j-1)} = \alpha^{i-1} b_j$ for $i = 1, \dots, r$ and $j = 1, \dots, m$. Then $\{e_1, \dots, e_n\}$ is a basis for $V \cong \mathbb{F}_q^n$.

For every $k = (A, \sigma) \in K = \Gamma L_m(\mathbb{F}_{q^r}/\mathbb{F}_q)$ such that $ke_1 = \lambda e_1, \lambda \neq 0$, we have $ke_2 = k(\alpha e_1) = \sigma(\alpha) \lambda e_1 = \sigma(\alpha) \lambda b_1$.

Assume that $m \geq 2$. Choose $u = e_1, v = e_2$ and $w = e_{r+1}$. Then, there exists some $k \in K$ such that $ke_1 = \lambda e_1$ and $ke_2 = \mu e_1 + \nu e_{r+1} = \mu b_1 + \nu b_2$ for $\lambda, \mu, \nu \in \mathbb{F}_q$ with $\lambda, \nu \neq 0$. That implies $\sigma(\alpha) \lambda b_1 = \mu b_1 + \nu b_2$ which leads to a contradiction. Thus, we may assume that $m = 1$ and $r = n \geq 3$.

We note that $\sigma(\alpha)$ cannot have an irreducible polynomial of degree 2 over \mathbb{F}_q , since then α would have such a polynomial as well, which contradicts $r \geq 3$.

Choose $u = e_1, v = e_2$ and $w = \sigma(\alpha)^2 e_1$. Since $r \geq 3, \sigma(\alpha)^2 \notin \mathbb{F}_q$ (otherwise $\sigma(\alpha)$ would have an irreducible polynomial of degree 2 over \mathbb{F}_q), so u and w are linearly independent over \mathbb{F}_q . Then, there exists some $k \in K$ such that $ke_1 = \lambda e_1$ and $ke_2 = \mu e_1 + \nu \sigma(\alpha)^2 e_1$, for $\lambda, \mu, \nu \in \mathbb{F}_q$ with $\lambda, \nu \neq 0$. Then, by comparison of coefficients of e_1 in the equality $\sigma(\alpha) \lambda e_1 = ke_2 = \mu e_1 + \nu \sigma(\alpha)^2 e_1$ we deduce that $\sigma(\alpha)$ has an irreducible polynomial of degree 2 over \mathbb{F}_q , which leads to a contradiction. \square

Proposition 4.10. *If $K \in \mathcal{C}_6(GL_n(\mathbb{F}_q))$ then $\theta_2|_K \supseteq \mathbb{1}_K$.*

Proof. Let $K \in \mathcal{C}_6(GL_n(\mathbb{F}_q))$ be the normalizer of some R as in [Definition 4.2](#), and assume that $\theta_2|_K \not\supseteq \mathbb{1}_K$. Let $r_1 \in R_1$ be a non-diagonal element. Choose some $v_1 \in V_1$ which is not an eigenvalue of r_1 , let $v_2 = r_1(v_1)$, and complete these to a basis of V_1 . Let $B = \{b_1, b_2, \dots\}$ be a basis of $V_2 \otimes \dots \otimes V_m$. Let $u \in R$.

Let $k \in K$. Then k is in the normalizer of R . Thus, $kR = Rk$. By $\theta_2|_K \not\supseteq \mathbb{1}_K$, there exists $\lambda, \mu, \nu \in \mathbb{F}_q$ with $\lambda, \nu \neq 0$ such that:

$$\begin{aligned} k(v_1 \otimes b_1) &= \lambda v_1 \otimes b_1 \text{ and} \\ k(v_2 \otimes b_1) &= \mu v_1 \otimes b_1 + \nu(v_2 \otimes b_1 + u). \end{aligned}$$

Then,

$$\begin{aligned} Rk(v_1 \otimes b_1) &= \{\lambda h(v_1 \otimes b_1) : h \in R\}, \\ kR(v_1 \otimes b_1) &= \{kh(v_1 \otimes b_1) : h \in R\}, \\ Rk(v_2 \otimes b_1) &= \{\mu h(v_1 \otimes b_1) + \nu h(v_2 \otimes b_1) + \nu h(u) : h \in R\} \text{ and} \\ kR(v_2 \otimes b_1) &= \{kh(v_2 \otimes b_1) : h \in R\} = \{kh(v_1 \otimes b_1) : h \in R\} \end{aligned}$$

with the last equality calculated by variable change, with $h' = h(r_1 \otimes I \otimes \cdots \otimes I)^{-1}$.

Comparing the various lines, we deduce

$$\{\lambda h(v_1 \otimes b_1): h \in R\} = \{\mu h(v_1 \otimes b_1) + \nu h(v_2 \otimes b_1) + \nu h(u): h \in R\}.$$

In particular, there exists some $h_0 \in R$ such that

$$\lambda h_0(v_1 \otimes b_1) = \mu v_1 \otimes b_1 + \nu v_2 \otimes b_1 + \nu u.$$

Denote $h_0 = \delta A \otimes C$ where $\delta \in \langle \omega \rangle$, $A \in R_1$ and $C \in R_2 \otimes \cdots \otimes R_m$. The coefficients of $v_2 \otimes b_1$ satisfy

$$\lambda \delta A_{2,1} C_{1,1} = \nu \implies A_{2,1} \neq 0.$$

Since $R_i \leq \langle \omega \rangle \wr S_r$, for every $s = 1$ or $3 \leq s \leq r$, $A_{s,1} = 0$. By comparing the coefficients of $v_1 \otimes b_1$, we get $\mu = \lambda \delta A_{1,1} C_{1,1} = 0$.

We get $h_0(v_1 \otimes b_1) = \delta A_{2,1} v_2 \otimes C b_1$. To finish, it is sufficient to choose (if $r = 2$, $2^m = n \geq 3 \implies m \geq 2$)

$$u = \begin{cases} v_3 \otimes b_1 & r \geq 3 \\ v_1 \otimes b_2, & r = 2, m \geq 2 \end{cases}$$

We get a contradiction to $\nu u = \lambda \delta A_{2,1} v_2 \otimes C b_1 - \nu v_2 \otimes b_1$. \square

Proposition 4.11. *If $K \in \mathcal{C}_8(GL_n(\mathbb{F}_q))$ then $\theta_2|_K \supseteq \mathbb{1}_K$.*

Proof. It is enough to find vectors u, v, w such that $\langle u, u \rangle = \langle u, v \rangle = 0$ and $\langle u, w \rangle \neq 0$. With such vectors, for every $\lambda, \mu, \nu \in \mathbb{F}_q$ with $\lambda, \nu \neq 0$,

$$\langle \lambda u, \mu u + \nu w \rangle = \lambda \nu \langle u, w \rangle \neq 0 = \langle u, v \rangle,$$

so there is no $k \in K$ such that $ku = \lambda u, kv = \mu u + \nu w$.

We define the vectors u, v, w as $u = e_1, w = f_1$ and

$$v = \begin{cases} f_2 & K = Sp_n(\mathbb{F}_q), U_{2m}(q^{\frac{1}{2}}) \text{ or } O_n^+(q) \\ x & K = U_{2m+1}(q^{\frac{1}{2}}), O_n^-(q) \text{ or } O_n^\circ(q) \end{cases}.$$

\square

From propositions 4.4-4.11, we conclude:

Corollary 4.12. *If $n \geq 3$, θ_2 marks every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$ and does not belong to $\mathcal{S}(GL_2(\mathbb{F}_q))$.*

For $n = 2$, $\mathcal{C}_8(GL_2(\mathbb{F}_q))$ is empty. Therefore, we can deduce:

Corollary 4.13. *The character θ_1 marks every subgroup of $GL_2(\mathbb{F}_q)$ that does not contain $SL_2(\mathbb{F}_q)$, and does not belong to $(\mathcal{C}_3 \cup \mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$.*

4.3 The Non-Geometric Subgroups for $n = 3$

Theorem 4.14 ([BHRD13, Table 8.4]). *$\mathcal{S}(SL_3(\mathbb{F}_q))$ are given in Table 3.*

By looking at the table, we see that there are finitely many possibilities for these groups. To understand the transitivity of their actions, we use the following immediate consequence of the orbit-stabilizer theorem.

Let G be a finite group, acting transitively on some set X . Let K be the kernel of the action, and let $x \in X$. Then

$$[G: K] = [G: G_x] [G_x: K] = |X| [G_x: K]. \quad (4.1)$$

In particular, $|X|$ divides $[G: K]$ and $|G|$.

Table 3: $\mathcal{S}(SL_3(\mathbb{F}_q))$

Case	$K' \leq SL_3(\mathbb{F}_q)$	Conditions on q	Order
(a)	$d \mathbb{P}SL_2(\mathbb{F}_7)$ $d = \gcd(q-1, 3)$	q is prime $q \equiv 1, 2, 4 \pmod{7}$ $q \neq 2$	$168d$
(b)	$3A_6$	q is prime $q \equiv 1, 4 \pmod{15}$	1080
(c)	$3A_6$	q is the square of a prime p $p \equiv 2, 3 \pmod{5}$ $p \neq 3$	1080

Recall that $\mathcal{S}(SL_n(\mathbb{F}_q)) = \{C \cap SL_n(\mathbb{F}_q) : C \in \mathcal{S}(GL_n(\mathbb{F}_q))\}$. We use (4.1) in the following lemma.

Lemma 4.15. $\forall K \in \mathcal{S}(GL_3(\mathbb{F}_q)) : \theta_2|_K \supseteq \mathbb{1}_K$.

Proof. By Theorem 4.14, the maximal subgroups of $SL_3(\mathbb{F}_q)$ have orders bounded by 1080.

Assume $\theta_2|_K \not\supseteq \mathbb{1}_K$. Then, K acts transitively on $X = \Phi(V)$, a set of size $(q^2+q+1)(q+1)$. By (4.1), $(q^2+q+1)(q+1) \mid |K|$.

Denote $K' := K \cap SL_3(\mathbb{F}_q) \in \mathcal{S}(SL_n(\mathbb{F}_q))$. By the second isomorphism theorem,

$$[K : K'] = [SL_3(\mathbb{F}_q)K : SL_3(\mathbb{F}_q)] \mid [GL_3(\mathbb{F}_q) : SL_3(\mathbb{F}_q)] = q - 1.$$

Therefore, $\frac{|K|}{c} \mid |K'|$, with $c = \gcd(|K|, q-1)$. Then, $\frac{(q^2+q+1)(q+1)}{c'} \mid |K'|$, with $c' = \gcd(|K|, q-1, (q^2+q+1)(q+1))$. This implies that $\frac{(q^2+q+1)(q+1)}{q-1} \leq |K'| \leq 1080$. Since the rational function is increasing with q , $q < 32$.

Then, it is easy to verify that for none of the possibilities for $|K'|$, $\frac{(q^2+q+1)(q+1)}{c'} \mid |K'|$, except for the case where $q = 2$. However, $q = 2$ is explicitly ruled out by the conditions for K . We deduce that $\theta_2|_K \supseteq \mathbb{1}_K$, for $K \in \mathcal{S}(GL_3(\mathbb{F}_q))$, for every prime power q . \square

We conclude:

Corollary 4.16. *The character θ_2 marks every subgroup of $GL_3(\mathbb{F}_q)$ that does not contain $SL_3(\mathbb{F}_q)$.*

5 Proofs of the Main Theorems for $n \geq 3$

In this section, we use the results from the previous sections to deduce the main theorems for $n \geq 3$. Since our proof of Theorem 1.6 for $GL_n(\mathbb{F}_q)$ uses results about $\mathbb{P}GL_3(\mathbb{F}_q)$, we prove that Theorem 1.6 for $GL_n(\mathbb{F}_q)$ implies Theorem 1.6 for $\mathbb{P}GL_n(\mathbb{F}_q)$ as well as Theorem 1.7, and only then prove Theorem 1.6 for $GL_n(\mathbb{F}_q)$.

5.1 The Special Linear Group

In this section, we assume Theorem 1.6 for $GL_n(\mathbb{F}_q)$, and prove Theorem 1.7 for $SL_n(\mathbb{F}_q)$.

Theorem 5.1 (e.g. [Leh73, Corollary 4.12]). *Let G be a finite group, let $K \trianglelefteq G$ and let $\chi \in \widehat{G}$. Assume that $G = KC$, where $C \leq G$ is cyclic and $K \cap C = 1$. Then, if $\chi|_K$ splits into b irreducible components, there are precisely $\frac{|C|}{b}$ distinct irreducible characters of G whose restriction to K equals $\chi|_K$.*

Remark By [Leh73, Proposition 4.11], the restricted components are all of the same degree.

Corollary 5.2. *Let $\chi^\mu \in \widehat{GL_n(\mathbb{F}_q)}$. Then, $\chi^\mu|_{SL_n(\mathbb{F}_q)} \in \widehat{SL_n(\mathbb{F}_q)}$.*

Proof. Let $\theta \in \widehat{\mathbb{F}_q^\times}$, and consider the character $\theta(\det(g))\chi^\mu(g) \in \widehat{GL_n(\mathbb{F}_q)}$. For every $g \in SL_n(\mathbb{F}_q)$, $\theta(\det(g))\chi^\mu(g) = \chi^\mu(g)$. Therefore $(\theta(\det)\chi^\mu)|_{SL_n(\mathbb{F}_q)} = \chi^\mu|_{SL_n(\mathbb{F}_q)}$.

Let C the cyclic group of diagonal matrices with 1 on the diagonal, except possibly in the top left cell. We have $|C| = q - 1$, $C \cap SL_n(\mathbb{F}_q) = 1$, $C SL_n(\mathbb{F}_q) = GL_n(\mathbb{F}_q)$ and C is a cyclic subgroup of $GL_n(\mathbb{F}_q)$.

It is easy to see that the characters $\theta(\det)\chi^\mu$ are all distinct by evaluating them on the elements of C (there exist characters χ of $GL_n(\mathbb{F}_q)$ for which $\theta(\det)\chi$ are not all distinct). As such, the restrictions of the $q - 1$ distinct irreducible characters $\theta(\det)\chi^\mu$ are equal to $\chi^\mu|_{SL_n(\mathbb{F}_q)}$. Therefore, by Theorem 5.1, $\chi^\mu|_{SL_n(\mathbb{F}_q)}$ splits into a single irreducible character, namely, it is irreducible itself. \square

By identifying $\chi^\mu := \chi^\mu|_{SL_n(\mathbb{F}_q)}$, using Corollary 5.2, we can deduce that the non-projective version of Theorem 1.6 implies the non-projective version of Theorem 1.7.

5.2 The Projective Groups

In this section, we assume the non-projective versions of the main theorems, and deduce the projective counterparts.

Lemma 5.3. *Let G and H be finite groups, with $p: G \twoheadrightarrow H$. Let $\chi \in \widehat{G}$, and let (π, V) be its corresponding representation. Define*

$$\ker(\chi) := \ker(\pi) = \{g \in G : \chi(g) = \chi(1)\}.$$

Assume that $\ker(\chi) \geq \ker(p)$. Then $\chi'(h) := \chi(p^{-1}(h))$ is well defined, and is an irreducible character of H .

Proof. Since $\ker(\pi) \geq \ker(p)$, there is a well defined representation π' of H with $\pi(g) = \pi'(p(g))$ for every $g \in G$. It is irreducible since $\text{Im}(\pi') = \text{Im}(\pi)$, which stabilizes no nontrivial subspace of V .

By taking the trace of the representations, with the trace of π' equal to χ' , the proposition follows. \square

Corollary 5.4. *Let $G' \leq G$ be groups, and let $\mathcal{X}(G) \subseteq \widehat{G}$ be a set that marks every subgroup of G that does not contain G' . Let $p: G \twoheadrightarrow H$, and assume that $\forall \chi \in \mathcal{X}(G) : \ker(\chi) \geq \ker(p)$. Define $H' := p(G')$ and*

$$\mathcal{X}(H) := \{g \mapsto \chi(p^{-1}(g)) : \chi \in \mathcal{X}(G)\} \subseteq \widehat{H}.$$

Then, $\mathcal{X}(H)$ marks every subgroup of H that does not contain H' .

Proof. Let $J \leq H$, such that $J \not\supseteq H'$, and let $K := p^{-1}(J)$.

Since K does not contain G' , there exists some $\chi \in \mathcal{X}(G)$ such that $\chi|_K \not\supseteq \mathbb{1}_K$. Denote by $\chi' \in \mathcal{X}(H)$ the character defined by $\chi'(g) = \chi(p^{-1}(g))$. We have

$$\langle \chi'|_J, \mathbb{1}_J \rangle_J = \frac{1}{|J|} \sum_{h \in J} \chi'(h) = \frac{1}{|J|} \frac{|J|}{|K|} \sum_{g \in K} \chi(g) = \langle \chi|_K, \mathbb{1}_K \rangle_K > 0$$

which implies that $\chi'|_J \not\supseteq \mathbb{1}_J$. \square

Lemma 3.5 implies that the kernels of the unipotent characters contain the center of $GL_n(\mathbb{F}_q)$, as they are spanned by the characters $\mathbb{1}_\lambda$, that do satisfy this. Thus, we may identify χ^μ with the corresponding characters as in Corollary 5.4.

Using Corollary 5.4, we deduce that the non-projective versions of Theorem 1.6 and Theorem 1.7 imply their projective counterparts.

5.3 $GL_n(\mathbb{F}_q)$

To prove the main theorems for $GL_n(\mathbb{F}_q)$, $n \geq 3$, we need to know the possibilities for subgroups of $GL_n(\mathbb{F}_q)$ that act 2-transitively on the projective space. The following theorem summarizes these possibilities:

Theorem 5.5 ([CK79]). *If $K \leq GL_n(\mathbb{F}_q)$, $n \geq 3$ and K is 2-transitive on $\mathbb{P}^{n-1}\mathbb{F}_q$, then either $K \geq SL_n(\mathbb{F}_q)$ or $K = A_7$, with $n = 4$ and $q = 2$.*

This theorem, [Corollary 4.16](#) and the results of [Section 5.2](#) show that θ_2 marks every subgroup of $PGL_3(\mathbb{F}_q)$ that is not $(\bar{e}_1, \bar{e}_2, \bar{e}_3)$ - $PGL_3(\mathbb{F}_q)$ -transitive, since $(\bar{e}_1, \bar{e}_2, \bar{e}_3)$ - $PGL_3(\mathbb{F}_q)$ -transitivity implies 2-transitivity.

Thus, using [Corollary 2.9](#), we see that $\forall n \geq 3$, $\theta_2 \circ \mathbb{1}_{(n-3)}$ marks every subgroup of $GL_n(\mathbb{F}_q)$ that is not (\bar{e}_1, \bar{e}_2) - $GL_n(\mathbb{F}_q)$ -transitive. Since $GL_n(\mathbb{F}_q)$ is 2-transitive with its projective action, we deduce that $(\theta_1 + \theta_2) \circ \mathbb{1}_{(n-3)}$ marks every subgroup of $GL_n(\mathbb{F}_q)$ that is not 2-transitive with the restricted action on $\mathbb{P}^{n-1}\mathbb{F}_q$.

Outside the exceptional case of the theorem, the subgroups of $GL_n(\mathbb{F}_q)$ that do not act 2-transitively on $\mathbb{P}^{n-1}\mathbb{F}_q$ are precisely those that do not contain $SL_n(\mathbb{F}_q)$, by [Theorem 5.5](#).

Thus, to find the irreducible characters mentioned in the main theorems (for all but $GL_4(\mathbb{F}_2)$), it remains to decompose $\theta_2 \circ \mathbb{1}_{(n-3)}$ to irreducible characters.

By the definition of θ_2 ,

$$\theta_2 \circ \mathbb{1}_{(n-3)} = (\mathbb{1}_{(1^3)} - \mathbb{1}_{(3)}) \circ \mathbb{1}_{(n-3)} = \mathbb{1}_{(n-3,1^3)} - \mathbb{1}_{(n-3,3)}.$$

These can be decomposed using [Lemma 3.5](#), to get:

$$\theta_2 \circ \mathbb{1}_{(n-3)} = \begin{cases} \chi^{(1^3)} + 2\chi^{(2,1)} & n = 3 \\ \chi^{(1^4)} + 3\chi^{(2,1^2)} + 2\chi^{(2^2)} + 2\chi^{(3,1)} & n = 4 \\ \chi^{(n-3,1^3)} + 2\chi^{(n-3,2,1)} + 3\chi^{(n-2,1^2)} + 2\chi^{(n-2,2)} + 2\chi^{(n-1,1)} & n \geq 5 \end{cases}$$

This concludes the proof of [Theorem 1.6](#).

5.4 $GL_4(\mathbb{F}_2)$

In this section, we study the exceptional case of $GL_4(\mathbb{F}_2)$. Since the underlying field is \mathbb{F}_2 , we have $GL_4(\mathbb{F}_2) = SL_4(\mathbb{F}_2) = PGL_4(\mathbb{F}_2) = PSL_4(\mathbb{F}_2)$.

The problem with this case is due to [Theorem 5.5](#). Note that $\theta_2 \circ \mathbb{1}_{(1)}$ marks every proper subgroup of $GL_4(\mathbb{F}_2)$ that is not the $A_7 \leq GL_4(\mathbb{F}_2)$ referenced in the theorem.

We wish to find a character $\psi \in \widehat{GL_4(\mathbb{F}_2)}$ such that $\psi|_{A_7} \supseteq \mathbb{1}_{A_7}$. For this goal, we use the following proposition to better understand the inclusion of A_7 in $GL_4(\mathbb{F}_2)$.

Proposition 5.6 (e.g. [DM96, Chapter 7.7]). *$GL_4(\mathbb{F}_2) \cong A_8$. Under this isomorphism, the 2-transitive A_7 is conjugate to the stabilizer of an element in the natural action of A_8 on 8 elements.*

Thus, we use the representation theory of A_8 . For that, we consider the representation theory for S_n (see, for instance, [Ful97]), and use [Theorem 5.1](#). This leads to the following proposition.

Proposition 5.7. *The characters of A_8 have degrees (with multiplicities):*

$$\{1, 7, 14, 20, 21, 21, 21, 28, 35, 45, 45, 56, 64, 70\}.$$

To prove the proposition, we use the following. See [Ful97] for information about standard tableaux.

Proposition 5.8 (e.g. [Ful97, Chapters 7.2 and 7.3]). *For every $n \in \mathbb{Z}_{\geq 1}$, $\widehat{S}_n = \{\chi^\lambda : \lambda \text{ is a partition of } n\}$, for some characters with $\dim(\chi^\lambda) = f^\lambda = \text{the number of standard tableaux of shape } \lambda$.*

In addition, $\text{sgn } \chi^\lambda = \chi^{\lambda'}$, with sgn being the sign representation.

Proof of Proposition 5.7. By substituting $n = 8$ and calculating the characters, the characters of S_n have degrees (with multiplicities)

$$\{1, 1, 7, 7, 14, 14, 20, 20, 21, 21, 28, 28, 35, 35, 42, 56, 56, 64, 64, 70, 70, 90\},$$

with every pair belonging to conjugate partitions.

By [Theorem 5.1](#), noting that $\chi^{\lambda'}|_{A_n} = (\text{sgn } \chi^\lambda)|_{A_n} = \chi^\lambda|_{A_n}$, we deduce that the characters of A_8 are precisely:

1. The restrictions of χ^λ for $\lambda \neq \lambda'$
2. Two summands of $\chi^\lambda|_{A_n}$ of equal dimensions for $\lambda = \lambda'$

Thus, we obtain the following degrees:

$$\{1, 7, 14, 20, 21, 21, 21, 28, 35, 45, 45, 56, 64, 70\}.$$

These are all of the characters of A_8 , as the sum of their squares is $|A_8|$. \square

The last thing we need to do to complete the proof of [Theorem 1.8](#) is understand the decomposition of $\text{Ind}_{A_7}^{GL_4(\mathbb{F}_2)}(\mathbb{1}_{A_7})$.

Proposition 5.9. $\text{Ind}_{A_7}^{GL_4(\mathbb{F}_2)}(\mathbb{1}_{A_7}) = \mathbb{1}_{GL_4(\mathbb{F}_2)} + \psi$, where $\psi \in \widehat{GL_4(\mathbb{F}_2)}$ is the single irreducible of dimension 7.

Proof. Let $\theta = \text{Ind}_{A_7}^{GL_4(\mathbb{F}_2)}(\mathbb{1}_{A_7})$. Observe that $\dim(\theta) = [A_8 : A_7] = 8$. By [Proposition 2.4](#), $\langle \theta, \mathbb{1}_{GL_4(\mathbb{F}_2)} \rangle = 1$. Thus, since $GL_4(\mathbb{F}_2) \cong A_8$ has a unique nontrivial character of degree at most 7, the claim follows. \square

Remark By calculating the degrees of the characters, we deduce that the character ψ is not a unipotent character.

We can now state the following proposition.

Proposition 5.10. The only nontrivial $\psi \in \widehat{GL_4(\mathbb{F}_2)}$ such that $\psi|_{A_7} \supseteq \mathbb{1}_{A_7}$ is the character of $GL_4(\mathbb{F}_2)$ of degree 7.

Proof. Let $\psi \in \widehat{GL_4(\mathbb{F}_2)}$ be the single irreducible character of dimension 7. By [Frobenius Reciprocity](#),

$$\langle \theta|_{A_7}, \mathbb{1}_{A_7} \rangle_{A_7} = \left\langle \theta, \text{Ind}_{A_7}^{GL_4(\mathbb{F}_2)}(\mathbb{1}_{A_7}) \right\rangle_{GL_4(\mathbb{F}_2)} = \langle \theta, \mathbb{1}_{GL_4(\mathbb{F}_2)} + \psi \rangle_{GL_4(\mathbb{F}_2)}.$$

So the only characters where the sum is non-zero are the trivial character and ψ . \square

This completes the proof of [Theorem 1.8](#)

6 The Main Theorems for $n = 2$

In this section, we prove the main theorems for $n = 2$. We find a character ψ_1 such that for every $K \in \mathcal{C}_3 \cup \mathcal{C}_6 \cup \mathcal{S}$, either $\psi_1|_K \supseteq \mathbb{1}_K$ or $\theta_1|_K \supseteq \mathbb{1}_K$. Then, by [Corollary 4.13](#), the non-projective case of [Theorem 1.9](#) follows.

The characters in the $n = 2$ case are slightly simpler than in the general case. We use the character tables in [Ada02], and refer the readers to them for definitions of the characters $\pi(\chi)$ and ω_0^+ . The two notations are related as follows, with $\alpha, \beta \in \widehat{\mathbb{F}_q^\times}$ and $\chi \in \widehat{\mathbb{F}_{q^2}^\times}$:

1. $\rho(\mu(\alpha, \beta)) = \alpha \circ \beta$
2. $\rho'(\alpha)$ is the linear character $\rho'(\alpha)(g) = \alpha(\det(g))$
3. $\bar{\rho}(\alpha) = \alpha \circ \alpha - \rho'(\alpha)$. In particular, $\bar{\rho}(\mathbb{1}_{\mathbb{F}_q^\times}) = \chi^{(1,1)} = \theta_1$.
4. $\pi(\chi)$ are additional characters of $GL_2(\mathbb{F}_q)$ called the cuspidal characters.

Definition 6.1. Define ψ_1 according to the following cases:

1. If $q = 3$ and G is $SL_2(\mathbb{F}_q)$ or $\mathbb{P}SL_2(\mathbb{F}_q)$, ψ_1 is ω_0^+ , as defined in the character table of these groups. ψ_1
2. Otherwise, if q is odd, ψ_1 is $\phi_1 := \pi(\phi)$, for $\phi \in \widehat{\mathbb{F}_{q^2}^\times}$, of order¹¹ $q + 1$ in the character ϕ_1 group.
3. If q is even and $q > 2$, ψ_1 is $\alpha \circ \bar{\alpha}$, where $\alpha \neq \mathbb{1}_{\mathbb{F}_q^\times}$.
4. If $q = 2$, ψ_1 is $\phi_1 := \pi(\phi)$, for any of the nontrivial $\phi \in \widehat{\mathbb{F}_4^\times}$. ϕ_1

By following the character tables, we see that the characters ϕ_1 and $\alpha \circ \bar{\alpha}$ do restrict irreducibly to the special groups, for $q \neq 3$. In the case of $q = 3$, ϕ_1 splits to two summands – one of which is the character ω_0^+ . In addition, the characters ω_0^+ , ϕ_1 and $\alpha \circ \bar{\alpha}$ project to the projective groups. Therefore, by claims similar to those made in Section 5.1 and Section 5.2, the theorems for the projective and the special groups follow from the case of $GL_2(\mathbb{F}_q)$.

6.1 $\mathcal{C}_3(GL_2(\mathbb{F}_q))$

The family $\mathcal{C}_3(GL_2(\mathbb{F}_q))$ is the family of stabilizers of the structure of \mathbb{F}_q^2 as a 1-dimensional vector space over \mathbb{F}_{q^2} , that is, elements that act on $\mathbb{F}_q^2 = \mathbb{F}_{q^2}$ as either multiplication by invertible elements of \mathbb{F}_{q^2} , or an automorphism of \mathbb{F}_{q^2} over \mathbb{F}_q . We use [Ada02] for the various character tables.

q is odd First, we handle the case when q is odd.

Since \mathbb{F}_q^\times is cyclic of even order, there exists some non square $\Delta \in \mathbb{F}_q^\times$. Therefore the group $\{x^2 : x \in \mathbb{F}_q^\times\}$ is a proper subgroup of \mathbb{F}_q^\times . This implies that $x^2 - \Delta$ is irreducible over \mathbb{F}_q .

Thus, we can choose the base $\{1, \delta\}$ for the field extension, where $\delta^2 = \Delta$. Since every $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$ respects the structure of \mathbb{F}_{q^2} , we deduce:

Proposition 6.2. *If q is odd and $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$, K is conjugate to*

$$\left\{ \begin{bmatrix} a & \Delta b \\ b & a \end{bmatrix}, \begin{bmatrix} a & -\Delta b \\ b & -a \end{bmatrix} \right\} \leq GL_2(\mathbb{F}_q).$$

We can conclude (since $Z(GL_2(\mathbb{F}_q)) \leq K$) that K has the following conjugacy classes (where a and b range over \mathbb{F}_q such that the matrices are invertible):

$$\left\{ \begin{bmatrix} a & \Delta b \\ b & a \end{bmatrix}, \begin{bmatrix} 0 & -\Delta b \\ b & 0 \end{bmatrix} c \text{ times}, \begin{bmatrix} a & 0 \\ 0 & -a \end{bmatrix} (q + 1 - c) \text{ times} \right\}.$$

Using the character tables, we can see that:

¹¹The order condition makes sure the character projects to the projective group

Proposition 6.3. *If q is odd and $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$, $\phi_1|_K \supseteq \mathbb{1}_K$.*

Proof.

$$\begin{aligned}
|K| \langle \phi_1|_K, \mathbb{1}_K \rangle_K &= (q-1) \sum_{x \in \mathbb{F}_q^\times} \phi(x) - \sum_{z \in \mathbb{F}_{q^2}^\times \setminus \mathbb{F}_q^\times} (\phi(z) + \phi(\bar{z})) \\
&\quad - c \sum_{x \in \mathbb{F}_q^\times} (\phi(\delta x) + \phi(-\delta x)) \\
&= (q+1) \sum_{x \in \mathbb{F}_q^\times} \phi(x) - 2 \sum_{z \in \mathbb{F}_{q^2}^\times} \phi(z) - 2c\phi(\delta) \sum_{x \in \mathbb{F}_q^\times} \phi(x) \\
&= (q+1+2c)(q-1)
\end{aligned}$$

Since $0 \leq c \leq q+1$, $|K| = 2(q^2-1)$, and $\langle \phi_1|_K, \mathbb{1}_K \rangle_K$ is an integer, we deduce that $c = \frac{q+1}{2}$. Thus, $\langle \phi_1|_K, \mathbb{1}_K \rangle_K = 1$. \square

This shows the desired result for the family $\mathcal{C}_3(GL_2(\mathbb{F}_q))$, for odd $q \neq 3$.

The case $q = 3$ While [Proposition 6.3](#) shows the desired result for $GL_2(\mathbb{F}_3)$, and projects to $PGL_2(\mathbb{F}_3)$, the special cases are slightly more complicated. In these cases, as can be seen in the character tables, ϕ_1 (both in the projective and non-projective cases) decomposes to two 1-dimensional characters called the oscillator characters (ω_o^\pm in [\[Ada02\]](#)). Either of these satisfies the desired result, and we arbitrarily choose ω_o^+ .

q is even Here, we handle the case when q is even.

If $q = q_0^2$, consider the trace morphism $\text{Tr}_{\mathbb{F}_q/\mathbb{F}_{q_0}}(x) = x^2 + x$, with its image \mathbb{F}_{q_0} . Choosing $\Delta \in \mathbb{F}_q \setminus \mathbb{F}_{q_0}$, the polynomial $x^2 + x - \Delta$ is irreducible over \mathbb{F}_q .

If $q \neq q_0^2$, \mathbb{F}_q is not an extension of \mathbb{F}_4 . As such, the polynomial $x^2 + x - 1$ is irreducible over \mathbb{F}_q . Choose $\Delta = 1$.

Thus, we can choose the basis $\{1, \delta\}$ for the field extension $\mathbb{F}_{q^2}/\mathbb{F}_q$, where $\delta^2 + \delta = \Delta$. Since every $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$ respects the structure of \mathbb{F}_{q^2} , we deduce:

Proposition 6.4. *If q is even and $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$, K is conjugate to*

$$\left\{ \begin{bmatrix} a & \Delta b \\ b & a+b \end{bmatrix}, \begin{bmatrix} a & a+\Delta b \\ b & a \end{bmatrix} \right\} \leq GL_2(\mathbb{F}_q).$$

Thus, we can conclude (since $Z(GL_2(\mathbb{F}_q)) \leq K$) that K has the following conjugacy classes (where a, b and x range over \mathbb{F}_q such that the matrices are invertible):

$$\left\{ \begin{bmatrix} a & \Delta b \\ b & a+b \end{bmatrix}, \begin{bmatrix} x & 1 \\ 0 & x \end{bmatrix} q+1 \text{ times} \right\}$$

Using the character tables, we can see that:

Proposition 6.5. *If $q \neq 2$ is even and $K \in \mathcal{C}_3(GL_2(\mathbb{F}_q))$, $\alpha \circ \bar{\alpha}|_K \supseteq \mathbb{1}_K$.*

Proof.

$$|K| \langle \alpha \circ \bar{\alpha}|_K, \mathbb{1}_K \rangle_K = 2(q+1) \sum_{x \in \mathbb{F}_q^\times} 1 = 2(q^2-1).$$

Dividing by $|K| = 2(q^2-1)$, we get $\langle \alpha \circ \bar{\alpha}|_K, \mathbb{1}_K \rangle_K = 1$. \square

By the character tables, if $\alpha \neq \mathbb{1}_{\mathbb{F}_q^\times}$, $\alpha \circ \bar{\alpha}$ is irreducible, and does restrict irreducibly to the special and projective cases. Such a choice exists for all even $q \neq 2$. This shows the desired result for the family $\mathcal{C}_3(GL_2(\mathbb{F}_q))$, for even $q \neq 2$.

Table 4: $(\mathcal{C}_6 \cup \mathcal{S})(SL_2(\mathbb{F}_q))$

Case	$K' \leq SL_2(\mathbb{F}_q)$	Conditions on q	Order
(a)	$2S_4$	q is prime $q \equiv \pm 1 \pmod{8}$	48
(b)	$2A_4$	q is prime $q \equiv \pm 3, 5, \pm 11, \pm 13, \pm 19 \pmod{40}$	24
(c)	$2A_5$	q is prime $q \equiv \pm 1 \pmod{10}$	120
(d)	$2A_5$	q is the square of a prime p $p \equiv \pm 3 \pmod{10}$	120

The case $q = 2$ In this case, a simple calculation shows that ψ_1 marks the single proper subgroup in $\mathcal{C}_3(GL_2(\mathbb{F}_2))$ which is of order 3.

6.2 $\mathcal{C}_6(GL_2(\mathbb{F}_q))$ and $\mathcal{S}(GL_2(\mathbb{F}_q))$

Theorem 6.6 ([BHRD13, Table 8.3]). $(\mathcal{C}_6 \cup \mathcal{S})(SL_2(\mathbb{F}_q))$ are given in Table 4.

We can see that the orders of these families are bounded. If $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$ does not act transitively on $\mathbb{P}^1\mathbb{F}_q$, then $\theta_1|_K \supseteq \mathbb{1}_K$, and our main claim follows.

As such, we use an argument about the transitivity of their action on $\mathbb{P}^1\mathbb{F}_q$ to bound q , namely:

Proposition 6.7. Assume $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$, and that it acts transitively on $\mathbb{P}^1\mathbb{F}_q$. Denote $K' := K \cap SL_2(\mathbb{F}_q)$. Then, $(q+1) \mid |K'|$.

Proof. By Table 4, we can see that q is odd. Let $J := K/Z(GL_2(\mathbb{F}_q))$ and $J' := K'/Z(SL_2(\mathbb{F}_q))$.

By (4.1), $(q+1) \mid |J|$. As $|K'|$ is given by Table 4, by the maximality of K' along with its order¹², $K' \geq Z(SL_2(\mathbb{F}_q))$. Thus, $|K'| = 2|J'|$. Then, lifting J' to $J \leq PGL_2(\mathbb{F}_q)$, we get either $|J| = |J'|$ or $|J| = 2|J'|$. In either case, $|J|$ divides $|K'|$. So we get that $(q+1) \mid |K'|$. \square

To ease the handling of cases (c) and (d), we use the following:

Proposition 6.8 ([Dic03, Sections 258-259]). $A_5 \leq PSL_2(\mathbb{F}_q)$ is, up to conjugation, generated by the following:

If $q \equiv \pm 1 \pmod{5}$

$$V_5 := \begin{bmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{bmatrix}, \text{ some } V_2 \text{ of order 2}$$

where ζ is a primitive root of unity of order 5.

If $q \equiv 0 \pmod{5}$

$$V_5 := \begin{bmatrix} 1 & \mu \\ 0 & 1 \end{bmatrix}, \text{ some } V_2 \text{ of order 2}$$

where $\mu \neq 0$.

Corollary 6.9. Assume $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$, K acts transitively on $\mathbb{P}^1\mathbb{F}_q$ and one of cases (c) or (d) holds. Then $(q+1) \mid 24$.

Proof. In these cases, the projective image of $K = 2A_5$ (which is the lifting of $A_5 \leq PSL_2(\mathbb{F}_q)$ to $PGL_2(\mathbb{F}_q)$) contains a subgroup of order 5 stabilizing \bar{e}_1 . As such, $5 \mid |K_{\bar{e}_1}|$

Since K acts transitively on $\mathbb{P}^1\mathbb{F}_q$, by (4.1), $(q+1) \mid |K_{\bar{e}_1}| = |K|$. Therefore, $5(q+1) \mid |2A_5| = 120$. This implies that $(q+1) \mid 24$. \square

¹²Otherwise, $K'Z(SL_2(\mathbb{F}_q)) = SL_2(\mathbb{F}_q)$. However, this implies that $2|K'| = |SL_2(\mathbb{F}_q)|$ (since q is odd, $|Z(SL_2(\mathbb{F}_q))| = 2$). None of the orders from Table 4 satisfies this condition.

Table 5: Generators and Characters

q	$K \cap SL_2(\mathbb{F}_q)$	Generators in $\mathbb{P}GL_2(\mathbb{F}_q)$		Character
5	$2A_4$	$\begin{bmatrix} 3 & 4 \\ 2 & 3 \end{bmatrix}$	$\begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix}$	ϕ_1
7	$2S_4$	$\begin{bmatrix} 2 & 6 \\ 5 & 5 \end{bmatrix}$	$\begin{bmatrix} 4 & 6 \\ 3 & 5 \end{bmatrix}$	ϕ_1
7	$2S_4$	$\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} 6 & 5 \\ 6 & 4 \end{bmatrix}$	ϕ_1
11	$2A_5$	$\begin{bmatrix} 8 & 7 \\ 6 & 4 \end{bmatrix}$	$\begin{bmatrix} 8 & 8 \\ 8 & 4 \end{bmatrix}$	ϕ_1
11	$2A_5$	$\begin{bmatrix} 3 & 9 \\ 5 & 8 \end{bmatrix}$	$\begin{bmatrix} 6 & 2 \\ 6 & 4 \end{bmatrix}$	ϕ_1
23	$2S_4$	$\begin{bmatrix} 8 & 2 \\ 18 & 18 \end{bmatrix}$	$\begin{bmatrix} 15 & 1 \\ 1 & 8 \end{bmatrix}$	ϕ_1
23	$2S_4$	$\begin{bmatrix} 1 & 15 \\ 17 & 8 \end{bmatrix}$	$\begin{bmatrix} 2 & 11 \\ 17 & 17 \end{bmatrix}$	ϕ_1
31	$2S_4$	$\begin{bmatrix} 11 & 19 \\ 7 & 4 \end{bmatrix}$	$\begin{bmatrix} 1 & 13 \\ 30 & 26 \end{bmatrix}$	θ_1
31	$2S_4$	$\begin{bmatrix} 5 & 25 \\ 18 & 6 \end{bmatrix}$	$\begin{bmatrix} 0 & 22 \\ 20 & 0 \end{bmatrix}$	θ_1
47	$2S_4$	$\begin{bmatrix} 34 & 10 \\ 3 & 13 \end{bmatrix}$	$\begin{bmatrix} 24 & 32 \\ 11 & 6 \end{bmatrix}$	θ_1
47	$2S_4$	$\begin{bmatrix} 11 & 2 \\ 29 & 36 \end{bmatrix}$	$\begin{bmatrix} 46 & 44 \\ 11 & 21 \end{bmatrix}$	θ_1

By using these [Proposition 6.7](#) and [Corollary 6.9](#), along with the conditions in [Table 4](#), we arrive at the following:

Proposition 6.10. *Only the following possibilities for $K \cap SL_2(\mathbb{F}_q)$ can act transitively on $\mathbb{P}^1\mathbb{F}_q$, for $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$:*

1. $2A_4 \leq SL_2(\mathbb{F}_5)$
2. $2S_4 \leq SL_2(\mathbb{F}_7)$
3. $2A_5 \leq SL_2(\mathbb{F}_{11})$
4. $2S_4 \leq SL_2(\mathbb{F}_{23})$
5. $2S_4 \leq SL_2(\mathbb{F}_{31})$
6. $2S_4 \leq SL_2(\mathbb{F}_{47})$

We do not claim these possibilities for K do act transitively on the projective space.

These can be checked by hand, using the character table in [\[Ada02\]](#), to arrive at:

Proposition 6.11. *For the possibilities listed in [Proposition 6.10](#), the different $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$, along with a set of generators (up to conjugation) and a character χ such that $\chi|_K \supseteq \mathbb{1}_K$, are given in [Table 5](#).*

Thus, either $\phi_1 \supseteq \mathbb{1}_K$ or $\theta_1 \supseteq \mathbb{1}_K$ every $K \in (\mathcal{C}_6 \cup \mathcal{S})(GL_2(\mathbb{F}_q))$. In addition, both characters restrict to the projective and special cases.

This completes the proof of [Theorem 1.9](#) and [Theorem 1.10](#).

7 Open Questions

Minimality of $\mathcal{X}(GL_n(\mathbb{F}_q))$ In [Theorem 1.6](#) we have shown that the set $\mathcal{X}(GL_n(\mathbb{F}_q)) \subseteq \widehat{GL_n(\mathbb{F}_q)}$ defined in [Section 1.1](#) suffices to mark every subset of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$. We have not shown that this set is indeed minimal. This raises the following question, and similar questions for the relatives of $GL_n(\mathbb{F}_q)$.

Question 7.1. Is there a set $\mathcal{X}(GL_n(\mathbb{F}_q)) \subseteq \widehat{GL_n(\mathbb{F}_q)}$ with at most 4 nontrivial irreducible characters that marks every subgroup of $GL_n(\mathbb{F}_q)$ that does not contain $SL_n(\mathbb{F}_q)$?

$\mathcal{X}(G)$ for other groups Given a group G , we denote by $c_{\min}(G)$ the minimal cardinality of a subset of nontrivial irreducible characters that marks every proper subgroup of G .

Question 7.2. Given a finite group G , what is $c_{\min}(G)$?

Question 7.3. Given a finite group G , what subsets of characters achieve $c_{\min}(G)$? Is there a unique minimal set?

Families of groups Given a family of finite groups, we consider $\max_G(c_{\min}(G))$, where the maximum ranges over all of the groups in the family.

Question 7.4. Given a family of finite groups, what is $\max_G(c_{\min}(G))$, where the maximum ranges over all of the groups in the family?

In particular, for the case of S_n , we prove in [Appendix A](#) that $\max_n(c_{\min}(S_n)) \leq 8$ for $n \geq 34$.

Question 7.5. What is the value of $\max_n(c_{\min}(S_n))$?

A Characters and Subgroups of S_n

In this section, we consider the group S_n . We use methods similar to those used in this paper to find a set of characters $\mathcal{X}(S_n)$ that marks every proper subgroup of S_n . Note that in [Example 1.5](#), we presented such a set with 12 elements, and the set we present in this section consists of 8 elements.

We introduce notation similar to the notation in the case of $GL_n(\mathbb{F}_q)$. Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_s)$, such that $\sum_{i=1}^s \alpha_i = n$. We denote the induction of the trivial character of $S_{\alpha_1} \times \dots \times S_{\alpha_s}$ to S_n by $\mathbb{1}_\alpha$.

Lemma A.1 (e.g. [\[Ful97, Chapter 7.3, Corollary 1\]](#)). *Let $\lambda = (\lambda_1, \dots, \lambda_l) \vdash n$. Then*

$$\mathbb{1}_\lambda = \sum_{\mu \vdash n} K_{\mu\lambda} \chi^\mu,$$

where $K_{\mu\lambda}$ are the Kostka numbers, and χ^μ are irreducible characters.

We define the characters $\chi^\mu \in \widehat{S}_n$ as these irreducible summands of $\mathbb{1}_\lambda$. Unlike the case of $GL_n(\mathbb{F}_q)$, these are all the irreducible characters of S_n . This definition agrees with the usual correspondence between partitions and irreducible characters (e.g. [\[Ful97\]](#)). χ^μ

We begin by considering the character $\mathbb{1}_{(n-3,1^3)}$ of S_n . It is the character induced by the permutation representation of S_n on 3-tuples of elements of $X_n = \{1, \dots, n\}$. Therefore, by [Corollary 2.6](#), $\mathbb{1}_{(n-3,1^3)} - \mathbb{1}_{(n)}$ marks every subgroup of S_n that does not act on X_n 3-transitively. By decomposing the character, we arrive at the following:

Proposition A.2. *Let $n \geq 6$ and let*

$$\mathcal{X}(S_n) = \left\{ \begin{array}{ccc} \chi^{(n-1,1)} & \chi^{(n-2,2)} & \chi^{(n-2,1^2)} \\ \chi^{(n-3,3)} & \chi^{(n-3,2,1)} & \chi^{(n-3,1^3)} \end{array} \right\} \subseteq \widehat{S}_n.$$

Then, $\mathcal{X}(S_n)$ marks every subgroup of S_n that is not 3-transitive under the usual action.

Therefore, it remains to consider the 3-transitive subgroups of S_n . By the classification of finite multiply transitive groups, we arrive at the following:

Proposition A.3. *Let $n \geq 25$. Then, the only possibilities for 3-transitive subgroups of S_n , along with the 3-transitive actions, are:*

1. *The groups S_n and A_n with the usual action on $\{1, \dots, n\}$.*

2. Groups $\mathbb{P}SL_2(\mathbb{F}_q) \leq G \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$ with the usual action on $\mathbb{P}^1\mathbb{F}_q$, for $q = n - 1$ a prime power. Such a group G is 3-transitive if and only if^{a3} for every $A \in \mathbb{P}GL_2(\mathbb{F}_q)$ there exists some $\sigma \in \text{Aut}(\mathbb{F}_q)$ such that $(A, \sigma) \in G$.
3. The affine linear groups $AGL_d(\mathbb{F}_2)$ with the usual action on \mathbb{F}_2^d , for $n = 2^d$ with $d \geq 5$.

Proof. The possibilities for 2-transitive groups can be seen in, e.g. [DM96, Chapter 7.7]. The only 3-transitive groups are either S_n or A_n with their usual action (listed in (1)), groups $\mathbb{P}SL_2(\mathbb{F}_q) \leq G \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$ with the usual action on $\mathbb{P}^1\mathbb{F}_q$, for $q = n - 1$ a prime power, or subgroups of $A\Gamma L_d(\mathbb{F}_q)$ with the action on \mathbb{F}_q^d . This can be seen in, e.g. [Cam81, Theorem 5.3].

$\mathbb{P}SL_2(\mathbb{F}_q) \leq G \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$

If $q = n - 1$ is a prime power, a group $\mathbb{P}SL_2(\mathbb{F}_q) \leq G \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$ with the usual action on $\mathbb{P}^1\mathbb{F}_q$ is 3-transitive if and only if given some distinct $x_1, x_2, x_3 \in \mathbb{P}^1\mathbb{F}_q$, for every distinct $y_1, y_2, y_3 \in \mathbb{P}^1\mathbb{F}_q$ there exists some $g \in G$ such that $gx_i = y_i$ for $i = 1, 2, 3$. In particular, denote by q_0 the characteristic of \mathbb{F}_q . Then \mathbb{F}_{q_0} is a subfield of \mathbb{F}_q fixed by the automorphisms in $\text{Aut}(\mathbb{F}_q)$.

In particular, if we choose distinct $x_1, x_2, x_3 \in \mathbb{P}^1\mathbb{F}_{q_0}$, we get $(A, \sigma)(x_i) = Ax_i$ for every $(A, \sigma) \in G \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$. Then, G is 3-transitive if and only if for every distinct $y_1, y_2, y_3 \in \mathbb{P}^1\mathbb{F}_q$ there exists some $(A, \sigma) \in G$ such that $Ax_i = y_i$ for $i = 1, 2, 3$.

Since $\mathbb{P}GL_2(\mathbb{F}_q)$ acts 3-transitively on $\mathbb{P}^1\mathbb{F}_q$ and is of order $q(q^2 - 1)$, it is sharply 3-transitive. Therefore, there is a bijection between distinct $y_1, y_2, y_3 \in \mathbb{P}^1\mathbb{F}_q$ and $A \in \mathbb{P}GL_2(\mathbb{F}_q)$ such that $Ax_i = y_i$ for $i = 1, 2, 3$. We denote $A = A(y_1, y_2, y_3)$.

To sum up, the following are equivalent:

1. G is 3-transitive.
2. For every distinct $y_1, y_2, y_3 \in \mathbb{P}^1\mathbb{F}_q$ there exists some $(A, \sigma) \in G$ such that $Ax_i = y_i$ for $i = 1, 2, 3$.
3. For every distinct $y_1, y_2, y_3 \in \mathbb{P}^1\mathbb{F}_q$ there exists some $\sigma \in \text{Aut}(\mathbb{F}_q)$ such that $(A(y_1, y_2, y_3), \sigma) \in G$.
4. For every $A \in \mathbb{P}GL_2(\mathbb{F}_q)$ there exists some $\sigma \in \text{Aut}(\mathbb{F}_q)$ such that $(A, \sigma) \in G$.

$G \leq A\Gamma L_d(\mathbb{F}_q)$ The only case remaining is where $G \leq A\Gamma L_d(\mathbb{F}_q)$. Assume G is 3-transitive and consider the stabilizer $G_0 \leq \Gamma L_d(\mathbb{F}_q)$, which acts 2-transitively on $\mathbb{F}_q^d \setminus \{0\}$. We use linearity to claim this implies that $q = 2$. Denote by q_0 the characteristic of \mathbb{F}_q , and let ℓ be the exponent, $q = q_0^\ell$.

$d \geq 2, q \geq 3$ There is no $g \in G_0$ with $g(e_1, ae_1) = (e_1, e_2)$, for $a \in \mathbb{F}_q, a \neq 0, 1$.

$d = 1$ Let $q = q_0^\ell$ with q_0 prime. Observe that $25 \leq n = q$. Let $a \in \mathbb{F}_q, a \neq 0, 1$. Consider the orbit $O_{\text{Aut}(\mathbb{F}_q)}(a)$, which is of size at most ℓ . Therefore, $|\mathbb{F}_q \setminus (\{0, 1\} \cup O_{\text{Aut}(\mathbb{F}_q)}(a))| = q - \ell - 2$. It is easy to see that $q \geq \ell + 3$ for $q \geq 25$. As such, we can choose b such that $b \in \mathbb{F}_q, b \neq 0, 1$ and b is not in the same orbit of $\text{Aut}(\mathbb{F}_q)$ as a . Then, there is no $g \in G_0$ with $g(e_1, ae_1) = (e_1, be_1)$.

Therefore, $q = 2$. Since $n \geq 25$, we get that $d \geq 5$. As such, G_0 acts 2-transitively on $\mathbb{F}_2^d \setminus \{0\} = \mathbb{P}^{d-1}\mathbb{F}_2$. By Theorem 5.5, $G_0 \geq SL_d(\mathbb{F}_2) = GL_d(\mathbb{F}_2) = \Gamma L_d(\mathbb{F}_2)$. Therefore, $G_0 = GL_d(\mathbb{F}_2)$ and $G = AGL_d(\mathbb{F}_2)$. \square

Proposition A.4. *Let $n \geq 25$ and $n \neq 33$. Then, $\mathbb{1}_{(n-4,4)} - \mathbb{1}_{(n)}$ marks every 3-transitive proper subgroup of S_n , except A_n .*

¹³For example, $\mathbb{P}SL_2(\mathbb{F}_q)$ is not 3-transitive when q is odd, while $\mathbb{P}GL_2(\mathbb{F}_q)$ is always 3-transitive.

Proof. Let K be one of the groups listed in [Proposition A.4](#). Note that $\mathbb{1}_{(n-4,4)}$ is the character of the permutation representation of S_n acting on Y_n , the collection subsets of $X_n = \{1, \dots, n\}$ of size 4.

By [Corollary 2.6](#), $\mathbb{1}_{(n-4,4)}$ marks precisely the subgroups of S_n that act transitively on Y_n . Thus, it is sufficient to show that K does not act transitively on Y_n .

$K \leq \mathbb{P}\Gamma L_2(\mathbb{F}_q)$ Let $q = q_0^\ell$, with q_0 prime. By [\(4.1\)](#), $|Y_n| \mid |K|$. Note that $|Y_n| = \binom{n}{4}$. Since $|K| \mid |\mathbb{P}\Gamma L_2(\mathbb{F}_q)|$, we get $\binom{q+1}{4} = |Y_{q+1}| \mid |\mathbb{P}\Gamma L_2(\mathbb{F}_q)| = \ell(q^3 - q)$, which is equivalent to $(q-2) \mid 24\ell$. It is easy to see that no $q = n-1 \geq 24$ except $q = 32$ (which is explicitly omitted) satisfies this equation.

$K = \text{AGL}_d(\mathbb{F}_2)$ We know that $d \geq 5$. It is easy to see that if x_1, x_2, x_3, x_4 satisfy $x_1 + x_2 = x_3 + x_4$ then $gx_1 + gx_2 = gx_3 + gx_4$ for every $g \in K$. As such, there exists no $g \in K$ such that $g\{0, e_1, e_2, e_1 + e_2\} = \{0, e_1, e_2, e_3\}$.

□

We use [Proposition A.2](#) and [Proposition A.4](#). Then, we decompose the character $\mathbb{1}_{(n-4,4)} - \mathbb{1}_{(n)} + \mathbb{1}_{(n-3,1^3)} - \mathbb{1}_{(n)}$ to irreducible characters and add the sign character (which is equal to $\chi^{(1^n)}$) to arrive at the following:

Corollary A.5. *Let $n \geq 34$ and let*

$$\mathcal{X}(S_n) = \left\{ \begin{array}{cccc} \chi^{(n-1,1)} & \chi^{(n-2,2)} & \chi^{(n-2,1^2)} & \chi^{(n-3,3)} \\ \chi^{(n-3,2,1)} & \chi^{(n-3,1^3)} & \chi^{(n-4,4)} & \chi^{(1^n)} \end{array} \right\} \subseteq \widehat{S_n}.$$

Then, $\mathcal{X}(S_n)$ marks every proper subgroup of S_n .

B Group Extensions

Throughout this section, given a finite group H , we denote by $c_{\min}(H)$ the minimal cardinality of a subset of nontrivial irreducible characters that marks every proper subgroup of H . To simplify the definitions, we say that a set $\mathcal{X}(H)$ achieves $c_{\min}(H)$ if $\mathcal{X}(H)$ is a subset of nontrivial irreducible characters of H that marks every proper subgroup of H with $|\mathcal{X}(H)| = c_{\min}(H)$.

$\mathcal{X}(H)$
achieves
 $c_{\min}(H)$

Let G be a finite group and let $N \trianglelefteq G$. We study the relations between $c_{\min}(G)$, $c_{\min}(G/N)$ and $c_{\min}(N)$.

B.1 $c_{\min}(G)$ and $c_{\min}(G/N)$

Denote by $f: G \twoheadrightarrow G/N$ the quotient epimorphism. If $\chi \in \text{Ch}(G/N)$, the character can be pulled back by f to obtain a character $f^*\chi$ of G , $f^*\chi(x) = \chi(f(x))$. It is easy to see that χ is irreducible if and only if $f^*\chi$ is irreducible, and that $f^*: \text{Ch}(G/N) \rightarrow \text{Ch}(G)$ is injective.

For every $K \leq G$ such that $K \geq N$, we denote the quotient epimorphism $K \twoheadrightarrow K/N$ by f_K , and the pull-back from characters of K/N to characters of K by f_K^* .

The following properties of the pull-back of an epimorphism are easy to verify:

Lemma B.1. *Let G be a finite group and let $N \trianglelefteq G$. Let $f: G \twoheadrightarrow G/N$ be the quotient epimorphism, let $\phi, \psi \in \text{Ch}(G/N)$, let $\theta \in \text{Ch}(K/N)$, and let $N \leq K \leq G$. Then:*

1. $\mathbb{1}_G = f^*\mathbb{1}_{G/N}$
2. $f^*(\phi + \psi) = f^*\phi + f^*\psi$

$$3. \langle \phi, \psi \rangle_{G/N} = \langle f^* \phi, f^* \psi \rangle_G$$

$$4. f^* \text{Ind}_{K/N}^{G/N}(\theta) = \text{Ind}_K^G(f^* \theta)$$

Proposition B.2. *Let G be a finite group, and let $N \trianglelefteq G$. Denote by $f: G \twoheadrightarrow G/N$ the quotient epimorphism. Let $K \leq G$ be some subgroup such that $K \geq N$. Let $\chi \in \widehat{G}$ be an irreducible character that satisfies $\chi|_K \supseteq \mathbb{1}_K$. Then $\chi \in \text{Im}(f^*)$.*

Proof. Since $\chi|_K \supseteq \mathbb{1}_K$, using [Frobenius Reciprocity](#) we deduce that

$$0 < \langle \chi|_K, \mathbb{1}_K \rangle_K = \langle \chi, \text{Ind}_K^G(\mathbb{1}_K) \rangle_G$$

which implies that $\text{Ind}_K^G(\mathbb{1}_K) \supseteq \chi$.

Let $\text{Ind}_{K/N}^{G/N}(\mathbb{1}_{K/N}) = \sum_i \phi_i$ for some $\phi_i \in \widehat{G/N}$, possibly with repetitions. We have $\text{Ind}_K^G(\mathbb{1}_K) = \text{Ind}_K^G(f^* \mathbb{1}_{K/N}) = f^* \text{Ind}_{K/N}^{G/N}(\mathbb{1}_{K/N}) = \sum_i f^* \phi_i$, which implies that all irreducible summands of $\text{Ind}_K^G(\mathbb{1}_K)$ are in $\text{Im}(f^*)$. In particular, $\chi \in \text{Im}(f^*)$. \square

Proposition B.3. *Let G be a finite group, and let $N \trianglelefteq G$. Denote by $f: G \twoheadrightarrow G/N$ the quotient epimorphism. Let $\mathcal{X}(G) \subseteq \widehat{G}$ be some set of nontrivial irreducible characters that marks every proper subgroup of G , and let*

$$\mathcal{X}(G/N) := \left\{ \chi \in \widehat{G/N} : f^* \chi \in \mathcal{X}(G) \right\}.$$

Then, $\mathcal{X}(G/N)$ marks every proper subgroup of G/N , and consists of nontrivial irreducible characters.

Proof. Let $J \leq G/N$. By the correspondence theorem, there exists some $K \leq G$ with $K \geq N$ and $f(K) = J$. Thus, there exists some $\chi \in \mathcal{X}(G)$ such that $\chi|_K \supseteq \mathbb{1}_K$.

By [Proposition B.2](#), $\chi \in \text{Im}(f^*)$. Therefore, there exists some $\chi' \in \widehat{G/N}$ such that $f^* \chi' = \chi$. This implies that $\chi' \in \mathcal{X}(G/N)$.

By [Lemma B.1](#) (with $G = K$), $\langle \chi'|_J, \mathbb{1}_J \rangle_J = \langle \chi|_K, \mathbb{1}_K \rangle_K > 0$.

To prove the characters in $\mathcal{X}(G/N)$ are nontrivial and irreducible, let $\chi \in \mathcal{X}(G/N)$. Since $f^* \chi \in \mathcal{X}(G)$, it is nontrivial and irreducible. By the properties of pull-backs, so is χ . \square

Remark This definition of $\mathcal{X}(G/N)$ is the same definition as the one made in [Section 5.2](#).

Corollary B.4.

$$c_{\min}(G/N) \leq c_{\min}(G).$$

Proof. Let $\mathcal{X}(G) \subseteq \widehat{G}$ be a set achieving $c_{\min}(G)$, let $f: G \twoheadrightarrow G/N$ be the quotient epimorphism, and let $\mathcal{X}(G/N)$ be as defined in [Proposition B.3](#). We get an injection

$$f^*|_{\mathcal{X}(G/N)}: \mathcal{X}(G/N) \rightarrow \mathcal{X}(G),$$

which implies

$$c_{\min}(G/N) \leq |\mathcal{X}(G/N)| \leq |\mathcal{X}(G)| = c_{\min}(G).$$

\square

Corollary B.5. *Let G be a finite group, and let $\mathcal{X}(G) \subseteq \widehat{G}$ be some set of nontrivial irreducible characters that marks every proper subgroup of G . Then, for every $N \trianglelefteq G$ there exist $c_{\min}\left(\frac{G}{N}\right)$ many characters $\chi \in \mathcal{X}(G)$ such that $\chi|_N = \dim(\chi)\mathbb{1}_N$.*

In particular, if $N \trianglelefteq G$ is proper, there exists at least one such character.

Proof. Let $N \trianglelefteq G$, denote by $f: G \rightarrow G/N$ the quotient epimorphism, and let $e_{G/N}$ be the identity element of G/N . We define $\mathcal{X}\left(\frac{G}{N}\right)$ as in [Proposition B.3](#). Then, $c_{\min}\left(\frac{G}{N}\right) \leq \left|\mathcal{X}\left(\frac{G}{N}\right)\right|$ and every character in $\mathcal{X}\left(\frac{G}{N}\right)$ is mapped to a different character in $\mathcal{X}(G)$ by the injection f^* . For every $\chi \in \mathcal{X}\left(\frac{G}{N}\right)$, the pull-back $f^*\chi \in \mathcal{X}(G)$ satisfies

$$f^*\chi(g) = \chi(f(g)) = \chi\left(e_{G/N}\right) = \dim(\chi),$$

for every $g \in N$. Thus, the claim follows.

If $N \trianglelefteq G$ is proper, G/N is not the trivial group and thus has the group $\{e_{G/N}\}$ as a proper subgroup. As such, $1 \leq c_{\min}\left(\frac{G}{N}\right)$, as every $\mathcal{X}\left(\frac{G}{N}\right)$ achieving it must contain at least one character to mark this proper subgroup. \square

B.2 Abelian Groups

In this section, we consider G which are finite abelian groups. These provide useful examples for [Section B.3](#).

Proposition B.6. *Let G be a finite abelian group, and let $\mathcal{X}(G) \subseteq \widehat{G}$ be a set achieving $c_{\min}(G)$ (since G is abelian, its elements are homomorphisms). Then, there is a bijection*

$$\{H \leq G: H \text{ is maximal}\} \leftrightarrow \mathcal{X}(G), H \mapsto \chi_H$$

where $\chi_H|_H = \mathbb{1}_H$ and $H = \ker(\chi_H)$.

Proof. Since G is abelian, every subgroup of G is normal. Thus, by [Corollary B.5](#), we can choose a character $\chi_H \in \mathcal{X}(G)$ for every maximal subgroup H that satisfies $\chi_H|_H = \dim(\chi_H)\mathbb{1}_H$. Since every character of an abelian group is linear, $\dim(\chi_H) = 1$ and $H \leq \ker(\chi_H)$. By the maximality of H and nontriviality of χ_H , $H = \ker(\chi_H)$. It remains to show that this choice is a bijection.

One-to-one Let $H, H' \leq G$ be maximal subgroups with $\chi_H = \chi_{H'}$. Taking the kernels of these characters, we get $H = H'$.

Onto The set $\{\chi_H: H \text{ is maximal}\}$ marks every maximal subgroup of G , and as such, it marks every proper subgroup of G (by the same argument as the one made in the beginning of [Section 4](#)). By the minimality of $c_{\min}(G) = |\mathcal{X}(G)|$, the mapping is onto. \square

Cyclic groups Let $C_n = \mathbb{Z}/_n\mathbb{Z}$ be the cyclic group of order n , for $n \in \mathbb{Z}_{\geq 1}$.

Proposition B.7. *For every $n \in \mathbb{Z}_{\geq 1}$, $c_{\min}(C_n)$ is precisely the number of different prime divisors of n .*

Proof. Let $g \in C_n$ be a generator. It is known that the subgroups of C_n are precisely $\{\langle g^k \rangle: k|n\}$, and all of these groups are different. It is easy to see that $\langle g^k \rangle$ is maximal if and only if k is prime. Thus, by [Proposition B.6](#), $c_{\min}(C_n) = |\{p \in \mathbb{Z}_{\geq 1}: p \text{ is prime}, p|n\}|$ \square

Corollary B.8. *There exist infinitely many finite groups G with $N \trianglelefteq G$ such that $c_{\min}(G) = a c_{\min}\left(\frac{G}{N}\right)$, for every $a \in \mathbb{Z}_{\geq 1}$.*

Proof. Let $G = C_{2^{m_n}}$, where n is odd, and has $a - 1$ distinct prime factors. For $N = C_n$, the quotient satisfies $G/N = C_{2^m}$. Then, $c_{\min}(C_{2^{m_n}}) = a = a c_{\min}(C_{2^m})$. \square

Corollary B.9. *Let $n \in \mathbb{Z}_{\geq 1}$. Then, $c_{\min}(GL_n(\mathbb{F}_q))$ is at least the number of prime divisors of $q - 1$.*

Proof. By [Corollary B.4](#) with $N = SL_n(\mathbb{F}_q)$, $c_{\min}(GL_n(\mathbb{F}_q)) \geq c_{\min}(C_{q-1})$. The claim follows by [Proposition B.7](#). \square

Vector spaces We continue with the additive groups of the vector spaces \mathbb{F}_p^ℓ , where p is prime and $\ell \in \mathbb{Z}_{\geq 1}$.

Proposition B.10. *Let p be prime and let $\ell \in \mathbb{Z}_{\geq 1}$. Then, $c_{\min}(\mathbb{F}_p^\ell) = \frac{p^\ell - 1}{p - 1}$.*

Proof. Denote by $f_w(x) = \sum_{i=1}^{\ell} x_i w_i$. A maximal subgroup of \mathbb{F}_p^ℓ is a $(\ell - 1)$ -dimensional subspace of \mathbb{F}_p^ℓ , namely, some $W_{\bar{w}} = \{x \in \mathbb{F}_p^\ell : f_w(x) = 0\} = \ker f_w$ where $0 \neq w \in \mathbb{F}_p^\ell$ and $\bar{w} \in \mathbb{P}^{\ell-1}\mathbb{F}_p$ is the point in the projective space represented by w . This shows a bijection between the maximal subgroups of \mathbb{F}_p^ℓ and $\mathbb{P}^{\ell-1}\mathbb{F}_p$. Thus,

$$c_{\min}(\mathbb{F}_p^\ell) = |\mathbb{P}^{\ell-1}\mathbb{F}_p| = \frac{p^\ell - 1}{p - 1}.$$

\square

B.3 $c_{\min}(G)$ and $c_{\min}(N)$

In this section, we study the relation between $c_{\min}(G)$ and $c_{\min}(N)$. We begin with the following proposition.

Proposition B.11. *For a finite group H , let $\mathcal{X}(H)$ be a set achieving $c_{\min}(H)$. Let G be a finite group, let $N \trianglelefteq G$, and let $f: G \twoheadrightarrow G/N$ be the quotient epimorphism. Define*

$$\overline{\mathcal{X}(G)} := \{\text{Ind}_N^G(\chi) : \chi \in \mathcal{X}(N)\} \cup \{f^*\phi : \phi \in \mathcal{X}(G/N)\}.$$

Then, $\overline{\mathcal{X}(G)}$ marks every proper subgroup of G .

Proof. Let $K \leq G$ and let $\chi \in \mathcal{X}(N)$. Denote by $\chi^{(t)}$ the character of N defined by $\chi^{(t)}(g) = \chi(t^{-1}gt)$, for every $t \in G$.

$$\begin{aligned} \langle \text{Ind}_N^G(\chi)|_K, \mathbb{1}_K \rangle_K &= \frac{1}{|K|} \sum_{k \in K} \text{Ind}_N^G(\chi)(k) \\ &= \frac{1}{|K| \cdot |N|} \sum_{k \in K} \left(\sum_{t \in G \text{ s.t. } t^{-1}kt \in N} \chi(t^{-1}kt) \right) \\ &= \frac{1}{|K| \cdot |N|} \sum_{t \in G} \left(\sum_{k \in K \text{ s.t. } t^{-1}kt \in N} \chi(t^{-1}kt) \right) \\ &= \frac{1}{|K| \cdot |N|} \sum_{t \in G} \sum_{k \in K \cap N} \chi(t^{-1}kt) \\ &= \frac{|N \cap K|}{|K| \cdot |N|} \sum_{t \in G} \langle \chi^{(t)}|_{K \cap N}, \mathbb{1}_{K \cap N} \rangle_{K \cap N} \\ &\geq \frac{|N \cap K|}{|K| \cdot |N|} \langle \chi|_{K \cap N}, \mathbb{1}_{K \cap N} \rangle_{K \cap N}. \end{aligned}$$

In particular, if $K \cap N \neq N$ there exists some $\chi \in \mathcal{X}(N)$ such that $\langle \text{Ind}_N^G(\chi)|_K, \mathbb{1}_K \rangle_K > 0$.

Therefore, $\overline{\mathcal{X}(G)}$ marks every subgroup of G that does not contain N , and we may assume that $K \geq N$.

In this case, there exists some $\phi \in \mathcal{X}(G/N)$ such that $\phi|_{K/N} \supseteq \mathbb{1}_{K/N}$. Then, $\langle f^* \phi|_K, \mathbb{1}_K \rangle_K = \langle \phi|_{K/N}, \mathbb{1}_{K/N} \rangle_{K/N} > 0$, which completes the proof. \square

The problem with this proposition is that the set $\overline{\mathcal{X}(G)}$ may not consist of irreducible characters. Therefore, we decompose the characters of $\overline{\mathcal{X}(G)}$ to get a bound on $c_{\min}(G)$.

Corollary B.12. *Let G be a finite group and let $N \trianglelefteq G$. Then*

$$c_{\min}(G/N) \leq c_{\min}(G) \leq [G:N] c_{\min}(N) + c_{\min}(G/N)$$

Proof. The left inequality follows from [Corollary B.4](#). To prove the right inequality it is sufficient to show, by [Proposition B.11](#), that for every $\chi \in \widehat{N}$ the induced character $\text{Ind}_N^G(\chi)$ decomposes to at most $[G:N]$ irreducible characters of G .

Let $\chi \in \widehat{N}$, and let ϕ be an irreducible summand of $\text{Ind}_N^G(\chi)$. By [Frobenius Reciprocity](#), $\langle \phi|_N, \chi \rangle_N = \langle \phi, \text{Ind}_N^G(\chi) \rangle_G > 0$. Therefore $\phi|_N \supseteq \chi$, which implies that $\dim(\phi) = \dim(\phi|_N) \geq \dim(\chi)$.

Assume that $\text{Ind}_N^G(\chi)$ decomposes to k irreducible characters of G , which we denote by ϕ_1, \dots, ϕ_k (these may be not distinct). By calculating the degrees:

$$[G:N] \dim(\chi) = \dim(\text{Ind}_N^G(\chi)) = \dim\left(\sum_{i=1}^k \phi_i\right) = \sum_{i=1}^k \dim(\phi_i) \geq k \dim(\chi)$$

which implies that $k \leq [G:N]$. \square

We show these inequalities are sharp by considering two examples.

Example B.13. Let $G = C_{p^\ell}$ and $N = C_{p^{\ell-1}} \trianglelefteq G$. By [Proposition B.7](#), every $\mathcal{X}(G)$ achieving $c_{\min}(G)$ consists of a single character, which we denote by χ . By the properties of $\mathcal{X}(G)$, when restricting χ to N , we get $\chi|_N = \mathbb{1}_N$, which cannot be a member of $\mathcal{X}(N)$. Then, by [Frobenius Reciprocity](#), χ cannot be the induction of a nontrivial irreducible character of N . In this case, $c_{\min}(G/N) = c_{\min}(G)$.

Example B.14. Let p be prime and let $\ell \in \mathbb{Z}_{\geq 1}$. Let $G = \mathbb{F}_p^\ell$ and $N \trianglelefteq G$. We use standard facts about G and [Proposition B.10](#).

1. $G \cong \mathbb{F}_p^\ell$, $c_{\min}(G) = \frac{p^\ell - 1}{p - 1}$
2. $N \cong \mathbb{F}_p^i$ for some $0 \leq i \leq \ell$, $c_{\min}(N) = \frac{p^i - 1}{p - 1}$
3. $G/N \cong \mathbb{F}_p^{\ell-i}$, $c_{\min}(G/N) = \frac{p^{\ell-i} - 1}{p - 1}$

From these we deduce: $c_{\min}(G) = [G:N] c_{\min}(N) + c_{\min}(G/N)$.

Acknowledgments

This paper forms part of an M.Sc. Thesis written by the author under the supervision of Dr. Doron Puder from Tel Aviv University. I deeply thank my thesis supervisor for introducing me to many fascinating topics in mathematics and showing me the beauty in them, as well as for his advice, guidance, patient readings of the various drafts and the time he has invested in the process of making this paper. I thank Professor David Soudry from Tel Aviv University, for sharing his knowledge about the general linear group. The research was supported by the Israel Science Foundation, ISF grant 1071/16.

References

- [Ada02] Jeffrey Adams. Character tables for $GL(2)$, $SL(2)$, $PGL(2)$ and $PSL(2)$ over a finite field. *Lecture Notes, University of Maryland*, 2002. [25](#), [26](#), [28](#)
- [Asc84] Michael Aschbacher. On the maximal subgroups of the finite classical groups. *Inventiones mathematicae*, 76(3):469–514, 1984. [16](#)
- [AT17] Scott Andrews and Nathaniel Thiem. The combinatorics of GL_n generalized Gelfand–Graev characters. *Journal of the London Mathematical Society*, 95(2):475–499, 2017. [13](#), [14](#)
- [BHRD13] John N. Bray, Derek F. Holt, and Colva M. Roney-Dougal. *The maximal subgroups of the low-dimensional finite classical groups*, volume 407. Cambridge university press, 2013. [20](#), [27](#)
- [Cam81] Peter J. Cameron. Finite permutation groups and finite simple groups. *Bulletin of the London Mathematical Society*, 13(1):1–22, 1 1981. [4](#), [30](#)
- [CK79] Peter J. Cameron and William M. Kantor. 2-transitive and antiflag transitive collineation groups of finite projective spaces. *Journal of Algebra*, 60(2):384–422, 1979. [4](#), [23](#)
- [CLR10] Pietro Caputo, Thomas Liggett, and Thomas Richthammer. Proof of Aldous spectral gap conjecture. *Journal of the American Mathematical Society*, 23(3):831–851, 2010. [3](#)
- [Dic03] Leonard Eugene Dickson. *Linear groups: with an exposition of the Galois field theory*. Courier Corporation, 2003. [27](#)
- [DM96] John D. Dixon and Brian Mortimer. *Permutation groups*, volume 163. Springer Science & Business Media, 1996. [4](#), [23](#), [30](#)
- [FH13] William Fulton and Joe Harris. *Representation theory: a first course*, volume 129. Springer Science & Business Media, 2013. [2](#), [8](#)
- [Ful97] William Fulton. *Young tableaux: with applications to representation theory and geometry*, volume 35. Cambridge University Press, 1997. [4](#), [23](#), [29](#)
- [Gre55] James A. Green. The characters of the finite general linear groups. *Transactions of the American Mathematical Society*, 80(2):402–447, 1955. [13](#), [14](#)
- [KL90] Peter B. Kleidman and Martin W. Liebeck. *The subgroup structure of the finite classical groups*, volume 129. Cambridge University Press, 1990. [16](#), [17](#)
- [Leh73] Gustav Isaac Lehrer. The characters of the finite special linear groups. *Journal of Algebra*, 26(3):564–583, 1973. [21](#), [22](#)
- [PP18] Ori Parzanchevski and Doron Puder. Aldous’ spectral gap conjecture for normal sets. *preprint arXiv:1804.02776*, 2018. [3](#)