

**CORRIGENDUM TO THE PROOF OF ONE THEOREM IN THE
ARTICLE “ON MULTIPLICATIVE SUBGROUPS IN DIVISION
RINGS” [J. ALGEBRA AND ITS APPLICATIONS VOL. 15,
NO. 3 (2016) 1550050 (16 PAGES)]**

BUI XUAN HAI^{1,2}

ABSTRACT. In this note, the correction to the proof of one theorem in some our previous paper will be given.

In the proof of [4, Theorem 3.4] there is the following mistake: in row -12 , page 1650050 – 11, we affirmed that by applying [3, Theorem 5], it follows that M is a (locally abelian)-by-finite group. But, in the reality, this argument is not valid as it was well remarked in recently accepted paper [1]. Then, in the main theorem of [1], the author has proved the similar result as our in [4, Theorem 3.4], but for more narrow class of division rings. In fact, he proved the result for centrally finite division rings, while our theorem holds for locally finite division rings. Firstly, we must confirm that [4, Theorem 3.4] is always true. Moreover, this result was generalized in [2, Theorem 3.1] for more general circumstance. Hence, [1, Theorem 4.1], and [4, Theorem 3.4] both are particular cases of [2, Theorem 3.1]. Our purpose in this note is to give the another proof of [4, Theorem 3.4]. The idea of this proof is very different from our in the proof of [2, Theorem 3.1], and has no any connection with the proof of [1, Theorem 4.1].

For the next use, we need some lemmas.

Lemma 1. *Let D be a locally finite division ring with the center F , and G a subgroup of D^* . If G' is radical over F , then G' is locally finite and either $G' \cong \text{SL}(2, 5)$ or G is solvable.*

Proof. In the first, we show that if an element $x \in D'$ is radical over F then x is periodic. Indeed, assume that $x = [x_1, y_1] \dots [x_k, y_k]$ and $x^n = a \in F$ for some positive integer n . The division subring $K = F(x_1, y_1, \dots, x_k, y_k)$ generated by $x_1, y_1, \dots, x_k, y_k$ over F in D is a finite dimensional vector space over F , say, of degree $m = [K : F]$. Hence,

$$1 = N_{K/F}(x)^n = N_{K/F}(x^n) = N_{K/F}(a) = a^m,$$

where $N_{K/F}$ denotes the norm of K to F . So $x^{nm} = 1$, and x is periodic. Therefore, G' is torsion. Since D is locally finite, G' can be viewed as locally linear group, so, by Schur's Theorem [5, p.154], G' is locally finite. The Amitsur's Theorem [9, 2.1.11, p.51] shows that the only unsolvable finite subgroup of a division ring is $\text{SL}(2, 5)$. Hence, if G' is not locally solvable, then $G' \cong \text{SL}(2, 5)$. Now, assume that G' is locally solvable. Then, by Zaleskii's result [11], G' is solvable, so is G . \square

Key words and phrases. maximal subgroups, subnormal subgroups, FC -elements.
2010 *Mathematics Subject Classification.* 16K20.

Lemma 2. *Let D be a division ring with center F , G a subnormal subgroup of D^* and assume that M is a maximal subgroup of G . If $[M : Z(M)] < \infty$, then M is abelian.*

Proof. Suppose that M is non-abelian and $\{x_1, \dots, x_m\}$ is a transversal of $Z(M)$ in M . Denote by E the subfield of D generated by $Z(M)$ over the prime subfield P of F , that is $E = P(Z(M))$, and put

$$K = \left\{ \sum_{i=1}^m a_i x_i \mid a_i \in E \right\}.$$

Clearly, K is a subring of D . Moreover, K is a division subring of D because it is a finite dimensional vector space over its center $E \subseteq Z(K)$. Also, we have $C_K(M) = Z(K)$. Since M is a maximal subgroup of G , either $M = K^* \cap G$ or $G \subseteq K^*$.

Case 1: $M = K^ \cap G$.*

In this case, M is subnormal in K^* . For $H = \langle x_1, \dots, x_m \rangle$, we have $M = HZ(M)$, hence $M' \leq H$, so H is normal in M . Therefore, H is a finitely generated subnormal subgroup of K^* , and by [7, Theorem 1], $H \subseteq Z(K)$. In particular, H is abelian, and this implies that M is abelian too, but this is a contradiction because by the assumption, M is non-abelian.

Case 1: $G \subseteq K^$.*

By Stuth's theorem, either $K \subseteq F$ or $K = D$. If $K \subseteq F$, then $M \subseteq F$ that is a contradiction. Hence, $K = D$, so D is centrally finite and $C_D(M) = Z(D) = F$. Take an arbitrary element $x \in G \setminus M$ and set $H = \langle x, x_1, \dots, x_m \rangle$. By maximality of M in G , we have $G = HZ(M)$. Hence, $G' \subseteq H$, so H is normal in G , and consequently, H is a finitely generated subnormal subgroup of D^* . By [7, Theorem 1], $H \subseteq F$. It follows $M \subseteq F$ that is impossible. \square

Theorem 3. ([4, Theorem 3.4]) *Let D be a locally finite division ring with center F and G a subnormal subgroup of D^* . Assume that M is a non-abelian maximal subgroup of G . If M contains no non-cyclic free subgroups, then $[D : F] < \infty$, $F(M) = D$, and there exists a maximal subfield K of D such that K/F is a Galois extension, $\text{Gal}(K/F) \cong M/K^* \cap G$ is a finite simple group, and $K^* \cap G$ is the FC-center. Also, $K^* \cap G$ is the Fitting subgroup of M .*

Proof. Assume that M does not contain non-cyclic free subgroups. We claim that $F(M) = D$. Indeed, if $F(M) \neq D$, then, by Stuth's theorem, $F(M)$ is not normalized by G . So, by maximality of M in G , we have $F(M) \cap G = M$. Thus, M is a non-abelian subnormal subgroup of $F(M)^*$ containing no non-cyclic free subgroup that contradicts to [3, Theorem 11]. Therefore, $F(M) = D$ as we have claimed. From the last equality, it follows $Z(M) = M \cap F$. Since D is locally finite, M is locally linear group. So, by Tit's Alternative [10], M is locally solvable-by-finite. By [9, 3.3.9, p.103], M is (locally solvable)-by-(locally finite). Note that the only unipotent element of a division ring is 1, so, by [9, 3.3.2, p.96], M is solvable-by-(locally finite). Therefore, there exists a solvable normal subgroup H of M such that M/H is locally finite. By [9, Theorem 5.5.1, Part a), p.199], there exists an abelian normal subgroup A of M such that H/A is locally finite. Hence, by [8, 14.3.1, p.429], M/A is locally finite. Thus, M is abelian-by-(locally finite).

We claim that there exists an abelian normal subgroup of M which is not contained in F . Indeed, if A is not contained in F , then A is such a subgroup. Otherwise, $A \subseteq F$ and it follows that M is radical over F because M/A is locally finite. By Lemma 1, M' is locally finite.

Now, let us consider three possible cases.

Case 1: $\text{char}(F) > 0$.

Since M' is locally finite, M' is abelian. Denotes by B the maximal abelian normal subgroup of M containing M' . The proof of [4, Theorem 3.3 (i)] shows in particular that B is a normal subgroup of M which is not contained in F .

Case 2: $\text{char}(F) = 0$ and $[D : F] < \infty$.

By Tits' Alternative [10], there exists a normal solvable subgroup S of M such that M/S is finite. By [6, Lemma 3], S contains a normal abelian subgroup H such that S/H is finite. Then, $B = \text{Core}_M(H)$ is a normal abelian subgroup of M of finite index. If $B \subseteq F$, then $B \subseteq Z(M)$, so it follows that $[M : Z(M)] < \infty$. Hence, in view of Lemma 2, M is abelian that is a contradiction.

Case 3: $\text{char}(F) = 0$ and $[D : F] = \infty$.

Assume that M' is not solvable. By Lemma 1, $M' \cong \text{SL}(2, 5)$. If $F(M') \cap G \subseteq M$, then $M' \triangleleft F(M') \cap G$ and this implies that M' is subnormal subgroup of $F(M')$. Since M' contains no non-cyclic free subgroup, in view of [3, Theorem 11], M' is contained in the center of $F(M')$. In particular, M' is abelian that is impossible because $M' \cong \text{SL}(2, 5)$. Thus, $F(M') \cap G \not\subseteq M$. Since M is maximal in G , $(F(M') \cap G)M = G$, so G normalizes $F(M')^*$. By Stuth's theorem, we have $F(M') = D$ that entails $[D : F] < \infty$, a contradiction. Therefore, M' is solvable. Let r be a natural number such that $M^{(r)} \not\subseteq F$ and $M^{(r+1)} \subseteq F$. Then, $M^{(r)}$ is nilpotent, so in view of [9, 2.5.2, p. 73], there is an abelian normal subgroup B of $M^{(r)}$ of finite index. Now, consider the division subring $D_1 = F(M^{(r)})$ of D . Clearly, $M^{(r)}$ normalizes D_1 . Suppose that G also normalizes D_1 . Since $M^{(r)} \not\subseteq F$, by Stuth's theorem, we have $F(M^{(r)}) = D_1 = D$. If $\{x_1, \dots, x_m\}$ is a transversal of B in $M^{(r)}$, then $F(M^{(r)}) = \sum_{i=1}^m x_i F(B)$. This implies the right dimension $[D : F(B)]_r$ is finite. By Double Centralizer Theorem, $[D : F] < \infty$, a contradiction. Hence, D_1 is not normalized by G . Then, by maximality of M in G , we have $D_1 \cap G \subseteq M$. This implies that $M^{(r)}$ is subnormal in D_1 , so $M^{(r)}$ is a nilpotent subnormal subgroup of D_1 . By Lemma [4, Lemma 3.1], $M^{(r)} \subseteq Z(D_1)$, so $M^{(r)}$ is an abelian normal subgroup of M which is not contained in F . Thus, we have proved that in all three cases, there exists an abelian normal subgroup of M which is not contained in F , as claimed. The claim is proved.

Now, assume that C is an abelian normal subgroup of M which is not contained in F . Then, $C \not\subseteq Z(M)$ because as we have noted above, $Z(M) = M \cap F$. Take an element $\alpha \in C \setminus Z(M)$. Since $Z(M) = M \cap F$ and $M/Z(M)$ is locally finite, α is algebraic over F . Set $K = F(\alpha^M)$ and $H = C_D(K)$. The condition $C \triangleleft M$ implies that the elements of α^M in the field $F(\alpha^M)$ have the same minimal polynomial over F , so $|\alpha^M| < \infty$, hence α is an FC -element. Therefore, we can apply [4,

Theorem 3.2] for α, K and H . In particular, $H^* \cap G$ is a subgroup of M . Clearly, $H^* \cap G$ is subnormal in H^* . So, by [3, Theorem 11], we have

$$H^* \cap G \subseteq Z(H) = C_H(H) \subseteq C_D(H) = C_D(C_D(K)) = K.$$

Therefore, all the requirements of [4, Theorem 3.2] are satisfied. The proof of the theorem is now complete. \square

REFERENCES

- [1] R. Fallah-Moghaddam, Free Subgroups in Maximal Subgroups of $SL_n(D)$, *J. Algebra Appl.*, doi:10.1142/S0219498821500717.
- [2] B. X. Hai, H. V. Khanh, Free subgroups in maximal subgroups of skew linear groups, *Internat. J. Algebra Comput.* **29**(3) (2019) 603–614.
- [3] Bui Xuan Hai and Nguyen Kim Ngoc, A note on the existence of non-cyclic free subgroups in division rings, *Archiv der Math.* **101** (2013) 437–443.
- [4] B. X. Hai and N. A. Tu, On multiplicative subgroups in division rings, *J. Algebra Appl.*, **15**(3) (2016) 1650050 (16 pages).
- [5] T. Y. Lam, *A First Course in Noncommutative Rings* (GMT **131**, Springer, 1991).
- [6] M. Mahdavi-Hezavehi, Free subgroups in maximal subgroups of $GL_1(D)$, *J. Algebra* **241** (2001) 720 – 730.
- [7] M. Mahdavi-Hezavehi, M. G. Mahmudi and S. Yasamin, Finitely generated subnormal subgroups of $GL_n(D)$ are central, *J. Algebra* **225** (2000) 517 – 521.
- [8] Derek J. S. Robinson, *A Course in the Theory of Groups* (Second Edition, Springer, 1995).
- [9] M. Shirvani and B. A. F. Wehrfritz, *Skew Linear Groups* (Cambridge Univ. Press, Cambridge, 1986).
- [10] J. Tits, Free subgroups in linear groups, *J. Algebra* **20** (1972) 250 – 270.
- [11] A. E. Zalesskii, The structure of several classes of matrix groups over a division ring (Russian), *Sibirsk. Mat. Z.* **8** (1967), 1284–1298; English translation in *Siberian Math. J.* **8** (1967) 978–988.

[1] UNIVERSITY OF SCIENCE, HO CHI MINH CITY, VIETNAM; [2] VIETNAM NATIONAL UNIVERSITY, HO CHI MINH CITY, VIETNAM.

E-mail address: bxhai@hcmus.edu.vn