

# Schmidt's subspace theorem for moving hypersurface targets in subgeneral position with index in algebraic variety

Tingbin Cao\* and Nguyen Van Thin†

## Abstract

Recently, Xie-Cao [15] obtained a Second Main Theorem for moving hypersurfaces located in subgeneral position with index which is extended the result of Ru [11]. By using some methods due to Son-Tan-Thin [13], Quang [9] and Xie-Cao [15], we shall give a Schmidt's Subspace Theorem for moving hypersurface targets in subgeneral position with index intersecting algebraic variety. Our result is an extension of the Schmidt's Subspace Theorem due to Son-Tan-Thin [13] and Quang [9].

*Keywords:* Diophantine approximation, Schmidt's Subspace Theorem.

Mathematics Subject Classification 2010. 11J68, 11J25, 11J97.

## 1 Some definitions and result

Let  $k$  be an algebraic number field of degree  $d$ . Denote  $M(k)$  by the set of places (i.e., equivalent classes of absolute values) of  $k$  and write  $M_\infty(k)$  for the set of Archimedean places. From  $v \in M(k)$ , we choose the normalized absolute value  $|\cdot|_v$  such that  $|\cdot|_v = |\cdot|$  on  $\mathbb{Q}$  (the standard absolute value) if  $v$  is archimedean, whereas for  $v$  non-archimedean  $|p|_v = p^{-1}$  if  $v$  lies above the rational prime  $p$ . Denote by  $k_v$  the completion of  $k$  with respect to  $v$  and by  $d_v = [k_v : \mathbb{Q}_v]$  the local degree. We put  $\| \cdot \|_v = |\cdot|_v^{d_v/d}$ . Then norm  $\| \cdot \|_v$  satisfies the following properties:

(i)  $\|x\|_v \geq 0$ , with equality if and only if  $x = 0$ ;

(ii)  $\|xy\|_v = \|x\|_v \|y\|_v$  for all  $x, y \in k$ ;

(iii)  $\|x_1 + \cdots + x_n\|_v \leq B_v^{n_v} \cdot \max\{\|x_1\|_v, \dots, \|x_n\|_v\}$  for all  $x_1, \dots, x_n \in k$ ,  $n \in \mathbb{N}$ ,

where  $n_v = d_v/d$ ,  $B_v = 1$  if  $v$  is non-archimedean and  $B_v = n$  if  $v$  is archimedean.

Moreover, for each  $x \in k \setminus \{0\}$ , we have the following product formula:

$$\prod_{v \in M(k)} \|x\|_v = 1.$$

---

\*The first author is supported by the National Natural Science Foundation of China (#11871260).

†Corresponding author: Nguyen Van Thin.

For  $v \in M(k)$ , we also extend  $\|\cdot\|_v$  to an absolute value on the algebraic closure  $\bar{k}_v$ .

For  $x \in k$ , the logarithmic height of  $x$  is defined by  $h(x) = \sum_{v \in M(k)} \log^+ \|x\|_v$ , where  $\log^+ \|x\|_v = \log \max\{\|x\|_v, 1\}$ . For  $\mathbf{x} = [x_0 : \cdots : x_M] \in \mathbb{P}^M(k)$ , we denote  $x = (x_0, \dots, x_M) \in k^{M+1}$ . We set  $\|x\|_v = \max_{0 \leq i \leq M} \|x_i\|_v$ , and define the logarithmic height of  $\mathbf{x}$  by

$$h(\mathbf{x}) = \sum_{v \in M(k)} \log \|x\|_v. \quad (1.1)$$

For a positive integer  $d$ , we set

$$\mathcal{T}_d := \{(i_0, \dots, i_M) \in \mathbb{N}_0^{M+1} : i_0 + \cdots + i_M = d\}.$$

Let  $Q$  be a homogeneous polynomial of degree  $d$  in  $k[x_0, \dots, x_M]$ . We write

$$Q = \sum_{I \in \mathcal{T}_d} a_I x^I.$$

A (fixed) hypersurface  $D$  of degree  $d$  in  $\mathbb{P}^M(k)$  is the zero set of a homogeneous polynomial  $Q$ , that is

$$D = \{(x_0 : \cdots : x_M) \in \mathbb{P}^M(k) : Q(x_0, \dots, x_M) = 0\}.$$

Set  $\|Q\|_v := \sum_{I \in \mathcal{T}_d} \|a_I\|_v$ . The height of  $Q$  (or  $D$ ) is denoted by  $h(Q)$  (or  $h(D)$ ) and given by

$$h(Q) := \sum_{v \in M(k)} \log \|Q\|_v.$$

Let  $D$  be a hypersurface which is defined by the zero of homogeneous polynomial  $Q$ . For each  $v \in M(k)$ , the Weil function  $\lambda_{D,v}$  is defined by

$$\lambda_{D,v}(\mathbf{x}) := \log \frac{\|x\|_v^d \cdot \|Q\|_v}{\|Q(x)\|_v}, \quad \mathbf{x} \in \mathbb{P}^M(k) \setminus D.$$

Let  $\Lambda$  be an infinite index set. We call a moving hypersurface  $D$  in  $\mathbb{P}^M(k)$  of degree  $d$ , indexed by  $\Lambda$  is collection of hypersurfaces  $\{D(\alpha)\}_{\alpha \in \Lambda}$  which are defined by the zero of homogeneous polynomials  $\{Q(\alpha)\}_{\alpha \in \Lambda}$  in  $k[x_0, \dots, x_M]$  respectively. Then, we can write  $Q = \sum_{I \in \mathcal{T}_d} a_I x^I$ , where  $a_I$ 's are functions from  $\Lambda$  into  $k$  having no common zeros points, and called it by moving homogeneous polynomial.

Throughout this paper, we consider an infinite index  $\Lambda$ ; a set  $\mathcal{Q} := \{D_1, \dots, D_q\}$  of moving hypersurfaces in  $\mathbb{P}^M(k)$ , indexed by  $\Lambda$ , which are defined by the zero of moving homogeneous polynomials  $\{Q_1, \dots, Q_q\}$  in  $k[x_0, \dots, x_M]$  respectively, an arbitrary projective variety  $V \subset \mathbb{P}^M(k)$  of dimension  $n$  generated by the homogeneous ideal  $\mathcal{I}(V)$ . We write

$$Q_j = \sum_{I \in \mathcal{T}_{d_j}} a_{j,I} x^I \quad (j = 1, \dots, q), \quad \text{where } d_j = \deg Q_j.$$

Let  $A \subset \Lambda$  be an infinite subset and denote by  $(A, a)$  each set-theoretic map  $a : A \rightarrow k$ . Denote by  $\mathcal{R}_A^0$  the set of equivalence classes of pairs  $(C, a)$ , where  $C \subset A$  is a subset with finite complement and  $a : C \rightarrow k$  is a map; and the equivalence relation is defined as follows:  $(C_1, a_1) \sim (C_2, a_2)$  if there exists  $C \subset C_1 \cap C_2$  such that  $C$  has finite complement in  $A$  and  $a_1|_C = a_2|_C$ . Then  $\mathcal{R}_A^0$  has an obvious ring structure. Moreover, we can embed  $k$  into  $\mathcal{R}_A^0$  as constant functions.

**Definition 1.1.** For each  $j \in \{1, \dots, q\}$ , we write  $\mathcal{T}_{d_j} = \{I_{j,1}, \dots, I_{j,M_{d_j}}\}$ , where  $M_{d_j} := \binom{d_j+M}{M}$ . A subset  $A \subset \Lambda$  is said to be coherent with respect to  $\mathcal{Q}$  if for every polynomial  $P \in k[x_{1,1}, \dots, x_{1,M_{d_1}}, \dots, x_{q,1}, \dots, x_{q,M_{d_q}}]$  that is homogeneous in  $x_{j,1}, \dots, x_{j,M_{d_j}}$  for each  $j = 1, \dots, q$ , either  $P(a_{1,I_{1,1}}(\alpha), \dots, a_{1,I_{1,M_{d_1}}}(\alpha), \dots, a_{q,I_{q,1}}(\alpha), \dots, a_{q,I_{q,M_{d_q}}}(\alpha))$  vanishes for all  $\alpha \in A$  or it vanishes for only finitely many  $\alpha \in A$ .

By [12, Lemma 2.1], there exists an infinite coherent subset  $A \subset \Lambda$  with respect to  $\mathcal{Q}$ . For each  $j \in \{1, \dots, q\}$ , we fix an index  $I_j \in \mathcal{T}_{d_j}$  such that  $a_{j,I_j} \not\equiv 0$  (this means that  $a_{j,I_j}(\alpha) \neq 0$  for all, but finitely many,  $\alpha \in A$ ), then  $\frac{a_{j,I}(\alpha)}{a_{j,I_j}(\alpha)}$  defines an element of  $\mathcal{R}_A^0$  for any  $I \in \mathcal{T}_{d_j}$ . This element given by the following function:

$$\{\alpha \in A : a_{j,I_j}(\alpha) \neq 0\} \rightarrow k, \quad \alpha \mapsto \frac{a_{j,I}(\alpha)}{a_{j,I_j}(\alpha)}.$$

Moreover, by coherent, the subring of  $\mathcal{R}_A^0$  generated over  $k$  by such all elements is an integral domain [8, page 3]. We define  $\mathcal{R}_{A,\mathcal{Q}}$  to be the field of fractions of that integral domain.

Denote by  $\mathcal{A}$  the set of all functions  $\{\alpha \in A : a_{j,I_j}(\alpha) \neq 0\} \rightarrow k$ ,  $\alpha \mapsto \frac{a_{j,I}(\alpha)}{a_{j,I_j}(\alpha)}$  and  $k_{\mathcal{Q}}$  the set of all formal finite sum  $\sum_{m=1}^s t_m \prod_{i=1}^s c_i^{n_i}$ , where  $t_m \in k$ ,  $c_i \in \mathcal{A}$ ,  $n_i \in \mathbb{N}$ .

Each pair  $(\widehat{b}, \widehat{c}) \in k_{\mathcal{Q}}^2$ , ( $\widehat{c}(\alpha) \neq 0$  for all, but finitely many,  $\alpha \in A$ ) defines a set-theoretic function, denoted by  $\frac{\widehat{b}}{\widehat{c}}$ , from  $\{\alpha : \widehat{c}(\alpha) \neq 0\}$  to  $k$ ,  $\frac{\widehat{b}}{\widehat{c}}(\alpha) := \frac{\widehat{b}(\alpha)}{\widehat{c}(\alpha)}$ . Denote by  $\widehat{\mathcal{R}}_{A,\mathcal{Q}}$  the set of all such functions. Each element  $a \in \mathcal{R}_{A,\mathcal{Q}}$  is a class of some functions  $\widehat{a}$  in  $\widehat{\mathcal{R}}_{A,\mathcal{Q}}$ . We call that  $\widehat{a}$  is a special representative of  $a$ . It is clear that for any two special representatives  $\widehat{a}_1, \widehat{a}_2$  of the same element  $a \in \mathcal{R}_{A,\mathcal{Q}}$ , we have  $\widehat{a}_1(\alpha) = \widehat{a}_2(\alpha)$  for all, but finitely many  $\alpha \in A$ . For a polynomial  $P := \sum_I a_I x^I \in \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$ , assume that  $\widehat{a}_I$  is a special representative of  $a_I$ . Then  $\widehat{P} := \sum_I \widehat{a}_I x^I$  is called a special representative of  $P$ . For each  $\alpha \in A$  such that all functions  $\widehat{a}_I$ 's are well defined at  $\alpha$ , we set  $\widehat{P}(\alpha) := \sum_I \widehat{a}_I(\alpha) x^I \in k[x_0, \dots, x_M]$ ; and we also say that the special representative  $\widehat{P}$  is well defined at  $\alpha$ . Note that each special representative  $\widehat{P}$  of  $P$  is well defined at all, but finitely many,  $\alpha \in A$ . If  $\widehat{P}_1, \widehat{P}_2$  are two special presentatives of  $P$ , then  $\widehat{P}_1(\alpha) = \widehat{P}_2(\alpha)$  for all, but finitely many  $\alpha$ .

**Definition 1.2.** A sequence of points  $x = [x_0 : \dots : x_M] : \Lambda \rightarrow V$  is said to be  $V$ -algebraically non-degenerate with respect to  $\mathcal{Q}$  if for each infinite coherent subset  $A \subset \Lambda$

with respect to  $\mathcal{Q}$ , there is no homogeneous polynomial  $P \in \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M] \setminus \mathcal{I}_{A,\mathcal{Q}}(V)$  such that  $\widehat{P}(\alpha)(x_0(\alpha), \dots, x_M(\alpha)) = 0$  for all, but finitely many,  $\alpha \in A$  for some (then for all) representative  $\widehat{P}$  of  $P$ , where  $\mathcal{I}_{A,\mathcal{Q}}(V)$  is the ideal in  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$  generated by  $\mathcal{I}(V)$ .

**Definition 1.3.** Let  $m \geq n$  be a positive integer. We say that the family moving hypersurfaces  $\mathcal{Q}$  is in  $m$ -strictly subgeneral position with respect to  $V$  if for any subset  $I \subset \{1, \dots, q\}$  with  $\#I \leq m + 1$ ,

$$\dim \left( \bigcap_{i \in I} D_i(\alpha) \cap V(\bar{k}) \right) \leq m - \#I$$

for all, but finitely many  $\alpha \in \Lambda$ , where  $\bar{k}$  is the algebraic closure of  $k$ .

From Definition 1.3, we get

**Remark 1.4.** Let  $m \geq n$  be a positive integer. The family moving hypersurfaces  $\mathcal{Q}$  is in  $m$ -strictly subgeneral position with respect to  $V$  implies that for each  $1 \leq j_0 < \dots < j_m \leq q, q \geq m + 1$ , the system of equations

$$Q_{j_i}(\alpha)(x_0, \dots, x_M) = 0, \quad 0 \leq i \leq m$$

has no solution  $(x_0, \dots, x_M)$  satisfying  $(x_0 : \dots : x_M) \in V(\bar{k})$ , for all, but finitely many  $\alpha \in \Lambda$ . It means that  $\mathcal{Q}$  is in  $m$ -subgeneral position with respect to  $V$ .

**Definition 1.5.** Let  $\kappa$  be a positive integer such that  $\kappa \leq \dim V$ . We say that the family moving hypersurfaces  $\mathcal{Q}$  is in  $m$ -strictly subgeneral position with index  $\kappa$  in  $V$  if  $\mathcal{Q}$  is in  $m$ -strictly subgeneral position with respect to  $V$  and for any subset  $I \subset \{1, \dots, q\}$  such that  $\#I \leq \kappa$ ,

$$\text{codim} \left( \bigcap_{i \in I} D_i(\alpha) \cap V(\bar{k}) \right) \geq \#I$$

for all, but finitely many  $\alpha \in \Lambda$ . Here we set  $\dim \emptyset = -\infty$ .

In 1997, Ru-Vojta [12] established the following Schmidt subspace theorem for the case of moving hyperplanes in projective spaces.

**Theorem A.** Let  $k$  be a number field and let  $S \subset M(k)$  be a finite set containing all archimedean places. Let  $\Lambda$  be an infinite index set and let  $\mathcal{H} := \{H_1, \dots, H_q\}$  be a set of moving hyperplanes in  $\mathbb{P}^M(k)$ , indexed by  $\Lambda$ . Let  $\mathbf{x} = [x_0 : \dots : x_M] : \Lambda \rightarrow \mathbb{P}^M(k)$  be a sequence of points. Assume that

- (i)  $\mathbf{x}$  is linearly nondegenerate with respect to  $\mathcal{H}$ , which mean, for each infinite coherent subset  $A \subset \Lambda$  with respect to  $\mathcal{H}$ ,  $x_0|_A, \dots, x_M|_A$  are linearly independent over  $\mathcal{R}_{A,\mathcal{H}}$ ,
- (ii)  $h(H_j(\alpha)) = o(h(\mathbf{x}(\alpha)))$  for all  $\alpha \in \Lambda$  and  $j = 1, \dots, q$  (i.e. for all  $\delta > 0$ ,  $h(H_j(\alpha)) \leq \delta h(\mathbf{x}(\alpha))$  for all, but finitely many,  $\alpha \in \Lambda$ ).

Then, for any  $\varepsilon > 0$ , there exists an infinite index subset  $A \subset \Lambda$  such that

$$\sum_{v \in S} \max_K \sum_{J \in K} \lambda_{H_j(\alpha), v}(\mathbf{x}(\alpha)) \leq (M + 1 + \varepsilon)h(\mathbf{x}(\alpha))$$

holds for all  $\alpha \in A$ . Here the maximum is taken over all subsets  $K$  of  $\{1, \dots, q\}$ ,  $\#K = M + 1$  such that  $H_j(\alpha), j \in K$  are linearly independent over  $k$  for each  $\alpha \in \Lambda$ .

One of the most important developments in recent years in Diophantine approximation is Schmidt's subspace theorems for fixed hypersurfaces of Corvaja-Zannier [4] and Evertse-Ferretti [6, 7]. Motivated by their paper, Ru [10, 11] obtained important results on the Second Main Theorem for fixed hypersurfaces. Later, Dethloff-Tan [5] generalized these Second Main Theorems of Ru to the case of moving results. Basing on method of Dethloff-Tan [5], Chen-Ru-Yan [2] and Le [8] extended Theorem A to the case of moving hypersurfaces in the projective spaces. In 2019, Dethloff-Tan [3] extended the result of Ru [11] for moving target. Via Vojta's dictionary, the Second Main Theorem in Nevanlinna theory corresponds to Schmidt's Subspace Theorem in Diophantine approximation. In 2018, Son-Tan-Thin [13] gave the counterpart of Second Main Theorem due to Dethloff-Tan [3] in Diophantine approximation. In 2018, Quang [9] obtained the Schmidt's subspace theorem for moving hypersurface in subgeneral position. In 2019, Xie-Cao [15] have been obtained the Second Main Theorem for moving hypersurfaces located in subgeneral position with index.

In this paper, by using some methods due to Son-Tan-Thin [13] and Quang [9], Xie-Cao [15], we give a Schmidt's Subspace Theorem for moving hypersurfaces in  $m$ -subgeneral position with index  $\kappa$  as follows:

**Theorem 1.6.** *Let  $k$  be a number field and let  $S \subset M(k)$  be a finite set containing all archimedean places. Let  $\mathbf{x} = [x_0 : \dots : x_M] : \Lambda \rightarrow V$  be a sequence of points. Assume that*

(i)  $\mathcal{Q}$  is in  $m$ -strictly subgeneral position with index  $\kappa$  in  $V$  and  $\mathbf{x}$  is  $V$ - algebraically nondegenerate with respect to  $\mathcal{Q}$ ;

par

(ii)  $h(D_j(\alpha)) = o(h(\mathbf{x}(\alpha)))$  for all  $\alpha \in \Lambda$  and  $j = 1, \dots, q$  (i.e. for all  $\delta > 0$ ,  $h(D_j(\alpha)) \leq \delta h(\mathbf{x}(\alpha))$  for all, but finitely many,  $\alpha \in \Lambda$ ).

Then, for any  $\varepsilon > 0$ , there exists an infinite index subset  $A \subset \Lambda$  such that

$$\sum_{v \in S} \sum_{j=1}^q d_j^{-1} \lambda_{D_j(\alpha), v}(\mathbf{x}(\alpha)) \leq \left( \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1) + \varepsilon \right) h(\mathbf{x}(\alpha))$$

holds for all  $\alpha \in A$ .

**Remark 1.7.** (i) By replacing  $Q_j$  by  $Q_j^{\frac{d}{d_j}}$ , where  $d = \text{lcm}\{d_1, \dots, d_q\}$ , in Theorem 1.6, we may assume that  $D_1, \dots, D_q$  have the same degree  $d$ .

(ii) By replacing  $Q_j = \sum_{I \in \mathcal{T}_d} a_{j,I} x^I$  by  $Q'_j = \sum_{I \in \mathcal{T}_d} \frac{a_{j,I}}{a_{j,I_j}} x^I$ , in Theorem 1.6, we may assume that  $Q_j \in \mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$ .

Our result is an extension of the Schmidt's Subspace Theorem due to Son-Tan-Thin [13] and Quang [9].

## 2 Some Lemmas

We write

$$Q_j = \sum_{I \in \mathcal{T}_d} a_{jI} x^I, \quad (j = 1, \dots, q).$$

Let  $A \subset \Lambda$  be an infinity coherent subset with respect to  $\mathcal{Q}$ . For each  $j \in \{1, \dots, q\}$ , we fix an index  $I_j \in \mathcal{T}_d$  such that  $a_{j, I_j} \neq 0$  (this means that  $a_{j, I_j}(\alpha) \neq 0$  for all but finitely many  $\alpha \in A$ ), then  $\frac{a_{jI}}{a_{jI_j}}$  defines an element of  $\mathcal{R}_A^0$  for any  $I \in \mathcal{T}_d$ . Set

$$Q'_j = \sum_{I \in \mathcal{T}_d} \frac{a_{jI}}{a_{jI_j}} x^I, \quad (j = 1, \dots, q).$$

Let  $t = (\dots, t_{jI}, \dots)$  be a family of variables. Set

$$\widetilde{Q}_j = \sum_{I \in \mathcal{T}_d} t_{jI} x^I \in k[t, x].$$

We have  $\widetilde{Q}_j(\dots, \frac{a_{jI}}{a_{jI_j}}(\alpha), \dots, x_0, \dots, x_M) = Q'_j(\alpha)(x_0, \dots, x_M)$  for all  $\alpha \in A$  outside a finite subset.

Assume that the ideal  $\mathcal{I}(V)$  of  $V$  is generated by homogeneous polynomials  $\mathcal{P}_1, \dots, \mathcal{P}_s$ . Since  $\mathcal{Q}$  is in  $m$ -subgeneral position on  $V$ , for each  $J := \{j_0, \dots, j_m\} \subset \{1, \dots, q\}$ , there is a subset  $A_J \subset A$  with finite complement such that for all  $\alpha \in A_J$ , the homogeneous polynomials

$$\mathcal{P}_1, \dots, \mathcal{P}_s, Q'_{j_0}(\alpha), \dots, Q'_{j_m}(\alpha) \in k[x_0, \dots, x_M]$$

have no common non-trivial solutions in  $\overline{k}^{M+1}$ . Denote by

$$k[t](\mathcal{P}_1, \dots, \mathcal{P}_s, \widetilde{Q}_{j_0}, \dots, \widetilde{Q}_{j_m})$$

the ideal in the ring of polynomials in  $x_0, \dots, x_M$  with coefficients in  $k[t]$  generated by  $\mathcal{P}_1, \dots, \mathcal{P}_s, \widetilde{Q}_{j_0}, \dots, \widetilde{Q}_{j_m}$ . A polynomial  $\widetilde{R}$  in  $k[t]$  is called an *inertia form* of the polynomials

$$\mathcal{P}_1, \dots, \mathcal{P}_s, \widetilde{Q}_{j_0}, \dots, \widetilde{Q}_{j_m}$$

if it has the following property:

$$x_i^N \cdot \widetilde{R} \in k[t](\mathcal{P}_1, \dots, \mathcal{P}_s, \widetilde{Q}_{j_0}, \dots, \widetilde{Q}_{j_m})$$

for  $i = 0, \dots, M$  and for some non-negative integer  $N$  (see e.g. [14]). It follows from the definition that the set  $\mathcal{I}$  of inertia forms of polynomials  $\mathcal{P}_1, \dots, \mathcal{P}_s, \tilde{Q}_{j_0}, \dots, \tilde{Q}_{j_m}$  is an ideal in  $k[t]$ .

It is well known that  $(s + m + 1)$  homogeneous polynomials

$$\mathcal{P}_i(x_0, \dots, x_M), \tilde{Q}_j(\dots, t_{jI}, \dots, x_0, \dots, x_M), i \in \{1, \dots, s\},$$

$j \in J$  have no common non-trivial solutions in  $x_0, \dots, x_M$  for special values  $t_{jI}^0$  of  $t_{jI}$  if and only if there exists an inertia form  $\tilde{R}_J^{t_{jI}^0}$  such that  $\tilde{R}_J^{t_{jI}^0}(\dots, t_{jI}^0, \dots) \neq 0$  (see e.g. [14, page 254]). For each  $\alpha \in A_J$ , choose  $\tilde{R}_J^\alpha \in \mathcal{I}$  with respect to the special values  $t_{jI}^\alpha := \frac{a_{jI}}{a_{jI_j}}(\alpha)$ . Set  $R_J^\alpha := \tilde{R}_J^\alpha(\dots, \frac{a_{jI}}{a_{jI_j}}, \dots)$ . Then  $R_J^\alpha$  is a special representative of an element  $\mathcal{R}_{A, \mathcal{Q}}$ . By construction, we have

$$R_J^\alpha(\alpha) \neq 0 \quad \text{for all } \alpha \in A_J \text{ (and hence, for all, but finitely many, } \alpha \in A). \quad (2.1)$$

Since  $k[t]$  is Noetherian,  $\mathcal{I}$  is generated by finite polynomials  $\tilde{R}_{J_1}, \dots, \tilde{R}_{J_p}$ . For each  $\alpha$ , we write  $\tilde{R}_J^\alpha = \sum_{\ell=1}^p \tilde{G}_\ell^\alpha \tilde{R}_{J_\ell}$ ,  $\tilde{G}_\ell^\alpha \in k[t]$ . We have that  $G_\ell^\alpha := \tilde{G}_\ell^\alpha(\dots, \frac{a_{jI}}{a_{jI_j}}, \dots)$ , and  $R_{J_\ell} := \tilde{R}_{J_\ell}(\dots, \frac{a_{jI}}{a_{jI_j}}, \dots)$  are special representatives of elements in  $\mathcal{R}_{A, \mathcal{Q}}$ . It is clear that  $R_J^\alpha = \sum_{\ell=1}^p G_\ell^\alpha R_{J_\ell}$ . Hence, by (2.1), we have

$$0 \neq R_J^\alpha(\alpha) = \sum_{\ell=1}^p G_\ell^\alpha(\alpha) R_{J_\ell}(\alpha)$$

for all, but finitely many  $\alpha \in A$ . Therefore, there is  $\ell_0 \in \{1, \dots, p\}$  such that

$$R_{J_{\ell_0}}(\alpha) \neq 0 \quad \text{for all, but finitely many } \alpha \in A. \quad (2.2)$$

Furthermore, by the definition of the inertia forms, there are a non-negative integer  $N$ , polynomials  $b_{i\ell} \in \mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$  with  $\deg b_{i j_k} = N - d$  and  $\deg b_{i\ell} = N - \deg P_\ell$  such that

$$R_{J_{\ell_0}} \cdot x_i^N = \sum_{k=0}^m b_{i j_k} \cdot Q'_{j_k} + \sum_{\ell=1}^s b_{i\ell} \cdot \mathcal{P}_\ell, \quad (2.3)$$

for all  $0 \leq i \leq M$ .

Let  $\mathbf{x} : \Lambda \rightarrow V \subset P^M(k)$  be a map. A map  $(C, a) \in \mathcal{R}_A^0$  is called small with respect to  $\mathbf{x}$  if and only if

$$h(a(\alpha)) = o(h(\mathbf{x}(\alpha))),$$

which means that, for every  $\varepsilon > 0$ , there exists a subset  $C_\varepsilon \subset C$  with finite complement such that  $h(a(\alpha)) \leq \varepsilon h(\mathbf{x}(\alpha))$  for all  $\alpha \in C_\varepsilon$ . We denote by  $\mathcal{K}_\mathbf{x}$  the set of all such small

maps. Then,  $\mathcal{K}_x$  is subring of  $\mathcal{R}_A^0$ . It is not an entire ring, however, if  $(C, a) \in \mathcal{K}_x$  and  $a(\alpha) \neq 0$  for all but finitely  $\alpha \in C$ , then we have  $(C \setminus \{\alpha : a(\alpha) = 0\}, \frac{1}{a}) \in \mathcal{K}_x$ . Denote by  $\mathcal{C}_x$  the set of all positive functions  $g$  defined over  $\Lambda$  outside a finite subset of  $\Lambda$  such that

$$\log^+(g(\alpha)) = o(h(\mathbf{x}(\alpha))).$$

Then  $\mathcal{C}_x$  is a ring. Moreover, if  $(C, a) \in \mathcal{K}_x$ , then for every  $v \in M(k)$ , the function  $\|a\|_v : C \rightarrow \mathbb{R}^+$  given by  $\alpha \mapsto \|a(\alpha)\|_v$  belongs to  $\mathcal{C}_x$ . Furthermore, if  $(C, a) \in \mathcal{K}_x$  and  $a(\alpha) \neq 0$  for all but finitely  $\alpha \in C$ , the function  $g : \{\alpha | a(\alpha) \neq 0\} \mapsto \frac{1}{\|a(\alpha)\|_v}$  also belongs to  $\mathcal{C}_x$ .

From (2.2) and (2.3), similarly to Lemma 2.2 in [8], under the assumption of Theorem 1.6 and Remark 1.7, we have the following result.

**Lemma 2.1.** *Let  $A \subset \Lambda$  be coherent with respect to  $\mathcal{Q}$ . Suppose that  $\mathcal{Q}$  is in  $m$ -subgeneral position with respect to  $V$ . Then for each  $J \subset \{1, \dots, q\}$  with  $|J| = m + 1$ , there are functions  $l_{1,v}, l_{2,v} \in \mathcal{C}_x$  such that*

$$l_{2,v}(\alpha) \|x(\alpha)\|_v^d \leq \max_{j \in J} \|Q_j(\alpha)(x(\alpha))\|_v \leq l_{1,v}(\alpha) \|x(\alpha)\|_v^d$$

for all  $\alpha \in A$  outside finite subset and all  $v \in S$ .

For each positive integer  $\ell$  and for each vector sub-space  $W$  in  $k[x_0, \dots, x_M]$  (or in  $\mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$ ), we denote by  $W_\ell$  the vector space consisting of all homogeneous polynomials in  $W$  of degree  $\ell$  (and of the zero polynomial).

By the usual theory of Hilbert polynomials, for  $N \gg 0$ , we have

$$\begin{aligned} H_V(N) &:= \dim_k \frac{k[x_0, \dots, x_M]_N}{\mathcal{I}(V)_N} \\ &= \dim_{\bar{k}} \frac{\bar{k}[x_0, \dots, x_M]_N}{\mathcal{I}(V(\bar{k}))_N} = \deg V \cdot \frac{N^n}{n!} + O(N^{n-1}). \end{aligned}$$

Using the method in the proof of [9, Lemma 3.1] and [15, Lemma 3.3], we get the result as follow:

**Lemma 2.2.** *Let  $Q_1, \dots, Q_{m+1}$  be homogeneous polynomials in  $\mathcal{R}_\Lambda^0[x_0, \dots, x_M]$  of the same degree  $d \geq 1$ , in  $m$ -subgeneral position with index  $\kappa$  in  $V$ . Then for an infinitely subset  $A \subset \Lambda$  which is coherent with respect to  $\{Q_1, \dots, Q_{m+1}\}$ , there exists  $n$  homogeneous polynomials  $P_1 = Q_1, \dots, P_\kappa = Q_\kappa, P_{\kappa+1}, \dots, P_{n+1}$  in  $\mathcal{R}_\Lambda^0[x_0, \dots, x_M]$  of the form*

$$P_t = \sum_{j=\kappa+1}^{m-n+t} c_{tj} Q_j, \quad c_{tj} \in k, \quad t = \kappa + 1, \dots, n + 1,$$

such that the system of equations

$$P_t(\alpha)(x_0, \dots, x_M) = 0, \quad 1 \leq t \leq n+1$$

has no solution  $(x_0, \dots, x_M)$  satisfying  $(x_0 : \dots : x_M) \in V(\bar{k})$ , for all  $\alpha \in \Lambda$  outside a finite subset of  $\Lambda$ .

*Proof.* We assume that  $Q_i$  ( $1 \leq i \leq m+1$ ) has the following form:

$$Q_i = \sum_{I \in \mathcal{T}_d} a_{i,I} x^I.$$

By the definition of  $m$ -strictly subgeneral position, there exists an infinite subset  $A'$  of  $A$  with finite complement such that, for all  $\alpha \in A'$ , the system of equations

$$Q_i(\alpha)(x_0, \dots, x_M) = 0, \quad 1 \leq i \leq m+1,$$

has only the trivial solution  $(0, \dots, 0)$  in  $\bar{k}$ . We note that  $A'$  is coherent with respect to  $\{Q_i\}_{i=1}^{m+1}$ . Replacing  $A$  by  $A'$  if necessary, we may assume that  $A' = A$ . We fix  $\alpha_0 \in A$ . For each homogeneous polynomial  $Q \in k[x_0, \dots, x_M]$ , we will denote

$$D^* = \{(x_0 : \dots : x_M) \in \mathbb{P}^M(\bar{k}) : Q(x_0, \dots, x_M) = 0\}.$$

Setting  $P_1 = Q_1, \dots, P_\kappa = Q_\kappa$ , we build  $P_{\kappa+1}, \dots, Q_{n+1}$  as follows. First we see that

$$\dim \left( \bigcap_{i=1}^t D_i^*(\alpha_0) \cap V(\bar{k}) \right) \leq m - t \text{ for } t = m - n + \kappa + 1, \dots, m + 1,$$

where  $\dim \emptyset = -\infty$  and for any  $1 \leq l \leq \kappa$ , we have

$$\dim \left( \bigcap_{j=1}^l D_j^*(\alpha_0) \cap V(\bar{k}) \right) = n - l.$$

Indeed, it is obvious  $\dim \left( \bigcap_{j=1}^\kappa D_j^*(\alpha_0) \cap V(\bar{k}) \right) \geq n - \kappa$ , and by the definition of  $m$ -strictly subgeneral position with index  $\kappa$ , we get  $\dim \left( \bigcap_{j=1}^l D_j^*(\alpha_0) \cap V(\bar{k}) \right) \leq n - l$ . In particular, we have

$$\dim \left( \bigcap_{j=1}^\kappa D_j^*(\alpha_0) \cap V(\bar{k}) \right) = n - \kappa.$$

**Step 1.** We will construct  $P_{\kappa+1}$  as follows. For each irreducible component  $\Gamma$  of dimension  $n - \kappa$  of  $\bigcap_{j=1}^\kappa D_j^*(\alpha_0) \cap V(\bar{k})$ , we put

$$V_{\Gamma} = \left\{ c = (c_{\kappa+1}, \dots, c_{m-n+\kappa+1}) \in k^{m-n+1}, \Gamma \subset D_c^*(\alpha_0), \text{ where } Q_c = \sum_{j=\kappa+1}^{m-n+\kappa+1} c_j Q_j \right\}.$$

Then  $V_{1\Gamma}$  is a subspace of  $k^{m-n+1}$ . Since

$$\dim \left( \bigcap_{i=1}^{m-n+\kappa+1} D_i^*(\alpha_0) \cap V(\bar{k}) \right) \leq n - \kappa - 1,$$

there exists  $i \in \{\kappa+1, \dots, m-n+\kappa+1\}$  such that  $\Gamma \not\subset D_i^*(\alpha_0)$ . Since the set of irreducible components of dimension  $n - \kappa$  of  $\bigcap_{j=1}^{\kappa} D_j^*(\alpha_0) \cap V(\bar{k})$  is at most countable, we have

$$k^{m-n+\kappa} \setminus \bigcup_{\Gamma} V_{1\Gamma} \neq \emptyset.$$

Hence there exists  $(c_{1(\kappa+1)}, \dots, c_{1(m-n+\kappa+1)}) \in k^{m-n+1}$  such that  $\Gamma \not\subset D_{\kappa+1}^*(\alpha_0)$  for all irreducible components of dimension  $n - \kappa$  of  $\bigcap_{j=1}^{\kappa} D_j^*(\alpha_0) \cap V(\bar{k})$ , where

$$P_{\kappa+1} = \sum_{j=\kappa+1}^{m-n+\kappa+1} c_{1(j)} Q_j.$$

This clearly implies that

$$\dim \left( \bigcap_{i=1}^{\kappa+1} D_i^*(\alpha) \cap V(\bar{k}) \right) \leq n - \kappa - 1. \quad (2.4)$$

Certainly, we have

$$\dim \left( \bigcap_{i=1}^{\kappa+1} D_i^*(\alpha) \cap V(\bar{k}) \right) \geq n - \kappa - 1. \quad (2.5)$$

From (2.4) and (2.5), we get

$$\dim \left( \bigcap_{i=1}^{\kappa+1} D_i^*(\alpha) \cap V(\bar{k}) \right) = n - \kappa - 1. \quad (2.6)$$

**Step 2.** We will construct  $P_{\kappa+2}$  as follows. For each irreducible component  $\Gamma'$  of dimension  $n - \kappa - 1$  of  $\bigcap_{j=1}^{\kappa+1} D_j^*(\alpha_0) \cap V(\bar{k})$ , we put

$$V_{1\Gamma'} = \left\{ c = (c_{\kappa+1}, \dots, c_{m-n+\kappa+2}) \in k^{m-n+2}, \Gamma' \subset D_c^*(\alpha_0), \text{ where } Q_c = \sum_{j=\kappa+1}^{m-n+\kappa+2} c_j Q_j \right\}.$$

Then  $V_{1\Gamma'}$  is a subspace of  $k^{m-n+2}$ . Since

$$\dim \left( \bigcap_{i=1}^{m-n+\kappa+2} D_i^*(\alpha_0) \cap V(\bar{k}) \right) \leq n - \kappa - 2,$$

there exists  $i \in \{\kappa+1, \dots, m-n+\kappa+2\}$  such that  $\Gamma \not\subset D_i^*(\alpha_0)$ . Since the set of irreducible components of dimension  $n - \kappa - 1$  of  $\bigcap_{j=1}^{\kappa+1} D_j^*(\alpha_0) \cap V(\bar{k})$  is at most countable, we have

$$k^{m-n+\kappa+1} \setminus \bigcup_{\Gamma'} V_{1\Gamma'} \neq \emptyset.$$

Hence there exists  $(c_{2(\kappa+1)}, \dots, c_{2(m-n+\kappa+2)}) \in k^{m-n+2}$  such that  $\Gamma' \not\subset D_{\kappa+2}^*(\alpha_0)$  for all irreducible components of dimension  $n - \kappa - 1$  of  $\bigcap_{j=1}^{\kappa} D_j^*(\alpha_0) \cap V(\bar{k})$ , where

$$P_{\kappa+2} = \sum_{j=\kappa+1}^{m-n+\kappa+2} c_{2(j)} Q_j.$$

This clearly implies that

$$\dim \left( \bigcap_{i=1}^{\kappa+2} D_i^*(\alpha_0) \cap V(\bar{k}) \right) \leq n - \kappa - 2$$

and we also have

$$\dim \left( \bigcap_{i=1}^{\kappa+2} D_i^*(\alpha_0) \cap V(\bar{k}) \right) = n - \kappa - 2.$$

Repeat again the above steps, after  $(n + 1 - \kappa)$ -th step, we get the hypersurface  $P_{\kappa+1}, \dots, P_{n+1}$  satisfying

$$\dim \left( \bigcap_{i=1}^t D_i^*(\alpha_0) \cap V(\bar{k}) \right) \leq n - t$$

for all  $t = \kappa + 1, \dots, n + 1$ . In particular,

$$\bigcap_{i=1}^{n+1} D_i^*(\alpha_0) \cap V(\bar{k}) = \emptyset. \quad (2.7)$$

We denote  $R$  by the inertia form of  $\mathcal{P}_1, \dots, \mathcal{P}_s, \tilde{P}_1, \dots, \tilde{P}_{n+1}$ . As in (2.2) and (2.7), we see that  $R(\alpha_0) \neq 0$ . By the property coherent of  $A$ , it implies that  $R(\alpha) \neq 0$  for all  $\alpha \in A$  outside a finite set. By property of inertia form, we have

$$\bigcap_{i=1}^{n+1} D_i^*(\alpha) \cap V(\bar{k}) = \emptyset \quad (2.8)$$

for all  $\alpha \in A$  outside a finite set.  $\square$

Note that  $c_{tj}$  does not depend on  $\Lambda$  for all  $t = 2, \dots, n + 1$  and  $j = 2, \dots, m + 1$ , then  $P_t[x_0, \dots, x_M] \in \mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$  for  $t = 1, \dots, n + 1$ . We denote  $\mathcal{I}_{A, \mathcal{Q}}(V)$  the ideal in  $\mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$  generated by the elements in  $\mathcal{I}(V)$ . It is clear that  $\mathcal{I}_{A, \mathcal{Q}}(V)$  is also the sub-vector space of  $\mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$  generated by  $\mathcal{I}(V)$ .

We use the lexicographic order in  $\mathbb{N}_0^n$  and for  $I = (i_1, \dots, i_n)$ , set  $\|I\| := i_1 + \dots + i_n$ .

**Definition 2.3.** For each  $I = (i_1, \dots, i_n) \in \mathbb{N}_0^n$  and  $N \in \mathbb{N}_0$  with  $N \geq d\|I\|$ , denote by  $\mathcal{L}_N^I$  the set of all  $\gamma \in \mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]_{N-d\|I\|}$  such that

$$P_1^{i_1} \dots P_n^{i_n} \gamma - \sum_{E=(e_1, \dots, e_n) > I} P_1^{e_1} \dots P_n^{e_n} \gamma_E \in \mathcal{I}_{A, \mathcal{Q}}(V)_N.$$

for some  $\gamma_E \in \mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]_{N-d\|E\|}$ .

Denote by  $\mathcal{L}^I$  the homogeneous ideal in  $\mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]$  generated by  $\bigcup_{N \geq d\|I\|} \mathcal{L}_N^I$ .

**Remark 2.4.** *i)  $\mathcal{L}_N^I$  is a  $\mathcal{R}_{A,\mathcal{Q}}$ -vector sub-space of  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_{N-d\|I\|}$ , and*

$$(\mathcal{I}(V), P_1, \dots, P_n)_{N-d\|I\|} \subset \mathcal{L}_N^I,$$

where  $(\mathcal{I}(V), P_1, \dots, P_n)$  is the ideal in  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$  generated by  $\mathcal{I}(V) \cup \{P_1, \dots, P_n\}$ .

*ii) For any  $\gamma \in \mathcal{L}_N^I$  and  $P \in \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_k$ , we have  $\gamma \cdot P \in \mathcal{L}_{N+k}^I$*

*iii)  $\mathcal{L}^I \cap \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_{N-d\|I\|} = \mathcal{L}_N^I$ .*

*iv)  $\frac{\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]}{\mathcal{L}^I}$  is a graded module over the graded ring  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$ .*

*v) If  $I_1 - I_2 = (i_{1,1} - i_{2,1}, \dots, i_{1,n} - i_{2,n}) \in N_0^n$ , then  $\mathcal{L}_N^{I_2} \subset \mathcal{L}_{N+d\|I_1\|-d\|I_2\|}^{I_1}$ . Hence  $\mathcal{L}^{I_2} \subset \mathcal{L}^{I_1}$ .*

Using Definition 2.3, by arguments as [13], we get the following result.

**Lemma 2.5.** [13] *For all  $N \gg 0$  divisible  $d$ , there are homogeneous polynomials  $\phi_1, \dots, \phi_{H_V(N)}$  in  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_N$  such that they form a basis of the  $\mathcal{R}_{A,\mathcal{Q}}$ -vector space  $\frac{\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_N}{\mathcal{I}_{A,\mathcal{Q}}(V)_N}$ , and*

$$\prod_{j=1}^{H_V(N)} \phi_j - (P_1 \cdots P_n)^{\frac{\deg V \cdot N^{n+1}}{d \cdot (n+1)!} - u(N)} \cdot P \in \mathcal{I}_{A,\mathcal{Q}}(V)_N,$$

where  $u(N)$  is a function satisfying  $u(N) \leq O(N^n)$ ,  $P \in \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$  is a homogeneous polynomials of degree

$$N \cdot H_V(N) - \frac{n \cdot \deg V \cdot N^{n+1}}{(n+1)!} + u(N) = \frac{\deg V \cdot N^{n+1}}{(n+1)!} + O(N^n).$$

### 3 Proof of Theorem 1.6

*Proof.* By [2, Lemma 2.1], there exists an infinite index subset  $A \subset \Lambda$  which is coherent with respect to  $\mathcal{Q}$ . By Remark 1.7, we may assume that the polynomials  $Q_j$ 's have the same degree  $d \geq 1$  and their coefficients belong to the field  $\mathcal{R}_{A,\mathcal{Q}}$ . By the fact that for any infinite subset  $B$  of  $A$ , then  $B$  is also coherent with respect to  $\mathcal{Q}$  and  $\mathcal{R}_{B,\mathcal{Q}} \subset \mathcal{R}_{A,\mathcal{Q}}$ , in our proof, we may freely pass to infinite subsets. For simplicity, we still denote these infinite subsets by  $A$ . We set

$$\mathcal{J} = \{(i_1, \dots, i_{m+1}); 1 \leq i_j \leq q \text{ for } j = 1, \dots, m+1, i_1, \dots, i_{m+1} \text{ distinct}\}.$$

For each  $J = (i_1, \dots, i_{m+1}) \in \mathcal{J}$ , we denote by  $P_{J,1}, \dots, P_{J,n+1}$  the moving homogeneous polynomials obtained in Lemma 2.2 with respect to the family of moving hypersurfaces  $\{Q_{i_1}, \dots, Q_{i_{m+1}}\}$ . It is easy to see that, for each  $v \in S$ , there exists a positive function  $h_v \in \mathcal{C}_x$  such that

$$\|P_{J,t}(x(\alpha))\|_v \leq h_v(\alpha) \max_{1 \leq j \leq m-n+t} \|Q_{i_j}(x(\alpha))\|_v \quad (3.1)$$

for all  $\alpha \in A$  outside a finite subset, for all  $t = \kappa + 1, \dots, n + 1$ . Here, the function  $h_v$  may choose common for all  $J \in \mathcal{J}$ . From the assumption, for each  $a \in \mathcal{R}_{A, \mathcal{Q}}$ , and  $v \in M(k)$ , we have, for all  $\alpha \in A$ ,

$$\log \|a(\alpha)\|_v \leq \sum_{v \in M(k)} \log^+ \|a(\alpha)\|_v = h(a(\alpha)) \leq o(h(x(\alpha))). \quad (3.2)$$

For each  $v \in S$ , and  $\alpha \in A$ , there exist a subset  $J(v, \alpha) = \{i_1(v, \alpha), \dots, i_q(v, \alpha)\}$  of  $\{1, \dots, q\}$  such that

$$\|Q_{i_1(v, \alpha)}(\alpha)(x(\alpha))\|_v \leq \|Q_{i_2(v, \alpha)}(\alpha)(x(\alpha))\|_v \dots \leq \|Q_{i_q(v, \alpha)}(\alpha)(x(\alpha))\|_v.$$

We have  $J(v, \alpha) = (i_1(v, \alpha), \dots, i_{m+1}(v, \alpha)) \subset \mathcal{J}$ . Denote by  $P_{J(v, \alpha), 1}, \dots, P_{J(v, \alpha), n+1}$  by the moving homogeneous polynomials obtained in Lemma 2.2 and by  $D_{J(v, \alpha), j}$  the hypersurface defined by  $P_{J(v, \alpha), j}$ ,  $j = 1, \dots, n + 1$ . Note that, we may freely pass to infinite subsets, then we may assume  $D_{J(v, \alpha), 1}, \dots, D_{J(v, \alpha), n+1}$  are in general position with respective  $V$ . Apply to Lemma 2.1 for  $\mathcal{Q} = \{D_{J(v, \alpha), 1}, \dots, D_{J(v, \alpha), n+1}\}$ , and using (3.1), there exist functions  $g_{0, v}^{J(v, \alpha)}, g_v^{J(v, \alpha)} \in \mathcal{C}_{\mathbf{x}}$  (depend on  $l_{2, v}$ ) such that

$$\|x(\alpha)\|_v^d \leq g_{0, v}^{J(v, \alpha)}(\alpha) \max_{1 \leq j \leq n+1} \|P_{J(v, \alpha), j}(\alpha)(x(\alpha))\|_v \leq g_v^{J(v, \alpha)}(\alpha) \|Q_{i_{m+1}}(\alpha)(x(\alpha))\|_v. \quad (3.3)$$

Therefore, from (3.3), we get

$$\begin{aligned} & \prod_{i=1}^q \frac{\|x(\alpha)\|_v^d}{\|Q_{i_j}(\alpha)(x(\alpha))\|_v} \quad (3.4) \\ & \leq (g_v^{J(v, \alpha)})^{q-m}(\alpha) \prod_{j=1}^m \frac{\|x(\alpha)\|_v^d}{\|Q_{i_j(v, \alpha)}(\alpha)(x(\alpha))\|_v} \\ & = (g_v^{J(v, \alpha)})^{q-m}(\alpha) \frac{\|x(\alpha)\|_v^{md}}{\prod_{j=1}^{\kappa} \|Q_{i_j(v, \alpha)}(\alpha)(x(\alpha))\|_v \prod_{j=\kappa+1}^{m-n+\kappa} \|Q_{i_j(v, \alpha)}(\alpha)(x(\alpha))\|_v} \\ & \quad \times \frac{1}{\prod_{j=m-n+\kappa+1}^m \|Q_{i_j(v, \alpha)}(\alpha)(x(\alpha))\|_v} \\ & \leq (g_v^{J(v, \alpha)})^{q-m}(\alpha) h_v^{n-\kappa-1}(\alpha) \frac{\|x(\alpha)\|_v^{md}}{\prod_{j=1}^n \|P_{J(v, \alpha), j}(\alpha)(x(\alpha))\|_v \prod_{j=\kappa+1}^{m-n+\kappa} \|Q_{i_j(v, \alpha)}(\alpha)(x(\alpha))\|_v}. \end{aligned}$$

From (3.4), we get

$$\sum_{i=1}^q \lambda_{D_{i_j}(\alpha), v}(\mathbf{x}(\alpha)) \leq \sum_{j=1}^n \lambda_{D_{J(v, \alpha), j}, v}(\mathbf{x}(\alpha)) + \sum_{j=1}^{m-n} \lambda_{D_{J(v, \alpha), j}, v}(\mathbf{x}(\alpha)) + o(h(\mathbf{x}(\alpha))) \quad (3.5)$$

if  $m - n \leq \kappa$  and

$$\begin{aligned}
& \sum_{i=1}^q \lambda_{D_i(\alpha),v}(\mathbf{x}(\alpha)) \\
& \leq \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + \sum_{j=\kappa+1}^{m-n+\kappa} \lambda_{D_{i_j(v,\alpha),v(\alpha)}}(\mathbf{x}(\alpha)) \\
& \leq \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + \sum_{j=\kappa+1}^{m-n+\kappa} \lambda_{D_{i_j(v,\alpha),v(\alpha)}}(\mathbf{x}(\alpha)) + o(h(\mathbf{x}(\alpha))) \\
& \leq \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + \frac{m-n}{\kappa} \sum_{j=\kappa+1}^{2\kappa} \lambda_{D_{i_j(v,\alpha),v(\alpha)}}(\mathbf{x}(\alpha)) + o(h(\mathbf{x}(\alpha))) \\
& = \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + \frac{m-n}{\kappa} \sum_{j=1}^{\kappa} \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + o(h(\mathbf{x}(\alpha))) \tag{3.6}
\end{aligned}$$

if  $m - n \geq \kappa$ . Combine (3.5) and (3.6), we get

$$\begin{aligned}
\sum_{i=1}^q \lambda_{D_i(\alpha),v}(\mathbf{x}(\alpha)) & \leq \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)) + o(h(\mathbf{x}(\alpha))) \\
& \quad + \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} \sum_{j=1}^n \lambda_{D_{J(v,\alpha),j,v}}(\mathbf{x}(\alpha)). \tag{3.7}
\end{aligned}$$

From (3.7), we obtain

$$\begin{aligned}
& \log \prod_{j=1}^q \frac{\|x(\alpha)\|_v^d}{\|Q_j(\alpha)(x(\alpha))\|_v} \\
& \leq \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \log \prod_{j=1}^n \frac{\|x(\alpha)\|_v^d}{\|P_{J(v,\alpha),j}(\alpha)(x(\alpha))\|_v} + o(h(\mathbf{x}(\alpha))).
\end{aligned}$$

This implies

$$\begin{aligned}
& \log \prod_{j=1}^q \|Q_j(\alpha)(x(\alpha))\|_v \tag{3.8} \\
& \geq \left( q - \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) n \right) d \log \|x(\alpha)\|_v \\
& \quad + \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \log \prod_{j=1}^n \|P_{J(v,\alpha),j}(\alpha)(x(\alpha))\|_v + o(h(\mathbf{x}(\alpha))).
\end{aligned}$$

By Lemma 2.5, there exist homogeneous polynomials  $\phi_1^{J(v,\alpha)}, \dots, \phi_{H_V(N)}^{J(v,\alpha)}$  (depend on  $J(v,\alpha)$ ) in  $\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_N$  and functions  $u(N), v(N)$  (common for all  $J(v,\alpha)$ ) such

that  $\{\phi_i^{J(v,\alpha)}\}$  a basis of  $\mathcal{R}_{A,\mathcal{Q}}$ - vector space  $\frac{\mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]_N}{\mathcal{I}_{A,\mathcal{Q}}(V)_N}$  and

$$\prod_{\ell=1}^{H_V(N)} \phi_\ell^{J(v,\alpha)} - (P_{J(v,\alpha),1} \dots P_{J(v,\alpha),n}) \frac{\deg V \cdot N^{n+1}}{d(n+1)!}^{-u(N)} P_{J(v,\alpha)} \in \mathcal{I}_{A,\mathcal{Q}}(V)_N,$$

where  $P_{J(v,\alpha)} \in \mathcal{R}_{A,\mathcal{Q}}[x_0, \dots, x_M]$  is a homogeneous polynomials of degree  $\frac{\deg V \cdot N^{n+1}}{(n+1)!} + v(N)$ ,  $v(N) = O(N^n)$ . Thus, for all  $\mathbf{x}(\alpha) \in V(k)$ , we have

$$\prod_{\ell=1}^{H_V(N)} \phi_\ell^{J(v,\alpha)}(\alpha)(x(\alpha)) = \left( \prod_{i=1}^n P_{J(v,\alpha),i}(\alpha)(x(\alpha)) \right) \frac{\deg V \cdot N^{n+1}}{d(n+1)!}^{-u(N)} P_{J(v,\alpha)}(\alpha)(x(\alpha)).$$

On the other hand, it is easy to see that there exist  $h_{J(v,\alpha)} \in \mathcal{C}_{\mathbf{x}}$  such that

$$\begin{aligned} & \|P_{J(v,\alpha)}(\alpha)(x(\alpha))\|_v \\ & \leq \|x(\alpha)\|_v^{\deg P_{J(v,\alpha)}} h_{J(v,\alpha)}(\alpha) \\ & = \|x(\alpha)\|_v^{\frac{\deg V \cdot N^{n+1}}{(n+1)!} + v(N)} h_{J(v,\alpha)}(\alpha). \end{aligned}$$

Therefore,

$$\begin{aligned} & \log \prod_{\ell=1}^{H_V(N)} \|\phi_\ell^{J(v,\alpha)}(\alpha)(x(\alpha))\|_v \\ & \leq \left( \frac{\deg V \cdot N^{n+1}}{d(n+1)!} - u(N) \right) \cdot \log \left\| \prod_{i=1}^n P_{J(v,\alpha),i}(\alpha)(x(\alpha)) \right\|_v \\ & \quad + \log^+ h_{J(v,\alpha)}(\alpha) + \left( \frac{\deg V \cdot N^{n+1}}{(n+1)!} + v(N) \right) \log \|x(\alpha)\|_v \end{aligned}$$

This implies that there are functions  $\omega_1(N), \omega_2(N) \leq O(\frac{1}{N})$  such that

$$\begin{aligned} & \log \left\| \prod_{i=1}^n P_{J(v,\alpha),i}(\alpha)(x(\alpha)) \right\|_v \\ & \geq \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \cdot \log \prod_{\ell=1}^{H_V(N)} \|\phi_\ell^{J(v,\alpha)}(\alpha)(x(\alpha))\|_v \\ & \quad - \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \log^+ h_{J(v,\alpha)}(\alpha) - (d + \omega_2(N)) \log \|x(\alpha)\|_v. \quad (3.9) \end{aligned}$$

By (3.8) and (3.9), we have

$$\begin{aligned}
& \log \prod_{j=1}^q \|Q_j(\alpha)(x(\alpha))\|_v \tag{3.10} \\
& \geq \left( q - \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1) \right) d \log \|x(\alpha)\|_v \\
& \quad + \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \right. \\
& \quad \times \log \prod_{\ell=1}^{H_V(N)} \|\phi_\ell^{J(v,\alpha)}(\alpha)(x(\alpha))\|_v \\
& \quad \left. - \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \log^+ h_{J(v,\alpha)}(\alpha) - \omega_2(N) \log \|x(\alpha)\|_v \right) + o(h(\mathbf{x}(\alpha))).
\end{aligned}$$

We fix homogeneous polynomials  $\Phi_1, \dots, \Phi_{H_V(N)} \in \mathcal{R}_{A, \{Q_j\}_{j=1}^q} [x_0, \dots, x_M]_N$  such that they form a basis of  $\mathcal{R}_{A, \mathcal{Q}}$ -vector space  $\frac{\mathcal{R}_{A, \mathcal{Q}}[x_0, \dots, x_M]_N}{\mathcal{I}_{A, \mathcal{Q}}(V)_N}$ . Then, there exist homogeneous linear polynomials

$$L_1^{J(v,\alpha)}, \dots, L_{H_V(N)}^{J(v,\alpha)} \in \mathcal{R}_{A, \mathcal{Q}}[y_1, \dots, y_{H_V(N)}]$$

such that they are linear independent over  $\mathcal{R}_{A, \{Q_j\}_{j=1}^q}$  and

$$\phi_\ell^{J(v,\alpha)} - L_\ell^{J(v,\alpha)}(\Phi_1, \dots, \Phi_{H_V(N)}) \in \mathcal{I}_{A, \mathcal{Q}}(V)_N,$$

for all  $\ell = 1, \dots, H_V(N)$ . It is clear that  $h(L_\ell^{J(v,\alpha)}(\beta)) = o(h(x(\beta)))$ ,  $\beta \in A$  and  $A$  is coherent with respect to  $\{L_\ell\}_{\ell=1}^{H_V(N)}$ .

We have,

$$\prod_{\ell=1}^{H_V(N)} \|\phi_\ell^{J(v,\alpha)}(\alpha)(x(\alpha))\|_v = \prod_{\ell=1}^{H_V(N)} \|L_\ell^{J(v,\alpha)}(\Phi_1, \dots, \Phi_{H_V(N)})(\alpha)(x(\alpha))\|_v. \tag{3.11}$$

We write

$$L_\ell^{J(v,\alpha)}(y_1, \dots, y_{H_V(N)}) = \sum_{s=1}^{H_V(N)} g_{\ell s} y_s, \quad g_{\ell s} \in \mathcal{R}_{A, \mathcal{Q}}.$$

Since  $L_1^{J(v,\alpha)}, \dots, L_{H_V(N)}^{J(v,\alpha)}$  are linear independent over  $\mathcal{R}_{A, \mathcal{Q}}$ , we have  $\det(g_{\ell s}) \neq 0 \in \mathcal{R}_{A, \mathcal{Q}}$ . Thus, due to coherent property of  $A$ ,  $\det(g_{\ell s})(\beta) \neq 0$  for all  $\beta \in A$ , outside a finite subset of  $A$ . By passing to an infinite subset if necessary, we may assume that  $L_1^{J(v,\alpha)}(\beta), \dots, L_{H_V(N)}^{J(v,\alpha)}(\beta)$  are linearly independent over  $k$  for all  $\beta \in A$ .

Now we consider the sequence of points  $F(\alpha) = [\Phi_1(x(\alpha)), \dots, \Phi_{H_V(N)}(x(\alpha))]$  from  $A$  to  $\mathbb{P}^{H_V(N)-1}(k)$  and moving hyperplanes  $\mathcal{L} := \{L_1^{J(v,\alpha)}, \dots, L_{H_V(N)}^{J(v,\alpha)}\}$  in  $\mathbb{P}^{H_V(N)-1}(k)$ , indexed

by  $A$ . We claim that  $F$  is linearly nondegenerate with respect to  $\mathcal{L}$ . Indeed, otherwise, then there is a linear form  $L \in \mathcal{R}_{B,\mathcal{L}}[y_1, \dots, y_{H_N(V)}]$  for some infinite coherent subset  $B \subset A$ , such that  $L(F)|_B \equiv 0$  in  $B$ , which contradicts to the assumption that  $x$  is algebraically nondegenerate with respect to  $\mathcal{Q}$ .

By Theorem A, for any  $\epsilon > 0$ , there is an infinite subset of  $A$  (common for all  $J(v, \alpha)$ ), denoted again by  $A$ , such that

$$\sum_{v \in S} \log \prod_{\ell=1}^{H_V(N)} \frac{\|F(\alpha)\|_v \|L_\ell^{J(v,\alpha)}(\alpha)\|_v}{\|L_\ell^{J(v,\alpha)}(\alpha)(F(\alpha))\|_v} \leq (H_V(N) + \epsilon)h(F(\alpha)), \quad (3.12)$$

for all  $\alpha \in A$ .

Combining with (3.10) and (3.12) we have

$$\begin{aligned} & \sum_{v \in S} \sum_{j=1}^q \log \frac{\|x(\alpha)\|_v^d}{\|Q_j(\alpha)(x(\alpha))\|_v} \\ & \leq \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1)d \sum_{v \in S} \log \|x(\alpha)\|_v + \omega_2(N) \sum_{v \in S} \log \|x(\alpha)\|_v \\ & \quad + \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \\ & \quad \times \sum_{v \in S} \log \prod_{\ell=1}^{H_V(N)} \frac{\|F(\alpha)\|_v \|L_\ell^{J(v,\alpha)}(\alpha)\|_v}{\|L_\ell^{J(v,\alpha)}(\alpha)(x(\alpha))\|_v} \\ & \quad - H_V(N) \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) \sum_{v \in S} \log \|F(\alpha)\|_v \\ & \quad + o(h(x(\alpha))). \end{aligned} \quad (3.13)$$

Since the above inequality is independent of the choice of components of  $\mathbf{x}(\alpha)$ , we can choose the components of  $\mathbf{x}(\alpha)$  being  $S$ -integers so that

$$\begin{aligned} \sum_{v \in S} \log \|x(\alpha)\|_v & = h(\mathbf{x}(\alpha)) + O(1), \text{ and} \\ \sum_{v \in S} \log \|F(\alpha)\|_v & \leq h(F(\alpha)) + O(1) \leq Nh(\mathbf{x}(\alpha)) + O(1). \end{aligned} \quad (3.14)$$

Combining with (3.13) and (3.14), we obtain

$$\begin{aligned}
& \sum_{v \in S} \sum_{j=1}^q \log \frac{\|x(\alpha)\|_v^d}{\|Q_j(\alpha)(x(\alpha))\|_v} \\
& \leq \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1) dh(\mathbf{x}(\alpha)) + \omega_2(N) h(\mathbf{x}(\alpha)) \\
& \quad + \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) (H_V(N) + \epsilon) h(F(\alpha)) \\
& \quad - H_V(N) \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) h(F(\alpha)) \\
& \quad + o(h(\mathbf{x}(\alpha))).
\end{aligned}$$

Hence, we obtain

$$\begin{aligned}
& \sum_{v \in S} \sum_{j=1}^q \log \frac{\|x(\alpha)\|_v^d \|Q_j(\alpha)\|_v}{\|Q_j(\alpha)(x(\alpha))\|_v} \tag{3.15} \\
& \leq \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1) dh(\mathbf{x}(\alpha)) + \omega_2(N) h(\mathbf{x}(\alpha)) \\
& \quad + \epsilon \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) \left( \frac{d(n+1)!}{\deg V \cdot N^{n+1}} + \omega_1(N) \right) h(F(\alpha)) + o(h(\mathbf{x}(\alpha))).
\end{aligned}$$

Here, we note that  $h(D_j(\alpha)) = o(h(\mathbf{x}(\alpha)))$  and there is only finite points  $\mathbf{x}$  such that  $h(\mathbf{x})$  is bounded, combining (3.14) and (3.15), for any  $\epsilon > 0$ , by our choice with  $\omega_1, \omega_2$  for  $N$  large enough, we get

$$\sum_{v \in S} \sum_{j=1}^q \log \frac{\|x(\alpha)\|_v^d \|Q_j(\alpha)\|_v}{\|Q_j(\alpha)(x(\alpha))\|_v} \leq \left( \left( \frac{m-n}{\max\{1, \min\{m-n, \kappa\}\}} + 1 \right) (n+1) + \epsilon \right) dh(\mathbf{x}(\alpha)),$$

for all  $\alpha \in A$  outside finite set. This completes the proof of Theorem 1.6.  $\square$

## References

- [1] Z. Chen, M. Ru, and Q. Yan, *The degenerated second main theorem and Schmidt's subspace theorem*, Sci. China Math. **55** (2012), 1367-1380.
- [2] Z. Chen, M. Ru, and Q. Yan, *Schmidt's subspace theorem with moving targets*, Internat. Math. Res. Notices **15**(2015), no. 1, 6305-6329, 2015.
- [3] G. Dethloff, and T. V. Tan, *Holomorphic curves into algebraic varieties intersecting moving hypersurface targets*, To appear in Acta Math. Vietnamica (2019), Doi: 10.1007/s40306-019-00336-3

- [4] P. Corvaja and U. Zannier, *On a general Thue's equation*, Amer. J. Math. **126** (2004), 1033-1055.
- [5] G. Dethloff and T. V. Tan, *A second main theorem for moving hypersurface targets*, Houston J. Math. **37** (2011), 79-111.
- [6] J. H. Evertse and R. G. Ferretti, *Diophantine inequalities on projective varieties*, Internat. Math. Res. Notices **25**(2002), 1295-1330.
- [7] J. H. Evertse and R. G. Ferretti, *A generalization of the subspace theorem with polynomials of higher degree*, Developments in Mathematics, **16** (2008), 175-198, Springer-Verlag, New York.
- [8] G. Le, *Schmidt's subspace theorem with moving hypersurfaces*, Int. J. Number Theory **11**(2015), 139-158.
- [9] S. D. Quang, *Schmidt's subspace theorem for moving hypersurface in subgeneral position*, Int. J. Number Theory **14**(2018), 103-121, 2018.
- [10] M. Ru, *A defect relation for holomorphic curves intersecting hypersurfaces*, Amer. J. Math. **126**(2004), 215-226.
- [11] M. Ru, *Holomorphic curves into algebraic varieties*, Ann of Math. **169**(2009), 255-267.
- [12] M. Ru and P. Vojta, *Schmidt's subspace theorem with moving targets*, Invent. Math. **127**(1997), 51-65.
- [13] N. T. Son, T. V. Tan and N. V. Thin, *Schmidt's subspace theorem for moving hypersurface targets*, J. Number Theory **186**(2018), 346-369.
- [14] O. Zariski, *Generalized weight properties of the resultant of  $n + 1$  polynomials in  $n$  indeterminates*, Trans. AMS **41**(1937), 249-265.
- [15] L. B. Xie and T. B. Cao, *Second Main Theorem for holomorphic curves into algebraic varieties intersecting moving hypersurfaces targets*, Arxiv: 1908.05844v2.

Tingbin Cao  
 Department of Mathematics  
 Nanchang University  
 Jiangxi 330031, P. R. China  
 e-mail: tbcao@ncu.edu.cn

Nguyen Van Thin  
 Department of Mathematics  
 Thai Nguyen University of Education

Luong Ngoc Quyen Street, Thai Nguyen city, Vietnam.  
e-mail: thinmath@gmail.com