

HALF-AUTOMORPHISM GROUP OF CHEIN LOOPS

GILIARD SOUZA DOS ANJOS*

Instituto de Matemática e Estatística - Universidade de São Paulo
 Rua do Matão, 1010, 05508-090, São Paulo - SP, Brazil
 giliards@ime.usp.br

Abstract

A bijection f of a loop L is a *half-automorphism* if $f(xy) \in \{f(x)f(y), f(y)f(x)\}$, for any $x, y \in L$. A half-automorphism is *nontrivial* when it is neither an automorphism nor an anti-automorphism. A *Chein loop* $L = G \cup Gu$ is a Moufang loop constructed from a group G and an element u of order 2 outside of G . In this paper, the half-automorphism group of finite Chein loops is described.

Keywords: Chein loop, half-isomorphism, half-automorphism, half-automorphism group.

1 Introduction

A loop $(L, *)$ consists of a nonempty set L with a binary operation $*$ on L such that for each $a, b \in L$ the equations $a * x = b$ and $y * a = b$ have unique solutions, and there is an identity element $1 \in L$ such that $1 * x = x = x * 1$, for any $x \in L$.

A *Moufang loop* is a loop that satisfies the identity $x(y(xz)) = ((xy)x)z$. Every Moufang loop is *diassociative*, that is, any two of its elements generate a group. The fundamental definitions and facts from loop theory and Moufang loops are found in [1, 12].

Let G be a group and u be an indeterminate. On $M(G, 2) = G \cup Gu$, define the operation $*$ by

$$g * h = gh, \quad g * (hu) = (hg)u, \quad (gu) * h = (gh^{-1})u, \quad (gu) * (hu) = h^{-1}g \quad (1)$$

for any $g, h \in G$. Then, $(M(G, 2), *)$ is a Moufang loop. Loops of this type are known as *Chein loops* [3]. Note that $M(G, 2)$ is associative if and only if G is abelian.

Let $(L, *)$ and (L', \cdot) be loops. A bijection $f : L \rightarrow L'$ is called a *half-isomorphism* if $f(x * y) \in \{f(x) \cdot f(y), f(y) \cdot f(x)\}$, for any $x, y \in L$. A half-isomorphism of a loop onto itself is a *half-automorphism*. We say that a half-isomorphism is *trivial* when it is either an isomorphism or an anti-isomorphism.

In 1957, Scott [13] showed that every half-isomorphism on groups is trivial. Recently, similar versions of this result were obtained for more structured classes of nonassociative loops. Kinyon, Stuhl and Vojtěchovský [8] generalized Scott's result to a more general subclass of Moufang loops. Gagola and Giuliani [5] proved that every half-automorphism of a finite simple Moufang loop is trivial. In [10], the authors showed that in a subclass of automorphic loops of odd order

*This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001

every half-automorphism is trivial. An *automorphic loop* is a loop in which every inner mapping is an automorphism [2].

Gagola and Giuliani also investigated loops that have nontrivial half-automorphisms. In [4], these authors established conditions for the existence of nontrivial half-automorphisms for Chein loops. A similar work was made for certain automorphic loops of even order in [9], where the authors also described the half-automorphism group of these loops. This work continues the investigation of nontrivial half-automorphisms in Chein loops initiated by Gagola and Giuliani. The aim is to describe the half-automorphism group of Chein loops.

This paper is organized as follows: Section 2 presents the definitions and basic results about half-automorphisms on Chein loops. In section 3, some properties of an important subgroup of the half-automorphism group of Chein loops are obtained. In section 4, the half-automorphism group of finite Chein loops is described.

2 Preliminaries

Here, the required definitions and basic results on half-automorphisms of Chein loops are stated.

First, a description of the automorphism group of Chein loops made by Grishkov, Zavarnitsine and Giuliani [7] is presented. Let G be a group and $L = M(G, 2)$. Denote the automorphism group of G by $Aut(G)$. For $\phi \in Aut(G)$ and $t \in G$, define $a_\phi, d_t : L \rightarrow L$ by:

$$a_\phi(gu^i) = (\phi(g))u^i, \quad i \in \{0, 1\}, \quad d_t(gu^i) = \begin{cases} g, & i = 0, \\ (tg)u, & i = 1. \end{cases}$$

Let H and H' be groups and $\sigma : H \rightarrow Aut(H')$ be a homomorphism. The set $H \times H'$ with the operation $(x, y) \cdot (x', y') = (xx', y\sigma_x(y'))$, where $\sigma_x = \sigma(x)$, is a semidirect product between H and H' , and it is denoted by $H \overset{\sigma}{\ltimes} H'$. If $\pi : Aut(G) \rightarrow Aut(G)$ is the identity mapping, we can construct the semidirect product $Aut(H) \overset{\pi}{\ltimes} H$, which is called the *holomorph of G* and denoted by $Hol(G)$.

Proposition 2.1. ([7, Lemma 1]) *Let $A = \{a_\phi \mid \phi \in Aut(G)\}$ and $D = \{d_t \mid t \in G\}$. Then:*

- a) $A, D \leq Aut(L)$, and $A \cap D = \{I\}$, where I is the identity mapping of L ,
- b) $a_\phi a_{\phi'} = a_{\phi\phi'}$, $d_t d_{t'} = d_{tt'}$, $(a_\phi)^{-1} = a_{\phi^{-1}}$, $(d_t)^{-1} = d_{t^{-1}}$, and $a_\phi d_t = d_{\phi(t)} a_\phi$,
- c) $A \cong Aut(G)$, $D \cong G$, and $AD \cong Hol(G)$.

If there exists an abelian subgroup $G_0 \leq G$ of index 2 such that $G = G_0 \cup G_0v$, where $v \notin G_0$, $v^2 = 1$ and $vgv = g^{-1}$, for all $g \in G_0$, then G is a *generalized dihedral group*. In this case, $L = M(G, 2) = G_0K$, where $K = \{1, u, v, w = uv = vu\}$ is isomorphic to the Klein group. For $\alpha \in Aut(K) \cong S_3$, we can define an automorphism s_α of L by $s_\alpha(gx) = g\alpha(x)$, for any $x \in K, g \in G_0$. Define $S = \{s_\alpha \mid \alpha \in Aut(K)\}$. Thus, $S \cong S_3$.

Theorem 2.2. ([7, Theorem 1]) *Let G be a group. If G is not a generalized dihedral group, then $Aut(M(G, 2)) \cong Hol(G)$. If $G = G_0 \cup G_0v$ is a generalized dihedral group and G_0 is not a group of exponent 2, then $Aut(M(G, 2)) = ADS$.*

Note that if $G = G_0 \cup G_0v$ is a generalized dihedral group and G_0 is not a group of exponent 2, then $AD \cap S = \{I, d_v\}$. Thus, $|Aut(M(G, 2))| = 3|Hol(G)| = 3|Aut(G)||G|$.

Denote the sets of the half-automorphisms and trivial half-automorphisms of the loop L by $Half(L)$ and $Half_T(L)$, respectively.

In any loop Q , it is easy to see that the composition of two half-automorphisms is also a half-automorphism, the identity mapping of Q is a half-automorphism and the inverse mapping of a trivial half-automorphism is also a trivial half-automorphism. Then, $Half_T(Q)$ is always a group, and it is called the *trivial half-automorphism group* of Q .

Let f be a half-automorphism of a diassociative loop M . In [8, Lemma 2.1], it was proved that if $x, y \in M$ commute, then $f(x)$ and $f(y)$ commute. Thus, f^{-1} is a half-automorphism by [10, Theorem 2.5]. Hence, $Half(M)$ is a group, and it is called the *half-automorphism group* of M .

We conclude that $Half(L)$ and $Half_T(L)$ are groups for every Chein loop L .

In the following, some results about half-automorphisms in Chein loops obtained by Gagola and Giuliani in [4] are presented.

Denote the order of the element x of a group G by $o(x)$ and the center of G by $\mathcal{Z}(G)$.

Theorem 2.3. ([4, Theorem 1]) *Let G be a finite nonabelian group and $L = M(G, 2)$. If L has a nontrivial half-automorphism, then there exists an element $x \in G$ such that:*

- a) $o(x) = 4$,
- b) $x^2 \in \mathcal{Z}(G)$,
- c) $x^{-1}gx = g^{-1}$ for any element $g \in G$ such that $o(g) \notin \{2, 4\}$.

Let G be a group and $L = M(G, 2)$. Define the set

$$H = \{\varphi \in Half(L) \mid \varphi(u) = u \text{ and } \varphi(g) = g, \forall g \in G\}.$$

Then, H is a subgroup of $Half(L)$. Analyzing the proof of [4, Lemma 5], this lemma can be rewritten as follows.

Proposition 2.4. ([4, Lemma 5]) *$Half(L) = Half_T(L)H$. In particular, if L has a nontrivial half-automorphism, then there exists a nontrivial half-automorphism in H .*

In the proof of Theorem 2.3, the authors in [4] proved the following result about the elements of H .

Proposition 2.5. ([4]) *Let $\varphi \in H$ and $y \in G$. If $o(y) \neq 4$, then $\varphi(yu) = yu$.*

3 The properties of the elements of H

In this section, G is considered as a finite nonabelian group such that $L = M(G, 2)$ has a nontrivial half-automorphism.

Denote the identity mapping of L by I . By Proposition 2.4, there exists $\varphi \in H$ such that $\varphi \neq I$.

Proposition 3.1. *Let $\varphi \in H \setminus \{I\}$. Then φ is a nontrivial half-automorphism.*

Proof. Since G is nonabelian and $\varphi(g) = g$, for all $g \in G$, the mapping φ is not an anti-automorphism. If φ is an automorphism, then $\varphi(gu) = \varphi(g) * \varphi(u) = gu$, for all $g \in G$, and hence $\varphi = I$. \square

Let $\varphi \in H$. Since φ is a half-automorphism, we have $\varphi(gu) \in \{\varphi(g) * \varphi(u), \varphi(u) * \varphi(g)\}$, for all $g \in G$. Then,

$$\varphi(gu) \in \{gu, g^{-1}u\}, \text{ for all } g \in G. \tag{2}$$

Since φ is a bijection, it follows that $\varphi^2 = I$. Thus, H is an abelian group of exponent 2. Denote by C_2^n the direct product of n cyclic groups of order 2. Hence, the following result is at hand.

Proposition 3.2. $H \cong C_2^r$, for some $r \geq 1$.

Let $\varphi \in H \setminus \{I\}$. Define the set $\gamma_\varphi = \{x \in G \mid \varphi(xu) = x^{-1}u, o(x) = 4\}$. By Theorem 2.3, $\gamma_\varphi \neq \emptyset$.

Proposition 3.3. Let $\varphi \in H \setminus \{I\}$. Then, for all $g, h \in G$ and $x \in \gamma_\varphi$:

- (a) If $g \notin \gamma_\varphi$, then $\varphi(gu) = gu$, $g^{-1} = x^{-1}gx$ and $o(xg) = 4$,
- (b) $x^{-1} \in \gamma_\varphi$,
- (c) $x^2 \in \mathcal{Z}(G)$,
- (d) If $o(g) \neq 4$, then $\gamma_\varphi g = g\gamma_\varphi = \gamma_\varphi$,
- (e) If $g, h \notin \gamma_\varphi$ and $hg \in \gamma_\varphi$, then $o(g) = 4$.

Proof. (a) Let $g \in G \setminus \gamma_\varphi$. By Proposition 2.5 and (2), $\varphi(gu) = gu$. Then,

$$g^{-1}x = \varphi((xu) * (gu)) \in \{\varphi(xu) * \varphi(gu), \varphi(gu) * \varphi(xu)\} = \{g^{-1}x^{-1}, xg\}.$$

Since $x \neq x^{-1}$, we have $g^{-1}x = xg$. Thus, $g^{-1} = xgx^{-1}$, and hence $g^{-1} = x^{-1}gx$. Furthermore, $xgxg = xgg^{-1}x = x^2$, and therefore $o(xg) = 4$.

(b) It follows from (2).

(c) Let $g \in G \setminus \gamma_\varphi$. By (a), $xgx^{-1} = g^{-1} = x^{-1}gx$. Hence, $x^2g = gx^2$.

Now, let $y \in \gamma_\varphi$. Then,

$$y^{-1}x = \varphi((xu) * (yu)) \in \{\varphi(xu) * \varphi(yu), \varphi(yu) * \varphi(xu)\} = \{yx^{-1}, xy^{-1}\}.$$

Thus, either $y^{-1}x = yx^{-1}$, or $y^{-1}x = xy^{-1}$. In both cases, we can conclude that $x^2y = yx^2$. Therefore, $x^2 \in \mathcal{Z}(G)$.

(d) Since $x \in \gamma_\varphi$, we have

$$\begin{aligned} \varphi(g) * \varphi(xu) &= g * (x^{-1}u) = (x^{-1}g)u, & \varphi(xu) * \varphi(g) &= (x^{-1}u) * g = (x^{-1}g^{-1})u \\ & & \text{and } \varphi(g * (xu)) &= \varphi((xg)u) = (xg)^\epsilon u, \end{aligned}$$

where $\epsilon \in \{-1, 1\}$. Thus, $(xg)^\epsilon \in \{x^{-1}g, x^{-1}g^{-1}\}$. If $\epsilon = 1$, then $xg = x^{-1}g^{-1}$ since $x \neq x^{-1}$, and so $g^2 = x^{-2} = x^2$. Hence, $o(g) = 4$, which is a contradiction. Therefore, $\epsilon = -1$. By (a), $g^{-1}x = xg \in \gamma_\varphi$.

Since $o(g^{-1}) \neq 4$, we conclude that $gx \in \gamma_\varphi$. The rest of the claim follows from the fact that G is finite.

(e) Let $g, h \in G \setminus \gamma_\varphi$ be such that $hg \in \gamma_\varphi$. Then,

$$\begin{aligned} \varphi(g) * \varphi(hu) &= g * (hu) = (hg)u, & \varphi(hu) * \varphi(g) &= (hu) * g = (hg^{-1})u \\ & & \text{and } \varphi(g * (hu)) &= \varphi((hg)u) = (hg)^{-1}u. \end{aligned}$$

Thus, $(hg)^{-1} \in \{hg, hg^{-1}\}$. Since $o(hg) = 4$, we have $g^{-1}h^{-1} = hg^{-1}$, and then $gh = h^{-1}g$. Hence, $hghg = h(h^{-1}g)g = g^2$ and we conclude that $o(g) = 4$.

□

Corollary 3.4. *There exists $y \in G$ such that:*

(a) $o(y) = 4$ and

(b) if $y = gh$, for some $g, h \in G$, then either $o(g) = 4$, or $o(h) = 4$.

Proof. Fix $\varphi \in H \setminus \{I\}$. Then, φ is a nontrivial half-automorphism and $\gamma_\varphi \neq \emptyset$. Let $y \in \gamma_\varphi$. Suppose there exist $g, h \in G$ such that $y = gh$, $o(g) \neq 4$ and $o(h) \neq 4$. Thus, $g, h \notin \gamma_\varphi$. By the item (e) of Proposition 3.3, $o(h) = 4$, which is a contradiction. \square

Suppose that $G = G_0 \cup G_0v$ is a generalized dihedral group. Note that $o(gv) = 2$, for any $g \in G_0$. Let $z \in G$ be of order 4. Then, $z \in G_0$ and $z = (z^{-1}v)(z^2v)$. Therefore, there is no element of G satisfying the condition of Corollary 3.4. Hence, the following result is at hand.

Corollary 3.5. *If G is a finite generalized dihedral group, then every half-automorphism of $M(2, G)$ is trivial.*

By Theorem 2.3, $|\mathcal{Z}(G)| \geq 2$. In the following result, this subgroup is described.

Theorem 3.6. $\mathcal{Z}(G) \cong C_2^m$, for some $m \geq 1$.

Proof. Fix $\varphi \in H \setminus \{I\}$. First, we will prove that $\mathcal{Z}(G) \cap \gamma_\varphi = \emptyset$.

Suppose there exists $z \in \mathcal{Z}(G) \cap \gamma_\varphi$. Let $g \in G \setminus \gamma_\varphi$. By the item (a) of Proposition 3.3, $g^{-1} = z^{-1}gz = g$. By (2), $\varphi(gu) = g^{-1}u$, for all $g \in G$. Consequently, for $g, h \in G$, we have

$$\varphi(gu * h) = (hg^{-1})u, \varphi(gu) * \varphi(h) = (g^{-1}h^{-1})u \text{ and } \varphi(h) * \varphi(gu) = (g^{-1}h)u.$$

Thus, either $hg = gh$, or $h^{-1} = ghg^{-1}$, for all $g, h \in G$. Since G is nonabelian, there exist $x, y \in G$ such that $xy \neq yx$, and then $y^{-1} = xyx^{-1}$. Since $x(yz) \neq yxz = (yz)x$, we have $(yz)^{-1} = xyzx^{-1} = y^{-1}z$, and hence $y^{-1} = z^2y^{-1}$. Therefore, $o(z) = 2$, a contradiction with $z \in \gamma_\varphi$. We conclude that $\mathcal{Z}(G) \cap \gamma_\varphi = \emptyset$.

Now, let $z \in \mathcal{Z}(G)$ and $x \in \gamma_\varphi$. By the item (a) of Proposition 3.3, $z^{-1} = x^{-1}zx = z$. Thus, $\mathcal{Z}(G)$ has exponent 2, and hence there exists $m \geq 1$ such that $\mathcal{Z}(G) \cong C_2^m$. \square

4 The half-automorphism group

We say that a loop Q has the *antiautomorphic inverse property* (AAIP) if it has two-sided inverses and satisfies the identity $(xy)^{-1} = y^{-1}x^{-1}$. In this type of loop, the *inversion mapping* $J : Q \rightarrow Q$, defined by $J(x) = x^{-1}$, is an anti-automorphism. Furthermore, it is easy to see that $J \in \mathcal{Z}(\text{Half}_T(Q))$.

Proposition 4.1. *Let Q be a nonabelian loop with the (AAIP). Then,*

$$\text{Half}_T(Q) = \langle J \rangle \text{Aut}(Q) \cong C_2 \times \text{Aut}(Q).$$

Proof. Since $J \in \mathcal{Z}(\text{Half}_T(Q))$, J has order 2 and $J \notin \text{Aut}(Q)$, it follows that $\langle J \rangle \text{Aut}(Q) \cong C_2 \times \text{Aut}(Q)$.

Denote the set of the anti-automorphisms of Q by $\text{Ant}(Q)$. Since Q is nonabelian, we have $\text{Half}_T(Q) = \text{Aut}(Q) \cup \text{Ant}(Q)$, where $\text{Aut}(Q) \cap \text{Ant}(Q) = \emptyset$. It therefore suffices to verify that $\text{Ant}(Q) = J \text{Aut}(Q)$.

Let $\varphi \in \text{Aut}(Q)$. Then, $J\varphi(xy) = J\varphi(y)J\varphi(x)$, for all $x, y \in G$, and so $J \text{Aut}(Q) \subset \text{Ant}(Q)$. On the other hand, let $\varphi \in \text{Ant}(Q)$. Then, $J\varphi(xy) = J\varphi(x)J\varphi(y)$, for all $x, y \in G$. Thus, $J\varphi \in \text{Aut}(Q)$, and hence $\varphi \in J \text{Aut}(Q)$. \square

Remark 4.2. If Q is an abelian loop, then $Half(Q) = Half_T(Q) = Aut(Q)$.

Every Moufang loop has the (AAIP). Thus, Theorem 2.2, Proposition 4.1 and Remark 4.2 give us a description of the trivial half-automorphism group of Chein loops.

From now on, we will suppose that the Chein loop $L = M(G, 2)$ is finite and has a nontrivial half-automorphism. Then, G is a finite nonabelian group that is not a generalized dihedral group.

By Theorem 2.2 and Propositions 2.4 and 4.1, $Half(L) = \langle J \rangle ADH$, where A and D were defined in section 2, and J is the inversion mapping of L .

Lemma 4.3. *Let $\varphi \in H$, $\phi \in Aut(G)$, and $t \in G$. Then:*

- (a) $(a_\phi)^{-1}\varphi a_\phi \in H$,
- (b) *If $t \in G - \gamma_\varphi$, then $(d_t)^{-1}\varphi d_t \in H$. If $t \in \gamma_\varphi$, then $d_t\varphi d_t \in H$.*

Proof. For $g \in G$, we have that $(a_\phi)^{-1}\varphi a_\phi(g) = \phi^{-1}\phi(g) = g$ and $(d_t)^{-1}\varphi d_t(g) = g = d_t\varphi d_t(g)$.

Consider $u \in L$. Thus, $(a_\phi)^{-1}\varphi a_\phi(u) = (a_\phi)^{-1}\varphi(u) = a_{\phi^{-1}}(u) = u$. If $t \in G \setminus \gamma_\varphi$, then $\varphi(tu) = tu$ by Proposition 3.3, and so $(d_t)^{-1}\varphi d_t(u) = d_{t^{-1}}(tu) = u$. If $t \in \gamma_\varphi$, then $\varphi(tu) = t^{-1}u$, and hence $d_t\varphi d_t(u) = d_t(t^{-1}u) = u$. \square

By Proposition 3.1, $H \cap AD = \{I\}$. Since $H, AD \leq Half(L)$, it follows that $DH, ADH \leq Half(L)$ by Lemma 4.3. In [8, Lemma 2.1], it was proved that $fJ = Jf$, for all $f \in Half(L)$. Thus, $J \in \mathcal{Z}(Half(L))$. Furthermore, it is clear that $J \notin ADH$. Consequently, we have

$$Half(L) \cong C_2 \times ADH. \quad (3)$$

Now, we will describe the group ADH . For $a_\phi \in A$, define $\sigma_\phi : DH \rightarrow DH$ by $\sigma_\phi(d_t\varphi) = a_\phi d_t\varphi (a_\phi)^{-1} = d_{\phi(t)}a_\phi\varphi(a_\phi)^{-1}$, where $\varphi \in H$ and $d_t \in D$. By Lemma 4.3, σ_ϕ is well-defined, so it is the restriction of an inner automorphism of ADH to DH . Therefore, $\sigma_\phi \in Aut(DH)$ and $\sigma_\phi\sigma_{\phi'} = \sigma_{\phi\phi'}$, for any $a_\phi, a_{\phi'} \in A$.

Define $\sigma : A \rightarrow Aut(DH)$ by $\sigma(a_\phi) = \sigma_\phi$. Then, the mapping σ is a homomorphism. Thus, we can construct the semidirect product $A \overset{\sigma}{\ltimes} DH$, where the operation is given by

$$(a_\phi, d_t\varphi) \overset{\sigma}{\cdot} (a_{\phi'}, d_{t'}\varphi') = (a_{\phi\phi'}, d_t\varphi d_{\phi(t')}a_{\phi'}\varphi'(a_\phi)^{-1}).$$

In ADH , we have that $d_t\varphi a_\phi d_{t'}\varphi' a_{\phi'}^{-1} = d_t\varphi d_{\phi(t')}a_{\phi'}\varphi'(a_\phi)^{-1}a_{\phi\phi'}$, and then the mapping $\psi : ADH \rightarrow A \overset{\sigma}{\ltimes} DH$, defined by $\psi(d_t\varphi a_\phi) = (a_\phi, d_t\varphi)$, is an isomorphism. Therefore, we established the following result.

Proposition 4.4. $ADH \cong A \overset{\sigma}{\ltimes} DH \cong Aut(G) \overset{\sigma}{\ltimes} DH$.

Now, we will describe the group DH .

Proposition 4.5. $Z = \{d_z \mid z \in \mathcal{Z}(G)\} \leq \mathcal{Z}(DH)$.

Proof. Let $z \in \mathcal{Z}(G)$ and $d_t \in D$. Then, $d_z d_t = d_{zt} = d_{tz} = d_t d_z$.

Now, let $\varphi \in H$. Since $o(z) \in \{1, 2\}$, $z\gamma_\varphi = \gamma_\varphi$ by the item (d) of Proposition 3.3. If $g \in \gamma_\varphi$, then $zg \in \gamma_\varphi$, and so $d_z\varphi(gu) = (zg^{-1})u = (g^{-1}z^{-1})u = \varphi d_z(gu)$. If $g \in G \setminus \gamma_\varphi$, then $zg \in G \setminus \gamma_\varphi$, and hence $d_z\varphi(gu) = (zg)u = \varphi d_z(gu)$.

Therefore, $Z \subset \mathcal{Z}(DH)$. Since $Z \cong \mathcal{Z}(G)$, we conclude that $Z \leq \mathcal{Z}(DH)$. \square

Since $Z \subset D$, it follows that $Z \cap H = \{I\}$. By Proposition 4.5, $ZH \leq DH$.

For $x \in G$, the mapping $I_x : G \rightarrow G$, defined by $I_x(g) = gxg^{-1}$, is an inner automorphism of G . Let $\mathcal{I}(G) = \{I_x \mid x \in G\}$ be the inner automorphism group of G . It is well known that $G/\mathcal{Z}(G) \cong \mathcal{I}(G)$.

Proposition 4.6. *We have:*

- a) $ZH \cong C_2^n$, where $|\mathcal{Z}(G)| = 2^m$, $|H| = 2^r$, and $n = m + r$,
- b) $ZH \triangleleft DH$,
- c) $DH/ZH \cong \mathcal{I}(G)$.

Proof. a) Since $Z \leq \mathcal{Z}(DH)$ and $Z \cap H = \{I\}$, we have $ZH \cong Z \times H$. The result follows from Proposition 3.2 and Theorem 3.6.

b) Let $z \in \mathcal{Z}(G)$, $t \in G$, and $\varphi, \varphi' \in H$. Then,

$$\varphi' d_z \varphi (\varphi')^{-1} = d_z \varphi' \varphi \varphi' = d_z \varphi \quad \text{and} \quad d_t d_z \varphi d_{t^{-1}} = d_z d_t \varphi d_{t^{-1}} = d_{zt^2} d_{t^{-1}} \varphi d_{t^{-1}}.$$

It is clear that $\varphi' d_z \varphi (\varphi')^{-1} \in ZH$. If $t \in \gamma_\varphi$, then $t^2 \in \mathcal{Z}(G)$ and $d_{t^{-1}} \varphi d_{t^{-1}} \in H$ by Proposition 3.3 and Lemma 4.3. If $t \notin \gamma_\varphi$, then $d_t \varphi d_{t^{-1}} \in H$ by Lemma 4.3. Thus, $d_t d_z \varphi d_{t^{-1}} \in ZH$, for all $t \in G$. Therefore, $ZH \triangleleft DH$.

c) Define $\psi : DH \rightarrow \mathcal{I}(G)$ by $\psi(d_t \varphi) = I_t$. It is clear that ψ is surjective.

Let $t, t' \in G$ and $\varphi, \varphi' \in H$. If $t' \in \gamma_\varphi$, then $t'^2 \in \mathcal{Z}(G)$, and hence $I_{t'}(g) = t'gt'^{-1} = t'^{-1}t'^2gt'^{-1} = t'^{-1}gt'^2t'^{-1} = I_{t'^{-1}}(g)$, for all $g \in G$. Since $d_{t'} \varphi d_{t'} \in H$, we have

$$\psi(d_t \varphi d_{t'} \varphi') = \psi(d_{tt'^{-1}} d_{t'} \varphi d_{t'} \varphi') = I_{tt'^{-1}} = I_t I_{t'^{-1}} = I_t I_{t'} = \psi(d_t \varphi) \psi(d_{t'} \varphi').$$

If $t' \notin \gamma_\varphi$, then $d_{t'^{-1}} \varphi d_{t'} \in H$. Thus,

$$\psi(d_t \varphi d_{t'} \varphi') = \psi(d_{tt'} d_{t'^{-1}} \varphi d_{t'} \varphi') = I_{tt'} = I_t I_{t'} = \psi(d_t \varphi) \psi(d_{t'} \varphi').$$

Hence, ψ is a homomorphism.

Suppose that $\psi(d_t \varphi) = I$. Then, $g = I_t(g) = tgt^{-1}$, for all $g \in G$. Thus, $t \in \mathcal{Z}(G)$, and hence $\text{Ker}(\psi) = ZH$. Therefore, $DH/ZH \cong \mathcal{I}(G)$. \square

For $t \in G$, define $\rho_t : ZH \rightarrow ZH$ by $\rho_t(d_z \varphi) = d_t d_z \varphi d_{t^{-1}}$, where $d_z \in Z$ and $\varphi \in H$. By the item (b) of Proposition 4.6, ρ_t is well-defined, and then it is the restriction of an inner automorphism of DH to ZH . Thus, $\rho_t \in \text{Aut}(ZH)$ and $\rho_t \rho_{t'} = \rho_{tt'}$, for any $t, t' \in G$.

Define $\rho : G \rightarrow \text{Aut}(ZH)$ by $\rho(t) = \rho_t$. Then, ρ is a homomorphism. Thus, we can construct the semidirect product $G \overset{\rho}{\ltimes} ZH$, where the operation is given by:

$$(t, \alpha) \overset{\rho}{\cdot} (t', \alpha') = (tt', \alpha d_t \alpha' d_{t^{-1}}) \quad (t, t' \in G, \alpha, \alpha' \in ZH)$$

We can now state and prove the main theorem of this paper.

Theorem 4.7. *Let G be a finite nonabelian group such that $M(G, 2)$ has a nontrivial half-automorphism. Then,*

$$\text{Half}(M(G, 2)) \cong C_2 \times (\text{Aut}(G) \overset{\sigma}{\ltimes} DH)$$

where $DH \cong (G \overset{\rho}{\ltimes} ZH)/W$, $ZH \cong C_2^n$ and $W = \{(z, d_z) \mid z \in \mathcal{Z}(G)\} \cong \mathcal{Z}(G)$. Furthermore, $|\text{Half}(M(G, 2))| = 2 |\text{Aut}(G)| |G| |H| = 2^{n+1} |\text{Aut}(G)| |\mathcal{I}(G)|$.

Proof. Define $\psi : G \overset{\rho}{\ltimes} ZH \rightarrow DH$ by $\psi((t, \alpha)) = \alpha d_t$. It is clear that ψ is surjective. Since $\alpha d_t \alpha' d_{t'} = \alpha d_t \alpha' d_{t^{-1}} d_{tt'}$, for all $\alpha, \alpha' \in ZH$ and $t, t' \in G$, it follows that ψ is a homomorphism.

Suppose that $\psi((t, d_z \varphi)) = I$, for some $t \in G$, $z \in \mathcal{Z}(G)$ and $\varphi \in H$. Then, $I = d_z \varphi d_t = \varphi d_{zt}$. Since $H \cap D = \{I\}$, we have $\varphi = I$ and $zt = 1$. Thus, $\text{Ker}(\psi) = \{(z, d_z) \mid z \in \mathcal{Z}(G)\}$. Therefore, $DH \cong (G \overset{\rho}{\ltimes} ZH)/W$, where $W = \text{Ker}(\psi)$.

For $(z, d_z), (z', d_{z'}) \in W$, we have that $(z, d_z) \overset{\rho}{\cdot} (z', d_{z'}) = (zz', d_{zz'})$. Hence, $W \cong \mathcal{Z}(G)$.

The rest of the claim follows from (3) and Propositions 4.4 and 4.6. \square

In the following, some examples of Chein loops that have nontrivial half-automorphisms are provided. The generators of H in these examples were presented in [4, Examples 8 and 9]. In each example, the order of H was obtained by using GAP [6] computing with the LOOPS package [11].

Example 4.8. Let $Q_8 = \{1, i, j, k, -1, -i, -j, -k\}$ be the quaternion group, where the operation is given by the rules: $i^2 = j^2 = k^2 = -1$, $(-1)^2 = 1$, $ij = k = -ji$, $jk = i = -kj$, and $ki = j = -ik$. The group H has order 8 and is generated by the nontrivial half-automorphisms

$$\varphi_x(gu^l) = \begin{cases} g, & \text{if } l = 0, \\ gu, & \text{if } l = 1 \text{ and } g \notin \{x, -x\}, \\ (-g)u, & \text{if } l = 1 \text{ and } g \in \{x, -x\}, \end{cases}$$

where $x \in \{i, j, k\}$. Since $|\text{Aut}(Q_8)| = 24$, we have $|\text{Half}(M(Q_8, 2))| = 3072$, $|\text{Half}_T(M(Q_8, 2))| = 384$ and $|\text{Aut}(M(Q_8, 2))| = 192$. Then, $M(Q_8, 2)$ has 2688 nontrivial half-automorphisms.

Example 4.9. Let $G = C_4 \times C_3$ be the nonabelian semidirect product between the cyclic groups of orders 4 and 3. In this case, $H = \{I, \varphi\}$, where φ is the nontrivial half-automorphism defined by

$$\varphi(gu^i) = \begin{cases} g, & \text{if } i = 0, \\ gu, & \text{if } i = 1 \text{ and } o(g) \neq 4, \\ (g^{-1})u, & \text{if } i = 1 \text{ and } o(g) = 4. \end{cases}$$

Since $|\text{Aut}(G)| = 12$, we have $|\text{Half}(M(G, 2))| = 576$, $|\text{Half}_T(M(G, 2))| = 288$ and $|\text{Aut}(M(G, 2))| = 144$. Then, $M(G, 2)$ has 288 nontrivial half-automorphisms.

Acknowledgments

Some calculations in this work have been made by using the LOOPS package [11] for GAP [6].

References

- [1] R.H. Bruck, *A Survey of Binary Systems*, Springer, 1971.
- [2] R.H. Bruck, L.J. Paige, *Loops whose inner mappings are automorphisms*, Ann. Math., **63**, (1956), 308–323.
- [3] O. Chein, *Moufang loops of small order I*, Trans. Amer. Math. Soc., **188**, (1974), 31–51.
- [4] S. Gagola III, M.L. Merlini Giuliani, *On half-automorphisms of certain Moufang Loops with even order*, J. Algebra, **386**, (2013), 131–141.

- [5] S. Gagola III, M.L. Merlini Giuliani, *Half-automorphisms of simple Moufang loops*, J. Algebra, **546**, (2020), 27–36.
- [6] The GAP Group, GAP - Groups, Algorithms, and Programming, Version 4.10.1, 2019. <http://www.gap-system.org>
- [7] A. Grishkov, A.V. Zavarnitsine, M.L. Merlini Giuliani, *Automorphism group of Chein loops*, Quasigroups Related Systems, **13**, (2005), 105 – 108.
- [8] M. Kinyon, I. Stuhl, P. Vojtěchovský, *Half-Isomorphisms of Moufang Loops*, J. Algebra, **450**, (2016), 152–161.
- [9] M.L. Merlini Giuliani, G. Souza dos Anjos, *Half-isomorphisms of dihedral automorphic loops*, Comm. Algebra, **48**, (2020), no. 3, 1150–1162.
- [10] M.L. Merlini Giuliani, G. Souza dos Anjos, *Lie automorphic loops under half-automorphisms*, J. Algebra Appl., DOI 10.1142/S0219498820502217.
- [11] G.P. Nagy, P. Vojtěchovský. LOOPS: Computing with quasigroups and loops in GAP, version 3.4.0, package for GAP, <https://cs.du.edu/~petr/loops/>
- [12] H.O. Pflugfelder, *Quasigroups and Loops: Introduction*, Sigma Series in Pure Math. **7**, Heldermann, 1990.
- [13] W.R. Scott, *Half-homomorphisms of groups*, Proc. Amer. Math. Soc., **8**, (1957), 1141–1144.