

Gröbner-Shirshov bases for the Coxeter groups of the types G_2, F_4, E_6, E_7 and E_8

JUN WANG

The author is mainly interested in the Gröbner-Shirshov bases of finite Coxeter groups. It is known that the finite Coxeter groups are classified in terms of Coxeter-Dynkin diagrams. Under the fixed order, it is worth mentioning that the presentation of the group determines the Gröbner-Shirshov bases of the group. In this paper, the author rearranges the generators, marks them on Coxeter-Dynkin diagrams, and gets a simple presentation of the Gröbner-Shirshov bases for Coxeter groups of types G_2, F_4, E_6 and E_7 . This article also gives the Gröbner-Shirshov basis of a Coxeter group of type E_8 .

[16S15; 20F55](#)

1 Introduction

Let $M = (m_{ij})$ be a symmetric $n \times n$ matrix over natural numbers, with ∞ such that $m_{ii} = 1, 2 \leq m_{ij} \leq \infty$ for $i \neq j$. The following presentation is in fact a presentation of a group which is called the Coxeter group $W = W(M)$:

$$W = \langle s_1, \dots, s_n \mid (s_i s_j)^{m_{ij}} = 1, \text{ where } 1 \leq i, j \leq n, m_{ij} \neq \infty \rangle$$

In 1935, Coxeter classified the finite Coxeter groups in terms of Coxeter-Dynkin diagrams [13]. For the purpose of understanding the structure of Coxeter groups, we focus on the noncommutative Gröbner bases theory, which is called the Gröbner-Shirshov bases theory. The effective notion stems from Shirshov's composition Lemma and his algorithm [17, 18] for Lie algebras, and independently from Buchberger's algorithm [11] of computing Gröbner bases for commutative algebras. The Gröbner-Shirshov bases for some Lie algebras and superalgebras were determined in [7, 1, 2, 3, 4, 5, 6]. In 2006, Bokut and Vesnin [9] applied Shirshov's method to some braid groups. Bokut considered the Coxeter groups of the types A_n, B_n, D_n in 2001 [8] and formulated a general hypothesis about Gröbner-Shirshov bases of any Coxeter group. In 2008, Lee and Denis [15] got the Gröbner-Shirshov bases for the Coxeter groups of types E_6 and E_7 . In 2009, Yuqun Chen and Cihua Liu [12] gave a counter-example for the general

hypothesis by Bokut in [8]. For non-crystallographic Coxeter groups, Jeong-Yup Lee and Dong-IL Lee [16] calculated the Gröbner-Shirshov bases in 2018.

In this paper, we focus on the Coxeter groups of types G_2, F_4, E_6, E_7 and E_8 . We rearrange the generators of Coxeter groups to reduce time in computing, get a simple presentation of Gröbner-Shirshov bases of the Coxeter groups of types G_2, F_4, E_6 and E_7 , and obtain the Gröbner-Shirshov basis of the Coxeter group of type E_8 . In Section 2, we introduce the Gröbner-Shirshov bases theory. The Gröbner-Shirshov bases for Coxeter groups of type G_2, F_4, E_6, E_7, E_8 will be set out in Section 3 and Section 4. The computation is based on the the Non Commutative Algebra Packages run under Mathematica made by Bill Helton and Mauricio de Oliveira at [14].

2 Preliminaries

Let K be a field, $K\langle X \rangle$ be the non-commutative polynomial algebra generated by X where $X = \{x_1, \dots, x_n\}$ is a finite set. Let $X^* = \{u | u = x_{i_1} \dots x_{i_m}, m \geq 0, i_1, \dots, i_m \in \{1, \dots, n\}\}$ be the monoid generated by X , and define the identity by the empty word which is denoted by 1. Denote the degree of a word $w \in X^*$ by $|w|$.

Definition 2.1 A total order σ on the set X^* is called well-ordering, if every non-empty subset of X^* has a minimal element under the order σ .

Definition 2.2 A well-ordering σ on the set X^* is called monomial, if for every $u, v \in X^*$,

$$u <_{\sigma} v \iff w_1 u w_2 <_{\sigma} w_1 v w_2, \forall w_1, w_2 \in X^*$$

Definition 2.3 The deg-lex order on the X^* is defined as follows: $\forall w_1 = x_{i_1} \dots x_{i_m}, w_2 = x_{j_1} \dots x_{j_s} \in X^*$,

$$w_1 <_{deg-lex} w_2 \iff \text{or } m < s, \text{ or } m = s, w_1 <_{lex} w_2$$

Example 2.4 For the set $X = \{x_1, x_2, x_3\}$, under the deg-lex order, we have

$$1 < x_1 < x_2 < x_3 < x_1^2 < x_1 x_2 < x_1 x_3 < x_2 x_1 < x_2^2 < x_2 x_3 < \dots$$

obviously, under the deg-lex order, X^* is a well-ordering set, and the deg-lex order is monomial.

Under the monomial order σ on X^* , every polynomial $f \in K\langle X \rangle$ can be uniquely represented as

$$f = \sum_i c_i w_i,$$

where $c_i \in K \setminus \{0\}$, $w_i \in X^*$, and satisfy that $w_i <_\sigma w_j$, if $i < j$.

Definition 2.5 • The monomial $c_1 w_1$ is called the leading term of f with respect to σ and denoted by $LT_\sigma(f)$.

- $c_1 \in K \setminus \{0\}$ is called the leading coefficient of f with respect to σ and denoted by $LC_\sigma(f)$.
- $w_1 \in X^*$ is called the leading word of f with respect to σ and denoted by $LW_\sigma(f)$ or \bar{f} . The polynomial f is called monic if $LC_\sigma(f) = 1$.

Definition 2.6 For $g \in K\langle X \rangle$, $w \in X^*$, the K -module endomorphism

$$r_g : K\langle X \rangle \rightarrow K\langle X \rangle$$

defined by

$$r_g(w) = \begin{cases} w - agb, & \text{if } \exists a, b \in X^*, \text{ s.t. } w = a\bar{g}b; \\ w, & \text{if } \forall a, b \in X^*, w \neq a\bar{g}b; \end{cases}$$

is called a reduction by g .

For a subset $S = \{g_1, \dots, g_n\} \in K\langle X \rangle$, $f \in K\langle X \rangle$ is called a monomial which can be reduced to 0 by S , if f can be reduced to 0 by a finite sequence of reduction $r_1, r_2, \dots, r_m (r_i \in \{r_g | g \in S\})$. Then denote it by $f \xrightarrow{S} 0$. If not, we denote it by $f \xrightarrow{S} \neq 0$.

Definition 2.7 Under a monomial order σ on X^* , given two monic polynomials $f, g \in K\langle X \rangle$, (f, g) defined by

$$(f, g) = \{fb - ag | a, b \in X^*, \bar{f}b = a\bar{g}, |\bar{f}| + |\bar{g}| > |\bar{f}b|\} \cup \{f - agb | a, b \in X^*, \bar{f} = a\bar{g}b\}$$

is called the composition set of f and g .

Remark Generally, (f, g) is not equal to (g, f) .

Under a monomial order σ , let S be a subset of $\{g \in K\langle X \rangle | LC(g) = 1\}$, the ideal generated by S is denoted by $\text{Ideal}(S)$.

Definition 2.8 S is called then Gröbner-Shirshov basis of $\text{Ideal}(S)$ under the monomial order σ , if $\forall f, g \in S$, all the elements of $(f, g), (g, f)$ can be reduced to 0 by S .

Now, we give the Shirshov algorithm about calculating the Gröbner-Shirshov basis. If $\forall f, g \in S$, all the elements of $(f, g), (g, f)$ can be reduced to 0 by S , then S is the Gröbner-Shirshov basis of $\text{Ideal}(S)$ under the monomial order σ ; If not, we add the elements of the composition set which can not be reduced to 0 by S , and we get the set S^1 . Repeat the process, we get $S \subset S^1 \subset S^2 \subset \dots$. If we get a set S^m satisfying that $\forall f, g \in S^m$, all the elements of $(f, g), (g, f)$ can be reduced to 0 by S^m , then we denote $S^m = S^c$, and have that $\text{Ideal}(S^c) = \text{Ideal}(S)$. The set S^c is exactly the Gröbner-Shirshov basis of $\text{Ideal}(S)$. This process is called *Shirshov algorithm*. It's worth to mention that m is not always finite and S^m is not always a finite set.

Definition 2.9 A Gröbner-Shirshov basis $S = \{f_1, \dots, f_n\} \subset K\langle X \rangle$ under the monomial order σ is called reduced if $\forall 1 \leq i, j \leq n, i \neq j, \bar{f}_i \neq a\bar{f}_j b$ for $\forall a, b \in X^*$.

Let $A = \text{sgp}\langle X | R \rangle$ be a semigroup presentation. Under a monomial order σ , $R = \{u_i = v_i | u_i, v_i \in X^*, v_i <_\sigma u_i\}$, corresponds to a subset $S = \{u_i - v_i | u_i, v_i \in X^*, v_i <_\sigma u_i\} \subset K\langle X \rangle$. The ideal generated by S is denoted by $\text{Ideal}(S)$. The set S^c got by Shirshov algorithm is called the Gröbner-Shirshov basis of the semigroup A .

Lemma 2.10 (Composition-Diamond Lemma [8]) S^c is a Gröbner-Shirshov basis if and only if the set

$$\{u \in X^* | u \neq a\bar{f}b, f \in S^c\}$$

of S^c -reduced words consists of a linear bases of the algebra $\langle X | R \rangle$ which is the factor-algebra of $K\langle X \rangle$ by the ideal generated by R .

We focus on finite Coxeter groups. By [10], the Coxeter-Dynkin diagram of any irreducible finite Coxeter group is isomorphic to one of the diagram in Figure 1.




Figure 1: The Coxeter-Dynkin diagrams of irreducible finite Coxeter groups

For the Coxeter groups of the types A_n, B_n, D_n , Bokut computed the Gröbner-Shirshov bases in [8]. We will set out the Gröbner-Shirshov bases for the Coxeter groups of the types G_2, F_4, E_6, E_7 and E_8 in the following sections.

It is significant that the presentation of group determines the computation of the Gröbner-Shirshov bases under the fixed order. In the following sections, we fix the deg-lex order and mark the generators x_i on Coxeter-Dynkin diagrams. Thus we get a new presentation of Coxeter groups, and get the Gröbner-Shirshov bases in simple forms.

3 Gröbner – Shirshov bases of the Coxeter groups of the type G_2, F_4

For the Coxeter group of type G_2 , we marked the generators on the Coxeter-Dynkin diagram of G_2 which is as Figure 2.




Figure 2: The Coxeter-Dynkin diagram of G_2

This means that the initial relations are

$$\begin{aligned}\theta_i^1 &= x_i^2 - 1, i = 1, 2 \\ \theta^2 &= (x_2x_1)^3 - (x_1x_2)^3.\end{aligned}$$

After the computation on Mathematica, the Gröbner-Shirshov basis of the Coxeter group of the type G_2 is as follows:

$$\begin{aligned}\theta_i^1 &= x_i^2 - 1, i = 1, 2 \\ \theta^2 &= (x_2x_1)^3 - (x_1x_2)^3.\end{aligned}$$

As the same, for the Coxeter group of type F_4 , we marked the generators on the Coxeter-Dynkin diagram of F_4 which is as Figure 3.




Figure 3: The Coxeter-Dynkin diagram of F_4

This means that the initial relations are

$$\theta_i^1 = x_i^2 - 1, i = 1, 2, 3, 4$$

$$\theta^2 = x_4x_3x_4 - x_3x_4x_3$$

$$\theta^3 = x_4x_2 - x_2x_4$$

$$\theta^4 = x_4x_1 - x_1x_4$$

$$\theta^5 = x_3x_2x_3x_2 - x_2x_3x_2x_3$$

$$\theta^6 = x_3x_1 - x_1x_3$$

$$\theta^7 = x_2x_1x_2 - x_1x_2x_1$$

After the computation on Mathematica, the Gröbner-Shirshov basis of the Coxeter group of the type F_4 is as follows:

$$\theta_i^1 = x_i^2 - 1, i = 1, 2, 3, 4$$

$$\theta^2 = x_4x_3x_4 - x_3x_4x_3$$

$$\theta^3 = x_4x_2 - x_2x_4$$

$$\theta^4 = x_4x_1 - x_1x_4$$

$$\theta^5 = x_3x_2x_3x_2 - x_2x_3x_2x_3$$

$$\theta^6 = x_3x_1 - x_1x_3$$

$$\theta^7 = x_2x_1x_2 - x_1x_2x_1$$

$$\alpha^1 = x_3x_2x_1x_3x_2x_1 - x_2x_3x_2x_3x_1x_2$$

$$\alpha^2 = x_4x_3x_2x_4 - x_3x_4x_3x_2$$

$$\alpha^3 = x_4x_3x_2x_3x_4x_3 - x_3x_4x_3x_2x_3x_4$$

$$\alpha^4 = x_4x_3x_2x_1x_4 - x_3x_4x_3x_2x_1$$

$$\alpha^5 = x_4x_3x_2x_1x_3x_4x_3 - x_3x_4x_3x_2x_3x_4x_1$$

$$\alpha^6 = x_4x_3x_2x_1x_3x_2x_4x_3x_2x_3 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2$$

$$\alpha^7 = x_4x_3x_2x_1x_3x_2x_4x_3x_2x_1x_3 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2x_1$$

$$\alpha^8 = x_4x_3x_2x_1x_3x_2x_3x_4x_3x_2x_3x_4 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2x_4x_3$$

$$\alpha^9 = x_4x_3x_2x_1x_3x_2x_3x_4x_3x_2x_1x_3x_4 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2x_1x_4x_3$$

$$\alpha^{10} = x_4x_3x_2x_1x_3x_2x_3x_4x_3x_2x_1x_3x_2x_4 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2x_1x_4x_3x_2$$

$$\alpha^{11} = x_4x_3x_2x_1x_3x_2x_3x_4x_3x_2x_1x_3x_2x_3x_4x_3 - x_3x_4x_3x_2x_3x_4x_1x_2x_3x_2x_1x_4x_3x_2x_3x_4$$

4 Gröbner–Shirshov bases of the Coxeter groups of the types E_6, E_7, E_8

For the Coxeter groups of types E_n , we marked the generators on the Coxeter-Dynkin diagram of E_n which is as Figure 4:




Figure 4: The Coxeter-Dynkin diagram of the Coxeter groups of types E_n

This means that for the Coxeter groups of the types E_n , we have the initial relations as follows:

$$\begin{aligned}
 \theta_i^1 &= x_i^2 - 1 & 7 - n \leq i \leq 5 \\
 \theta_i^2 &= x_{i+1}x_i x_{i+1} - x_i x_{i+1} x_i & 7 - n \leq i \leq 3 \\
 \theta_{i,j}^3 &= x_i x_j - x_j x_i & 7 - n \leq j + 1 < i \leq 4 \\
 \theta_i^4 &= x_5 x_i - x_i x_5 & i \neq 6, 5, 3 \\
 \theta^5 &= x_5 x_3 x_5 - x_3 x_5 x_3 \\
 \varepsilon^1 &= x_6^2 - 1 \\
 \varepsilon^2 &= x_6 x_4 x_6 - x_4 x_6 x_4 \\
 \varepsilon_i^3 &= x_6 x_i - x_i x_6 & i \neq 4, 6
 \end{aligned}$$

Let $n = 6, 7, 8$, they are initial relations of finite Coxeter groups of types E_6, E_7 and E_8 . We denote that

$$\begin{aligned}
 x_{i,j} &= x_i x_{i-1} x_{i-2} \dots x_j, \quad i > j; & x_{ii} &= x_i, & x_{i,i+1} &= 1; \\
 x'_{i,j} &= x_i x_{i-2} x_{i-3} \dots x_j, \quad i > j + 1; & x'_{i,i-1} &= x_i, & x'_{i,i} &= 1.
 \end{aligned}$$

And we obtain the following Gröbner-Shirshov bases via the computation on Mathematica:

The Gröbner-Shirshov basis of the Coxeter group of the type E_6 is as follows:

$$\begin{aligned}
\theta_i^1 &= x_i^2 - 1 & 1 \leq i \leq 5 \\
\theta_i^2 &= x_{i+1}x_i x_{i+1} - x_i x_{i+1} x_i & 1 \leq i \leq 3 \\
\theta_{i,j}^3 &= x_i x_j - x_j x_i & 1 \leq j+1 < i \leq 4 \\
\theta_i^4 &= x_5 x_i - x_i x_5 & i \neq 3, 5, 6 \\
\theta^5 &= x_5 x_3 x_5 - x_3 x_5 x_3 \\
\varepsilon^1 &= x_6^2 - 1 \\
\varepsilon^2 &= x_6 x_4 x_6 - x_4 x_6 x_4 \\
\varepsilon_i^3 &= x_6 x_i - x_i x_6 & i \neq 4, 6 \\
\alpha_{i,j}^1 &= x_{i+1,j} x_{i+1} - x_i x_{i+1,j} & 1 \leq j < i \leq 3 \\
\alpha_{5,j}^2 &= x'_{5,j} x_5 - x_3 x'_{5,j} & 1 \leq j \leq 2 \\
\beta_i^1 &= x'_{5,i} x_{4,i} - x_4 x'_{5,i} x_{4,i+1} & 1 \leq i \leq 3 \\
\beta_i^2 &= x'_{5,i} x_4 x_5 - x_3 x'_{5,i} x_4 & 1 \leq i \leq 3 \\
\beta_{i,j}^3 &= x'_{5,i} x_{4,j} x'_{5,j} - x_3 x'_{5,i} x_{4,j} x'_{5,j+1} & 1 \leq i < j \leq 3 \\
\eta_i &= x'_{6,i} x'_{5,i} - x_5 x'_{6,i} x'_{5,i+1} & 1 \leq i \leq 4 \\
\xi_{i,j} &= x'_{6,i} x'_{5,j} x_6 - x_4 x'_{6,i} x'_{5,j} & 1 \leq i < j \leq 5 \\
\lambda_{i,j,k} &= x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k} - x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k+1} & 1 \leq i < j < k \leq 4 \\
\nu_{1,2,3,l} &= x'_{6,1} x'_{5,2} x_{4,3} x_5 x'_{6,l} x_5 - x_4 x'_{6,1} x'_{5,2} x_{4,3} x_5 x'_{6,l} & 1 \leq l \leq 3
\end{aligned}$$

The Gröbner-Shirshov basis of the Coxeter group of the type E_7 is as follows:

$$\begin{aligned}
\theta_i^1 &= x_i^2 - 1 && 0 \leq i \leq 5 \\
\theta_i^2 &= x_{i+1}x_i x_{i+1} - x_i x_{i+1} x_i && 0 \leq i \leq 3 \\
\theta_{i,j}^3 &= x_i x_j - x_j x_i && 0 \leq j+1 < i \leq 4 \\
\theta_i^4 &= x_5 x_i - x_i x_5 && i \neq 3, 5, 6 \\
\theta^5 &= x_5 x_3 x_5 - x_3 x_5 x_3 \\
\varepsilon^1 &= x_6^2 - 1 \\
\varepsilon^2 &= x_6 x_4 x_6 - x_4 x_6 x_4 \\
\varepsilon_i^3 &= x_6 x_i - x_i x_6 && i \neq 4, 6 \\
\alpha_{i,j}^1 &= x_{i+1,j} x_{i+1} - x_i x_{i+1,j} && 0 \leq j < i \leq 3 \\
\alpha_{5,j}^2 &= x'_{5,j} x_5 - x_3 x'_{5,j} && 0 \leq j \leq 2 \\
\beta_i^1 &= x'_{5,i} x_{4,i} - x_4 x'_{5,i} x_{4,i+1} && 0 \leq i \leq 3 \\
\beta_i^2 &= x'_{5,i} x_4 x_5 - x_3 x'_{5,i} x_4 && 0 \leq i \leq 3 \\
\beta_{i,j}^3 &= x'_{5,i} x_{4,j} x'_{5,j} - x_3 x'_{5,i} x_{4,j} x'_{5,j+1} && 0 \leq i < j \leq 2 \\
\eta_i &= x'_{6,i} x'_{5,i} - x_5 x'_{6,i} x'_{5,i+1} && 0 \leq i \leq 4 \\
\xi_{i,j} &= x'_{6,i} x'_{5,j} x_6 - x_4 x'_{6,i} x'_{5,j} && 0 \leq i < j \leq 5 \\
\lambda_{i,j,k} &= x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k} - x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k+1} && 0 \leq i < j < k \leq 4 \\
\nu_{i,j,k,l} &= x'_{6,i} x'_{5,j} x_{4,k} x_5 x'_{6,l} x_5 - x_4 x'_{6,i} x'_{5,j} x_{4,k} x_5 x'_{6,l} && 0 \leq i < j < k \leq 3, 0 \leq l < k \\
\mu_{0,1,2,l,m} &= x'_{6,0} x'_{5,2} x_{4,2} x'_{5,3} x'_{6,l} x'_{5,m} x_4 - x_4 x'_{6,0} x'_{5,1} x_{4,2} x'_{5,3} x'_{6,l} x'_{5,m} && 0 \leq l < m \leq 2 \\
f &= x'_{6,0} x'_{5,1} x_{4,2} x'_{5,3} x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{5,l} x_{4,m} x'_{6,m} \\
&\quad - x_4 x'_{6,0} x'_{5,1} x_{4,2} x'_{5,3} x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{5,l} x_{4,m} x'_{6,m+1} && 0 \leq i < j < k < l \leq 5, 0 \leq l < m \leq 5
\end{aligned}$$

The Gröbner-Shirshov basis of the Coxeter group of the type E_8 is as follows:

$$\begin{aligned}
\theta_i^1 &= x_i^2 - 1 & -1 \leq i \leq 5 \\
\theta_i^2 &= x_{i+1}x_i x_{i+1} - x_i x_{i+1} x_i & -1 \leq i \leq 3 \\
\theta_{i,j}^3 &= x_i x_j - x_j x_i & -1 \leq j+1 < i \leq 4 \\
\theta_i^4 &= x_5 x_i - x_i x_5 & i \neq 3, 5, 6 \\
\theta^5 &= x_5 x_3 x_5 - x_3 x_5 x_3 \\
\varepsilon^1 &= x_6^2 - 1 \\
\varepsilon^2 &= x_6 x_4 x_6 - x_4 x_6 x_4 \\
\varepsilon_i^3 &= x_6 x_i - x_i x_6 & i \neq 4, 6 \\
\alpha_{i,j}^1 &= x_{i+1,j} x_{i+1} - x_i x_{i+1,j} & -1 \leq j < i \leq 3 \\
\alpha_{5,j}^2 &= x'_{5,j} x_5 - x_3 x'_{5,j} & -1 \leq j \leq 2 \\
\beta_i^1 &= x'_{5,i} x_{4,i} - x_4 x'_{5,i} x_{4,i+1} & -1 \leq i \leq 3 \\
\beta_i^2 &= x'_{5,i} x_4 x_5 - x_3 x'_{5,i} x_4 & -1 \leq i \leq 3 \\
\beta_{i,j}^3 &= x'_{5,i} x_{4,j} x'_{5,j} - x_3 x'_{5,i} x_{4,j} x'_{5,j+1} & -1 \leq i < j \leq 2 \\
\eta_i &= x'_{6,i} x'_{5,i} - x_5 x'_{6,i} x'_{5,i+1} & -1 \leq i \leq 4 \\
\xi_{i,j} &= x'_{6,i} x'_{5,j} x_6 - x_4 x'_{6,i} x'_{5,j} & -1 \leq i < j \leq 5 \\
\lambda_{i,j,k} &= x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k} \\
&\quad - x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{6,k+1} & -1 \leq i < j < k \leq 4 \\
\nu_{i,j,k,l} &= x'_{6,i} x'_{5,j} x_{4,k} x_5 x'_{6,l} x_5 \\
&\quad - x_4 x'_{6,i} x'_{5,j} x_{4,k} x_5 x'_{6,l} & -1 \leq i < j < k \leq 3, -1 \leq l \leq k \\
\mu_{0,1,2,l,m} &= x'_{6,0} x'_{5,2} x_{4,2} x'_{5,3} x'_{6,l} x'_{5,m} x_4 \\
&\quad - x_4 x'_{6,0} x'_{5,1} x_{4,2} x'_{5,3} x'_{6,l} x'_{5,m} & -1 \leq l < m \leq 2 \\
\delta_{i,j,k,l,m,p} &= x'_{6,i} x'_{5,j} x_{4,k} x'_{5,3} x_4 x'_{6,l} x'_{5,m} x_4 x'_{6,p} x_{6,k+3} \\
&\quad - x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{5,3} x_4 x'_{6,l} x'_{5,m} x_4 x'_{6,p} x_{6,k+4} \\
& & -1 \leq i < j < k \leq 2, -1 \leq l < m < p \leq k+2 \\
\rho_{i,j,k,l,m} &= x'_{6,i} x'_{5,j} x_{4,k} x'_{5,2} x'_{6,l} x'_{5,m} x_4 x'_{6,p} x_{5,k+3} \\
&\quad - x_4 x'_{6,i} x'_{5,j} x_{4,k} x'_{5,2} x'_{6,l} x'_{5,m} x_4 x'_{6,p} x_{5,k+4} \\
& & -1 \leq i < j < k \leq 1, -1 \leq l < m < p \leq k+2 \\
\varphi_1 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,3} x_4 x'_{6,l} x'_{5,m} x_4 x'_{5,p} x'_{6,q} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,3} x_4 x'_{6,l} x'_{5,m} x_4 x'_{5,p} x'_{6,q} x_{6,q+1} & -1 \leq l < m < p < q \leq 4 \\
\varphi_2 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_4 x'_{6,l} x'_{5,m} x_4 x'_{5,p} x'_{6,q} x_{6,p} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_4 x'_{6,l} x'_{5,m} x_4 x'_{5,p} x'_{6,q} x_{6,p+1} & -1 \leq l < m < p < q \leq 4
\end{aligned}$$

$$\begin{aligned}
f_1 &= x'_{6,i_1} x'_{5,i_2} x_{4,2} x'_{5,3} x_4 x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_6 \\
&\quad - x_4 x'_{6,i_1} x'_{5,i_2} x_{4,2} x'_{5,3} x_4 x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} && -1 \leq i_1 < i_2 \leq 1, -1 \leq j_1 < \dots < j_4 \leq 4 \\
f_2 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{6,j_2} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{6,j_2} + 1 && -1 \leq j_1 < \dots < j_4 \leq 4 \\
f_3 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{6,j_1} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{6,j_1} + 1 && -1 \leq j_1 < \dots < j_4 \leq 4 \\
f_4 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_3} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_3} + 1 && -1 \leq j_1 < \dots < j_5 \leq 4 \\
f_5 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_2} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_2} + 1 && -1 \leq j_1 < \dots < j_5 \leq 4 \\
f_6 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,3} x_4 x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_4} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,3} x_4 x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_4} + 1 && -1 \leq j_1 < \dots < j_5 \leq 4 \\
f_7 &= x'_{6,i_1} x'_{5,i_2} x_{4,2} x'_{5,3} x_4 x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_5} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_5} + 1 && -1 \leq i_1 < i_2 \leq 1, -1 \leq j_1 < \dots < j_5 \leq 4 \\
f_8 &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_1} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x_{4,j_5} x'_{6,j_1} + 1 && -1 \leq j_1 < \dots < j_5 \leq 4 \\
f_9 &= x'_{6,i_1} x'_{5,i_2} x_{4,2} x'_{5,3} x_4 x'_{5,j_1} x'_{5,j_2} x_{4,3} x'_{5,k_1} x_5 \\
&\quad - x_4 x'_{6,i_1} x'_{5,i_2} x_{4,2} x'_{5,3} x_4 x'_{5,j_1} x'_{5,j_2} x_{4,3} x'_{5,k_1} x_5 && -1 \leq i_1 < i_2 \leq 1, -1 \leq k_1 \leq 3 \\
f_{10} &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,i_4} x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,k_1} x'_{5,i_4} \\
&\quad - x_4 x'_{6,i_1} x'_{5,i_2} x_{4,1} x'_{5,i_4} x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,k_1} x'_{5,i_4} + 1 && -1 \leq 1 < i_4 \leq 3, -1 \leq k_1 < i_4 \\
f_{11} &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,k_1} x'_{5,1} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,k_1} x'_{5,2} && -1 \leq k_1 \leq 0 \\
f_{12} &= x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{5,0} \\
&\quad - x_4 x'_{6,-1} x'_{5,0} x_{4,1} x'_{5,2} x_{4,3} x'_{5,j_1} x'_{5,j_2} x_{4,j_3} x'_{5,j_4} x'_{5,0} &&
\end{aligned}$$

Remark For the Coxeter-Dynkin diagram of E_n which is as Figure 4, let $n = 9$, it is actually the affine Coxeter group \widetilde{E}_8 . The presentation of group E_n we defined can also be used to calculate the Gröbner-Shirshov basis of affine Coxeter group \widetilde{E}_8 via Mathematica.

Acknowledgements

The author would like to thank Professor X.Zhao and Doctor X.Liu for their guidances, useful discussions in writing up this paper.

References

- [1] **L A Bokut, Y Chen, A Obul**, *Some new results on Gröbner-Shirshov bases for Lie algebras and around*, Internat. J. Algebra Comput. 28 (2018) 1403–1423
- [2] **L A Bokut, Y Fong, W-F Ke**, *Gröbner-Shirshov bases and composition lemma for associative conformal algebras: an example*, from “Combinatorial and computational algebra (Hong Kong, 1999)”, Contemp. Math. 264, Amer. Math. Soc., Providence, RI (2000) 63–90
- [3] **L A Bokut, Y Fong, W-F Ke**, *Composition-diamond lemma for associative conformal algebras*, J. Algebra 272 (2004) 739–774
- [4] **L A Bokut, S-J Kang, K-H Lee, P Malcolmson**, *Gröbner-Shirshov bases for Lie superalgebras and their universal enveloping algebras*, J. Algebra 217 (1999) 461–495
- [5] **L A Bokut, A A Klein**, *Serre relations and Gröbner-Shirshov bases for simple Lie algebras. I, II*, Internat. J. Algebra Comput. 6 (1996) 389–400, 401–412
- [6] **L A Bokut, A A Klein**, *Gröbner-Shirshov bases for the exceptional Lie algebras E_6, E_7, E_8* , from “Algebras and combinatorics (Hong Kong, 1997)”, Springer, Singapore (1999) 37–46
- [7] **L Bokut, P Malcolmson**, *Gröbner-Shirshov bases for quantum enveloping algebras*, Israel J. Math. 96 (1996) 97–113
- [8] **L A Bokut, L-S Shiao**, *Gröbner-Shirshov bases for Coxeter groups*, Comm. Algebra 29 (2001) 4305–4319 Special issue dedicated to Alexei Ivanovich Kostrikin
- [9] **L Bokut, A Vesnin**, *Gröbner-Shirshov bases for some braid groups*, J. Symbolic Comput. 41 (2006) 357–371
- [10] **N Bourbaki**, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin (2002) Translated from the 1968 French original by Andrew Pressley
- [11] **B Buchberger**, *An algorithm for finding the basis elements of the residue class ring of a zero dimensional polynomial ideal*, J. Symbolic Comput. 41 (2006) 475–511 Translated from the 1965 German original by Michael P. Abramson
- [12] **Y Chen, C Liu**, *Gröbner-Shirshov bases for Coxeter groups I*, arXiv 0910.0096 (2009)
- [13] **H S M Coxeter**, *The Complete Enumeration of Finite Groups of the Form $R_i^2 = (R_i R_j)^{k_{ij}} = 1$* , Journal of the London Mathematical Society s 1-10 (1935) 21–25

- [14] **B Helton, M de Oliveira**, *NCAgebra*, <https://github.com/NCAgebra/NC>
- [15] **D Lee**, *Gröbner-Shirshov bases and normal forms for the Coxeter groups E_6 and E_7* , from “Advances in algebra and combinatorics”, World Sci. Publ., Hackensack, NJ (2008) 243–255
- [16] **J-Y Lee, D-i Lee**, *Gröbner-Shirshov bases for non-crystallographic Coxeter groups*, Acta Crystallogr. Sect. A 75 (2019) 584–592
- [17] **A I Shirshov**, *Some algorithm problems for Lie algebras*, Sibirsk. Mat. Ž. 3 (1962) 292–296
- [18] **A I Shirshov**, *Selected works of A. I. Shirshov*, Contemporary Mathematicians, Birkhäuser Verlag, Basel (2009) Translated from the Russian by Murray Bremner and Mikhail V. Kotchetov, Edited by Leonid A. Bokut, Victor Latyshev, Ivan Shestakov and Efim Zelmanov

Capital Normal University

School of Mathematical Science, Capital Normal University, 105 West Third Ring Road North, Haidian District, Beijing, 100048, China

wjun@cnu.edu.cn