

REGULARIZED PERIODS AND THE GLOBAL GAN–GROSS–PRASAD CONJECTURE : THE CASE OF $U(n + 2r) \times U(n)$

JAEHO HAAN

ABSTRACT. In this paper, we introduce regularized trilinear periods on certain non-reductive groups. It has two direct applications. Firstly, it enables us to define the regularized Bessel periods and the regularized Fourier-Jacobi periods for all classical and metaplectic groups. Secondly, by using the properties of the regularized Fourier-Jacobi periods, we can prove one direction of the full Gan–Gross–Prasad conjecture on skew-hermitian unitary groups.

CONTENTS

1.	Introduction	1
2.	General notation	6
3.	Automorphic forms on $N \rtimes G'$	7
4.	The GGP triplets	9
5.	Mixed truncation	12
6.	Regularized trilinear periods	16
7.	Jacquet modules corresponding to Fourier-Jacobi characters	22
8.	Eisenstein series and residual representation	25
9.	Reciprocal non-vanishing of the Fourier–Jacobi periods	26
10.	Proof of the Main Theorem	35
	References	36

1. Introduction

1.1. General introduction. The study of special L -values is a pivotal domain in modern number theory, because special L -values encode a considerable amount of arithmetic information. Among all other special values, central L -values are of particular interest as the Birch and Swinnerton-Dyer conjecture and its generalization imply.

In 1992, Gross and Prasad [16] proposed a fascinating conjecture on the relationship between automorphic period integrals and central L -values of certain tensor product L -functions on special orthogonal groups. Subsequently, together with Gan in 2012 [18], they generalized the conjecture to all classical

Date: November 3, 2021.

Key words and phrases. Gan-Gross-Prasad conjecture, Arthur truncation, regularized periods, Rankin-Selberg L -function, central L -values.

groups and metaplectic groups, which is now referred to as the global *Gan–Gross–Prasad* (GGP) conjecture.

There are two types of periods in the global GGP conjecture: the Bessel periods and the Fourier–Jacobi periods. The Bessel periods are periods of automorphic forms on orthogonal groups or hermitian unitary groups and the Fourier–Jacobi periods are those of automorphic forms on metaplectic-symplectic groups or skew hermitian unitary groups. We briefly recall the definition of both periods at least for (skew) hermitian unitary groups.

Let E and F be quadratic extensions of number fields with adèle rings \mathbb{A}_E and \mathbb{A}_F , respectively. Let ω be the non-trivial quadratic character of $F^\times \backslash \mathbb{A}_F^\times$ associated with E/F by the global class field theory. Fix a unitary character μ of $E^\times \backslash \mathbb{A}_E^\times$ such that $\mu|_{\mathbb{A}_F^\times} = \omega$. Sometimes, we view μ as a character of $GL_n(\mathbb{A}_E)$ and in that case, it refers to $\mu \circ \det$. We also fix a nontrivial continuous character ψ of $E \backslash \mathbb{A}_E$. For $\epsilon \in \{\pm\}$, let $W_m \subset W_n$ be non-degenerate m and n -dimensional ϵ -Hermitian spaces over E such that

- $\epsilon \cdot (-1)^{\dim W_m^\perp} = -1$;
- W_m^\perp in W_n is a split space.

Let G_n, G_m be the isometry groups of W_n, W_m , respectively. Regard G_m as a subgroup of G_n , which acts trivially on the orthogonal complement of W_m in W_n . Let $N_{n,r}$ be some unipotent subgroup of G_n on which G_m acts through conjugation. Denote $H = N_{n,r} \rtimes G_m$. When $n - m = 2r + 1$, there is an essentially unique generic character $\chi' : N_{n,r}(\mathbb{A}_F) \backslash N_{n,r}(\mathbb{A}_F) \rightarrow \mathbb{C}^\times$ that is stabilized by $G_m(\mathbb{A}_F)$ and hence can be extended to a character of $H(F) \backslash H(\mathbb{A}_F)$. When $n - m = 2r$, there is a global generic Weil representation $\nu_{\psi^{-1}, \mu^{-1}, W_m}$ of $H(\mathbb{A}_F)$ realized on Schwartz space $\mathcal{S} = \mathcal{S}(Y(\mathbb{A}_F))$, where Y is a Lagrangian F -space of $\text{Res}_{E/F} W_m$.

For each $f \in \mathcal{S}$, the associated theta series is defined by

$$\Theta_{\psi^{-1}, \mu^{-1}, W_m}(h, f) = \sum_{x \in Y(F)} (\nu_{\psi^{-1}, \mu^{-1}, W_m}(h)f)(x), \quad h \in H(\mathbb{A}_F).$$

The space of theta series gives an $H(F)$ -invariant automorphic realization of $\nu_{\psi^{-1}, \mu^{-1}, W_m}$. From now on, we denote $G(F) \backslash G(\mathbb{A}_F)$ by $[G]$ for each algebraic group G defined over F .

Let π_1, π_2 be two irreducible cuspidal automorphic representations of $G_n(\mathbb{A}_F)$ and $G_m(\mathbb{A}_F)$, respectively. We regard H as a subgroup of G_n through the map $(n, g) \rightarrow ng$. Depending on whether $n - m$ is odd or even, the *Bessel periods* and the *Fourier–Jacobi periods* are defined to be the following integrals:

- If $n - m$ is odd, for $\varphi_1 \in \pi_1, \varphi_2 \in \pi_2$, the Bessel period for φ_1, φ_2 is defined by

$$\mathcal{B}_{\chi'}(\varphi_1, \varphi_2) := \int_{[N_{n,r} \rtimes G_m]} \varphi_1(ng) \varphi_2(g) \chi'^{-1}((n, g)) dndg.$$

- If $n - m$ is even, for $\varphi_1 \in \pi_1, \varphi_2 \in \pi_2, f \in \nu_{\psi^{-1}, \mu^{-1}, W_m}$, the Fourier–Jacobi period φ_1, φ_2 is defined by

$$\mathcal{FJ}_{\psi, \mu}(\varphi_1, \varphi_2, f) := \int_{[N_{n,r} \rtimes G_m]} \varphi_1(ng) \varphi_2(g) \Theta_{\psi^{-1}, \mu^{-1}, W_m}((n, g), f) dndg.$$

The GGP conjecture [18, Conjecture 24.1] predicts that for an irreducible tempered cuspidal representation $\pi_1 \boxtimes \pi_2$ of $G_n \times G_m$, the following two conditions are equivalent in each case. In what follows, BC stands for the standard base change and the L -functions are the Rankin–Selberg L -functions.

- **Bessel case**
 - (i) $\mathcal{B}_{\chi'}$ is non-vanishing,

(ii) $\text{Hom}_{H(F_v)}(\pi_{1,v} \boxtimes \pi_{2,v}, \mathbb{C}) \neq 0$ for all places v and $L(\frac{1}{2}, BC(\pi_1) \times BC(\pi_2)) \neq 0$.

• **Fourier-Jacobi case**

- (i) $\mathcal{FJ}_{\psi, \mu}$ is non-vanishing,
- (ii) $\text{Hom}_{H(F_v)}(\pi_{1,v} \boxtimes \pi_{2,v} \boxtimes \nu_{\psi_v^{-1}, \mu_v^{-1}, W_m}, \mathbb{C}) \neq 0$ for all places v and $L(\frac{1}{2}, BC(\pi_1) \times BC(\pi_2) \otimes \mu^{-1}) \neq 0$.

In their influential paper [28], Ichino–Ikeda formulated a refined GGP conjecture expressing the Bessel periods explicitly in terms of special values for orthogonal groups of co-rank 1, i.e., the $SO(n+1) \times SO(n)$ case. Thereafter, Liu [44] extended their conjecture to higher co-rank cases for both orthogonal groups and hermitian unitary groups and Xue [56] formulated the corresponding refined conjecture for metaplectic-symplectic groups. The refined GGP conjecture implies the (i) \rightarrow (ii) direction of the original GGP conjecture.

After the formulation of both the original and refined GGP conjectures, there has been huge progress towards the low co-rank (i.e., co-rank ≤ 1) GGP conjectures until recently. Wei Zhang was the first who made a breakthrough toward the low co-rank GGP conjectures. Using the relative trace formula of Jacquet and Rallis [32], he proved the original and refined GGP conjectures for the co-rank 1 hermitian unitary groups, i.e., $U(n+1) \times U(n)$ under two local assumptions ([61, 62]). Shortly thereafter, following a similar idea of W. Zhang, H. Xue [57, 55] proved the original and refined GGP conjectures for the co-rank 0 skew-hermitian unitary groups, i.e., $U(n) \times U(n)$, under the similar local assumptions. Those two local assumptions arise when one applies a simple version of the Jacquet–Rallis relative trace formula. The first condition has now been completely dropped in the recent work of R. Beuzart-Plessis, Y. Liu, W. Zhang and X. Zhu [9], which proves the original and refined GGP conjectures for $U(n+1) \times U(n)$ in the stable case. However, to remove the second condition, which forces $BC(\pi_1) \boxtimes BC(\pi_2)$ to be cuspidal, was regarded very hard for a long time because it would require the fine spectral expansion of the Jacquet–Rallis relative trace formula. In a striking recent paper [7], R. Beuzart-Plessis, Pierre-Henri Chaudouard and M. Zydor settled the original GGP conjecture and refined GGP conjectures for $U(n+1) \times U(n)$ in all endoscopic cases circumventing the fine spectral expansion of the Jacquet–Rallis relative trace formula. Their proof combines many results (including [9]) concerning the Jacquet–Rallis relative trace formula in a very clever way. As a consequence, the original and refined GGP conjectures for $U(n+1) \times U(n)$ have been completely settled. However, it is not clear whether such an approach using the Jacquet–Rallis’s relative trace formula is available for other classical groups other than unitary groups.

On the other hand, there has been a different approach towards the low co-rank GGP conjectures. Inspired by the construction of the regularized period integral of Jacquet–Lapid–Rogawski [31, 43], Ichino–Yamana [30] and Yamana [58] invented regularized period integrals for co-rank 1 hermitian unitary groups, i.e. $U(n+1) \times U(n)$ and for co-rank 0 metaplectic–symplectic groups, i.e. $Mp(2n) \times Sp(2n)$, respectively. By computing the regularized period involving certain residual Eisenstein series, they proved the (i) \rightarrow (ii) direction of the original GGP conjecture in such low co-rank cases without any local assumptions. It is also worthwhile to mention that the Ichino–Yamana’s mixed truncation for $GL(n+1) \times GL(n)$ (resp. $U(n+1) \times U(n)$) invented in [29] (resp. [30]) was also used in [7].

For the higher co-rank (i.e., co-rank > 1) GGP conjecture, there has been a significant advancement in the Bessel case recently. In their seminal paper, under some hypothesis, Jiang and Zhang [34, Theorem 5.7] proved the (i) \rightarrow (ii) direction of the original full GGP conjecture in all cases and the (ii) \rightarrow (i) direction in some special cases for both orthogonal groups and hermitian unitary groups. Their method crucially uses the *reciprocal non-vanishing of Bessel periods* [34, Theorem 5.3] to establish the global non-vanishing of the twisted automorphic descent which extends the automorphic descent developed by Ginzburg–Rallis–Soudry [22]. Since the proof of [34, Theorem 5.3] heavily depends on the unproven general calculation

[34, Theorem 4.8] of the unramified local zeta integrals defined using local Bessel functionals, their proof of the (i) \rightarrow (ii) direction of the GGP conjecture is also conditional on such hypothetical computation. Though some special cases of [34, Theorem 4.8] were carried out in several papers, for example, [23], [38], [47], [48], [49], the general computation, which is required to prove [34, Theorem 5.7], does not seem to be easy. (As written in [35], since 2018, Jiang, Zhang and Soudry have been preparing a paper [33], which treats the computation of them in full generality for quasi-split classical groups. Their preprint, however, has not appeared so [34, Theorem 5.7] is still conditional.) We also mention that Furusawa–Morimoto [12],[13] recently proved the original and refined GGP conjecture for certain special irreducible tempered representations of $SO(2n+1) \times SO(2)$.

In the Fourier–Jacobi case, on the other hand, the most well-known result for the higher co-rank GGP conjecture is the one by Ginzburg–Jiang–Rallis on metaplectic-symplectic groups [15] which has been recently supplemented with [14]. In that paper, by combining the Arthur truncation method and the Rankin-Selberg method, they proved the (i) \rightarrow (ii) direction of the original GGP conjecture for arbitrary even co-rank metaplectic-symplectic groups, i.e., $Mp(2n+2r) \times Sp(2n)$ under a stability assumption. The assumption forces the Langlands functorial transfer of $\pi_1 \boxtimes \pi_2$ to their corresponding general linear groups to be cuspidal. Since this assumption is critical in their argument, it appears hard to remove it while adopting their approach. (See the remarks after [34, Theorem 5.7].) On the other hand, it is plausible to apply the similar approach used in [34] to the Fourier–Jacobi case. Indeed, as it is written in [36], Jiang and Zhang have been preparing a paper [37] concerning the Fourier–Jacobi case for metaplectic-symplectic groups since 2015. However no preprint seems available yet, and in any case, the results would be conditional on the non-existent results in [33] if they use the same method as in [34].

In contrast to the metaplectic-symplectic case, very little is known for non-equal rank skew-hermitian unitary groups. To the best of the author’s knowledge, Gelbart–Rogawski’s paper [21] which studied the Fourier-Jacobi period of $U(3) \times U(1)$ is the only work that deals with non-equal rank skew-hermitian groups.

The main goal of this paper is to prove the (i) \rightarrow (ii) direction of the original GGP conjecture for skew-hermitian unitary groups, i.e., $U(n+2r) \times U(n)$ without any hypothesis. To do this, we first define regularized trilinear periods which involve an integration on a unipotent subgroup. This enables us to construct the regularized Bessel and Fourier-Jacobi periods for all classical and metaplectic groups. In particular, when $r = 0$, these are exactly the same regularized periods introduced in [30] and [58], respectively. By exploiting several properties of regularized Fourier–Jacobi periods, we prove the following:

Main Theorem. *Let $n - m = 2r$ for some non-negative integer r and π_1, π_2 be irreducible tempered cuspidal automorphic representations of $G_n(\mathbb{A}_F)$ and $G_m(\mathbb{A}_F)$ respectively. If there are $\varphi_1 \in \pi_1, \varphi_2 \in \pi_2$ and $f \in \nu_{W_m}$ such that*

$$\mathcal{FJ}_{\psi, \mu}(\varphi_1, \varphi_2, f) \neq 0,$$

then $L(\frac{1}{2}, BC(\pi_1) \times BC(\pi_2) \otimes \mu^{-1}) \neq 0$.

We remark on the proof of the main theorem.

Remark 1.1. Based on the works of Arthur ([4]), Mok ([42]) and Kaletha–Minguez–Shin–White ([39]) on the endoscopic classification for unitary groups, the irreducible tempered representations π_1, π_2 have the following properties:

- (i) the weak base changes $BC(\pi_1)$ and $BC(\pi_2)$ exist.
- (ii) $BC(\pi_1)$ is decomposed as the isobaric sum $\sigma_1 \boxplus \cdots \boxplus \sigma_t$, where σ_i ’s are irreducible cuspidal automorphic representations of general linear groups such that the (twisted) Asai L -function $L(s, \sigma_i, As^{(-1)^{n-1}})$ has a pole at $s = 1$ for all $1 \leq i \leq t$.

(iii) The local normalized intertwining operator is holomorphic for all z with $\operatorname{Re}(z) \geq \frac{1}{2}$.

In the course of proving our main theorem, we use the above three properties in a crucial manner.

Remark 1.2. The proof of our main theorem, to some extent, is in the same vein with [15] in that it uses the relation of non-vanishing of central L -values with the existence of the residual Eisenstein series. However, comparing with the proof of [15], there are two main innovations in our approach. First, to compute the Fourier–Jacobi period integrals involving residual Eisenstein series, Ginzburg, Jiang and Rallis [15] used the Arthur truncation method to make sense of a divergent integral involving Eisenstein series. In their approach using the Arthur truncation, the stability assumption on π_1, π_2 becomes indispensable. However, if we use the mixed truncation method to compute the regularized Fourier–Jacobi period involving residual Eisenstein series, we don’t need such a stability assumption on π_1, π_2 .

Secondly, though we took a different truncation from G–J–R’s, in order to relate the Fourier–Jacobi periods of residual Eisenstein series with the original Fourier–Jacobi periods, we reach the almost same place in G–J–R’s, that is, [15, Proposition 5.3]. (Compare with Lemma 9.7). To prove the direction (i) \rightarrow (ii) of the GGP conjecture, G–J–R employed the isobaric sum of $FL(\pi_1)$ (the Langlands functorial transfer of π_1). However, this choice of the isobaric sum makes the proof of [15, Proposition 5.3] very difficult. Around 2013, some experts found a serious gap in the original proof of [15, Proposition 5.3]. Thereafter, in 2019, G–J–R, together with B. Liu, uploaded their corrected proof of [15, Proposition 5.3] to arXiv. ([14].) But their corrected proof is rather involved and roundabout. In this paper, we overcome such difficulty by using the isobaric sum of $BC(\pi_2)$ instead of $BC(\pi_1)$. As a consequence, it shifts the difficulty of proving Lemma 9.7 for the isobaric sum of $BC(\pi_1)$ to that of Lemma 9.4 for the isobaric sum of $BC(\pi_2)$. However, the latter difficulty is manageable. Our arguments streamline the discussion substantially and are much easier to understand. (See Remark 9.8.)

Remark 1.3. Theorem 9.9 bears a resemblance to [34, Theorem 5.3], the reciprocal non-vanishing of Bessel periods. However, while the hypothesis on the unramified calculation of certain local zeta integrals is crucially used in the proof of [34, Theorem 5.3], we derive Theorem 9.9 without such a hypothesis. We also mention that Theorem 9.9 has a potential that can be used to prove the (ii) \rightarrow (i) direction of the GGP conjecture as demonstrated in [34, Cor. 6.11].

This paper is organized as follows. In Section 2, we fix some general notation and in Section 3, we give a definition of automorphic forms on certain non-reductive groups modeled after Jacobi forms. In Section 4, we explain the GGP triplets to which our mixed truncation is designed and in Section 5, we define the mixed truncation operator which sends an automorphic form on a certain non-reductive group to a rapidly decreasing automorphic form on a reductive group. Using this mixed truncation, we can define the regularized trilinear period integrals and the regularized Bessel and Fourier–Jacobi periods in Section 6. In Section 7, after introducing the Jacquet module corresponding to Fourier–Jacobi character, we prove a lemma which is required to prove a vanishing property of regularized periods involving certain residual Eisenstein series. In Section 8, we explain the definition of (residual) Eisenstein series and review [30, Proposition 5.3] which manifests the correlation of analytic properties of Eisenstein series with the behavior of the associated Rankin–Selberg L -functions. In Section 9, we prove various lemmas required to establish Theorem 9.9, which claims the reciprocal non-vanishing of the Fourier–Jacobi periods. By using Theorem 9.9, we prove our main theorem in Section 10.

We stress that separate approaches need to be taken to prove Lemma 7.1 and for Lemma 9.4 according to whether $r = 0$ or $r > 0$ because the relevant twisted Jacquet modules are totally different. Therefore, we cannot reduce the proof of our main theorem to a single one for all ranks. This is in contrast to the case of the local GGP conjecture, where one can reduce the local GGP for $r > 0$ to that of $r = 0$. (Compare with [18, Theorem 19.1].)

After completing this paper, we became aware that Gan, Gross and Prasad extended their original conjecture to non-tempered cases in their recent paper. In [17, Conjecture 9.1], they formulated the general global (resp. local) GGP conjecture for automorphic representations in global (resp. local) A -packets. However, automorphic representations in a non-tempered A -packet are not necessarily cuspidal. Therefore, to formulate the general GGP conjecture including non-tempered representations, one needs to regularize both $\mathcal{B}_{\chi'}$ and $\mathcal{FJ}_{\psi, \mu}$. For this reason, they formulated [17, Conjecture 9.1] assuming the existence of certain regularized period integrals in each case. With our regularized Bessel and Fourier–Jacobi period integrals in Definition 6.5, we can now state their general conjecture more explicitly. (See Remark 6.6.) We expect that properties of the regularized periods established in this paper will help prove the general GGP conjecture.

1.2. Some related problems. The methods used in this paper may be applicable to other problems. We discuss some of them here.

- (i) Using the same method of this paper, we can prove the (i) \rightarrow (ii) direction of the original GGP conjecture for higher co-rank metaplectic-symplectic groups without the stability assumption appearing in [15]. Since we defined the regularized Bessel periods in this paper, a similar approach is also possible to cover the Bessel cases. As our approach does not require the calculation of the unramified local zeta integrals, it will improve the result of D. Jiang and L. Zhang [34, Theorem 5.7]. Our subsequent papers [24], [26] are devoted to them.
- (ii) In [7], R. Beuzart-Plessis, P-H. Chaudouard and M. Zydor calculated a certain term appearing in the coarse spectral expansion of the Jacquet–Rallis trace formula for linear groups in terms of the Rankin–Selberg periods of Eisenstein series and the Flicker–Rallis intertwining periods. In their computation, the Ichino–Yamana’s mixed truncation for $(GL(n+1) \times GL(n))$ in [30] is crucially used to define certain truncated integrals involving a relevant kernel function and Eisenstein series. Since this computation is one of the key innovation of their paper, to extend their result to attack the higher co-rank GGP conjecture for unitary groups $(U(n+2r+1) \times U(n))$, the mixed truncation for $(GL(n+2r+1) \times GL(n))$ seems to be necessary. Though such a mixed truncation for higher co-rank general linear groups can also be defined similarly, its very definition and proof are slightly different from those of classical groups. For this reason, we deal with the construction of the mixed truncation for linear groups separately in our future work [27].
- (iii) Our strategy also works for some important non-tempered cases of the GGP conjecture. In [25], we will study the regularized Bessel periods involving Eisenstein series and establish the (i) \rightarrow (ii) direction of the non-tempered GGP conjecture in this case.

2. General notation

Throughout the paper, we fix some general notation and convention. Let F be a number field with adèle ring \mathbb{A}_F . For a place v of F , denote by F_v the localization of F at v . Let G be a connected reductive algebraic group over F and fix a minimal F -parabolic subgroup P_0 of G with a Levi decomposition $P_0 = M_0U_0$. We call any F -parabolic subgroup $P = MU$ of G which contains P_0 *standard*. Fix a maximal compact subgroup $K = \prod_v K_v$ of $G(\mathbb{A}_F)$ which satisfies $G(\mathbb{A}_F) = P_0(\mathbb{A}_F)K$ and for every standard parabolic subgroup $P = MU$ of G ,

$$P(\mathbb{A}_F) \cap K = (M(\mathbb{A}_F) \cap K)(U(\mathbb{A}_F) \cap K)$$

and $M(\mathbb{A}_F) \cap K$ is still maximal compact in $M(\mathbb{A}_F)$ (see [41, I.1.4]).

We give the Haar measure on $U(\mathbb{A}_F)$ so that the volume of $[U]$ is 1 and give the Haar measure on K so that the total volume of K is 1. We also choose Haar measures on $M(\mathbb{A}_F)$ for all Levi subgroups M of G compatibly with respect to the Iwasawa decomposition.

Let \mathbf{G} be a topological group that is a finite central covering of $G(\mathbb{A}_F)$, i.e., there exists a surjective morphism $pr : \mathbf{G} \rightarrow G(\mathbb{A}_F)$, whose kernel is a finite subgroup of the center of \mathbf{G} and pr a topological covering. It is known that $U_0(\mathbb{A}_F)$ lifts canonically into \mathbf{G} (see [41, Appendix I]). A result of Weil [54] says that $G(F)$ lifts uniquely into $\mathbf{G}(\mathbb{A}_F)$. For the unipotent subgroup U of any parabolic subgroup of G (or a subgroup S of $G(F)$), we use the same notation U (or S) to denote their image of such lifting. For a subgroup J of $G(\mathbb{A}_F)$, we use the boldface \mathbf{J} to denote $pr^{-1}(J)$. Note that parabolic subgroups of \mathbf{G} are $\mathbf{P} = \mathbf{M}U$. Sometimes, we view a function defined on J as a function on \mathbf{J} by composing them with pr .

Let $\mathcal{A}_P(\mathbf{G})$ be the space of automorphic forms on $U(\mathbb{A}_F)P(F)\backslash\mathbf{G}(\mathbb{A}_F)$, i.e., smooth, \mathbf{K} -finite and \mathfrak{z} -finite functions on $U(\mathbb{A}_F)P(F)\backslash\mathbf{G}(\mathbb{A}_F)$ of moderate growth, where \mathfrak{z} is the center of the universal enveloping algebra of the complexified Lie algebra of $G(F \otimes_{\mathbb{Q}} \mathbb{R})$. When $P = G$, we simply write $\mathcal{A}(\mathbf{G})$ for $\mathcal{A}_G(\mathbf{G})$. For a cuspidal automorphic representation ρ of $\mathbf{M}(\mathbb{A}_F)$, write $\mathcal{A}_P^\rho(\mathbf{G})$ for the subspace of functions $\phi \in \mathcal{A}_P(\mathbf{G})$ such that for all $k \in \mathbf{K}$, the function $m \rightarrow \delta_P(m)^{-\frac{1}{2}} \cdot \phi(mk)$ belongs to the space of ρ . (Here, δ_P is the modulus function of $P(\mathbb{A}_F)$ (see [41, I.2.17]).)

3. Automorphic forms on $N \rtimes \mathbf{G}'$

In this section, we extend the definition of automorphic forms from reductive groups to certain non-reductive groups. Let G' be a connected reductive group defined over F . We will use similar notations for subgroups of G' by adding a prime $'$ on those of G . Let N be a unipotent F -group which admits a G' -action in the category of algebraic groups over F . We denote this action by $\sigma : G' \rightarrow \text{Aut}(N)$. Using σ , we consider the semi-direct product $N \rtimes \mathbf{G}'$ and define automorphic forms on $N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F)$ generalizing the properties of Jacobi forms in [51, Proposition 4.1.2].

Let $\|\cdot\|$ be a norm function on $G'(\mathbb{A}_F)$ as in [41, Section I.2.2].

Definition 3.1. For a function $\varphi : N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F) \rightarrow \mathbb{C}$, we say that φ is an *automorphic form* if

- $\varphi((\delta, \gamma) \cdot (n, ug)) = \varphi(n, g)$ for $u \in U'(\mathbb{A}_F)$ and $(\delta, \gamma) \in N(F) \rtimes P'(F)$;
- φ is smooth;
- φ is of moderate growth;
- φ is right \mathbf{K}' -finite;
- φ is \mathfrak{z}' -finite.

Here, φ is said to be of *moderate growth* if there is a positive integer m and constant C such that

$$|\varphi(n, g)| < C\|g\|^m$$

for all $(n, g) \in N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F)$.

We denote by $\mathcal{A}_{P'}(N \rtimes \mathbf{G}')$ the space of automorphic forms on $N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F)$ satisfying the above properties. When $P' = G'$, we simply write $\mathcal{A}(N \rtimes \mathbf{G}')$ for $\mathcal{A}_{G'}(N \rtimes \mathbf{G}')$. Note that if N is the trivial group $\mathbf{1}$, $\mathcal{A}_{P'}(\mathbf{1} \rtimes \mathbf{G}')$ is just $\mathcal{A}_{P'}(\mathbf{G}')$.

For $\varphi \in \mathcal{A}(N \rtimes \mathbf{G}')$, define $\varphi_{P'} : N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F) \rightarrow \mathbb{C}$ by

$$\varphi_{P'}(n, g) := \int_{[U']} \varphi(n, ug) du \quad \text{for } (n, g) \in N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F).$$

It is obvious that $\varphi_{P'} \in \mathcal{A}_{P'}(N \rtimes \mathbf{G}')$. If we regard $\varphi \in \mathcal{A}(\mathbf{1} \rtimes \mathbf{G}')$ as $\varphi \in \mathcal{A}(\mathbf{G}')$, then $\varphi_{P'}$ is the usual constant term of $\varphi \in \mathcal{A}(\mathbf{G}')$ along P' defined by

$$\varphi_{P'}(g) = \int_{[U']} \varphi(ug) du.$$

Like automorphic forms on \mathbf{G}' , we can decompose automorphic forms on $N \rtimes \mathbf{G}'$. To describe it precisely, we introduce some notations.

Write $X(G')$ for the group of F -rational characters of G' . Let $(\mathfrak{a}'_0)^*$ be the \mathbb{R} -vector space spanned by the lattice $X(M'_0)$ and $\mathfrak{a}'_0 = \text{Hom}((\mathfrak{a}'_0)^*, \mathbb{R})$ its dual space. The canonical pairing on $(\mathfrak{a}'_0)^* \times \mathfrak{a}'_0$ is denoted by $\langle \cdot, \cdot \rangle$. Let Δ'_0 and $(\Delta'_0)^\vee = \{\alpha^\vee : \alpha \in \Delta'_0\}$ be the sets of simple roots and simple coroots in $(\mathfrak{a}'_0)^*$ and \mathfrak{a}'_0 respectively. Write $(\hat{\Delta}'_0)^\vee$ and $\hat{\Delta}'_0$ for the dual bases of Δ'_0 and $(\Delta'_0)^\vee$, respectively. (In other words, $(\hat{\Delta}'_0)^\vee$ and $\hat{\Delta}'_0$ are the set of coweights and weights, respectively.) For a standard parabolic subgroup $P' = M_{P'}U_{P'}$ of G' , write $A_{M_{P'}}$ for the maximal split torus in the center of $M_{P'}$ and $\mathfrak{a}_{P'}^* = X(M_{P'}) \otimes_{\mathbb{Z}} \mathbb{R}$ and $\mathfrak{a}_{P'}$ for its dual space.

Let $Q' \subset P'$ be a pair of standard parabolic subgroups of G' . There is a canonical injection $\mathfrak{a}_{P'} \hookrightarrow \mathfrak{a}_{Q'}$ and a canonical surjection $\mathfrak{a}_{Q'} \rightarrow \mathfrak{a}_{P'}$ induced by two inclusion maps $A_{M_{P'}} \hookrightarrow A_{M_{Q'}}$ and $M_{Q'} \hookrightarrow M_{P'}$. So we have a canonical decomposition

$$\mathfrak{a}_{Q'} = \mathfrak{a}_{Q'}^{P'} \oplus \mathfrak{a}_{P'}, \quad \mathfrak{a}_{Q'}^* = (\mathfrak{a}_{Q'}^{P'})^* \oplus \mathfrak{a}_{P'}^*.$$

In particular, if we take $Q' = P'_0$, we have a decomposition

$$\mathfrak{a}_0 = \mathfrak{a}_0^{P'_0} \oplus \mathfrak{a}_{P'_0}, \quad \mathfrak{a}_0^* = (\mathfrak{a}_0^{P'_0})^* \oplus \mathfrak{a}_{P'_0}^*.$$

Let $\Delta_{P'} \subset \Delta'_0$ be the set of non-trivial restrictions of simple roots to $\mathfrak{a}_{P'}^*$. We denote by $\Delta_{Q'}^{P'}$ the subset of $\Delta_{Q'}$ appearing in the root decomposition of the Lie algebra of the unipotent radical $U_{Q'} \cap M_{P'}$. Then for $H \in \mathfrak{a}_{P'}$, $\langle \alpha, H \rangle = 0$ for all $\alpha \in \Delta_{Q'}^{P'}$ and so $\Delta_{Q'}^{P'} \subset (\mathfrak{a}_{Q'}^{P'})^*$. Note that $\Delta_{P'}^{G'} = \Delta_{P'}$. For any $\alpha \in \Delta_{Q'}^{P'}$, there is an $\tilde{\alpha} \in \Delta'_0$ whose restriction to $(\mathfrak{a}_{Q'}^{P'})^*$ is α . Write α^\vee for the projection of $\tilde{\alpha}^\vee$ to $\mathfrak{a}_{Q'}^{P'}$. Define

$$(\Delta_{Q'}^{P'})^\vee = \{\alpha^\vee \mid \alpha \in \Delta_{Q'}^{P'}\}.$$

Define $\hat{\Delta}_{Q'}^{P'} \subset (\mathfrak{a}_{Q'}^{P'})^*$ and $(\hat{\Delta}^\vee)_{Q'}^{P'} \subset \mathfrak{a}_{Q'}^{P'}$ to be the dual bases of $(\Delta_{Q'}^{P'})^\vee$ and $\Delta_{Q'}^{P'}$ respectively. Thus, $\hat{\Delta}_{Q'}^{P'}$ is the set of simple weights and $(\hat{\Delta}^\vee)_{Q'}^{P'}$ is the set of coweights. We simply write $\hat{\Delta}_{P'}^\vee$ for $(\hat{\Delta}^\vee)_{P'}^{G'}$ and $\hat{\Delta}_{P'}$ for $\hat{\Delta}_{P'}^{G'}$, respectively.

Let $\tau_{Q'}^{P'}$ be the characteristic function of the subset

$$\{H \in \mathfrak{a}'_0 : \langle \alpha, H \rangle > 0 \text{ for all } \alpha \in \Delta_{Q'}^{P'}\} \subset \mathfrak{a}'_0$$

and let $\hat{\tau}_{Q'}^{P'}$ be the characteristic function of the subset

$$\{H \in \mathfrak{a}'_0 : \langle \varpi, H \rangle > 0 \text{ for all } \varpi \in \hat{\Delta}_{Q'}^{P'}\} \subset \mathfrak{a}'_0.$$

Since these two functions depend only on the projection of \mathfrak{a}'_0 to $\mathfrak{a}_{Q'}^{P'}$, we also regard them as functions on $\mathfrak{a}_{Q'}^{P'}$. For simplicity, write $\mathfrak{a}_{P'}$ for $\mathfrak{a}_{P'}^{G'}$ and $\tau_{P'}$, $\hat{\tau}_{P'}$ for $\tau_{P'}^{G'}$, $\hat{\tau}_{P'}^{G'}$, respectively.

For each parabolic subgroup $P' = M'U'$ of G' , we have a height map

$$H_{P'} : G'(\mathbb{A}_F) \rightarrow \mathfrak{a}_{P'}$$

characterized by the following properties (see [1, page 917]):

- $|\chi|(m) = e^{\langle \chi, H_{P'}(m) \rangle}$ for all $\chi \in X(M')$ and $m \in M'(\mathbb{A}_F)$;

- $H_{P'}(umk) = H_{P'}(m)$ for all $u \in U'(\mathbb{A}_F), m \in M'(\mathbb{A}_F), k \in K'$.

The restriction of $H_{P'}$ on $M'(\mathbb{A}_F)$ is a surjective homomorphism. For a pair of parabolic subgroups $Q' \subseteq P'$ of G' , it is easy to check that the projection of $H_{Q'}(g) \in \mathfrak{a}_{Q'}$ to $\mathfrak{a}_{P'}$ is $H_{P'}(g)$ for all $g \in G'(\mathbb{A}_F)$.

Denote the kernel of $H_{P'}|_{M'(\mathbb{A}_F)}$ by $M'(\mathbb{A}_F)^1$ and the connected component of 1 in $A_{M'}(\mathbb{R})$ by $A_{M'}(\mathbb{R})^0$. Write A'_0 for $A_{M'_0}(\mathbb{R})^0$. Then $M'(\mathbb{A}_F)$ is the direct product of the normal subgroup $M'(\mathbb{A}_F)^1$ with $A_{M'}(\mathbb{R})^0$, and $H_{P'}$ gives an isomorphism between $A_{M'}(\mathbb{R})^0$ and $\mathfrak{a}_{P'}$. Denote the inverse of this map by $X \rightarrow e^X$. We fix a Haar measure on $\mathfrak{a}_{P'}$. Write ρ'_0 for the half sum of positive roots in $(\mathfrak{a}'_0)^*$ and denote by $\rho_{P'}, \rho_{Q'}$ the projection of ρ'_0 to $(\mathfrak{a}_{P'})^*, (\mathfrak{a}_{Q'})^*$, respectively. Recall that $e^{2\langle \rho_{P'}, H_{P'}(p) \rangle} = \delta_{P'}(p)$ for $p \in P'(\mathbb{A}_F)$. We use the similar notation for G by removing the prime '.

The following is a generalization of [41, Lemma I.3.2] and can be proved in the same way.

Proposition 3.2. *An automorphic form $\phi \in \mathcal{A}_{P'}(N \rtimes \mathbf{G}')$ admits a finite decomposition*

$$(3.1) \quad \phi((n, ue^X mk)) = \sum_i Q_i(X) \phi_i((n, mk)) e^{\langle \lambda_i + \rho_{P'}, X \rangle}$$

for $u \in U'(\mathbb{A}_F), X \in \mathfrak{a}_{P'}, m \in M'(\mathbb{A}_F)^1$ and $k \in K'$, where $\lambda_i \in \mathfrak{a}_{P'}^* \otimes_{\mathbb{R}} \mathbb{C}$, $Q_i \in \mathbb{C}[\mathfrak{a}_{P'}]$ and $\phi_i \in \mathcal{A}_{P'}(N \rtimes \mathbf{G}')$ satisfies $\phi_i(n, e^X g) = \phi_i(n, g)$ for $X \in \mathfrak{a}_{P'}$ and $g \in \mathbf{G}'$.

We denote the finite set of exponents λ_i appearing in the above decomposition of ϕ by $\mathcal{E}_{P'}(\phi)$. When $N = \mathbf{1}$, the above proposition is nothing but [41, Lemma I.3.2].

For $\varphi \in \mathcal{A}(N \rtimes \mathbf{G}')$, define $\varphi_{P'} : N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F) \rightarrow \mathbb{C}$ by

$$\varphi_{P'}(n, g) := \int_{[U']} \varphi(n, ug) du \quad \text{for } (n, g) \in N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F).$$

Then it is obvious that $\varphi_{P'} \in \mathcal{A}_{P'}(N \rtimes \mathbf{G}')$. If we regard $\varphi \in \mathcal{A}(\mathbf{1} \rtimes \mathbf{G}')$ as $\varphi \in \mathcal{A}(\mathbf{G}')$, then $\varphi_{P'}$ is the usual constant term of $\varphi \in \mathcal{A}(\mathbf{G}')$ along P' defined by

$$\varphi_{P'}(g) = \int_{[U']} \varphi(ug) du.$$

As we shall see in the next section, we are mainly interested in the situation $N \rtimes \mathbf{G}' \subseteq G$. In this case, we will consider the restriction of the constant terms of functions in $\mathcal{A}(\mathbf{G}) \otimes \mathcal{A}(N \rtimes \mathbf{G}')$ to $N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F)$. Therefore, we introduce the following notation to denote it.

For $\phi \in \mathcal{A}(\mathbf{G})$, $\phi' \in \mathcal{A}(N \rtimes \mathbf{G}')$ and a parabolic subgroup $\tilde{S} = S_1 \times S'_2$ of $\mathbf{G} \times \mathbf{G}'$, a function $(\phi \otimes \phi')_{\tilde{S}} : N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F) \rightarrow \mathbb{C}$ is defined by

$$(\phi \otimes \phi')_{\tilde{S}}(n, g) = \phi_{S_1}(ng) \phi'_{S'_2}(n, g).$$

4. The GGP triplets

From now on, we focus on special triplets of algebraic groups (G, G', N) which appear in the setting of the GGP conjecture.

Let E be either F or a quadratic extension of F with adèle ring \mathbb{A}_E . For a place v of F , denote by E_v the localization of E at v . Fix a non-trivial additive character ψ of $F \backslash \mathbb{A}_F$ and the quadratic character ω of \mathbb{A}_F^\times associated to E/F by the global class field theory. (When $E = F$, ω is just the trivial character.) Fix a character μ of $E^\times \backslash \mathbb{A}_E^\times$ such that $\mu|_{\mathbb{A}_F^\times} = \omega$. Write $|\cdot|_{E_v}$ for the normalized absolute valuation on E_v and for $x \in \mathbb{A}_E^\times$, put $|x|_E = \prod_v |x|_{E_v}$. When $E = F$, we simply write $|\cdot|_v$ for $|\cdot|_{F_v}$ and $|\cdot|$ for $|\cdot|_F$.

Occasionally, we regard ψ as the character of $E \setminus \mathbb{A}_E$ by composition with the trace map $\text{Tr}_{E/F}$. We also view μ and $|\cdot|_E$ as the characters of $GL_n(\mathbb{A}_E)$ by composition with the determinant map. Sometimes, we denote $\mu \circ \det_{GL_n}$ by $\mu(n)$.

Fix $\epsilon \in \{\pm\}$. Let $(W_n, (\cdot, \cdot)_n)$ be a non-degenerate n -dimensional ϵ -Hermitian space over E and W_m its a m -dimensional subspace. Let W_m^\perp be the orthogonal complement of W_m in W_n and $(\cdot, \cdot)_m$ be the restriction of $(\cdot, \cdot)_n$ to W_m . We further assume that $(\cdot, \cdot)_m$ is also non-degenerate and

- $\epsilon \cdot (-1)^{\dim W_m^\perp} = -1$;
- W_m^\perp in W_n is a split space.

When $n - m = 2r$, the second condition means that there exist r -dimensional isotropic subspaces X, X^* of W_m^\perp such that $W_m^\perp = X \oplus X^*$. When $n - m = 2r + 1$, it means that there exist r -dimensional isotropic subspaces X, X^* and a non-isotropic hermitian line E in W_m^\perp such that E is orthogonal to $X \oplus X^*$.

Let k, k' be Witt indices of W_n, W_m , respectively. When $n - m = 2r$ (resp. $n - m = 2r + 1$), it is easy to check $k = k' + r$ (resp. $k' + r \leq k \leq k' + r + 1$). Let G_n be the isometry group of $(W_n, (\cdot, \cdot)_n)$ and G_m that of $(W_m, (\cdot, \cdot)_m)$. We identify G_m as the subgroup of G_n which acts trivially on W_m^\perp . We fix a norm $\|\cdot\|_{G_n}$ on $G_n(\mathbb{A}_F)$ and define a norm $\|\cdot\|_{G_m}$ on $G_m(\mathbb{A}_F)$ by

$$\|g'\|_{G_m} := \|g'\|_{G_n}, \quad g' \in G_m(\mathbb{A}_F).$$

Until section 6, we fix n, m and use the notation G and G' to denote G_n and G_m , respectively.

We fix maximal totally isotropic subspaces $X'_{k'}, Y'_{k'}$ of W_m , in duality, with respect to $(\cdot, \cdot)_m$. Fix also their complete flags

$$\begin{aligned} 0 &= X'_0 \subset X'_1 \subset \cdots \subset X'_{k'}, \\ 0 &= Y'_0 \subset Y'_1 \subset \cdots \subset Y'_{k'}, \end{aligned}$$

in order that for all $1 \leq i \leq k'$, X'_i and Y'_i are in dual with respect to $(\cdot, \cdot)_m$. Let $P'_0 = M'_0 U'_0$ be the parabolic subgroup of G' which stabilizes the above complete flag of X'_k with the Levi subgroup M'_0 , which stabilizes the complete flag of $Y'_{k'}$. Let $A_{M'_0}$ be the maximal split torus in M'_0 and call any F -parabolic subgroup P' of G' which contains P'_0 *standard*. We also fix a maximal compact subgroup $K' = \prod_v K'_v$ of $G'(\mathbb{A}_F)$ which satisfies $G'(\mathbb{A}_F) = P'_0(\mathbb{A}_F)K'$ and for every standard parabolic subgroup $P' = M'U'$ of G' ,

$$P'(\mathbb{A}_F) \cap K' = (M'(\mathbb{A}_F) \cap K')(U'(\mathbb{A}_F) \cap K')$$

and $M'(\mathbb{A}_F) \cap K'$ is still maximal compact in $M'(\mathbb{A}_F)$ (see [41, I.1.4]).

Let $P(\overline{X'_{k'}})$ be the parabolic subgroup of G which stabilizes the same complete flag of $X'_{k'}$. Fix a complete flag \overline{X} of X and let $N_{n,r}$ be the unipotent radical of the parabolic subgroup of G which stabilizes \overline{X} . Let $P_0 = M_0 U_0$ be a minimal F -parabolic subgroup of G contained in $P(\overline{X'_{k'}})$ and stabilizes \overline{X} . Note that $N_{n,r} \subseteq U_0$. We also fix a maximal compact subgroup K of $G(\mathbb{A}_F)$ containing K' as we choose K' in $G'(\mathbb{A}_F)$. Note that the map $P \rightarrow P \cap G'$ gives a bijection between the set of F -parabolic subgroups of G which contains $P(\overline{X'_{k'}})$ and the set of standard parabolic subgroups of G' . Throughout the rest of the paper, parabolic subgroup of G (resp. parabolic subgroup of G') always means a standard parabolic subgroup of G containing $P(\overline{X'_{k'}})$ (resp. a standard parabolic subgroup of G'). When P is a parabolic subgroup of G , we use the letter P' to denote $P \cap G'$ and vice versa. In particular, $(P(\overline{X'_{k'}}))' = P'_0$.

Since $\mathfrak{a}_{0'}$ (resp. \mathfrak{a}_0) is isomorphic to $\mathbb{R}^{k'}$ (resp. \mathbb{R}^k), we regard functions on $\mathfrak{a}_{0'}$ as those of \mathfrak{a}_0 by composing them with the projection. Since \mathfrak{a}_P and $\mathfrak{a}_{P'}$ are isomorphic, we identify them and fix Euclidean norms $\|\cdot\|_P$ on \mathfrak{a}_P and $\|\cdot\|_{P'}$ on $\mathfrak{a}_{P'}$ so that they are same by the identification of \mathfrak{a}_P and $\mathfrak{a}_{P'}$.

Let I be either G or G' . Let $I(\mathbb{A}_F)^1 = \cap_{\chi \in X(H)} \ker(|\chi|)$ and $I(F \otimes_{\mathbb{Q}} \mathbb{R})^1 = I(\mathbb{A}_F)^1 \cap I(F \otimes_{\mathbb{Q}} \mathbb{R})$, where $X(I)$ is the additive group of homomorphisms from I to $GL(1)$ defined over F . Then $I(F) \subset I(\mathbb{A}_F)^1$. When $E = F$ and $\epsilon = -1$, I is the symplectic group. Then it has a non-trivial double cover, which is called the metaplectic group. We define the topological group \mathbf{I} and the surjective morphism $pr : \mathbf{I} \rightarrow I$ according to (E, ϵ) as follows:

$$\mathbf{I} = \begin{cases} \text{a metaplectic double cover of } I, & E = F, \epsilon = -1; \\ I, & \text{otherwise,} \end{cases}$$

$$pr = \begin{cases} \text{a double covering map whose kernel is } \{\pm 1\}, & E = F, \epsilon = -1; \\ \text{identity map,} & \text{otherwise.} \end{cases}$$

The letters P, Q (resp. P', Q') are reserved for parabolic subgroups of G (resp. G') and M, L (resp. M', L') for their Levi subgroups and U, V (resp. U', V') for their unipotent radicals. We fix the minimal F -parabolic subgroup $P_0 \times P'_0$ of $G \times G'$ and call $P \times Q'$ a parabolic subgroup of $G \times G'$ if P and Q' are parabolic subgroup of G and G' respectively. To distinguish the notation of parabolic subgroups of $G \times G'$ with those of G , we use the letter \tilde{S} for parabolic subgroups of $G \times G'$. Let θ be the involution on the set of parabolic subgroups of $G \times G'$ sending $P \times Q'$ to $Q \times P'$. Then the map $P \rightarrow P \times P'$ gives a bijection between parabolic subgroups of G and θ -invariant parabolic subgroups of $G \times G'$. We use the letter \bar{P} to denote $P \times P'$. Let N be a unipotent F -subgroup of G which admits a G' -action $\sigma : G' \rightarrow \text{Aut}(N)$ such that $N \rtimes G'$ is a F -subgroup of G .

Now we define a function $(\phi \otimes \phi')_{P,N} : \mathbf{G}'(\mathbb{A}_F) \rightarrow \mathbb{C}$ as

$$(\phi \otimes \phi')_{P,N}(g) := \int_{[N]} (\phi \otimes \phi')_{\bar{P}}(n, g) dn.$$

Remark 4.1. When N is the trivial group $\mathbf{1}$, for every $\phi \in \mathcal{A}(\mathbf{G})$ and $\phi' \in \mathcal{A}(\mathbf{1} \rtimes \mathbf{G}')$, $(\phi \otimes \phi')_{P,\mathbf{1}} = \phi_P \cdot \phi'_{P'}$.

Proposition 4.2. For $\phi \in \mathcal{A}(\mathbf{G}), \phi' \in \mathcal{A}(N \rtimes \mathbf{G}')$, $(\phi \otimes \phi')_{P,N}$ is a $U'(\mathbb{A}_F)P'(F)$ -invariant smooth function on \mathbf{G}' for any parabolic subgroup P of G .

Proof. The smoothness readily follows from those of ϕ and ϕ' . To prove the left $U'(\mathbb{A}_F)P'(F)$ -invariance, choose an arbitrary $v \in U'(\mathbb{A}_F)$ and $p \in P'(F)$. Then $(\phi \otimes \phi')_{P,N}(apg)$ equals

$$\int_{[N]} \int_{[U']} \int_{[U]} \varphi(nu_1vpg) \varphi'(n, u_2vpg) du_1 du_2 dn = \int_{[N]} \int_{[U']} \int_{[U]} \varphi(nu_1pg) \varphi'(n, u_2pg) du_1 du_2 dn.$$

(We used the change of variables $u_1 \rightarrow u_1v^{-1}, u_2 \rightarrow u_2v^{-1}$.) For each $i = 1, 2$, write $u'_i = p^{-1}u_i p$. Since $p \in G'(F) \subseteq G'(\mathbb{A}_F)^1 \subseteq G(\mathbb{A}_F)^1$, we have $du'_i = du_i$ because $\delta_{P'}(p^{-1}) = \delta_P(p^{-1}) = 1$. We also write $n' = \sigma(p^{-1})n$. Then $dn' = \delta_{\sigma}(p^{-1})dn$, where

$$\delta_{\sigma}(p^{-1}) = |\det(d\sigma(p^{-1}); \text{Lie}(N))|$$

with $d\sigma(p^{-1}) : \text{Lie}(N) \rightarrow \text{Lie}(N)$ being the differential of $\sigma(p^{-1})$. Since $p \in G'(F) \subset G'(\mathbb{A}_F)^1$, we have $\delta_{\sigma}(p^{-1}) = 1$. Thus

$$\int_{[N]} \int_{[U']} \int_{[U]} \varphi(nu_1pg) \varphi'(n, u_2pg) du_1 du_2 dn$$

equals

$$\int_{[N]} \int_{[U']} \int_{[U]} \varphi(pn'u'_1g) \varphi'((1, p) \cdot (n', u'_2g)) du'_1 du'_2 dn'$$

and since φ is $N(F) \times \mathbf{G}'(F)$ -invariant, the last integral equals

$$\int_{[N]} \int_{[U']} \int_{[U]} \varphi(n' u'_1 g) \varphi(n', u'_2 g) du'_1 du'_2 dn' = (\phi \otimes \phi')_{P,N}(g).$$

This shows that $(\phi \otimes \phi')_{P,N}$ is a left $U'(\mathbb{A}_F)P'(F)$ -invariant function. \square

5. Mixed truncation

In this section, we introduce the mixed truncation which is an essential ingredient to define regularized periods.

Let $T' \in \mathfrak{a}'_0$. For $\phi_1 \in \mathcal{A}(\mathbf{G})$, $\phi_2 \in \mathcal{A}(N \times \mathbf{G}')$, we define their mixed truncation by

$$\Lambda_m^{T'}(\phi_1 \otimes \phi_2)(g) = \sum_{P'} (-1)^{\dim \mathfrak{a}_{P'}} \sum_{\gamma \in P'(F) \backslash \mathbf{G}'(F)} (\phi_1 \otimes \phi_2)_{P,N}(\gamma g) \hat{\tau}_{P'}(H_{P'}(\gamma g) - T'), \quad g \in \mathbf{G}'(\mathbb{A}_F).$$

More generally, we define a partial mixed truncation by

$$\Lambda_m^{T',P'}(\phi_1 \otimes \phi_2)(g) = \sum_{Q' \subset P'} (-1)^{\dim \mathfrak{a}_{Q'}^{P'}} \sum_{\delta \in Q'(F) \backslash P'(F)} (\phi_1 \otimes \phi_2)_{Q,N}(\delta g) \hat{\tau}_{Q'}^{P'}(H_{Q'}(\delta g) - T').$$

Then $\Lambda_m^{T',P'}(\phi_1 \otimes \phi_2)$ is a function on $P'(F) \backslash \mathbf{G}'(\mathbb{A}_F)$.

For any two parabolic groups $Q' \subseteq P'$, Langlands' combinatorial lemma asserts that

$$(5.1) \quad \sum_{Q' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{Q'}^{R'}} \hat{\tau}_{Q'}^{R'} \tau_{R'}^{P'} = \begin{cases} 1 & \text{if } P' = Q' \\ 0 & \text{otherwise.} \end{cases}$$

We put

$$\Gamma_{Q'}^{P'}(H', X) = \sum_{Q' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \cdot \tau_{Q'}^{R'}(H') \hat{\tau}_{R'}^{P'}(H' - X), \quad H', X \in \mathfrak{a}'_0.$$

Note that this function depends only on the projection of H' and X onto $\mathfrak{a}_{Q'}^{P'}$ and is compactly supported on $\mathfrak{a}_{Q'}^{P'}$.

The following lemma is a consequence of Langlands' combinatorial lemma.

Lemma 5.1. *For $H', X \in \mathfrak{a}'_0$,*

$$(5.2) \quad \hat{\tau}_{Q'}^{P'}(H' - X) = \sum_{Q' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \cdot \hat{\tau}_{Q'}^{R'}(H') \Gamma_{R'}^{P'}(H', X),$$

$$(5.3) \quad \tau_{Q'}^{P'}(H' - X) = \sum_{Q' \subseteq R' \subseteq P'} \Gamma_{Q'}^{R'}(H' - X, -X) \cdot \tau_{R'}^{P'}(H').$$

Proof. Note that

$$\begin{aligned} \sum_{Q' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \cdot \hat{\tau}_{Q'}^{R'}(H') \Gamma_{R'}^{P'}(H', X) &= \sum_{Q' \subseteq R' \subseteq P'} \sum_{R' \subseteq S' \subseteq P'} (-1)^{\dim \mathfrak{a}_{R'}^{S'}} \cdot \hat{\tau}_{Q'}^{R'}(H') \tau_{R'}^{S'}(H') \hat{\tau}_{S'}^{P'}(H' - X) \\ &= \sum_{Q' \subseteq S' \subseteq P'} (-1)^{\dim \mathfrak{a}_{Q'}^{S'}} \cdot \hat{\tau}_{S'}^{P'}(H' - X) \cdot \left(\sum_{Q' \subseteq R' \subseteq S'} (-1)^{\dim \mathfrak{a}_{Q'}^{R'}} \cdot \hat{\tau}_{Q'}^{R'}(H') \tau_{R'}^{S'}(H') \right). \end{aligned}$$

By (5.1), it equals to $\hat{\tau}_{Q'}^{P'}(H' - X)$. This proves (5.2).

On the other hand,

$$\begin{aligned}
 & \sum_{Q' \subseteq R' \subseteq P'} \Gamma_{Q'}^{R'}(H' - X, -X) \cdot \tau_{R'}^{P'}(H') \\
 &= \sum_{Q' \subseteq R' \subseteq P'} \sum_{Q' \subseteq S' \subseteq R'} (-1)^{\dim \mathfrak{a}_{S'}^{R'}} \cdot \tau_{Q'}^{S'}(H' - X) \cdot \hat{\tau}_{S'}^{R'}(H') \tau_{R'}^{P'}(H') \\
 &= \sum_{Q' \subseteq S' \subseteq P'} \tau_{Q'}^{S'}(H' - X) \sum_{S' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{S'}^{R'}} \cdot \hat{\tau}_{S'}^{R'}(H') \tau_{R'}^{P'}(H').
 \end{aligned}$$

By (5.1), it equals to $\tau_{Q'}^{P'}(H' - X)$. This proves (5.3). □

The mixed truncation has the following property.

Lemma 5.2. *Let $\phi \in \mathcal{A}(\mathbf{G})$, $\phi' \in \mathcal{A}(N \rtimes \mathbf{G}')$. Then for all $g \in \mathbf{G}'(\mathbb{A}_F)$,*

$$\Lambda_m^{T'+S',P'}(\phi \otimes \phi')(g) = \sum_{Q' \subseteq P'} \sum_{\delta \in Q'(F) \backslash P'(F)} \Lambda_m^{T',Q'}(\phi \otimes \phi')(\delta g) \cdot \Gamma_{Q'}^{P'}(H_Q(\delta g) - T', S').$$

Proof. Using (5.2), $\Lambda_m^{T'+S',P'}(\phi \otimes \phi')(g)$ equals

$$\begin{aligned}
 & \sum_{Q' \subseteq P'} (-1)^{\dim \mathfrak{a}_{Q'}^{P'}} \cdot \sum_{\delta \in Q'(F) \backslash P'(F)} (\phi_1 \otimes \phi_2)_{Q,N}(\delta g) \\
 & \times \left(\sum_{Q' \subseteq R' \subseteq P'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \cdot \hat{\tau}_{Q'}^{R'}(H_{Q'}(\delta g) - T') \Gamma_{R'}^{P'}(H_{R'}(\delta g) - T', S') \right).
 \end{aligned}$$

Write $\delta = \delta_1 \delta_2$, where $\delta_1 \in Q'(F) \backslash R'(F)$, $\delta_2 \in R'(F) \backslash P'(F)$. Then $H_{R'}(\delta g) = H_{R'}(\delta_2 g)$ and thus

$$\begin{aligned}
 & \Lambda_m^{T'+S',P'}(\phi \otimes \phi')(g) \\
 &= \sum_{R' \subseteq P'} \sum_{\delta_2 \in R'(F) \backslash P'(F)} \left(\sum_{Q' \subseteq R'} \sum_{\delta_1 \in Q'(F) \backslash R'(F)} (-1)^{\dim \mathfrak{a}_{Q'}^{R'}} \cdot (\phi_1 \otimes \phi_2)_{Q,N}(\delta_1 \delta_2 g) \cdot \hat{\tau}_{Q'}^{R'}(H_{Q'}(\delta_1 \delta_2 g) - T') \right) \\
 & \times \Gamma_{R'}^{P'}(H_{R'}(\delta_2 g) - T', S') \\
 &= \sum_{R' \subseteq P'} \sum_{\delta_2 \in R'(F) \backslash P'(F)} \Lambda_m^{T',R'}(\phi \otimes \phi')(\delta_2 g) \cdot \Gamma_{R'}^{P'}(H_{R'}(\delta_2 g) - T', S').
 \end{aligned}$$

This completes the proof. □

For any two parabolic subgroups $Q' \subseteq P'$, put

$$\sigma_{Q'}^{P'} = \sum_{Q' \subseteq R'} (-1)^{\dim \mathfrak{a}_{Q'}^{R'}} \cdot \tau_{P'}^{R'} \hat{\tau}_{R'}.$$

Using the binomial theorem, it is easy to check that for any parabolic subgroup $P'_1 \supseteq P'$,

$$(5.4) \quad \tau_{P'_1}^{P'_1} \hat{\tau}_{P'_1} = \sum_{P'_1 \subseteq R'} \sigma_{P'_1}^{R'}.$$

For a parabolic subgroup $P = UM$ (resp. $P' = U'M'$) of G (resp. G'), write $\mathbf{M}^1 = pr^{-1}(M^1)$ (resp. $(\mathbf{M}')^1 = pr^{-1}((M')^1)$). Let w be a compact subset of $U_0(\mathbb{A}_F)\mathbf{M}_0(\mathbb{A}_F)^1$ and $t_0 \in \mathfrak{a}_0$. Note that $\mathbf{M}_0(\mathbb{A}_F)$ is the direct product of the normal subgroup $\mathbf{M}_0(\mathbb{A}_F)^1$ with A_0 . Put

$$A_0(t_0) = \{a \in A_0 \mid \langle \beta, H_{P_0}(a) - t_0 \rangle > 0 \text{ for all } \beta \in \Delta_0^P\}.$$

For any parabolic subgroup P of G and $T \in \mathfrak{a}_0$, define a Siegel set and a truncated Siegel set relative to P by

$$\begin{aligned} \mathcal{S}^P &= \mathcal{S}^P(t_0) = \{x = pak \mid p \in w, a \in A_0(t_0), k \in \mathbf{K}\}, \\ \mathcal{S}^P(T) &= \mathcal{S}^P(t_0, T) = \{x \in \mathcal{S}^P(t_0) \mid \langle \alpha, H_{P_0}(x) - T \rangle \leq 0 \text{ for all } \alpha \in \hat{\Delta}_0^P\}, \end{aligned}$$

respectively. When $P = G$, we simply write $\mathcal{S} = \mathcal{S}^G$ and $\mathcal{S}(T) = \mathcal{S}^G(T)$. Let $F^P(g, T)$ be the characteristic function on $P(F)\mathcal{S}^P(T)$ in $\mathbf{G}(\mathbb{A}_F)$. We take $w', \mathcal{S}^{P'}, \mathcal{S}^{P'}(T')$ and $F^{P'}(g, T')$ similarly. We choose w, w', t_0, t'_0 so as to $\mathbf{G}(\mathbb{A}_F) = P(F)\mathcal{S}^P$ and $\mathbf{G}'(\mathbb{A}_F) = P'(F)\mathcal{S}^{P'}$.

The following lemma, which is a generalization of [30], Lemma 2.1], is crucial to prove the rapidly decreasing property of $\Lambda_m^{T', P'}$.

Lemma 5.3. *For any w', t'_0 , we can choose w and t_0 so that $\mathcal{S}^{P'} \subset \mathcal{S}^P$.*

Proof. When $k = k' + r$, the lemma is obvious because $P'_0 = P_0 \cap G'$. Suppose $k > k' + r$ and we fix w', t'_0 . Since $A'_0 \subseteq A_0$, we can choose $t_0 \in \mathfrak{a}_0$ such that $A'_0(t'_0) \subseteq A_0(t_0)$. We shall find a compact subset $w \subseteq U_0(\mathbb{A}_F)\mathbf{M}_0(\mathbb{A}_F)^1$ such that $w'A'_0(t'_0)\mathbf{K}' \subseteq wA_0(t_0)\mathbf{K}$.

Choose arbitrary $g \in w'A'_0(t'_0)\mathbf{K}'$ and decompose it $g = u'm'a'k'$, where $u' \in U'_0, m' \in \mathbf{M}'_0(\mathbb{A}_F)^1, a' \in A'_0(t'_0), k' \in \mathbf{K}'$. Note that $U'_0 \subseteq U_0, A'_0(t'_0) \subseteq A_0(t_0)$ and $\mathbf{K}' \subseteq \mathbf{K}$, though P'_0 is not contained in P_0 . Let $W_{m, k'}$ be the orthogonal complement of $X'_{k'} + Y'_{k'}$ in W_m and $G'_{k'}$ its isometry group. Since $(M'_0)^1 = (A_{M'_0})^1 \times G'_{k'}$ and $A_{M'_0} \subseteq A_{M_0}$, we may assume $m' \in \mathbf{G}'_{k'}(\mathbb{A}_F)$. Let $W_{n, k'}$ be the orthogonal complement of $X'_{k'} + Y'_{k'}$ in W_n and put $G_{k'}$ its isometry group. Since $k > k' + r$, $G_{k'} \cap P_0$ is a proper parabolic subgroup of $G_{k'}$. Since $G'_{k'} \subseteq G_{k'}$, using the Iwasawa decomposition of $G_{k'}$ with respect to $G_{k'} \cap P_0$, we can decompose m' in $\mathbf{G}_{k'}$ as $u_0 m_0 k_0$, where $u_0 \in \mathbf{G}_{k'} \cap U_0, m_0 \in \mathbf{G}_{k'} \cap \mathbf{M}_0, k_0 \in \mathbf{G}_{k'} \cap \mathbf{K}$. Since $\mathbf{G}_{k'}$ and A'_0 are commutative, $u'm'a'k' = u'u_0 m_0 k_0 a' k' = u'u_0 m_0 a' k_0 k$. This shows that we can find w as desired. \square

Now, we prove the rapidly decreasing property of $\Lambda_m^{T', P'}$.

Lemma 5.4. *Let $\phi \in \mathcal{A}(\mathbf{G}), \phi' \in \mathcal{A}(N \rtimes \mathbf{G}')$. Then for every positive integer N_1 and $k \in \mathbf{K}'$,*

$$\sup_{m \in \mathcal{S}^{P'} \cap \mathbf{M}'(\mathbb{A}_F)^1} |\Lambda_m^{T', P'}(\phi \otimes \phi')(mk)| \cdot \|m\|_{G'}^{N_1} < \infty.$$

Proof. It is enough to show when $P' = G'$ and $k = 1$. Fix $T' \in \mathfrak{a}'_0$ and for each parabolic subgroup P' of G' , choose $\mathcal{S}^{P'}$ so that $\mathbf{G}'(\mathbb{A}_F) = P'(F)\mathcal{S}^{P'}$. [1, Lemma 6.4] says that

$$(5.5) \quad \sum_{P'_1 \subseteq P'} \sum_{\delta_1 \in P'_1(F) \backslash P'(F)} F^{P'_1}(\delta_1 x, T') \tau_{P'_1}^{P'}(H_{P'_1}(\delta_1 x) - T') = 1.$$

By substituting (5.5) into the definition of $\Lambda_m^{T'}(\phi_1 \otimes \phi_2)(g)$, we have

$$\begin{aligned} &\Lambda_m^{T'}(\phi_1 \otimes \phi_2)(g) \\ &= \sum_{P'} (-1)^{\dim \mathfrak{a}_{P'}} \sum_{\delta \in P'(F) \backslash G'(F)} (\phi_1 \otimes \phi_2)_{P, N}(\delta g) \hat{\tau}_{P'}(H_{P'}(\delta g) - T') \end{aligned}$$

$$\begin{aligned}
 &= \sum_{P'_1 \subseteq P'} (-1)^{\dim \mathfrak{a}_{P'}} \sum_{\delta \in P'(F) \backslash G'(F)} \left(\sum_{\delta_1 \in P'_1(F) \backslash P'(F)} F^{P'_1}(\delta_1 \delta g, T') \tau_{P'_1}^{P'}(H_{P'_1}(\delta_1 \delta g) - T') \right) \\
 &\times (\phi_1 \otimes \phi_2)_{P,N}(\delta g) \hat{\tau}_{P'}(H_{P'}(\delta g) - T') \\
 &= \sum_{P'_1 \subseteq P'} (-1)^{\dim \mathfrak{a}_{P'}} \sum_{\delta \in P'_1(F) \backslash G'(F)} F^{P'_1}(\delta g, T') (\phi_1 \otimes \phi_2)_{P,N}(\delta g) \tau_{P'_1}^{P'}(H_{P'_1}(\delta g) - T') \hat{\tau}_{P'}(H_{P'}(\delta g) - T').
 \end{aligned}$$

(In the last equality, we used the fact $(\phi_1 \otimes \phi_2)_{P,N}(\delta g) \hat{\tau}_{P'}(H_{P'}(\delta g) - T') = (\phi_1 \otimes \phi_2)_{P,N}(\delta_1 \delta g) \hat{\tau}_{P'}(H_{P'}(\delta_1 \delta g) - T')$.)

Using (5.4), we have

$$\begin{aligned}
 \Lambda_m^{T'}(\phi_1 \otimes \phi_2)(g) &= \sum_{P'_1 \subseteq P' \subseteq Q'} (-1)^{\dim \mathfrak{a}_{P'}} \sum_{\delta \in P'_1(F) \backslash G'(F)} F^{P'_1}(\delta g, T') (\phi_1 \otimes \phi_2)_{P,N}(\delta g) \sigma_{P'_1}^{Q'}(H_{P'_1}(\delta g) - T') \\
 &= \sum_{P'_1 \subseteq Q'} \sum_{\delta \in P'_1(F) \backslash G'(F)} F^{P'_1}(\delta g, T') \sigma_{P'_1}^{Q'}(H_{P'_1}(\delta g) - T') \left(\int_{[N]} \phi_{P_1, Q}(n, \delta g) dn \right)
 \end{aligned}$$

where

$$\phi_{P_1, Q}(n, g) = \sum_{P_1 \subseteq R \subseteq Q} (-1)^{\dim \mathfrak{a}_R} \cdot (\phi \otimes \phi')_{\bar{R}}(n, g).$$

By applying the similar argument in the proof of [43, Lemma 8.2.1(1)], one can show that

$$(-1)^{\dim \mathfrak{a}_{P_1}} \phi_{P_1, Q}(n, g) = \sum_{\bar{P}_1 \subseteq \bar{S} \subseteq \bar{Q}} (-1)^{\dim \mathfrak{a}_{\bar{S}}} \phi_{\bar{S}, \bar{Q}}(n, g),$$

where $\tilde{S} = (S_1, S_2)$ runs over all parabolic subgroups of $G \times G'$ between \bar{P}_1 and \bar{Q} such that $S_1 \cap S_2 = P_1$ and

$$\phi_{\bar{S}, \bar{Q}}(n, g) = \sum_{\tilde{S} \subseteq \tilde{R} \subseteq \bar{Q}} (-1)^{\dim \mathfrak{a}_{\tilde{R}}} \cdot (\phi \otimes \phi')_{\tilde{R}}(n, g).$$

Note that

$$|\phi_{\bar{S}, \bar{Q}}(n, g)| = |\phi_{S_1, Q}(ng)| \cdot |\phi'_{S'_2, Q}(n, g)|$$

where

$$\phi_{S_1, Q}(h) = \sum_{S_1 \subseteq R \subseteq Q} (-1)^{\dim \mathfrak{a}_R} \phi_R(h), \quad h \in \mathbf{G}$$

and

$$\phi'_{S'_2, Q'}(n, g) = \sum_{S'_2 \subseteq R' \subseteq Q'} (-1)^{\dim \mathfrak{a}_{R'}} \phi'_{R'}(n, g), \quad (n, g) \in N \rtimes \mathbf{G}'.$$

Choose $\delta \in G'(F)$ and $g \in \mathcal{S}'$ such that $F^{P'_1}(\delta g, T') \sigma_{P'_1}^{Q'}(H_{P'_1}(\delta g) - T') \neq 0$. Note that $\sigma_{P'_1}^{Q'}$ is a characteristic function by [1, Lemma 6.1]. Therefore, by Lemma 5.3, we can take $T \in \mathfrak{a}_0$ positive enough such that the projection of T to $\mathfrak{a}_{P_k'} \simeq \mathfrak{a}'_0$ is T' and $P'(F)\mathcal{S}^{P'}(T') \subseteq P(F)\mathcal{S}^P(T)$ and $F^{P_1}(\delta g, T) \sigma_{P_1}^Q(H_{P_1}(\delta g) - T) \neq 0$. The property (iv) of the heights function in [41, I.2.2] shows that there are positive constants c_1, ϵ such that

$$e^{\|H_{P_1}(a)\|_{P_1}} = e^{\|H_{P'_1}(a)\|_{P'_1}} > c_1 \|a\|^\epsilon \quad \text{for all } a \in A_{P'_1}(\mathbb{R})^0.$$

If we apply the same arguments in [2, page 92-96] to \mathbf{G}' and \mathbf{G} , respectively, there exists a positive constant $c_{P'_1, Q'}$ such that for all $g \in \mathcal{S}'$ and $\delta \in G'(F)$ with $F^{P'_1}(\delta g, T')\sigma_{P'_1}^{Q'}(H_{P'_1}(\delta g) - T') \neq 0$,

$$\|g\|_G^{N_2\epsilon} \cdot \int_{[N]} |\phi_{S_1, Q}(n\delta g)| dn < c_{P'_1, Q'} \quad , \quad \|g\|_{G'}^{N_2\epsilon} \cdot \int_{[N]} |\phi'_{S'_2, Q'}(n, \delta g)| dn < c_{P'_1, Q'}$$

for all positive integer N_2 . Furthermore, there are positive constant $c_{T'}, N_3$ such that

$$\sum_{P'_1 \subseteq Q'} \sum_{\delta \in P'_1(F) \setminus G'(F)} F^{P'_1}(\delta g, T')\sigma_{P'_1}^{Q'}(H_{P'_1}(\delta g) - T') < c_{T'} \|g\|_{G'}^{N_3} \quad , \quad g \in \mathbf{G}'(\mathbb{A}_F)^1.$$

(See [2, page 96-97].)

Thus if we choose N_2 so that $2N_2\epsilon \geq N_1 + N_3$, it follows

$$\sup_{g \in \mathcal{S}' \cap \mathbf{G}'(\mathbb{A}_F)^1} |\Lambda_m^{T'}(\phi \otimes \phi')(g)| \cdot \|g\|_{G'}^{N_1} < \infty.$$

This completes the proof. \square

6. Regularized trilinear periods

In this section, we shall define the regularized trilinear periods using the mixed truncation introduced in the previous section. We keep the same notations and conventions as in Section 2.

Let \mathbf{C} be the kernel of pr and \mathbf{H} be either \mathbf{G} or $N \rtimes \mathbf{G}'$ or \mathbf{G}' . Then for any $f \in \mathcal{A}(\mathbf{H})$, there is a character $\chi_f : \mathbf{C} \rightarrow \{\pm 1\}$ such that

$$f(\xi g) = \chi_f(\xi) f(g) \quad (\xi \in \mathbf{C}, g \in \mathbf{H}).$$

When $(E, \epsilon) = (F, -1)$, such χ_f is either trivial or the non-trivial quadratic character. We call f *genuine* if χ_f is non-trivial. When $(E, \epsilon) \neq (F, -1)$, χ_f is always trivial because \mathbf{C} is the trivial group.

Let $\chi = \{\chi_1, \chi_2, \chi_3\}$ be a character of \mathbf{C}^3 such that $\chi_1(\xi)\chi_2(\xi)\chi_3(\xi) = 1$ for all $\xi \in \mathbf{C}$. Let $\mathcal{A}(\mathbf{G}, \mathbf{G}')_\chi$ denote the space of triplets $(\phi_1, \phi_2, \phi_3) \in \mathcal{A}(\mathbf{G}) \times \mathcal{A}(N \rtimes \mathbf{G}') \times \mathcal{A}(\mathbf{G}')$ which satisfy

$$\phi_i(\xi g) = \chi_i(\xi) \phi_i(g) \quad (\xi \in \mathbf{C}, g \in \mathbf{G}', i = 1, 2, 3).$$

For $(\phi_1, \phi_2, \phi_3) \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_\chi$, we consider the following integral

$$(6.1) \quad \int_{G'(F) \setminus G'(\mathbb{A}_F)^1} \Lambda_m^{T'}(\phi \otimes \phi')(g) \phi''(g) dg.$$

By the assumption of χ , the integrand is well defined on $G'(\mathbb{A}_F)$ and thanks to Lemma 5.4, this integral converges absolutely.

Proposition 6.1. *The integral in (6.1) is a function of the form $\sum_\lambda p_\lambda(T') e^{\langle \lambda, T' \rangle}$, where p_λ is a polynomial in T' and λ can be taken from the set*

$$\bigcup_{P'} \{ \lambda_1 + \lambda_2 + \lambda_3 + \rho_P \mid \lambda_1 \in \mathcal{E}_P(\phi_P), \lambda_2 \in \mathcal{E}_{P'}(\phi_{P'}), \lambda_3 \in \mathcal{E}_{P'}(\phi''_{P'}) \}.$$

Proof. Since $\Gamma_{P'}^{G'}(X - T', S')$ is compactly supported, using Lemma 5.4, we integrate the equation in Lemma 5.2 against $\phi''(g)$ over $G'(F) \setminus G'(\mathbb{A}_F)^1$ to obtain

$$\int_{G'(F) \setminus G'(\mathbb{A}_F)^1} \Lambda_m^{T'+S'}(\phi \otimes \phi')(g) \phi''(g) dg$$

$$\begin{aligned}
 &= \sum_{P'} \int_{G'(F) \backslash G'(\mathbb{A}_F)^1} \left(\sum_{\delta \in P'(F) \backslash G'(F)} \Lambda_m^{T', P'}(\phi \otimes \phi')(\delta g) \phi''(g) \cdot \Gamma_{P'}^{G'}(H_{P'}(\delta g) - T', S') \right) dg \\
 &= \sum_{P'} \int_{P'(F) \backslash G'(\mathbb{A}_F)^1} \Lambda_m^{T', P'}(\phi \otimes \phi')(g) \phi''(g) \cdot \Gamma_{P'}^{G'}(H_{P'}(g) - T', S') dg \\
 &= \sum_{P'} \int_{K'} \int_{M'(F) \backslash M'(\mathbb{A}_F)^1} \int_{\mathfrak{a}_{P'}}^{\#} \Lambda_m^{T', P'}(\phi \otimes \phi')(e^X mk) \phi''_{P'}(e^X mk) \cdot \Gamma_{P'}^{G'}(X - T', S') e^{-2\langle \rho_{P'}, X \rangle} dX dm dk.
 \end{aligned}$$

(In the second equality, we used $\phi''(g) = \phi''(\delta g)$ for all $\delta \in G(F)$. The symbol $\int_{\mathfrak{a}_{P'}}^{\#}$ in the last equality represents the regularized integral of a polynomial exponential function over a cone in a vector space. See [31, Sec 2].)

The inner integral over $\mathfrak{a}_{P'}$ can be expressed in terms of the Fourier transform of $\Gamma_{P'}^{G'}(\cdot, T')$ evaluated at $\lambda_1 + \lambda_2 + \lambda_3 + \rho_P$ for $\lambda_1 \in \mathcal{E}_P(\phi_P)$, $\lambda_2 \in \mathcal{E}_{P'}(\phi_{P'})$, $\lambda_3 \in \mathcal{E}_{P'}(\phi''_{P'})$. Thus by [2, Lemma 2.2], the $\#$ -integral is a polynomial exponential function in S' whose exponents are $\{\lambda_1 + \lambda_2 + \lambda_3 + \rho_P \mid \lambda_1 \in \mathcal{E}_P(\phi_P), \lambda_2 \in \mathcal{E}_{P'}(\phi_{P'}), \lambda_3 \in \mathcal{E}_{P'}(\phi''_{P'})\}_{Q \supset P'}$. \square

Definition 6.2. Let $\mathcal{A}_0(\mathbf{G}, \mathbf{G}')_{\chi}$ be the subspace of triplets in $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}$ such that the polynomial corresponding to the zero exponent of (6.1) is constant. For $(\phi, \phi', \phi'') \in \mathcal{A}_0(\mathbf{G}, \mathbf{G}')_{\chi}$, we define its regularized period $\mathcal{P}(\phi, \phi', \phi'')$ as its value $p_0(T')$. We also write

$$\mathcal{P}(\phi, \phi', \phi'') = \int_{G'(F) \backslash G'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{G, N}(g) \phi''(g) dg.$$

For a parabolic subgroup P' of G' , we set the $\#$ -integral

$$\mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'') = \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', P'}(\phi \otimes \phi')(g) \phi''_{P'}(g) \tau_{P'}(H_{P'}(g) - T') dg$$

as the triple integral

$$\int_{K'} \int_{M'(F) \backslash M'(\mathbb{A}_F)^1} \int_{\mathfrak{a}_{P'}}^{\#} \Lambda_m^{T', P'}(\phi \otimes \phi')(e^X mk) \phi''_{P'}(e^X mk) \tau_{P'}(X - T') e^{-2\langle \rho_{P'}, X \rangle} dX dm dk.$$

Since the functions $m \rightarrow \Lambda_m^{T', P'}(\phi \otimes \phi')(mk)$ are rapidly decreasing, $\mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ exists if and only if

$$\langle \lambda_1 + \lambda_2 + \lambda_3 + \rho_P, \omega^{\vee} \rangle \neq 0 \text{ for all } \omega^{\vee} \in (\hat{\Delta}^{\vee})_{P'}, \lambda_1 \in \mathcal{E}_P(\phi_P), \lambda_2 \in \mathcal{E}_{P'}(\phi_{P'}), \lambda_3 \in \mathcal{E}_{P'}(\phi''_{P'})$$

by [31, Lemma 3].

Let $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$ be the space of all triplets $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}$ such that

$$\langle \lambda_1 + \lambda_2 + \lambda_3 + \rho_P, \omega^{\vee} \rangle \neq 0 \text{ for all } \omega^{\vee} \in (\hat{\Delta}^{\vee})_{P'}, \lambda_1 \in \mathcal{E}_P(\phi_P), \lambda_2 \in \mathcal{E}_{P'}(\phi_{P'}), \lambda_3 \in \mathcal{E}_{P'}(\phi''_{P'})$$

for all proper parabolic subgroups P' of G' .

Proposition 6.3. *Let $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$. Then the following statements hold.*

- (i) $\sum_{P'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ is independent of T' .
- (ii) $\mathcal{P}(\phi, \phi', \phi'') = \sum_{P'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$. It says $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^* \subset \mathcal{A}_0(\mathbf{G}, \mathbf{G}')_{\chi}$.
- (iii) The regularized period \mathcal{P} is a right $N(\mathbb{A}_F) \rtimes G'(\mathbb{A}_F)^1$ -invariant linear functional on $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$.

(iv) If $g \rightarrow (\phi \otimes \phi')_{G,N}(g)\phi''(g)$ is integrable over $G'(F)\backslash G'(\mathbb{A}_F)^1$, then

$$\mathcal{P}(\phi, \phi', \phi'') = \int_{G'(F)\backslash G'(\mathbb{A}_F)^1} (\phi \otimes \phi')_{G,N}(g) \cdot \phi''(g) dg.$$

Proof. The proof of (i) is somewhat similar with [29, Proposition 3.3]. By the assumption, $\sum_{P'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ is well-defined. Choose a regular $S' \in \mathfrak{a}'_0$. Using the Lemma 5.2 and the decomposition of automorphic forms (3.1), we can write $\mathcal{P}_{P'}^{T'+S'}(\phi, \phi', \phi'')$ as

$$\begin{aligned} & \int_{P'(F)\backslash G'(\mathbb{A}_F)^1}^{\#} \sum_{\substack{Q' \\ Q' \subseteq P'}} \sum_{\delta \in Q'(F)\backslash P'(F)} \Lambda_m^{T', Q'}(\phi \otimes \phi')(\delta g) \phi''_{P'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(\delta g) - T', S') \tau_{P'}(H_{P'}(g) - T' - S') dg \\ &= \int_{Q'(F)\backslash G'(\mathbb{A}_F)^1}^{\#} \sum_{\substack{Q' \\ Q' \subseteq P'}} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{P'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(g) - T', S') \tau_{P'}(H_{P'}(g) - T' - S') dg \\ &= \sum_{\substack{Q' \\ Q' \subseteq P'}} \int_{K'} \int_{(Q' \cap M')(F)\backslash M'(\mathbb{A}_F)^1} \int_{\mathfrak{a}_{P'}}^{\#} \left(\Lambda_m^{T', Q'}(\phi \otimes \phi') \cdot \phi''_{P'} \right) (e^X mk) \Gamma_{Q'}^{P'}(H_{Q'}(m) - T', S') \\ & \quad \times \tau_{P'}(X' - T' - S') e^{-2\langle \rho_{P'}, X \rangle} dX dmdk. \end{aligned}$$

Using the Iwasawa decomposition of M' with respect to $Q' \cap M' = U'_{Q'} M'_{Q'}$, write $m = ve^Y lk'$ where

$$v \in (U_{Q'} \cap M')(\mathbb{A}_F), Y \in \mathfrak{a}_{Q'}^{P'}, l \in M'_{Q'}(\mathbb{A}_F)^1, k' \in (K \cap M')(\mathbb{A}_F).$$

Since $U'_{Q'} = U'(U'_{Q'} \cap M)$ and $\phi''_{P'}(e^X mk) = \phi''_{P'}(ue^X mk)$ for all $u \in U'(\mathbb{A}_F)$,

$$\int_{[U'_{Q'} \cap M']} \phi''_{P'}(e^X ve^Y lk'k) dv = \int_{[U']} \int_{[U'_{Q'} \cap M']} \phi''_{P'}(uv'e^{X+Y} lk'k) dv' du = \phi''_{Q'}(e^{X+Y} lk'k).$$

Thus $\mathcal{P}_{P'}^{T'+S'}(\phi, \phi', \phi'')$ equals the sum over $Q' \subseteq P'$ of

$$\begin{aligned} & \int_{K'} \int_{[M'_{Q'}]} \int_{\mathfrak{a}_{Q'}^{P'}}^{\#} \int_{\mathfrak{a}_{P'}}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(e^{X+Y} lk) \phi''_{Q'}(e^{X+Y} lk) \Gamma_{Q'}^{P'}(Y - T', S') \tau_{Q'}(X - T' - S') e^{-2\langle \rho_{Q'}, X+Y \rangle} dX dY dmdk \\ &= \int_{K'} \int_{[M'_{Q'}]} \int_{\mathfrak{a}_{Q'}}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(e^X lk) \phi''_{Q'}(e^X lk) \Gamma_{Q'}^{P'}(X - T', S') \tau_{P'}(X - T' - S') e^{-2\langle \rho_{Q'}, X \rangle} dX dmdk. \end{aligned}$$

(We incorporated the double $\#$ -integral into a single $\#$ -integral over $\mathfrak{a}_{Q'}$ using [31, (14)].)

Thus if we change two sums over P' and Q' in $\sum_{P'} \mathcal{P}_{P'}^{T'+S'}(\phi, \phi', \phi'')$ and use (5.3),

$$\begin{aligned} \sum_{P'} \mathcal{P}_{P'}^{T'+S'}(\phi, \phi', \phi'') &= \sum_{Q'} \int_{K'} \int_{[M'_{Q'}]} \int_{\mathfrak{a}_{Q'}}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(e^X lk) \phi''_{Q'}(e^X lk) \tau_{Q'}(X - T') e^{-2\langle \rho_{Q'}, X \rangle} dX dmdk \\ &= \sum_{Q'} \mathcal{P}_{Q'}^{T'}(\phi, \phi', \phi'') = \sum_{P'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi''). \end{aligned}$$

This proves (i).

Next we prove (ii). Applying the similar argument in the proof of Proposition 6.1 to M' instead of G' , we see that $\mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ is also a polynomial exponential function in T' . Among all $\mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$,

the zero exponent appears only in the term $P' = G'$ by the assumption. Since $\sum_{P'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ is independent of T' , other exponents are cancelled out except for the zero exponents. This proves (ii).

Next we prove (iii). The linearity and the $N(\mathbb{A}_F)$ -invariance is obvious from the definition of \mathcal{P} . Thus we are sufficient to check $G'(\mathbb{A}_F)^1$ -invariance of \mathcal{P} . For any $x \in \mathbf{G}'(\mathbb{A}_F)$ and a function f on $\mathbf{G}(\mathbb{A}_F)$ or $N(\mathbb{A}_F) \rtimes \mathbf{G}'(\mathbb{A}_F)$ or $\mathbf{G}'(\mathbb{A}_F)$, write f_x for the right translation of f by x .

For $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$, it is easy to see that

$$(\phi_x \otimes \phi'_x)_{P,N} = ((\phi \otimes \phi')_{P,N})_x.$$

The same argument just before [31, Theorem 9] shows that for arbitrary $\varphi \in \mathcal{A}_{P'}(\mathbf{G}')$,

$$\mathcal{E}_{P'}(\varphi) = \mathcal{E}_{P'}(\varphi_x).$$

Thus $\mathcal{E}_{P'}((\phi_x \otimes \phi'_x)_{P,N}) = \mathcal{E}_{P'}((\phi \otimes \phi')_{P,N})$ and so $(\phi_x, \phi'_x, \phi''_x) \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$.

By (ii), it is enough to show that for any $x \in G'(\mathbb{A}_F)^1$,

$$(6.2) \quad \mathcal{P}(\phi, \phi', \phi'') = \sum_{P'} \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', P'}(\phi_x \otimes \phi'_x)(g) (\phi_x)''_{P'}(g) \tau_{P'}(H_{P'}(g) - T') dg.$$

Fix $x \in G'(\mathbb{A}_F)^1$ and set $F_{P'}(g) = \Lambda_m^{T', P'}(\phi_{x^{-1}} \otimes \phi'_{x^{-1}})(gx)$. By applying the same argument in [31, page 193], we have

$$\begin{aligned} & \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', P'}(\phi_x \otimes \phi'_x)(g) \phi''_{x, P'}(g) \tau_{P'}(H_{P'}(g) - T') dg \\ & \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} F_{P'}(g) (\phi_{x^{-1}})''_{P'}(gx) \tau_{P'}(H_{P'}(gx) - T') dg = \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} F_{P'}(g) \phi''_{P'}(g) \tau_{P'}(H_{P'}(gx) - T') dg \end{aligned}$$

because $(\phi_{x^{-1}})''_{P'}(gx) = \phi''_{P'}(g)$.

For $h \in G'(\mathbb{A}_F)$, let $K(h) \in K'$ be any element such that $hK(h)^{-1} \in P'_0(\mathbb{A}_F)$. Then using the same argument in [31, page 193-194], we have

$$F_{P'}(g) = \sum_{\substack{Q' \\ Q' \subseteq P'}} \sum_{\delta \in Q'(F) \backslash P'(F)} \Lambda_m^{T', Q'}(\phi \otimes \phi')(\delta g) \Gamma_{Q'}^{P'}(H_{Q'}(\delta g) - T', -H_{Q'}(K(\delta g)x)).$$

Applying the similar argument in the proof of (ii), we have

$$\begin{aligned} & \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \sum_{\delta \in Q'(F) \backslash P'(F)} \Lambda_m^{T', Q'}(\phi \otimes \phi')(\delta g) \phi''_{P'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(\delta g) - T', -H_{Q'}(K(\delta g)x)) \tau_{P'}(H_{P'}(gx) - T') dg \\ & = \int_{Q'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{Q'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(g) - T', -H_{Q'}(K(g)x)) \tau_{P'}(H_{P'}(gx) - T') dg. \end{aligned}$$

Note that $H_{Q'}(gx) - H_{Q'}(g) = H_{Q'}(K(g)x)$. By putting $H = H_{Q'}(g) - T'$, $X = H_{Q'}(K(g)x)$ in (5.3),

$$\begin{aligned} & \sum_{\substack{P' \\ Q' \subseteq P'}} \int_{Q'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{Q'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(g) - T', -H_{Q'}(K(g)x)) \tau_{P'}(H_{P'}(gx) - T') dg \\ & = \int_{Q'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{Q'}(g) \tau_{Q'}(H_{Q'}(g) - T') dg. \end{aligned}$$

By changing the order of integration and summation, we have

$$\begin{aligned}
& \sum_{P'} \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', P'}(\phi_x \otimes \phi'_x)(g) (\phi_x)''_{P'}(g) \tau_{P'}(H_{P'}(g) - T') dg \\
&= \sum_{P'} \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} F_{P'}(g) \phi''_{P'}(g) \tau_{P'}(H_{P'}(gx) - T') dg \\
&= \sum_{Q'} \sum_{\substack{P' \\ Q' \subseteq P'}} \int_{Q'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{Q'}(g) \Gamma_{Q'}^{P'}(H_{Q'}(g) - T', -H_{Q'}(K(g)x)) \tau_{P'}(H_{P'}(gx) - T') dg \\
&= \sum_{Q'} \int_{Q'(F) \backslash G'(\mathbb{A}_F)^1}^{\#} \Lambda_m^{T', Q'}(\phi \otimes \phi')(g) \phi''_{Q'}(g) \tau_{Q'}(H_{Q'}(g) - T') dg \\
&= \mathcal{P}(\phi, \phi', \phi'')
\end{aligned}$$

and so we proved (6.2) as required. (Changing the order of integration and summation in the third and fourth equality is justified by [5, Lemma 2.1]. We refer the reader to [31, page 194, 196] for the detail.)

The last assertion (iv) essentially follows from the fact that

$$\int_{G'(F) \backslash G'(\mathbb{A}_F)^1} (\phi \otimes \phi')_{G, N}(g) \phi''(g) F^{G'}(g, T') dg - \int_{G'(F) \backslash G'(\mathbb{A}_F)^1} \Lambda_m^{T'}(\phi \otimes \phi')(g) \phi''(g) dg \rightarrow 0$$

as $T' \rightarrow \infty$ in the positive Weyl chamber. (Here $F^{G'}(g, T')$ is the function defined just before Lemma 5.3.) Since the proof is almost same with that of [30, Proposition 3.8, Cor. 3.10], we omit the detail. \square

Let $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^{**}$ be the subspace of all triplets $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$ such that

$$\langle \lambda_1 + \lambda_2 + \lambda_3 + \rho_P, \omega^{\vee} \rangle \neq 0 \quad \text{for all } \omega^{\vee} \in (\hat{\Delta}^{\vee})_{Q'}^{P'}, \lambda_1 \in \mathcal{E}_Q(\phi_Q), \lambda_2 \in \mathcal{E}_{Q'}(\phi_{Q'}), \lambda_3 \in \mathcal{E}_{Q'}(\phi''_{Q'})$$

for all pairs of parabolic subgroups $Q' \subset P'$ of G' . Clearly $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^{**} \subset \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^*$.

If $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^{**}$, then the regularized integral

$$\begin{aligned}
& \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{P, N}(g) \phi''_{P'}(g) \hat{\tau}_{P'}(H_{P'}(g) - T') dg \\
&= \int_{K'} \int_{M'(F) \backslash M'(\mathbb{A}_F)^1} \int_{\mathfrak{a}_{P'}}^{\#} (\phi \otimes \phi')_{P, N}(e^X mk) \phi''_{P'}(e^X mk) \hat{\tau}_{P'}(X - T') e^{-2\langle \rho_{P'}, X \rangle} dX dm dk
\end{aligned}$$

is well-defined for all proper parabolic subgroups P' of G' by [31, Lemma 3].

Proposition 6.4. *If $(\phi, \phi', \phi'') \in \mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi}^{**}$, then*

$$\begin{aligned}
& \int_{G'(F) \backslash G'(\mathbb{A}_F)^1} \Lambda_m^{T'}(\phi \otimes \phi')(g) \phi''(g) dg \\
&= \sum_{P'} (-1)^{\dim \mathfrak{a}_{P'}} \int_{P'(F) \backslash G'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{P, N}(g) \cdot \phi''_{P'}(g) \hat{\tau}_{P'}(H_{P'}(g) - T') dg.
\end{aligned}$$

Proof. We prove this by induction on the split rank of G' . Assume that it holds for groups $M_{P'}$, where $M_{P'}$ is the Levi subgroup of a proper parabolic subgroup P' of G' .

By Proposition 6.3 (ii), we can write

$$\int_{G'(F) \backslash G'(\mathbb{A}_F)^1} \Lambda_m^{T'}(\phi \otimes \phi')(g) \phi''(g) dg = \mathcal{P}(\phi, \phi', \phi'') - \sum_{P' \subsetneq G'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'').$$

Using the similar argument in [31, Theorem 10], we can show that $\mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'')$ is equal to

$$\sum_{R' \subset P'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \int_{R'(F) \backslash R'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{R,N}(g) \cdot \phi''_{R'}(g) \hat{\tau}_{R'}^{P'}(H_{R'}(g) - T') \tau_{P'}(H_{P'}(g) - T') dg$$

from our induction hypothesis. (See also [59, Proposition 3.5].)

Thus

$$\begin{aligned} & \sum_{P' \subsetneq G'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'') \\ &= \sum_{R' \subset P' \subsetneq G'} (-1)^{\dim \mathfrak{a}_{R'}^{P'}} \int_{R'(F) \backslash R'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{R,N}(g) \cdot \phi''_{R'}(g) \hat{\tau}_{R'}^{P'}(H_{R'}(g) - T') \tau_{P'}(H_{P'}(g) - T') dg \\ &= \sum_{R' \subsetneq G'} -(-1)^{\dim \mathfrak{a}_{R'}} \int_{R'(F) \backslash R'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{R,N}(g) \cdot \phi''_{R'}(g) \hat{\tau}_{R'}(H_{R'}(g) - T') dg \end{aligned}$$

by Langlands' combinatorial lemma. Thus we proved

$$\mathcal{P}(\phi, \phi', \phi'') - \sum_{P' \subsetneq G'} \mathcal{P}_{P'}^{T'}(\phi, \phi', \phi'') = \sum_{P'} (-1)^{\dim \mathfrak{a}_{P'}} \int_{P'(F) \backslash P'(\mathbb{A}_F)^1}^* (\phi \otimes \phi')_{P,N}(g) \cdot \phi''_{P'}(g) \hat{\tau}_{P'}(H_{P'}(g) - T') dg,$$

as desired. \square

We close this section by giving the definition of the *regularized Bessel periods* and the *regularized Fourier-Jacobi periods* for all classical and metaplectic groups.

Let $\nu_{\psi^{-1}, \mu^{-1}}$ be a global generic Weil representation of $(N_{n,r} \rtimes \mathbf{G}')(\mathbb{A}_F)$ and $\Theta_{\psi^{-1}, \mu^{-1}, W_m}(\cdot)$ a theta series functional which gives a realization of $\nu_{\psi^{-1}, \mu^{-1}}$ in $\mathcal{A}(N_{n,r} \rtimes \mathbf{G}')$. Let χ' be a generic automorphic character of $N_{n,r}(\mathbb{A}_F)$, that is stable under the conjugate action of G' in G . We view χ' as an automorphic character of $(N_{n,r} \rtimes \mathbf{G}')(\mathbb{A}_F)$, which is trivial on G' .

Definition 6.5. Let $\varphi_1 \in \mathcal{A}(\mathbf{G})$, $\varphi_2 \in \mathcal{A}(\mathbf{G}')$ and $f \in \nu_{\psi^{-1}, \mu^{-1}, W_m}$. Suppose that $(\varphi_1, \chi', \varphi_2)$ and $(\varphi_1, \Theta_{\psi^{-1}, \mu^{-1}, W_m}(f), \varphi_2)$ belong to $\mathcal{A}(\mathbf{G}, \mathbf{G}')_{\chi'}^*$.

When $n - m = 2r + 1$, the **regularized Bessel period** for (ϕ_1, ϕ_2) is defined by

$$\mathcal{B}_{\chi'}(\varphi_1, \varphi_2) := \mathcal{P}(\varphi_1, \chi', \varphi_2).$$

When $n - m = 2r$, the **regularized Fourier-Jacobi period** for (ϕ_1, ϕ_2, f) is defined by

$$\mathcal{FJ}_{\psi, \mu}(\varphi_1, \varphi_2, f) := \mathcal{P}(\varphi_1, \Theta_{\psi^{-1}, \mu^{-1}, W_m}(f), \varphi_2).$$

Remark 6.6. When $N = 1$, Remark 4.1 and Proposition 6.3 (ii) show that \mathcal{P} is exactly the same regularized period functional defined in [30] and [58] in each case. Proposition 6.3 (iv) shows that the above definitions coincide with those of the Bessel period and Fourier-Jacobi period defined on cusp forms. Therefore we can state [17, Conjecture 9.1] more concretely with our regularized periods.

7. Jacquet modules corresponding to Fourier-Jacobi characters

Starting this section, we prove one implication in the global GGP conjecture for skew hermitian spaces. In this section, we shall prove a lemma which plays a key role in proving Lemma 9.4, which asserts the vanishing of the regularized Fourier-Jacobi periods involving residual representations.

Throughout the rest of the paper, we assume that E/F is a quadratic extension number fields. Then for a place v of F ,

$$E_v = \begin{cases} F_v \oplus F_v, & \text{when } v \text{ splits in } E \\ \text{quadratic extension of } F_v, & \text{when } v \text{ non-splits in } E. \end{cases}$$

Let $\sigma \in \text{Gal}(E_v/F_v)$ be the non-trivial element. If $E_v = F_v \oplus F_v$, σ is defined by

$$\sigma(x, y) = (y, x).$$

Since we shall only consider the local situation, we suppress v from the notation until the end of this section.

Let $\psi : F \rightarrow \mathbb{C}^\times$ be an additive character and $\mu : E^\times \rightarrow \mathbb{C}^\times$ a multiplicative character defined similarly as in the global case. Let W_n be a free left E -module of rank n which has a skew-hermitian structure (\cdot, \cdot) and G_n be its unitary group. Let k be the dimension of a maximal totally isotropic subspace of W_n and we assume $k > 0$. We fix maximal totally isotropic subspaces X and X^* of W_n , in duality, with respect to (\cdot, \cdot) . Fix a complete flag in X

$$0 = X_0 \subset X_1 \subset \cdots \subset X_k = X,$$

and choose a basis $\{e_1, e_2, \dots, e_j\}$ of X_j such that $\{e_1, \dots, e_j\}$ is a basis of X_j for $1 \leq j \leq k$. Let $\{f_1, f_2, \dots, f_k\}$ be the basis of X^* which is dual to the fixed basis of X , i.e., $(e_i, f_j) = \delta_{ij}$ for $1 \leq i, j \leq k$, where $\delta_{i,j}$ denotes the Kronecker delta. We write X_j^* for the subspace of X^* spanned by $\{f_1, f_2, \dots, f_j\}$ and W_{n-2j} for the orthogonal complement of $X_j + X_j^*$ in W_n .

For each $1 \leq j \leq k$, denote by $P_{n,j}$ the parabolic subgroup of G_n stabilizing X_j , by $U_{n,j}$ its unipotent radical and by $M_{n,j}$ the Levi subgroup of $P_{n,j}$ stabilizing X_j^* . Then $M_{n,j} \simeq GL(X_j) \times G_{n-2j}$. (Here, we regard $GL(X_j) \simeq GL_j(E)$ as the subgroup of $M_{n,j}$ which acts as the identity map on W_{n-2j} .) Let P_0 be the minimal parabolic subgroup of G_n which stabilizes the above complete flag of X . We also fix a good maximal compact subgroup K_n of G_n with respect to P_0 (see [41, I. 1.4]).

For $0 \leq j \leq k$, write $N_{n,j}$ (resp. \mathcal{N}_j) for the unipotent radical of the parabolic subgroup of G_n (resp., $GL(X_j)$) stabilizing the flag $\{0\} = X_0 \subset X_1 \subset \cdots \subset X_j$. If we regard \mathcal{N}_j as a subgroup of $M_{n,j} \simeq GL(X_j) \times G_{n-2j}$, it acts on $U_{n,j}$ and so $N_{n,j} = U_{n,j} \rtimes \mathcal{N}_j$. When $j = 0$, \mathcal{N}_0 denotes the trivial group.

For any $0 \leq j < k$, $(\text{Res}_{E/F}(W_{n-2j}), \text{Tr}_{E/F}(\cdot, \cdot))$ is a nondegenerate symplectic F -space of F -dimension $2(n-2j)$. Let $\mathcal{H}_{W_{n-2j}} = \text{Res}_{E/F}(W_{n-2j}) \oplus F$ be the Heisenberg group associated to $(\text{Res}_{E/F}(W_{n-2j}), \text{Tr}_{E/F}(\cdot, \cdot))$ and $\Omega_{\psi^{-1}, \mu^{-1} W_{n-2j}}$ be the Weil representation of $\mathcal{H}_{W_{n-2j}} \rtimes G_{n-2j}$ with respect to ψ^{-1}, μ^{-1} . Then since $N_{n,j-1} \backslash N_{n,j} \simeq \mathcal{H}_{W_{n-2j}}$ and $\mathcal{N}_j \subseteq N_{n,j-1}$, we can pull back $\Omega_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$ to $N_{n,j} \rtimes G_{n-2j}$ and denote it by the same symbol $\Omega_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$. When E is a field, $\text{Tr}_{E/F}$ and $\text{N}_{E/F}$ denote the usual trace and norm map, respectively. When $E = F \oplus F$, define

$$\text{Tr}_{E/F}(x, y) = x + y, \quad \text{N}_{E/F}(x, y) = xy, \quad (x, y) \in E.$$

When $j \geq 2$, we define a character $\lambda_j : \mathcal{N}_j \rightarrow \mathbb{C}^\times$ by

$$\lambda_j(n) = \psi(\text{Tr}_{E/F}((ne_2, f_1) + (ne_3, f_2) + \cdots + (ne_j, f_{j-1}))), \quad n \in \mathcal{N}_j.$$

When $j = 0, 1$, λ_j denotes the trivial character. Put $\nu_{\psi^{-1}, \mu^{-1}, W_{n-2j}} = \lambda_j \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$ and denote $H_{n,j} = N_{n,j} \rtimes G_{n-2j}$. We can embed $H_{n,j}$ into $G_n \times G_{n-2j}$ by inclusion on the first factor and projection on the second factor. Then $\nu_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$ is a smooth representation of $H_{n,j} = N_{n,j} \rtimes G_{n-2j}$ and up to conjugation of the normalizer of $H_{n,j}$ in $G_n \times G_{n-2j}$, it is uniquely determined by ψ modulo $N_{E/F}(E)^\times$ and μ . We shall denote by $\omega_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$ the restriction of $\nu_{\psi^{-1}, \mu^{-1}, W_{n-2j}}$ to G_{n-2j} .

For $0 \leq l \leq k-1$, we define a character ψ_l of $N_{n,l+1}$, which factors through the quotient $n: N_{n,l+1} \rightarrow U_{n,l+1} \backslash N_{n,l+1} \simeq \mathcal{N}_{l+1}$, by setting

$$\psi_l(u) = \lambda_{l+1}(n(u)).$$

Denote by q the cardinality of the residue field of F and write $q_E = q^2$.

For an irreducible smooth representation π' of G_n , write $J_{\psi_l}(\pi' \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_{n-2l-2}})$ for the Jacquet module of $\pi' \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_{n-2l-2}}$ with respect to the group $N_{n,l+1}$ and its character ψ_l . It is a representation of the unitary group G_{n-2l-2} . We use the symbol Ind to denote the normalized induction functor, ind for the unnormalized induction functor and c-ind for compactly supported induction.

Lemma 7.1. *Let n, m, a positive integers such that $2r = n - m \geq 0$. Let \mathcal{E} be a smooth representation of G_{m+2a} of finite length and σ, π and Σ be irreducible smooth representations of $GL(X_a), G_n$ and G_{n+2a} , respectively. Then*

(i) If $r > 0$,

$$\begin{aligned} & \text{Hom}_{N_{n+2a,r} \rtimes G_{m+2a}} (\Sigma \otimes \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C}) \\ & \simeq \text{Hom}_{G_{n+2a}} \left(J_{\psi_{r-1}^{-1}} (\Sigma \otimes \Omega_{\psi^{-1}, \nu^{-1}, W_{m+2a}}), \mathcal{E}^\vee \right) \end{aligned}$$

(ii) If $r = 0$,

$$\begin{aligned} & \dim_{\mathbb{C}} \text{Hom}_{G_{m+2a}} \left(\text{Ind}_{P_{m+2a,a}}^{G_{m+2a}} (\sigma | \cdot |_E^s \boxtimes \pi) \otimes \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C} \right) \\ & \leq \dim_{\mathbb{C}} \text{Hom}_{G_m} (J_{\psi_{a-1}^{-1}} (\mathcal{E} \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m}), \pi^\vee) \end{aligned}$$

except for finitely many q_E^{-s} .

Proof. Since the case $E = F \oplus F$ is quite similar, we give the proof only when E is a field.

We first prove (i). By the Frobenius reciprocity,

$$\begin{aligned} & \text{Hom}_{N_{n+2a,r} \rtimes G_{m+2a}} (\Sigma \otimes \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C}) \\ & = \text{Hom}_{N_{n+2a,r} \rtimes G_{m+2a}} (\Sigma \otimes \lambda_r \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C}) \\ & \simeq \text{Hom}_{G_{m+2a}} \left(J_{\psi_{r-1}^{-1}} (\Sigma \otimes \Omega_{\psi^{-1}, \nu^{-1}, W_{m+2a}}) \otimes \mathcal{E}, \mathbb{C} \right) \\ & \simeq \text{Hom}_{G_{m+2a}} \left(J_{\psi_{r-1}^{-1}} (\Sigma \otimes \Omega_{\psi^{-1}, \nu^{-1}, W_{m+2a}}), \mathcal{E}^\vee \right). \end{aligned}$$

Next, we prove (ii). The proof is quite similar with [58, Lemma 4.1]. By the Frobenius reciprocity,

$$\text{Hom}_{G_{m+2a}} \left(\text{Ind}_{P_{m+2a,a}}^{G_{m+2a}} (\sigma | \cdot |_E^s \boxtimes \pi) \otimes \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C} \right)$$

is isomorphic to

$$(7.1) \quad \text{Hom}_{P_{m+2a,a}} \left((\sigma | \cdot |_E^{s-s_0} \boxtimes \pi) \otimes \omega_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C} \right)$$

where s_0 is defined so that $\delta_{P_{m+2a,a}}^{\frac{1}{2}} = | \cdot |_E^{s_0}$. For $1 \leq i \leq a$, write $\mathcal{P}_{a,i}$ for the subgroup of $GL(X_a)$ which stabilizes the flag $X_{a-i} \subset X_{a-i+1} \subset \cdots \subset X_{a-1}$ and fixes e_j modulo X_{j-1} for $a-i+1 \leq j \leq a$. Put

$Q_{a,i} = (\mathcal{P}_{a,i} \times G_m) \rtimes U_{m+2a,a}$. Using a similar argument to the proof of [18, Theorem 16.1], we have an exact sequence

(7.2)

$$0 \rightarrow \text{c-ind}_{Q_{a,1}}^{P_{m+2a,a}} \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \rightarrow \omega_{\psi^{-1}, \mu^{-1}, W_{m+2a}} |_{P_{m+2a,a}} \rightarrow \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \rightarrow 0,$$

where c-ind is the unnormalized compact induction functor. Since \mathcal{E} is admissible, we have

$$(7.3) \quad \begin{aligned} & \text{Hom}_{P_{m+2a,a}} \left((\sigma | \cdot | \frac{1}{2} \boxtimes \pi) \otimes (\mu^{-1} | \cdot | \frac{1}{2} \boxtimes \Omega_{\psi^{-1}, \mu^{-1}, W_m}) \otimes \mathcal{E}, \mathbb{C} \right) \\ & \simeq \text{Hom}_{M_{m+2a,a}} \left(\left((\sigma \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \pi) \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \right) \otimes J_{P_{m+2a,a}}(\mathcal{E}), \mathbb{C} \right), \end{aligned}$$

where $J_{P_{m+2a,a}}$ is the Jacquet functor with respect to $U_{m+2a,a}$ and its trivial character. Since the central character of any irreducible subquotient of $J_{P_{m+2a,a}}(\mathcal{E})|_{GL(X_a)} \otimes \sigma \mu^{-1} | \cdot | \frac{1}{2} \boxtimes$ is not trivial for almost all q_E^{-s} , the Hom space in (7.3) is zero for almost all q_E^{-s} . By tensoring the exact sequence (7.2) with $\mathcal{E} \otimes (\sigma | \cdot | \frac{1}{2} \boxtimes \pi)$ and applying the $\text{Hom}_{P_{m+2a,a}}(-, \mathbb{C})$ functor, we can regard (7.1) as a subspace of

$$\begin{aligned} & \text{Hom}_{P_{m+2a,a}} \left((\sigma | \cdot | \frac{1}{2} \boxtimes \pi) \otimes (\text{c-ind}_{Q_{a,1}}^{P_{m+2a,a}} \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \Omega_{\psi^{-1}, \mu^{-1}, W_m}) \otimes \mathcal{E}, \mathbb{C} \right) \\ & \simeq \text{Hom}_{Q_{a,1}} \left((\sigma | \cdot | \frac{1}{2} \boxtimes \pi) \otimes (\delta_{P_{m+2a,a}} \delta_{Q_{a,1}}^{-1} \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \Omega_{\psi^{-1}, \mu^{-1}, W_m}) \otimes \mathcal{E}, \mathbb{C} \right) \\ & \simeq \text{Hom}_{Q_{a,1}} \left((\sigma \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \pi) \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \right) \otimes \mathcal{E}, \mathbb{C} \end{aligned}$$

for almost all s , where s_1 is defined so that $\delta_{P_{m+2a,a}} \delta_{Q_{a,1}}^{-1} | \cdot | \frac{1}{2} \boxtimes \pi = | \cdot | \frac{1}{2} \boxtimes \pi$. (Here we have used the Frobenius reciprocity law again.)

Note that $\mathcal{P}_{a,1}$ is a mirabolic subgroup of $GL(X_a)$. By the Bernstein and Zelevinsky's result (See [8, Sec. 3.5]), the restriction of σ to $\mathcal{P}_{a,1}$ has a filtration

$$\{0\} = \sigma_{a+1} \subset \sigma_a \subset \cdots \subset \sigma_2 \subset \sigma_1 = \sigma$$

such that $\sigma_j / \sigma_{j+1} \simeq (\Phi^+)^{j-1} \Psi^+ \sigma^{(j)}$ for $1 \leq j \leq a$. (For the definition of Φ^+ and Ψ^+ , see [8, Sec. 3].) Let $X_{a,i}$ be the subspace of X_a generated by $e_{a-i+1}, e_{a-i+2}, \dots, e_a$ and put $\mathcal{N}_{a,i}$ by $\mathcal{N}_a \cap GL(X_{a,i})$. Then for $1 \leq j \leq a$,

$$(\Phi^+)^{j-1} \Psi^+ \sigma^{(j)} \simeq \text{c-ind}_{\mathcal{P}_{a,j}}^{\mathcal{P}_{a,1}} | \cdot | \frac{1}{2} \boxtimes \sigma^{(j)} \boxtimes \lambda_a |_{\mathcal{N}_{a,j}}.$$

By applying the Frobenius reciprocity law, we have

$$\text{Hom}_{Q_{a,1}} \left(\left((\Phi^+)^{j-1} \Psi^+ \sigma^{(j)} \mu^{-1} | \cdot | \frac{1}{2} \boxtimes \pi \right) \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \right) \otimes \mathcal{E}, \mathbb{C} \right)$$

(7.4)

$$\simeq \text{Hom}_{GL(X_{a-j}) \times (\mathcal{N}_{a,j} \times G_m) \rtimes U_{m+2j,j}} \left((| \cdot | \frac{j+1}{2} \boxtimes \pi \mu^{-1} \sigma^{(j)} \boxtimes \lambda_a |_{\mathcal{N}_{a,j}} \boxtimes (\pi \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m})) \otimes J_{P_{m+2a,a-j}}(\mathcal{E}), \mathbb{C} \right)$$

for some $s'_1 \in \mathbb{R}$ (depending on j). For $1 \leq j \leq a-1$, the central character of any irreducible subquotient of $(| \cdot | \frac{j+1}{2} \boxtimes \pi \mu^{-1} \sigma^{(j)}) \otimes J_{P_{m+2a,a-j}}(\mathcal{E})|_{GL(X_{a-j})}$ is not trivial for almost all q_E^{-s} . Thus the Hom space in (7.4) is zero for almost all q_E^{-s} unless $j = a$, in which case it is isomorphic to

$$\begin{aligned} & \text{Hom}_{Q_{a,a}} ((\psi_{a-1} \boxtimes \pi) \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m} \otimes \mathcal{E}, \mathbb{C}) \\ & \simeq \left(\text{Hom}_{G_m} (J_{\psi_{a-1}^{-1}}(\mathcal{E} \otimes \Omega_{\psi^{-1}, \mu^{-1}, W_m}), \pi^\vee) \right)^{\oplus t}, \end{aligned}$$

where $t = \dim_{\mathbb{C}} \sigma^{(a)}$. Since $t \leq 1$, our claim is proved. \square

Remark 7.2. Lemma 7.1 can be used to control the size of the Hom space

$$\mathrm{Hom}_{N_{n+2a,r} \times G_{m+2a}} \left(\mathrm{Ind}_{P_{n+2a,a}}^{G_{n+2a}} (\sigma | \cdot |_E^s \boxtimes \pi) \otimes \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} \otimes \mathcal{E}, \mathbb{C} \right)$$

in terms of the relevant Jacquet modules. When $r > 0$, the existence of $N_{n+2a,r}$ in this Hom space becomes essential because without it the Hom space would have infinite dimension. This leads us to distinguish the cases of $r > 0$ and $r = 0$ separately.

8. Eisenstein series and residual representation

In this section, we introduce the residual Eisenstein series representations. For an irreducible cuspidal automorphic representation π of $G_n(\mathbb{A}_F)$ and σ of $GL_a(\mathbb{A}_E)$, write $L(s, \sigma \times \pi)$ for the Rankin-Selberg L -function $L(s, \sigma \times BC(\pi))$. We also write $L(s, \sigma, As^+)$ for the Asai L -function of σ and $L(s, \sigma, As^-)$ for the μ -twisted Asai L -function $L(s, \sigma \otimes \mu, As^+)$ (cf., [18, Section 7]).

For $g \in G_{n+2a}(\mathbb{A}_F)$, write $g = m_g u_g k$ for $m_g \in M_{n+2a,a}(\mathbb{A}_F)$, $u_g \in U_{n+2a,a}(\mathbb{A}_F)$, $k_g \in K$. Since $M_{n+2a,a} \simeq GL(X_a) \times G_n$, we decompose $m_g = m_1 m_2$, where $m_1 \in GL(X_a)(\mathbb{A}_E)$, $m_2 \in G_n(\mathbb{A}_F)$. Define $d(g) = |\det m_1|_{\mathbb{A}_E}$. Then for $\phi \in \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a})$, the associated Eisenstein series with ϕ is defined by

$$E(g, \phi, z) = \sum_{P_{n+2a,a}(F) \backslash G_{n+2a}(F)} \phi(\gamma g) d(\gamma g)^z.$$

From the theory of Eisenstein series, it converges absolutely when $\mathrm{Re}(z)$ is sufficiently large and admits a meromorphic continuation to the whole complex plane. [41, IV.1.8].

Remark 8.1. For $\phi \in \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a})$, we can uniquely extend it to a flat section $\phi_z \in \mathcal{A}_{P_{n+2a,a}}^{\sigma | \cdot |^z \boxtimes \pi}(G_{n+2a})$ which satisfying $\phi_z(k) = \phi(k)$ for $k \in K$. Then $E(g, \phi, z) = E(g, \phi_z, 0)$ and the map $\phi_z \rightarrow E(g, \phi_z, 0)$ gives an automorphic realization of the induced representation $\mathrm{Ind}_{P_{n+2a,a}(\mathbb{A}_F)}^{G_{n+2a}(\mathbb{A}_F)} (\sigma | \cdot |_E^z \boxtimes \pi)$.

One of the properties of tempered representations (i.e., Remark 1.1 (iii)) plays an essential role in the proof of the following proposition.

Proposition 8.2 ([30], Proposition 5.3, [34], Proposition 5.2). *Let π be an irreducible tempered cuspidal automorphic representation of $G_n(\mathbb{A}_F)$ and σ an irreducible cuspidal automorphic representation of $GL_a(\mathbb{A}_E)$. For $\phi \in \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a})$, the Eisenstein series $E(\phi, z)$ has at most a simple pole at $z = \frac{1}{2}$ and $z = 1$. Moreover, it has a pole at $z = \frac{1}{2}$ as ϕ varies if and only if $L(s, \sigma \times \pi^\vee)$ is non-zero at $s = \frac{1}{2}$ and $L(s, \sigma, As^{(-1)^{n-1}})$ has a pole at $s = 1$. Furthermore, it has a pole at $z = 1$ as ϕ varies if and only if $L(s, \sigma \times \pi^\vee)$ has a pole at $s = 1$.*

Thanks to Proposition 8.2, we can define the residues of the Eisenstein series to be the limits

$$\mathcal{E}^0(\phi) = \lim_{z \rightarrow \frac{1}{2}} (z - \frac{1}{2}) E(\phi, z), \quad \mathcal{E}^1(\phi) = \lim_{z \rightarrow 1} (z - 1) E(\phi, z) \quad \text{for } \phi \in \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a}).$$

For $i = 0, 1$, let $\mathcal{E}^i(\sigma, \pi)$ be the residual representation of $G_{n+2a}(\mathbb{A}_F)$ generated by $\mathcal{E}^i(\phi)$ for $\phi \in \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a})$. Since the residual representation is isomorphic to the image of some relevant intertwining operator, the residue map $\mathcal{E}^i : \mathcal{A}_{P_{n+2a,a}}^{\sigma \boxtimes \pi}(G_{n+2a}) \rightarrow \mathcal{E}^i(\sigma, \pi)$ is G_{n+2a} -equivariant.

Remark 8.3. As we mentioned in Remark 1.1 (ii), we can write $BC(\pi)$ as an isobaric sum of the form $\sigma_1 \boxplus \cdots \boxplus \sigma_t$, where $\sigma_1, \dots, \sigma_t$ are distinct irreducible cuspidal automorphic representations of some general linear groups such that the (twisted) Asai L -function $L(s, \sigma_i, As^{(-1)^{n-1}})$ has a pole at $s = 1$. Since $L(s, \sigma \times \pi^\vee) = \prod_{i=1}^t L(s, \sigma \times \sigma_i^\vee)$, Proposition 8.2 implies that $\mathcal{E}^1(\sigma, \pi)$ is non-zero if and only if $\sigma \simeq \sigma_i$ for some $1 \leq i \leq t$.

Remark 8.4. Let $c: E \rightarrow E$ be the non-trivial element in $\text{Gal}(E/F)$ and use the same notation for the automorphism of $GL_n(E)$ induced by $c: E \rightarrow E$. For a representation σ of $GL_n(\mathbb{A}_E)$, define $\sigma^c := \sigma \circ c$. Note that $L(s, \sigma, As^\pm)$ are nonzero at $s = 1$ by [50, Theorem 5.1]. Thus if $L(s, \sigma, As^{(-1)^{n-1}})$ has a pole at $s = 1$, the Rankin–Selberg L -function

$$L(s, \sigma \times \sigma^c) = L(s, \sigma, As^+) \cdot L(s, \sigma, As^-)$$

has a simple pole at $s = 1$ and so $\sigma^c \simeq \sigma^\vee$.

Remark 8.5. Let π be an irreducible tempered cuspidal automorphic representation of $G_n(\mathbb{A}_F)$ and σ an irreducible cuspidal automorphic representation of $GL_a(\mathbb{A}_E)$. Let \mathcal{E} be an automorphic representation of G_{m+2a} . Then for $\phi \in \text{Ind}_{P_{n+2a,a}}^{G_{n+2a}}(\sigma \boxtimes \pi)$, $f \in \nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}$ and $\varphi \in \mathcal{E}$, $\mathcal{P}(E(\phi, z), \Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(f), \varphi)$ is well defined for all $z \in \mathbb{C}$ but the poles of $E(\phi, z)$. By the meromorphy of the Eisenstein series $E(\phi, z)$ and Proposition 6.3 (ii), $\mathcal{P}(E(\phi, z), \Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(f), \varphi)$ extends to a meromorphic function on the whole complex plane with possible poles contained in the set of poles of $E(\phi, z)$.

9. Reciprocal non-vanishing of the Fourier–Jacobi periods

In this section, we prove Theorem 9.9 asserting the reciprocal non-vanishing of the Fourier–Jacobi periods, which plays a crucial role to prove our main theorem. To prove it, we require some lemmas.

Let m, a be positive integers and r non-negative integer. Write $n = m + 2r$ and let $(W_{n+2a}, (\cdot, \cdot))$ be a skew-hermitian spaces over E of dimension $n + 2a$. Let X, X^* be both r -dimensional isotropic subspaces of W_{n+2a} which are in dual with respect to (\cdot, \cdot) and $(W_{m+2a}, (\cdot, \cdot))$ be its orthogonal complement. Then one has the polar decomposition $W_{n+2a} = X \oplus W_{m+2a} \oplus X^*$.

Suppose that $(W_{m+2a}, (\cdot, \cdot))$ also has the polar decomposition $W_{m+2a} = Y_a \oplus W_m \oplus Y_a^*$, where Y_a, Y_a^* are both a -dimensional isotropic subspaces of W_{m+2a} which are in dual with respect to (\cdot, \cdot) .

For $h \in GL(Y_a(\mathbb{A}_F))$, $b \in \text{Hom}(W_m, Y_a)$, $c \in \text{Hom}(Y_a^*, Y_a)$, we define $h^* \in GL(Y_a^*(\mathbb{A}_E))$, $b^* \in \text{Hom}(Y_a^*, W_m)$, $c^* \in \text{Hom}(Y_a, Y_a^*)$ by requiring that

$$(h^*x, y) = (x, hy), \quad (b^*x, z) = (x, bz), \quad (c^*y, x) = (y, cx) \quad \text{for all } x \in Y_a^*(\mathbb{A}_E), y \in Y_a(\mathbb{A}_E), z \in W_m(\mathbb{A}_E).$$

Let $(\rho_a, \mathcal{S}(Y_a^*(\mathbb{A}_E)))$ be the Heisenberg representation of $\mathcal{H}_{Y_a \oplus Y_a^*}(\mathbb{A}_F)$ with respect to ψ^{-1} and (ρ^0, \mathcal{S}^0) be the fixed Heisenberg representation of $\mathcal{H}_{W_m}(\mathbb{A}_F)$ with respect to ψ^{-1} . Then $\mathcal{S} = \mathcal{S}(Y_a^*(\mathbb{A}_E)) \otimes \mathcal{S}^0$ provides a mixed model for both the Heisenberg representation ρ of $\mathcal{H}_{W_{m+2a}}(\mathbb{A}_F)$ and the global Weil representation $\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}} = \otimes \nu_{\psi_v^{-1}, \mu_v^{-1}, W_{m+2a}}$ of $H_{n+2a, r}(\mathbb{A}_F)$. (Here, we regard \mathcal{S} as a space of functions on Y_a^* whose values in \mathcal{S}^0 .)

Using [[40], Theorem 3.1], we can describe the (partial) $N_{n, r}(\mathbb{A}_F) \times P_{m+2a, a}(\mathbb{A}_F)$ -action of the global Weil representation $\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}$ on \mathcal{S} as follows: (see also [[20], page 35])

For $f \in \mathcal{S}$ and $y \in Y_a^*(\mathbb{A}_E)$,

$$(9.1) \quad (\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(n)f)(y) = \nu_{\psi^{-1}, \mu^{-1}, W_m}(n)(f(y)), \quad n \in N_{n, r}(\mathbb{A}_F),$$

$$(9.2) \quad (\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(g_0)f)(y) = \nu_{\psi^{-1}, \mu^{-1}, W_m}(g_0)(f(y)), \quad g_0 \in U(W_m),$$

$$(9.3) \quad (\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(h_0)f)(y) = \mu^{-1}(h_0) |\det(h_0)|_{\mathbb{A}_E}^{\frac{1}{2}} \cdot f(h_0^*y), \quad h_0 \in GL(Y_a(\mathbb{A}_E)),$$

$$(9.4) \quad (\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(b)f)(y) = \rho^0(b^*y, 0)(f(y)), \quad b \in \text{Hom}(W_m, Y_a),$$

$$(9.5) \quad (\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(c)f)(y) = \psi^{-1}\left(\frac{\text{Tr}_{E/F}(cy, y)}{2}\right) \cdot f(y), \quad c \in \text{Herm}(Y_a^*, Y_a),$$

$$(9.6) \quad (\rho(x+x', 0)f)(y) = \psi^{-1}\left(\text{Tr}_{E/F}(y, x) + \frac{\text{Tr}_{E/F}(x', x)}{2}\right) \cdot f(y+x'), \quad x \in Y_a(\mathbb{A}_E), x' \in Y_a^*(\mathbb{A}_E)$$

where

$$\text{Herm}(Y_a^*, Y_a) = \{c \in \text{Hom}(Y_a^*, Y_a) \mid c^* = -c\}$$

and we identified $\text{Herm}(Y_a^*, Y_a)$ and $\text{Hom}(W_m, Y_a)$ as subgroups of $U_{m+2a,a}$ via the canonical isomorphism $U_{m+2a,a} \simeq \text{Hom}(W_m, Y_a) \times \text{Herm}(Y_a^*, Y_a)$.

By choosing a theta functional $\theta_1 : \mathcal{S}^0 \rightarrow \mathbb{C}$, we define theta functions associated to \mathcal{S} as follows:

For $f \in \mathcal{S}(Y_a^*(\mathbb{A}_E)) \otimes \mathcal{S}^0$, its associated theta function $\Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(\cdot, f)$ is defined by

$$\Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(h, f) = \sum_{y \in Y_a(E)} \theta_1((\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(h)f)(y)),$$

where $h = ((u, n), g) \in ((U_{n+2a,r} \rtimes \mathcal{N}_r) \times G_{m+2a})(\mathbb{A}_F) = (N_{n+2a,r} \times G_{m+2a})(\mathbb{A}_F) = H_{n+2a,r}(\mathbb{A}_F)$.

Then $\Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(f) \in \mathcal{A}(H_{n+2a,r})$ and the space of these theta functions $\{\Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(f) \mid f \in \mathcal{S}\}$ is a realization of the global Weil representation $\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}$ as an automorphic representation of $H_{n+2a,r}(\mathbb{A}_F)$. By one of main results of Weil [54], the global Weil representation has a unique (up to scaling) automorphic realization. Thus we can choose θ_1 so that this theta function coincides with the one we defined in Section 1.

Since we have fixed μ, ψ , we simply write $\nu_{W_{m+2a}}$ for $\nu_{\psi^{-1}, \mu^{-1}, W_{m+2a}}$ and its associated theta function $\Theta_{\psi^{-1}, \mu^{-1}, W_{m+2a}}(f)$ as $\Theta_{W_{m+2a}}(f)$. Write $\Theta_{P_{m+2a,a}}(f)$ for the constant term of $\Theta_{W_{m+2a}}(f)$ along $P_{m+2a,a}$.

From now on, G_k denotes the isometry group of W_k for various k .

Lemma 9.1 (cf. [58], Lemma 6.2)]. For $h \in H_{n+2a,r}(\mathbb{A}_F)$, $f \in \mathcal{S}(Y_a^*(\mathbb{A}_E)) \otimes \mathcal{S}^0$,

$$\Theta_{P_{m+2a,a}}(h, f) = \theta_1((\nu_{W_{m+2a}}(h)f)(0)).$$

Proof. Using the isomorphism $U_{m+2a,a} \simeq \text{Hom}(W_m, Y_a) \times \text{Herm}(Y_a^*, Y_a)$, we take Haar measure $du = db dc$ on $U_{m+2a,a}$ for $u = bc$ with $b \in \text{Hom}(W_m, Y_a)$, $c \in \text{Herm}(Y_a^*, Y_a)$. For $y \in Y_a^*(\mathbb{A}_E)$, $\psi^{-1}\left(\frac{\text{Tr}_{E/F}(cy, y)}{2}\right)$ is the trivial character of $c \in \text{Herm}(Y_a^*, Y_a)$ if and only if $y = 0$. Therefore by (9.5),

$$\int_{\text{Herm}(Y_a^*, Y_a)} \Theta_{W_{m+2a}}((1, c)h, f) dc = \text{meas}(\text{Herm}(Y_a^*, Y_a)) \cdot \theta_1((\nu_{W_{m+2a}}(h)f)(0)).$$

By (9.4), $(\nu_{W_{m+2a}}((1, b)h)(f))(0) = \nu_{W_{m+2a}}(h)(f)(0)$ for all $b \in \text{Hom}(W_m, Y_a)$ and so the claim follows. \square

Remark 9.2. Let e be the identity element of $H_{n+2a,r}(\mathbb{A}_F) = N_{n+2a,r}(\mathbb{A}_F) \times G_{m+2a}(\mathbb{A}_F)$. Observe that

$$\Theta_{P_{m+2a,a}}(e, f) = \theta_1(f(0))$$

and every Schwartz function in \mathcal{S}^0 can be obtained as evaluation at $0 \in Y_a^*(\mathbb{A}_F)$ of some Schwartz function in \mathcal{S} . So we can regard theta functions in ν_{W_m} as the evaluation at e of the constant term of theta functions in $\nu_{W_{m+2a}}$ along $P_{m+2a,a}$. This observation will be used in the proof of Lemma 9.7.

Remark 9.3. From (9.3), we have

$$\Theta_{P_{m+2a,a}}((u, hg), f) = \mu^{-1}(h) |\det(h)|_{\mathbb{A}_E}^{\frac{1}{2}} \cdot \Theta_{P_{m+2a,a}}((u, g), f)$$

for $h \in GL(Y_a(\mathbb{A}_E))$. Thus from the Lemma 9.1 and (9.1), (9.2), we see that the constant terms of theta functions $\{\Theta_{P_{m+2a,a}}(\cdot, f)\}_{f \in \mathcal{S}}$ belong to the induced representation

$$\text{ind}_{GL_a(\mathbb{A}_E) \times (N_{n,r}(\mathbb{A}_F) \times G_m(\mathbb{A}_F))}^{N_{n+2a,r}(\mathbb{A}_F) \times G_{m+2a}(\mathbb{A}_F)} \mu^{-1} \cdot |\cdot|_{\mathbb{A}_E}^{\frac{1}{2}} \boxtimes \nu_{W_m}.$$

The following vanishing lemma plays an essential role in proving Theorem 9.9.

Lemma 9.4. *Let σ be an irreducible cuspidal automorphic representation of $GL_a(\mathbb{A}_E)$, π_1, π_2 irreducible tempered cuspidal automorphic representations of $G_n(\mathbb{A}_F)$ and $G_m(\mathbb{A}_F)$, respectively. Write $BC(\pi_2)$ as an isobaric sum $\sigma_1 \boxplus \cdots \boxplus \sigma_t$, where $\sigma_1, \dots, \sigma_t$ are distinct irreducible cuspidal automorphic representations of the general linear groups such that the (twisted) Asai L -function $L(s, \sigma_i, As^{(-1)^{m-1}})$ has a pole at $s = 1$. Assume that $\sigma \simeq \sigma_i$ for some $1 \leq i \leq t$. Then*

$$\mathcal{P}(E(\phi, z), \Theta_{W_{m+2a}}(f), \varphi) = 0$$

for all $\phi \in \mathcal{A}_{P_{n+2a,a}}^{\mu \cdot \sigma^\vee \boxtimes \pi_1}(G_{n+2a})$, $\varphi \in \mathcal{E}^1(\sigma, \pi_2)$ and $f \in \nu_{W_{m+2a}}$.

Proof. Since $\sigma \simeq \sigma_i$, the residue $\mathcal{E}^1(\sigma, \pi_2)$ is non-zero by Remark 8.3.

By Proposition 6.3 (iii) and Remark 8.1, the linear functional $\mathcal{P}(E(\cdot, z), \Theta_{W_{m+2a}}(\cdot), \cdot)$ belongs to

$$(9.7) \quad \text{Hom}_{N_{n+2a,r}(\mathbb{A}_F) \times G_{m+2a}(\mathbb{A}_F)} \left(\text{Ind}_{P_{n+2a,a}}^{G_{n+2a}} (\mu \cdot |z|_E \cdot \sigma^\vee \boxtimes \pi_1) \otimes \nu_{W_{m+2a}} \otimes \mathcal{E}^1(\sigma, \pi_2), \mathbb{C} \right).$$

Since $z \mapsto \mathcal{P}(E(\phi, z), \Theta_{W_{m+2a}}, \varphi)$ is a meromorphic function on \mathbb{C} by Remark 8.5, if we prove that the Hom space (9.7) is zero for almost all q_E^{-z} , then $\mathcal{P}(E(\phi, z), \Theta_{W_{m+2a}}, \varphi)$ is identically zero. (Note that $G_{m+2a}(\mathbb{A}_F) = G_{m+2a}(\mathbb{A}_F)^1$.)

We first prove the irreducibility of $\mathcal{E}^1(\sigma, \pi_2)$. Since the cuspidal support of the residues in $\mathcal{E}^1(\sigma, \pi_2)$ consists only of $|\sigma| \cdot |z|_E^{-1} \boxtimes \pi_2$, the residues are square integrable by [41, Lemma I.4.11]. Thus $\mathcal{E}^1(\sigma, \pi_2)$ is a unitary quotient of $\text{Ind}_{P_{m+2a,a}(\mathbb{A}_F)}^{G_{m+2a}(\mathbb{A}_F)} (|\sigma| \cdot |z|_E \boxtimes \pi_2)$. Since the Langlands quotient is the unique irreducible quotient, it is isomorphic to $\mathcal{E}^1(\sigma, \pi_2)$.

We fix a finite place v of F which splits in E and the local v -components of $\psi, \mu, \sigma, \pi_1, \pi_2$ and $\mathcal{E}^1(\sigma, \pi_2)$ are unramified. We drop the subscript v from the notation. Note that $G_n(F) \simeq GL_n(F)$ and $GL_a(E) \simeq GL_a(F) \times GL_a(F)$. Since μ is trivial on F^\times , we can write $\mu = (\mu_1^{-1}, \mu_1)$ for some unitary character μ_1 of F^\times . Let B_k be the standard Borel subgroup of GL_k . Recall that an irreducible generic unramified representation of $GL_k(F)$ is an irreducible principal series representation. From now on, we suppress the field F from the notation.

Since $\sigma^c \simeq \sigma^\vee$ by Remark 8.4, we can write $\sigma \simeq \sigma_0 \boxtimes (\sigma_0)^\vee$ where $\sigma_0 \simeq \text{Ind}_{B_a}^{GL_a}(\chi_1 \boxtimes \cdots \boxtimes \chi_a)$, and since σ is an isobaric summand of $BC(\pi_2)$, we can assume

$$\pi_2 \simeq \text{Ind}_{B_m}^{GL_m}(\chi_1 \boxtimes \cdots \boxtimes \chi_a \boxtimes \chi_{a+1} \boxtimes \cdots \boxtimes \chi_m),$$

where χ_i 's are unramified characters of F^\times such that $|\chi_i| = 1$. (Note that π_2 is tempered.)

For a sequence $\mathbf{a} = (a_1, a_2, \dots, a_t)$ of positive integers whose sum is k , we denote by $\mathcal{P}_{\mathbf{a}} = \mathcal{M}_{\mathbf{a}} \mathcal{U}_{\mathbf{a}}$ the standard parabolic subgroup of GL_k whose Levi subgroup $\mathcal{M}_{\mathbf{a}}$ is isomorphic to $GL_{a_1} \times \cdots \times GL_{a_t}$.

Write $\sigma_1 = \text{Ind}_{B_a}^{GL_a}(\mu_1 \chi_1 \boxtimes \cdots \boxtimes \mu_1 \chi_a)$. Note that

$$(9.8) \quad \text{Ind}_{P_{n+2a,a}}^{G_{n+2a}}(\mu \cdot \sigma^\vee | \cdot |_E^z \boxtimes \pi_1) \simeq \text{Ind}_{\mathcal{P}_{(a,a,n)}}^{GL_{n+2a}}(\sigma_1^\vee | \cdot |^z \boxtimes \sigma_1^\vee | \cdot |^{-z} \boxtimes \pi_1),$$

$$(9.9) \quad \text{Ind}_{P_{m+2a,a}}^{G_{m+2a}(F)}(\sigma | \cdot |_E \boxtimes \pi_2) \simeq \text{Ind}_{\mathcal{P}_{(a,m,a)}}^{GL_{m+2a}}(\sigma_0 | \cdot | \boxtimes \pi_2 \boxtimes \sigma_0 | \cdot |^{-1}).$$

Write

$$\Sigma = \text{Ind}_{\mathcal{P}_{(3,\dots,3,1,\dots,1)}}^{GL_{m+2a}}(\chi_1(\det_{GL_3}) \boxtimes \cdots \boxtimes \chi_a(\det_{GL_3}) \boxtimes \chi_{a+1} \boxtimes \cdots \boxtimes \chi_m).$$

Since $|\chi_i| = 1$ for all $1 \leq i \leq m$, Σ is irreducible by [60, Theorem 4.2]. It is not so difficult to show that Σ is an irreducible unramified subquotients of (9.9) and so $\mathcal{E}^1(\sigma, \pi_2) \simeq \Sigma$. Therefore, we are sufficient to prove that

$$(9.10) \quad \text{Hom}_{N_{n+2a,r} \times G_{m+2a}} \left(\text{Ind}_{\mathcal{P}_{(a,n,a)}}^{GL_{n+2a}}(\sigma_1^\vee | \cdot |^z \boxtimes \pi_1 \boxtimes \sigma_1^\vee | \cdot |^{-z}) \otimes \nu_{W_{m+2a}} \otimes \Sigma, \mathbb{C} \right)$$

is zero for almost all $q_E^{-z} = q^{-2z}$. Before proving this, we introduce some notations.

Given a partition $\tilde{l} = (l_1, l_2, l_3)$ of k with $l_1, l_3 \geq 0$ and $l_2 \geq 2$, let

$$\mathcal{N}_{\tilde{l}} = \left\{ v = \begin{pmatrix} z_1 & y_1 & x \\ & I_{l_2} & y_2 \\ & & z_2 \end{pmatrix} \in GL_k(F) \mid z_1 \in \mathcal{N}_{l_1}, z_2 \in \mathcal{N}_{l_3} \right\}$$

and define its character $\psi_{\tilde{l}}$ by

$$\psi_{\tilde{l}}(v) = \psi \left(\sum_{i=2}^{l_1} (z_1)_{i-1,i} \right) \cdot \psi \left(\sum_{i=2}^{l_3} (z_2)_{i-1,i} \right) \cdot \psi((y_1)_{l_1,1} + (y_2)_{l_2,1}).$$

For a smooth representation τ of GL_k and $0 \leq i \leq k$, let $\tau^{(i)}$ (reps. $\tau_{(i)}$) be the representation of GL_{k-i} (resp. $\mathcal{P}_{k-i-1,1}$) defined by some Jacquet module of τ as in [22, p.87]. (Note that $\tau^{(i)}$ is nothing but the i -th Bernstein-Zelevinsky derivative of τ if we use the unnormalized Jacquet functors in the definition of Bernstein-Zelevinsky derivative as in [6].)

Let w_k be the $k \times k$ matrix, which has 1 along the main anti-diagonal, and zeroes elsewhere. Denote by τ^* , the representation of GL_k , acting in the space of τ , by $\tau^*(g) = \tau(g^*)$, where $g^* = w_k \cdot {}^t g^{-1} \cdot w_k^{-1}$.

Put

$$\tau^{[i]} := [(\tau^*)^{(i)}]^*, \quad \tau_{[i]} := [(\tau^*)_{(i)}]^*.$$

For a character χ of F^\times , the Bernstein-Zelevinski derivative of $\chi(k) := \chi(\det_{GL_k})$ is given by

$$(\chi(k))^{(t)} = \begin{cases} \chi(k-t) & \text{if } t \leq 1, \\ 0 & \text{if } t \geq 2 \end{cases}$$

and

$$(\chi(k))_{(t)} = 0, \quad \text{for } t \geq 1.$$

For a partition (m_1, \dots, m_t) of k and characters μ_i 's, let $\tau = \text{Ind}_{\mathcal{P}_{(m_1, \dots, m_t)}}^{GL_k} \mu_1(m_1) \otimes \cdots \otimes \mu_t(m_t)$. Using the Leibniz rule of the Bernstein-Zelevinsky derivative, it is easy to see that for every non-negative integer j ,

$$(9.11) \quad | \cdot |^{\frac{1-j}{2}} \cdot \tau^{(j)} \simeq \bigoplus_{\substack{i_1 + \cdots + i_t = j \\ 0 \leq i_1, \dots, i_t \leq 1}} \text{Ind}_{\mathcal{P}_{(m_1-i_1, \dots, m_t-i_t)}}^{GL_{k-j}} | \cdot |^{\frac{1-i_1}{2}} \cdot \mu_1(m_1 - i_1) \otimes \cdots \otimes | \cdot |^{\frac{1-i_t}{2}} \cdot \mu_t(m_t - i_t)$$

and

$$(9.12) \quad |\cdot|^{-\frac{j-1}{2}} \cdot \tau[j] \simeq \bigoplus_{\substack{i_1+\dots+i_t=j \\ 0 \leq i_1, \dots, i_t \leq 1}} \text{Ind}_{\mathcal{P}_{(m_1-i_1, \dots, m_t-i_t)}}^{GL_{k-j}} |\cdot|^{-\frac{i_1-1}{2}} \cdot \mu_1(m_1-i_1) \otimes \dots \otimes |\cdot|^{-\frac{i_t-1}{2}} \cdot \mu_t(m_t-i_t)$$

up to semisimplification. (See [15, (5.34)])

To prove that (9.10) is zero for almost all q^{-2z} , we divide the cases $r > 0$ and $r = 0$ separately.

We first deal with the case $r > 0$.

Let $\tilde{r} = (r-1, m+2a+2, r-1)$ be a partition of $n+2a$. Then by Lemma 7.1,

$$(9.10) \simeq \text{Hom}_{GL_{m+2a}} \left(J_{\psi_{\tilde{r}}}^{-1} (\text{Ind}_{\mathcal{P}_{(a,a,n)}}^{GL_{n+2a}} (\sigma_1^\vee |\cdot|^z \boxtimes \sigma_1^\vee |\cdot|^{-z} \boxtimes \pi_1) \otimes \Omega_{W_{m+2a}}), \Sigma^\vee \right).$$

Put $\tau_{1,z} = \text{Ind}_{\mathcal{P}_{(a,a)}}^{GL_{2a}} (\sigma_1^\vee |\cdot|^z \boxtimes \sigma_1^\vee |\cdot|^{-z})$. By applying [22, Theorem 6.5] and [22, Proposition 6.11] with $\tau_1 = \tau_{1,z}$ and $\tau_2 = \pi_1$, we can decompose $J_{\psi_{\tilde{r}}}^{-1} (\text{Ind}_{\mathcal{P}_{(2a,n)}}^{GL_{n+2a}} (\tau_{1,z} \otimes \pi_1) \otimes \Omega_{W_{m+2a}})$ as

(9.13)

$$\bigoplus_{\max\{0, 2a-(r-1)\} \leq s \leq 2a} \text{Ind}_{\mathcal{P}_{(s, m+2a-s)}}^{GL_{m+2a}} \left(\mu_1 |\cdot|^{\frac{1-(2a-s)}{2}} (\tau_{1,z})^{(2a-s)} \boxtimes |\cdot|^{\frac{2a-s}{2}} J_{\psi_{(r-1, m+2a+2-s, r-1-2a+s)}}^{-1} (\pi_1 \otimes \Omega_{W_{m+2a-s}}) \right)$$

(9.14)

$$\bigoplus_{0 \leq s < \max\{2a-r, r-1\}} \text{Ind}_{\mathcal{P}_{(2a-r-s-1, m+r+1+s)}}^{GL_{m+2a}} \left(|\cdot|^{\frac{s-r+1}{2}} J_{\psi_{(s, 2a-r-s+1, r-1)}}^{-1} (\tau_{1,z} \otimes \Omega_{W_{2a-r-s-1}}) \boxtimes \mu_1 |\cdot|^{\frac{r-s-2}{2}} (\pi_1)^{[r-s-1]} \right)$$

(9.15)

$$\bigoplus \begin{cases} \text{Ind}_{\mathcal{P}_{(2a-r, n-r)}}^{GL_{m+2a}} \left(\mu_1 \otimes \left(|\cdot|^{\frac{1-r}{2}} (\tau_{1,z})^{(r)} \boxtimes |\cdot|^{\frac{r-1}{2}} (\pi_1)^{[r]} \right) \right), & \text{if } r < 2a+1 \\ 0, & \text{otherwise,} \end{cases}$$

up to semisimplification.

Since π_1 is tempered, by (9.12) and [8, Theorem 2.9], (9.14) and (9.15) have no common composition factor with Σ^\vee for almost all q^{-2z} . Therefore, we are enough to show that $\text{Hom}_{GL_{m+2a}} ((9.13), \Sigma^\vee) = 0$ for almost all q^{-2z} .

Suppose $\text{Hom}_{GL_{m+2a}} ((9.13), \Sigma^\vee) \neq 0$ for infinitely many q^{-2z} . Then there exists an index s_0 whose corresponding summand in (9.13) has common composition factor with Σ^\vee for infinitely many q^{-2z} . Write $\pi_1 \simeq \text{Ind}_{B_n}^{GL_n} (\chi'_1 \boxtimes \dots \boxtimes \chi'_n)$, where χ'_i 's are unramified characters of F^\times such that $|\chi'_i| = 1$.

For $1 \leq i \leq n-1$, put $\pi_{1,i} \simeq \text{Ind}_{B_{n-i}}^{GL_{n-i}} (\chi'_{i+1} \boxtimes \dots \boxtimes \chi'_n)$ and $\tilde{r}_{s_0} = (r-1, m+2a-s_0+2, r-1-2a+s_0)$. By applying [22, Theorem 6.5] and [22, Proposition 6.11] with $\tilde{l} = \tilde{r}_{s_0}, j = 1, \tau_1 = \chi'_1$ and $\tau_2 = \pi_{1,1}$, we can decompose $|\cdot|^{\frac{2a-s_0}{2}} J_{\psi_{\tilde{r}_{s_0}}}^{-1} (\pi_1 \otimes \Omega_{W_{m+2a-s_0}})$ as

(9.16)

$$\bigoplus_{t=0,1} \text{Ind}_{\mathcal{P}_{(t, m+2a-s_0-t)}}^{GL_{m+2a-s_0}} \left(\mu_1 |\cdot|^{\frac{t}{2}} (\chi'_1)^{(1-t)} \boxtimes |\cdot|^{\frac{2a-s_0+(1-t)}{2}} J_{\psi_{(r-1, m+2a-s_0+2-t, r-1-2a+s_0-1+t)}}^{-1} (\pi_{1,1} \otimes \Omega_{W_{m+2a-s_0-t}}) \right)$$

(9.17)

$$\bigoplus \begin{cases} \mu_1 |\cdot|^{\frac{r-1}{2}} (\pi_{1,1})^{[r]}, & \text{if } r-1-2a+s_0 = 0 \\ 0, & \text{otherwise,} \end{cases}$$

up to semisimplification. (Note that the second term in [22, (6.30)] does not contribute.)

Since $\pi_{1,1}$ is tempered, by (9.12) and [8, Theorem 2.9],

$$\mathrm{Hom}_{GL_{m+2a}} \left(\mathrm{Ind}_{\mathcal{P}_{(s_0, m+2a-s_0)}}^{GL_{m+2a}} \left(\mu_1 | \cdot |^{\frac{1-(2a-s_0)}{2}} (\tau_{1,z})^{(2a-s_0)} \boxtimes (9.17) \right), \Sigma^\vee \right) = 0$$

and so

$$\mathrm{Hom}_{GL_{m+2a}} \left(\mathrm{Ind}_{\mathcal{P}_{(s_0, m+2a-s_0)}}^{GL_{m+2a}} \left(\mu_1 | \cdot |^{\frac{1-(2a-s_0)}{2}} (\tau_{1,z})^{(2a-s_0)} \boxtimes (9.16) \right), \Sigma^\vee \right)$$

is zero for infinitely many q^{-2z} . Following the similar argument in the proof of [22, Theorem 6.7], if we continue this procedure repeatedly, we see that there exist i -tuple integers $\bar{t} = (t_1, \dots, t_i)$ such that

$$\mathrm{Hom}_{GL_{m+2a}} \left(\mathrm{Ind}_{\mathcal{P}_{(s_0, m+2a-s_0)}}^{GL_{m+2a}} \left(\mu_1 | \cdot |^{\frac{1-(2a-s_0)}{2}} (\tau_{1,z})^{(2a-s_0)} \boxtimes \pi_{\bar{t}} \right), \Sigma^\vee \right) \neq 0$$

for infinitely many q^{-2z} .

(Here, $\pi_{\bar{t}}$ is the representation of GL_{m+2a-s_0} parabolically induced from the representation of

$$\mu_1 \cdot \left(| \cdot |^{\frac{t_1}{2}} (\chi'_1)^{(1-t_1)} \boxtimes \dots \boxtimes | \cdot |^{\frac{t_i}{2}} (\chi'_i)^{(1-t_i)} \boxtimes | \cdot |^{\frac{r-1}{2}} J_{\psi_{(r-1, m+2a-s_0+2-\sum_{k=1}^i t_k, 0)}}^{-1}} \left(\pi_{1, i+1} \otimes \Omega_{W_{m+2a-s_0-\sum_{k=1}^i t_k}} \right) \right).$$

Applying the similar computation in the proof of [22, Theorem 6.8] (see also [22, Theorem 6.7]), we can decompose $| \cdot |^{\frac{r-1}{2}} J_{\psi_{(r-1, m+2a-s_0+2-\sum_{k=1}^i t_k, 0)}}^{-1}} \left(\pi_{1, i+1} \otimes \Omega_{W_{m+2a-s_0-\sum_{k=1}^i t_k}} \right)$ as

$$\bigoplus_{j \geq 0} \mathrm{Ind}_{\mathcal{P}_{(1, \dots, 1, m+2a-s_0-\sum_{k=1}^i t_k-j)}}^{GL_{m+2a-s_0-\sum_{k=1}^i t_k}} \left(1 \boxtimes \dots \boxtimes 1 \boxtimes | \cdot |^{\frac{z_j-1}{2}} (\pi_{1, i+1})^{[z_j]} \right),$$

up to semisimplification. (Here, the exponents z_j depend on j, t_1, \dots, t_i .)

Since $\mathrm{Hom}_{GL_{m+2a}} \left((9.13), \Sigma^\vee \right) \neq 0$ for infinitely many q^{-2z} , there exists some index j_0 such that the representation of GL_{m+2a} parabolically induced from

$$\mu_1 \cdot \left(| \cdot |^{\frac{1-(2a-s_0)}{2}} (\tau_{1,z})^{(2a-s_0)} \boxtimes | \cdot |^{\frac{t_1}{2}} (\chi'_1)^{(1-t_1)} \boxtimes \dots \boxtimes | \cdot |^{\frac{t_i}{2}} (\chi'_i)^{(1-t_i)} \boxtimes 1 \boxtimes \dots \boxtimes 1 \boxtimes | \cdot |^{\frac{z_{j_0}-1}{2}} (\pi_{1, i+1})^{[z_{j_0}]} \right)$$

has common composition factor with Σ^\vee for infinitely many q^{-2z} . However, by (9.11) and (9.12), it is impossible because π_1, π_2 are tempered. Therefore, $\mathrm{Hom}_{GL_{m+2a}} \left((9.13), \Sigma^\vee \right) = 0$ for almost all q^{-2z} and we proved our claim when $r > 0$.

Next we shall prove that (9.10) is zero for almost all q^{-2z} when $r = 0$.

Write $\tilde{a} = (a-1, m+2, a-1)$. By Lemma 7.1, we are enough to show that

$$\mathrm{Hom}_{GL_m} \left(J_{\psi_{\tilde{a}}}^{-1} \left(\mathrm{Ind}_{\mathcal{P}_{(3, m+2a-3)}}^{GL_{m+2a}} (\chi \circ \det_{GL_3} \boxtimes \tau) \otimes \Omega_{W_m} \right) \otimes \pi_1, \mathbb{C} \right) = 0$$

for a character χ of F^\times and a smooth representation τ of GL_{m+2a-3} .

We apply [22, Theorem 6.5] and [22, Proposition 6.11] with $\tilde{l} = \tilde{a}, j = 3, \tau_1 = \chi(3)$ and $\tau_2 = \tau$ to compute $J_{\psi_{\tilde{a}}}^{-1} \left(\mathrm{Ind}_{\mathcal{P}_{(3, m+2a-3)}}^{GL_{m+2a}} (\chi(3) \boxtimes \tau) \otimes \Omega_{W_m} \right)$. Since the Jacquet modules of $\chi(3)$ with respect to non-trivial characters $\psi_{(1,2,0)}, \psi_{(0,2,1)}$ and $\psi_{(0,3,0)}$ are zero, the second term in [22, (6.30)] does not contribute. Thus $J_{\psi_{\tilde{a}}}^{-1} \left(\mathrm{Ind}_{\mathcal{P}_{(3, m+2a-3)}}^{GL_{m+2a}} (\chi(3) \boxtimes \tau) \otimes \Omega_{W_m} \right)$ is decomposed as

$$(9.18) \quad \bigoplus_{2 \leq s \leq 3} \mathrm{Ind}_{\mathcal{P}_{(s, m-s)}}^{GL_m} (\rho_{1,s} \boxtimes \rho_{2,s})$$

up to semisimplification. (Here, $\rho_{1,s}$ is some 1-dimensional representation of GL_s and $\rho_{2,s}$ is some smooth representation of GL_{m-s} .)

By the Frobenius reciprocity,

$$(9.19) \quad \mathrm{Hom}_{GL_m} \left(\mathrm{Ind}_{\mathcal{P}_{s,m-s}}^{GL_m} (\rho_{1,s} \boxtimes \rho_{2,s}) \otimes \pi_1, \mathbb{C} \right) \simeq \mathrm{Hom}_{GL_s \times GL_{m-s}} \left((\rho_{1,s} \boxtimes \rho_{2,s}) \otimes J_{\mathcal{P}_{s,m-s}}(\pi_1), \mathbb{C} \right),$$

where $J_{\mathcal{P}_{s,m-s}}$ is the Jacquet functor with respect to $\mathcal{P}_{s,m+2a-s}$. Since π_1 is the irreducible principal series, by [60, Theorem 1.2], every irreducible subquotient of $J_{\mathcal{P}_{s,m-s}}(\pi_1)$ should be a tensor product of irreducible principal series representation of GL_s and that of GL_{m-s} . However, since $\rho_{1,s}$ is one-dimensional, (9.19) is zero. This proves our claim when $r = 0$. \square

From now on, we simply write P (resp. P_a, M_a, K_a) for $P_{n+2a,a}$ (resp. $P_{m+2a,a}, M_{m+2a,a}, K_{m+2a}$). Note that $N_{n+2a,r} \simeq (\mathrm{Hom}(Y_a, X) \times \mathrm{Hom}(Y_a^*, X)) \rtimes N_{n,r}$ and $N_{n+2a,r} \cap P = \mathrm{Hom}(Y_a^*, X) \rtimes N_{n,r}$. Denote $\mathrm{Hom}(Y_a, X) \times \mathrm{Hom}(Y_a^*, X)$ by V and let dv be the Haar measure of V such that $dn' = dvdn$, where dn' (resp. dn) is the Haar measures of $N_{n+2a,r}$ (resp. $N_{n,r}$). The following remark is an easy observation which will be used in the proof of Lemma 9.6,

Remark 9.5. For $x \in M_{m+2a,a}$, $xVx^{-1} = V$ and the Jacobian $\frac{d(xvx^{-1})}{dv} = 1$.

Lemma 9.6. *With the same notation as in Lemma 9.4, assume $\sigma \simeq \sigma_i$ for some $1 \leq i \leq t$. If $\phi \in \mathcal{A}_P^{\mu \cdot \sigma^\vee \boxtimes \pi_1}(G_{n+2a})$, $\varphi \in \mathcal{E}^1(\sigma, \pi_2)$ and $f \in \nu_{W_{m+2a}}$, then*

$$\begin{aligned} & \mathcal{FJ}_{\psi,\mu}(\mathcal{E}^0(\phi), \varphi, f) = \\ & \int_{K_a} \int_{V(\mathbb{A}_F)} \left(\int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \left(\int_{[N_{n,r}]} \phi(nmvk) \cdot \Theta_{P_a}(f)((n, m) \cdot (v, k)) dn \right) \varphi_{P_a}(mk) dm \right) \cdot e^{\frac{H_P(vk)}{2}} dvdk. \end{aligned}$$

Proof. The proof is similar with [58, Lemma 6.5]. Using the residue theorem, write

$$\mathcal{E}^0(\phi)(g) = \frac{1}{2\pi i} \int_{|z - \frac{1}{2}| = \epsilon} E(g, \phi, z) dz$$

for some small $\epsilon > 0$. If we apply the similar argument introduced by Arthur in [3, pp. 47-48], the complex line integral can be interchanged with the mixed truncation operator. Then by Fubini's theorem, we have

$$(9.20) \quad \begin{aligned} & \int_{G_{m+2a}(F) \backslash G_{m+2a}(\mathbb{A}_F)} \Lambda_m^T(\mathcal{E}^0(\phi) \otimes \Theta_{W_{m+2a}}(f))(g) \cdot \varphi(g) dg \\ &= \frac{1}{2\pi i} \cdot \int_{|z - \frac{1}{2}| = \epsilon} \int_{G_{m+2a}(F) \backslash G_{m+2a}(\mathbb{A}_F)} \Lambda_m^T(E(\phi, z) \otimes \Theta_{W_{m+2a}}(f))(g) \cdot \varphi(g) dg dz \\ &= \lim_{z \rightarrow \frac{1}{2}} \left(z - \frac{1}{2} \right) \cdot \int_{G_{m+2a}(F) \backslash G_{m+2a}(\mathbb{A}_F)} \Lambda_m^T(E(\phi, z) \otimes \Theta_{W_{m+2a}}(f))(g) \cdot \varphi(g) dg. \end{aligned}$$

Since $\mathcal{FJ}_{\psi,\mu}(\mathcal{E}^0(\phi), \varphi, f)$ is the zero coefficient of the left-hand side, we compute the zero coefficient of the right-hand side. By Proposition 6.4 and Lemma 9.4,

$$\begin{aligned} & \int_{G_{m+2a}(F) \backslash G_{m+2a}(\mathbb{A}_F)} \Lambda_m^T(E(\phi, z) \otimes \Theta_{W_{m+2a}}(f))(g) \cdot \varphi(g) dg \\ &= - \int_{P_a(F) \backslash G_{m+2a}(\mathbb{A}_F)}^* (E(\phi, z) \otimes \Theta_{W_{m+2a}}(f))_{P_a, N_{n+2a,r}} \cdot \varphi_{P_a}(g) \cdot \hat{\tau}_{P_a}(H_{P_a}(g) - T) dg. \end{aligned}$$

Note that $E_P(g, \phi, z) = \phi(g)d(g)^z + (M(z)\phi)(g)d(g)^{-z}$, where $M(z)$ is the relevant intertwining operator. Therefore,

$$\begin{aligned} & \int_{[N_{n+2a,r}]} E_P(\phi, z)(lg) \cdot \Theta_{P_a}(f)(l, g) dl \\ &= \int_{[N_{n+2a,r}]} \phi(lg) \cdot \Theta_{P_a}(f)(l, g) \cdot d(lg)^z dl + \int_{[N_{n+2a,r}]} (M(z)\phi)(lg) \cdot \Theta_{P_a}(f)(l, g) \cdot d(lg)^{-z} dl. \end{aligned}$$

Since the second term does not contribute to the zero coefficient in (9.20), we only consider the first term.

Then by Lemma 9.1 and Remark 9.5,

$$\begin{aligned} & - \int_{P_a(F) \backslash G_{m+2a}(\mathbb{A}_F)}^* \left(\int_{[N_{n+2a,r}]} \phi(lg) \cdot \Theta_{P_a}(f)(l, g) \cdot d(lg)^z dl \right) \cdot \varphi_{P_a}(g) \cdot \hat{\tau}_{P_a}(H_{P_a}(g) - T) dg \\ &= - \int_{K_a} \int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \int_T \int_{[V]} \int_{[N_{n,r}]} \phi(nvmk) \cdot \Theta_{P_a}(f)(nv, mk) e^{z(X+H_P(vmk))} \varphi_{P_a}(mk) \cdot e^{-\frac{X}{2}} dndvdXdmdk \\ &= - \int_T e^{(z-\frac{1}{2})X} dX \cdot \int_{K_a} \int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \int_{[V]} \int_{[N_{n,r}]} \phi(nmvk) \cdot \Theta_{P_a}(f)((n, m)(v, k)) \varphi_{P_a}(mk) e^{zH_P(vk)} dndvdmdk. \end{aligned}$$

Since

$$- \int_T e^{(z-\frac{1}{2})X} dX = \frac{e^{(z-\frac{1}{2})T}}{z - \frac{1}{2}},$$

the zero coefficient of

$$\lim_{z \rightarrow \frac{1}{2}} \left(z - \frac{1}{2} \right) \cdot \int_{G_{m+2a}(F) \backslash G_{m+2a}(\mathbb{A}_F)} \Lambda_m^T(E(g, \phi, z) \otimes \Theta_{W_{m+2a}}(f))(g) \cdot \varphi(g) dg$$

is equal to

$$\begin{aligned} & \lim_{z \rightarrow \frac{1}{2}} \left(z - \frac{1}{2} \right) \cdot \mathcal{P}(E(\phi, z), \Theta_{W_{m+2a}}(f), \varphi) + \\ & \int_{K_a} \int_{[V]} \left(\int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \left(\int_{[N_{n,r}]} \phi(nmvk) \cdot \Theta_{P_a}(f)((n, m) \cdot (v, k)) dn \right) \varphi_{P_a}(mk) dm \right) \cdot e^{\frac{H_P(vk)}{2}} dvdk. \end{aligned}$$

Since $\mathcal{P}(E(\phi, z), \Theta_{W_{m+2a}}(f), \varphi) = 0$ by Lemma 9.4, it completes the proof. \square

Lemma 9.7. *With the same notation as in Lemma 9.4, we assume $\sigma \simeq \sigma_i$ for some $1 \leq i \leq t$. Then the following conditions are equivalent.*

- (i) *The Fourier–Jacobi period functional $\mathcal{FJ}_{\psi, \mu}(\pi_1, \pi_2, \nu_{W_m})$ is nonzero*
- (ii) *There exists $\phi \in \mathcal{A}_{P_a}^{\mu, \sigma^\vee \boxtimes \pi_1}(G_{m+2a})$, $\varphi \in \mathcal{E}^1(\sigma, \pi_2)$ and $f \in \nu_{W_{m+2a}}$ such that*

(9.21)

$$\int_{K_a} \int_{[V]} \left(\int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \left(\int_{[N_{n,r}]} \phi(nmvk) \cdot \Theta_{P_a}(f)((n, m) \cdot (v, k)) dn \right) \varphi_{P_a}(mk) dm \right) \cdot e^{\frac{H_P(vk)}{2}} \neq 0$$

Remark 9.8. The above lemma seems similar with [15, Proposition 5.3] appearing in the proof of the one direction of the global GGP for metaplectic–symplectic case. Experts found that there is a technical gap in the proof of [15, Proposition 5.3] when the co-rank is not zero. To be more precise, they confused that the image of a certain residual intertwining operator would be the full induced representation and so they chose a vector in the images of the operator whose support is good. However, since the image of such operator is not the full induced representation, it is highly non-trivial to choose such a vector in

the images of the operator. To circumvent such difficulty, we used the isobaric sum of $BC(\pi_2)$ instead of $BC(\pi_1)$ in Lemma 9.4. It has one advantage and one disadvantage. The advantage is that it does not require to choose a special vector in the image of residual intertwining operator in the proof of Lemma 9.7. The disadvantage is that comparing with the approach using the isobaric sum of $BC(\pi_1)$, the proof of Lemma 9.4 becomes hard because one needs to compute explicitly the relevant twisted Jacquet-modules corresponding to Fourier-Jacobi characters not only for its first occurrence but almost all occurrences. After having proved Lemma 9.4 using the Leibniz rule of these Jacquet-modules, we can prove the following lemma without too much difficulty using the inductive structure of the constant terms of theta series.

Proof. The proof of the (i) \rightarrow (ii) direction basically follows the arguments in the proof of [58, Lemma 6.5]. However, our situation is a bit complicated because it includes an integration on V . Therefore, a more careful treatment is required to control such an additional integral.

Put

$$\Pi = \mu| \cdot |_{\mathbb{E}}^{\frac{1}{2}} \cdot \sigma^\vee \boxtimes \pi_1, \quad \Pi' = \sigma| \cdot |_{\mathbb{E}}^{-1} \boxtimes \pi_2, \quad \Pi'' = \mu^{-1}| \cdot |_{\mathbb{E}}^{\frac{1}{2}} \boxtimes \nu_{\psi^{-1}, \mu^{-1}, W_m}.$$

We define a functional on $\Pi \boxtimes \Pi' \boxtimes \Pi''$ by

$$l(\eta \boxtimes \eta' \boxtimes \eta'') = \int_{M_a(F) \backslash M_a(\mathbb{A}_F)^1} \left(\int_{[N_{n,r}]} \eta(nm) \eta'((n, m)) \eta''(m) dn \right) \eta'(m) dm.$$

Let $(\Pi \boxtimes \Pi' \boxtimes \Pi'')^\infty$ be the canonical Casselman-Wallach globalization of $\Pi \boxtimes \Pi' \boxtimes \Pi''$ realized in the space of smooth automorphic forms without the $K_{M_{n+2a}} \times K_{M_{m+2a}} \times K_{M_{m+2a}}$ -finiteness condition, where $K_{M_{i+2a}} = K_{i+2a} \cap M_{i+2a,a}$ for $i = n, m$ (cf. [10], [53, Chapter 11]). Since cusp forms are bounded, l can be uniquely extended to a continuous functional on $(\Pi \boxtimes \Pi' \boxtimes \Pi'')^\infty$ and denote it by the same notation. Our assumption enables us to choose $\eta \in \Pi, \eta' \in \Pi'$ and $\eta'' \in \Pi''$ so that $l(\eta \boxtimes \eta' \boxtimes \eta'') \neq 0$. We may assume that η, η' and η'' are pure tensors. By [45, 52], the functional l is a product of local functionals $l_v \in \text{Hom}_{M_{a,v}}((\Pi_v \boxtimes \Pi'_v \boxtimes \Pi''_v)^\infty, \mathbb{C})$, where we set $(\Pi_v \boxtimes \Pi'_v \boxtimes \Pi''_v)^\infty = \Pi_v \boxtimes \Pi'_v \boxtimes \Pi''_v$ if v is finite. Then we have $l_v(\eta_v \boxtimes \eta'_v \boxtimes \eta''_v) \neq 0$.

Denote by 1 (resp. e) the identity element of G_{n+2a} (resp. G_{m+2a}). Choose $\varphi' \in \mathcal{E}^1(\sigma, \pi_2)$ and $f \in \nu_{W_{m+2a}}$ such that

- (i) $\delta_{P_a}^{-\frac{1}{2}} \cdot \varphi'_{P_a} = \boxtimes_v \varphi'_v, \Theta_{P_a}^\psi(f) = \boxtimes_v f_v;$
- (ii) $\varphi'_v(e) = \eta_v, f_v((1, e)) = \eta''_v.$

(We can choose such f by Remark 9.2 and Remark 9.3.)

Let $U_{n+2a,a}^-$ be the unipotent radical of the opposite parabolic subgroup P^- of P and α_v be a smooth function on $U_{n+2a,a,v}^-$, which has a compact support. We can define a section ϕ_v by requiring

$$\phi_v(muu_-) = \delta_{P_v}(m)^{\frac{1}{2}} \alpha_v(u_-) \cdot (\Pi_v)^\infty(m) \eta_v, \quad m \in M_{n+2a,a,v}, u_- \in U_{n+2a,a,v}^-.$$

Since $N_{n+2a,r} \simeq (\text{Hom}(Y_a, X) \times \text{Hom}(Y_a^*, X)) \times N_{n,r}$, we regard elements of $\text{Hom}(Y_a, X)$ and $\text{Hom}(Y_a^*, X)$ as elements of $N_{n+2a,r}$ via the natural inclusion maps. By Remark 9.2 and (9.6), for every $p_1 \in \text{Hom}(Y_a, X)_v$,

$$f_v((p_1, e)) = f_v((1, e)) = \eta''_v.$$

Using the Iwasawa decomposition of $G_{n+2a,v}$ with respect to P_v , it is easy to check that $\text{Hom}(Y_a, X)_v$ is contained in $U_{n+2a,a,v}^-$. By taking the support of α_v sufficiently small near e , we have

$$\int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1) \boxtimes \eta'_v \boxtimes f_v((p_1, e))) \cdot e^{\frac{H_P(p_1)}{2}} dp_1 = \int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1) \boxtimes \eta'_v \boxtimes \eta''_v) \cdot e^{\frac{H_P(p_1)}{2}} dp_1 \neq 0.$$

On the other hand, there is a small neighborhood N_v of 0 in $\text{Hom}(Y_a^*, X)_v$ such that

$$\int_{\text{Hom}(Y_a^*, X)_v} \int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1 p_2) \boxtimes \eta'_v \boxtimes f_v((p_1 p_2, e))) \cdot e^{\frac{H_P(p_1 p_2)}{2}} \cdot \chi_{N_v}(p_2) dp_1 dp_2 \neq 0.$$

(Here, χ_{N_v} is the characteristic function on N_v .)

By taking the support of the Schwartz function f sufficiently small neighborhood of $0 \in Y_a^*$, we may assume that the support of f_v is contained in N_v .

Then

$$(9.22) \quad \begin{aligned} & \int_{\text{Hom}(Y_a^*, X)_v} \int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1 p_2) \boxtimes \eta'_v \boxtimes f_v((p_1 p_2, e))) \cdot e^{\frac{H_P(p_1 p_2)}{2}} dp_1 dp_2 \\ &= \int_{\text{Hom}(Y_a^*, X)_v} \int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1 p_2) \boxtimes \eta'_v \boxtimes f_v((p_1 p_2, e))) \cdot e^{\frac{H_P(p_1 p_2)}{2}} \cdot \chi_{N_v}(p_2) dp_1 dp_2, \end{aligned}$$

and so (9.22) is nonzero.

Put

$$I(\phi_v) = \int_{K_{a,v}} \int_{\text{Hom}(Y_a^*, X)_v} \int_{\text{Hom}(Y_a, X)_v} l_v(\phi_v(p_1 p_2 k) \boxtimes \varphi_v(k) \boxtimes f_v((p_1 p_2, k))) \cdot e^{\frac{H_P(p_1 p_2 k)}{2}} dp_1 dp_2 dk.$$

Since $P_{a,v} \cdot U_{a,v}^-$ is an open dense subset of $G_{m+2a,v}$, we can rewrite the local integral as

$$I(\phi_v) = \int_{U_{a,v}^-} \int_{\text{Hom}(Y_a^*, X)_v} \int_{\text{Hom}(Y_a, X)_v} \alpha_v(u_-) l_v(\phi_v(p_1 p_2 u_-) \boxtimes \eta'_v \boxtimes f_v((p_1 p_2, u_-))) \cdot e^{\frac{H_P(p_1 p_2 u_-)}{2}} dp_1 dp_2 du_-.$$

We can choose α_v to be supported in a small neighborhood of e so that $I(\phi_v) \neq 0$. Since I is continuous and $K_{n+2a,v}$ -finite vectors are dense in the induced representation, we can choose $K_{n+2a,v}$ -finite function ϕ_v such that $I(\phi_v) \neq 0$. This completes the proof of the (i) \rightarrow (ii) direction.

The proof of (ii) \rightarrow (i) direction is almost immediate. From what we have seen in the above, we see that the Fourier–Jacobi period integral $\mathcal{FJ}_{\psi,\mu}(\pi_1, \pi_2, \nu_{W_m})$ is just the inner period integral in (9.21). Therefore, if $\mathcal{FJ}_{\psi,\mu}(\pi_1, \pi_2, \nu_{W_m}) = 0$, the integral (9.21) is always zero. \square

By combining Lemma 9.6 with Lemma 9.7, we get the following reciprocal non-vanishing theorem.

Theorem 9.9. *With the same notation as in Lemma 9.4, assume $\sigma \simeq \sigma_i$ for some $1 \leq i \leq t$. Then $\mathcal{FJ}_{\psi,\mu}(\pi_1, \pi_2, \nu_{W_m}) \neq 0$ is equivalent to $\mathcal{FJ}_{\psi,\mu}(\mathcal{E}^0(\mu \cdot \sigma^\vee, \pi_1), \mathcal{E}^1(\sigma, \pi_2), \nu_{W_{m+2a}}) \neq 0$.*

10. Proof of the Main Theorem

Proof. Since π_2 is tempered, $BC(\pi_2)$ is an isobaric sum of the form $\sigma_1 \boxplus \cdots \boxplus \sigma_t$, where $\sigma_1, \dots, \sigma_t$ are distinct irreducible cuspidal automorphic representations of general linear groups such that the (twisted) Asai L -function $L(s, \sigma_i, As^{(-1)^{m-1}})$ has a pole at $s = 1$. Then for each $1 \leq i \leq t$, $L(s, \mu^{-1} \cdot \sigma_i, As^{(-1)^m})$ has a pole at $s = 1$. On the other hand, $\mathcal{E}^0(\mu \cdot \sigma_i^\vee, \pi_1)$ is nonzero by Theorem 9.9. Thus by Proposition 8.2, we have $L(\frac{1}{2}, BC(\pi_1^\vee) \times \mu \cdot \sigma_i^\vee) \neq 0$ and so $L(\frac{1}{2}, BC(\pi_1) \times \mu^{-1} \sigma_i) \neq 0$ by the functional equation. Thus

$$L\left(\frac{1}{2}, BC(\pi_1) \times BC(\pi_2) \otimes \mu^{-1}\right) = \prod_{i=1}^t L\left(\frac{1}{2}, BC(\pi_1) \times \mu^{-1} \sigma_i\right) \neq 0$$

and so our claim is proved. \square

Acknowledgements. The author would like to express his deepest gratitude to Prof. Hiraku Atobe for his suggestion this nice problem and many helpful discussion. Without him, this work would not see the light of the world. The definition of automorphic forms on non-reductive groups in Section 3 is due to him. He also found critical errors on the proof of Lemma 7.1 and Lemma 9.4 in our earlier draft. Taking this opportunity, we highly appreciate him for his significant contribution on this paper.

We are also much grateful to Prof. Shunsuke Yamana. The huge influence of his paper [58] is evident to the reader. Besides being the author of the beautiful paper which inspired this work, he kindly answered the author's many questions on his paper. We also would like to thank to Prof. Erez Lapid for his explanation on some part of his proof about [43, Lemma 8.2.1(1)]. We express our sincere thanks to them for their endless kindness they have shown to us. We would like to thank Prof. Dipendra Prasad and Prof. Sug Woo Shin for many helpful comments on our draft.

Thanks are also to Prof. Yeansu Kim and Prof. Sanghoon Kwon. As running 5-months reading seminar on 'Arthur trace formula' with them, the author could get familiar with the Arthur truncation and gain some ideas how to prove various lemmas in Section 3. He also would like to thank to Dr. Wan Lee and Dr. Chulhee Lee for useful discussion.

We sincerely thank to anonymous referee for their careful reading and helpful comments, which improved the exposition of the paper.

Most part of this paper was written when the author was a postdoc at KIAS. We thank KIAS and Yonsei university for their generous supports. The author was supported by KIAS Individual Grant(SP037002) at Korea Institute for Advanced Study, the POSCO Science Fellowship of POSCO TJ Park Foundation and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2020R1F1A1A01048645).

REFERENCES

- [1] J. Arthur, *A trace formula for reductive groups, I: Terms associated to classes in $G(\mathbb{Q})$* , Duke Math. J., **45** No. 1. (1978), 911–952.
- [2] J. Arthur, *A trace formula for reductive groups. II: Applications of a truncation operator*, Compositio Math. **40** (1980), 87–121.
- [3] J. Arthur, *On the inner product of truncated Eisenstein series*, Duke Math. J. **49** (1982), 35–70.
- [4] J. Arthur, *The endoscopic classification of representations: orthogonal and symplectic groups*, Colloquium Publications **61**, American Mathematical Society, (2013)
- [5] J. Arthur, *The trace formula in invariant form*, Ann. of Math. (2) **114** (1981), 1–74.
- [6] J. Bernstein and A. Zelevinsky, *Representations of the group $GL(n, F)$, where F is a non-archimedean local field*, Russian Math. Surveys, **31** (1976), 1–68.
- [7] R. Beuzart-Plessis, Pierre-Henri Chaudouard, M. Zydor, *The global Gan-Gross-Prasad conjecture for unitary groups: the endoscopic case*, <https://arxiv.org/pdf/2007.05601.pdf>
- [8] J. Bernstein and A. Zelevinsky, *Induced representations of reductive p -adic groups I*, Ann. Sci. Ec. Norm. Super. (4) **10** (1977), 441–472.
- [9] R. Beuzart-Plessis, Yifeng Liu, Wei Zhang, Xinwen Zhu, *Isolation of cuspidal spectrum, with application to the Gan-Gross-Prasad conjecture*, <https://arxiv.org/pdf/1912.07169.pdf>
- [10] W. Casselman, *Canonical extensions of Harish-Chandra modules to representations of G* , Canad. J. Math. **41** (1989), 385–438.
- [11] J. Cogdell, I. Piatetski-Shapiro and F. Shahidi, *Functoriality for the classical groups*, Publ. Math. Inst. Hautes Etudes Sci. **99** (2004), 163–233.
- [12] M. Furusawa, K. Morimoto, *On special Bessel periods and the Gross-Prasad conjecture for $SO(2n+1) \times SO(2)$* , Math. Ann., **368**(1-2), (2017), 561–586.
- [13] M. Furusawa, K. Morimoto, *Refined global Gross-Prasad conjecture on special Bessel periods and Boecherer's conjecture*, J. Eur. Math. Soc., **23**(4), (2021), 1295 - 1331.
- [14] D. Ginzburg, D. Jiang, B. Liu, S.Rallis, *Erratum to "On the Non-vanishing of the Central Value of the Rankin-Selberg L -functions"*, preprint 2019, <https://arxiv.org/pdf/1905.02644.pdf>

- [15] D. Ginzburg, D. Jiang, S. Rallis, *On the nonvanishing of the central critical value of the Rankin-Selberg L -functions*, J. Amer. Math. Soc., **17**, (2004), no. 3, 679-722.
- [16] B. Gross and D. Prasad, *On the decomposition of a representation of $SO(n)$ when restricted to $SO(n-1)$* , Cand. J. Math. **44**, (1992), 974-1002.
- [17] W. T. Gan, B. Gross and D. Prasad, *Branching laws for classical groups: The non-tempered case*, Compositio Mathematica, **156**, Issue 11, (2020), 2298-2367.
- [18] W. T. Gan, B. Gross and D. Prasad, *Symplectic local root numbers, central critical L -values, and restriction problems in the representation theory of classical groups*, Astérisque **346** (2012), 1–110.
- [19] W. T. Gan, B. Gross and D. Prasad, *Restrictions of representations of classical groups: examples*, Astérisque **346** (2012), 111–170.
- [20] W. T. Gan and A. Ichino, *The Gross-Prasad conjecture and local theta correspondence*, Invent. Math. **206** (2016), 705–799.
- [21] S.S. Gelbart, J.D. Rogawski, *L -functions and Fourier-Jacobi coefficients for the unitary group $U(3)$* , Invent. Math. **105**(3), (1991), 445-472.
- [22] D. Ginzburg, S. Rallis and D. Soudry, *The descent map from automorphic representations of $GL(n)$ to classical groups*, World Scientific, Hackensack (2011).
- [23] D. Ginzburg, Piatetski-Shapiro, S. Rallis, *L functions for the orthogonal group*, Mem. Amer. Math. Soc. **128** (1997), no. 611.
- [24] J. Haan, *The Bessel periods : the orthogonal and unitary groups cases*, in preparation.
- [25] J. Haan, *The Bessel periods of Eisenstein series*, in preparation.
- [26] J. Haan, *The Fourier–Jacobi periods : the case of $Mp(2n+2r) \times Sp(2n)$* , in preparation.
- [27] J. Haan, *The regularized trilinear periods for general linear groups*, in preparation.
- [28] A. Ichino and T. Ikeda, *On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture*, Geometric Functional Analysis. **19** (5) (2010), 1378–1425.
- [29] A. Ichino and S. Yamana, *Period of automorphic forms: The case of $(GL_{n+1} \times GL_n, GL_n)$* , Compositio. Math. (151) (2015), 665–712.
- [30] A. Ichino and S. Yamana, *Period of automorphic forms: The case of $(U_{n+1} \times U_n, U_n)$* , J. Reine Angew. Math. **19** (5) (2019), 1–38.
- [31] H. Jacquet, E. Lapid, J. Rogawski, *Periods of automorphic forms*, J. Amer. Math. Soc., **12** (1999), 173–240.
- [32] H. Jacquet and S. Rallis, *On the Gross-Prasad conjecture for unitary groups. in On certain L -functions*, Clay Math. Proc. **13**, American Mathematical Society, Providence (2011), 205-264.
- [33] D. Jiang, D. Soudry, L. Zhang, *The unramified computation of Rankin-Selberg integrals for quasi-split classical groups: Bessel model case*, in preparation.
- [34] D. Jiang, L. Zhang, *Arthur parameters and cuspidal automorphic modules of classical groups*, Ann. of Math. (2) **191** (2020), 739-827.
- [35] D. Jiang, L. Zhang, *Arthur parameters and cuspidal automorphic modules of classical groups*, <https://arxiv.org/pdf/1508.03205v3.pdf>.
- [36] D. Jiang, L. Zhang, *Automorphic Integral Transforms for Classical group II: Twisted Descents*, in Representation Theory, Number Theory, and Invariant Theory, J. Cogdell et al. (eds.), Progress in Mathematics **323** (2017), 303-335.
- [37] D. Jiang, L. Zhang, *Arthur parameters and cuspidal automorphic modules of classical groups II: Fourier-Jacobi case*, (2015), in preparation
- [38] D. Jiang, L. Zhang, *A product of tensor product L -functions for classical groups of hermitian type*, Geom. Funct. Anal. **24** (2014), no. 2, 552-609.
- [39] T. Kaletha, A. Minguez, S. W. Shin and P.-J. White, *Endoscopic classification of representations: Inner forms of unitary groups*, preprint 2014, <https://arxiv.org/pdf/1409.3731.pdf>.
- [40] S. S. Kudla, *Splitting metaplectic covers of dual reductive pairs*, Israel J. Math. **87** (1994), 361–401.
- [41] C. Mœglin and J. L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Tracts in Math. **113**, Cambridge University Press, Cambridge (1995).
- [42] C. P. Mok, *Endoscopic classification of representations of quasi-split unitary groups*, Mem. Amer. Math. Soc. **235** (2015), no. 1108.
- [43] E. Lapid and J. Rogawski, *Periods of Eisenstein series: the Galois side*, Duke Math. J. **120**(1) (2003), 153–226.
- [44] Y. Liu, *Refined global Gan-Gross-Prasad conjecture for Bessel periods*, J. Reine Angew. Math. (717) (2016), 133-194.
- [45] Y. Liu and B. Sun, *Uniqueness of Fourier-Jacobi models: the Archimedean case*, J. Funct. Anal. **265** (2013), 3325–3340.
- [46] W. Luo, Z. Rudnick, P. Sarnak, *On the generalized Ramanujan conjecture for $GL(n)$* , Proc. Sym. Pure Math., **66**, Vol. II, (1999), 301-311.

- [47] D. Soudry, *Rankin-Selberg integrals, the descent method, and Langlands functoriality*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, (2006), 1311-1325.
- [48] D. Soudry, *The unramified computation of Rankin-Selberg integrals expressed in terms of Bessel models for split orthogonal groups: Part I*, Israel J. Math., **222**, (2017), no. 2, 711–786.
- [49] D. Soudry, *The unramified computation of Rankin-Selberg integrals expressed in terms of Bessel models for split orthogonal groups: Part II*, J. Number Theory., **186**, (2018), 62-102
- [50] F. Shahidi, *On certain L-functions*, Amer. J. Math., **103** (2), (1981), 297–355.
- [51] R. Berndt, Ralf Schmidt, *Elements of the Representation Theory of the Jacobi group*, Progress in Mathematics, Vol. 163, Birkhauser Verlag, Basel, (1998)
- [52] B. Sun, *Multiplicity one theorems for Fourier-Jacobi models*, Amer. J. Math., **134** (6) (2012), 1655–1678.
- [53] N. Wallach, *Real Reductive Groups vol. II*, Pure and Applied Mathematics, Volume **132** (Academic Press, Inc., Boston, MA, (1992). xiv+454 pp.
- [54] A. Weil, *Sur certains groupes d'opérateurs unitaires*, Acta Math, **111** (1964), 143–211.
- [55] H. Xue, *Fourier-Jacobi periods and the central critical value of Rankin-Selberg L-functions*, Israel J. Math., **212**, (2016), no. 2, 547-633.
- [56] H. Xue, *Refined global Gan-Gross-Prasad conjecture for Fourier-Jacobi periods on symplectic groups*, Compositio. Math. (**153**) (2017), no. 1, 68–131.
- [57] H. Xue, *The Gan-Gross-Prasad conjecture for $U(n) \times U(n)$* , Adv. Math., **262** (2014), 1130-1191.
- [58] S.Yamana, *Period of automorphic forms: The trilinear case*, J. Inst. Math. Jussieu., **17**(1), (2018), 59–74.
- [59] S.Yamana, *Periods of residual automorphic forms*, J. Funct. Anal. **268** (2015), 1078–1104.
- [60] A. Zelevinsky, *Induced representations of reductive p-adic groups. II: On irreducible representations of $GL(n)$* , Ann. Sci. Ec. Norm. Super. (4) **13** (1980), 165-210.
- [61] W. Zhang, *Fourier transform and the global Gan-Gross-Prasad conjecture for unitary groups*, Ann. of Math., (2) **180** (2014), 971-1049.
- [62] W. Zhang, *Automorphic period and the central value of Rankin-Selberg L-function*, J. Amer. Math. Soc., **27** (2014), 541-612.

DEPARTMENT OF MATHEMATICS YONSEI UNIVERSITY 50 YONSEI-RO, SEODAEMUN-GU, SEOUL 03722, REPUBLIC OF KOREA

Email address: jaehoahan@gmail.com