

Degree bounds for projective division fields associated to elliptic modules with a trivial endomorphism ring

Alina Carmen Cojocaru and Nathan Jones

ABSTRACT. Let k be a global field, let A be a Dedekind domain with $\text{Quot}(A) = k$, and let K be a finitely generated field. Using a unified approach for both elliptic curves and Drinfeld modules M defined over K and having a trivial endomorphism ring, with $k = \mathbb{Q}$, $A = \mathbb{Z}$ in the former case and k a global function field, A its ring of functions regular away from a fixed prime in the latter case, for any nonzero ideal $\mathfrak{a} \triangleleft A$ we prove best possible estimates in the norm $|\mathfrak{a}|$ for the degrees over K of the subfields of the \mathfrak{a} -division fields of M fixed by scalars.

1. Introduction

In the theory of elliptic modules – elliptic curves and Drinfeld modules – division fields play a fundamental role; their algebraic properties (e.g., ramification, degree, and Galois group structure) are intimately related to properties of Galois representations and are essential to global and local questions about elliptic modules themselves. Among the subfields of the division fields of an elliptic module, those fixed by the scalars are of special significance. For example, as highlighted in [Ad01, Chapter 5] and [CoDu04, Section 3], in the case of an elliptic curve E defined over \mathbb{Q} and a positive integer a , the subfield J_a of the a -division field $\mathbb{Q}(E[a])$ fixed by the scalars of $\text{Gal}(\mathbb{Q}(E[a]/\mathbb{Q})) \leq \text{GL}_2(\mathbb{Z}/a\mathbb{Z})$ is closely related to the modular curve $X_0(a)$ which parametrizes cyclic isogenies of degree a between elliptic curves; indeed, J_a may be interpreted as the splitting field of the modular polynomial $\Phi_a(X, j(E))$ (see [We08, Section 69] and [Co89, Section 11.C] for the properties of the modular polynomials $\Phi_a(X, Y)$). The arithmetic properties of the family of fields $(J_a)_{a \geq 1}$ are closely related to properties of the reductions $E(\text{mod } p)$ of E modulo primes p , including the growth of the order of the Tate-Shafarevich group of the curve $E(\text{mod } p)$ when viewed as constant over its own function field (see [CoDu04]) and the growth of the absolute discriminant of the endomorphism ring of the curve $E(\text{mod } p)$ when viewed over the finite field \mathbb{F}_p (see [CoF19]). An essential ingredient when

Key words and phrases: Elliptic curves, Drinfeld modules, division fields, Galois representations

2010 Mathematics Subject Classification: Primary 11G05, 11G09, 11F80

A.C.C. was partially supported by a Collaboration Grant for Mathematicians from the Simons Foundation under Award No. 318454.

deriving properties about $E(\text{mod } p)$ from the fields J_a is the growth of the degrees $[J_a : \mathbb{Q}]$. The goal of this article is to prove best possible estimates in a for the degrees of such fields in the unified setting of elliptic curves and Drinfeld modules with a trivial endomorphism ring.

To state our main result, we proceed as in [Br10] and fix: k a global field, A a Dedekind domain with $\text{Quot}(A) = k$, K a finitely generated field, and M a (G_K, A) -module of rank $r \geq 2$, where $G_K = \text{Gal}(K^{\text{sep}}/K)$ denotes the absolute Galois group of K . Specifically, M is an A -module with a continuous G_K -action that commutes with the A -action and with the property that, for any ideal $0 \neq \mathfrak{a} \triangleleft A$, the \mathfrak{a} -division submodule

$$M[\mathfrak{a}] := \{x \in M : \alpha x = 0 \forall \alpha \in \mathfrak{a}\}$$

has A -module structure

$$M[\mathfrak{a}] \simeq_A (A/\mathfrak{a})^r.$$

The (G_K, A) -module structure on M gives rise to a compatible system of Galois representations $\rho_{\mathfrak{a}} : G_K \rightarrow \text{GL}_r(A/\mathfrak{a})$ and to a continuous representation

$$\rho : G_K \rightarrow \text{GL}_r(\hat{A}),$$

where $\hat{A} := \varprojlim_{\mathfrak{a} \triangleleft A} A/\mathfrak{a}$. Associated to these representations we have the \mathfrak{a} -division fields $K_{\mathfrak{a}} := (K^{\text{sep}})^{\text{Ker } \rho_{\mathfrak{a}}}$, for which we distinguish the subfields $J_{\mathfrak{a}}$ fixed by the scalars $\{\lambda I_r : \lambda \in (A/\mathfrak{a})^{\times}\} \cap \text{Gal}(K_{\mathfrak{a}}/K)$ (with $\text{Gal}(K_{\mathfrak{a}}/K)$ viewed as a subgroup of $\text{GL}_r(A/\mathfrak{a})$).

Denoting by $\hat{\rho}_{\mathfrak{a}} : G_K \rightarrow \text{PGL}_r(A/\mathfrak{a})$ the composition of the representation $\rho_{\mathfrak{a}}$ with the canonical projection $\text{GL}_r(A/\mathfrak{a}) \rightarrow \text{PGL}_r(A/\mathfrak{a})$, we observe that $J_{\mathfrak{a}} = (K^{\text{sep}})^{\text{Ker } \hat{\rho}_{\mathfrak{a}}}$ and we deduce that $[J_{\mathfrak{a}} : K] \leq |\text{PGL}_r(A/\mathfrak{a})|$. Our main result provides a lower bound for $[J_{\mathfrak{a}} : K]$ of the same order of growth as $|\text{PGL}_r(A/\mathfrak{a})|$, as follows:

THEOREM 1. *We keep the above setting and assume that*

$$(1) \quad \left| \text{GL}_r(\hat{A}) : \rho(G_K) \right| < \infty.$$

Then, for any ideal $0 \neq \mathfrak{a} \triangleleft A$,

$$(2) \quad |\mathfrak{a}|^{r^2-1} \ll_{M,K} [J_{\mathfrak{a}} : K] \leq |\mathfrak{a}|^{r^2-1},$$

where $|\mathfrak{a}| := |A/\mathfrak{a}|$.

By specializing the above general setting to elliptic curves and to Drinfeld modules, we obtain:

COROLLARY 2. *Let K be a finitely generated field with $\text{char } K = 0$ and let E/K be an elliptic curve over K with $\text{End}_{\overline{K}}(E) \simeq \mathbb{Z}$. Then, for any integer $a \geq 1$, the degree $[J_a : K]$ of the subfield J_a of the a -division field $K_a := K(E[a])$ fixed by the scalars of $\text{Gal}(K(E[a])/K)$ satisfies*

$$(3) \quad a^3 \ll_{E,K} [J_a : K] \leq a^3.$$

COROLLARY 3. *Let k be a global function field, let ∞ be a fixed place of k , let A be the ring of elements of k regular away from ∞ , let K be a finitely generated A -field with $A\text{-char } K = 0$ (i.e. $k \subseteq K$), and let $\psi : A \rightarrow K\{\tau\}$ be a Drinfeld A -module over K of rank $r \geq 2$ with $\text{End}_{\overline{K}}(\psi) \simeq A$. Then, for any ideal $0 \neq \mathfrak{a} \triangleleft A$, the degree $[J_{\mathfrak{a}} : K]$ of the subfield $J_{\mathfrak{a}}$ of the \mathfrak{a} -division field $K_{\mathfrak{a}} := K(\psi[\mathfrak{a}])$ fixed by the scalars of $\text{Gal}(K(\psi[\mathfrak{a}])/K)$ satisfies*

$$(4) \quad |\mathfrak{a}|^{r^2-1} \ll_{\psi, K} [J_{\mathfrak{a}} : K] \leq |\mathfrak{a}|^{r^2-1}.$$

The proof of Theorem 1 relies on consequences of assumption (1), on applications of Goursat’s Lemma, as well as on vertical growth estimates for open subgroups of GL_r . Specializing to elliptic curves and to Drinfeld modules, assumption (1) is essentially Serre’s Open Image Theorem [Se72] and, respectively, Pink-Rütsche’s Open Image Theorem [PiRu09]. Variations of these open image theorems also hold for elliptic curves and Drinfeld modules with nontrivial endomorphism rings. While these complementary cases are treated unitarily in [Br10] when investigating the growth of torsion, when investigating the growth of $[J_{\mathfrak{a}} : K]$ they face particularities whose treatment we relegate to future work.

We emphasize that the upper bound in Theorem 1 always holds and does not necessitate assumption (1). In contrast, the lower bound in Theorem 1 is intimately related to assumption (1). Indeed, one consequence of (1) is that there exists an ideal $\mathfrak{a}(M, K) \trianglelefteq A$, which (a priori) depends on M and K and which has the property that, for any prime ideal $\mathfrak{l} \nmid \mathfrak{a}(M, K)$, $\text{Gal}(J_{\mathfrak{l}}/K) \simeq \text{PGL}_r(A/\mathfrak{l})$. Then, for such an ideal \mathfrak{l} , the lower bound in (2) follows immediately. The purpose of Theorem 1 is to prove similar lower bounds for *all* ideals $\mathfrak{a} \triangleleft A$.

The dependence of the lower bound in (2) on M (which also includes dependence on r) and on K is an important topic related to the uniform boundedness of the torsion of M ; while we do not address it in the present paper, we refer the reader to [Br10] and [Jo19] for related discussions and for additional references.

The fields $J_{\mathfrak{a}}$ play a prominent role in a multitude of problems, such as in deriving non-trivial upper bounds for the number of non-isomorphic Frobenius fields associated to an elliptic curve and, respectively, to a Drinfeld module (see [CoDa08ec] and [CoDa08dm]); in investigating the discriminants of the endomorphism rings of the reduction of an elliptic curve and, respectively, of a Drinfeld module (see [CoFi19] and [CoPa20]); and in proving non-abelian reciprocity laws for primes and, respectively, for irreducible polynomials (see [DuTo02], [CoPa15], and [GaPa19]). For such applications, an essential piece of information is the growth of the degree $[J_{\mathfrak{a}} : K]$ as a function of the norm $|\mathfrak{a}|$. For example, Corollary 2 is a key ingredient in proving that, for any elliptic curve E/\mathbb{Q} with $\text{End}_{\overline{\mathbb{Q}}}(E) \simeq \mathbb{Z}$, provided the Generalized Riemann Hypothesis holds for the division fields of E , there exists a set of primes p of natural density 1 with the property that the absolute discriminant of the imaginary quadratic order $\text{End}_{\mathbb{F}_p}(E)$ is as close as possible to its natural upper bound; see [CoFi19, Theorem 1]. Similarly, Corollary 3 is a key ingredient in proving that, denoting by \mathbb{F}_q the finite field with q elements and assuming that q is odd, for any generic Drinfeld module $\psi : \mathbb{F}_q[T] \rightarrow \mathbb{F}_q(T)\{\tau\}$ of rank 2 and with $\text{End}_{\overline{\mathbb{F}_q(T)}}(\psi) \simeq \mathbb{F}_q[T]$, there exists a set of prime

ideals $\mathfrak{p} \triangleleft \mathbb{F}_q[T]$ of Dirichlet density 1 with the property that the norm of the discriminant of the imaginary quadratic order $\text{End}_{\mathbb{F}_p}(\psi)$ is as close as possible to its natural upper bound; see [CoPa20, Theorem 6]. We expect that Theorem 1 will be of use to other arithmetic studies of elliptic modules.

Notation. In what follows, we use the standard \ll , \gg , and \asymp notation: given suitably defined real functions h_1, h_2 , we say that $h_1 \ll h_2$ or $h_2 \gg h_1$ if h_2 is positive valued and there exists a positive constant C such that $|h_1(x)| \leq Ch_2(x)$ for all x in the domain of h_1 ; we say that $h_1 \asymp h_2$ if h_1, h_2 are positive valued and $h_1 \ll h_2 \ll h_1$; we say that $h_1 \ll_D h_2$ or $h_2 \gg_D h_1$ if $h_1 \ll h_2$ and the implied \ll -constant C depends on priorly given data D ; similarly, we say that $h_1 \asymp_D h_2$ if the implied constant in either one of the \ll -bounds $h_1 \ll h_2 \ll h_1$ depends on priorly given data D . We also use the standard divisibility notation for ideals in a Dedekind domain. In particular, given two ideals $\mathfrak{a}, \mathfrak{b}$, we write $\mathfrak{a} \mid \mathfrak{b}^\infty$ if all the prime ideal factors of \mathfrak{a} are among the prime ideal factors of \mathfrak{b} (with possibly different exponents). Further notation will be introduced over the course of the paper as needed.

2. Goursat's Lemma and variations

In this section we recall Goursat's Lemma on fibered products of groups (whose definition we also recall shortly) and detail the behavior of such fibered products under intersection.

LEMMA 4. (*Goursat's Lemma*)

Let G_1, G_2 be groups and for $i \in \{1, 2\}$ denote by $\pi_i : G_1 \times G_2 \rightarrow G_i$ the projection map onto the i -th factor. Let $G \leq G_1 \times G_2$ be a subgroup and assume that $\pi_1(G) = G_1, \pi_2(G) = G_2$. Then there exist a group Γ and a pair of surjective group homomorphisms $\psi_1 : G_1 \rightarrow \Gamma, \psi_2 : G_2 \rightarrow \Gamma$ such that

$$G = G_1 \times_\psi G_2 := \{(g_1, g_2) \in G_1 \times G_2 : \psi_1(g_1) = \psi_2(g_2)\}.$$

PROOF. See [Ri76, Lemma 5.2.1]. □

We call $G_1 \times_\psi G_2$ the *fibered product of G_1 and G_2 over $\psi := (\psi_1, \psi_2)$* . The next lemma details what happens when we intersect such a fibered product with a subgroup of the form $H_1 \times H_2$ defined by subgroups $H_1 \leq G_1$ and $H_2 \leq G_2$.

It is clear that

$$(H_1 \times H_2) \cap (G_1 \times_\psi G_2) = H_1 \times_\psi H_2 := \{(h_1, h_2) \in H_1 \times H_2 : \psi_1(h_1) = \psi_2(h_2)\}.$$

However, this representation does not specify the restricted common quotient inside Γ . In particular, it can be the case that the fibered product $H := H_1 \times_\psi H_2$ does *not* satisfy $\pi_i(H) = H_i$ for each $i \in \{1, 2\}$. The following lemma clarifies this situation.

LEMMA 5. Let G_1, G_2 be groups, let $\psi_1 : G_1 \rightarrow \Gamma, \psi_2 : G_2 \rightarrow \Gamma$ be surjective group homomorphisms onto a group Γ , and let $G_1 \times_\psi G_2$ be the associated fibered product. Furthermore, let $H_1 \leq G_1, H_2 \leq G_2$ be

subgroups. Define the subgroup

$$\Gamma_H := \psi_1(H_1) \cap \psi_2(H_2) \leq \Gamma.$$

Then

$$(5) \quad (H_1 \times H_2) \cap (G_1 \times_\psi G_2) = (H_1 \cap \psi_1^{-1}(\Gamma_H)) \times_\psi (H_2 \cap \psi_2^{-1}(\Gamma_H))$$

and the canonical projection maps

$$\pi_1 : (H_1 \cap \psi_1^{-1}(\Gamma_H)) \times_\psi (H_2 \cap \psi_2^{-1}(\Gamma_H)) \longrightarrow H_1 \cap \psi_1^{-1}(\Gamma_H),$$

$$\pi_2 : (H_1 \cap \psi_1^{-1}(\Gamma_H)) \times_\psi (H_2 \cap \psi_2^{-1}(\Gamma_H)) \longrightarrow H_2 \cap \psi_2^{-1}(\Gamma_H)$$

are surjective.

PROOF. We first establish (5). Since the containment “ \supseteq ” is immediate, we only need to establish “ \subseteq .” Let $(h_1, h_2) \in (H_1 \times H_2) \cap (G_1 \times_\psi G_2)$, i.e. $h_1 \in H_1$, $h_2 \in H_2$, and $\psi_1(h_1) = \psi_2(h_2)$. From the definition of Γ_H , it follows that $\psi_1(h_1) = \psi_2(h_2) \in \Gamma_H$. Thus $(h_1, h_2) \in (H_1 \cap \psi_1^{-1}(\Gamma_H)) \times_\psi (H_2 \cap \psi_2^{-1}(\Gamma_H))$, establishing (5).

To see why the projection map

$$(6) \quad \pi_1 : H_1 \cap \psi_1^{-1}(\Gamma_H) \longrightarrow H_1 \cap \psi_1^{-1}(\Gamma_H)$$

is surjective, fix $h_1 \in H_1 \cap \psi_1^{-1}(\Gamma_H)$ and set $\gamma := \psi_1(h_1) \in \Gamma_H$. By the definition of Γ_H , we find $h_2 \in H_2$ with $\psi_2(h_2) = \gamma$. Thus $(h_1, h_2) \in (H_1 \cap \psi_1^{-1}(\Gamma_H)) \times_\psi (H_2 \cap \psi_2^{-1}(\Gamma_H))$ and $\pi_1(h_1, h_2) = h_1$, proving the surjectivity of π_1 in (6). The surjectivity of π_2 is proved similarly. \square

3. Proof of Theorem 1

In this section we prove Theorem 1. We will make use of the following notation:

$$G := \rho(G_K) \leq \mathrm{GL}_r(\hat{A});$$

for any ideal $0 \neq \mathfrak{a} \triangleleft A$, we write

$$G(\mathfrak{a}) := \rho_{\mathfrak{a}}(G_K) \leq \mathrm{GL}_r(A/\mathfrak{a});$$

for any subgroup $H \leq \mathrm{GL}_r(A/\mathfrak{a})$, we write

$$\mathrm{Scal}_H := H \cap \{\alpha I_r : \alpha \in (A/\mathfrak{a})^\times\}.$$

With this notation, we see that $J_{\mathfrak{a}} = K(E[\mathfrak{a}])^{\mathrm{Scal}_{G(\mathfrak{a})}}$.

To prove the theorem, let $0 \neq \mathfrak{a} \triangleleft A$ be a fixed arbitrary ideal. The proof of the upper bound is an immediate consequence to the injection $\mathrm{Gal}(J_{\mathfrak{a}}/K) \hookrightarrow \mathrm{PGL}_r(A/\mathfrak{a})$ defined by $\hat{\rho}_{\mathfrak{a}}$. Indeed, using that

$$|\mathrm{PGL}_r(A/\mathfrak{a})| = \frac{1}{|(A/\mathfrak{a})^\times|} |\mathrm{GL}_r(A/\mathfrak{a})|,$$

$$|(A/\mathfrak{a})^\times| = |\mathfrak{a}| \prod_{\mathfrak{p}|\mathfrak{a}} \left(1 - \frac{1}{|\mathfrak{p}|}\right),$$

and

$$|\mathrm{GL}_r(A/\mathfrak{a})| = |\mathfrak{a}|^{r^2} \prod_{\substack{\mathfrak{p}|\mathfrak{a} \\ \mathfrak{p} \text{ prime}}} \left(1 - \frac{1}{|\mathfrak{p}|}\right) \left(1 - \frac{1}{|\mathfrak{p}|^2}\right) \cdots \left(1 - \frac{1}{|\mathfrak{p}|^r}\right)$$

(see [Br10, Lemma 2.3, p. 1244] for the latter), we obtain that

$$[J_{\mathfrak{a}} : K] \leq |\mathrm{PGL}_r(A/\mathfrak{a})| = |\mathfrak{a}|^{r^2-1} \prod_{\substack{\mathfrak{p}|\mathfrak{a} \\ \mathfrak{p} \text{ prime}}} \left(1 - \frac{1}{|\mathfrak{p}|^2}\right) \cdots \left(1 - \frac{1}{|\mathfrak{p}|^r}\right) \leq |\mathfrak{a}|^{r^2-1}.$$

The proof of the lower bound relies on several consequences to assumption (1), as well as on applications of Goursat's Lemma 4 and its variation Lemma 5, as detailed below.

Thanks to (1), there exists an ideal $\mathfrak{m} = \mathfrak{m}_{M,K} \trianglelefteq A$ such that

$$(7) \quad G = \pi^{-1}(G(\mathfrak{m})),$$

where $\pi : \mathrm{GL}_r(\hat{A}) \rightarrow \mathrm{GL}_r(A/\mathfrak{m})$ is the canonical projection. We take \mathfrak{m} to be the smallest such ideal with respect to divisibility and we write its unique prime ideal factorization as $\mathfrak{m} = \prod_{\mathfrak{p}^{v_{\mathfrak{p}}(\mathfrak{m})} || \mathfrak{m}} \mathfrak{p}^{v_{\mathfrak{p}}(\mathfrak{m})}$, where each exponent satisfies $v_{\mathfrak{p}}(\mathfrak{m}) \geq 1$.

With the ideal \mathfrak{m} in mind, we write the arbitrary ideal \mathfrak{a} uniquely as

$$(8) \quad \mathfrak{a} = \mathfrak{a}_1 \mathfrak{a}_2,$$

where

$$(9) \quad \mathfrak{a}_1 | \mathfrak{m}^{\infty},$$

$$(10) \quad \mathrm{gcd}(\mathfrak{a}_2, \mathfrak{m}) = 1.$$

For future use, we record that

$$(11) \quad \mathrm{gcd}(\mathfrak{a}_1, \mathfrak{a}_2) = 1.$$

We also write the ideal \mathfrak{a}_1 uniquely as

$$\mathfrak{a}_1 = \mathfrak{a}_{1,1} \mathfrak{a}_{1,2},$$

where $\mathfrak{a}_{1,1} = \prod_{\substack{\mathfrak{p}^{e_{\mathfrak{p}}} || \mathfrak{a}_{1,1} \\ e_{\mathfrak{p}} > v_{\mathfrak{p}}(\mathfrak{m})}} \mathfrak{p}^{e_{\mathfrak{p}}}$ and $\mathfrak{a}_{1,2} = \prod_{\substack{\mathfrak{p}^{f_{\mathfrak{p}}} || \mathfrak{a}_{1,1} \\ f_{\mathfrak{p}} \leq v_{\mathfrak{p}}(\mathfrak{m})}} \mathfrak{p}^{f_{\mathfrak{p}}}$. Note that

$$(12) \quad \mathrm{gcd}(\mathfrak{a}_{1,1}, \mathfrak{a}_{1,2}) = 1,$$

$$(13) \quad \mathfrak{a}_{1,1} | \mathfrak{m}^{\infty}$$

and

$$(14) \quad \mathfrak{a}_{1,2} | \mathfrak{m}.$$

Under the isomorphism of the Chinese Remainder Theorem, we deduce from (7) that

$$(15) \quad G(\mathfrak{a}) \simeq G(\mathfrak{a}_1) \times \mathrm{GL}_r(A/\mathfrak{a}_2)$$

and, consequently, that there exist group isomorphisms

$$(16) \quad \mathrm{Scal}_{G(\mathfrak{a})} \simeq \mathrm{Scal}_{G(\mathfrak{a}_1)} \times \mathrm{Scal}_{\mathrm{GL}_r(A/\mathfrak{a}_2)},$$

$$(17) \quad G(\mathfrak{a})/\mathrm{Scal}_{G(\mathfrak{a})} \simeq (G(\mathfrak{a}_1)/\mathrm{Scal}_{G(\mathfrak{a}_1)}) \times \mathrm{PGL}_r(A/\mathfrak{a}_2).$$

Next, applying Lemma 4 to the groups $G = G(\mathfrak{a}_1)$, $G_1 = G(\mathfrak{a}_{1,1})$, and $G_2 = G(\mathfrak{a}_{1,2})$, we deduce that there exist a group Γ and surjective group homomorphisms $\psi_1 : G(\mathfrak{a}_{1,1}) \rightarrow \Gamma$, $\psi_2 : G(\mathfrak{a}_{1,2}) \rightarrow \Gamma$, which give rise to a group isomorphism

$$(18) \quad G(\mathfrak{a}_1) \simeq G(\mathfrak{a}_{1,1}) \times_{\psi} G(\mathfrak{a}_{1,2}).$$

Furthermore, applying Lemma 5 to the subgroups $H_1 = \mathrm{Scal}_{G(\mathfrak{a}_{1,1})}$ and $H_2 = \mathrm{Scal}_{G(\mathfrak{a}_{1,2})}$, we deduce that there exists a group isomorphism

$$(19) \quad (\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \times \mathrm{Scal}_{G(\mathfrak{a}_{1,2})}) \cap (G(\mathfrak{a}_{1,1}) \times_{\psi} G(\mathfrak{a}_{1,2})) \simeq (\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \cap \psi_1^{-1}(\Gamma_{\mathrm{Scal}})) \times_{\psi} (\mathrm{Scal}_{G(\mathfrak{a}_{1,2})} \cap \psi_2^{-1}(\Gamma_{\mathrm{Scal}})),$$

where

$$\Gamma_{\mathrm{Scal}} := \psi_1(\mathrm{Scal}_{G(\mathfrak{a}_{1,1})}) \cap \psi_2(\mathrm{Scal}_{G(\mathfrak{a}_{1,2})}) \leq \Gamma.$$

From (17) we derive that

$$(20) \quad [J_{\mathfrak{a}} : K] = |G(\mathfrak{a})/\mathrm{Scal}_{G(\mathfrak{a})}| = |G(\mathfrak{a}_1)/\mathrm{Scal}_{G(\mathfrak{a}_1)}| \cdot |\mathrm{PGL}_r(A/\mathfrak{a}_2)|.$$

Then, using (18) and (19), we derive that

$$(21) \quad \begin{aligned} |G(\mathfrak{a}_1)/\mathrm{Scal}_{G(\mathfrak{a}_1)}| &= \frac{|G(\mathfrak{a}_{1,1}) \times_{\psi} G(\mathfrak{a}_{1,2})|}{|(\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \times \mathrm{Scal}_{G(\mathfrak{a}_{1,2})}) \cap G(\mathfrak{a}_1)|} \\ &= \frac{|G(\mathfrak{a}_{1,1}) \times_{\psi} G(\mathfrak{a}_{1,2})|}{|(\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \cap \psi_1^{-1}(\Gamma_{\mathrm{Scal}})) \times_{\psi} (\mathrm{Scal}_{G(\mathfrak{a}_{1,2})} \cap \psi_2^{-1}(\Gamma_{\mathrm{Scal}}))|} \\ &= \frac{|G(\mathfrak{a}_{1,1})|}{|\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \cap \psi_1^{-1}(\Gamma_{\mathrm{Scal}})|} \cdot \frac{|\Gamma_{\mathrm{Scal}}|}{|\Gamma|} \cdot \frac{|G(\mathfrak{a}_{1,2})|}{|\mathrm{Scal}_{G(\mathfrak{a}_{1,2})} \cap \psi_2^{-1}(\Gamma_{\mathrm{Scal}})|} \\ &= \frac{|G(\mathfrak{a}_{1,1})|}{|\mathrm{Scal}_{G(\mathfrak{a}_{1,1})} \cap \psi_1^{-1}(\Gamma_{\mathrm{Scal}})|} \cdot \frac{|\psi_1(\mathrm{Scal}_{G(\mathfrak{a}_{1,1})}) \cap \psi_2(\mathrm{Scal}_{G(\mathfrak{a}_{1,2})})|}{|\Gamma|} \cdot \frac{|G(\mathfrak{a}_{1,2})|}{|\mathrm{Scal}_{G(\mathfrak{a}_{1,2})} \cap \psi_2^{-1}(\Gamma_{\mathrm{Scal}})|}. \end{aligned}$$

Recalling (14), we deduce that the last two factors above are bounded, from above and below, by constants depending on \mathfrak{m} , hence on M and K :

$$(22) \quad \frac{|\psi_1(\mathrm{Scal}_{G(\mathfrak{a}_{1,1})}) \cap \psi_2(\mathrm{Scal}_{G(\mathfrak{a}_{1,2})})|}{|\Gamma|} \cdot \frac{|G(\mathfrak{a}_{1,2})|}{|\mathrm{Scal}_{G(\mathfrak{a}_{1,2})} \cap \psi_2^{-1}(\Gamma_{\mathrm{Scal}})|} \asymp_{M,K} 1.$$

It remains to analyze the first factor in (21). For this, consider the canonical projection

$$\pi_{1,1} : \mathrm{GL}_r(A/\mathfrak{a}_{1,1}) \rightarrow \mathrm{GL}_r(A/\mathrm{gcd}(\mathfrak{a}_{1,1}, \mathfrak{m}))$$

and, upon recalling (7), observe that

$$(23) \quad G(\mathfrak{a}_{1,1}) = \pi_{1,1}^{-1}(G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})))$$

and

$$(24) \quad \text{Ker } \pi_{1,1} \subseteq \text{Ker } \psi_1.$$

Thus the subgroups $G(\mathfrak{a}_{1,1}) \leq \text{GL}_r(A/\mathfrak{a}_{1,1})$ and $G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})) \leq \text{GL}_r(A/\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))$, together with the group Γ , fit into a commutative diagram

$$\begin{array}{ccc} G(\mathfrak{a}_{1,1}) & \xrightarrow{\pi_{1,1}} & G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})) \\ & \searrow \psi_1 & \downarrow \rho \\ & & \Gamma \end{array}$$

in which the vertical map ρ is some surjective group homomorphism and the horizontal map $\pi_{1,1}|_{G(\mathfrak{a}_{1,1})}$ is $\left(\frac{|\mathfrak{a}_{1,1}|}{|\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})|}\right)^{r^2}$ to 1. Furthermore, the subgroups $\psi_1^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\mathfrak{a}_{1,1})} \leq \text{Scal}_{\text{GL}_r(A/\mathfrak{a}_{1,1})} \simeq (A/\mathfrak{a}_{1,1})^\times$ and $\rho^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))} \leq \text{Scal}_{\text{GL}_r(A/\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))} \simeq (A/\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))^\times$, together with the group Γ_{Scal} , fit into the commutative diagram

$$\begin{array}{ccc} \psi_1^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\mathfrak{a}_{1,1})} & \xrightarrow{\pi_{1,1}} & \rho^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))} \\ & \searrow \psi_1 & \downarrow \rho \\ & & \Gamma_{\text{Scal}} \end{array}$$

in which the horizontal map $\pi_{1,1}|_{\psi_1^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\mathfrak{a}_{1,1})}}$ is $\frac{|\mathfrak{a}_{1,1}|}{|\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})|}$ to 1. We deduce that

$$(25) \quad |G(\mathfrak{a}_{1,1})| = \left(\frac{|\mathfrak{a}_{1,1}|}{|\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})|}\right)^{r^2} |G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))| \asymp_{M,K} |\mathfrak{a}_{1,1}|^{r^2}$$

and

$$(26) \quad |\psi_1^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\mathfrak{a}_{1,1})}| = \frac{|\mathfrak{a}_{1,1}|}{|\gcd(\mathfrak{a}_{1,1}, \mathfrak{m})|} |\rho^{-1}(\Gamma_{\text{Scal}}) \cap \text{Scal}_{G(\gcd(\mathfrak{a}_{1,1}, \mathfrak{m}))}| \asymp_{M,K} |\mathfrak{a}_{1,1}|.$$

Putting together (20), (21), (22), (25), and (26), we obtain that

$$(27) \quad [J_{\mathfrak{a}} : K] \asymp_K |\mathfrak{a}_{1,1}|^{r^2-1} |\text{PGL}_r(A/\mathfrak{a}_2)|.$$

To conclude the proof, observe that

$$\begin{aligned} |\text{PGL}_r(A/\mathfrak{a}_2)| &= |\mathfrak{a}_2|^{r^2-1} \prod_{\substack{\mathfrak{p}|\mathfrak{a}_2 \\ \mathfrak{p} \text{ prime}}} \left(1 - \frac{1}{|\mathfrak{p}|^2}\right) \cdots \left(1 - \frac{1}{|\mathfrak{p}|^r}\right) \\ &\geq |\mathfrak{a}_2|^{r^2-1} \prod_{\substack{\mathfrak{p} \\ \mathfrak{p} \text{ prime}}} \left(1 - \frac{1}{|\mathfrak{p}|^2}\right) \cdots \left(1 - \frac{1}{|\mathfrak{p}|^r}\right) \\ &\gg_{r,K} |\mathfrak{a}_2|^{r^2-1}, \end{aligned}$$

which, combined with (27), (8) and (14), gives

$$[J_{\mathfrak{a}} : K] \asymp_K \frac{|\mathfrak{a}_1|^{r^2-1}}{|\mathfrak{a}_{1,2}|^{r^2-1}} |\mathrm{PGL}_r(A/\mathfrak{a}_2)| \gg_{r,K} \frac{|\mathfrak{a}_1 \mathfrak{a}_2|^{r^2-1}}{|\mathfrak{a}_{1,2}|^{r^2-1}} \gg_{M,K} |\mathfrak{a}|^{r^2-1}.$$

4. Proof of Corollaries 2 and 3

First consider the setting of Corollary 2: K a finitely generated field with $\mathrm{char} K = 0$ and E/K an elliptic curve over K with $\mathrm{End}_{\overline{K}}(E) \simeq \mathbb{Z}$. This is the specialization to the setting of Theorem 1 to $k = \mathbb{Q}$, $A = \mathbb{Z}$, K as above, and $M = E$. In this case, $r = 2$ and assumption (1) holds thanks to an extension of Serre’s Open Image Theorem for elliptic curves over number fields [Se72, Théorème 3, p. 299] as explained in [Br10, Theorem 3.2, p. 1248]. Corollary 2 follows.

Next consider the setting of Corollary 3: k a global function field, ∞ a fixed place of k , A the ring of elements of k regular away from ∞ , K a finitely generated A -field with $\mathrm{char} K = \mathrm{char} k$ and A -char $K = 0$, and $\psi : A \rightarrow K\{\tau\}$ a Drinfeld A -module over K of rank $r \geq 2$ with $\mathrm{End}_{\overline{K}}(\psi) \simeq A$. This is the specialization to the setting of Theorem 1 to k, A, K as above, and $M = \psi$. In this case, assumption (1) holds thanks to Pink-Rütsche’s Open Image Theorem for Drinfeld modules [PiRu09, Theorem 0.1, p. 883]. Corollary 3 follows.

References

- [Ad01] C. Adelmann, *The decomposition of primes in torsion point fields*, Lecture Notes in Mathematics, 1761, Springer-Verlag, Berlin, 2001.
- [Br10] F. Breuer, *Torsion bounds for elliptic curves and Drinfeld modules*, Journal of Number Theory 130 (2010), no. 5, pp. 1241–1250.
- [CoDa08ec] A.C. Cojocaru and C. David, *Frobenius fields for elliptic curves*, American Journal of Mathematics 130 (2008), no. 6, pp. 1535–1560.
- [CoDa08dm] A.C. Cojocaru and C. David, *Frobenius fields for Drinfeld modules of rank 2*, Compositio Mathematica 144 (2008), no. 4, pp. 827–848.
- [CoDu04] A.C. Cojocaru and W. Duke, *Reductions of an elliptic curve and their Tate-Shafarevich groups*, Mathematische Annalen 329 (2004), no. 3, pp. 513–534.
- [CoFi19] A.C. Cojocaru and M. Fitzpatrick, *The absolute discriminant of the endomorphism ring of most reductions of a non-CM elliptic curve is maximal*, preprint 2019.
- [CoPa15] A.C. Cojocaru and M. Papikian, *Drinfeld modules, Frobenius endomorphisms, and CM-liftings*, Int. Math. Res. Not. IMRN 2015, no. 17, pp. 7787–7825.
- [CoPa20] A.C. Cojocaru and M. Papikian, *The growth of the discriminant of the endomorphism ring of the reduction of a rank 2 generic Drinfeld module*, preprint 2020.
- [Co89] D.A. Cox, *Primes of the form $x^2 + ny^2$. Fermat, class field theory and complex multiplication*, John Wiley & Sons, Inc., New York, 1989.
- [DuTo02] W. Duke and Á. Tóth, *The splitting of primes in division fields of elliptic curves*, Experimental Mathematics 11 (2002), no. 4, pp. 555–565.
- [GaPa19] S. Garai and M. Papikian, *Endomorphism rings of reductions of Drinfeld modules*, Journal of Number Theory (2019), DOI: 10.1016/j.jnt.2019.02.008.

- [Jo19] N. Jones, *A bound for the conductor of an open subgroup of GL_2 associated to an elliptic curve*, preprint 2019, available at <https://arxiv.org/abs/1904.10431>
- [PiRu09] R. Pink and E. Rütsche, *Adelic openness for Drinfeld modules in generic characteristic*, *Journal of Number Theory* 129 (2009), no. 4, pp. 882–907.
- [Ri76] K. Ribet, *Galois action on division points of Abelian varieties with real multiplications*, *American Journal of Mathematics* 98 (1976), no. 3, pp. 751–804.
- [Se72] J-P. Serre, *Propriétés galoisiennes des points d'ordre fini des courbes elliptiques*, *Inventiones Mathematicae* 15 (1972), pp. 259–331.
- [We08] H. Weber, *Lehrbuch der Algebra*, Band III, 1908 (reprint AMS Chelsea, New York, 2001).

(Alina Carmen Cojocaru)

- DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S MORGAN ST, 322 SEO, CHICAGO, 60607, IL, USA;
- INSTITUTE OF MATHEMATICS “SIMION STOILOW” OF THE ROMANIAN ACADEMY, 21 CALEA GRIVITEI ST, BUCHAREST, 010702, SECTOR 1, ROMANIA

E-mail address, Alina Carmen Cojocaru: cojocaru@uic.edu

(Nathan Jones)

- DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S MORGAN ST, 322 SEO, CHICAGO, 60607, IL, USA;

E-mail address, Nathan Jones: ncjones@uic.edu