

A generalization of modular moonshine to composite order

Satoru Urano

Institute of Mathematics, University of Tsukuba

March 24, 2021

Abstract

We introduce a generalization of Brauer character to allow arbitrary finite length modules over discrete valuation rings. We show that the generalized super Brauer character of Tate cohomology is a linear combination of trace functions. Using this result, we find a counterexample to a conjecture of Borcherds about vanishing of Tate cohomology for Fricke elements of the Monster.

Contents

1	Introduction	1
	Notation	2
2	Definitions and basic notions	3
3	Modular moonshine for elements of prime order	6
4	Tate cohomology and ramified extensions	8
5	p-Brauer character	10

Acknowledgements

References

1 Introduction

Conway and Norton's Monstrous Moonshine conjecture [7] stated the existence of a graded representation $V = \bigoplus_{n \in \mathbb{Z}} V_n$ such that for any element g of the monster group \mathbb{M} , the McKay-Thompson series $T_g(\tau) = \sum_{n \in \mathbb{Z}} \text{Tr}(g|V_n)q^{n-1}$ ($q = e^{2\pi i\tau}$) is a Hauptmodul for some genus 0 congruence subgroup of $SL_2(\mathbb{R})$. Frenkel, Lepowsky, and Meurman [8] constructed a well-behaved graded representation as a vertex algebra, and Borcherds [2] showed that this vertex algebra satisfied Conway and Norton's conjecture.

The original modular moonshine conjectures of Ryba asserted the existence of vertex algebras gV over finite fields with finite group actions, such that the graded Brauer characters are genus zero modular functions. He also suggested a construction of the vertex algebra gV (see Section 3) for an element $g \in \mathbb{M}$ of prime order p and of conjugacy class pA , in terms of a self-dual integral form of the monster vertex algebra whose existence was proved later by Carnahan [5]. Borcherds and Ryba reinterpreted the modular moonshine conjectures in terms of Tate

cohomology with coefficients in V . They proved that $\hat{H}^1(g, V) = 0$ for any Fricke element g of odd prime order, and Borcherds proved that if h is a p -regular element of $C_{\mathbb{M}}(g)$, then $\widehat{\text{Tr}}(h|\hat{H}^i(g, V)) = \sum_{n \in \mathbb{Z}} \widehat{\text{Tr}}(h|\hat{H}^i(g, V_n))q^{n-1}$ ($i = 0, 1$) is a linear combination of Hauptmoduls for any non-Fricke element g of odd prime order [3],[4]. He conjectured that Tate cohomology $\hat{H}^1(g, V)$ for any Fricke element vanishes.

It is natural to consider Tate cohomology of V for elements of composite order, but there are no results in the literature. Tate cohomology does not yield vector spaces over a field when the element has composite order, but Brauer characters are only defined for such objects. Specifically, Tate cohomology yields torsion modules over the ground ring. We can't use the Brauer character for such modules because it is defined only for representations defined over a field and p -regular elements. So, we need to define a generalized Brauer character.

In section 2, we introduce some terminology in this paper. In section 3, we summarize the modular moonshine conjecture for elements of prime order. In section 4, we consider ramified extensions R_p of \mathbb{Z}_p and calculate Tate cohomology. In section 5, to generalize the modular moonshine conjectures, we define a generalized Brauer character and show that the generalized super Brauer character of Tate cohomology is a linear combination of traces. This is a generalization of Proposition 2.2 of [4] to the composite order case. Using this result, we give a counterexample to a conjecture of Borcherds about vanishing of Tate cohomology. We propose a weaker conjecture about vanishing of Tate cohomology.

Notation

A^G	The largest submodule of A on which G acts trivially.
Aut	The automorphism group of something.
\mathbb{C}	The complex numbers.
$C_G(g)$	The centralizer of g in G .
$\text{Cent}(G)$	The center of G .
End V	The endomorphisms on V .
\hat{f}	The discrete Fourier transform of f .
\mathbb{F}_p	A finite field of order p .
$\text{Frac}(R)$	The fraction field of R .
Frob_p	The Frobenius endomorphism.
g	An element of G , usually of order N .
$\langle g_1, g_2, \dots \rangle$	The group generated by $g_1, g_2, \dots \in G$.
G	A group, often cyclic group of order N and generated by g .
h	An N -regular element of G .
\mathbb{H}	The upper half plane of \mathbb{C} .
$\text{Hom}(A, B)$	The homomorphisms from A to B .
$\hat{H}^i(G, A)$	A Tate cohomology group of G with coefficients in the G -module A .
$\hat{H}^i(g, A)$	A Tate cohomology group $\hat{H}^i(\langle g \rangle, A)$.
$\hat{H}^*(g, A)$	The sum of the Tate cohomology groups $\hat{H}^0(g, A)$ and $\hat{H}^1(g, A)$.
I	The kernel of the natural map from $\mathbb{Z}_p[G]$ to \mathbb{Z}_p .
id_V	The identity map on V .
Im	The image of a map.
k	The residue field R/\mathfrak{m}_R with prime characteristic p .
Ker	The kernel of a map.
\mathfrak{m}_R	A maximal ideal of R .
\mathbb{M}	The monster simple group.
M_{24}	A Mathieu group.

N	The order of g .
Nr	The norm map.
$O_p(G)$	The largest normal p -subgroup of the finite group G .
p	A prime, often a prime factor of N .
\mathbb{Q}	The rational numbers.
\mathbb{R}	The real numbers.
R	A discrete valuation ring.
R_p	A finite extension of the p -adic integers.
SL_2	A special linear group.
Sym	The symmetric algebra of something.
$\text{Stab}_G(\lambda)$	The stabilizer subgroup of G with respect to λ .
$T_g(\tau)$	The McKay-Thompson series of g .
$\text{Tr}(g A)$	The trace of g on a module A .
$\widetilde{\text{Tr}}(h A)$	The Brauer character of h on a module A .
$\widetilde{\text{Tr}}_p(h A)$	A p -Brauer character of h on a module A .
v	A discrete valuation.
V	An integral form of the monster vertex algebra.
gV	A modular vertex algebra or superalgebra $\hat{H}^*(g, V)$.
$V[[z]]$	The space of formal power series with coefficients in V .
$V((z))$	The space of formal Laurent series with coefficients in V .
V_Λ	An integral form of the Leech lattice vertex algebra.
\mathbb{Z}	The integers.
\mathbb{Z}_p	The p -adic integers.
ζ_N	A primitive N -th root of unity.
Λ	The Leech lattice.
μ	The Möbius function.
φ	Euler's totient function.

2 Definitions and basic notions

In this section, we introduce some terminology in this paper and will give brief definitions.

Definition 2.1. ([1]) Let R be a commutative ring. A vertex algebra over R is an R -module V equipped with a distinguished element $\mathbf{1} \in V$, a distinguished linear transformation $T : V \rightarrow V$, and a left multiplication map $Y(\cdot, z) : V \rightarrow (\text{End } V)[[z, z^{-1}]]$, written $Y(a, z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$ satisfying the following conditions:

- For any $a \in V$, $Y(\mathbf{1}, z) = id_V$, and $Y(a, z)\mathbf{1} \in a + zV[[z]]$.
- For any $a, b \in V$, $Y(a, z)b \in V((z))$, i.e. $a_n b = 0$ for n sufficiently large.
- (The Jacobi identity) For any $a, b \in V$,

$$\begin{aligned} x^{-1} \delta \left(\frac{y-z}{x} \right) Y(a, y) Y(b, z) - x^{-1} \delta \left(\frac{z-y}{-x} \right) Y(b, z) Y(a, y) \\ = z^{-1} \delta \left(\frac{y-x}{z} \right) Y(Y(a, x)b, z), \end{aligned}$$

where $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$, called the formal delta function.

- For any $a \in V$, $[T, Y(a, z)] = \frac{d}{dz} Y(a, z)$.

$\mathbf{1}$ is called a vacuum vector and $Y(\cdot, z)$ is called a vertex operator.

Definition 2.2. The Leech lattice Λ is the unique free abelian group of rank 24 equipped with a symmetric bilinear form $\langle \cdot, \cdot \rangle : \Lambda \times \Lambda \rightarrow \mathbb{Z}$ that satisfies the following conditions:

- (even) For all $a \in \Lambda$, $\langle a, a \rangle \in 2\mathbb{Z}$.
- (positive definite) For all nonzero elements $a \in \Lambda$, $\langle a, a \rangle > 0$.
- (no roots) For all $a \in \Lambda$, $\langle a, a \rangle \neq 2$.
- (unimodular) If we choose a basis (a_1, \dots, a_{24}) of Λ , then we may form the Gram matrix $(\langle a_i, a_j \rangle)_{1 \leq i, j \leq 24}$ and its determinant is 1.

Definition 2.3. A Lie algebra over a commutative ring R is an R -module \mathfrak{g} equipped with an R -bilinear map $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying the following conditions:

- For any $x \in \mathfrak{g}$, $[x, x] = 0$.
- For any $x, y, z \in \mathfrak{g}$, $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0$.

We can construct a vertex algebra for any even positive definite lattice of finite rank [1]. Let V_Λ be an integral form of the vertex algebra constructed by Λ . Λ has a unique nonsplit central extension $\hat{\Lambda}$ such that if we denote a lifting of $a \in \Lambda$ by $e^a \in \hat{\Lambda}$, then $e^a e^b = \pm e^{a+b}$ and $e^a e^b = (-1)^{\langle a, b \rangle} e^b e^a$ so we have an exact sequence

$$0 \rightarrow \{\pm 1\} \rightarrow \hat{\Lambda} \rightarrow \Lambda \rightarrow 0.$$

The group of automorphisms of $\hat{\Lambda}$ that preserve the inner product on Λ is a nonsplit extension $2^{24} \cdot \text{Aut}(\Lambda)$ of the group $\text{Aut}(\Lambda)$, and this group acts on V_Λ faithfully.

A lattice vertex algebra is described by using notions of affine Lie algebras and twisted group rings. In this paper, we explain them briefly.

We consider a Lie algebra $\mathfrak{h} = \Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$ with $[a, b] = 0$ for any $a, b \in \mathfrak{h}$. Then we take the Lie algebra $\hat{\mathfrak{h}} = \mathfrak{h} \otimes \mathbb{Q}[t, t^{-1}] \oplus \mathbb{Q}c$ satisfying relations $[c, \hat{\mathfrak{h}}] = 0$ and $[x \otimes t^m, y \otimes t^n] = \langle x, y \rangle m \delta_{m+n, 0} c$, where $\delta_{i, j}$ is the Kronecker delta. Let $\hat{\mathfrak{h}}^\pm$ be a subalgebra $\mathfrak{h} \otimes t^{\pm 1} \mathbb{Q}[t^{\pm 1}]$ and $\hat{\mathfrak{h}}^0$ be a subalgebra $\mathfrak{h} \otimes 1$. We view \mathbb{Q} as $\hat{\mathfrak{h}}$ -module which $\hat{\mathfrak{h}}^{\geq 0}$ acts as 0 and c acts as 1.

We view \mathbb{Q} as $\{\pm 1\}$ -module which ± 1 acts as ± 1 . The twisted group ring $\mathbb{Q}\{\Lambda\}$ of Λ is defined to be $\mathbb{Q}[\hat{\Lambda}] \otimes_{\mathbb{Q}[\{\pm 1\}]} \mathbb{Q}$ and is spanned by elements $\iota(e^\lambda) := e^\lambda \otimes 1$ for $e^\lambda \in \hat{\Lambda}$. Then we have $e^a \cdot \iota(e^\lambda) = \iota(e^a e^\lambda)$ and $\{\pm 1\} \cdot \iota(e^\lambda) = \pm \iota(e^\lambda)$, hence $\mathbb{C}\{\Lambda\}$ is $\hat{\Lambda}$ -module.

$V_\Lambda \otimes \mathbb{Q}$ is isomorphic to $\text{Sym}_{\mathbb{Q}}(\hat{\mathfrak{h}}^-) \otimes \mathbb{Q}\{\Lambda\}$. $\hat{\Lambda}$ and $\hat{\mathfrak{h}}$ act on $V_\Lambda \otimes \mathbb{Q}$ as following:
For any $v \otimes \iota(e^\lambda) \in V_\Lambda$, $h \in \mathfrak{h}$, $n \neq 0$, and $e^a \in \hat{\Lambda}$, writing $h \otimes t^n$ by $h(n)$,

$$\begin{aligned} h(n) \cdot (v \otimes \iota(e^\lambda)) &= h(n)v \otimes \iota(e^\lambda), \quad h(0) \cdot (v \otimes \iota(e^\lambda)) = \langle h, \lambda \rangle (v \otimes \iota(e^\lambda)), \\ c \cdot (v \otimes \iota(e^\lambda)) &= v \otimes \iota(e^\lambda), \\ e^a \cdot (v \otimes \iota(e^\lambda)) &= v \otimes \iota(e^{a+\lambda}). \end{aligned}$$

$V_\Lambda \otimes \mathbb{Q}$ is graded by Λ , called Λ -degree, and we have $V_\Lambda \otimes \mathbb{Q} = \bigoplus_{\lambda \in \Lambda} \text{Sym}_{\mathbb{Q}}(\hat{\mathfrak{h}}^-) \otimes \mathbb{Q}\iota(e^\lambda)$.

Let $\mathbf{1}$ be a vacuum vector of V_Λ and let $Y(\cdot, z)$ be a vertex operator on V_Λ . We define the operator $D^{(n)}$ by $Y(v, z)\mathbf{1} = \sum_{n \in \mathbb{Z}} D^{(n)}(v)z^n$ for any $v \in V_\Lambda$. The operator $D^{(n)}$ has the properties $D^{(0)}(v) = v$, $D^{(n)}D^{(m)} = \binom{n+m}{n} D^{(n+m)}$, $D^{(n)} = 0$ if $n < 0$, and $D^{(n)}(vw) = \sum_{m \in \mathbb{Z}} D^{(m)}(v)D^{(n-m)}(w)$ for any $v, w \in V_\Lambda$. The derivation D is a element of $\hat{\mathfrak{h}}$ and has Λ -degree 0 i.e. $D^{(n)}(v)$ has the same Λ -degree as v .

Suppose that we choose a basis (a_1, \dots, a_{24}) of Λ and let p be a prime number. The subring $V_{\Lambda, 0}$ is generated as a commutative ring by $e^{-a_j} D^{(i)}(e^{a_j})$ for $i \geq 1$ and $1 \leq j \leq 24$ [4]. Let M be the free \mathbb{Z} -submodule of $\hat{\mathfrak{h}}_{\mathbb{Q}}$ given by the basis $\{e^{-a_j} D^{(i)}(e^{a_j})\}_{i \geq 1, 1 \leq j \leq 24}$. Then V_Λ is isomorphic to $\text{Sym}_{\mathbb{Z}}(M) \otimes \mathbb{Z}\{\Lambda\}$.

Definition 2.4. Let R be a subring of \mathbb{C} . An R -vertex algebra V is an R -form of the monster vertex algebra if $V \otimes_R \mathbb{C} \cong V^\natural$, where V^\natural is the monster vertex algebra over \mathbb{C} .

Definition 2.5. Let G be a group and N be an integer. An element h of G is N -regular if $(|h|, N) = 1$.

Definition 2.6. Let R be a discrete valuation ring and \mathfrak{m} the maximal ideal of R . Suppose that $k = R/\mathfrak{m}$ has prime characteristic p and that the fraction field K of R has characteristic 0. The Brauer character of a $k[h]$ -module M for a p -regular element h is defined by

$$\widetilde{\text{Tr}}(h|M) = \text{Tr}(h|\widetilde{M} \otimes_R K),$$

where \widetilde{M} is any R -free $R[h]$ -module satisfying $M \cong \widetilde{M} \otimes_{R[h]} k[h]$.

Definition 2.7. Suppose that G is a group acting on an abelian group A . Let Nr be the norm map $A \rightarrow A; a \mapsto \sum_{g \in G} g(a)$. Then, the i -th Tate cohomology $\hat{H}^i(G, A)$ is defined to be the following:

$$\hat{H}^i(G, A) = \begin{cases} H_{-i-1}(G, A) & (i \leq -2) \\ \text{Ker}(Nr)/\langle ga - a | a \in A, g \in G \rangle & (i = -1) \\ A^G/\text{Im}(Nr) & (i = 0) \\ H^i(G, A) & (i \geq 1) \end{cases},$$

where $H_i(G, A)$ is the i -th homology, and $H^i(G, A)$ is the i -th cohomology.

It has the following properties.

(1) If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is a short exact sequence of G -modules, then we have a long exact sequence

$$\dots \rightarrow \hat{H}^{i-1}(G, C) \rightarrow \hat{H}^i(G, A) \rightarrow \hat{H}^i(G, B) \rightarrow \hat{H}^i(G, C) \rightarrow \hat{H}^{i+1}(G, A) \rightarrow \dots$$

(2) $\bigoplus_{i \in \mathbb{Z}} \hat{H}^i(G, A)$ is $|G|$ -torsion module. In particular, if multiplication by $|G|$ is an isomorphism on A , then $\hat{H}^i(G, A)$ vanishes for all $i \in \mathbb{Z}$.

Moreover, if G is a cyclic group generated by g , then we have the following properties.

(3) $\hat{H}^i(g, A) \cong \hat{H}^{i+2}(g, A)$ for any $i \in \mathbb{Z}$.

(4) $\hat{H}^*(g, A) = \hat{H}^0(g, A) \oplus \hat{H}^1(g, A)$ has more or less a super version of any G -invariant algebraic structure on A . In particular, if V is a vertex algebra, then $\hat{H}^*(g, V)$ is a vertex superalgebra by the composite map

$$\hat{H}^*(g, V) \otimes \hat{H}^*(g, V) \rightarrow \hat{H}^*(g, V \otimes V) \rightarrow \hat{H}^*(g, V((z))) \rightarrow \hat{H}^*(g, V)((z)).$$

(5) If A is a finite module, then $|\hat{H}^0(g, A)| = |\hat{H}^1(g, A)|$.

Definition 2.8. Let \mathbb{H} be the upper half plane of \mathbb{C} . A Hauptmodul is a holomorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ satisfying the following conditions:

- f is invariant under some discrete subgroup Γ_f of $SL_2(\mathbb{R})$
- f generates the meromorphic function field on \mathbb{H}/Γ_f

Definition 2.9. A monster element g is Fricke if there is some $N \geq 1$ s.t. $T_g(\frac{-1}{N\tau}) = T_g(\tau)$ for any $\tau \in \mathbb{H}$.

Definition 2.10. Let R be a discrete valuation ring of mixed characteristic $(0, p)$ i.e. R has characteristic 0 and R/\mathfrak{m}_R has characteristic p . Let S be the ring of integers of a finite extension of $\text{Frac}(R)$. Then $\mathfrak{m}_S^e = \mathfrak{m}_R S$ for some positive integer e , called the ramification index. When $e = 1$, S is called an unramified extension of R . On the other hand, when $e \neq 1$, S is called a ramified extension of R .

Definition 2.11. Let R be a commutative ring with prime characteristic p . The Frobenius endomorphism $\text{Frob}_p : R \rightarrow R$ is defined by $\text{Frob}_p(r) = r^p$ for all $r \in R$.

If R is a finite field, the Frobenius endomorphism is an automorphism.

Definition 2.12. Let A be a finite abelian group, A^* be $\text{Hom}(A, \mathbb{C}^\times)$ and $f : A \rightarrow \mathbb{C}$ be a function. The discrete Fourier transform $\hat{f} : A^* \rightarrow \mathbb{C}$ is defined by $\hat{f}(\chi) = \sum_{a \in A} f(a)\chi(a)$ for any $\chi \in A^*$.

There is an inverse discrete Fourier transform, it is described by the following

$$f(a) = \frac{1}{|A|} \sum_{\chi \in A^*} \hat{f}(\chi) \overline{\chi}(a).$$

3 Modular moonshine for elements of prime order

In this section, we summarize modular moonshine for elements of prime order [3],[4].

Ryba proposed the modular moonshine conjecture. The existence of a self-dual integral form of the monster vertex algebra was not proved when he suggested it, but Carnahan proved it [5].

Theorem 3.1. (The original modular moonshine conjecture)

Suppose that an element $g \in \mathbb{M}$ has prime order p and lies in conjugacy class pA , and that V is a self-dual integral form of the monster vertex algebra. Let V^g be the set of vectors fixed by g . We consider the following vertex algebra with characteristic p :

$${}^gV = \frac{V^g/pV^g}{(V^g/pV^g) \cap (V^g/pV^g)^\perp}.$$

For ${}^gV = \bigoplus_{n \in \mathbb{Z}} {}^gV_n$, if h is a p -regular element of the centralizer $C_{\mathbb{M}}(g)$ of g , then

$$\sum_{n \in \mathbb{Z}} \widetilde{\text{Tr}}(h|{}^gV_n) q^{n-1} = T_{gh}(\tau).$$

Let G be a finite group including an element g of prime order. Borchers and Ryba noticed that this definition of gV yields $\hat{H}^0(g, V)$. They redefined gV to be $\hat{H}^*(g, V)$ and reinterpreted the conjectures in terms of Tate cohomology. To prove the conjectures, they used the fact that there are exactly 3 indecomposable finitely generated modules over the group ring $\mathbb{Z}_p[G]$ which are free as \mathbb{Z}_p -modules; there are $\mathbb{Z}_p, \mathbb{Z}_p[G]$ and I , which is the kernel of the natural map $\mathbb{Z}_p[G] \rightarrow \mathbb{Z}_p$. The Tate cohomology of these is given as following:

- $\hat{H}^0(g, \mathbb{Z}_p) = \mathbb{Z}/p\mathbb{Z}, \hat{H}^1(g, \mathbb{Z}_p) = 0.$
- $\hat{H}^0(g, \mathbb{Z}_p[G]) = \hat{H}^1(g, \mathbb{Z}_p[G]) = 0.$
- $\hat{H}^0(g, I) = 0, \hat{H}^1(g, I) = \mathbb{Z}/p\mathbb{Z}.$

By the following lemma, the Tate cohomology for any \mathbb{Z} -module is identified with the Tate cohomology for \mathbb{Z}_p -module.

Lemma 3.2. (Lemma 2.1. of [4])

If A is a free $\mathbb{Z}[1/n]$ -module acted on by a p -group G with $(p, n) = 1$, then the natural map from $\hat{H}^i(G, A)$ to $\hat{H}^i(G, A \otimes \mathbb{Z}_p)$ is an isomorphism for any $i \in \mathbb{Z}$.

The next proposition is the main tool for calculating the Brauer character of a modular moonshine vertex super algebra.

Proposition 3.3. (Proposition 2.2. of [4])

Let G be a finite group. Suppose that $g \in \text{Cent}(G)$ has prime order p and that $h \in G$ is a p -regular element. Let A be a finitely generated \mathbb{Z} -free $\mathbb{Z}[\langle g, h \rangle]$ -module. Then $\hat{H}^*(g, A) = \hat{H}^0(g, A) - \hat{H}^1(g, A)$ is a virtual representation of $\langle h \rangle$, and

$$\widetilde{\text{Tr}}(h|\hat{H}^*(g, A)) = \text{Tr}(gh|A).$$

We have the following assertion by applying Proposition 3.3 to an element of the monster group and the monster vertex algebra.

Corollary 3.4. Suppose that g is an element of \mathbb{M} of odd prime order. Let V be an integral form of the monster vertex algebra. If $h \in C_{\mathbb{M}}(g)$ is p -regular, then $\sum_{n \in \mathbb{Z}} \widetilde{\text{Tr}}(h|\hat{H}^*(g, V_n))q^{n-1}$ is equal to the McKay-Thompson series $T_{gh}(\tau)$. In particular, it is a Hauptmodul for some genus 0 subgroup of $SL_2(\mathbb{R})$.

Borcherds and Ryba proved that $\hat{H}^1(g, V) = 0$ for any Fricke element g of odd prime order. There are two main methods for proving $\hat{H}^1(g, V) = 0$.

First, when g has small order, we can transfer a vanishing statement from V_{Λ} , using a decomposition of V into submodules isomorphic to pieces of V_{Λ} [4].

Lemma 3.5. (Lemma 4.1. of [4])

If A is a permutation module of a finite group G over a ring R with no $|G|$ torsion, then $\hat{H}^{-1}(G, A) = 0$.

(A free R -module A acted on by a group G is a permutation module if it has a basis which is closed under the action of G .)

Lemma 3.6. (Lemma 4.2. of [4])

The set of permutation modules is closed under taking sums, tensor products, and symmetric powers.

Lemma 3.7. (Lemma 4.5. and Theorem 4.6. of [4])

If g is an element of odd prime order in M_{24} viewed as a subgroup of $\text{Aut}(\Lambda)$, then $\hat{H}^1(g, \Lambda) = 0$ and $\hat{H}^1(g, V_{\Lambda}) = 0$.

Theorem 3.8. (Theorem 4.7. of [4])

Suppose that g is an element of the monster of type $3A, 3C, 5A, 7A, 11A$, or $23A$. Then $\hat{H}^1(g, V) = 0$.

Remark 3.9. Similarly, for an element g of the monster type $2A$, they proved $\hat{H}^1(g, V) = 0$ by showing that $\hat{H}^1(g, \Lambda)$ and $\hat{H}^1(g, V_{\Lambda})$ vanish. The method of proof is same as the case of odd prime order, but slightly more complicated.

Borcherds and Ryba don't use the assumption that g has prime order in proofs of Lemma 3.7 and Theorem 3.8. However, this assumption is applied to use Lemma 3.2. Hence, we will show that Lemma 3.2 holds without the assumption that G is a p -group, and that $\hat{H}^1(g, V) = 0$ for an element g of the monster of type $15A$ or $21A$ in Section 4.

Second, when g has large order, it suffices to show that the coefficients $c_g^-(n) = \dim \hat{H}^0(g, V_n) + \dim \hat{H}^1(g, V_n)$ are equal to the coefficients $c_g^+(n) = \text{Tr}(g|V_n) = \dim \hat{H}^0(g, V_n) - \dim \hat{H}^1(g, V_n)$ [3]. This was proved by combining Hodge theory, an improved "no-ghost" theorem, and some explicit computation.

Borcherds proved that $\widetilde{\text{Tr}}(h|\hat{H}^i(g, V)) = \sum_{n \in \mathbb{Z}} \widetilde{\text{Tr}}(h|\hat{H}^i(g, V_n))q^{n-1}$ ($i = 0, 1$) is a linear combination of Hauptmoduls for any non-Fricke element g of odd prime order [3],[4].

Theorem 3.10. (Theorem 5.3. of [4])

If g is an element of the monster of type $2B$, then $\hat{H}^0(g, V_n)$ vanishes if n is odd, and $\hat{H}^1(g, V_n)$ vanishes if n is even.

Theorem 3.11. (Theorem 6.1. of [3])

If g is an element of the monster of type $3B, 5B, 7B$, and $13B$ and σ is the element of order 2 generating the center of $C_{\mathbb{M}}(g)/O_p(C_{\mathbb{M}}(g))$, then $O_p(C_{\mathbb{M}}(g))$ acts trivially on ${}^9V = \hat{H}^0(g, V) \oplus \hat{H}^1(g, V)$ and σ fixes $\hat{H}^0(g, V)$ and acts as -1 on $\hat{H}^1(g, V)$.

We have the following complete description of $\widetilde{\text{Tr}}(h|\hat{H}^i(g, V))$ ($i = 0, 1$).

$$\widetilde{\text{Tr}}(h|\hat{H}^0(g, V)) = \begin{cases} T_{gh}(\tau) & (g \in pA, 3C) \\ \frac{T_{gh}(\tau) + T_{gh}(\tau + 1/2)}{2} & (g \in 2B) \\ \frac{T_{gh}(\tau) + T_{gh\sigma}(\tau)}{2} & (g \in pB, 2|(p-1)) \end{cases},$$

$$\widetilde{\text{Tr}}(h|\hat{H}^1(g, V)) = \begin{cases} 0 & (g \in pA, 3C) \\ \frac{-T_{gh}(\tau) + T_{gh}(\tau + 1/2)}{2} & (g \in 2B) \\ \frac{-T_{gh}(\tau) + T_{gh\sigma}(\tau)}{2} & (g \in pB, 2|(p-1)) \end{cases},$$

where $\sigma \in C_{\mathbb{M}}(g)/O_p(C_{\mathbb{M}}(g))$ is an involution which acts as 1 on $\hat{H}^0(g, V)$ and as -1 on $\hat{H}^1(g, V)$.

4 Tate cohomology and ramified extensions

Let R be a discrete valuation ring. Let $g \in G$ be an element of order N , $h \in G$ be an N -regular element and A be an $R[g]$ -module.

Proposition 4.1. *If a commutative ring homomorphism $R \rightarrow S$ is flat, then $\hat{H}^*(g, A \otimes_R S) \cong \hat{H}^*(g, A) \otimes_R S$.*

Proof. Let Nr be a norm map as in Definition 2.7. Then, $\hat{H}^0(g, A) \cong \text{Ker}(g-1)/\text{Im}(Nr)$. We have an exact sequence of R -modules $0 \rightarrow \text{Im}(Nr) \rightarrow \text{Ker}(g-1) \rightarrow \hat{H}^0(g, A) \rightarrow 0$. If $R \rightarrow S$ is flat, the sequence of S -modules $0 \rightarrow \text{Im}(Nr) \otimes S \rightarrow \text{Ker}(g-1) \otimes S \rightarrow \hat{H}^0(g, A) \otimes S \rightarrow 0$ is exact. We consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Im}(Nr \otimes id_S) & \longrightarrow & \text{Ker}((g-1) \otimes id_S) & \longrightarrow & \hat{H}^0(g, A \otimes S) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \psi_0 \\ 0 & \longrightarrow & \text{Im}(Nr) \otimes S & \longrightarrow & \text{Ker}(g-1) \otimes S & \longrightarrow & \hat{H}^0(g, A) \otimes S \longrightarrow 0 \end{array}$$

Then, we have $\text{Im}(Nr \otimes id_S) \cong \text{Im}(Nr) \otimes S$ and $\text{Ker}((g-1) \otimes id_S) \cong \text{Ker}(g-1) \otimes S$. Indeed, we can take the isomorphisms $Nr \otimes id_S(x \otimes s) \mapsto Nr(x) \otimes s$ and $x \otimes s \mapsto x \otimes s$, respectively. Also, the above diagram is commutative. Hence, by the Five Lemma, ψ_0 is an isomorphism.

Like the degree 0 case, we have an exact sequence $0 \rightarrow \text{Im}(g-1) \rightarrow \text{Ker}(Nr) \rightarrow \hat{H}^1(g, A) \rightarrow 0$. As well above, we consider the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Im}((g-1) \otimes id_S) & \longrightarrow & \text{Ker}(Nr \otimes id_S) & \longrightarrow & \hat{H}^1(g, A \otimes S) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \psi_1 \\ 0 & \longrightarrow & \text{Im}(g-1) \otimes S & \longrightarrow & \text{Ker}(Nr) \otimes S & \longrightarrow & \hat{H}^1(g, A) \otimes S \longrightarrow 0 \end{array}$$

Then, we have $\text{Im}((g-1) \otimes id_S) \cong \text{Im}(g-1) \otimes S$ and $\text{Ker}(Nr \otimes id_S) \cong \text{Ker}(Nr) \otimes S$. Indeed, we can take the isomorphisms $(g-1) \otimes id_S(x \otimes s) \mapsto (gx-x) \otimes s$ and $x \otimes s \mapsto x \otimes s$, respectively. By the Five Lemma, ψ_1 is an isomorphism. \square

Using the above proposition, we will prove that the 1st Tate cohomology for an element of the monster type 15A or 21A vanishes. To do this, we will show the following lemma.

Lemma 4.2. *If A is a free $\mathbb{Z}[1/q]$ -module acted on by a group G with $|G| = p^k q$ and $(p, q) = 1$, then the natural map from $\hat{H}^i(G, A)$ to $\hat{H}^i(G, A \otimes \mathbb{Z}_p)$ is an isomorphism for any $i \in \mathbb{Z}$.*

Proof. We show that multiplication by $|G|$ is an isomorphism on $\mathbb{Z}_p/\mathbb{Z}[1/q]$. For any $p^l \frac{a}{b} \in \text{Ker}(|G|)$, where l is a non-negative integer and p, a, b are pairwise coprime, we have $|G| \cdot p^l \frac{a}{b} = p^{l+k} \frac{aq}{b} \in \mathbb{Z}[1/q]$. It follows that b is a product of divisors of q and then $p^l \frac{a}{b} \in \mathbb{Z}[1/q]$. Hence, $|G|$ is injective. For any $\sum_{i=0}^{\infty} a_i p^i \in \mathbb{Z}_p/\mathbb{Z}$, where $a_i \in \{0, \dots, p-1\}$, we have $\sum_{i=0}^{\infty} a_i p^i \equiv \sum_{i=k}^{\infty} a_i p^i \pmod{\mathbb{Z}}$. Hence, we have $|G| \cdot \frac{1}{q} \sum_{i=0}^{\infty} a_{k+i} p^i = \sum_{i=k}^{\infty} a_i p^i$ and then $|G|$ is surjective on \mathbb{Z}_p/\mathbb{Z} . By the surjectivity of $\mathbb{Z}_p/\mathbb{Z} \rightarrow \mathbb{Z}_p/\mathbb{Z}[1/q]$, $|G|$ is surjective on $\mathbb{Z}_p/\mathbb{Z}[1/q]$. By property (2) of Tate cohomology, $\hat{H}^i(G, A \otimes \mathbb{Z}_p/\mathbb{Z}[1/q]) = 0$ for all $i \in \mathbb{Z}$. For the exact sequence

$$0 \rightarrow A \rightarrow A \otimes \mathbb{Z}_p \rightarrow A \otimes (\mathbb{Z}_p/\mathbb{Z}[1/q]) \rightarrow 0,$$

taking the Tate cohomology, we have a long exact sequence because of property (1) of Tate cohomology. It follows that $\hat{H}^i(G, A) \cong \hat{H}^i(G, A \otimes \mathbb{Z}_p)$ for any $i \in \mathbb{Z}$. \square

Lemma 4.3. *If g is an element of odd order in M_{24} viewed as a subgroup of $\text{Aut}(\Lambda)$, then $\hat{H}^1(g, \Lambda) = 0$ and $\hat{H}^1(g, V_\Lambda) = 0$.*

Proof. Let $|g| = p^k q$, where p is a prime and $(p, q) = 1$. By Proposition 4.1 and Lemma 4.2, we have $\hat{H}^1(g, \Lambda) \otimes \mathbb{Z}[1/q] \cong \hat{H}^1(g, \Lambda \otimes \mathbb{Z}[1/q]) \cong \hat{H}^1(g, \Lambda \otimes_{\mathbb{Z}[1/q]} \mathbb{Z}_p)$. By the same argument as the proof of Lemma 4.5 of [4], $\hat{H}^1(g, \Lambda \otimes_{\mathbb{Z}[1/q]} \mathbb{Z}_p) = 0$. Hence, $\hat{H}^1(g, \Lambda) = 0$.

Similarly, we have $\hat{H}^1(g, V_\Lambda) \otimes \mathbb{Z}[1/q] \cong \hat{H}^1(g, V_\Lambda \otimes_{\mathbb{Z}[1/q]} \mathbb{Z}_p)$. We can decompose $V_\Lambda \otimes \mathbb{Z}_p$ into the following:

$$V_\Lambda \otimes \mathbb{Z}_p = \bigoplus_{\text{orbits}_{\langle g \rangle \lambda}} \left(\bigoplus_{g^r \lambda \in \langle g \rangle \lambda} V_{\Lambda, g^r \lambda} \otimes \mathbb{Z}_p \right),$$

where $\bigoplus_{g^r \lambda \in \langle g \rangle \lambda} V_{\Lambda, g^r \lambda} \otimes \mathbb{Z}_p \cong \mathbb{Z}_p[\langle g \rangle / \text{Stab}_{\langle g \rangle}(\lambda)] \otimes_{\mathbb{Z}_p} (V_{\Lambda, 0} \otimes \mathbb{Z}_p)$ as $\mathbb{Z}_p[g]$ -modules.

By the proof of Theorem 4.6 of [4], $V_{\Lambda, 0} \otimes \mathbb{Z}_p$ is a permutation module. By Lemma 3.6, $\bigoplus_{g^r \lambda \in \langle g \rangle \lambda} V_{\Lambda, g^r \lambda} \otimes \mathbb{Z}_p$ is a permutation module. By Lemma 3.6 and Lemma 3.5, $V_\Lambda \otimes \mathbb{Z}_p$ is a permutation module and we have $\hat{H}^1(g, V_\Lambda) = 0$. \square

Theorem 4.4. *Suppose that g is an element of the monster of type 15A or 21A. Then $\hat{H}^1(g, V) = 0$.*

Proof. In the same way as the proof of Theorem 4.7 of [4], this statement follows from Lemma 4.3. \square

Let p be a prime factor of N . Now, we consider a ramified extension $R_p = \mathbb{Z}_p[\zeta_{N|h}]$ of \mathbb{Z}_p . We define rank 1 R -free $R_p[\langle g, h \rangle]$ -modules by

$$R_{n,m} = \{cv_{n,m} | c \in R_p, gv_{n,m} = \zeta_N^n v_{n,m}, hv_{n,m} = \zeta_{|h|}^m v_{n,m}\}.$$

Any $R_p[\langle g, h \rangle]$ -module that is R_p -free of finite rank is an extension of $\{R_{n,m}\}_{0 \leq n \leq N-1, 0 \leq m \leq |h|-1}$. Hence, it suffices to consider the Tate cohomology of $R_{n,m}$.

Lemma 4.5. For any $0 \leq n \leq N-1, 0 \leq m \leq |h|-1$,

$$\hat{H}^0(g, R_{n,m}) = \begin{cases} R_p/NR_p & (n=0) \\ 0 & (n \neq 0) \end{cases},$$

$$\hat{H}^1(g, R_{n,m}) = \begin{cases} 0 & (n=0) \\ R_p/(1-\zeta_N^n)R_p & (n \neq 0) \end{cases},$$

where h acts as $\zeta_{|h|}^m$ on R_p/NR_p and $R_p/(1-\zeta_N^n)R_p$.

Proof. $R_{0,m}^g = R_{0,m}, R_{n,m}^g = 0$ ($n \neq 0$). Hence, $\hat{H}^0(g, R_{0,m}) = R_p/NR_p$,

$\hat{H}^0(g, R_{n,m}) = 0$ ($n \neq 0$), because the norm map $R_{n,m} \rightarrow R_{n,m}$ is $v_{n,m} \mapsto \sum_{i=0}^{N-1} g^i v_{n,m}$.

By $\sum_{i=0}^{N-1} g^i v_{0,m} = Nv_{0,m}$, the kernel of the norm map on $R_{0,m}$ is 0. Hence, $\hat{H}^1(g, R_{0,m}) = 0$.
By $\sum_{i=0}^{N-1} g^i v_{n,m} = \sum_{i=0}^{N-1} \zeta_N^{ni} v_{n,m} = 0$ ($n \neq 0$), the kernel of the norm map on $R_{n,m}$ is $R_{n,m}$.
Therefore, by the definition of Tate cohomology, $\hat{H}^1(g, R_{n,m}) = R_p/(1-\zeta_N^n)R_p$ ($n \neq 0$). \square

5 p -Brauer character

Let p be a prime factor of N . Let R be a discrete valuation ring of mixed characteristic $(0, p)$, $v : R \rightarrow \mathbb{Z}_{\geq 0} \cup \{\infty\}$ be a surjective valuation on R and $K = \text{Frac}(R)$.

We can't use the Brauer character for $R_p/(1-\zeta_N^n)R_p$ because this is not a vector space over a field. To solve this, we will define a p -Brauer character.

Definition 5.1. Let M be a finite length $R[h]$ -module. Suppose that M has composition series $0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_n = M$. We define the p -Brauer character of M for an N -regular element h by

$$\widetilde{\text{Tr}}_p(h|M) = \frac{1}{v(p)} \sum_{i=1}^n \text{Tr}(h|\widetilde{M}_i \otimes K),$$

where \widetilde{M}_i is an R -free $R[h]$ -module satisfying $M_i/M_{i-1} \cong \widetilde{M}_i \otimes_{R[h]} R/\mathfrak{m}_R[h]$.

Lemma 5.2. This definition does not depend on our choices of M_i and \widetilde{M}_i .

Proof. For $M_i/M_{i-1} \cong \widetilde{M}_i \otimes_{R[h]} R/\mathfrak{m}_R[h] \cong \widetilde{M}'_i \otimes_{R[h]} R/\mathfrak{m}_R[h]$, we have $\text{Tr}(h|\widetilde{M}_i \otimes K) = \widetilde{\text{Tr}}(h|M_i/M_{i-1}) = \text{Tr}(h|\widetilde{M}'_i \otimes K)$. By the Jordan-Hölder theorem, the composition series of M is unique up to permutation and isomorphism. That is, for any two composition series $0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_n = M, 0 = M'_0 \subsetneq M'_1 \subsetneq \cdots \subsetneq M'_{n'} = M$, we have $n = n'$ and some permutation σ of $\{1, \dots, n\}$ s.t. $M_i/M_{i-1} \cong M'_{\sigma(i)}/M'_{\sigma(i)-1}$. Hence, $\sum_{i=1}^n \widetilde{\text{Tr}}(h|M_i/M_{i-1}) = \sum_{i=1}^n \widetilde{\text{Tr}}(h|M'_{\sigma(i)}/M'_{\sigma(i)-1}) = \sum_{i=1}^n \widetilde{\text{Tr}}(h|M'_i/M'_{i-1})$. Therefore the p -Brauer character is well-defined. \square

Note that it does depend on a choice of a prime factor of N .

The p -Brauer character has the following properties.

Lemma 5.3. The p -Brauer character is additive on short exact sequences.

(“additive” means that for any short exact sequence $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$, $\widetilde{\text{Tr}}_p$ satisfies $\widetilde{\text{Tr}}_p(h|M') = \widetilde{\text{Tr}}_p(h|M) + \widetilde{\text{Tr}}_p(h|M'')$.)

Proof. Let $0 \rightarrow M \xrightarrow{\varphi} M' \xrightarrow{\psi} M'' \rightarrow 0$ be an exact sequence of finite length $R[h]$ -modules. Suppose that M has length m and a composition series $0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_n = M$. We use analogous notation for M' and M'' . Then the length m' of M' is $m + m''$ and

$$0 = \varphi(M_0) \subsetneq \varphi(M_1) \subsetneq \cdots \subsetneq \varphi(M_m) = \psi^{-1}(M_0'') \subsetneq \psi^{-1}(M_1'') \subsetneq \cdots \subsetneq \psi^{-1}(M_{m''}'') = M'$$

is a composition series of M' . By injectivity of φ , $M_i \cong \varphi(M_i)$ for any $i \in \{0, \dots, m\}$ and by surjectivity of ψ , $\psi^{-1}(M_j'')/\text{Ker}(\psi) \cong M_j''$ for any $j \in \{0, \dots, m''\}$. Hence, we have $\varphi(M_i)/\varphi(M_{i-1}) \cong M_i/M_{i-1}$ and

$$\psi^{-1}(M_j'')/\psi^{-1}(M_{j-1}'') \cong \left(\psi^{-1}(M_j'')/\text{Ker}(\psi) \right) / \left(\psi^{-1}(M_{j-1}'')/\text{Ker}(\psi) \right) \cong M_j''/M_{j-1}''.$$

Therefore,

$$\begin{aligned} \widetilde{\text{Tr}}_p(h|M') &= \frac{1}{v(p)} \sum_{i=1}^m \widetilde{\text{Tr}}(h|\varphi(M_i)/\varphi(M_{i-1})) + \frac{1}{v(p)} \sum_{j=1}^{m''} \widetilde{\text{Tr}}(h|\psi^{-1}(M_j'')/\psi^{-1}(M_{j-1}'')) \\ &= \frac{1}{v(p)} \sum_{i=1}^m \widetilde{\text{Tr}}(h|M_i/M_{i-1}) + \frac{1}{v(p)} \sum_{j=1}^{m''} \widetilde{\text{Tr}}(h|M_j''/M_{j-1}'') \\ &= \widetilde{\text{Tr}}_p(h|M) + \widetilde{\text{Tr}}_p(h|M''). \end{aligned}$$

□

Lemma 5.4. *Let S be the ring of integers of a finite extension of $\text{Frac}(R)$ and A be a finite length $R[h]$ -module. Then $\widetilde{\text{Tr}}_p(h|A \otimes_R S) = \widetilde{\text{Tr}}_p(h|A)$.*

Proof. It suffices to consider the case A is a simple $R[h]$ -module on which h acts as $\zeta_{|h|}^m$. Now, $\widetilde{\text{Tr}}_p(h|A) = \frac{1}{v_R(p)} \text{Tr}(h|\widetilde{A} \otimes \text{Frac}(R))$, where v_R is a discrete valuation on R , \widetilde{A} is an R -free $R[h]$ -module satisfying $A \cong \widetilde{A} \otimes_{R[h]} R/\mathfrak{m}_R[h]$ and \mathfrak{m}_R is a maximal ideal of R . We have $A \otimes_R S \cong \widetilde{A} \otimes_{R[h]} R/\mathfrak{m}_R[h] \otimes_R S \cong \widetilde{A} \otimes_{R[h]} S/\mathfrak{m}_R S[h]$.

Let v_S be a discrete valuation on S and \mathfrak{m}_S be a maximal ideal of S . If S is an unramified extension, then $\mathfrak{m}_S = \mathfrak{m}_R S$ and $v_S(p) = v_R(p)$. Hence,

$$\widetilde{\text{Tr}}_p(h|A \otimes_R S) = \frac{1}{v_S(p)} \text{Tr}(h|\widetilde{A} \otimes_R S \otimes \text{Frac}(S)) = \frac{1}{v_R(p)} \text{Tr}(h|\widetilde{A} \otimes \text{Frac}(R)) = \widetilde{\text{Tr}}_p(h|A).$$

If S is a ramified extension, then there is a positive integer $k > 1$ s.t. $\mathfrak{m}_S^k = \mathfrak{m}_R S$ and $v_S(p) = kv_R(p)$. Hence, we have $A \otimes_R S = \widetilde{A} \otimes_{R[h]} S/\mathfrak{m}_S^k[h]$ and a composition series $0 \subsetneq \widetilde{A} \otimes S/\mathfrak{m}_S \subsetneq \cdots \subsetneq \widetilde{A} \otimes S/\mathfrak{m}_S^k$. Its composition factors are all isomorphic to $\widetilde{A} \otimes_{R[h]} S/\mathfrak{m}_S[h]$. Therefore,

$$\widetilde{\text{Tr}}_p(h|A \otimes_R S) = \frac{1}{v_S(p)} \sum_{i=1}^k \text{Tr}(h|\widetilde{A} \otimes_R S \otimes \text{Frac}(S)) = \frac{1}{kv_R(p)} k \text{Tr}(h|\widetilde{A} \otimes \text{Frac}(R)) = \widetilde{\text{Tr}}_p(h|A).$$

□

By Proposition 4.1 and 5.4, any finitely generated \mathbb{Z} -free $\mathbb{Z}[\langle g, h \rangle]$ -module base changes to an $R_p[\langle g, h \rangle]$ -module that is R_p -free of finite rank and the p -Brauer character of the Tate cohomology is unchanged. By Lemma 5.3, it suffices to consider the p -Brauer character for the Tate cohomology of $R_{n,m}$.

We will calculate the Tate cohomology of $R_{n,m}$ by using the p -Brauer character. To do this, we use the following lemma:

Lemma 5.5. (Chapter III, Lemma 3 of [6])

Let $q = p^k$ for a positive integer k . Then $pR = (1 - \zeta_q)^{\varphi(q)} R$.

For a positive integer s and $R = \mathbb{Z}_p[\zeta_{p^s}]$, a discrete valuation $v(p)$ is given by $\varphi(p^s)$ and maximal ideal of R is $(1 - \zeta_{p^s})R$.

Lemma 5.6. *Suppose that the prime factorization of N is $\prod p_i^{n_i}$. Then,*

$$\widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, R_{n,m})) = \begin{cases} n_i \zeta_{|h|}^m & (n = 0) \\ \frac{-\zeta_{|h|}^m}{\varphi(p_i^l)} & (|g^n| = p_i^l, 1 \leq l \leq n_i) \\ 0 & (\text{otherwise}) \end{cases}.$$

Proof. Now, h acts as $\zeta_{|h|}^m$ on $\hat{H}^*(g, R_{n,m})$ and $v(p) = \varphi(p_i^{n_i})$. By $\widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, R_{n,m})) = \widetilde{\text{Tr}}_{p_i}(h|\hat{H}^0(g, R_{n,m})) - \widetilde{\text{Tr}}_{p_i}(h|\hat{H}^1(g, R_{n,m}))$ and Lemma 4.5, it suffices to calculate the p_i -Brauer characters of R_{p_i}/NR_{p_i} and $R_{p_i}/(1 - \zeta_N^n)R_{p_i}$.

When $n = 0$, by Lemmas 4.5 and 5.5, $R_{p_i}/NR_{p_i} \cong R_{p_i}/p_i^{n_i}R_{p_i} \cong R_{p_i}/(1 - \zeta_{p_i^{n_i}})^{n_i \varphi(p_i^{n_i})}R_{p_i}$. This module has a composition series $0 \subsetneq R_{p_i}/(1 - \zeta_{p_i^{n_i}})R_{p_i} \subsetneq \cdots \subsetneq R_{p_i}/(1 - \zeta_{p_i^{n_i}})^{n_i \varphi(p_i^{n_i})}R_{p_i}$ and its composition factors are all isomorphic to $R_{p_i}/(1 - \zeta_{p_i^{n_i}})R_{p_i}$. Hence, $\widetilde{\text{Tr}}_{p_i}(h|R_{p_i}/NR_{p_i}) = n_i \zeta_{|h|}^m$.

When $|g^n| = p_i^l$, ζ_N^n is a p_i^l -th primitive root of unity. By Lemma 5.5, we have $(1 - \zeta_{p_i^l})^{\varphi(p_i^l)} = (1 - \zeta_{p_i^{n_i}})^{\varphi(p_i^{n_i})}$. Hence, by Lemma 4.5, we have $R_{p_i}/(1 - \zeta_{p_i^l})R_{p_i} \cong R_{p_i}/(1 - \zeta_{p_i^{n_i}})^{\varphi(p_i^{n_i})/\varphi(p_i^l)}R_{p_i}$. This module has the length $\varphi(p_i^{n_i})/\varphi(p_i^l)$ and its composition factors are all isomorphic to $R_{p_i}/(1 - \zeta_{p_i^{n_i}})R_{p_i}$. Calculating the p_i -Brauer character, we have $\widetilde{\text{Tr}}_{p_i}(h|R_{p_i}/(1 - \zeta_{p_i^l})R_{p_i}) = 1/\varphi(p_i^l)$.

When $|g^n| = p_i^k q$, where $q \neq 1, (q, p_i) = 1$ and $0 \leq k \leq n_i$, ζ_N^n is a $p_i^k q$ -th primitive root of unity. Let π be a surjective homomorphism $R_{p_i} \rightarrow \mathbb{F}_{p_i}[\zeta_N^{p_i^k q}]; \sum_{j=0}^{\infty} c_j p_i^j \mapsto c_0, \zeta_N^{p_i^k q} \mapsto \zeta_N^{p_i^k q}$. Then $\text{Ker}(\pi)$ is the maximal ideal of R_{p_i} and Frob_{p_i} on $\mathbb{F}_{p_i}[\zeta_N^{p_i^k q}]$ is an automorphism. We have $\pi(\zeta_{p_i^k q}) = \text{Frob}_{p_i}^{-k}(\zeta_{p_i^k q}) \neq 1$. Hence, $1 - \zeta_N^n \notin \text{Ker}(\pi)$. $1 - \zeta_N^n$ is a unit of R_{p_i} because every element of a local ring not included in a maximal ideal is a unit. Therefore, $R_{p_i}/(1 - \zeta_N^n)R_{p_i} = 0$. \square

We generalize Proposition 3.3 (Proposition 2.2. of [4]) using the p -Brauer character.

Theorem 5.7. *Let G be a finite group. Suppose that $g \in \text{Cent}(G)$ has order N and that $h \in G$ is an N -regular element. Let A be a finitely generated R -free $R[\langle g, h \rangle]$ -module. Then $\hat{H}^*(g, A) = \hat{H}^0(g, A) - \hat{H}^1(g, A)$ is a virtual representation of $\langle h \rangle$, and*

$$\begin{aligned} \widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, A)) &= \sum_{k=1}^{N-1} a_{k,p_i} \text{Tr}(g^k h|A), \\ a_{k,p_i} &= \begin{cases} \frac{1}{\sum_{(p_i, d)=1, d|N} \varphi(N/d)} \left(n_i - l - \frac{n_i - l - 1}{p_i} \right) & ((k, p_i^{n_i}) = p_i^l, 0 \leq l \leq n_i - 1) \\ 0 & ((k, p_i^{n_i}) = p_i^{n_i}) \end{cases}, \end{aligned}$$

where the prime factorization of N is $\prod p_i^{n_i}$.

Proof. Suppose $\widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, A)) = \sum_{k=0}^{N-1} a_{k,p_i} \text{Tr}(g^k h|A)$. It suffices to take $A = R_{n,m}$. By an inverse discrete Fourier transform and Lemma 5.6, we have

$$\begin{aligned} a_{k,p_i} &= \frac{1}{N} \sum_{b=0}^{N-1} \widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, R_{b,m})) \overline{\text{Tr}(g^k h|R_{b,m})} \\ &= \frac{1}{N} \left(n_i + \sum_{|g^b|=p_i} \frac{-1}{\varphi(p_i)} \zeta_N^{-kb} + \sum_{|g^b|=p_i^2} \frac{-1}{\varphi(p_i^2)} \zeta_N^{-kb} + \cdots + \sum_{|g^b|=p_i^{n_i}} \frac{-1}{\varphi(p_i^{n_i})} \zeta_N^{-kb} \right). \end{aligned}$$

Let $(k, p_i^{n_i}) = p_i^l$. If $s > l$, then $\sum_{|g^b|=p_i^s} \zeta_N^{-kb}$ is a sum of primitive p_i^{s-l} -th roots of unity and equals an integer multiple of the Möbius function $\mu(p_i^{s-l})$. If $s \leq l$, then we have $\sum_{|g^b|=p_i^s} \zeta_N^{-kb} = \varphi(p_i^s)$.

Hence, when $(k, p_i^{n_i}) = p_i^{n_i}$, we have $a_{k, p_i} = \frac{1}{N}(n_i - 1 - \dots - 1) = 0$.

When $(k, p_i^{n_i}) = p_i^l$, because $(N/p_i^{n_i}) = \sum_{d|(N/p_i^{n_i})} \varphi(N/p_i^{n_i}d)$, we have

$$\begin{aligned} a_{k, p_i} &= \frac{1}{N} \left(n_i - 1 - \dots - 1 + \frac{p_i^l}{\varphi(p_i^{l+1})} \right) \\ &= \frac{(n_i - l)\varphi(p_i^{l+1}) + p_i^l}{N\varphi(p_i^{l+1})} \\ &= \frac{p_i^{l+1}(n_i - l) - p_i^l(n_i - l - 1)}{p_i^{l+1}\varphi(p_i^{n_i})(N/p_i^{n_i})} \\ &= \frac{1}{\sum_{(p_i, d)=1, d|N} \varphi(N/d)} \left(n_i - l - \frac{n_i - l - 1}{p_i} \right). \end{aligned}$$

□

Corollary 5.8. *Suppose that g is an element of \mathbb{M} of order $N = \prod p_i^{n_i}$. Let V be an integral form of the monster vertex algebra. If $h \in C_{\mathbb{M}}(g)$ is N -regular, then $\sum_{n \in \mathbb{Z}} \widetilde{\text{Tr}}_{p_i}(h|\hat{H}^*(g, V_n \otimes \mathbb{Z}_{p_i}))q^{n-1}$ is a linear combination of Hauptmoduls and is denoted by $\sum_{d|N} a_{d, p_i} \varphi(N/d) T_{g^d h}(\tau)$.*

Proof. By Theorem 5.7 and the fact that McKay-Thompson series depend only on conjugacy classes. □

We propose the following conjecture about the graded p -Brauer character for Tate cohomology;

Conjecture 5.9. *For any element $g \in \mathbb{M}$ of order N with prime factor p and N -regular element $h \in C_{\mathbb{M}}(g)$, $\sum_{n \in \mathbb{Z}} \widetilde{\text{Tr}}_p(h|\hat{H}^i(g, V_n \otimes \mathbb{Z}_p))q^{n-1}$ ($i = 0, 1$) is a linear combination of Hauptmoduls.*

From Corollary 5.8, we can give a counterexample to a conjecture of Borchers about vanishing of Tate cohomology $\hat{H}^1(g, V)$.

Theorem 5.10. *Let V be an integral form of the monster vertex algebra. Then there exists a Fricke element g s.t. $\hat{H}^1(g, V) \neq 0$.*

Proof. Let g be an element of the monster in class 8A. Then g is a Fricke element. By Corollary 5.8, we have

$$\sum_{n \in \mathbb{Z}} \text{length}(\hat{H}^*(g, V_n))q^{n-1} = a_{1,2}\varphi(8)T_g(\tau) + a_{2,2}\varphi(4)T_{g^2}(\tau) + a_{4,2}\varphi(2)T_{g^4}(\tau).$$

Comparing coefficients of q^2 , we have $\text{length}(\hat{H}^0(g, V_3)) - \text{length}(\hat{H}^1(g, V_3)) = -256$. This means $\hat{H}^1(g, V_3) \neq 0$. Hence, $\hat{H}^1(g, V) \neq 0$. □

We propose a weaker version of Borchers's conjecture about vanishing of Tate cohomology $\hat{H}^1(g, V)$.

Conjecture 5.11. *Let V be an self-dual integral form of the monster vertex algebra. If g is a Fricke element such that g^i is a Fricke element for any $i > 0$, then $\hat{H}^1(g, V) = 0$.*

Acknowledgements

I would like to thank Scott Carnahan for many helpful comments and advice.

References

- [1] R. E. Borcherds, “Vertex algebras, Kac-Moody algebras, and the monster”, Proc. Natl. Acad. Sci. USA. Vol. 83, 3068-3071, (1986).
- [2] R. E. Borcherds, “Monstrous moonshine and monstrous Lie superalgebras”, Invent. Math. 109, 405-444, (1992).
- [3] R. E. Borcherds, “Modular Moonshine III”, Duke Math. Journal Vol. 93 No. 1, 129-154, (1998).
- [4] R. E. Borcherds, A.J. E. Ryba, “Modular Moonshine II”, Duke Math. Journal Vol. 83 No. 2, 435-459, (1996).
- [5] S. Carnahan, “A self-dual integral form of the moonshine module”, SIGMA Vol. 15, 030, 36 pages, (2019).
- [6] J. M. S. Cassels and A. Fröhlich, “Algebraic Number Theory”, Academic Press, (1967).
- [7] J. H. Conway, S. Norton, “Monstrous moonshine”, Bull. London. Math. Soc. 11, 308-339, (1979).
- [8] I. B. Frenkel, J. Lepowsky, A. Meurman, “Vertex operator algebras and monster”, Academic Press, (1988).
- [9] J. Lepowsky, H. Li, “Introduction to Vertex Operator Algebras and Their Representations”, Progress in Mathematics, Vol. 227, Birkhäuser, Boston, (2004).