

MULTIPLICATIVE FUNCTIONS WITH

$$f(p + q - n_0) = f(p) + f(q) - f(n_0)$$

POO-SUNG PARK

ABSTRACT. Let n_0 be 1 or 3. If a multiplicative function f satisfies $f(p + q - n_0) = f(p) + f(q) - f(n_0)$ for all primes p and q , then f is the identity function $f(n) = n$ or a constant function $f(n) = 1$.

1. INTRODUCTION

In 2016 Chen, Fang, Yuan, and Zheng showed that if a multiplicative function f satisfies $f(p + q + n_0) = f(p) + f(q) + f(n_0)$ with $1 \leq n_0 \leq 10^6$ then f is the identity function provided $f(p_0) \neq 0$ for some prime p_0 [1]. This is a variation of Spiro's paper in 1992 in which she dealt multiplicative functions satisfying $f(p + q) = f(p) + f(q)$ [7]. She called the set of primes an *additive uniqueness set* for multiplicative functions f with $f(p_0) \neq 0$ for some prime p_0 .

A natural question follows about n_0 being negative for Chen et al.'s paper. It is natural to consider the condition $f(p + q - n_0) = f(p) + f(q) - f(n_0)$ with $n_0 = 1, 2, 3$ because a multiplicative function is defined on positive integers.

The author already studied a multiplicative function satisfying $f(p + q - 2) = f(p) + f(q) - f(2)$, which also yields that the set of numbers 1 less than primes is an additive uniqueness set for multiplicative functions [5].

In this article we classify multiplicative functions satisfying $f(p + q - n_0) = f(p) + f(q) - f(n_0)$ with $n_0 = 1, 3$. For consistency we state the classification for $n_0 = 2$ as well.

Theorem 1.1. *If a multiplicative function f satisfies $f(p + q - 1) = f(p) + f(q) - f(1)$ for all primes p and q , then f is the identity function $f(n) = n$ or a constant function $f(n) = 1$.*

Theorem 1.2 ([5]). *If a multiplicative function f satisfies $f(p+q-2) = f(p) + f(q) - f(2)$ for all primes p and q , then f is the identity function $f(n) = n$, a constant function $f(n) = 1$, or $f(n) = 0$ for $n \geq 2$ unless n is odd and squareful.*

Theorem 1.3. *If a multiplicative function f satisfies $f(p+q-3) = f(p)+f(q)-f(3)$ for all primes p and q , then f is the identity function $f(n) = n$ or a constant function $f(n) = 1$.*

Theorem 1.2 for $n_0 = 2$ has one more option. We give a proof for Theorem 1.3. The proof of Theorem 1.1 is similar and the proof of Theorem 1.2 is given in [5, §4].

2. LEMMAS

Lemma 2.1. *Assume a multiplicative function f satisfies $f(p+q-3) = f(p) + f(q) - f(3)$ for all primes p and q . Then, $f(n) = 1$ or $f(n) = n$ for $n = 2, 3, 5, 7$, and 11.*

Proof. Note that $f(1) = 1$ and the equalities

$$\begin{aligned} f(1) &= f(2+2-3) = f(2) + f(2) - f(3), \\ f(7) &= f(5+5-3) = f(5) + f(5) - f(3), \\ f(10) &= f(2)f(5) \\ &= f(11+2-3) = f(11) + f(2) - f(3), \\ f(11) &= f(7+7-3) = f(7) + f(7) - f(3), \\ f(15) &= f(3)f(5) \\ &= f(11+7-3) = f(11) + f(7) - f(3). \end{aligned}$$

Let $a = f(2)$, $b = f(3)$, $c = f(5)$, $d = f(7)$, $e = f(11)$. Then,

$$\begin{aligned} (1) \quad & 1 = 2a - b \\ (2) \quad & d = 2c - b \\ (3) \quad & ac = e + a - b \\ (4) \quad & e = 2d - b \\ (5) \quad & bc = e + d - b. \end{aligned}$$

The equation (3) becomes

$$ac = 4c - 7a + 4$$

by the equations (1), (2), and (4). Also, the equation (5) becomes

$$2ac = 7c - 10a + 5.$$

So, $c = 4a - 3$ and we obtain an equation $a^2 - 3a + 2 = 0$.

Thus, $a = 1$ or $a = 2$ and it follows that

$$\begin{aligned} f(2) = 1, \quad f(3) = 1, \quad f(5) = 1, \quad f(7) = 1, \quad f(11) = 1; \\ f(2) = 2, \quad f(3) = 3, \quad f(5) = 5, \quad f(7) = 7, \quad f(11) = 11. \end{aligned}$$

□

Lemma 2.2. *The results in Lemma 2.1 can be extended up to n odd and $n < 10^{10}$.*

Proof. We use induction. Let n be odd and $11 < n < 10^{10}$.

If n is prime, then $n = 6k - 1$ or $n = 6k + 1$. Suppose $n = 6k - 1$. Note that

$$f(n+4) = f(6k+3) = f(n+7-3) = f(n) + f(7) - f(3).$$

Since $6k+3$ can be factored into the product of two smaller integers, $f(6k+3) = 1$ or $f(6k+3) = 6k+3$ by induction hypothesis. Thus, $f(n) = 1$ or $f(n) = n$ when $n = 6k - 1$ is prime.

Similarly, if n is a prime of the form $6k + 1$, then $f(n) = 1$ or $f(n) = n$ by

$$f(n+2) = f(6k+3) = f(n+5-3) = f(n) + f(5) - f(3).$$

If n is not a prime, n is either a product of two relatively prime integers or a power of a prime. The first case is easy by the multiplicity of f . So the second case remains.

Now, assume that n is a power of a prime with exponent ≥ 2 . Then, $n + 3$ is even and can be written as a sum of two primes p and q with $5 \leq p, q < n$ by the numerical verification of the Goldbach Conjecture up to 4×10^{18} [4].

Then, since $f(n) = f(n + 3 - 3) = f(p + q - 3) = f(p) + f(q) - f(3)$, we obtain that $f(n) = 1$ or $f(n) = n$ by the induction hypothesis. \square

Indeed, those can be extended up to $n \leq 4 \times 10^{18} - 3$.

Lemma 2.3. *The results in Lemma 2.1 can be extended up to n even and $n < 10^{10}$.*

Proof. It is enough to investigate $f(2^r)$ with $r \leq 33$. Note that $k \cdot 2^r + 1$ with $k < 2^r$ is called Proth number. If a Proth number is prime, it is called a Proth prime. It is verified that there exists an odd integer $k \leq 4141$ such that $k \cdot 2^r + 1$ is a Proth prime for $1 \leq r \leq 1000$ in The On-Line Encyclopedia of Integer Sequences (OEIS, <https://oeis.org/A057778>), although the infinitude of Proth primes is not yet proved [6].

Then, $k \cdot 2^r + 1$ is an odd prime and

$$f(k) f(2^r) = f((k \cdot 2^r + 1) + 2 - 3) = f(k \cdot 2^r + 1) + f(2) - f(3).$$

Thus, we are done by Lemmas 2.1 and 2.2. \square

If the Goldbach Conjecture and the infinitude of Proth primes for all exponents were proved, Theorem 1.3 could be easily proved. But, neither of them has not yet been proved, so that we need other strategy. In the following lemma, $v_p(n)$ means the exponent of p in the prime factorization of n when p is a prime and n is a positive integer. The set H was defined by Spiro and the numerical verification of the Goldbach Conjecture was up to 2×10^{10} at that time. We would call the set H in the lemma the *Spiro set*.

Lemma 2.4. *Let*

$$H = \{n \mid v_p(n) \leq 1 \text{ if } p > 1000; v_p(n) \leq \lfloor 9 \log_p 10 \rfloor - 1 \text{ if } p < 1000\}.$$

For any integer $m > 10^{10}$, there is an odd prime $q \leq m - 1$ such that $m + q \in H$.

Proof. This lemma is the consequence of [1, Lemma 2.4] which follows the proof of [7, Lemma 5]. \square

Lemma 2.5 ([2, 3, 8]). *Almost every even positive integer is expressible as the sum of two primes.*

Lemma 2.6. *The restricted function $f|_H$ is the identity function or a constant function on H .*

Proof. Assume $f(n) = n$ for $n = 2, 3, 5, 7, 11$. If $n < 10^{10}$, then $f(n) = n$ from Lemmas 2.2 and 2.3. Let $n \in H$ with $n \geq 10^{10}$ and assume that $f(m) = m$ for all $m \in H$ with $m < n$. If n is not a prime power, then $f(n) = f(a)f(b)$ with $(a, b) = 1$ and $a, b > 1$. Since $f(a) = a$ and $f(b) = b$ by the induction hypothesis, $f(n) = n$.

Now, if n is a prime power, then n is a prime by the definition of H . If $n = 6k - 1$, then consider $n + 7 - 3 = 6k + 3$. Since

$$f(n + 4) = f(6k + 3) = f(n + 7 - 3) = f(n) + f(7) - f(3)$$

and $6k + 3$ can be factored into the product of two smaller integers, $f(n) = n$.

Similarly, if $n = 6k + 1$, then

$$f(n + 2) = f(6k + 3) = f(n + 5 - 3) = f(n) + f(5) - f(3)$$

yields $f(n) = n$.

By the same reasoning, we can conclude that $f(n) = 1$ if $f(2) = f(3) = f(5) = f(7) = f(11) = 1$.

\square

Lemma 2.7 ([7, Lemma 7]). *For any positive integer n , put*

$$H_n = \begin{cases} \{mn : m \in H, (m, n) = 1\} & \text{if } 2 \mid n; \\ \{2mn : 2m \in H, (m, n) = 1\} & \text{if } 2 \nmid n. \end{cases}$$

Then H_n satisfies the following properties:

- (1) *Every element of H_n is even.*
- (2) *The set H_n has positive lower density.*

3. PROOFS OF THEOREMS

Let us start to prove Theorem 1.3. Suppose that there exists n for which $f(n) \neq n$. For $kn \in H_n$, we have that

$$f(kn) = f(k) f(n) = k f(n).$$

If $f(kn) = kn$, then $f(n) = f(kn)/k = kn/k = n$, which contradicts. So $f(kn) \neq kn$ for every $kn \in H_n$.

But, if $kn + 3$ with k odd can be represented as a sum of two primes p and q , then

$$f(kn) = f(p + q - 3) = f(p) + f(q) - f(3) = p + q - 3 = kn.$$

Thus, this implies that there exist many counterexamples to the Goldbach Conjecture whose density is positive. But, this contradicts Lemma 2.5. Therefore, $f(n) = n$ for all n .

We can prove Theorem 1.1 in the similar way. First, we have that

$$f(3) = f(2 + 2 - 1) = f(2) + f(2) - f(1),$$

$$f(5) = f(3 + 3 - 1) = f(3) + f(3) - f(1),$$

$$\begin{aligned} f(6) &= f(2) f(3) \\ &= f(5 + 2 - 1) = f(5) + f(2) - f(1). \end{aligned}$$

Let $a = f(2)$, $b = f(3)$, and $c = f(5)$. Then,

$$b = 2a - 1, \quad c = 2b - 1, \quad ab = c + a - 1.$$

Thus,

$$a(2a - 1) = (2(2a - 1) - 1) + a - 1$$

and it becomes

$$a^2 - 3a + 2 = 0.$$

Hence, $a = 1$ or $a = 2$.

Next, we should check $f(2^r)$ as in Lemma 2.3. We can use $k \cdot 2^r - 1$ instead of $k \cdot 2^r + 1$. The list of prime $k \cdot 2^r - 1$ with $0 \leq r \leq 10000$ is in OEIS (<https://oeis.org/A126717>). See also [6].

ACKNOWLEDGMENT

The author would like to thank the Korea Institute for Advanced Study(KIAS) for its support and hospitality.

REFERENCES

- [1] Y.-G. Chen, J.-H. Fang, P. Yuan, and Y. Zheng, On multiplicative functions with $f(p + q + n_0) = f(p) + f(q) + f(n_0)$, *J. Number Theory* **165** (2016), 270–289.
- [2] N. Čudakov, On the Goldbach’s problem, *Dokl. Akad. Nauk SSSR* (1937), 331–334.
- [3] T. Estermann, On the Goldbach’s problem: Proof that almost all even positive integers are sums of two primes, *Proc. London Math. Soc. (2)* **44** (1938), 307–314.
- [4] T. Oliveira e Silva, S. Herzog, and S. Pardi, Empirical verification of the even the Goldbach conjecture and computation of prime gaps up to $4 \cdot 10^{18}$, *Math. Comp.* **83** (2014) 2033–2060.
- [5] P.-S. Park, Additive uniqueness of PRIMES $- 1$ for multiplicative functions, *Int. J. Number Theory*, to appear.
- [6] Proth Search Page, <http://www.prothsearch.com>
- [7] C. A. Spiro, Additive uniqueness sets for arithmetic functions, *J. Number Theory* **42** (1992), 232–246.
- [8] J. G. van der Corput, Sur l’hypothèse de the Goldbach, *Proc. Akad. Wet. Amsterdam* **41** (1938), 76–80.

E-mail address: pspark@kyungnam.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION, KYUNGNAM UNIVERSITY, CHANGWON-SI, 51767,
REPUBLIC OF KOREA